Aberrant MNX1 Expression Associated with t(7;12)(q36;p13) Pediatric Acute Myeloid

Leukemia Induces the Disease Through Altering Histone Methylation

Ahmed Waraky[1,2], Anders Östlund[1], Tina Nilsson[2], Dieter Weichenhan[3], Pavlo Lutsik[3], Marion

Bähr[3], Joschka Hey[3], Jenni Adamsson[2], Mohammad Hamdy Abdelrazak Morsy [4], Susann Li[2],

Linda Fogelstrand[1,2], Christoph Plass[4] and Lars Palmqvist[1,2*]

[1]Department of Laboratory Medicine, Institute of Biomedicine, University of Gothenburg, and

[2]Department of Clinical Chemistry, Sahlgrenska University Hospital, Gothenburg, Sweden.

[3]Division of Cancer Epigenomics, German Cancer Research Center (DKFZ), Heidelberg,

Germany.

[4]Department of Laboratory Medicine, Division of Pathology, Karolinska Institute, Stockholm,

Sweden.

*Corresponding author

Lars Palmqvist MD, PhD

E-mail: lars.palmqvist@clinchem.gu.se

Institute of Biomedicine, University of Gothenburg

Sahlgrenska University Hospital,

SE-413 45 Gothenburg, Sweden

Phone: +46-31-3427829, Fax: +46-31-828458

Acknowledgment

We thank Mohamad Ali and Akram Mendez for help with bioinformatic analysis. Tova

Johansson and Hanna Brissman for help with animals. This work was supported by grants from

the Swedish Cancer Society (CAN2017/461), the Swedish Childhood Cancer Foundation

(PR2014-0125, PR2019-0013 and TJ2019-0053, TJ2022-0017), Wilhelm och Martina

Lundgrens Fond, Assar Gabrielsson Fond and Västra Götalandsregionen (ALFGBG-431881).

The German Funding Agency (DFG) with funding for Collaborative Research Center 1074,

Project B11N to CP. The computations were enabled by resources in project [SNIC 2021/22-

754] provided by the Swedish National Infrastructure for Computing (SNIC) at UPPMAX,

partially funded by the Swedish Research Council through grant agreement no. 2018-05973.

Authors contribution

31   LP, AÖ and AW designed the research study. AW, AÖ, TN, DW, PL, JH, JA, SL, MHM

32   performed laboratory work and result analysis. AW, LF, CP and LP analyzed the combined

33   data and wrote the paper.

Conflict of interest

35   The authors declare no conflict of interest associated with this paper.

Data Availability Statement

37   The data that support the findings of this study are available in the supplementary material of

38   this article, from Gene Expression Omnibus (GEO) database, accession number GSE182168,

39   GSE202137 and GSE205698 and from PRIDE/ProteomeXchange, accession number

40   PDXD034416. Further details and other data that support the findings of this study are

41   available from the corresponding author upon request.

ABSTRACT

Certain subtypes of acute myeloid leukemia (AML) in children have inferior outcome. One of these has a translocation t(7;12)(q36;p13) leading to a *MNX1*::*ETV6* fusion along with high expression of MNX1. Here we identified the transforming event in this AML and possible ways of treatment. Only MNX1 was able to induce AML in mice, with similar gene expression and pathway enrichment to t(7;12) AML patient data. Importantly, this leukemia was only induced in immune incompetent mice using fetal but not adult hematopoietic stem and progenitor cells. The restriction in transforming capacity to cells from fetal liver is in alignment with t(7;12)(q36;p13) AML being mostly seen in infants. Expression of MNX1 led to increased histone 3 lysine 4 mono-, di- and trimethylation, reduction in H3K27me3, accompanied with changes in genome-wide chromatin accessibility and genome expression, likely mediated through MNX1 interaction with the methionine cycle and methyltransferases. MNX1 expression increased DNA damage, depletion of the Lin-/Sca1+/c-Kit+ population and skewing toward the myeloid lineage. These effects, together with leukemia development, was prevented by pretreatment with the S-adenosylmethionine analog Sinefungin. In conclusion, we have shown the importance of MNX1 in development of AML with t(7;12), supporting a rationale for targeting MNX1 and downstream pathways.

INTRODUCTION

Non-random cytogenetic aberrations are often involved in the development of acute myeloid leukemia (AML) and several aberrations can serve as diagnostic markers, prognosis predictors, and impact the choice of therapy [1]. In AML diagnosed in children under the age of 24 months, a chromosomal translocation t(7;12)(q36;p13) with poor prognosis has been reported [2]. There have been contradictory results on the incidence of t(7;12) but recent studies suggest the frequency in children <24 months to be 5-7% [3, 4]. Similarly, different results have been reported regarding the prognosis where recent studies show 20%-43% 3-year event-free survival but with a high relapse rate, ranging from 57-80% [3, 4]. However, the understanding of the mechanisms behind the leukemia transformation of t(7;12) AML remains poorly understood.

The chromosomal break points in t(7;12) have consistently been found to be located close to the Motor neuron and pancreas homeobox 1 (*MNX1*) gene on chromosome 7, and in introns 1 or 2 in the *ETV6* gene in chromosome 12 [5]. The translocation leads to *MNX1* gene activation and in most reported cases also to a *MNX1::ETV6* fusion transcript consisting of exon1 of *MNX1* transcript variant 1 spliced to the remaining ETV6 exons, depending on the location of the break point in ETV6 [6].

MNX1, also known as Homeobox HB9 (*HLXB9*), belongs to the homeobox domain family of transcription factors, with previous studies showing the importance of *MNX1* in motor neuron development [7], pancreas development [8, 9] and in hereditary sacral agenesis [10]. *ETV6,* also known as *TEL* belongs to the ETS-family transcription factors. *ETV6* encodes a transcriptional repressor that plays a critical role in embryonic development and hematopoiesis, where it is essential for normal HSC function and generation of thrombocytes by megakaryocytes [11].

Translocations involving the chromosomal region of 12p13 that result in the rearrangements of the *ETV6* gene are one of the most observed chromosomal abnormalities in human leukemia

with more than 30 reported translocations. These chromosomal translocations can induce

leukemias through the ectopic expression of a proto-oncogene in the vicinity of a chromosomal

translocation [12] or the constitutive activation of the partner protein [13]. In addition, the formation

of ETV6 fusion proteins can result in the modification of the original functions of the

transcription factor [14], or loss of function of the fusion gene, affecting *ETV6* and the partner

gene [15].

The role of the MNX1::ETV6 fusion protein in the development of AML with t(7;12) has not

been established. It is also unclear whether the driver of leukemogenesis is the MNX1::ETV6

fusion protein or overexpression of MNX1. The aim of this study was to assess the

transformation capacity and the molecular mechanism of the MNX1::ETV6 fusion and the

ectopic expression of MNX1 *in vitro* and *in vivo* using murine transplantation models.

MATERIAL AND METHODS (Complete materials and methods in supplement)

Plasmid constructions

The *MNX1, ETV6* and MNX1::ETV6 fusion sequences are listed in Supplement table 1. An HA-tag (36bp) is introduced at the 5' end and via a linker sequence of 24 bp attached to the separate gene sequences where the first ATG is removed. These constructs were cloned into the MSCV-IRES-GFP and MSCV-IRES-YFP vectors (Takara, Japan) #634401 under control of the viral LTR promoter.

Generation of transduced murine bone marrow (BM) cells and transplantation assays

Mice were bred and maintained at the Gothenburg university laboratory for experimental biomedicine Animal Facility (Gothenburg, Sweden) in an SPF environment. Establishment and characterization of BM cell lines following transduction of BM cells with *MNX1, MNX1::ETV6, ETV6 or* empty vector Ctrl have been done as described previously [16]. In brief, BM cell lines were established from BM cells previously treated with 5-fluorouracil (5-FU) from 8-12 weeks old (C57Bl/6) mice (Charles River Laboratories, Inc., Wilmington, Massachusetts, USA) for three days for adult bone marrow (ABM) or from fetal liver at embryonic days 14.5 (E14.5) and maintained in liquid culture (Dulbecco's modified Eagle's medium supplemented with 18% fetal bovine serum, 10 ng/mL human interleukin [IL]-6, 6 ng/mL murine IL-3, and 50 ng/mL murine stem cell factor). All culture media were obtained from Sigma and recombinant growth factors from Peprotech. To generate *MNX1, MNX1::ETV6, ETV6 or* empty vector Ctrl BM cell lines, the BM cells were transduced by cocultivation on irradiated (4000 cGy) E86 producers (ATCC, Manassas, VA, USA) for a period of 2 days in the presence of 5 μg/mL protamine sulfate (Sigma). Cells were sorted for GFP$^+$ and/or YFP$^+$ expression by flow cytometry analysis (fluorescence-activated cell sorting [FACS]) FACSAria (BD Biosciences) and maintained in culture for 5-7 days post transduction

before transplantation in mice. Lethally (8.5 Gy) or sublethally (5.5 Gy) radiated 8-12 weeks

old C57Bl/6 mice received the equivalent of 0.6×10^6 rescue BM cells and/or $0.8\text{-}1\times10^6$

transduced cells via tail vein injection. For immunocompromised NOD.Cg-$Kit^{W\text{-}}$

$^{41J}\,Tyr\,^{+}\,Prkdc^{scid}\,Il2rg^{tm1Wjl}$/ThomJ (*NBSGW*) *mice (*The Jackson Laboratory, Bar Harbor, ME,

USA*)*, cells were transplanted either with no radiation, or after lethal dose of (1.6 Gy) and

sublethal dose of (0.9 Gy). Donor-derived engraftment and reconstitution were monitored by

flow cytometry analysis for GFP$^+$ and/or YFP$^+$ expression in the peripheral blood of the

transplants every 2 weeks. Mice were sacrificed using Isoflurane (Baxter, Deerfield, IL,

USA). Blood counts were analyzed on a Sysmex KX-21 Hematology Analyzer (Sysmex,

Norderstedt, Germany).

Statistical analysis

Two-sided Student's t-test were used for comparisons between different groups in all

experiments, unless stated otherwise. Log-rank test was used to compare survival between mice

groups. Mann-Whitney two-tailed U-test was used for comparison between t(7;12) patients and

normal human bone marrow from Target cohort.

Ethics Statement

All animal experiments have been accepted by the Jordbruksverket and the animal ethics

committee in Gothenburg, Dnr 5.8.18-17008/2021.

RESULTS

MNX1 *induces acute myeloid leukemia in hematopoietic cells of fetal origin*

To investigate the leukemogenic potential of t(7;12), we transduced primary murine (C57BL/6) hematopoietic stem and progenitor cells (HSPC) from either adult bone marrow after 5-FU stimulation (ABM-HSPC) or fetal liver cells at E14.5 (FL-HSPC) with retroviral vectors for expression of the *MNX1::ETV6* fusion, *MNX1*, *ETV6* or empty vector (Ctrl). Expression of *MNX1*, *ETV6* and *MNX1::ETV6* was confirmed in both FL-HSPC and ABM-HSPC (Supplement figure 1a-c). The transduced cells were transplanted into lethally irradiated C57BL/6 mice with rescue bone marrow. Neither *MNX1* overexpression nor *MNX1::ETV6* in ABM-HSPC or FL-HSCP were able to induce leukemia in these mice, as assessed by survival, WBC, hemoglobin, blood smears and spleen size (Supplement figure 2a-d). However, when increasing the percentage of MNX1 chimerism through sublethal radiation with no rescue BM 20% of the mice exhibited signs of malignant transformation after transplantation with FL-HSPC, including high WBCs count, severe anemia and enlarged spleen (Supplement figure 2e-f & 3a-b). To further examine the leukemogenic potential of *MNX1* overexpression and the *MNX1::ETV6* fusion, we repeated the experiment using immune-compromised non-irradiated NOD.Cg-Kit^{W-41J} Tyr^{+} $Prkdc^{scid}$ $Il2rg^{tm1Wjl}$/ThomJ (NBSGW) mice. Within 12-18 weeks after transplant, the *MNX1* mice showed clear signs of leukemia including pallor, weight loss, severe anemia, leukocytosis with a high percentage of *MNX1*-transduced cells, elevated blast cells in blood and bone marrow, enlarged spleen and liver infiltrated by leukemia (Figure 1a-d). Cells from bone marrow showed predominant expression of the c-Kit protein, but not the stem cell marker SCA-1 or more differentiated myeloid markers such as Mac-1, Ly6G1 and Ly6C1, suggesting a poorly differentiated myeloid leukemia (Figure 1d; Supplement figure 3d and 4a). To rule out that the leukemia development in immunocompromised mice was not due to enhanced transplantation efficiency, we reduced the chimerism of *MNX1* cells in NBSGW mice

through the usage of lethal radiation and rescue bone marrow. These mice also developed leukemia but with a slightly longer latency than mice that were sublethally irradiated without rescue BM (Supplement figure 3d). Acute leukemia induction by MNX1 was confirmed by leukemia development after secondary transplant of BM from mice with primary leukemia, both in non-irradiated mice and in sublethally irradiated mice receiving rescue BM (Supplement figure 5a). When NBSGW mice were transplanted with ABM-HSPC transduced with *MNX1*, there were no signs of leukemia six months after the transplant (Figure 1e; Supplement figure 5c-d). Taken together, leukemogenesis was achieved by MNX1 only in the immunocompromised setting with fetal origin of leukemic cells.

MNX1 alters differentiation in favor of myeloid lineage while increasing proliferation and colony replating capacity of the cells

In order to characterize this leukemia model, *in vitro* FL-HSPC (r-FL) transduced with *MNX1*::*ETV6*, *MNX1*, *ETV6* or empty retroviral vector control were assessed for their immunophenotype. Both *MNX1* and *MNX1*::*ETV6* altered differentiation in favor of myeloid lineage, with *MNX1* showing the most prominent effects. *MNX1* increased Mac-1 and Ly6C+ cells, accompanied by depletion of the Lin-/Sca1+/c-kit+ (LSK) population, while *MNX1*::*ETV6* only increased Ly6C+ cells (Figure 2a-b; Supplement figure 4b-d). Ectopic expression of MNX1 reduced the progenitor MEP population without significantly affecting CMP or GMP (Supplement figure 5a). Additionally, *MNX1* increased GEMM colonies with a concomitant reduction in BFU colonies and increased both CFU replating and proliferation capacity (Figure 2c-d; Supplement figure 6f-h). In ABM-HSPC *in vitro* cells (r-ABM), *MNX1* had similar effects but to a lower extent, and *MNX1*::*ETV6* fusion had no effects (Supplement figure 6 c-e).

MNX1 induces DNA damage

To investigate the molecular pathway through which *MNX1* is mediating its leukemogenic effect, differential gene expression between *MNX1* FL-HSCP leukemic NSG BM cells and FL-HSPC transduced with empty vector (Ctrl) was assessed with RNA-sequencing (Figure 2e Supplement figure 7). Gene ontology biological (GO) pathway and gene set enrichment (GSEA) analyses revealed that the highest enriched pathways in *MNX1* cells involved DNA damage, cell cycle, chromatin organization, methylation of histones, metabolic processes, megakaryocyte, and myeloid cell differentiation pathways (Figure 2e). Similar results were seen when using BM taken from mice transplanted with FL-HSCP with empty vector as control (Supplement figure 8). The effect on DNA damage was confirmed through examination of γH2AX foci induction. *MNX1* and to a lower extent *MNX1::ETV6* induced a higher number of γH2AX foci, indicative of higher DNA damage, in *in vitro* FL-HSPC as well as in ABM-HSPC (r-Fl & r-ABM) (Figure 3a-b; Supplement figure 9a-b). In both *MNX1* and *MNX1*::*ETV6* transduced FL-HSPC (r-FL), the DNA damage was accompanied with a transient G1 cell cycle accumulation and fewer cells in S-phase (Figure 3c; Supplement figure 9c). No such effect was observed in ABM-HSPC (r-ABM), where the G1 cells were replaced by a Sub-G1 peak suggestive of apoptotic cells after 3-4 weeks of transduction (Supplement figure 9d). Using Annexin-V and DAPI staining a 3.5-4-fold increase in apoptosis was induced by *MNX1* in ABM-HSPC (r-ABM), but not seen in FL-HSPC (r-FL) (Figure 3d-e). Following up the consequences of such pronounced DNA damage, we showed that BM from NSG leukemia mice exhibited an increase in the amount of DNA or a hyperploidy in comparison with the BM from Ctrl mice as indicated by increased DNA index (Supplement figure 10a-b). An interesting candidate for mediating DNA damage identified in several differentially regulated pathways was the centrosomal protein 164 (*Cep164*) (Supplement table 2). Increased expression of *Cep164* was confirmed with qPCR in both *MNX1* transduced FL-HSPC and leukemia BM

(Supplement figure 10c). Furthermore, MNX1 increased binding to the promoter of *Cep164*

was detected by ChIP-qPCR (Supplement figure 10c), along with altered binding of histone

modifications that can change CEP164 expression namely H3K4me3 and H3K27me3 in the

same promoter region (Supplement figure 10d).

MNX1 alters histone modifications globally affecting chromatin accessibility

Since MNX1 induced histone modifications H3K4me3 and H3K27me3 at the *Cep164*

promoter, we assessed global histone modifications induced by MNX1. Western blotting

showed that MNX1 altered H3K4me3, H3K27me3, and both mono- and di-methylation of the

H3K4 *in vitro* and in BM from NSG leukemic mice (Figure 4a; Supplement figure 10e). Co-

immunoprecipitation experiments (Co-IP) confirmed the association of MNX1 to histone H3

(Supplement figure 11a). Antibody-guided chromatin tagmentation sequencing (ACT-Seq) was

then used for mapping genome-wide distribution of the histone modifications (Figure 4b).

H3K27me3 histone modification showed fewer accessible regions in MNX1 (BM from NSG

leukemic mice) compared to Ctrl (FL-HSPC transduced with empty vector) (FDR $\leq$ 0.05, log2

fold change $\geq$ |1|), involving mainly distal intergenic regions followed by promoter and other

intronic regions (Figure 4b). The consequences of these histone modifications on chromatin

accessibility were investigated with ATAC-Seq. MNX1 leukemic BM cells exhibited increased

number of differentially accessible chromatin regions in comparison with Ctrl FL-HSPC (FDR

$\leq$ 0.05, log2 fold change $\geq$ |1|) (Figure 4c), mainly involving promoters, followed by distal

intergenic and intronic regions, very similar to the pattern seen for H3K27me3 (Figure 4b).

Pathway analysis of genes annotated to the differentially accessible regions from ATAC-Seq

(FDR $\leq$ 0.05, log2 fold change $\geq$ |1|) revealed similar pathway enrichment to the RNA-Seq,

with high enrichment in metabolic pathways, myeloid cell differentiation, erythrocyte

differentiation, cytoskeleton organization, cell cycle process and apoptotic cell process

(Supplement figure 11b). Further analysis through integrating ATAC-Seq data with in-silico

predicted transcription factor binding sites by DiffTF package, revealed 130 differentially

activated sites (FDR < 0.05) between MNX1 leukemic BM cells and Ctrl FL-HSPC (Table 1,

Supplement figure 11c). These were enriched in pathways of myeloid/erythrocyte

differentiation, cellular metabolism, G1/S-phase transition of cell cycle, histone methylation,

DNA methylation and DNA damage response (Table1, Supplement figure 11c). A significant

correlation between the accessible chromatin regions identified by ATAC-Seq and

differentially expressed genes by RNA-Seq was shown at r_S=0.47 and (p ≤ 0.01) (Figure 4d).

In conclusion, MNX1 induces global histone modifications that are affecting chromatin

accessibility and inducing differential gene regulation.

MNX1 alters the methylation of histone H3

To understand the mechanism for the MNX1-induced global histone modifications, we

performed mass spectrometry analysis, to study proteins in association with MNX1. MNX1

associated proteins were pulled down after Co-IP from *in vitro* cells with anti-HA antibody.

Pathway analysis using STRING for protein-protein interactions for the identified proteins

revealed a high enrichment for methylation pathway proteins MAT2A, MAT2B and AHCY,

(Figure 5a, Table 2), in addition to several S-adenosylmethionine (SAM) dependent methyl

transferases and their downstream targets (Supplement table 3). Co-IP confirmed the

association of MNX1 to MAT2A and AHCY both *in vitro* and in NSG BM leukemic cells

(Figure 5b; Supplement figure 12a), with no effect on protein expression as shown in parallel

with western blot (Supplement figure 12b). Furthermore, MNX1 overexpression increased the

concentration of S-adenosylhomocysterin (SAH) and reduced free methionine in both FL cells

and leukemia NSG BM cells (Supplement figure 12c-e). In support of a MNX1 role in

methylation pathway and in altering histone methylation, MNX1 pulled down with Co-IP and

incubated with recombinant Histone H3 resulted in methylation of Histone H3 (Figure 5c-d,

Supplement figure 12f-g).

MNX1 induced leukemia in mice and human pediatric t(7;12) AML show similar gene expression and pathway enrichment

To validate the similarity between the AML developed in our mouse model with human t(7;12) AML, we used our RNA-sequencing data on leukemic cells from mice and retrieved RNA-sequencing data from pediatric t(7;12) AML patient samples from the children's oncology group (COG)–National Cancer Institute (NCI) TARGET AML initiative data set. Differential gene expression in mouse leukemia was determined by comparing *MNX1* FL-HSCP leukemic NSG BM cells and FL-HSPC transduced with empty vector and differential gene expression in human AML was determined by comparing pediatric t(7;12) AML patient samples with normal human BM. Comparing the differential gene expression between the mouse AML with MNX1 expression and the t(7;12) patient data revealed close to 50% overlapping differential gene expression (Figure 5e; Supplement table 4). These included increased expression of the genes *MNX1*, *cKIT*, *CEP164*, *AHCY, MAT2A* and *MAT2B* (Supplement figure 13). Pathway enrichment analysis of the t(7;12) patient using gene set enrichment (GSEA) analyses revealed several common enriched pathways including; DNA damage, H3K4methylation, monocyte differentiation, apoptotic processes and metabolic processes (Figure 5f and supplement figure 14). Interestingly, only the H3K4me3 methylation, and no other histone methylation pathway was enriched in the GSEA analysis from t(7;12) AML patients. Thus, our data highlight the biological significance for MNX1 expression in pediatric t(7;12) AML and suggesting a mechanism where MNX1 is mediating its oncogenic effects through aberrant methylation resulting in histone modifications resulting in leukemia transformation.

The effects of MNX1 on methylation are crucial for leukemogenesis

To confirm the importance of this effect for MNX1-induced leukemia, we used the natural nucleoside analogue of SAM, Sinefungin, which acts as a competitor and accordingly a pan-

methyltranferase inhibitor[17]. Adding 5µM of Sinefungin to MNX1-transduced FL-HSPC

partially or completely prevented the effects of MNX1 on histone modifications (Figure 6a,

Supplement figure 15a), DNA damage induction (Figure 6b; Supplement figure 15b), cell cycle

distribution (Supplement figure 15c), myeloid differentiation and LSK depletion (Figure 6c-d).

Furthermore, when MNX1 FL-HSPC was pretreated *in vitro* with Sinefungin and then

transplanted into NBSGW mice there were no signs of leukemia development (Figure 6e;

Supplement figure 15d-e), despite maintained high MNX1 expression (Supplement figure 15f)

and continuous presence of viable transplanted cells in blood (Supplement figure 16a) and. To

investigate the effect of Sinefungin treatment on MNX1 induced differential gene expression,

RNA sequencing was done before and after treatment. Differentially expressed genes (FDR $\leq$

0.05, log2 fold change $\geq |1.5|$) from MNX1 cells after treatment (MNX+S) clustered with

MNX1 (Figure 6f), and most of the differentially expressed genes by MNX1 overexpression

remained at similar level after treatment (Figure 6f), with limited number of genes altered in

MNX1 differential gene expression after treatment (Supplement figure 12b)

DISCUSSION

The t(7;12) has only been reported in children diagnosed with AML before the age of 24 months. The function of this translocation in inducing infant leukemia and the reason for its absence in adult leukemia is unknown. In the current study using a murine model, we showed that ectopic expression of MNX1 rather than the *MNX1*::*ETV6* fusion was able to initiate and drive leukemogenesis. Our data suggest a mechanism through which MNX1 is mediating the leukemogenic effect through aberrant methylation that results in histone modifications and DNA damage.

The malignant transformation mediated by MNX1 overexpression in our mouse model matched the criteria for AML [18], compatible with an AML without maturation. This mouse MNX1 driven leukemia had a high degree of differentially expressed gene that overlap with the genes gene expression signature and pathway enrichment that is seen in human AML with t(7;12). It has also been shown that MNX1 clearly has oncogenic properties in both human and murine hematopoietic cells by inducing a myeloid biased perturbed hematopoietic differentiation and premature senescence [19]. This fits well with the properties seen in our mouse leukemia model with both block in differentiation and induced cell cycle arrest. Similar gene expression and pathway enrichment has also been shown between human cells with an engineered t(7;12) translocation, which results in high expression of MNX1, and human t(7;12) AML which suggests a common gene expression program induced by MNX1 in hematopoietic cells [20, 21]. Our data showed that ectopic expression of MNX1 was able to induce AML using HSPC from fetal origin but not from adult bone marrow. One possible reason for this was the dramatic induction of apoptosis seen in the hematopoietic progenitor cells from adult bone marrow, prohibiting leukemic transformation. The higher susceptibility for apoptosis and DNA damage induced by MNX1 is concordant with the presence of naturally occurring DNA damage in the adult stem cells [22]. Possibly, the balance between fetal and adult stem cell programs affects the

transforming ability of the cells upon overexpression of MNX1. Lin28b has been shown as a

key regulator of the self-renewing capacity characteristic of fetal but not adult hematopoietic

stem cells [23]. *LIN28B* is, together with *MNX1*, a signature gene expressed in all pediatric t(7;12)

AML and not seen in other AML subtypes [24]. Lin28b was seen expressed in our MNX1 induced

mouse leukemia (Supplement table 2), which suggests that transformation by MNX1 might be

dependent on Lin28b for its transforming effects or perhaps more likely indicates that the fetal

hematopoietic program is needed for MNX1 oncogenesis. However, the expression of *Lin28b*

in our leukemia and *LIN28B* in human t(7;12) AML is contradictory to the finding that Lin28b

suppresses MLL-ENL fusion driven leukemogenesis, also typically seen in infants leukemia [25].

But what is intriguing is that the MLL-ENL fusion's potential to initiate leukemia development

peaks during neonatal development and drops dramatically in adult mice [25]. The importance of

the development stage of cells that can be transformed into leukemia including the leukemia

phenotype as well as intrinsic properties of progenitor populations have been shown in several

studies[26-28]. Other factors that differ between fetal and adult hematopoetic stem cells and that

might affect their propensity for transformation are metabolic demand and cell cycle profile [29-

31]. Even though MNX1 overexpression induced leukemia in the cells of fetal origin, the fusion

MNX1::*ETV6* by itself did not induce leukemia. This finding is in line with the previously

reported inability to induce transformation with *MNX1*::*ETV6* (*HLXB9/TEL*) *in vitro* and the

paucity of transgenic mouse models of AML with *MNX1*::*ETV6* [32]. In our study, the

development of leukemia induced by MNX1 expression was primarily seen in

immunocompromised NSG mice. Thus, the adaptive B- and T-cell immune response might be

enough to eradicate cells with overexpression of MNX1 and prevent leukemia development.

This may be yet another clue to the development of AML with t(7;12) in very young children,

typically before six months of age when the immune system is still under development [4-6].

Our studies of MNX1 expression *in vitro* revealed an induction of DNA damage both in FL-

HSPC and ABM-HSPC, which could contribute to the observed skewed differentiation towards

myeloid lineage evident by the depletion of LSK and MEP population and increased Mac1+

and Ly6C+ population. The influence of DNA damage in stem cells on differentiation was first

demonstrated in melanocyte stem cells, where ionizing radiation triggered differentiation into

mature melanocytes [33]. In hematopoietic stem cells, DNA damage can induce differentiation

towards lymphoid or myeloid lineage and may be the reason for the skewing towards myeoloid

differentiation in the aging hematopoietic system [34-37].

We found MNX1 to associate with members of the methionine cycle, including MAT2A,

AHCY and MAT2B, in addition to several downstream SAM-dependent methyl transferases.

Methionine is an essential amino acid that is converted to the universal methyl donor SAM,

which upon the donation of its methyl group is converted to SAH. This reaction is catalyzed by

methionine adenosyl transferases (MAT). SAM is used as a cofactor in most methylation

reactions and provides the activated methyl group for methylation of proteins, including

histones, DNA, RNA and lipids. These methylation events are highly dependent on methionine

metabolism, with alterations in methionine showing profound effects on DNA and histone

methylation [38, 39]. The role of another homeodomain protein (MSX1) in recruiting

methyltransferase to regulate gene expression and chromatin structure through histone

modifications has been in the differentiation of myoblasts [40, 41]. During neural development,

MNX1 binding to loci on chromatin are enriched for H3K4me1 and H3K4me3 [42]. Therefore,

the binding and association of MNX1 with methyltranferases and members of methionine cycle,

and the subsequent change in chromatin structure and histone modifications might be the

physiological role for MNX1 during differentiation [43, 44]. We conclude that the abnormal

expression of MNX1 and subsequent effect on the methionine cycle and chromatin structure in

hematopoietic cells acts as the driver of leukemia transformation, supported by the inhibition

of the phenotype by the SAM analog Sinefungin.

In conclusion, our results provide the biological and clinical significance for MNX1 as an

epigenetic regulator in pediatric t(7;12) AML. Given that many epigenetic modifications are

chemically reversible, the inhibition of MNX1 ectopic expression or its downstream effects in

t(7;12) could provide a foundation for alternative treatment options to improve outcome.

REFERENCES

1.   Lagunas-Rangel FA, Chavez-Valencia V, Gomez-Guijosa MA, Cortes-Penagos

C. Acute Myeloid Leukemia-Genetic Alterations and Their Clinical Prognosis. Int J Hematol

Oncol Stem Cell Res. 2017;11(4):328-339.

2.   von Bergh AR, van Drunen E, van Wering ER, et al. High incidence of

t(7;12)(q36;p13) in infant AML but not in infant ALL, with a dismal outcome and ectopic

expression of HLXB9. Genes, chromosomes & cancer. 2006;45(8):731-739.

3.   Bolouri H, Farrar JE, Triche T, Jr., et al. The molecular landscape of pediatric

acute myeloid leukemia reveals recurrent structural alterations and age-specific mutational

interactions. Nature medicine. 2018;24(1):103-112.

4.   Espersen ADL, Noren-Nystrom U, Abrahamsson J, et al. Acute myeloid leukemia

(AML) with t(7;12)(q36;p13) is associated with infancy and trisomy 19: Data from Nordic

Society for Pediatric Hematology and Oncology (NOPHO-AML) and review of the literature.

Genes, chromosomes & cancer. 2018;57(7):359-365.

5.   Beverloo HB, Panagopoulos I, Isaksson M, et al. Fusion of the homeobox gene

HLXB9 and the ETV6 gene in infant acute myeloid leukemias with the t(7;12)(q36;p13).

Cancer Res. 2001;61(14):5374-5377.

6.   Tosi S, Mostafa Kamel Y, Owoka T, Federico C, Truong TH, Saccone S.

Paediatric acute myeloid leukaemia with the t(7;12)(q36;p13) rearrangement: a review of the

biological and clinical management aspects. Biomark Res. 2015;3(21.

7.   Thaler J, Harrison K, Sharma K, Lettieri K, Kehrl J, Pfaff SL. Active suppression

of interneuron programs within developing motor neurons revealed by analysis of

homeodomain factor HB9. Neuron. 1999;23(4):675-687.

8.   Li H, Arber S, Jessell TM, Edlund H. Selective agenesis of the dorsal pancreas in

mice lacking homeobox gene Hlxb9. Nature genetics. 1999;23(1):67-70.

9.   Harrison KA, Thaler J, Pfaff SL, Gu H, Kehrl JH. Pancreas dorsal lobe agenesis

and abnormal islets of Langerhans in Hlxb9-deficient mice. Nat Genet. 1999;23(1):71-75.

10.   Ross AJ, Ruiz-Perez V, Wang Y, et al. A homeobox gene, HLXB9, is the major

locus for dominantly inherited sacral agenesis. Nat Genet. 1998;20(4):358-361.

11.   Hock H, Shimamura A. ETV6 in hematopoiesis and leukemia predisposition.

Semin Hematol. 2017;54(2):98-104.

12.   Rawat VP, Cusan M, Deshpande A, et al. Ectopic expression of the homeobox

gene Cdx2 is the transforming event in a mouse model of t(12;13)(p13;q12) acute myeloid

leukemia. Proc Natl Acad Sci U S A. 2004;101(3):817-822.

13.   Golub TR, Barker GF, Lovett M, Gilliland DG. Fusion of PDGF receptor beta to

a novel ets-like gene, tel, in chronic myelomonocytic leukemia with t(5;12) chromosomal

translocation. Cell. 1994;77(2):307-316.

14.   Zelent A, Greaves M, Enver T. Role of the TEL-AML1 fusion gene in the

molecular pathogenesis of childhood acute lymphoblastic leukaemia. Oncogene.

2004;23(24):4275-4283.

15.   De Braekeleer E, Douet-Guilbert N, Morel F, Le Bris MJ, Basinko A, De

Braekeleer M. ETV6 fusion genes in hematological malignancies: a review. Leuk Res.

2012;36(8):945-961.

16.   Arabanian LS, Johansson P, Staffas A, et al. The endothelin receptor type A is a

downstream target of Hoxa9 and Meis1 in acute myeloid leukemia. Leukemia research.

2018;75(61-68.

17.   Zhang J, Zheng YG. SAM/SAH Analogs as Versatile Tools for SAM-Dependent

Methyltransferases. ACS Chem Biol. 2016;11(3):583-597.

18.   Kogan SC, Ward JM, Anver MR, et al. Bethesda proposals for classification of

nonlymphoid hematopoietic neoplasms in mice. Blood. 2002;100(1):238-245.

19. Ingenhag D, Reister S, Auer F, et al. The homeobox transcription factor HB9

induces senescence and blocks differentiation in hematopoietic stem and progenitor cells.

Haematologica. 2019;104(1):35-46.

20. Ragusa D, Ciciro Y, Federico C, et al. Engineered model of t(7;12)(q36;p13)

AML recapitulates patient-specific features and gene expression profiles. Oncogenesis.

2022;11(1):50.

21. Nilsson T, Waraky A, Ostlund A, et al. An induced pluripotent stem cell

t(7;12)(q36;p13) acute myeloid leukemia model shows high expression of MNX1 and a block

in differentiation of the erythroid and megakaryocytic lineages. Int J Cancer. 2022;151(5):770-

782.

22. Biechonski S, Yassin M, Milyavsky M. DNA-damage response in hematopoietic

stem cells: an evolutionary trade-off between blood regeneration and leukemia suppression.

Carcinogenesis. 2017;38(4):367-377.

23. Copley MR, Babovic S, Benz C, et al. The Lin28b-let-7-Hmga2 axis determines

the higher self-renewal potential of fetal haematopoietic stem cells. Nat Cell Biol.

2013;15(8):916-925.

24. Balgobind BV, Van den Heuvel-Eibrink MM, De Menezes RX, et al. Evaluation

of gene expression signatures predictive of cytogenetic and molecular subtypes of pediatric

acute myeloid leukemia. Haematologica. 2011;96(2):221-230.

25. Okeyo-Owuor T, Li Y, Patel RM, et al. The efficiency of murine MLL-ENL-

driven leukemia initiation changes with age and peaks during neonatal development. Blood

Adv. 2019;3(15):2388-2399.

26. Ugale A, Norddahl GL, Wahlestedt M, et al. Hematopoietic stem cells are

intrinsically protected against MLL-ENL-mediated transformation. Cell Rep. 2014;9(4):1246-

1255.

27. Rowe RG, Lummertz da Rocha E, Sousa P, et al. The developmental stage of the

hematopoietic niche regulates lineage in MLL-rearranged leukemia. The Journal of

experimental medicine. 2019;216(3):527-538.

28. Copley MR, Eaves CJ. Developmental changes in hematopoietic stem cell

properties. Exp Mol Med. 2013;45(e55.

29. Manesia JK, Xu Z, Broekaert D, et al. Highly proliferative primitive fetal liver

hematopoietic stem cells are fueled by oxidative metabolic pathways. Stem Cell Res.

2015;15(3):715-721.

30. Bowie MB, Kent DG, Dykstra B, et al. Identification of a new intrinsically timed

developmental checkpoint that reprograms key hematopoietic stem cell properties. Proc Natl

Acad Sci U S A. 2007;104(14):5878-5882.

31. Chen C, Yu W, Tober J, et al. Spatial Genome Re-organization between Fetal and

Adult Hematopoietic Stem Cells. Cell Rep. 2019;29(12):4200-4211 e4207.

32. Wildenhain S, Ruckert C, Rottgers S, et al. Expression of cell-cell interacting

genes distinguishes HLXB9/TEL from MLL-positive childhood acute myeloid leukemia.

Leukemia. 2010;24(9):1657-1660.

33. Inomata K, Aoto T, Binh NT, et al. Genotoxic stress abrogates renewal of

melanocyte stem cells by triggering their differentiation. Cell. 2009;137(6):1088-1099.

34. Mandal PK, Rossi DJ. DNA-damage-induced differentiation in hematopoietic

stem cells. Cell. 2012;148(5):847-848.

35. Beerman I, Bhattacharya D, Zandi S, et al. Functionally distinct hematopoietic

stem cells modulate hematopoietic lineage potential during aging by a mechanism of clonal

expansion. Proc Natl Acad Sci U S A. 2010;107(12):5465-5470.

36.   Cho RH, Sieburg HB, Muller-Sieburg CE. A new mechanism for the aging of

hematopoietic stem cells: aging changes the clonal composition of the stem cell compartment

but not individual stem cells. Blood. 2008;111(12):5553-5561.

37.   Scheffold A, Baig AH, Chen Z, et al. Elevated Hedgehog activity contributes to

attenuated DNA damage responses in aged hematopoietic cells. Leukemia. 2020;34(4):1125-

1134.

38.   Kaiser P. Methionine Dependence of Cancer. Biomolecules. 2020;10(4):

39.   Sanderson SM, Gao X, Dai Z, Locasale JW. Methionine metabolism in health and

cancer: a nexus of diet and precision medicine. Nat Rev Cancer. 2019;19(11):625-637.

40.   Wang J, Abate-Shen C. The MSX1 homeoprotein recruits G9a methyltransferase

to repressed target genes in myoblast cells. PLoS One. 2012;7(5):e37647.

41.   Wang J, Kumar RM, Biggs VJ, et al. The Msx1 Homeoprotein Recruits Polycomb

to the Nuclear Periphery during Development. Dev Cell. 2011;21(3):575-588.

42.   Sun M-a, Ralls S, Wu W, et al. Homeobox transcription factor MNX1 is crucial

for restraining the expression of pan-neuronal genes in motor neurons. bioRxiv.

2021;2021.2008.2007.455331.

43.   Leotta CG, Federico C, Brundo MV, Tosi S, Saccone S. HLXB9 gene expression,

and nuclear location during in vitro neuronal differentiation in the SK-N-BE neuroblastoma

cell line. PLoS One. 2014;9(8):e105481.

44.   Dalgin G, Ward AB, Hao le T, Beattie CE, Nechiporuk A, Prince VE. Zebrafish

mnx1 controls cell fate choice in the developing endocrine pancreas. Development.

2011;138(21):4597-4608.

Table 1. In-silico predicted transcription factor binding sites by DiffTF package from ATAC-

Seq showing top differentially activated (FDR < 0.05) transcription factors (TFs) binding

between MNX1 BM from leukemia mice and FL-HSPC Ctrl cells

Up-regulated TFs	weighted_mean Difference	pvalueAdj	Down regulated TFs	weighted_mean Difference	pvalueAdj
Tal1.a	1.294844532	9.30E-79	Fosl2	-0.421333369	1.84E-24
Gata1	0.567577232	1.24E-20	Jun	-0.415408916	2.58E-23
Gata3	0.550998432	1.12E-19	Cebpe	-0.39280302	0.000108
Gata2	0.540274293	1.01E-16	Batf	-0.387629056	1.57E-17
Alx1	0.371423122	0.023026	Fosl1	-0.384618862	4.57E-21
Nf2l2	0.288226747	2.11E-17	Cebpa	-0.353311343	3.14E-06
Zeb1	0.232871599	9.60E-41	Gfi1	-0.349686909	8.73E-08
Tbx5	0.231419764	1.76E-09	Fosb	-0.339609157	5.76E-15
Mafk.s	0.22701829	3.01E-10	Sta5a	-0.339203846	2.94E-09
Nfe2	0.218904994	4.77E-11	Jund	-0.337147615	8.92E-16
Gata6	0.213067314	2.28E-11	Cebpd	-0.323368792	0.000182
Foxo3	0.202788032	0.000274	Gfi1b	-0.308823273	8.94E-12

513 **Table 2.** Identified proteins in top enriched pathways after Co-IP of proteins in complex with

514 MNX1 and mass spectrometry analysis. As determined by STRING using reactome data set

Pathway	Matching proteins
Association with TriC/CCT	Cct8,Cct6a,Tcp1,Cct4
Methylation	Mat2b,Ahcy,Mat2a,Trmt112
RHO GTPases activate PAKs	Ppp1cb,Myh9,Myh11,Myh10,Myh14
Citric acid cycle (TCA cycle)	Cs,Mdh2,Idh3g,Dld
Pyruvate metabolism	Pdk3,Slc16a3,Vdac1,Ldha,Dld
Actin dynamics for phagocytic	Actr2,Arpc2,Myh9,Cyfip2,Arpc3,Myo5a,Actr3
Translation initiation complex	Pabpc1,Eif3e,Rps21,Eif2s3x,Rps12,Rps28
RHO GTPase Effectors	Actr2,Mapk14,Arpc2,Ppp1cb,Myh9,Pfn1,Aurkb,S100a8,Tuba1b,Bub3,Myh11,Cyfip2,Arpc3,Myh10,Ywhaq,Myh14,Kif2a,S100a9,Actr3
Cellular Senescence	Mapk14,Asf1a,Lmnb1,Txn1,H2afz,Hist1h1b,Hist2h2be,Eed
Mitotic Anaphase	Psmd11,Aurkb,Psmd6,Lmnb1,Smc3,Psmd13,Chmp4b,Smc1a,Tuba1b,Rcc1,Bub3,Kif2a
M Phase	Psmd11,Aurkb,Psmd6,Lmnb1,Smc3,Psmd13,Chmp4b,H2afz,Smc1a,Smc4,Tuba1b,Rcc1,Bub3,Hist2h2be,Csnk2a1,Smc2,Kif2a
Metabolism of proteins	Pabpc1,Srp14,Psmd11,Rab10,Ddx5,Aurkb,Psmd6,Vdac2,Ero1l,Eif3e,Mrpl21,Smc3,Psmd13,Cct8,Srp9,Txn1,Trappc3,Cct6a,Cmas,Rab27a,Rpl22l1,Smc1a,Rpl37,Mat2b,Pdia6,Fn1,Rps21,Eif2s3x,Hist2h2ab,Rps12,Tuba1b,Ctbp1,Rpl15,Rpl36,Rpl34,Lgals1,Hist2h2be,Eef1g,Csnk2a1,Vdac1,Wdr5,Rps28,Trmt112,Tcp1,Copa,Gnb4,Ctsd,Cct4,Nop58,Tceb1
Metabolism of RNA	Pabpc1,U2af2,Psmd11,Prpf8,Snrpd3,Ddx5,Psmd6,Wdr46,Psmd13,Rpl22l1,Rpl37,Utp15,Wdr43,Rps21,Rcl1,Rps12,Rpl15,Rpl36,Rpl34,Srsf2,Nhp2l1,Snrpb,Rps28,Srsf3,Ddx6,Hnrnpd,Apobec3,Nop58

 FIGURE LEGENDS

Figure 1. MNX1 induces AML in immunocompromised mice. Kaplan-Meier survival curve

of (NBSGW) mice transplanted with FL-HSPC after retrovirus transduction with either ectopic

expression of *MNX1* (green), *ETV6* (purple), *MNX1::ETV6* fusion (red) or empty vector control

(blue). Results from the transplanted mice (control n = 11, *MNX1*: n = 12, *ETV6* and

MNX1::ETV6 n = 4) were analyzed using the log-rank test (***: significant at $P \leq 0.01$, ns:

non-significant) (a). Mice were euthanized when showing sign of disease or at the end of the

experiment and analyzed for spleen weight, white blood cell number and hemoglobin

concentration (b). Quantification of flow cytometry analysis of BM cells from (NBSGW) mice

with c-Kit expression showing all events. GFP/YFP were used as indicative for MNX1

expression (c). Representative images of H- and E-stained formaldehyde fixed spleen and liver

sections from control and MNX1 mice (d; left-panels). Representative images of Giemsa-

stained peripheral blood smears from Ctrl and MNX1 mice (d; right-panels) (d). Kaplan-Meier

survival curves of (NBSGW) mice transplanted with ABM cells with either ectopic expression

of *MNX1*, *ETV6*, *MNX1::ETV6* fusion or empty vector control after sub-lethal radiation (0.9

Gy) with no rescue bone marrow. Results of control and transfected mice (control n = 5, *MNX1*:

n = 8, *ETV6* and *MNX1::ETV6* n = 3) were analyzed using the log-rank test (ns: non-significant)

(e). Data represents mean ± SD of at least three experiments and is considered significant using

two-sided student t-test (**) at $p \leq 0.01$ and (*) at $p \leq 0.05$, or non-significant (ns) at $p > 0.05$.

FL: Fetal liver, BM: bone marrow, ABM: Adult Bone Marrow, Ctrl: empty vector control,

Fusion: *MNX1::ETV6* fusion, HGB: Hemoglobin, LSK: Lin⁻Sca-1⁺c-Kit⁺, NSG: NOD.Cg-

$Kit^{W-41J} Tyr^{+} Prkdc^{scid} Il2rg^{tm1Wjl}$/ThomJ (NBSGW), H&E: hematoxylin and eosin. GFP: Green

fluorescence protein, YFP: Yellow fluorescence protein.

Figure 2. MNX1 alters differentiation and proliferation of FL-HSPC. Quantification of the

flowcytometry analysis of *in vitro* FL-HSPC (r-FL) transduced with ectopic expression of

MNX1, *ETV6*, *MNX1::ETV6* fusion or empty vector control with the indicated antibodies

represented as percentage of population (a-b). Number of CFU colonies after transduction of

FL-HSPC with replating for five consecutive weeks (c). MTT proliferation assay of transduced

invitro FL-HSPC (r-FL) cells (d). Gene set enrichment analysis (GSEA) using the Gene

ontology biological pathways (GO) gene set showing normalized enrichment score for

pathways with NOM p-value ≤ 0.05 from leukemia BM cells with *MNX1* ectopic expression in

comparison with FL-HSPC with empty vector control (e). Data represents mean $\pm$ SD of at least

three experiments and is considered significant using two-sided student t-test (**) at $p \leq 0.01$

and (*) at $p \leq 0.05$.

HSPC: hematopoietic stem and progenitor cells, FL: fetal liver, Ctrl: empty vector control,

Fusion: *MNX1::ETV6* fusion, LSK: Lin$^-$Sca-1$^+$c-Kit$^+$, lin+: lineage positive, r-FL: retroviral

transduced *in vitro* FL cells, CFU: colony forming unit.

Figure 3. MNX1 induces DNA damage. Immunofluorescence of *in vitro* FL-HSPC (r-FL)

cells stained with γH2AX –FITC antibody (green) and counterstained with DAPI (blue) (a).

Quantification of the number of γH2AX foci/cell. At least 50 cells were counted (b).

Quantification of cell cycle distribution from flowcytometry analysis represented as fold

difference relative to control (c). Representative dot plots from the flow cytometry analysis of

FL-HSPC (r-FL) and ABM-HSPC (r-ABM) after double staining with Annexin/V and DAPI

for apoptotic analysis (d). Quantification of the flowcytometry analysis represented as fold

difference relative to control (e). Data represents mean $\pm$ SD of at least three experiments and

is considered significant using two-sided student t-test (**) at $p \leq 0.01$ and (*) at $p \leq 0.05$.

FL: fetal liver, ABM: adult bone marrow, Ctrl: empty vector control, Fusion: *MNX1::ETV6*

fusion, r-FL: retroviral *invitro* FL cells, r-BM: retroviral *in vitro* ABM cells e.A: early

apoptosis, l-.A: late apoptosis.

Figure 4. MNX1 alters histone methylation. Image quantification of western blot analysis of

H3K4me1, H3K4me2, H3K4me3 and H3K27me3 as fold difference relative to loading control

(a). Heatmap of differentially bound H3K27me3 from BM of leukemic mice (MNX1) versus

control FL-HSPC transduced with empty vector (Ctrl) as determined by ACT-Seq and

annotation of different enriched regions for H3K27me3 ACT-Seq (b). Results were considered

at the log fold-change cutoff (logFC) of $\geq |1|$ and false discovery rate (FDR) of ≤ 0.05. Heatmap

of differentially accessible regions from BM of leukemic mice (MNX1) versus control FL-

HSPC transduced with empty vector (Ctrl) as determined by ATAC-Seq and annotation of

different enriched regions for ATAC-Seq (c). Results were considered at the log fold-change

cutoff (logFC) of $\geq |1|$ and false discovery rate (FDR) of ≤ 0.05. Scatter plot showing the

correlation between differentially expressed genes from RNA-Seq expression analysis and

genes with differentially accessible regions from ATAC-Seq analysis (d). Results were

considered at the log fold-change cutoff (logFC) of $\geq |1|$ and false discovery rate (FDR) of

≤ 0.05.

Figure 5. MNX1 associates with proteins from the methionine cycle and show similar gene

expression and pathway enrichment with human t(7;12) AML. Pathway enrichment

analysis for the identified proteins after Co-IP and mass spectrometry using STRING with

reactome data set, showing strength of the pathway as Log10 observed proteins/background in

the respective pathway (a). The binding of AHCY and MAT2A was detected by

immunoprecipitation of MNX1 using HA-antibody followed by western blot using AHCY and

MAT2A antibodies. ETV6 with HA tag was used as a negative control. Total protein input used

as loading control (b). Recombinant Histone H3.1and H3.3 were subjected to an in vitro

methyltransferase reaction using MNX1 complex pulled down with HA antibody from FL-

HSPC transduced with MNX1(M), in comparison with FL-HSPC with empty vector control

(C), in the presence of SAM and DTT. The reactions were terminated by boiling in SDS sample

buffer. Separation of samples in 12% SDS-PAGE was followed by immunoblotting (IB) with

mono, di and tri methyl-lysines antibody. Reblotting was made for detection of HA and total

Histone H3. As indicated, negative controls were obtained by omitting Histone H3, or the pulled

down immune complex (c). Quantification of the in vitro methyltransferase reaction (d). Data

represents mean ± SD of at least three experiments and is considered significant using two-

sided student t-test (**) at $p \leq 0.01$ and (*) at $p \leq 0.05$. Graph showing differentially expressed

genes with same or opposite regulation when comparing genes expression data from t(7;12)

AML patients from the TARGET database and RNA-seq data from BM from mice with MNX1

induced leukemia. Differentially expressed genes were selected based on a LogFC ±1 and FDR

≤ 0.05 (e). Gene set enrichment analysis (GSEA) using the Gene ontology biological pathways

(GO), showing normalized enrichment score for common pathways between t(7;12) AML

patients and the MNX1 induced leukemia in mice with NOM p-value ≤ 0.05 (b).

r-FL: fetal liver, Ctrl: empty vector control, WB: Western blot, IP: immunoprecipitation. SAM:

S-Adenosyl methionine, DTT: Dithiothreitol.

Figure 6. Sinefungin rescued MNX1-induced phenotype. Protein expression levels of

H3K4me3, H3k27me3, H3K4me1 and H3K4me2 in *invitro* FL-HSPC (r-FL) cells with either

MNX1 overexpression or empty vector (Ctrl) after treatment with vehicle or 5μM Sinefungin

(Ctrl+S, MNX1+S) as determined by Western blotting. Actin served as loading control (a).

Quantification of the Western blot analysis in Supplement figure 9a. Quantification of the

number of γH2AX foci/cell. At least 50 cells were counted. Data represented as fold difference

relative to control (b). Quantification of flow cytometry analysis of the cells with the indicated

antibodies. Data represented as fold difference relative to control (c and d). Data represents

mean ± SD of at least three experiments and is considered significant using two-sided student

t-test between MNX1 and MNX1+S (**) at $p \leq 0.01$ and (*) at $p \leq 0.05$. Kaplan-Meier survival

curves of (NBSGW) mice transplanted with FL cells after retrovirus transduction with either

ectopic expression of MNX1 or empty vector (Ctrl) after treatment with vehicle or 5μM

Sinefungin (MNX1+S, Ctrl+S). Results of MNX1 (n = 7) and Sinefungin treated MNX1 cells

(MNX1+S) (n = 6) were analyzed using the log-rank test (***: significant at $P \leq 0.01$) (e).

LogFC heat map of downregulated (blue) and upregulated (red) differentially expressed genes

of FL cells with *MNX1* ectopic expression (MNX) in comparison with FL cells with empty

vector (Ctrl), with 5μM Sinefungin (Ctrl+S, MNX+S) or without treatment (Ctrl, MNX1),

showing clustering and similarity between the samples. Results were considered at the log fold-

change cutoff (LogFC) $\geq$ |1.5| and FDR $\leq$ 0.05 (f).

a

Ctrl Ctrl+S MNX MNX+S KDa

H3K27me3 — 17

H3K4me3 — 17

H3K4me1 — 17

H3K4me2 — 17

Actin — 50

b

Fold difference relative to control

Ctrl Ctrl+S MNX1 MNX1+S

*

c

Fold difference relative to control

Mac1
CD115
Ly6G
Ly6C

Ctrl Ctrl+S MNX1 MNX1+S

*

d

Fold difference relative to control

Ctrl Ctrl+S MNX1 MNX1+S

**

e

Survival (%)

weeks

Ctrl, n=7
MNX1, n=7
Ctrl+S, n=7
MNX+S, n=6

***

f

Ctrl+S_3 Ctrl_3 Ctrl_2 Ctrl_1 Ctrl+S_2 Ctrl+S_1 MNX+s_3 MNX_3 MNX+s_2 MNX+s_1 MNX_2 MNX_1