Mnx1 Induces Leukemia Transformation Through Altering Histone Methylation in a Model of Pediatric Acute Myeloid Leukemia with t(7;12)(q36;p13)

Ahmed Waraky[1,2], Anders Östlund[1], Tina Nilsson[2], Dieter Weichenhan[4], Pavlo Lutsik[4], Marion Bähr[4], Joschka Hey[4], Jenni Adamsson[2], Mohammad Hamdy Abdelrazak Morsy [3], Susann Li[2], Linda Fogelstrand[1,2], Christoph Plass[4] and Lars Palmqvist[1,2]*

[1]Department of Laboratory Medicine, Institute of Biomedicine, University of Gothenburg, and

[2]Department of Clinical Chemistry, Sahlgrenska University Hospital, Gothenburg, Sweden.

[3]Department of laboratory medicine, division of Pathology, Karolinska institutet, Stockholm, Sweden.

[4]Division of Cancer Epigenomics, German Cancer Research Center (DKFZ), Heidelberg, Germany.

*Corresponding author

Lars Palmqvist MD, PhD

E-mail: lars.palmqvist@clinchem.gu.se

Institute of Biomedicine, University of Gothenburg

Sahlgrenska University Hospital,

SE-413 45 Gothenburg, Sweden

Phone: +46-31-3427829, Fax: +46-31-828458

ABSTRACT

Certain subtypes of acute myeloid leukemia (AML) in children have still inferior outcome. One of these AML subtypes has a translocation t(7;12)(q36;p13), always leading to high expression of MNX1 and often to *MNX1*::*ETV6* fusion expression. Here we identified the transforming event in this AML and possible ways to target them. Only MNX1 was able to induce AML in mice, and this was observed using hematopoietic stem and progenitor cells derived from fetal origin but not from adult bone marrow. The restriction in the transforming capacity to cells from fetal liver origin is in concordance with the fact that t(7;12)(q36;p13) AML is mostly restricted to infants. Ectopic expression of MNX1 led to increase of H3K4me1, H3K4me2 and H3K4me3, reduction in H3K27me3, accompanied with changes in genome-wide chromatin accessibility and genome expression, likely mediated through MNX1 interaction with the methionine cycle and different methyltransferases. MNX1 expression resulted in increased DNA damage, depletion of the Lin-/Sca1+/c-Kit+ population and skewing toward the myeloid lineage. These effects, together with leukemia development, was prevented by the S-adenosylmethionine analog Sinefungin. In conclusion, we have shown the importance of MNX1 in leukemia development in AML with t(7;12), supporting a rationale for targeting MNX1 and downstream pathways.

INTRODUCTION

Non-random cytogenetic aberrations are often involved in the development of acute myeloid leukemia (AML) and several aberrations can serve as diagnostic markers, prognosis predictors, and impact the choice of therapy (1). In AML diagnosed in children under the age of 24 months, a chromosomal translocation t(7;12)(q36;p13) with poor prognosis has been reported (2). There have been contradictory results on the incidence of t(7;12) but recent studies suggest the frequency in children <24 months to be 5-7% (3, 4). Similarly, different results have been reported regarding the prognosis where recent studies show 20%-43% 3-year event-free survival but with a high relapse rate, ranging from 57-80% (3, 4). However, the understanding of the mechanisms behind the leukemia transformation of t(7;12) AML remains poorly understood.

The chromosomal break points in t(7;12) have consistently been found to be located close to the Motor neuron and pancreas homeobox 1 (*MNX1)* gene on chromosome 7, and in introns 1 or 2 in the *ETV6* gene in chromosome 12 (5). The translocation leads to *MNX1* gene activation and in most reported cases also to a *MNX1::ETV6* fusion transcript consisting of exon1 of *MNX1* transcript variant 1 spliced to the remaining ETV6 exons, depending on the location of the break point in ETV6 (6).

MNX1, also known as Homeobox HB9 (*HLXB9*), belongs to the homeobox domain family of transcription factors, with previous studies showing the importance of *MNX1* in motor neuron development (7), pancreas development (8, 9) and in hereditary sacral agenesis (10). *ETV6,* also known as *TEL* belongs to the ETS-family transcription factors. *ETV6* encodes a transcriptional repressor that plays a critical role in embryonic development and hematopoiesis, where it is essential for normal HSC function and generation of thrombocytes by megakaryocytes (11).

Translocations involving the chromosomal region of 12p13 that result in the rearrangements of

the *ETV6* gene are one of the most commonly observed chromosomal abnormalities in human

leukemia with more than 30 reported translocations. These chromosomal translocations can

induce leukemias through the ectopic expression of a proto-oncogene in the vicinity of a

chromosomal translocation (12) or the constitutive activation of the partner protein (13). In

addition, the formation of ETV6 fusion proteins can result in the modification of the original

functions of the transcription factor (14), or loss of function of the fusion gene,

affecting *ETV6* and the partner gene (15).

The role of the MNX1::ETV6 fusion protein in the development of AML with t(7;12) has not

been established. It is also unclear whether the driver of leukemogenesis is the MNX1::ETV6

fusion protein or overexpression of MNX1. The aim of this study was to assess the

transformation capacity and the molecular mechanism of the MNX1::ETV6 fusion and the

ectopic expression of MNX1 *in vitro* and *in vivo* using murine transplantation models.

 RESULTS

 MNX1 *induces acute myeloid leukemia in hematopoietic cells of fetal origin*

To investigate the leukemogenic potential of t(7;12), we transduced primary murine (C57BL/6)

hematopoietic stem and progenitor cells (HSPC) from either adult bone marrow after 5-FU

stimulation (ABM-HSPC) or fetal liver cells at E14.5 (FL-HSPC) with retroviral vectors for

expression of the *MNX1::ETV6* fusion, *MNX1*, *ETV6* or empty vector (Ctrl). Expression of

MNX1, *ETV6* and *MNX1*::*ETV6* was confirmed in both FL-HSPC and ABM-HSPC

(Supplement figure 1a-c). The transduced cells were transplanted into lethally irradiated

C57BL/6 mice with rescue bone marrow. Neither *MNX1* overexpression nor *MNX1::ETV6* in

ABM-HSPC or FL-HSCP were able to induce leukemia in these mice, as assessed by survival,

WBC, hemoglobin, blood smears and spleen size (Supplement figure 2a-d). However, when

increasing the percentage of MNX1 chimerism through sublethal radiation with no rescue BM

20% of the mice exhibited signs of malignant transformation after transplantation with FL-

HSPC, including high WBCs count, severe anemia and enlarged spleen (Supplement figure 2e-

f & 3a-b). To further examine the leukemogenic potential of *MNX1* overexpression and the

MNX1::*ETV6* fusion, we repeated the experiment using immune-compromised non-irradiated

NOD.Cg-Kit^{W-41J} Tyr^{+} $Prkdc^{scid}$ $Il2rg^{tm1Wjl}$/ThomJ (NBSGW) mice. Within 12-18 weeks after

transplant, the *MNX1* mice showed clear signs of leukemia including pallor, weight loss,

hunched posture, severe anemia, leukocytosis with a high percentage of *MNX1*-transduced

cells, elevated blast cells in blood and bone marrow, enlarged spleen and liver infiltrated by

leukemia (Figure 1a-c). Cells from bone marrow showed predominant expression of the

myeloid marker c-Kit but not the stem cell marker SCA-1 or more differentiated myeloid

markers such as MAC-1, LY6G1 and LY6C1, suggesting a poorly differentiated myeloid

leukemia (Figure 1d-e). To rule out that the leukemia development in immunocompromised

mice was not due to enhanced transplantation efficiency, we reduced the chimerism of *MNX1*

cells in NBSGW mice through the usage of lethal radiation and rescue bone marrow. These

mice also developed leukemia but with a slightly longer latency than mice that were sublethally

irradiated without rescue BM (Supplement figure 3d). Acute leukemia induction by MNX1

was confirmed by leukemia development after secondary transplant of BM from mice with

primary leukemia, both in non-irradiated mice and in sublethally irradiated mice receiving

rescue BM (Supplement figure 3e-f). When NBSGW mice were transplanted with ABM-HSPC

transduced with *MNX1*, there were no signs of leukemia six months after the transplant

(Supplement figure 4). Taken together, in line with t(7;12) AML occurring solely in infants,

leukemogenesis was achieved by MNX1 only in the immunocompromised setting with fetal

origin of leukemic cells.

115 *MNX1 alters differentiation in favor of myeloid lineage while increasing proliferation and*
colony replating capacity of the cells

In order to characterize this infant leukemia model, FL-HSPC transduced with *MNX1::ETV6*,

MNX1, *ETV6* or empty retroviral vector control were assessed for their immunophenotype.

Both *MNX1* and *MNX1::ETV6* altered differentiation in favor of myeloid lineage, with *MNX1*

showing the most prominent effects. *MNX1* increased Mac-1 and Ly6C+ cells, accompanied

by depletion of the Lin-/Sca1+/c-kit+ (LSK) population, while *MNX1::ETV6* only increased

Ly6C+ cells (Figure 2a-c). Ectopic expression of MNX1 reduced the progenitor MEP

population without affecting CMP or GMP (Supplement figure 5a). Additionally, *MNX1*

increased GEMM colonies with a concomitant reduction in BFU colonies and increased both

CFU replating and proliferation capacity (Figure 2d-f; Supplement figure 5e-f). In ABM-

HSPC, *MNX1* had similar effects but to a lower extent, and *MNX1::ETV6* fusion had no effects

(Supplement figure 5 b-d).

MNX1 induces DNA damage

To investigate the molecular pathway through which *MNX1* is mediating its leukemogenic

effect, differential gene expression between *MNX1* FL-HSCP leukemic BM cells and FL-HSPC

transduced with empty vector (Ctrl) was assessed with RNA-sequencing (Figure 3a-d,

Supplement figure 6b). Gene ontology biological (GO) pathway and gene set enrichment

(GSEA) analyses revealed that the highest enriched pathways in *MNX1* cells involved DNA

damage, cell cycle, chromatin organization, methylation of histones, metabolic processes,

megakaryocyte, and myeloid cell differentiation pathways (Figure 3d; Supplement figure 6b).

Similar results were seen when using BM taken from mice transplanted with FL-HSCP with

empty vector as control (Supplement figure 6c-e). The effect on DNA damage was confirmed

through examination of γH2AX foci induction. *MNX1* and to a lower extent *MNX1::ETV6*

induced a higher number of γH2AX foci, indicative of higher DNA damage, in FL-HSPC as

well as in ABM-HSPC (Figure 4a; Supplement figure 7a). In both *MNX1* and *MNX1::ETV6*

*t*ransduced FL-HSPC, the DNA damage was accompanied with a transient G1 cell cycle

accumulation and fewer cells in S-phase (Figure 4b, Supplement figure 7b). No such effect was

observed in ABM-HSPC, where the G1 cells were replaced by a Sub-G1 peak suggestive of

apoptotic cells after 3-4 weeks of transduction (Supplement figure 7b). Using Annexin-V and

DAPI staining a 3.5-4-fold increase in apoptosis was induced by *MNX1* in ABM-HSPC, but

not seen in FL-HSPC (Figure 4c). Following up the consequences of such pronounced DNA

damage, we showed that BM from leukemia mice exhibited an increase in the amount of DNA

or a hyperploidy in comparison with the BM from Ctrl mice as indicated by increased DNA

index (Figure 4d, supplement figure 7c).

An interesting candidate for mediating DNA damage identified in several differentially

regulated pathways was the centrosomal protein 164 (*Cep164*). Increased expression of *Cep164*

was confirmed with qPCR in both *MNX1* transduced FL-HSPC and leukemia BM (Figure 5a).

Furthermore, MNX1 increased binding to the promoter of *Cep164* was detected by ChIP-qPCR

(Figure 5b), along with increased binding to histone modifications that can alter CEP164

expression namely H3K4me3 and H3K27me3 (Supplement figure 7d).

MNX1 alters histone modifications globally affecting chromatin accessibility

Since MNX1 induced histone modifications H3K4me3 and H3K27me3 at the *Cep164*

promoter, we assessed global histone modifications. Western blotting showed that MNX1

altered H3K4me3, H3K27me3, and both mono- and di-methylation of the H3K4 (Figure 5c,

supplement figure 7e), and co-immunoprecipitation experiments (Co-IP) confirmed the

association of MNX1 to histone H3 (Supplement figure 7f). Antibody-guided chromatin

tagmentation sequencing (ACT-Seq) was then used for mapping genome-wide distribution of

the histone modifications (Figure 5d). H3K27me3 histone modification showed fewer

accessible regions in MNX1 compared to Ctrl in FL-HSPC (FDR ≤ 0.05, log2 fold change ≥

|1|), involving mainly distal intergenic regions followed by promoter and other intronic regions

(Figure 5d). The consequences of these histone modifications on chromatin accessibility were

investigated with ATAC-Seq. MNX1 leukemic BM cells exhibited increased number of

differentially accessible chromatin regions in comparison with Ctrl FL-HSPC (FDR ≤ 0.05,

log2 fold change ≥ |1|) (Figure 5e), mainly involving promoters, followed by distal intergenic

and intronic regions, very similar to the pattern seen for H3K27me3 (Figure 5d-e). Pathway

analysis of genes annotated to the differentially accessible regions from ATAC-Seq (FDR ≤

0.05, log2 fold change ≥ |1|) revealed similar pathway enrichment to the RNA-Seq, with high

enrichment in metabolic pathways, myeloid cell differentiation, erythrocyte differentiation,

cytoskeleton organization, cell cycle process and apoptotic cell process (Supplement figure 7g).

Further analysis through integrating ATAC-Seq data with in-silico predicted transcription

factor binding sites by DiffTF package, revealed 130 differentially activated sites (FDR < 0.05)

between MNX1 leukemic BM cells and Ctrl FL-HSPC (Table 1, Supplement figure 8a). These

were enriched in pathways of myeloid/erythrocyte differentiation, cellular metabolism, G1/S-phase transition of cell cycle, histone methylation, DNA methylation and DNA damage response (Table1, Supplement figure 8a). A significant correlation between the accessible chromatin regions identified by ATAC-Seq and differentially expressed genes by RNA-Seq was shown at r_S=0.47 and (p ≤ 0.01) (Supplement figure 8b). In conclusion, MNX1 induces global histone modifications that are affecting chromatin accessibility and inducing differential gene regulation.

The effects of MNX1 on methylation are crucial for leukemogenesis

In order to understand the mechanism for the MNX1-induced global histone modifications, we performed mass spectrometry analysis, to study proteins in association with MNX1. MNX1 associated proteins were pulled down after Co-IP with anti-HA antibody. Pathway analysis using STRING for protein-protein interactions for the identified proteins revealed a high enrichment for methylation pathway proteins MAT2A, MAT2B and AHCY, (Figure 5f, Table 2), in addition to several SAM-dependent methyl transferases and their downstream targets (Supplement table 1). Co-IP confirmed the association of MNX1 to MAT2A and AHCY (Figure 5g, supplement figure 8c), with no effect on protein expression as shown in parallel with Western blot (Supplement figure 8d). Furthermore, MNX1 overexpression increased the concentration of SAH and reduced free methionine in both FL cells and leukemia BM cells (Figure 5h; Supplement figure 8e-f). In support of MNX1 role in methylation pathway and in altering histone methylation, MNX1 pulled down with Co-IP and incubated with recombinant Histone H3 resulted in methylation of Histone H3 (Figure 5i, Supplement figure 8g). To confirm the importance of this interaction for MNX1-induced leukemia, we used the natural nucleoside analogue of S-adenosylmethionine Sinefungin, which acts as a SAM competitor and

accordingly a pan-methyltranferase inhibitor (16). Adding 5µM of Sinefungin to MNX1-

transduced FL-HSPC partially or completely prevented the effects of MNX1 on histone

modifications (Figure 6a, Supplement figure 9a), DNA damage induction (Figure 6b;

Supplement figure 9b), cell cycle distribution (Supplement figure 9c), myeloid differentiation

and LSK depletion (Figure 6c-d). Furthermore, when MNX1 FL-HSPC treated *in vitro* with

Sinefungin were transplanted into NBSGW mice there were no signs of leukemia development

(Figure 6e; Supplement figure 9d-e), despite the continuous presence of viable transplanted

cells in the blood (Supplement figure 9g) and maintained MNX1 expression (Supplement figure

9f). To confirm that Sinefungin treatment prevented leukemia transformation through inhibition

of methyltransferase activity in complex with MNX1 and not through the gene expression

downstream of methylation pathway, RNA sequencing was done before and after treatment.

Differentially expressed genes (FDR ≤ 0.05, log2 fold change ≥ |1.5|) from MNX1 cells after

treatment (MNX+S) clustered with MNX1 (Figure 6f), and most of the differentially expressed

genes by MNX1 overexpression remained at similar level after treatment (Supplement figure

9h). Taken together, our results showed the effect of MNX1 on methylation is crucial for

leukemogenesis.

DISCUSSION

The t(7;12) rearrangement has only been reported in children diagnosed with AML before the

age of 24 months. The function of this translocation in inducing infant leukemia and the reason

for its absence in adult leukemia is unknown. In the current study using a murine model, we

showed that ectopic expression of MNX1 rather than the *MNX1*::*ETV6* fusion was able to

initiate and drive leukemogenesis. Our data suggest a mechanism through which MNX1 is

mediating the leukemogenic effect through aberrant methylation that results in histone

modifications and DNA damage.

The malignant transformation mediated by MNX1 overexpression in our mouse model matched

the criteria for AML (17), compatible with an AML without maturation where the development

of leukemia was rapidly fatal in both primary transplanted mice as well as in secondary

recipients. Our data showed that ectopic expression of MNX1 was only able to induce AML

using HSPC from fetal origin and not from adult bone marrow, primarily in

immunocompromised mice. This is interesting in light of the clinical observation that this AML

subtype is only found in very young children with peak at six months where the immune system

is still under development (4, 6). One possible explanation for the incapability of MNX1 ectopic

expression to induce leukemia from adult progenitor hematopoietic cells was further suggested

in our study, as the expression of MNX1 in these cells resulted in dramatic apoptotic induction,

which was not observed in hematopoietic progenitor cells from fetal origin. This could be due

to the difference in gene expression, metabolic demand, cell cycle profiles between adult and

fetal stem cells (18-20), the different hematopoietic lineage and progenitor population between

fetal and adult stem cells (21), or due to the presence of naturally occurring DNA damage in

the adult stem cells (22), thus making the cells more susceptible for apoptotic induction with

additional DNA damage induced by MNX1.

Studying the mechanism of MNX1 *in vitro* revealed an induction in DNA damage both in FL-

HSPC and ABM-HSPC, which could contribute to the observed skewed differentiation towards

myeloid lineage evident by the depletion of LSK and MEP population and increased Mac1+

and Ly6C+ population. The notion that DNA damage in stem cells might influence

differentiation was first demonstrated in melanocyte stem cells, where irreparable DNA

damage, caused by ionizing radiation triggered differentiation into mature melanocytes (23).

With respect to hematopoietic stem cells, DNA damage was found to induce differentiation

towards lymphoid lineage (24), or myeloid lineage (25). This led to the assumption that DNA

damage might be responsible for the skewed differentiation in the aging hematopoietic system

towards myelopoiesis (24-27). The consequence of this DNA damage was evident in the BM

from the leukemia mice with increased DNA content associated with hyperploidy. This shows

a resemblance to the clinical presentation where t(7;12) is associated with gain of chromosomal

material, typically trisomy 8 and 19 (3, 4).

Molecularly we found MNX1 to associate with members of the methionine cycle, including

MAT2A, AHCY and MAT2B, in addition to several downstream SAM-dependent methyl

transferases. Methionine is an essential amino acid that is converted to the universal methyl

donor S-adenosyl-methionine (SAM), which upon the donation of its methyl group is converted

to S-adenosyl-homocysteine (SAH). This reaction is catalyzed by methionine adenosyl

transferases (MAT). SAM is used as a cofactor in most methylation reactions and provides the

activated methyl group for methylation of proteins, including histones, DNA, RNA and lipids.

These methylation events are highly dependent on methionine metabolism, with alterations in

methionine showing profound effects on DNA and histone methylation (28, 29). Here, the

consequences effect of MNX1 on methylation and methionine cycle was observed through

altered histone methylation, namely H3K4me mono-, di- and trimethylation and H3K27me3,

which resulted in altered chromatin accessibility. The role of MNX1 in methylation/methyl-

transferase was shown by; a decline in the free methionine concentration; elevated SAH

concentration; the association of MNX1 to MAT2A and AHCY in addition to several

downstream SAM-dependent methyl transferase; the global alteration of histone methylation

H3K4me and H2K27me3; methylation of recombinant Histone H3 by MNX1 complex *invitro;*

pathway analysis of RNA-seq and ATAC-Seq which showed enrichment in the methylation

pathway, in addition to protein metabolism, a downstream pathway of methionine cycle (29,

30) and nucleic acid/lipid metabolism a downstream pathway of the methionine-closely linked

folate cycle (31, 32). This is in addition to the complete or partial prevention of MNX1

phenotype through the usage of the SAM/SAH analog Sinefungin (33, 34).

The role of homeodomain proteins in recruiting methyltransferase to regulate gene expression

and chromatin structure through histone modifications has been shown previously as a

mechanism of regulating the differentiation of myoblasts by MSX1 (35, 36). This suggests that

the binding and association of MNX1 with methyltranferases and members of methionine cycle,

and the subsequent change in chromatin structure and histone modifications, could be the

physiological role for MNX1 during differentiation (37, 38), and that the abnormal expression

of MNX1 in hematopoietic cells might act as the driver of leukemia transformation. This is

further supported by recent findings showing MNX1 binding to loci on chromatin to be

enriched for H3K4me1 and H3K4me3 during neural development (39).

In conclusion, our results provide the biological and clinical significance for MNX1 as an

epigenetic regulator in pediatric t(7;12) AML. Given that many epigenetic modifications are

chemically reversible, the inhibition of MNX1 ectopic expression or its downstream effects in

t(7;12) could provide a foundation for alternative treatment options to improve outcome.

MATERIAL AND METHODS (Complete materials and methods in supplement)

Plasmid constructions

The *MNX1*, *ETV6* and MNX1::ETV6 fusion sequences are listed in Supplement table 2. An HA-tag (36bp) is introduced at the 5' end and via a linker sequence of 24 bp attached to the separate gene sequences where the first ATG is removed. These constructs were cloned into the MSCV-IRES-GFP and MSCV-IRES-YFP vectors (Takara, Japan) #634401 under control of the viral LTR promoter.

Generation of transduced murine bone marrow (BM) cells and transplantation assays

Mice were bred and maintained at the Gothenburg university laboratory for experimental biomedicine Animal Facility (Gothenburg, Sweden) in an SPF environment. Establishment and characterization of BM cell lines following transduction of BM cells with *MNX1, MNX1::ETV6, ETV6 or* empty vector Ctrl have been done as described previously (40). In brief, BM cell lines were established from BM cells previously treated with 5-fluorouracil (5-FU) from 8-12 weeks old (C57Bl/6) mice (Charles River Laboratories, Inc., Wilmington, Massachusetts, USA) for three days for adult bone marrow (ABM) or from fetal liver at embryonic days 14.5 (E14.5) and maintained in liquid culture (Dulbecco's modified Eagle's medium supplemented with 18% fetal bovine serum, 10 ng/mL human interleukin [IL]-6, 6 ng/mL murine IL-3, and 50 ng/mL murine stem cell factor). All culture media were obtained from Sigma and recombinant growth factors from Peprotech. To generate *MNX1, MNX1::ETV6 , ETV6 or* empty vector Ctrl BM cell lines, the BM cells were transduced by cocultivation on irradiated (4000 cGy) E86 producers (ATCC, Manassas, VA, USA) for a period of 2 days in the presence of 5 μg/mL protamine sulfate (Sigma). Cells were sorted for GFP$^+$ and/or YFP$^+$ expression by flow cytometry analysis (fluorescence-activated cell sorting [FACS]) FACSAria (BD Biosciences) and maintained in culture for 5-7 days post transduction before

transplantation in mice. Lethally (8.5 Gy) or sublethally (5.5 Gy) radiated 8-12 weeks old

C57Bl/6 mice received the equivalent of 0.6×10^6 rescue BM cells and/or $0.8\text{-}1 \times 10^6$ transduced

cells via tail vein injection. For immunocompromised NOD.Cg-$Kit^{W\text{-}41J}$ Tyr^{+} $Prkdc^{scid}$ $Il2rg^{tm1Wjl}$/ThomJ (*NBSGW*) *mice (*The Jackson Laboratory, Bar Harbor, ME,

USA)*, cells were transplanted either with no radiation, or after lethal dose of (1.6 Gy) and

sublethal dose of (0.9 Gy). Donor-derived engraftment and reconstitution were monitored by

flow cytometry analysis for GFP[+] and/or YFP[+] expression in the peripheral blood of the

transplants every 2 weeks. Mice were sacrificed using Isoflurane (Baxter, Deerfield, IL,

USA). Blood counts were analyzed on a Sysmex KX-21 Hematology Analyzer (Sysmex,

Norderstedt, Germany).

Immunophenotyping and flowcytometry

Cells were collected 1-2 weeks after transduction. Then they were washed twice with PBS,

blocked with 0.1% FBS, mouse BD FC Block #553142 (BD Biosciences) in PBS for 10 mins

RT. Then incubated with the antibodies for 20 minutes at RT, washed and resuspended in PBS.

The following antibodies were purchased from BD Biosciences. Sca1-V450#560653, Cd3e-

V450#560801, Ter119-V450#560504, CD45R-V450#560473, Ly6G/Ly6C-V450#560454,

Mac1-V450#560455, Ly6G-PeCy7#560601, B220-V450#560472, CD127-V450#561205,

cKit-APC#553356, Sca1-PeCy7#558162, CD45-APCCy7#561037, Ter119-APC#561033,

CD8a-APCCy7#561967, CD4-APCCy7#561830. The following from (ThermoFisher

Scientific). CD115-APC#17-1152-80, CD64-APCCy7#47-0641-82.

For cell cycle analysis, the proportion of cells in G_1 and G_2 phases were measured by staining

with DAPI (BD Biosciences, #564901) as described previously (41). Briefly, cells were washed

with PBS, and stained DAPI solution 1:100 containing RNase A (250 µg/ml) and 0.25% Triton

X100 for 15 min at room temperature. The samples were processed using FACSAria (BD

Biosciences), and the data were analyzed with the Cell Quest software (BD Biosciences) or with flowing software v 2.5.1 (Turku bioscience center, Finland).

Apoptosis analysis was studied using Annexin V/DAPI method (Annexin V-FLUOS staining kit, Roche, Mannheim, Germany) according to the manufacturer's protocol using AnexinV-APC antibody#550474 (BD Biosciences) and DAPI. Flow cytometric analysis was immediately performed using FACSAria.

Statistical analysis

Two-sided Student's t-test were used for comparisons between different groups in all experiments. Log-rank test was used to compare survival between mice groups.

ACKNOWLEDGMENT

We thank Mohamad Ali and Akram Mendez for help with bioinformatic analysis. Tova Johansson and Hanna Brissman for help with animals. This work was supported by grants from the Swedish Cancer Society (20 0925 PjF), the Swedish Childhood Cancer Foundation (PR2021-0025) and Sahlgrenska University Hospital (ALFGBG-431881) to LP and Swedish Childhood Cancer Foundation (TJ2022-0017), Wilhelm och Martina Lundgrens Fond and Assar Gabrielsson Fond to AW. The German Funding Agency (DFG) with funding for Collaborative Research Center 1074, Project B11N to CP. The computations were enabled by resources in project [SNIC 2021/22-754] provided by the Swedish National Infrastructure for Computing (SNIC) at UPPMAX, partially funded by the Swedish Research Council through grant agreement no. 2018-05973.

LP, AÖ and AW designed the research study. AW, AÖ, TN, DW, PL, JH, JA, SL, MHM performed laboratory work and result analysis. AW, LF, CP and LP analyzed the combined data and wrote the paper.

Conflict of interest

The authors declare no conflict of interest associated with this paper.

Data Availability Statement

The data that support the findings of this study are available in the supplementary material of this article, from Gene Expression Omnibus (GEO) database, accession number GSE182168, GSE202137 and GSE205698 and from PRIDE/ProteomeXchange, accession number PDXD034416. Further details and other data that support the findings of this study are available from the corresponding author upon request.

Ethics Statement

All animal experiments have been accepted by the Jordbruksverket and the animal ethics

committee in Gothenburg, Dnr 5.8.18-17008/2021.

REFERENCES

1.   Lagunas-Rangel FA, Chavez-Valencia V, Gomez-Guijosa MA, Cortes-Penagos C. Acute Myeloid Leukemia-Genetic Alterations and Their Clinical Prognosis. Int J Hematol Oncol Stem Cell Res. 2017;11(4):328-39.

2.   von Bergh AR, van Drunen E, van Wering ER, van Zutven LJ, Hainmann I, Lonnerholm G, et al. High incidence of t(7;12)(q36;p13) in infant AML but not in infant ALL, with a dismal outcome and ectopic expression of HLXB9. Genes Chromosomes Cancer. 2006;45(8):731-9.

3.   Bolouri H, Farrar JE, Triche T, Jr., Ries RE, Lim EL, Alonzo TA, et al. The molecular landscape of pediatric acute myeloid leukemia reveals recurrent structural alterations and age-specific mutational interactions. Nature medicine. 2018;24(1):103-12.

4.   Espersen ADL, Noren-Nystrom U, Abrahamsson J, Ha SY, Pronk CJ, Jahnukainen K, et al. Acute myeloid leukemia (AML) with t(7;12)(q36;p13) is associated with infancy and trisomy 19: Data from Nordic Society for Pediatric Hematology and Oncology (NOPHO-AML) and review of the literature. Genes, chromosomes & cancer. 2018;57(7):359-65.

5.   Beverloo HB, Panagopoulos I, Isaksson M, van Wering E, van Drunen E, de Klein A, et al. Fusion of the homeobox gene HLXB9 and the ETV6 gene in infant acute myeloid leukemias with the t(7;12)(q36;p13). Cancer Res. 2001;61(14):5374-7.

6.   Tosi S, Mostafa Kamel Y, Owoka T, Federico C, Truong TH, Saccone S. Paediatric acute myeloid leukaemia with the t(7;12)(q36;p13) rearrangement: a review of the biological and clinical management aspects. Biomark Res. 2015;3:21.

7.   Thaler J, Harrison K, Sharma K, Lettieri K, Kehrl J, Pfaff SL. Active suppression

of interneuron programs within developing motor neurons revealed by analysis of

homeodomain factor HB9. Neuron. 1999;23(4):675-87.

8.   Li H, Arber S, Jessell TM, Edlund H. Selective agenesis of the dorsal pancreas in

mice lacking homeobox gene Hlxb9. Nature genetics. 1999;23(1):67-70.

9.   Harrison KA, Thaler J, Pfaff SL, Gu H, Kehrl JH. Pancreas dorsal lobe agenesis

and abnormal islets of Langerhans in Hlxb9-deficient mice. Nat Genet. 1999;23(1):71-5.

10.   Ross AJ, Ruiz-Perez V, Wang Y, Hagan DM, Scherer S, Lynch SA, et al. A

homeobox gene, HLXB9, is the major locus for dominantly inherited sacral agenesis. Nat

Genet. 1998;20(4):358-61.

11.   Hock H, Shimamura A. ETV6 in hematopoiesis and leukemia predisposition.

Semin Hematol. 2017;54(2):98-104.

12.   Rawat VP, Cusan M, Deshpande A, Hiddemann W, Quintanilla-Martinez L,

Humphries RK, et al. Ectopic expression of the homeobox gene Cdx2 is the transforming event

in a mouse model of t(12;13)(p13;q12) acute myeloid leukemia. Proc Natl Acad Sci U S A.

2004;101(3):817-22.

13.   Golub TR, Barker GF, Lovett M, Gilliland DG. Fusion of PDGF receptor beta to

a novel ets-like gene, tel, in chronic myelomonocytic leukemia with t(5;12) chromosomal

translocation. Cell. 1994;77(2):307-16.

14.   Zelent A, Greaves M, Enver T. Role of the TEL-AML1 fusion gene in the

molecular pathogenesis of childhood acute lymphoblastic leukaemia. Oncogene.

2004;23(24):4275-83.

15.   De Braekeleer E, Douet-Guilbert N, Morel F, Le Bris MJ, Basinko A, De

Braekeleer M. ETV6 fusion genes in hematological malignancies: a review. Leuk Res.

2012;36(8):945-61.

16. Zhang J, Zheng YG. SAM/SAH Analogs as Versatile Tools for SAM-Dependent

Methyltransferases. ACS Chem Biol. 2016;11(3):583-97.

17. Kogan SC, Ward JM, Anver MR, Berman JJ, Brayton C, Cardiff RD, et al.

Bethesda proposals for classification of nonlymphoid hematopoietic neoplasms in mice. Blood.

2002;100(1):238-45.

18. Manesia JK, Xu Z, Broekaert D, Boon R, van Vliet A, Eelen G, et al. Highly

proliferative primitive fetal liver hematopoietic stem cells are fueled by oxidative metabolic

pathways. Stem Cell Res. 2015;15(3):715-21.

19. Bowie MB, Kent DG, Dykstra B, McKnight KD, McCaffrey L, Hoodless PA, et

al. Identification of a new intrinsically timed developmental checkpoint that reprograms key

hematopoietic stem cell properties. Proc Natl Acad Sci U S A. 2007;104(14):5878-82.

20. Chen C, Yu W, Tober J, Gao P, He B, Lee K, et al. Spatial Genome Re-

organization between Fetal and Adult Hematopoietic Stem Cells. Cell Rep. 2019;29(12):4200-

11 e7.

21. Copley MR, Eaves CJ. Developmental changes in hematopoietic stem cell

properties. Exp Mol Med. 2013;45:e55.

22. Biechonski S, Yassin M, Milyavsky M. DNA-damage response in hematopoietic

stem cells: an evolutionary trade-off between blood regeneration and leukemia suppression.

Carcinogenesis. 2017;38(4):367-77.

23. Inomata K, Aoto T, Binh NT, Okamoto N, Tanimura S, Wakayama T, et al.

Genotoxic stress abrogates renewal of melanocyte stem cells by triggering their differentiation.

Cell. 2009;137(6):1088-99.

24. Mandal PK, Rossi DJ. DNA-damage-induced differentiation in hematopoietic

stem cells. Cell. 2012;148(5):847-8.

25.   Beerman I, Bhattacharya D, Zandi S, Sigvardsson M, Weissman IL, Bryder D, et

al. Functionally distinct hematopoietic stem cells modulate hematopoietic lineage potential

during aging by a mechanism of clonal expansion. Proc Natl Acad Sci U S A.

2010;107(12):5465-70.

26.   Cho RH, Sieburg HB, Muller-Sieburg CE. A new mechanism for the aging of

hematopoietic stem cells: aging changes the clonal composition of the stem cell compartment

but not individual stem cells. Blood. 2008;111(12):5553-61.

27.   Scheffold A, Baig AH, Chen Z, von Lohneysen SE, Becker F, Morita Y, et al.

Elevated Hedgehog activity contributes to attenuated DNA damage responses in aged

hematopoietic cells. Leukemia. 2020;34(4):1125-34.

28.   Kaiser P. Methionine Dependence of Cancer. Biomolecules. 2020;10(4).

29.   Sanderson SM, Gao X, Dai Z, Locasale JW. Methionine metabolism in health and

cancer: a nexus of diet and precision medicine. Nat Rev Cancer. 2019;19(11):625-37.

30.   Parkhitko AA, Jouandin P, Mohr SE, Perrimon N. Methionine metabolism and

methyltransferases in the regulation of aging and lifespan extension across species. Aging Cell.

2019;18(6):e13034.

31.   Zheng Y, Cantley LC. Toward a better understanding of folate metabolism in

health and disease. J Exp Med. 2019;216(2):253-66.

32.   Zarou MM, Vazquez A, Vignir Helgason G. Folate metabolism: a re-emerging

therapeutic target in haematological cancers. Leukemia. 2021;35(6):1539-51.

33.   Fuller RW, Nagarajan R. Inhibition of methyltransferases by some new analogs

of S-adenosylhomocysteine. Biochem Pharmacol. 1978;27(15):1981-3.

34.   Couture JF, Hauk G, Thompson MJ, Blackburn GM, Trievel RC. Catalytic roles

for carbon-oxygen hydrogen bonding in SET domain lysine methyltransferases. J Biol Chem.

2006;281(28):19280-7.

35.   Wang J, Abate-Shen C. The MSX1 homeoprotein recruits G9a methyltransferase

to repressed target genes in myoblast cells. PLoS One. 2012;7(5):e37647.

36.   Wang J, Kumar RM, Biggs VJ, Lee H, Chen Y, Kagey MH, et al. The Msx1

Homeoprotein Recruits Polycomb to the Nuclear Periphery during Development. Dev Cell.

2011;21(3):575-88.

37.   Leotta CG, Federico C, Brundo MV, Tosi S, Saccone S. HLXB9 gene expression,

and nuclear location during in vitro neuronal differentiation in the SK-N-BE neuroblastoma

cell line. PLoS One. 2014;9(8):e105481.

38.   Dalgin G, Ward AB, Hao le T, Beattie CE, Nechiporuk A, Prince VE. Zebrafish

mnx1 controls cell fate choice in the developing endocrine pancreas. Development.

2011;138(21):4597-608.

39.   Sun M-a, Ralls S, Wu W, Demmerle J, Jiang J, Miller C, et al. Homeobox

transcription factor MNX1 is crucial for restraining the expression of pan-neuronal genes in

motor neurons. bioRxiv. 2021:2021.08.07.455331.

40.   Arabanian LS, Johansson P, Staffas A, Nilsson T, Rouhi A, Fogelstrand L, et al.

The endothelin receptor type A is a downstream target of Hoxa9 and Meis1 in acute myeloid

leukemia. Leuk Res. 2018;75:61-8.

41.   Waraky A, Akopyan K, Parrow V, Stromberg T, Axelson M, Abrahmsen L, et al.

Picropodophyllin causes mitotic arrest and catastrophe by depolymerizing microtubules via

insulin-like growth factor-1 receptor-independent mechanism. Oncotarget. 2014;5(18):837992.

 FIGURE LEGENDS

 Figure 1. MNX1 induces AML in immunocompromised mice. Kaplan-Meier survival curve

 of (NBSGW) mice transplanted with FL-HSPC after retrovirus transduction with either ectopic

 expression of *MNX1* (green), *ETV6* (purple), *MNX1::ETV6* fusion (red) or empty vector Ctrl

 (blue). Results from the transplanted mice (control n = 11, *MNX1*: n = 12, *ETV6* and

 MNX1::ETV6 n = 4) were analyzed using the log-rank test (***: significant at $P \leq 0.01$, ns:

 non-significant) (a). Mice were euthanized at the end of the experiment and analyzed for spleen

 weight, white blood cell number and hemoglobin concentration (b). Representative images of

 H and E-stained formaldehyde fixed spleen and liver sections from Ctrl and MNX1 mice (e;

 left-panels). Representative images of Giemsa-stained peripheral blood smears from Ctrl and

 MNX1 mice (e; right-panels) (c). Representative dot plots from the flow cytometry analysis for

 the bone marrow of the euthanized mice at the end of the experiment with the indicated

 antibodies (d and e). Data represent mean ± SD of at least three experiments and is considered

 significant using two-sided student t-test (**) at $p \leq 0.01$ and (*) at $p \leq 0.05$, or non-significant

 (ns) at $p > 0.05$. FL: Fetal liver, BM: bone marrow, Ctrl: control empty vector, Fusion:

 MNX1::ETV6 fusion, HGB: Hemoglobin, LSK: Lin⁻Sca-1⁺c-Kit⁺, NSG: NOD.Cg-*Kit^{W-41J} Tyr*

 $^{+}$ Prkdc^{scid} Il2rg^{tm1Wjl}/ThomJ (NBSGW), HandE: hematoxylin and eosin.

 Figure 2. MNX1 alters differentiation and proliferation of FL-HSPC. Representative dot

 plots from the flow cytometry analysis of FL-HSPC transduced with ectopic expression of

 MNX1, *ETV6*, *MNX1::ETV6* fusion or empty vector with the indicated antibodies (let panel, a-

 c). Blue population represents the positive population. Quantification of the flowcytometry

 analysis represented as percentage of population (right panel, a-c). CFU two weeks after

 transduction of FL-HSPC (d), or weekly number of colonies with relating for five weeks (e).

 MTT proliferation assay of transduced FL-HSPC (f). Data represent mean ± SD of at least three

experiments and is considered significant using two-sided student t-test (**) at $p \leq 0.01$ and (*)

at $p \leq 0.05$. HSPC: hematopoietic stem and progenitor cells, FL: fetal liver, Ctrl: control empty

vector, Fusion: *MNX1::ETV6* fusion, LSK: Lin$^-$Sca-1$^+$c-Kit$^+$), lin+: lineage positive, CFU:

colony forming unit, GM: granulocyte, macrophage, GEMM: granulocyte, erythroid,

macrophage, megakaryocyte, BFU: Burst-forming unit.

Figure 3. Downstream targets of MNX1 overexpression. LogFC heat map of downregulated

(blue) and upregulated (red) differentially expressed genes of leukemia BM cells with *MNX1*

ectopic expression in comparison with FL-HSPC with empty vector (Ctrl). Results were

considered at the log fold-change cutoff (logFC) of $\geq |1|$ and false discovery rate (FDR) of $\leq$

0.05 (a). PCA scatter plot of normalized gene expression using variance Stabilizing

Transformation (b). Volcano plot of differentially expressed genes (c). Gene set enrichment

analysis (GSEA) using the Gene ontology biological pathways (GO) gene set showing

normalized enrichment score for pathways with NOM p-value ≤ 0.05 (d). LogFC: Log fold

change, PCA: principal component analysis, FL: fetal liver, BM: Bone marrow, Ctrl: control

empty vector, Fusion: *MNX1::ETV6* fusion, NSG: NOD.Cg-*Kit*$^{W-41J}$ *Tyr* $^+$ *Prkdc*^{scid}

Il2rg^{tm1Wjl}/ThomJ (NBSGW), PCA: principal component analysis, LogFC: Log fold change,

FDR: false recovery date, ChIP: chromatin immunoprecipitation.

Figure 4. MNX1 induces DNA damage. Immunofluorescence of FL-HSPC stained with

γH2AX –FITC antibody (green) and counterstained with DAPI (blue) (left panel, a).

Quantification of the number of γH2AX foci/cell. At least 50 cells were counted (right panel,

a). Representative histogram from the flow cytometry analysis of FL-HSPC and ABM-HSPC

after staining with DAPI for cell cycle distribution (left panel, b). Quantification of the

flowcytometry analysis represented as fold difference relative to control (right panel, b). Representative dot plots from the flow cytometry analysis of FL-HSPC and ABM-HSPC after double staining with Annexin/V and DAPI for apoptotic analysis (upper panel, c). Quantification of the flowcytometry analysis represented as fold difference relative to control (lower panel, c). Representative histogram from the flow cytometry analysis of BM from leukemia NBSGW mice transduced with MNX1-FL cells (yellow) in comparison with NBSGW mice transduced with empty vector Ctrl (red) after staining with DAPI for DNA content, numbers indicated median DNA values at G1 phase and G2 phase of cell cycle. (d). Quantification of DNA content in supplement figure 7c. Data represent mean ± SD of at least three experiments and is considered significant using two-sided student t-test (**) at $p \leq 0.01$ and (*) at $p \leq 0.05$. FL: fetal liver, ABM: adult bone marrow, Ctrl: control empty vector, Fusion: *MNX1::ETV6* fusion, e.A: early apoptosis, l.A: late apoptosis.

Figure 5. MNX1 alters histone methylation. mRNA expression of *Cep164* determined by quantitative real time qPCR, normalized to *Hprt* and represented as fold difference relative to control from both in vitro FL-HSPC (MNX1-FL) and BM from leukemia mice (MNX1-BM) (a). Quantification of the binding of MNX1 to the promoter regions of *Cep164* in FL-HSPC with either MNX1 or empty ctrl determined by ChIP-qPCR (b). Western blot analysis of H3K4me1, H3K4me2, H3K4me3 and H3K27me3 in the transduced FL-HSPC. Histone 3 and actin used as loading controls (c). Heatmap of differentially bound H3K27me3 from BM of leukemic mice (MNX1) versus control FL-HSPC transduced with empty vector (Ctrl) as determined by ACT-Seq (left panel, d). Annotation of different enriched regions for H3K27me3 ACT-Seq (right panel, d). Results were considered at the log fold-change cutoff (logFC) of $\geq$ |1| and false discovery rate (FDR) of ≤ 0.05. Heatmap of differentially accessible regions from BM of leukemic mice (MNX1) versus control FL-HSPC transduced with empty vector (Ctrl)

as determined by ATAC-Seq (left panel, e). Annotation of different enriched regions for ATAC-Seq (right panel, e). Results were considered at the log fold-change cutoff (logFC) of ≥ |1| and false discovery rate (FDR) of ≤0.05. Pathway enrichment analysis for the identified proteins after Co-IP and mass spectrometry using STRING with reactome data set, showing strength of the pathway as Log10 observed proteins/background in the respective pathway (f). The binding of AHCY and MAT2A was detected by immunoprecipitation of MNX1 using HA-antibody followed by western blot using AHCY and MAT2A antibodies. ETV6 with HA tag was used as a negative control. Total protein input used as loading control (g). Methionine concentration was determined by LC-ESI-MSMS from FL-HSPC expressing MNX1 (MNX1-FL) and leukemia BM cells (MNX1-BM), normalized against total protein concentration and represented as fold difference relative to control (h). Recombinant Histone H3.1and H3.3 were subjected to an invitro methyltransferase reaction using MNX1 complex pulled down with HA antibody from FL-HSPC transduced with MNX1(M), in comparison with FL-HSPC with empty vector (C), in the presence of SAM and DTT. The reactions were terminated by boiling in SDS sample buffer. Separation of samples in 12% SDS-PAGE was followed by immunoblotting (IB) with mono, di and tri methyl-lysines antibody. Reblotting was made for detection of HA and total Histone H3. As indicated, negative controls were obtained by omitting Histone H3, or the pulled down immune complex (left panel, i). Quantification of the invitro methyltransferase reaction (right panel, i). Data represent mean ± SD of at least three experiments and is considered significant using two-sided student t-test (**) at $p \leq 0.01$ and (*) at $p \leq 0.05$. FL: fetal liver, BM: Bone marrow, Ctrl: control empty vector, Fusion: *MNX1::ETV6* fusion, WB: Western blot, ChIP: chromatin immunoprecipitation, SAM: S-Adenosyl methionine , DTT: Dithiothreitol.

Figure 6. Sinefungin rescued MNX1-induced phenotype.   Protein expression levels of

H3K4me3, H3k27me3, H3K4me1 and H3K4me2 in FL-HSPC with either MNX1

overexpression or empty vector (Ctrl) after treatment with vehicle or 5μM Sinefungin (Ctrl+S,

MNX1+S) as determined by Western blotting. Actin served as loading control (a).

Quantification of the Western blot analysis in Supplement figure 9a. Quantification of the

number of γH2AX foci/cell. At least 50 cells were counted. Data represented as fold difference

relative to control (b). Quantification of flow cytometry analysis of the cells with the indicated

antibodies. Data represented as fold difference relative to control (c and d). Data represent mean

$\pm$ SD of at least three experiments and is considered significant using two-sided student t-test

(**) at $p \leq 0.01$ and (*) at $p \leq 0.05$. Kaplan-Meier survival curves of (NBSGW) mice

transplanted with FL cells after retrovirus transduction with either ectopic expression of MNX1

or empty vector (Ctrl) after treatment with vehicle or 5μM Sinefungin (MNX1+S, Ctrl+S).

Results of MNX1 (n = 7) and Sinefungin treated MNX1 cells (MNX1+S) (n = 6) were analyzed

using the log-rank test (***: significant at $P \leq 0.01$) (e). Heatmap of sample distance and

similarity matrix for the differentially expressed genes with either ectopic expression of MNX1

or empty vector (Ctrl) with or without 5μM Sinefungin (Ctrl+S, MNX1+S), showing clustering

of cells and similarity between the samples. Results were considered at the log fold-change

cutoff (LogFC) $\geq |1.5|$ and FDR ≤ 0.05 (f). Schematic diagram for the proposed model of action

of MNX1 (g).

Table 1. In-silico predicted transcription factor binding sites by DiffTF package from ATAC-

Seq showing top differentially activated (FDR < 0.05) transcription factors (TFs) binding

between MNX1 BM from leukemia mice and FL-HSPC Ctrl cells

Up-regulated TFs	weighted_mean Difference	pvalueAdj	Down regulated TFs	weighted_mean Difference	pvalueAdj
Tal1.a	1.294844532	9.30E-79	Fosl2	-0.421333369	1.84E-24
Gata1	0.567577232	1.24E-20	Jun	-0.415408916	2.58E-23
Gata3	0.550998432	1.12E-19	Cebpe	-0.39280302	0.000108
Gata2	0.540274293	1.01E-16	Batf	-0.387629056	1.57E-17
Alx1	0.371423122	0.023026	Fosl1	-0.384618862	4.57E-21
Nf2l2	0.288226747	2.11E-17	Cebpa	-0.353311343	3.14E-06
Zeb1	0.232871599	9.60E-41	Gfi1	-0.349686909	8.73E-08
Tbx5	0.231419764	1.76E-09	Fosb	-0.339609157	5.76E-15
Mafk.s	0.22701829	3.01E-10	Sta5a	-0.339203846	2.94E-09
Nfe2	0.218904994	4.77E-11	Jund	-0.337147615	8.92E-16
Gata6	0.213067314	2.28E-11	Cebpd	-0.323368792	0.000182
Foxo3	0.202788032	0.000274	Gfi1b	-0.308823273	8.94E-12

 Table 2. Identified proteins in top enriched pathways after Co-IP of proteins in complex with

616 MNX1 and mass spectrometry analysis. As determined by STRING using reactome data set

Pathway	Matching proteins
Association with TriC/CCT	Cct8,Cct6a,Tcp1,Cct4
Methylation	Mat2b,Ahcy,Mat2a,Trmt112
RHO GTPases activate PAKs	Ppp1cb,Myh9,Myh11,Myh10,Myh14
Citric acid cycle (TCA cycle)	Cs,Mdh2,Idh3g,Dld
Pyruvate metabolism	Pdk3,Slc16a3,Vdac1,Ldha,Dld
Actin dynamics for phagocytic	Actr2,Arpc2,Myh9,Cyfip2,Arpc3,Myo5a,Actr3
Translation initiation complex	Pabpc1,Eif3e,Rps21,Eif2s3x,Rps12,Rps28
RHO GTPase Effectors	Actr2,Mapk14,Arpc2,Ppp1cb,Myh9,Pfn1,Aurkb,S100a8,Tuba1b,Bub3,Myh11,Cyfip2,Arpc3,Myh10,Ywhaq,Myh14,Kif2a,S100a9,Actr3
Cellular Senescence	Mapk14,Asf1a,Lmnb1,Txn1,H2afz,Hist1h1b,Hist2h2be,Eed
Mitotic Anaphase	Psmd11,Aurkb,Psmd6,Lmnb1,Smc3,Psmd13,Chmp4b,Smc1a,Tuba1b,Rcc1,Bub3,Kif2a
M Phase	Psmd11,Aurkb,Psmd6,Lmnb1,Smc3,Psmd13,Chmp4b,H2afz,Smc1a,Smc4,Tuba1b,Rcc1,Bub3,Hist2h2be,Csnk2a1,Smc2,Kif2a
Metabolism of proteins	Pabpc1,Srp14,Psmd11,Rab10,Ddx5,Aurkb,Psmd6,Vdac2,Ero1l,Eif3e,Mrpl21,Smc3,Psmd13,Cct8,Srp9,Txn1,Trappc3,Cct6a,Cmas,Rab27a,Rpl22l1,Smc1a,Rpl37,Mat2b,Pdia6,Fn1,Rps21,Eif2s3x,Hist2h2ab,Rps12,Tuba1b,Ctbp1,Rpl15,Rpl36,Rpl34,Lgals1,Hist2h2be,Eef1g,Csnk2a1,Vdac1,Wdr5,Rps28,Trmt112,Tcp1,Copa,Gnb4,Ctsd,Cct4,Nop58,Tceb1
Metabolism of RNA	Pabpc1,U2af2,Psmd11,Prpf8,Snrpd3,Ddx5,Psmd6,Wdr46,Psmd13,Rpl22l1,Rpl37,Utp15,Wdr43,Rps21,Rcl1,Rps12,Rpl15,Rpl36,Rpl34,Srsf2,Nhp2l1,Snrpb,Rps28,Srsf3,Ddx6,Hnrnpd,Apobec3,Nop58

Figure 1

Figure 2

a

Heatmap of RNA-seq from BM of NSG mice.
LogFC 1 and FDR ≤ 0.05

MNX1   Ctrl

b

PCA of RNA-seq from BM of NSG mice

condition
● Ctrl
● MNX

c

d

Gene enrichment analysis of RNA-Seq using Gene ontology gene set

PYRIDINE_CONTAINING_COMPOUND_BIOSYNTHETIC_PROCESS
PORPHYRIN_CONTAINING_COMPOUND_METABOLIC_PROCESS
MITOTIC_G2_M_TRANSITION_CHECKPOINT
NUCLEAR_BODY_ORGANIZATION
NUCLEOSOME_POSITIONING
NEGATIVE_REGULATION_OF_DNA_REPAIR
DNA_DEALKYLATION
MITOTIC_DNA_INTEGRITY_CHECKPOINT_SIGNALING
MITOTIC_G2_DNA_DAMAGE_CHECKPOINT_SIGNALING
HISTONE_EXCHANGE
SNRNA_TRANSCRIPTION
NUCLEOSOME_ORGANIZATION
REGULATION_OF_DOUBLE_STRAND_BREAK_REPAIR_VIA_NHEJ
NUCLEOSOME_ASSEMBLY
CENTROMERE_COMPLEX_ASSEMBLY
REGULATION_OF_MEGAKARYOCYTE_DIFFERENTIATION
CHROMOSOME_CONDENSATION
MITOTIC_INTRA_S_DNA_DAMAGE_CHECKPOINT_SIGNALING
CHROMATIN_REMODELING
HISTONE_MRNA_CATABOLIC_PROCESS
HISTONE_H3_K4_METHYLATION
IMMUNE_RESPONSE_TO_TUMOR_CELL
SMOOTH_MUSCLE_CELL_APOPTOTIC_PROCESS
REGULATION_OF_MYELINATION
REGULATION_OF_PHOSPHOLIPID_METABOLIC_PROCESS
NEGATIVE_REGULATION_OF_NIK_NF_KAPPAB_SIGNALING
REGULATION_OF_MYELOID_CELL_APOPTOTIC_PROCESS
REGULATION_OF_VIRAL_TRANSCRIPTION
POLYAMINE_BIOSYNTHETIC_PROCESS
SKELETAL_MUSCLE_TISSUE_REGENERATION
NEGATIVE_REGULATION_OF_TYPE_I_INTERFERON_PRODUCTION
NEGATIVE_REGULATION_OF_NERVOUS_SYSTEM_PROCESS

NES (normalized enrichment score), NOM p-val ≤ 0.05

Figure 3

Figure 4

Figure 5

Figure 6