Asymmetric retinal direction tuning predicts optokinetic eye movements across stimulus conditions

Scott C. Harris[1,2]* and Felice A. Dunn[1]*

1. Department of Ophthalmology, University of California, San Francisco, San Francisco, CA 94143, USA
2. Neuroscience Graduate Program, University of California San Francisco, San Francisco, CA 94158, USA
*Corresponding Authors, Scott.Harris@ucsf.edu, Felice.Dunn@ucsf.edu

ABSTRACT

Across species, the optokinetic reflex (OKR) stabilizes vision during self-motion. OKR occurs when ON direction-selective retinal ganglion cells (oDSGCs) detect slow, global image motion on the retina. How oDSGC activity is integrated centrally to generate behavior remains unknown. Here, we discover mechanisms that contribute to motion-encoding in vertically-tuned oDSGCs, and leverage these findings to empirically define signal transformation between retinal output and vertical OKR behavior. Specifically, we demonstrate that motion encoding in vertically-tuned oDSGCs is contrast-sensitive and asymmetric for oDSGC types that prefer opposite directions. These phenomena arise from the interplay between spike threshold nonlinearities and differences in synaptic input weights, including shifts in the balance of excitation and inhibition. In behaving mice, these neurophysiological observations, along with a central subtraction of oDSGC outputs, accurately predict the direction and magnitude of vertical OKR across stimulus conditions. Thus, asymmetric synaptic input across competing sensory input channels can critically shape behavior.

INTRODUCTION

From humans to insects, a wide range of organisms depend on visual cues to navigate the world. When these animals respond to incoming visual information by enacting motor plans, however, relative motion is created between the eye and the visual scene. Such motion, termed "retinal slip," has the potential to corrupt subsequent vision and threaten survival.

To compensate for this possibility, the optokinetic reflex (OKR) is a highly conserved visual behavior that performs retinal image stabilization across an impressive range of species (including invertebrates[1], reptiles, amphibians, fish, birds, and all mammals, including mice and humans, reviewed by ref. [2]). OKR consists of visually-evoked, compensatory eye movements (or head movements in some species) that offset the slow, global image motion generated by self-movement[3]. In most species, OKR is also well known to change across stimulus conditions. Adjustments to stimulus contrast, color, location, velocity, or spatial frequency, for example, can elicit distinct OKR patterns[4–11]. Nonetheless, neurophysiological mechanisms for such phenomena are largely missing.

Anatomically, the pathways that underlie OKR in mammals are well defined (reviewed by refs. [2,3,12,13]). Starting in the retina, a dedicated class of biological motion detectors known as ON direction-selective retinal ganglion cells (oDSGCs) encode the slow global image motion that occurs during retinal slip. Classic work identified three types of oDSGCs that each maximize their firing rate in response to a particular cardinal direction of stimulus motion (i.e., up/superior, down/inferior, and nasal/anterior)[14,15]. A recent study also identified a fourth oDSGC type in mouse that encodes temporal/posterior motion[16]. As signals from oDSGCs depart the retina, they avoid typical retinal projection nuclei such as the superior colliculus and lateral geniculate nucleus. Instead, axons from vertically- and horizontally-preferring oDSGCs course along dedicated retinofugal tracts to a set of midbrain nuclei known collectively as the accessory optic system. Vertically-tuned oDSGCs terminate exclusively in, and comprise the sole retinal inputs to, the

medial and lateral terminal nuclei (MTN and LTN)[12,17–22] (but see Discussion and ref. [23]) - though

in mice, LTN is engulfed within MTN[17,21,24,25]. Here, their inputs are likely integrated into a single

velocity signal that reflects the vertical component of retinal slip. This information is then relayed

deeper into the brainstem where corresponding eye movements are enacted. Likewise,

horizontally-tuned oDSGCs exclusively target the nucleus of the optic tract/dorsal terminal

nucleus (NOT/DTN) where their signals are similarly integrated to ultimately generate the

horizontal component of OKR[17,24]. While correlations have been recognized between the

physiological properties of oDSGCs and OKR - particularly a matched speed tuning[26] - little is

known about how signals from multiple oDSGC types are integrated to generate OKR across

varying stimulus conditions[12,27].

Here, we reveal a general mechanism by which OKR may be generated across a range

of stimulus statistics. To do so, we focus primarily on a single parameter that is known to affect

OKR: motion direction. In many species - including both humans[28–31] (but see ref. [32]) and mice[21]

- OKR is more robust in response to superior than inferior motion. We aim to illuminate the

biological mechanism underlying this asymmetry in order to reveal more general processes by

which OKR may be differentially generated across stimulus conditions. Focusing on the direction-

sensitivity of vertical OKR is methodologically strategic in that (1) it limits the source of possible

neurophysiological mechanisms to functions of only the vertical OKR pathway, and (2) unlike

horizontal OKR, which relies on interhemispheric communication and mirror-image motion signals

from each eye[2,17,33], vertical OKR can be studied unilaterally. Our results indicate that the

behavioral asymmetry between superior and inferior OKR is linked to differences in the direction

tuning properties of oDSGCs that prefer superior and inferior motion. These physiological

differences arise from a shift in the balance of excitatory and inhibitory (E/I) synaptic inputs across

cell types, along with nonlinear transformations associated with spike thresholding. More

generally, we demonstrate that the tuning properties of vertically-tuned oDSGCs are uniquely

sensitive to such changes in synaptic input weights and show how this sensitivity can account for

changes to OKR across multiple stimulus conditions in behaving mice. As a whole, this work

reveals a neurophysiological mechanism underlying gaze stabilization and natural vision

throughout the seeing world.

RESULTS

Superior OKR is more robust than inferior OKR

Across species, superior motion generates a more robust OKR than inferior motion (e.g.

cat[33–35], chicken[36], monkey[37], human[28–31]). In mice, this phenomenon has been reported

anecdotally in juvenile animals[21]. To investigate if an asymmetry between superior and inferior

OKR exists in adult mice, we designed a behavioral rig to accurately evoke and quantify vertical

OKR in head-fixed animals (**Fig 1a-b**, Methods)[38]. Eye movements were measured in response

to vertically-drifting, full-field gratings that have been used previously to evoke OKR[19,21,24,39].

Across all adult mice, superior- and inferior-drifting gratings generated distinct, but reproducible

eye movements: while superior gratings elicited repetitive slow, upward drifting eye movements

("slow nystagmus") interleaved with frequent resetting saccades in the opposite direction ("fast

nystagmus") (**Fig 1c**), inferior gratings tended to reliably drive only an initial slow nystagmus

immediately following stimulus onset, after which fast nystagmuses were infrequent and eye

position changed minimally (**Fig 1d**). To quantify these differences, we isolated periods of slow

and fast nystagmus post hoc by extracting saccadic eye movements from the raw eye position

trace. Superior stimuli elicited more frequent fast nystagmuses than did inferior stimuli (**Fig 1e**).

In addition, the total distance traveled during slow nystagmus was greater for superior stimuli (**Fig

1f**). These results demonstrate that vertical OKR is asymmetric in adult mice.

Despite the stark asymmetry between superior and inferior OKR in response to

unidirectional drifting gratings, quantifying OKR gain (ratio of eye velocity to stimulus velocity)

under these conditions presented challenges due to variability in the OKR waveform across

stimulus directions (**Fig 1c-d**). To better quantify gain, we designed a second stimulus in which a

grating oscillated sinusoidally between superior and inferior motion while retaining a constant

average position. In every animal subject, this oscillating grating evoked sequential superior and

inferior slow nystagmuses that were phase-locked to the stimulus (**Fig 1g-h, j**). Removing fast

nystagmuses from the eye trace (**Fig 1h**) revealed that gain tended to be higher during the

superior phase compared to the inferior phase of individual oscillations (**Fig 1i**). This bias was

reflected by an average offset of vertical eye position in the superior direction over the course of

a single stimulus oscillation (**Fig 1j**). Taken together, these results demonstrate that superior

motion drives a more robust OKR than inferior motion.

Superior and Inferior oDSGCs have distinct direction tuning properties

While behavioral asymmetries between superior and inferior OKR could arise anywhere

along the vertical OKR pathway, a plausible neurophysiological substrate is at the level of the

retina where the pathways that encode superior and inferior motion have yet to converge. Further,

because the ganglion cells that encode superior motion ("Superior oDSGCs") and inferior motion

("Inferior oDSGCs") together serve as an input bottleneck for the remainder of the pathway, any

physiological asymmetry between these cell types will propagate to behavior unless specifically

corrected for by subsequent circuitry (see Discussion). Thus, we hypothesized that the behavioral

asymmetry between superior and inferior OKR may result from physiological differences between

Superior and Inferior oDSGCs.

To probe oDSGCs involved in vertical OKR, we made central injections of a fluorescent

retrograde tracer into their central target, MTN (**Fig 2a, S1**). This approach labeled an average of

667±16 retinal ganglion cells (RGCs; N=17 retinas), distributed evenly across each quadrant of the contralateral retina (**Fig 2b, S1**). We investigated the direction tuning properties of MTN-projecting RGCs by making cell-attached recordings from labeled cells while presenting a drifting bar stimulus that moved slowly (10°/s) across the retina in 8 directions (**Fig 2c**). Importantly, this stimulus was comparable to the grating stimuli used to evoke vertical OKR in behaving animals, i.e. equivalent cycle width, wavelength, and speed to the unidirectional gratings. The majority (94.76%) of retrogradely labeled cells were direction-selective and preferred either dorsal-to-ventral (n=116 of 286) [i.e., Superior oDSGCs, because these cells detect superior motion in visual space after accounting for inversion of the image by the eye's optics (**Fig 2d**)] or ventral-to-dorsal (i.e., Inferior oDSGCs, n=155 of 286) motion on the retina (**Fig 2e**). In agreement, mosaic analyses indicated that retrogradely labeled cells likely consisted of two RGC types (**Fig S1**). Both Superior and Inferior oDSGCs invariably had baseline firing rates of approximately 0hz (spikes/s: Sup. 0.0134±0.006; Inf. 0.0338±0.016; p=0.13 Sup. vs. Inf.). Nonetheless, Superior oDSGCs tended to produce more total spikes than Inferior oDSGCs in response to the drifting bar stimulus (**Fig 2f**), and the total areas of their tuning curves were greater (**Fig 2g**). Similarly, we noticed prominent topographic differences within cell types, with MTN-projecting RGCs in ventral retina spiking more than those in dorsal retina. The firing rates of Superior and Inferior oDSGCs covaried across the retina, however, and Superior oDSGCs tended to spike more than Inferior oDSGCs in every retinal quadrant (**Fig S2**).

We wondered if the difference in response magnitude between Superior and Inferior oDSGCs could be explained by a simple scaling difference of their tuning curves (i.e., same shape, different size), or if it was instead associated with an asymmetry in tuning curve shape. To answer this question, we first normalized the tuning curve of each cell in our data set to the magnitude of its response in the preferred direction, and then rotated it such that the preferred direction of each cell was aligned (**Fig 2h**). If the tuning curves of Superior oDSGCs were simply scaled versions of those of Inferior oDSGCs, then this normalization and alignment procedure

should eliminate any apparent differences between cell types. Instead, however, we found that

the normalized tuning curves of Superior oDSGCs had greater widths at 50% response magnitude

(**Fig 2i**) and total areas (**Fig 2j**), indicating that their tuning curves were broader than those of

Inferior oDSGCs. In agreement, circular Gaussian fits (see Methods) of Superior oDSGC tuning

curves were consistently wider than those of Inferior oDSGCs (**Fig 2k**). To understand how this

difference in tuning curve width might affect the ability of Superior and Inferior oDSGCs to encode

motion, we quantified the direction selectivity index (DSI) of each cell (magnitude of the vector

sum divided by the scalar sum, see Methods). In agreement, the DSIs of Superior oDSGCs were

lower than those of Inferior oDSGCs (**Fig 2l**). Finally, we tested if these asymmetries persisted

across stimulus conditions by using a drifting bar stimulus with 5-fold lower contrast. As before,

Superior oDSGCs spiked more, had broader tuning curves, and were less direction-selective

compared to Inferior oDSGCs in response to the low contrast bar (**Fig 2g, j, l insets**).

Together, these data indicate that asymmetries in vertical OKR are concomitant with

prominent physiological asymmetries between Superior and Inferior oDSGCs: Superior oDSGCs

not only produce more total spikes than Inferior oDSGCs, but also have broader tuning curves

and are less direction-selective. Further, tuning curve size (i.e., area of the unnormalized tuning

curve) and width (i.e., area of the normalized tuning curve) were positively correlated on a cell-

by-cell basis (**Fig 2m**), indicating that asymmetries in these metrics could arise from a common

mechanism.

Superior oDSGCs receive more excitatory input than Inferior oDSGCs

We sought to determine the source of tuning curve asymmetries between Superior and

Inferior oDSGCs. As in the other primary class of direction-selective retinal ganglion cell, the ON-

OFF DSGCs (ooDSGCs)[40–43], oDSGCs inherit the bulk of their direction selectivity via greater

inhibition from starburst amacrine cells (SACs) in response to null direction stimuli[44–46] (reviewed

by refs. [27,47,48]). Therefore, we postulated that the difference in spike tuning curve width between

Superior and Inferior oDSGCs may be the result of asymmetric inhibitory inputs between the two

cell types.

To investigate this possibility, we made whole-cell voltage-clamp recordings at the

reversal potential for excitation to isolate inhibitory inputs to Superior and Inferior oDSGCs in

response to the drifting bar stimulus (**Fig 3a-b**). Across cells, we found no significant difference

in the magnitude of inhibitory postsynaptic currents (IPSCs) between Superior and Inferior

oDSGCs (**Fig 3c-d**). IPSCs to Superior oDSGCs, however, were slightly more direction-selective

than those to Inferior oDSGCs (**Fig 3e**). Alone, sharper IPSC direction tuning in Superior oDSGCs

implies sharper spike tuning curves. Our cell-attached data directly contradict this prediction (**Fig**

2). Thus, asymmetries between Superior and Inferior oDSGC spike tuning curves cannot be

explained by the canonical model of retinal direction selectivity involving inhibition.

Since inhibitory inputs failed to account for the differences between Superior and Inferior

oDSGCs, we wondered whether excitatory inputs could better explain the asymmetries in their

spike tuning curves. To test this possibility, we made voltage-clamp recordings at the reversal

potential for inhibition to isolate excitatory postsynaptic currents (EPSCs) during the drifting bar

stimulus. For every direction of stimulus motion, Superior oDSGCs had larger EPSCs than Inferior

oDSGCs (**Fig 3f-i**). EPSCs to Superior cells were also less direction-selective than those to

Inferior oDSGCs (**Fig 3j**). In agreement, the ratio of the peak EPSC to the peak IPSC (E/I) was

greater for Superior oDSGCs than for Inferior oDSGCs (**Fig 3k, S3**), though not different in

direction-selectivity (**Fig 3l)**. In addition to examining the magnitude of inputs, we also found no

difference in the relative timing of peak EPSCs and IPSCs across cell types (not shown). Based

on these results, the difference in spike tuning curve size and shape between Superior and Inferior

oDSGCs may be related to a corresponding shift in the balance of E/I, due in particular to an

asymmetry in the amount of net excitation that each cell type receives.

To test whether this difference in the magnitude of excitatory input to Superior and Inferior

oDSGCs generalized across stimulus types, we measured the spike responses and membrane

currents of both cell types in response to a full-field light increment (**Fig S4**). As with the drifting

bar, the light increment elicited more total spikes in Superior than in Inferior oDSGCs (**Fig S4f-h**).

We also observed a positive correlation between the magnitude of a cell's increment response

and the area of both its tuning curve (Sup: R^2=0.75, p=1.57x10^{-21}; Inf: R^2=0.73, p=2.13x10^{-26}) and

normalized tuning curve (Sup: R^2=0.23, p=0.013; Inf: R^2=0.47, p=1.18x10^{-9}). Under voltage-clamp

conditions, Superior oDSGCs had greater peak EPSCs and excitatory charge transfers than

Inferior oDSGCs in response to the increment (**Fig S4m-o**). However, there was no difference in

the peak IPSCs or inhibitory charge transfers between cell types (**Fig S4i-k**). Further, we found a

strong correlation between the maximum firing rate of a cell to the increment and the magnitude

of its EPSC, but not IPSC (**Fig S4l, p**). These results demonstrate that Superior oDSGCs generate

more spikes than Inferior oDSGCs across multiple stimuli, and, further, that this increased spike

output is consistently associated with differences in the amount of excitatory, but not inhibitory,

input.

Postsynaptic differences may account for shifts in E/I

Differences in the postsynaptic currents of Superior and Inferior oDSGCs could result from

asymmetries in presynaptic wiring and/or the postsynaptic properties of each oDSGC type. Serial

block-face electron microscopy[40,49,50] and analysis of dendritic stratification[21] have not yet

provided evidence of presynaptic wiring differences between oDSGCs with different preferred

directions. Thus, we investigated possible postsynaptic asymmetries that might account for the

differences in how Superior and Inferior oDSGCs encode motion.

We first analyzed the morphological features of Superior and Inferior oDSGCs by filling

cells of both types with dye (**Fig 4a**). Convex hull analysis revealed that the dendritic fields of

Superior oDSGCs covered a larger total area than those of Inferior oDSGCs (**Fig 4b**). Sholl

analysis, however, showed similar dendritic complexity between cell types (**Fig 4c**). To explore

whether synaptic differences exist between cell types, we stained for the excitatory postsynaptic

density scaffolding protein PSD-95[51] (**Fig 4d-g**) and the inhibitory postsynaptic scaffolding protein

gephyrin[52,53] (**Fig 4h-k**). These assays revealed no difference in the number or density of synaptic

puncta between cell types. However, Superior oDSGCs had significantly larger excitatory, but not

inhibitory, puncta (**Fig 4g, k**). This finding is consistent with greater amounts of excitatory input to

Superior oDSGCs.

To complement these morphological differences, electrophysiological recordings also

revealed a number of intrinsic differences between Superior and Inferior oDSGCs. First, the

membrane capacitances of Superior oDSGCs were greater than those of Inferior oDSGCs (pF,

Sup: 76.84±7.41; Inf: 52.47±4.59; p=0.001). Sequentially recording both the spike output and

EPSCs of individual oDSGCs to the full-field increment stimulus also revealed that Superior

oDSGCs had lower spike-to-EPSC ratios (**Fig S4p**). This observation also tended to be true for

each direction of the drifting bar stimulus (**Fig S3d-aa**). In agreement, the input resistances of

Superior oDSGCs were lower than those of Inferior oDSGCs (GOhms, Sup: 0.181±0.021; Inf:

0.319±0.033; p=0.004). These phenomena are each consistent with the larger size of Superior

oDSGCs relative to Inferior oDSGCs. We found no significant differences in other intrinsic

properties including the resting membrane potential (mV, Sup: -54.6±1.1; Inf: -54.7±1.5; p=0.86)

and spike threshold (mV, Sup: -43.7±1.17; Inf: -45.0±1.8; p=0.70). Together, these data indicate

that Superior oDSGCs are less intrinsically excitable than Inferior oDSGCs, but that this

asymmetry is outweighed by counteracting discrepancies in the magnitude of excitatory input to

each cell type.

Untuned excitation broadens spike tuning curves

That Superior oDSGCs receive relatively more excitation than Inferior oDSGCs explains

their greater spike output (**Fig 2f-g; S4**). Less obvious, however, is whether this difference in

excitatory input can also account for the wider tuning curves of Superior oDSGCs (**Fig 2h-I**). A

debate remains over whether excitatory inputs to DSGCs are directionally tuned[48,50,54–58]

(reviewed by ref. [27]). While our results indicate that MTN-projecting RGCs might receive different

amounts of excitation based on stimulus direction (**Fig 3j**), the majority of directionally tuned

inputs were inhibitory (**Fig 3m**). Further, the apparent direction selectivity of EPSCs is likely

partially attributable to imprecise space clamp[48,56] (but see ref. [55]). Thus, the extent to which the

tuning curves of Superior and Inferior oDSGCs are shaped by directionally asymmetric excitation

is unclear, and we remain agnostic on this point. Instead, we focus on the more pronounced

observation that Superior oDSGCs receive greater excitation than Inferior oDSGCs across

stimulus directions (**Fig 3h-i, k, S3**), regardless of the extent to which this excitation is tuned. In

the following experiments, we investigate the relationship between spike tuning curve shape and

the overall amount of excitation to an oDSGC. We ask whether and how different magnitudes of

excitatory input to Superior and Inferior oDSGCs can explain their difference in tuning curve width.

To test how the magnitude of excitation, even when directionally untuned, to an oDSGC

changes the shape of its tuning curve, we measured the spikes of Superior and Inferior oDSGCs

in current-clamp (**Fig 5a-b, h**) while injecting constant amounts of either positive (to add ~6mV,

"depolarizing") or negative (to subtract ~6mV, "hyperpolarizing") current across stimulus

directions. Importantly, the depolarizing current injections were small enough such that every cell

in our dataset retained a baseline firing rate of 0hz. This approach allowed us to investigate how

providing a cell with more or less directionally-untuned excitation influences the shape of its spike

tuning curve. Indeed, we found that tuning curves measured under depolarizing conditions were

not only larger (**Fig 5i**), but also wider than those measured under hyperpolarizing conditions on

a cell-by-cell basis (**Fig 5a-d, j**). In agreement, both Superior and Inferior oDSGCs were less

direction-selective under depolarizing relative to hyperpolarizing conditions (**Fig 5k**). Together,

these data demonstrate that increasing the amount of excitation to an oDSGC broadens its tuning

curve.

Interestingly, comparing these results to those recorded extracellularly in cell-attached

recordings revealed an additional nuance. Depolarizing current injections widened only the tuning

curves of Inferior oDSGCs (**Fig 5c-d, top rows of 5e-f**). In contrast, hyperpolarizing injections

sharpened the tuning curves of both cell types (**Fig 5c-d, bottom rows of 5e-f**). One possible

explanation is that while excitation generally broadens tuning curve width, greater excitatory input

minimally affects Superior oDSGCs (**Fig 5c**) because they are already positioned closer to an

upper limit on this phenomenon. Nonetheless, our results highlight a causal relationship between

tuning curve width and the amount of untuned excitation that an oDSGC receives.

Spike threshold plays a dominant role in setting tuning curve width

Two complementary mechanisms could explain how untuned excitation broadens tuning

curve width. In the first mechanism, excitation influences the nonlinear transformation between

synaptic input and spike output that is introduced by a neuron's spike threshold (**Fig 5g**). More

specifically, thresholding may sharpen a neuron's spike tuning curve relative to the tuning of

underlying membrane fluctuations by clamping spike output at zero in response to subthreshold

membrane responses that are likely to occur for null direction stimuli[59]. When the amount of

untuned excitatory input increases, however, membrane fluctuations more readily surpass the

spike threshold for all stimulus directions, thereby broadening the spike tuning curve. In the

second mechanism, untuned excitation directly circularizes the tuning curve of underlying

membrane potentials by increasing null direction responses proportionally more than preferred

direction responses. This "additive" contribution would then be inherited by the spike tuning

curve[60].

We first tested whether thresholding effects contribute to the width of oDSGC tuning

curves. To do so, we isolated the underlying subthreshold voltages (Vm) from our current-clamp

recordings that also contained spikes (**Fig 5l**). Three independent observations about these Vm

tuning curves each suggests that thresholding critically influences spike tuning curve width. First,

unlike for spikes, depolarizing and hyperpolarizing current injections did not affect the width of Vm

tuning curves (**Fig 5n-o, t-u**). The total areas of Vm tuning curves were, however, slightly larger

under the hyperpolarizing condition, likely due to the marginally greater driving force on excitatory

conductances in this setting (**Fig 5m**). Second, the Vm tuning curves of Superior oDSGCs were

larger in magnitude than those of Inferior oDSGCs, but also narrower and more direction-selective

(**Fig 5m-o arrowheads**). This result was anticipated from our voltage-clamp recordings which

indicated that inhibition is more direction-selective in Superior oDSGCs (**Fig 3e**). It also suggests,

however, that the spike tuning curve width of oDSGCs is not directly inherited from that of the

underlying membrane potentials, and instead reflects the interplay between Vm magnitude and

spike threshold. Third, comparing the width of Vm tuning curves to those measured

simultaneously in spikes revealed that the spike tuning curves were substantially sharper than

those of the subthreshold voltages for both depolarizing and hyperpolarizing injections (**Fig 5p-**

s). The difference between the shapes of the spike and Vm tuning curves was smaller for the

depolarizing condition, however, because in this condition the majority of stimulus directions

elicited Vm responses that eclipsed the spike threshold (**Fig 5v-w**). Together, these three results

each corroborate the notion that thresholding plays a prominent role in setting spike tuning curve

width relative to the amount of untuned excitation that an oDSGC receives.

To test the extent to which excitation broadens oDSGC tuning curves through additive

effects, we recomputed Vm tuning curves after including a constant, additive offset that reflected

the average current injection supplied during depolarizing and hyperpolarizing injections. These

offset-corrected Vm tuning curves were significantly wider under depolarizing conditions than they

were under hyperpolarizing conditions (**Fig S5**). However, the observation that (uncorrected) Vm

tuning curves — in which putative additive differences *between* cell types are expected to persist

— were more sharply tuned for Superior than Inferior oDSGCs (**Fig 5n-o arrowheads**) indicates

that thresholding has a greater influence on spike tuning curves than does additive excitation. In

agreement, current injections caused significantly larger changes in spike tuning curves than in

offset-corrected Vm tuning curves (**Fig 5t-u**). These results demonstrate that while untuned

excitation likely has some additive effect, complementary thresholding has greater influence over

oDSGC spike tuning curves.

A parallel conductance model recapitulates the influence of thresholding on oDSGC

direction tuning

Together, our current injection experiments suggest that (1) spike thresholding plays a

prominent role in setting the width of oDSGC tuning curves, and (2) additive effects are likely

present but do not dominate. To independently test these findings, we built a parallel conductance

model of an oDSGC based on empirically measured parameters from a separate set of cells to

those used for the current injection experiments. Among these parameters were 8 separate

inhibitory conductances for each stimulus direction, and a single, directionally-untuned excitatory

conductance (**Fig 6a-b**). We then tested how manipulating the gain of the excitatory conductance

affected tuning curves generated from either spikes or the peak subthreshold membrane

potentials (**Fig 6c-f**) (see Methods).

As in our empirical data, we found that increasing the gain of untuned excitation to the

model oDSGC increased the total area of both spike and Vm tuning curves (**Fig 6d**). However,

while the spike tuning curve rapidly widened and became less direction-selective with increasing

excitatory gain, the shape of the Vm tuning curve was much less dependent on excitatory gain

(**Fig 6e-f**). The stark difference between the spike and Vm trajectories can only be attributed to

thresholding effects. On the other hand, the shallow slope of the Vm curve reflects the additive

contribution of excitation. Consistent with our physiological results, thresholding played a critical

role in setting the model oDSGC's tuning curve width, whereas additive effects were relatively

minor. Further, the prior observation that depolarizing current injections influenced Superior

oDSGCs less than Inferior oDSGCs (**Fig 5c-f**) is supported by the diminishing marginal effect of

additional excitatory gain on spike tuning curve width. We also noticed that the average

empirically measured normalized tuning curve area and DSI from our cell-attached recordings

(**Fig 6e-f circles**) fell within the regime where these metrics were steeply dependent on excitatory

gain. In this regime, thresholding effects render small changes in synaptic inputs particularly

influential for oDSGC direction tuning.

Thresholding effects produce contrast-sensitive direction tuning in oDSGCs

The dependence of oDSGC tuning on thresholding predicts that any stimulus which

influences the magnitude of synaptic inputs may also alter tuning curve shape. We tested this

hypothesis by comparing the tuning curves of Superior and Inferior oDSGCs in response to high

contrast (stimuli used in all previous figures) and low contrast (5-fold dimmer) drifting bars. For

most cells, low contrast bars elicited fewer total spikes (**Fig 7a-b**). However, as for high contrast

stimuli, Superior oDSGCs tended to spike more and have broader tuning curves than Inferior

oDSGCs in response to low contrast bars (**Fig 2g, j, l, insets**). Interestingly, cells of both types

had sharper spike tuning curves in response to low, compared to high, contrast stimuli (**Fig 7c-**

d). To test whether this contrast-sensitivity could be attributed to thresholding effects, we

measured the tuning curves of subthreshold membrane potentials. While the area of Vm tuning

curves was greater under high contrast conditions (**Fig 7e**), Vm tuning curve width and direction

selectivity did not change with contrast (**Fig 7f-g**). In agreement, the fraction of spike tuning curves

that sharpened under low contrast was significantly different than the equivalent fraction of Vm

tuning curves that followed the same trend (normalized area: p=0.0019; DSI: p=0.0046; 2-sided

Fisher's exact) (**Fig 7c-d, f-g**), and Vm tuning curves were less affected by stimulus contrast than

the tuning curves of simultaneously measured spikes (**Fig 7h-i**). These results indicate that

thresholding is critical to setting oDSGC spike tuning curve width across stimulus contrasts, just

as it is across cell types.

Our finding that oDSGCs are contrast-sensitive departs from previous work which has

shown that direction tuning in ooDSGCs is contrast-invariant[61–63]. Contrast-invariance in

ooDSGCs appears to rely on the stability of E/I across contrasts. For this reason, we tested

whether the contrast-sensitivity of oDSGCs was associated with a mutable E/I by sequentially

measuring excitatory and inhibitory synaptic currents for preferred and null direction stimuli across

contrasts. Our data show that E/I in neither the preferred nor null direction changes systematically

with contrast (**Fig 7j-k**). Further, the fraction of spike tuning curves that sharpened under low

contrast conditions was significantly greater than the fraction of E/I ratios that were lower under

low contrast conditions (normalized area: PD p=0.0132, ND p=0.0132; DSI: PD p=0.0044, ND

p=0.0044; 2-sided Fisher's exact) (**Fig 7c-d, j-k**). These results suggest that as in ooDSGCs, E/I

in oDSGCs is relatively stable across contrasts. However, the spike tuning curves of oDSGCs are

nonetheless contrast-sensitive. Thus, stable E/I alone is insufficient to maintain the contrast-

invariance of spikes. In the case of oDSGCs, changes to the absolute magnitude of synaptic

inputs across contrasts, along with thresholding effects, appear to trump the contrast-invariance

of E/I.

To further test the extent to which the magnitude of synaptic inputs can affect oDSGC

tuning curves even with stable E/I, we revisited our parallel conductance model of an exemplary

oDSGC. We modeled oDSGC responses while changing the gain of excitatory and inhibitory

conductances together, thereby keeping E/I constant while nevertheless adjusting the total

magnitude of E and I. As in our empirical data, we found that lowering the gain of E and I together

to simulate responses to low contrast stimuli resulted in sharper spike tuning curves. The width

of Vm tuning curves, however, remained relatively contrast-invariant (**Fig 7I**). These results

recapitulate our empirical findings and indicate that the spike threshold nonlinearity influences

spike tuning curve width across contrasts. Other stimulus parameters that affect the magnitude

of excitation, inhibition, or both are also likely to modulate the direction coding properties of

oDSGCs.

A subtraction algorithm predicts vertical OKR from oDSGC activity

Having established the tuning properties of Superior and Inferior oDSGCs, we asked

whether the asymmetries between these cell types, along with their contrast-sensitivities, could

explain how vertical OKR changes across stimulus conditions. Cross-species work has

established that Superior and Inferior oDSGCs are likely centrally integrated by a subtraction

algorithm[3,22,64]. In this model, OKR is predicted on the basis of the *difference* in spike rate between

Superior and Inferior oDSGCs, rather than by the absolute spike rate of either cell type. A number

of observations support this model: (1) stimuli that activate both Superior and Inferior oDSGCs

[e.g. a full-field increment of light (**Fig S4**)] do not elicit OKR, (2) stimuli that differentially activate

Superior and Inferior oDSGCs (e.g. drifting gratings) maximally drive OKR (**Fig 1, 2**), (3) Superior

and Inferior oDSGCs project to separate MTN subnuclei[21,22] that are mutually connected by

inhibitory interneurons (i.e., a simple differentiating circuit)[22,65], (4) MTN neurons prefer either

superior or inferior motion[21,22,64,66–68], but their preferred and null directions are not 180° apart;

instead, they correspond to the preferred directions of opposing oDSGC types[64,67,68], which may

differ by less than 180° [15] (but see ref. [16]), and (5) MTN neurons maintain moderate baseline spike

rates that are not only augmented by preferred direction stimuli, but also diminished by null

direction stimuli[64,66–68]. Together, along with the simplicity of the vertical OKR pathway and its

isolation from other visual circuits, these lines of evidence all point to a circuit motif in which MTN

neurons encode the difference in spike rate between Superior and Inferior oDSGCs. Thus,

superior OKR likely occurs when Superior oDSGCs spike significantly more than Inferior

oDSGCs, and vice versa. The robustness of OKR is putatively related to the magnitude of the

spike rate difference (**Fig 8a**).

We used our empirically recorded electrophysiology data from Superior and Inferior

oDSGCs to generate hypotheses for how OKR gain would change across stimulus conditions

(**Fig 8b**). To do so, distributions of Superior and Inferior oDSGC firing rates were compared for

superior and inferior drifting bars at high and low contrast (i.e., four stimulus conditions) (**Fig 8c-**

f). We inferred that the relative gain under each stimulus condition would be related to the

corresponding difference in spike rate between Superior and Inferior oDSGCs under that same

condition. For instance, gain in response to a high contrast superior grating was predicted by the

difference between the preferred direction responses of Superior oDSGCs and the null direction

responses of Inferior oDSGCs to high contrast bars. Behavioral predictions about how gain would

change across stimulus conditions were made by comparing bootstrapped distributions of such

differences (**Fig 8g-h, k-l**). Importantly, these inferences constitute linear predictions of gain.

Allowing for the possibility that downstream circuitry incorporates additional monotonic

nonlinearities, a linear prediction is consistent with behavior so long as it predicts gain changes

in the correct *direction* (but not magnitude) across stimulus conditions.

The asymmetries in the tuning curves of Superior and Inferior oDSGCs resulted in four

first-order predictions for how OKR gain changes across stimulus conditions: (1) for superior

stimuli, gain is greater in response to high contrast stimuli than to low contrast stimuli (**Fig 8g**),

(2) for inferior stimuli, gain is greater in response to high contrast stimuli than to low contrast

stimuli (**Fig 8h**), (3) at high contrast, gain is greater in response to superior stimuli than to inferior

stimuli (**Fig 8k**), and (4) at low contrast, gain is greater in response to superior stimuli than to

inferior stimuli (**Fig 8l**). We also generated two additional second-order hypotheses that rely on

an approximately linear relationship between differences in oDSGC output and gain in the tested

regime: (5) gain is higher under high contrast, regardless of stimulus direction (**Fig 8o, s**), and (6)

behavioral asymmetries in response to superior and inferior stimuli diminish with decreasing

contrast (**Fig 8p, s**). Thus, our physiological data generate clear predictions for how the tuning

properties of oDSGCs affect OKR across stimulus conditions.

We next tested these predictions in behaving mice. OKR in the superior and inferior

directions was measured in response to high and low contrast oscillating gratings. Our results

revealed that all six of our linear predictions were consistent with behavior (**Fig 8i-j, m-n, q-r, u,**

S6). Most notably, gain decreased with contrast (**Fig 8q, u**), and the asymmetry between superior

and inferior OKR that we originally noticed under high contrast conditions diminished in response

to low contrast stimuli (**Fig 8r, u**). This latter observation is consistent with our linear prediction

that the net change in eye position over the course of a single oscillation would be greater for high

contrast stimuli (**Fig 8p, s**). While this result may be related to the overall decrease in gain under

low contrast, it is certainly not *necessitated* by that phenomenon and relies on further subtleties

in the relationship between Superior and Inferior oDSGC responses at both contrasts. Indeed,

permuting the behavioral predictions by scrambling which cellular responses were assigned to

superior/inferior motion and high/low contrast revealed that only 5 permutations (out of 256

possibilities, 1.95%) accurately matched our behavioral results. Consistent with these findings,

the relationship between our cellular linear predictions and behavioral results were fit well by a

simple monotonic function that was approximately linear within the measured regime (**Fig 8t**).

Taken together, these results provide a neurophysiological explanation for how vertical OKR

changes across multiple stimulus conditions and reveal that the circuit and cellular properties that

shape oDSGC tuning curves have direct and predictable consequences for behavior.

DISCUSSION

Our results depict a neurophysiological mechanism by which vertical OKR changes across stimulus conditions. We demonstrate that superior and inferior OKR are asymmetric in adult mice (**Fig 1**), and show how this behavioral phenomenon can be traced to novel asymmetries in the direction tuning properties of Superior and Inferior oDSGCs (**Fig 2**). Mechanistically, a shift in the balance of excitatory and inhibitory inputs across cell types influences direction tuning, primarily through an effect associated with spike thresholding (**Fig 3-6**). Similar thresholding effects also confer contrast-sensitivity of the spike tuning curve, even when E/I is contrast-invariant (**Fig 7**). Together, these cellular properties accurately predict how vertical OKR changes in response to stimulus direction and contrast (**Fig 8**).

Directional asymmetries in OKR are common across species. Besides the vertical asymmetries that we investigated here, horizontal OKR is asymmetric in many organisms[2,69]. This asymmetry manifests nearly universally as higher OKR gain in response to temporal-to-nasal (anterior) motion than to nasal-to-temporal (posterior) motion, and often only when stimuli are viewed monocularly. It is tempting to speculate that these horizontal asymmetries may similarly be linked to direction tuning in the retina. While anterior-preferring oDSGCs are critical to horizontal OKR[12,24,39], posterior-preferring oDSGCs were only recently identified in rodents and display definitively weaker direction tuning properties than their anterior-preferring counterparts[16]. Further, a subtraction mechanism between horizontally tuned oDSGCs may also underlie horizontal OKR[33]. However, at least two confounds obscure a connection between the tuning properties of Anterior and Posterior oDSGCs and asymmetries in horizontal OKR. First, oDSGCs in the left and right eyes, and their contralateral central targets (NOT/DTN), encode different absolute directions of stimulus motion (reflection occurs over the sagittal body axis). To compensate, signals are compared across eyes/hemispheres prior to behavior[33], and, in some species, NOT/DTN receives descending, often binocular, inputs from visual cortex[2,13,70,71].

Second, recent studies have suggested that ooDSGCs could be involved in horizontal OKR[17,23].

Asymmetries in the horizontal version of the behavior may thus depend on more complex

considerations.

Analogous confounds are less concerning when considering the mechanisms that could

account for asymmetries in vertical OKR. For one, stimuli that induce vertical OKR, such as the

gratings used here, are perceived identically by both eyes. Interhemispheric communication is

unlikely to influence behavior under such conditions. Though signals are probably also exchanged

between the horizontal and vertical OKR pathways[3,13,18], these channels likely play a minimal role

in shaping OKR to purely vertical stimuli[33]. Finally, while it has been suggested that a vertically-

tuned ooDSGC may also project to MTN[23], this cell type was not clearly revealed by either

anatomical or electrophysiological analyses during our retrograde labeling experiments (**Fig S1,**

S4). Instead, we find that two populations of ganglion cells project to MTN, and that these cells

can be classified as Superior and Inferior oDSGCs (**Fig S1**). These results are consistent with

characterizations of MTN-projecting RGCs across many species[20,21,72–74]. Thus, unlike for

horizontal OKR, asymmetries in vertical OKR can be explained more simply by the physiology of

oDSGCs.

Despite the possibility that asymmetries in vertical and horizontal OKR are influenced by

separate mechanisms, the two phenomena could share a common ethological function. Optic

flow associated with forward locomotion typically includes a large posterior component. Thus, it

has been suggested that posterior OKR is less reliable than anterior OKR in order to mitigate

lateral eye movements that might otherwise occur when an animal walks forward[75]. Similar

reasoning may explain the asymmetry between superior and inferior OKR[28,35,37]. Indeed, recent

work in freely moving mice demonstrated that optic flow has a stronger inferior than superior

component[76]. Further, optic flow was largest in dorsal retina, which could explain our additional

observation that stronger stimuli were required to activate Superior and Inferior oDSGCs alike in

dorsal relative to ventral retina (**Fig S2**). Thus, the cellular and behavioral asymmetries identified

in this study may serve to mitigate the influence of natural locomotion on OKR. Future work should

address this possibility by mapping OKR gain as a function of the extent to which various stimuli

reflect locomotion-associated optic flow.

From a physiological perspective, our results provide the first evidence that Superior and

Inferior oDSGCs encode motion asymmetrically. This finding coincides with previous work

demonstrating genetic and anatomical differences between oDSGC types. Multiple transgenic

lines are known to label either Superior or Inferior oDSGCs, but not both[18,20,21,74]. The axons of

Superior and Inferior oDSGCs take separate retinofugal tracts and project to different MTN

subnuclei in mice[21]. Additional differences exist between vertically- and horizontally-tuned

oDSGCs[12,24]. The competitive advantage of directionally-asymmetric OKR may have thereby

shaped the development of many differences between oDSGC types.

The asymmetric spike tuning of Superior and Inferior oDSGCs is associated with

differences in the magnitude of excitatory synaptic input to each cell type (**Fig 3**). Three synaptic

partners are primary candidates for the source of this asymmetry: (1) glutamatergic input from

bipolar cells (types 5 and 7), (2) cholinergic input from SACs, and (3) glutamatergic input from

VGluT3 amacrine cells[44–46,49,50,58,77,78]. Glutamatergic conductances in oDSGCs rely on AMPARs

and NMDARs, whereas cholinergic conductances rely on nicotinic AChRs[79]. These conductances

have been studied primarily in ooDSGCs and deserve further characterization in oDSGCs.

Interestingly, VGluT3 amacrine cells also corelease glycine, which cancels oDSGC spiking in

response to high-velocity stimuli[49,57,77,78]. Asymmetries at the VGluT3-oDSGC synapse could

coincide with differences in the speed tuning properties of Superior and Inferior oDSGCs. A

combination of genetic, optical, and pharmacological manipulations could distinguish among

these possible sources of asymmetric tuning.

Further insight into how oDSGCs encode motion can be gained by comparing their tuning

properties to those of the more comprehensively studied ooDSGCs. While prior work has focused

on speed tuning as the primary difference between oDSGCs and ooDSGCs[14,26,49,57,78], our results

reveal an additional difference between these two classes of DSGCs that has been previously

overlooked: oDSGCs are contrast-sensitive (**Fig 7**), whereas ooDSGCs are not[62]. In ooDSGCs,

preservation of E/I across contrasts is apparently critical for retaining contrast-invariant direction

tuning[61,63]. Counterintuitively, we find that E/I in oDSGCs is also relatively stable across contrasts.

Therefore, stability of E/I is not alone sufficient for contrast-invariant spike tuning; independence

from thresholding effects is also required. Indeed, thresholding modulates oDSGC direction tuning

following changes to either E/I (**Fig 5-6**) or the absolute magnitude of E and I (**Fig 7**). Evidently,

the influence of thresholding constitutes a major difference in the mechanisms that govern how

oDSGCs and ooDSGCs encode motion.

That E/I is contrast-insensitive in both oDSGCs and ooDSGCs also indicates some extent

of shared circuitry between these cell types. Contrast-invariance of E/I in ooDSGCs relies on

postsynaptic NMDA conductances and constrains possible presynaptic wiring motifs[61,63].

oDSGCs likely share many of these features. Indeed, serial block-face electron microscopy has

confirmed that oDSGCs and ooDSGCs share many of their presynaptic partners[40,43,49,50,54].

Nonetheless, differences in the intrinsic properties between and within DSGC classes, including

dependence on thresholding, could magnify the impact of subtle circuit differences on spike

output.

Finally, our results show that vertical OKR is predicted by a simple subtraction between

the outputs of Superior and Inferior oDSGCs (**Fig 8**). It is interesting that asymmetries in oDSGCs

are not apparently corrected by downstream circuitry, considering that normalization operations

pervade the nervous system[80]. One explanation is that there is no simple compensatory solution

to normalize the responses of Superior and Inferior oDSGCs because multiple stimulus

parameters (e.g., stimulus direction and contrast) simultaneously affect the asymmetry

magnitude. On the other hand, the ethological advantage of asymmetric OKR may have provided

sufficient evolutionary pressure to allow asymmetries between Superior and Inferior oDSGCs to

propagate to behavior when they might otherwise have been compensated. Regardless, a linear

subtraction of oDSGC outputs offers an accurate and parsimonious explanation of vertical OKR.

Moreover, this algorithm fits well with both the anatomy and physiology of MTN and the isolation

of the vertical OKR pathway from other visual circuits[13,21,22,64–68].

Similar subtraction algorithms are prevalent across the animal kingdom. In the mammalian

retina, such a computation confers both the spatial center-surround[81–83] and chromatic

dichotomy[84–86] of receptive fields. In Drosophila, spatially offset antennae allow accurate

estimation of wind velocity by differentiation of signals across two input sites[87]. Similar

mechanisms likely underlie tropotaxic orienting behaviors that also rely on spatially offset

receptors, including arthropod antennae[88], reptile forked tongues[89], and mammalian ears[90]. The

ubiquity of this circuit motif may reflect an efficient solution for integrating complementary

information streams. Our results highlight how such circuits can be influenced by subtle

asymmetries across input channels. Further investigation will determine whether diverse sensory

systems rely on asymmetric inputs to adaptively change behavior across stimulus conditions.

METHODS

Statistics: All metrics reported in the text refer to population mean±SEM unless otherwise

specified. Nonparametric hypothesis tests were used to compute significance values wherever

possible. Mann-Whitney U tests were used for instances in which the test type is not specified.

Wilcoxon signed-rank and Fisher's exact tests were used where specified in the text and/or figure

legends. All tests were two-sided. R^2 values are Pearson correlation coefficients. Significance

markings are as follows: not significant (N.S.) $p \geq 0.05$, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

Animals: Adult wildtype C57BL/6 mice between the ages of postnatal day P60 and P100 of both

sexes were used for all experiments. Animals were kept on a 12 hour dark-12 hour light cycle with

continuous access to food and water. All experiments were performed in accordance with

protocols approved by the University of California, San Francisco Institutional Animal Care and

Use Program.

Behavior rig: To accurately evoke and measure OKR, we custom designed a behavior rig that

was capable of presenting full-field, binocular stimuli to behaving mice. The design of the rig was

based on Denman et al., 2017[38]. Briefly, an acrylic hemisphere (diameter=24", California Quality

Plastics) was covered with a custom paint that had 50% diffuse reflectivity between 350 and

750nm (Twilight Labs) in order to limit reflections within the hemisphere. Stimuli were emitted

from a DLP projector with peak emission at 405nm (LightCrafter through EKB Technologies) and

were reflected onto the hemisphere via a silver-coated brass hemisphere ("convex mirror",

diameter=6", Wagner). Stimuli were built using Psychopy (www.psychopy.org) and a custom

wrapper to manage their sequential presentation and alignment with eye tracking videos. The

wrapper and stimuli are both available at https://github.com/ScottHarris17/Bassoon. Aberrations

in the projection were corrected by applying a manually-fit spherical morph to all stimuli

(Meshmapper, www.paulbourke.net). Blackout curtains surrounded the rig to minimize light

contamination.

Unidirectional OKR stimuli: Unidirectional sinusoidal gratings were presented in groups of 6

consecutive epochs. Each epoch consisted of 20s of a static grating, followed by 60s of a grating

drifting either directly upward or directly downward, and an additional 20s of a static grating. The

6 total epochs consisted of 3 upward and 3 downward epochs that were randomly interleaved. All

gratings moved at 10°/s and had a spatial frequency of 0.15 cycles/°. The brightest part of the

grating evoked 1.38×10^3 S-cone photoisomerizations/s and the darkest part of the grating evoked

25.9 S-cone photoisomerizations/s. M-cone photoisomerizations were 60% those of S-cone.

Oscillating OKR stimuli: Oscillating sinusoidal gratings were presented in groups of 3

consecutive epochs. Each epoch consisted of 20s of a static grating, followed by 120s of

oscillation, and an additional 20s of a static grating. During the oscillation, the grating velocity was

modulated sinusoidally up and down. The oscillation had an amplitude of 20°, a period of 15s,

and a phase shift of 0°. 8 oscillations were completed over the course of one epoch. All gratings

had a spatial frequency of 0.15 cycles/°. The intensities of high contrast oscillating gratings were

equivalent to those used for unidirectional OKR stimuli and had 5-fold greater Michelson contrast

than low contrast oscillating gratings. High and low contrast gratings had the same mean

luminance.

Eye tracking: Prior to eye tracking experiments, animals underwent stereotaxic surgery for

implantation of a custom head-fixing apparatus. After surgery, animals were given 7 days to

recover. Animals were then gradually habituated to the behavior rig by spending increasing

amounts of time head-fixed on the rig for five consecutive days prior to the beginning of

experiments.

Eye tracking experiments were run over the course of up to 3 days per animal. Each

animal spent no more than 30 minutes per day on the rig. During experiments, mice were head-

fixed in the center of the hemisphere, which filled the entirety of their visual field. Eye movements

were recorded using a GigE camera (Photonfocus), an infrared filter, and a hot mirror (Edmund

Optics) that allowed the camera to be positioned outside of the animal's field of view. Infrared

LEDs (880nm) were mounted on the top and side of the camera to generate corneal reflections

that marked the meridian and equator of the eye, respectively. StreamPix software (NorPix) was

used to capture video of the eye and align it to stimuli via TTL signals.

After completion of the experiments, Deeplabcut (www.deeplabcut.org) was used to train

a neural network to locate the pupil and corneal reflection on each video frame. The two

dimensional pupil location was then translated to angular eye position for every recording frame

using the methods described by Stahl et al., 2000, 2004[91,92] and Zoccolan et al., 2010[93]. In short,

prior to experiments, we calibrated the eye tracking system for each animal by repeatedly

swinging the camera ±6° in the horizontal plane and measuring the relative position of the pupil

and corneal reflection. This process was repeated across 5 different luminances to fit a linear

regression between pupil size and angular eye position. The meridian and equator of the eye

were measured by turning on the top- and side-aligned infrared LEDs in sequence. During

experiments, only the top LED was on. Angular eye position was then calculated as $\phi =$

$arcsin(\frac{\Delta x}{\sqrt{R_{p0}^2 - \Delta y^2}})$ [Eq. 1] and $\theta = arcsin(\frac{\Delta y}{R_{p0}})$ [Eq. 2], where ϕ is the horizontal eye position, θ

is the vertical eye position, Δx is the horizontal distance measured between the eye center and

the pupil, Δy is the vertical distance measured between the eye center and the pupil, and R_{p0} is

the radius of rotation between the pupil and the eye's center, which was computed empirically for

each pupil diameter for each animal. For all stimuli, fast and slow phases of nystagmus were

separated on the basis of eye velocity and acceleration using custom MATLAB scripts. For all

analyses in this report, we consider only θ, the vertical eye position, and $\Delta\theta/\Delta t$, the vertical

component of eye velocity.

By measuring the vertical eye velocity ($\Delta\theta/\Delta t$) in response to the static gratings (multiple

contrasts) that occurred before the onset of all stimuli, we also computed a baseline median eye

drift of 0.0787°/s (IQR: 4.1792°/s; p=0.015) in the ventral direction across animals. This drift may

reflect either the error associated with the estimation of the eye's center during our calibration

process, or a natural biological eye drift associated with our rig. It is not possible to disambiguate

between these possibilities. Since the eye position was near neutral for the time in which this drift

was calculated, we baseline subtracted it from the eye position traces for all oscillating grating

stimuli (for which the average eye position was also approximately neutral). For unidirectional

gratings, we were unable to calculate an appropriate drift for baseline subtraction since the eye

position was not often near neutral during these stimuli (**Fig S6**).

Retrograde labeling: MTN-projecting retinal ganglion cells were labeled via stereotaxic injection

of red fluorescent retrobeads (Lumafluor) into MTN. Prior to surgery, mice were anesthetized by

IP injection of ketamine/xylazine and 5% isoflurane inhalation. Once fully anesthetized, as

assessed by absence of the pedal reflex, animals were transferred to a sterile heated surface and

the eyes were covered with a lubricating ointment. 2% isoflurane was administered continuously

to maintain anesthesia. Fur was removed prior to incision and lidocaine (<7mg/kg) was injected

locally under the scalp. After incision, animals' heads were leveled by aligning bregma and

lambda in the horizontal plane. A burr hole was drilled at A/P: 0.00mm, M/L: 0.85mm from bregma.

All injections were performed into the right MTN. A glass needle filled with retrobeads (diluted 1:3

in di-water) and connected to a Hamilton syringe was lowered into the burr hole at an angle of

30° A/P to a depth of 5.36mm below the surface of the brain. After 10 minutes, an injection of

400nL was made at a rate of 5nL/s. After injection, the needle was left in place for an additional

10 minutes before removal. The scalp was sutured and animals recovered in a heated cage.

Analgesics [buprenorphine (0.05-0.1mg/kg) and NSAIDs (5-10mg/kg)] were delivered via

subcutaneous injection immediately after animals awoke from anesthesia, again 12 hours later,

and a third time the following morning. Animal health was monitored for three days after surgery

and additional analgesics were administered as required. Labeling of retinal ganglion cells in the

contralateral eye was typically observed as soon as 48 hours following surgery and did not

increase with time.

Empirical mosaic analysis: All mosaic analyses occurred ≥2 days after injection of the

retrograde tracer into MTN. Retinas were dissected, fixed in 4% PFA for 20 minutes, and flat-

mounted onto a microscope slide with a spacer. One widefield fluorescent image was taken of

each retina. The location of each labeled cell in the image was determined using custom MATLAB

scripts. Only retinas with near complete labeling (determined as greater than 500 identified RGCs)

were included in analyses. The retina perimeter, optic nerve head, and dorsal-ventral axis were

manually measured. These points were used to define a normalized polar coordinate system that

allowed for the comparison of cell locations and densities across retinas. Density recovery profiles

were calculated using the methods described by Rodieck, 1991[94].

Mosaic models: To model spatial distributions of single and multiple mosaics, we randomly

generated mosaics in a model circular retina that had an equivalent radius to that of the average

whole-mount retina used for empirical density recovery profile estimation. "Cell bodies" were

modeled as circles with radius $15\mu m$ and scattered randomly across the model retina so long as

the following conditions were met: (1) No two cell bodies can overlap in space (i.e., cells must

form a 'monolayer'), and (2) Adjacent cells that are members of the same mosaic must obey a

(noisy) exclusion zone that is set by the mosaic coverage factor (number of cells/retina area).

Coverage factors were changed systematically such that the total number of cells across all

mosaics - regardless of the number of mosaics being modeled - always approximated the number

of retrogradely labeled cells per retina in our empirical data set. Density recovery profiles were

computed as described for empirical data.

Electrophysiology tissue preparation: All electrophysiology experiments occurred ≥2 days

after injection of the retrograde tracer into MTN. Prior to electrophysiology experiments, mice

were dark-adapted for ≥12 hours. Mice were then euthanized via cervical dislocation and the left

eye was enucleated (contralateral to the right MTN injection). Retina dissections occurred in the

dark using infrared converters and warmed bicarbonate-based Ames solution, equilibrated with

95% O_2/5% CO_2. Brains were simultaneously harvested and fixed in 4% PFA for imaging and to

confirm that the retrograde tracer was properly injected into MTN. Retinas were whole mounted,

keeping track of orientation, and continuously perfused at 10mL/min with freshly equilibrated

Ames heated to 35°C throughout the course of experiments.

Retinal location of recorded cells: At the beginning of electrophysiology experiments, the center and radius of the retina were measured: two-dimensional coordinates of 8 standardized points around the perimeter of the retina were noted. The center of the retina was estimated by computing the median of the circumcenters of all unique triangles that could be formed from these 8 points. The radius of the retina was estimated by finding the median radius of the circles that circumscribed these same triangles. In all cases, the convex hull of the whole-mount retina was well approximated by a circle and the retina center estimation was near the optic nerve head. The coordinates of each recorded cell were computed in reference to the retina center. Cell locations were combined across retinas by normalizing to the estimated radius in a polar coordinate system.

Identification of MTN-projecting RGCs: An NIR light source (950nm, Sutter) was used to visualize the tissue for the majority of the experiment. To identify retrogradely-labeled ganglion cells, a green epifluorescent light was turned on briefly (~1-3s) prior to recording from each cell. This light evoked a moderate number of spikes in oDSGCs, but did not compromise their responses to other stimuli. At least 1 minute of darkness was provided between the offset of the epifluorescent light and the beginning of subsequent experiments.

Electrophysiology: Patch electrodes were pulled from borosilicate glass (Sutter) to 3-5MOhm resistance using a Narishige puller. A MultiClamp 700B Amplifier (Axon Instruments) was used for all recordings. Cell-attached experiments were performed using electrodes filled with HEPES buffered Ames. Voltage-clamp experiments were performed using fresh electrodes filled with cesium methane sulfonate[95]. Current-clamp experiments were performed using fresh electrodes filled with potassium aspartate[96]. A subset of cells were recorded in both cell-attached and whole-cell configurations. In these cases, cell-attached recordings were performed first. Voltage-clamp and current-clamp recordings were never made from the same cell. Electrophysiology

experiments were conducted using Symphony DAS (https://symphony-das.github.io/) and light

stimuli were constructed and presented using Stage (https://stage-vss.github.io/).

Light increment stimulus: Light increments were the first stimulus to be presented, and were

often additionally interleaved between other stimuli. Light increments were delivered from an LED

with peak emission at 405nm from darkness for 1s, and had an intensity of 8.6×10^4 S-cone

photoisomerizations/s. M-cone photoisomerizations were 79% those of S-cone. The LED spot

had diameter 500μm or 300μm (no significant difference was observed in the responses to either

spot size), was centered on the cell body of each recorded cell, and was focused on the

photoreceptor layer of the retina.

Drifting bar stimulus: Drifting bars were presented from a DLP projector with peak at 405nm

(LightCrafter through EKB Technologies, same model as used for behavioral experiments). The

native optics of the projector were replaced with neutral density filters and optics to focus stimuli

on the photoreceptor layer of the retina via a condenser. The projector covered a rectangular area

of $427 \times 311\mu$m which was centered on the soma of each recorded cell. Drifting bars had a width

of 3.2° and moved at 10°/s using a conversion factor of 31μm/° [97]. Bar height was limited only by

the area covered by the projector. High contrast bars measured 2.4×10^4 S-Cone

photoisomerizations/s and were presented on top of a background of 124 S-cone

photoisomerizations/s. M-cone photoisomerizations were 74% those of S-cone. See below for the

specifications of low contrast bars. For tuning curve estimation, bars moved in 8 directions

separated by 45°. The presented sequence of stimulus directions was randomized for each

recording. Tuning curves were estimated by mean measurements taken over 5 repetitions per

stimulus direction.

Tuning curve area: Tuning curve area was calculated by dividing the area under the curve of the

linear tuning curve by 360°. For clarity, this metric is also referred to as the "linear tuning curve

area."

Preferred direction: The preferred direction was calculated as the direction of the vector sum of

responses to all 8 stimulus directions. Thus, the preferred direction was not necessarily equivalent

to the single stimulus direction that evoked the largest response. The null direction was defined

as 180° away from the preferred direction.

Normalized tuning curves: Normalized tuning curves were calculated by first determining the

cell's preferred direction (see above), and then dividing the response in all 8 stimulus directions

to the response in that preferred direction. For cases in which the preferred direction did not match

a stimulus direction that was specifically probed, the preferred direction response was estimated

by a linear interpolation of the two neighboring probed directions. The area of the normalized

tuning curve (abbreviated as the "normalized area") was calculated by dividing the area under the

curve of the linear normalized tuning curve by 360°. The area of the normalized tuning curve is

always greater than 0, but has no upper bound - though it tended to fall below 1. Larger values

indicate a wider tuning curve. Perfectly circular tuning curves take a value of 1.

Direction selectivity index: The direction selectivity index (DSI) was calculated as the

magnitude of the vector sum divided by the scalar sum of responses to all 8 stimulus directions.

The direction selectivity index ranges between 0 and 1, with larger values indicating sharper

tuning curves.

Von Mises fit: Tuning curves were fit to the Von Mises function by minimizing the sum of squared

residuals. The Von Mises function is a circular analog of the Gaussian curve defined as $f(x) =$

$\frac{e^{\kappa cos(x-\mu)}}{2\pi I_0(\kappa)}$ [Eq. 3], where μ is the center of the curve, $1/\kappa$ controls the width of the curve, and I_0 is

the modified Bessel function of the first kind, of order 0. A larger κ value indicates a sharper fit.

Retinal ganglion cell classification: Retrogradely-labeled retinal ganglion cells were classified

as either Superior or Inferior oDSGCs if they had a direction selectivity index of greater than 0.05

and a preferred direction more than 30° off of the temporal-nasal axis, as calculated by spike

outputs measured in the cell-attached configuration. The vast majority of recorded cells met these

criteria, but those that did not were excluded from further analyses (**Fig S1**). All recorded cells in

our data set were dominated by ON responses to a light increment (**Fig S4**).

Current injections: Depolarizing or hyperpolarizing currents were continuously injected while

measuring voltages across stimulus directions in the current-clamp configuration. The magnitude

of current injections changed subtly from cell to cell depending on resting membrane potential,

input resistance, and spike threshold. On average, depolarizing current injections increased the

membrane potential by ~6mV (to ~-48mV), whereas hyperpolarizing current injections decreased

the membrane potential by ~6mV (to ~-60mV). Depolarizing injections were always small enough

such that the new resting membrane potential remained below spike threshold and each cell's

baseline firing rate was 0hz. The order of depolarizing and hyperpolarizing injections was random

across cells.

Isolation of spikes and subthreshold voltages: From current-clamp recordings, the onset and

offset of each action potential were determined using the first and second derivatives of the

voltage trace and a fixed minimum refractory period. The subthreshold voltage was then linearly

interpolated between action potential onsets and offsets.

Subthreshold voltage tuning curves: The maximum voltage deflection from baseline was used

to determine subthreshold membrane potential tuning curves. Values were averaged over 5

repetitions for each stimulus direction. Tuning curve metrics were calculated as for spikes.

Electrophysiology at low contrast: Low contrast drifting bars had an intensity of 0.5×10^4 S-cone

photoisomerizations/s (~5-fold dimmer than high contrast bars) from the same background of 124

S-cone photoisomerizations/s. All other stimulus parameters were equivalent across contrasts. In

a subset of cells, low contrast bars failed to elicit spikes for every stimulus direction. In such cases,

the cell's spike tuning curve area was set to 0, the area of its normalized tuning curve was set to

0, and its direction selectivity index was set to 1. We chose this convention because of the

observation that, across cells, as the total number of spikes approached 0, the area of the

normalized tuning curve approached 0 and the direction selectivity index approached 1. This

pattern also fits the prediction made by our leaky integrate-and-fire model (**Fig 7I**). Cells with no

responses under low contrast were classified as Superior or Inferior on the basis of their

responses to high contrast stimuli.

Immunohistochemistry: Individual ganglion cells were filled with either Lucifer yellow or biocytin

during electrophysiology experiments. Retinas were subsequently fixed in 2% paraformaldehyde

for 20 minutes at room temperature. The following protocol was used to enhance for the cell fills

[anti-Lucifer yellow (Life Technologies A5750) and/or streptavidin-488 (Thermo Fisher S11223)],

label synaptic puncta [anti-postsynaptic density (PSD-95, UC Davis NeuroMab 75-028), anti-

Gephyrin (Synaptic Systems 147 111)], and stain for cholinergic starburst amacrine cells [anti-

choline acetyltransferase (ChAT, Millipore AB144P)]: blocking serum (1 day), primary antibody

incubation (5 days), rinse 3x in PBS, secondary antibody (Jackson ImmunoResearch) incubation

(1 day), rinse 3x in PBS. Retinas were mounted with a spacer in vectashield (Vector labs) and

under a coverslip.

Imaging: Individual ganglion cells with known direction selectivity and their associated synaptic puncta were imaged on a confocal microscope (Leica SP8). Ganglion cells were imaged with a 40x objective (NA 1.3) at a resolution of 0.102 x 0.102 x 0.3μm.

Image analysis: Confocal images were first median filtered in 3 dimensions (Fiji). Ganglion cell dendrites were reconstructed using the filament function in Imaris (Oxford Instruments). Convex polygons, dendritic branch numbers, and total dendritic length were obtained from the filaments. Excitatory PSD-95 and inhibitory gephyrin puncta were identified and quantified within the filament mask of the ganglion cell dendrites (ObjectFinder, https://lucadellasantina.github.io/ObjectFinder/). Quantification of the puncta density from the center to periphery and along dendritic depth were also obtained in ObjectFinder.

Electrical properties of oDSGCs: The resting membrane potential, spike threshold, and input resistance of oDSGCs were all measured during whole-cell current-clamp recordings. Resting membrane potential was taken as the initial membrane voltage immediately after establishing intracellular access. Spike threshold and input resistance were both calculated by injecting a slow ramp of current. Spike threshold was the average voltage at which a cell initiated its first action potential in response to the ramp. Input resistance was calculated from the average slope of the I-V response below spike threshold. Both metrics were averaged over at least 5 repetitions of the ramp stimulus per cell. The membrane capacitance was measured during voltage-clamp recordings using the built-in capacitance calculator from the MultiClamp 700B Amplifier (Axon Instruments).

Parallel conductance model: We implemented a parallel conductance model in MATLAB to build the model oDSGC (adopted from Antoine et al., 2019[98]). Excitatory (Gex) and inhibitory (Gin)

conductances were calculated at each time point for each direction of stimulus motion using

Ohm's law: $G = I/(Vhold - Erev)$ [Eq. 4], where I was the mean current trace recorded in

voltage-clamp across both Superior and Inferior cells, $Vhold$ was the holding potential, and $Erev$

was the reversal potential for either excitation or inhibition. A liquid junction potential of 5mV was

subtracted from $Vhold$ and $Erev$. Because our model called for directionally untuned excitation,

but the recorded excitatory conductances were slightly different for each direction of stimulus

motion (likely due to space-clamp error), we used an identical excitatory conductance for all

directions of stimulus motion that was equal to the maximum conductance at each time point

across all recorded directions (space-clamp errors reduce empirically recorded excitatory currents

in voltage-clamp mode). The excitatory conductance was then multiplied by a gain value to

achieve a final time series for Gex. Next, we used the equation

$C\frac{dV}{dt} = Gex(Eex - Vm) + Gin(Ein - Vm) + Grest(Erest - Vm)$ [Eq. 5]

to determine the membrane potential at each time point. C was the median capacitance of

Superior and Inferior oDSGCs as measured during whole-cell recordings. $Grest$ was the

reciprocal of the median input resistance, which we calculated by injecting a slow ramp of current

in a subset of recorded cells and determining the average slope of the I-V response below spike

threshold. $Erest$ was the median resting membrane potential. Vm was calculated at each point

in time by initializing it at $Erest$, and then determining each subsequent value using Euler's

method and an integration time step of 1ms. The peak change in this Vm value above $Erest$ was

used to construct the Vm tuning curves in the version of the model without a spiking component.

In the version of the model with a spiking component, we again solved for Vm at every

time point using Euler's method. In this case, however, whenever Vm surpassed the threshold

potential, a "spike" was counted and a 3ms pause corresponding to the spike time and refractory

period was initiated, after which Vm was reset to $Erest$ and the process continued. The threshold

value was fixed such that the the normalized area and direction selectivity index of the model

oDSGC's spike tuning curve matched the median empirical values for these metrics (taken from

cell-attached recordings) when the excitatory gain was set to 1.0. This resulted in a threshold

value of -46.2mV, which was 2.9mV more negative than the median empirically recorded spike

threshold.

Because resetting Vm to $Erest$ after each spike also changed the driving forces for

excitation and inhibition - and therefore possibly the shape of resulting spike tuning curves - we

also simulated spike responses by assuming that the number of spikes produced to a given

stimulus was linearly proportional to the amount of time that Vm (calculated without a spiking

mechanism in place) spent above spike threshold (data not shown). Results from this model were

not substantively different than those from the model in which a refractory period was included

and Vm was reset to $Erest$ after each spike.

This same model was used to test the contrast-dependence of spike and Vm tuning

curves. In this case, both the excitatory and each of the 8 inhibitory conductance time series were

multiplied by the gain value to calculate Gex and Gin, respectively.

Behavioral predictions from oDSGC activity: Predictions for OKR gain were calculated on the

basis of the difference in median firing rates between Superior and Inferior oDSGC populations.

The preferred and null directions of each cell in our dataset were computed in response to a high

contrast drifting bar. Gain predictions were made under each stimulus condition by (1) resampling

from distributions of oDSGC responses for that condition 10,000 times, (2) computing the

difference (i.e., delta) between the median preferred and null direction responses on each iteration

for the appropriate cell types, and (3) using the median of these bootstrapped distributions of delta

as an estimate of relative gain.

Computing the preferred and null direction of each cell has the advantage of controlling

for the fact that the observed preferred direction of individual oDSGCs can change based on

retinotopic location when the retina is flatmounted[16]. Assigning the preferred direction to a

constant stimulus direction across cells (e.g. dorsal-to-ventral motion on the retina for Superior

oDSGCs) did not change the behavioral predictions (data not shown).

Empirical nonlinearity: oDSGC spike outputs were related to OKR gain by computing an

empirical nonlinearity using the Hill equation, defined as $g(r) = g_{min} + \frac{g_{max}-g_{min}}{1+(\frac{r_{1/2}}{r})^n}$ [Eq. 6], where

g_{min} is the minimum OKR gain value, g_{max} is the maximum OKR gain value, $r_{1/2}$ is the ganglion

cell response (defined as the difference between Superior and Inferior oDSGC firing rates) that

corresponds to 50% of g_{max}, and n sets the steepness of the curve. g_{min} was fixed at 0 such that

the nonlinearity would predict a gain of 0 for an oDSGC response of 0. g_{max} was fixed at 1.0 such

that the nonlinearity would not predict gains above 1.0. The empirical nonlinearity was then

determined by using a least-squares fit of Eq. 6 to the four measured data points (oDSGC

responses and OKR gain values measured for superior/inferior stimuli at high/low contrast).

ACKNOWLEDGEMENTS

We would like to thank Connie Chen and Jeremiah John for technical assistance; Kevin Bender,

David Copenhagen, Daniel Denman, Tonatiuh Garcia Ruiz, Jonathan Horton, Jeanette Hyer,

David Kastner, Joo Yeun Lee, Yien Ming-Kuo, Satoru Miura, Yvonne Ou, Luca Della Santina,

Massimo Scanziani, Manuel Solino, and Alfred Yu for helpful discussions; David Berson, Guy

Bouvier, and Fred Rieke for comments on the manuscript. This work was supported by NIH

through F31 EY-033225 (S.C.H.), R01 EY-029772 (F.A.D.), R01 EY-030136 (F.A.D.), and the

Vision Core Grant P30 EY-002162 (UCSF), internally by the Moritz-Heyman Discovery Fellowship

(S.C.H.) and the Kavli Institute for Neuroscience (UCSF), and through foundation grants from the

McKnight Scholar Award (F.A.D.), Research to Prevent Blindness (Unrestricted Grant), and That

Man May See.

AUTHOR CONTRIBUTIONS

990 Both authors conceptualized the project, designed experiments, and constructed experimental

991 rigs. S.C.H. conducted and analyzed the behavioral and electrophysiological experiments and

992 performed the modeling. F.A.D. conducted and analyzed the confocal imaging. Both authors

993 wrote the manuscript and acquired funding and resources.

COMPETING INTERESTS

998 The authors declare no competing interests.

Figure 1

Figure 1. Superior and inferior OKR are asymmetric in adult mice. (a) Schematic of behavioral setup to elicit the vertical optokinetic reflex (OKR). The mouse is situated so that one eye is centered in a hemisphere. Stimuli are projected onto the hemisphere's concave surface via reflection off of a convex mirror. Eye movements are tracked using an infrared-sensitive camera and corneal reflection (see Methods). **(b)** Example video frames demonstrating that the eye traverses between superior, neutral, and inferior positions in the presence of vertically-drifting sinusoidal gratings. Red arrows mark the infrared corneal reflection. **(c-d)** Vertical angle of the eye across an example epoch in response to full-contrast **(c)** superior and **(d)** inferior drifting gratings (10°/s). For each epoch, a continuous 60s stimulus was flanked by 20s of a static grating (shaded regions). Ticks above the plots mark the time of fast nystagmus either in the superior (magenta) or inferior (gray) direction. Examples from one animal. **(e)** Rate of vertical fast nystagmus for superior and inferior stimuli. Horizontal line represents median, box boundaries are the IQR, whiskers represent most extreme observation within 1.5xIQR. **(f)** Cumulative vertical distance traveled during slow nystagmus in response to superior (magenta) and inferior (gray) drifting gratings after removing fast nystagmuses (mean±SEM). **(g-h)** Eye position in response to a grating oscillating sinusoidally along the vertical axis. For each epoch, a continuous 120s stimulus, consisting of 8 oscillation cycles, was flanked by 20s of a static grating (shaded regions). Stimulus position in lavender. **(g)** Raw vertical eye position (blue) including the presence of fast nystagmuses (ticks). **(h)** Eye position after removing fast nystagmuses (green). **(i)** Average gain of slow nystagmus during the superior versus inferior phase of individual oscillations. Each small dot is a single oscillation. The region of magenta [or gray] indicates that gain was greater for the superior [or inferior] phase of the oscillation. Points that fall on the line indicate equivalent gain for both stimulus directions. Large point and whiskers represent univariate medians and 95% confidence intervals (via bootstrapping), respectively. Significance value indicates whether the points tend to fall unevenly on one side of the unity line (2-sided signed-rank). **(j)** Eye position (green) and stimulus position (lavender) averaged across all oscillations and all animals

1025   (mean±SEM). Starting eye position is normalized to 0° at cycle onset. The average ending eye

1026   position is displaced in the superior direction (2-sided signed-rank). N=5 mice for all experiments;

1027   n=number of trials. See also Fig S6.

Figure 2

Figure 2. Superior and Inferior oDSGCs have asymmetric spike tuning curves. **(a)**

Schematic illustrating unilateral bead injections into medial terminal nucleus (MTN) to retrogradely

label ganglion cells in the contralateral retina. **(b)** Flat-mount retina with retrogradely labeled,

MTN-projecting ganglion cells. **(c)** Drifting bar stimulus (3.2° x limiting projector dimension, 10°/s,

2.4×10^4 S-cone photoisomerization/s). **(d)** Definitions of superior (magenta) and inferior (gray)

motion in visual space and on the retina. Directions are inverted by the lens. **(e)** Cell-attached

spikes from labeled retinal ganglion cells in a flat-mount retina in response to a bar drifting in 8

directions. Spike responses and average tuning curves from an example Superior (left, magenta)

and Inferior (right, gray) oDSGC. Mean spike counts are presented as the distance from the origin,

marked by concentric circles. Numbers on circles indicate spike counts. Dashed lines represent

the preferred direction of each cell, calculated as the direction of the vector sum of all responses.

Coordinates are in retinal space. **(f)** Population tuning curves across all Superior and Inferior

oDSGCs (mean±SEM). Left: polar plots [as in (e)] aligned by rotating the tuning curves of Superior

cells by 180°. Right: linear representation of the same data (referred to as the "linear tuning

curve"). CW: clockwise; CCW: counterclockwise. 0° represents directly superior/inferior motion.

(g) Histograms of the area under the curve of the linear tuning curve of every cell. Inset shows

the same metric for a stimulus at 5-fold lower contrast. **(h)** Population mean (±SEM) normalized

tuning curves - computed by normalizing and aligning (at 0°) the response of each cell to its

response in the preferred direction. Vertical dashed lines indicate the point at which mean

response magnitudes drop below 50% of the preferred direction response. **(i)** Distributions of the

distance (in degrees) from each cell's preferred direction to the point at which its response

magnitude crosses the 50% threshold [dashed lines in (h)]. Horizontal line represents median,

box boundaries are the IQR, whiskers represent most extreme observation within 1.5xIQR. Points

represent individual cells. **(j)** Histograms of the area under the curve of the normalized tuning

curve [as in (h)] of every cell (abbreviated as "normalized area"). A larger normalized area

indicates a broader tuning curve (see Methods). Inset shows the same metric for a stimulus at 5-

fold lower contrast. **(k)** Histograms of the kappa parameter for the Von Mises fit of the tuning curve

of every cell (see Methods). A smaller kappa value indicates that the fit is wider. **(l)** Histograms of

the direction selectivity index (DSI, vector sum, see Methods) of every cell. Inset shows the same

metric for a stimulus at 5-fold lower contrast. **(m)** Linear tuning curve area, i.e. "magnitude," [as

in (g)] versus area of the normalized tuning curve, i.e., "width," [as in (j)] were positively correlated

on a cell-by-cell basis for both Superior and Inferior oDSGCs. For all histograms, medians of

Superior (magenta) and Inferior (gray) oDSGC distributions are indicated by arrows. *p<0.05,

**p<0.01, ***p<0.001. See also Fig S1, S2, S3.

a Inhibition
V-clamp (Inh)
600pA

b V-clamp (Inh)
100pA
1s
600pA

c Peak
N.S.
600
pA
300
-180 -90 0 90 180
n=19 Sup.
20 Inf.

d Charge Transfer
N.S.
400
pC
200
-180 -90 0 90 180
Aligned Stimulus Direction (deg)

e
DSI
0.2
0.1
*
p=0.027
Superior   Inferior

f Excitation
V-clamp (Exc)
300pA

g V-clamp (Exc)
100pA
1s
300pA

h Peak
* * * * * * * * * * *
240
|pA|
120
-180 -90 0 90 180
n=17 Sup.
17 Inf.

i Charge Transfer
* * * * * * * * *
120
|pC|
60
-180 -90 0 90 180
Aligned Stimulus Direction (deg)

j
DSI
0.2
0.1
*
p=0.039
Superior   Inferior

k E/I Magnitude
1
Tuning Curve Area
0.5
**
p=5.3x10⁻³
n=17 Sup., 17 Inf.
Sup.   Inf.

l E/I Shape
N.S.
DSI
0.2
0.1
p=0.39
Sup.   Inf.

m E vs. I Shape
***
DSI
0.2
0.1
p=3.19x10⁻⁴
Inh.   Exc.

Figure 3

Figure 3. Superior oDSGCs receive similar inhibitory inputs but greater excitatory inputs compared to Inferior oDSGCs. (a) Inhibitory currents measured from an exemplar Superior oDSGC under voltage-clamp at +10mV in response to a bar drifting in 8 directions. **(b)** Same as (a) for an exemplar Inferior oDSGC. **(c-d)** Population responses for inhibitory currents. Stimulus direction is on the abscissa and **(c)** peak inhibitory current or **(d)** total inhibitory charge transfer is on the ordinate for Superior (magenta) and Inferior (gray) oDSGCs (mean±SEM). Stimulus directions are aligned across cell types, where 0° indicates directly superior (for Superior oDSGCs) or inferior (for Inferior oDSGCs) motion. Positive directions are clockwise. **(e)** Distributions of the direction selectivity index for peak inhibitory currents in individual Superior and Inferior oDSGCs. **(f)** Excitatory currents measured from an exemplar Superior oDSGC under voltage-clamp at -60mV in response to a bar drifting in 8 directions. Same cell as in (a). **(g)** Same as (f) for an exemplar Inferior oDSGC. Same cell as in (b). **(h-i)** Population responses for excitatory currents. Aligned stimulus direction is on the abscissa and **(h)** peak excitatory current or **(d)** total excitatory charge transfer is on the ordinate for Superior (magenta) and Inferior (gray) oDSGCs (mean±SEM). **(j)** Distributions of the direction selectivity index for peak excitatory currents in individual Superior and Inferior oDSGCs. **(k-l)** The ratio of the peak excitatory current to the peak inhibitory current (E/I) was calculated for each stimulus direction for cells in which both metrics were recorded. **(k)** Distributions of the linear tuning curve area of E/I. Arbitrary units (E/I/360°). **(l)** Distributions of the direction selectivity index for E/I. **(m)** Direction selectivity index for peak inhibitory (blue) and excitatory (yellow) currents collapsed across Superior and Inferior oDSGCs. For box plots, horizontal line represents median, box boundaries are IQR, and whiskers represent most extreme observation within 1.5xIQR. *p<0.05, **p<0.01, ***p<0.001. See also Fig S3, S4.

a Superior   Inferior

Ganglion cell   ChAT

b Dendritic Field Area

***

p=5.60x10⁻⁵
n=27 Sup., 25 Inf.

Superior   Inferior

c Sholl Analysis

n=13 Sup., 12 Inf.

d Lucifer yellow   PSD-95

e Density

n=13 Sup., 11 Inf.

f Count

N.S.

p=0.10   n=13 Sup. 12 Inf.

g Volume

**

p=0.001

h Lucifer yellow   Gephyrin

i

n=11 Sup., 6 Inf.

j N.S.

p=0.25   n=11 Sup. 6 Inf.

k N.S.

p=0.22

Figure 4

Figure 4. Superior oDSGCs have greater dendritic fields and larger excitatory postsynaptic sites. (a) Confocal images of a Superior (left) and Inferior (right) oDSGC filled with dye. Convex polygons are drawn around the tips of their dendrites. (Bottom) Side views of different Superior and Inferior oDSGCs filled and stained for acetylcholinesterase (ChAT) bands. Both cell types have dendrites that stratify in the ON and OFF ChAT bands, with the majority of dendrites in the ON sublamina. **(b)** Convex polygon areas across cells. **(c)** Sholl analysis indicating the number of dendritic crossings as a function of radial distance from the soma (mean±SEM). **(d, h)** Ganglion cells with immunostaining for **(d)** excitatory postsynaptic scaffolding protein PSD-95 or **(h)** inhibitory postsynaptic scaffolding protein gephyrin. (Bottom) Magnification of a stretch of dendrites with labeled puncta. **(e, i)** Quantification of puncta density as a function of distance along the dendrite for **(e)** PSD-95 and **(i)** gephyrin (mean±SEM). **(f, j)** Total number of puncta within each ganglion cell for **(f)** PSD-95 and **(j)** gephyrin. **(g, k)** Quantification of average puncta volume for **(g)** PSD-95 and **(k)** gephyrin. For box plots, horizontal line represents median, box boundaries are IQR, and whiskers represent most extreme observation within 1.5xIQR. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. See also Fig S3, S4.

Figure 5

Figure 5. The spike threshold differentiates the tuning properties of Superior and Inferior

oDSGCs. (a-b) Exemplar Inferior oDSGC in whole-cell current-clamp during **(a)** depolarizing and

(b) hyperpolarizing current injection in response to a bar moving in 8 directions. **(c-d)** Mean

(±SEM) normalized tuning curves (aligned and normalized to the response of each cell in its

preferred direction) for **(c)** Superior and **(d)** Inferior oDSGCs under conditions of depolarizing

(green) and hyperpolarizing (purple) current injection. Dotted lines indicate the average

normalized spike tuning curves of each population from cell-attached recordings (as in Fig 2h).

(e) For Superior oDSGCs, histograms of the area of the normalized tuning curve (left) and

direction selectivity index (right) under conditions of no current injection (i.e. cell-attached,

magenta), depolarizing injection (top, green), and hyperpolarizing injection (bottom, purple). **(f)**

Same as (e) for Inferior oDSGCs. Gray histograms represent cell-attached recordings. **(g)**

Illustration of the influence of untuned excitation on the tuning curve through additive (yellow) and

thresholding (red) effects. The blue area indicates the membrane potential and the dashed red

line indicates the spike threshold. **(h, l)** Example whole-cell current-clamp recording in which **(h)**

spikes and **(l)** subthreshold voltages (Vm) have been separated. **(i-k, m-o)** Scatter plots of tuning

curve metrics during hyperpolarizing (abscissa) and depolarizing (ordinate) current injections for

(i-k) spikes and **(m-o)** subthreshold voltages measured simultaneously: **(i, m)** linear tuning curve

area, **(j, n)** area of the normalized tuning curve, **(k, o)** direction selectivity index. Regions of green

[or purple] indicate that the metric is greater during depolarizing [or hyperpolarizing] injections.

Points that fall on the line indicate equivalent metrics under the two conditions. Individual cells are

shown as dots (Superior in magenta, Inferior in gray). Red and blue circles represent univariate

medians (collapsed across cell type) and whiskers indicate 95% confidence intervals determined

via bootstrapping. Significance values indicate whether the data tend to fall unevenly on one side

of the unity line (2-sided signed-rank). Arrowheads in (m-o) represent the median of Superior

(magenta) and Inferior (gray) oDSGCs along the line of slope unity, and associated significance

values indicate comparison between Superior and Inferior oDSGCs (2-sided rank-sum). **(p-s)**

Tuning curve metrics for spikes (abscissa) and simultaneously measured subthreshold voltages

(ordinate) under **(p-q)** depolarizing and **(r-s)** hyperpolarizing conditions: **(p, r)** area of the

normalized tuning curve, and **(q, s)** direction selectivity index. Significance values indicate

whether the data tend to fall unevenly on one side of the unity line (2-sided signed-rank). **(t-u)**

Residuals from the unity line for individual cells from the plots in **(t)** (j) and (n) and **(u)** (k) and (o).

Dashed line indicates unity (i.e. no difference across depolarizing and hyperpolarizing conditions).

(v-w) Residuals from the unity line for individual cells from the plots in **(v)** (p) and (r) and **(w)** (q)

and (s). Dashed line indicates unity (i.e., no difference between spikes and subthreshold

voltages). Comparisons in (t-w) are made between conditions (2-sided rank-sum). For box plots,

the blue line represents median, box boundaries are IQR, and whiskers represent most extreme

observation within 1.5xIQR. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. See also Fig S5.

a Conductances

Excitation
— all directions

Inhibition
— 0
— 45
— 90 (PD)
— 135
— 180
— 225
— 270
— 315

1nS
1s

b Membrane Potentials

Gain = 1

Threshold

1s

c Preferred vs Null Direction Responses

V_m Spikes
PD
ND

d Tuning Curve Area

e Normalized Area

I

f Direction Selectivity Index

I

Figure 6

Figure 6. A leaky integrate-and-fire model demonstrates how untuned excitation contributes to direction tuning. An exemplar oDSGC was modeled using parameters recorded directly from cells, including directionally tuned inhibitory conductances for each drifting bar direction, and a single, untuned excitatory conductance (see Methods). **(a)** Inhibitory (pastel colors) and excitatory (yellow) conductances of the model oDSGC in response to bars moving in 8 directions. **(b)** The leaky integrate-and-fire model uses the empirically measured parameters to model membrane potential across bar directions, shown here for the case in which the excitatory gain (i.e. a multiplication factor applied to the excitatory conductance) is set to 1.0. The red dotted line indicates the spike threshold. **(c)** Preferred (solid lines) and null (dotted lines) direction responses of subthreshold voltages (blue) and spikes (red) across a range of excitatory gains. Values are normalized to the maximum preferred direction response for each metric. The yellow column indicates the regime in which the null direction elicits zero spikes but the preferred direction has nonzero spikes. **(d-f)** Directional tuning properties as a function of excitatory gain for subthreshold voltages (blue) and spikes (red): **(d)** area of the linear tuning curve (normalized to the maximum area for each metric), **(e)** area of the normalized tuning curve, and **(f)** direction selectivity index. Orange dots [and error bars] in (e) and (f) correspond to the empirically measured median [and 95% confidence intervals determined via bootstrapping] for each metric from cell-attached recordings, collapsed across cell types.

Figure 7

Figure 7. Stimulus contrast modulates the spike tuning curves of oDSGCs. (a) Example cell-attached tuning curves from a Superior oDSGC at high (green) and low (tan) contrasts. **(b-d)** Metrics of **(b)** linear tuning curve area, **(c)** area of the normalized tuning curve, and **(d)** direction selectivity index from spike responses to high (abscissa) and low (ordinate) contrast bars moving in 8 directions. The region of tan [or green] indicates the metric is greater under low [or high] contrast conditions. Points on the line indicate equivalent metrics under the two conditions. Individual cells are represented by dots. Superior (magenta) and Inferior (gray) oDSGCs maintain their relative differences even under low contrast (see Fig 2). Red circles represent univariate medians (collapsed across cell type). Whiskers indicate 95% confidence intervals determined via bootstrapping. Significance values indicate whether the data tend to fall unevenly on one side of the unity line (2-sided signed-rank). **(e-g)** Metrics of **(e)** linear tuning curve area, **(f)** area of the normalized tuning curve, and **(g)** direction selectivity index from subthreshold voltages in response to high (abscissa) and low (ordinate) contrast bars moving in 8 directions. Blue circles represent univariate medians (collapsed across cell type). Whiskers indicate 95% confidence intervals determined via bootstrapping. **(h-i)** Residuals from the unity line for **(h)** area of the normalized tuning curve and **(i)** direction selectivity index under high and low contrast conditions for simultaneously measured spikes and subthreshold voltages. Dashed lines indicate unity (i.e. no difference between high and low contrast). Blue lines represent medians, box boundaries are IQR, and whiskers represent most extreme observation within 1.5xIQR. Comparisons are made between spike and Vm distributions (2-sided signed-rank). **(j-k)** E/I ratios for high and low contrast bars moving in **(j)** the preferred and **(k)** the null direction of each cell. **(l)** Direction selectivity index generated from a leaky integrate-and-fire model of an oDSGC under different scale factors applied jointly to excitatory and inhibitory inputs (constant E/I). *$p<0.05$, **$p<0.01$, ***$p<0.001$

Figure 8

Figure 8. Ganglion cell responses predict behavioral OKR gain across stimulus directions and contrasts. **(a)** Schematic of the putative computation between oDSGCs and OKR, consisting of a subtraction between Superior and Inferior oDSGC outputs and a nonlinearity. **(b)** Legend and summary for panels (c-r). Cellular Data (gray box): The responses of Superior (magenta) and Inferior (gray) oDSGCs in their preferred (PD) and null (ND) directions at high (left column) and low (right column) contrasts to superior (top row) and inferior (bottom row) drifting stimuli. Linear predictions (blue boxes): the differences between responses in the preferred and null directions, represented by Δ, provide predictions of the relative OKR gain for stimuli of different directions and contrasts. Behavioral results (orange boxes): arrows represent how vertical OKR gain changes with stimulus direction and contrast. **(c-f)** Cellular Data: Superior (magenta) and Inferior (gray) oDSGC spike rates for superior (top row) and inferior (bottom row) stimuli at high (left column) and low (right column) contrast. Brackets above show the difference between the median responses of the two populations of oDSGCs. **(g-h)** Linear predictions for how behavioral OKR gain changes across high (green) and low (tan) contrasts for **(g)** superior and **(h)** inferior stimuli. Predictions are calculated via iteratively bootstrapping the difference of medians from the distributions shown in (c-f). **(i-j)** Behavioral results of OKR gain for **(i)** superior and **(j)** inferior motion at high (green) and low (tan) contrast. **(k-l)** Linear predictions for how behavioral OKR gain changes between superior (magenta) and inferior (gray) stimulus directions at **(k)** high and **(l)** low contrast. **(m-n)** Behavioral results of OKR gain at **(m)** high and **(n)** low contrast for superior (magenta) and inferior (gray) stimuli (2-sided signed-rank). **(o)** Linear predictions for behavioral OKR gain at high (green) and low (tan) contrast, irrespective of stimulus direction. Gain is predicted to decrease with contrast. **(p)** Linear prediction for how the asymmetry between superior and inferior OKR changes with contrast. The asymmetry is predicted to be greater at high (green) compared to low (tan) contrast. **(q)** Behavioral results of OKR gain reveal greater gain at high (green) compared to low (tan) contrast, irrespective of stimulus direction. **(r)** Behavioral results of OKR gain reveal that the asymmetry between superior and inferior OKR is greater at high (green)

compared to low (tan) contrast. **(s)** Summary of the linear predictions for eye position across a

single stimulus oscillation cycle. Two second-order predictions indicate that (1) gain decreases

with contrast and (2) the superior/inferior asymmetry decreases with contrast. **(t)** Relationship

between the cellular predictions and the corresponding behavioral OKR gains, i.e. nonlinearity,

for superior (magenta points) and inferior (black points) stimuli. Points indicate univariate medians

and error bars show 95% confidence intervals determined via bootstrapping. Least-squares fit of

a Hill equation to the data (red line), including a maximum value of 1.0 and a fixed point at the

origin. **(u)** Summary of behavioral results showing the mean eye position across a single

oscillation cycle for high (green, same data as in Fig 1j) and low (tan) contrast oscillating gratings.

Stimulus position in lavender. Behavioral OKR shows higher gain and greater asymmetry at high

contrast. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$. See also Fig S6.

REFERENCES

1. Zeil, J., Nalbach, G. & Nalbach, H.-O. Spatial Vision in a Flat World: Optical and Neural Adaptations in Arthropods. in *Neurobiology of Sensory Systems* (eds. Singh, R. N. & Strausfeld, N. J.) 123–137 (Springer US, 1989). doi:10.1007/978-1-4899-2519-0_10.

2. Masseck, O. A. & Hoffmann, K.-P. Comparative Neurobiology of the Optokinetic Reflex. *Ann. N. Y. Acad. Sci.* **1164**, 430–439 (2009).

3. Simpson, J. I. The Accessory Optic System. *Annu. Rev. Neurosci.* **7**, 13–41 (1984).

4. Collewijn, H. Optokinetic eye movements in the rabbit: Input-output relations. *Vision Res.* **9**, 117–132 (1969).

5. Donaghy, M. The contrast sensitivity, spatial resolution and velocity tuning of the cat's optokinetic reflex. *J. Physiol.* **300**, 353–365 (1980).

6. Gravot, C. M., Knorr, A. G., Glasauer, S. & Straka, H. It's not all black and white: visual scene parameters influence optokinetic reflex performance in Xenopus laevis tadpoles. *J. Exp. Biol.* **220**, 4213–4224 (2017).

7. Knorr, A. G., Gravot, C. M., Glasauer, S. & Straka, H. Image motion with color contrast suffices to elicit an optokinetic reflex in Xenopus laevis tadpoles. *Sci. Rep.* **11**, 8445 (2021).

8. Leguire, L. E. *et al.* Contrast sensitivity of optokinetic nystagmus. *Vision Res.* **31**, 89–97 (1991).

9. Cahill, H. & Nathans, J. The Optokinetic Reflex as a Tool for Quantitative Analyses of Nervous System Function in Mice: Application to Genetic and Drug-Induced Variation. *PLOS ONE* **3**, e2055 (2008).

10. Dehmelt, F. A. *et al.* Spherical arena reveals optokinetic response tuning to stimulus location, size, and frequency across entire visual field of larval zebrafish. *eLife* **10**, e63355 (2021).

11. Shimizu, N. *et al.* Distribution of optokinetic sensitivity across the retina of mice in relation to

eye orientation. *Neuroscience* **168**, 200–208 (2010).

12. Dhande, O. S., Stafford, B. K., Lim, J.-H. A. & Huberman, A. D. Contributions of Retinal Ganglion Cells to Subcortical Visual Processing and Behaviors. *Annu. Rev. Vis. Sci.* **1**, 291–328 (2015).

13. Giolli, R. A., Blanks, R. H. I. & Lui, F. The accessory optic system: basic organization with an update on connectivity, neurochemistry, and function. in *Progress in Brain Research* (ed. Büttner-Ennever, J. A.) vol. 151 407–440 (Elsevier, 2006).

14. Oyster, C. W. The analysis of image motion by the rabbit retina. *J. Physiol.* **199**, 613–635 (1968).

15. Oyster, C. W. & Barlow, H. B. Direction-Selective Units in Rabbit Retina: Distribution of Preferred Directions. *Science* **155**, 841–842 (1967).

16. Sabbah, S. *et al.* A retinal code for motion along the gravitational and body axes. *Nature* **546**, 492–497 (2017).

17. Dhande, O. S. *et al.* Genetic Dissection of Retinal Inputs to Brainstem Nuclei Controlling Image Stabilization. *J. Neurosci.* **33**, 17797–17813 (2013).

18. Lilley, B. N. *et al.* Genetic access to neurons in the accessory optic system reveals a role for Sema6A in midbrain circuitry mediating motion perception. *J. Comp. Neurol.* **527**, 282–296 (2019).

19. Sun, L. O. *et al.* Functional Assembly of Accessory Optic System Circuitry Critical for Compensatory Eye Movements. *Neuron* **86**, 971–984 (2015).

20. Yonehara, K. *et al.* Expression of SPIG1 Reveals Development of a Retinal Ganglion Cell Subtype Projecting to the Medial Terminal Nucleus in the Mouse. *PLoS ONE* **3**, (2008).

21. Yonehara, K. *et al.* Identification of Retinal Ganglion Cells and Their Projections Involved in Central Transmission of Information about Upward and Downward Image Motion. *PLoS ONE* **4**, (2009).

22. Togt, C. van der, Want, J. van der & Schmidt, M. Segregation of direction selective neurons

and synaptic organization of inhibitory intranuclear connections in the medial terminal

nucleus of the rat: An electrophysiological and immunoelectron microscopical study. *J.*

Comp. Neurol. **338**, 175–192 (1993).

23. Kay, J. N. *et al.* Retinal Ganglion Cells with Distinct Directional Preferences Differ in

    Molecular Identity, Structure, and Central Projections. *J. Neurosci.* **31**, 7753–7762 (2011).

24. Osterhout, J. A., Stafford, B. K., Nguyen, P. L., Yoshihara, Y. & Huberman, A. D. Contactin-

    4 Mediates Axon-Target Specificity and Functional Development of the Accessory Optic

    System. *Neuron* **86**, 985–999 (2015).

25. Pak, M. W. *et al.* Retinopretectal and accessory optic projections of normal mice and the

    OKN-defective mutant mice beige, beige-J, and pearl. *J. Comp. Neurol.* **258**, 435–446

    (1987).

26. Oyster, C. W., Takahashi, E. & Collewijn, H. Direction-selective retinal ganglion cells and

    control of optokinetic nystagmus in the rabbit. *Vision Res.* **12**, 183–193 (1972).

27. Wei, W. Neural Mechanisms of Motion Processing in the Mammalian Retina. *Annu. Rev.*

    Vis. Sci. **4**, 165–192 (2018).

28. Takahashi, M., Sakurai, S. & Kanzaki, J. Horizontal and Vertical Optokinetic Nystagmus in

    Man. *ORL J Otorhinolaryngol Relat Spec* **40**, 43–52 (1978).

29. Hainline, L., Lemerise, E., Abramov, I. & Turkel, J. Orientational asymmetries in small-field

    optokinetic nystagmus in human infants. *Behav. Brain Res.* **13**, 217–230 (1984).

30. Murasugi, C. M. & Howard, I. P. Up-down asymmetry in human vertical optokinetic

    nystagmus and afternystagmus: contributions of the central and peripheral retinae. *Exp.*

    Brain Res. **77**, 183–192 (1989).

31. van den Berg, A. V. & Collewijn, H. Directional asymmetries of human optokinetic

    nystagmus. *Exp. Brain Res.* **70**, 597–604 (1988).

32. Knapp, C. M., Proudlock, F. A. & Gottlob, I. OKN Asymmetry in Human Subjects: A

    Literature Review. *Strabismus* **21**, 37–49 (2013).

33. Hoffmann, K.-P. & Fischer, W. H. Directional effect of inactivation of the nucleus of the optic tract on optokinetic nystagmus in the cat. *Vision Res.* **41**, 3389–3398 (2001).

34. Evinger, C. & Fuchs, A. F. Saccadic, smooth pursuit, and optokinetic eye movements of the trained cat. *J. Physiol.* **285**, 209–229 (1978).

35. Grasse, K. L. & Cynader, M. S. The effect of visual cortex lesions on vertical optokinetic nystagmus in the cat. *Brain Res.* **455**, 385–389 (1988).

36. Wallman, J. & Velez, J. Directional asymmetries of optokinetic nystagmus: developmental changes and relation to the accessory optic system and to the vestibular system. *J. Neurosci.* **5**, 317–329 (1985).

37. Takahashi, M. & Igarashi, M. Comparison of Vertical and Horizontal Optokinetic Nystagmus in the Squirrel Monkey. *ORL* **39**, 321–329 (1977).

38. Denman, D. J., Siegle, J. H., Koch, C., Reid, R. C. & Blanche, T. J. Spatial Organization of Chromatic Pathways in the Mouse Dorsal Lateral Geniculate Nucleus. *J. Neurosci.* **37**, 1102–1116 (2017).

39. Yonehara, K. *et al.* Congenital Nystagmus Gene FRMD7 Is Necessary for Establishing a Neuronal Circuit Asymmetry for Direction Selectivity. *Neuron* **89**, 177–193 (2016).

40. Briggman, K. L., Helmstaedter, M. & Denk, W. Wiring specificity in the direction-selectivity circuit of the retina. *Nature* **471**, 183–188 (2011).

41. Fried, S. I., Münch, T. A. & Werblin, F. S. Mechanisms and circuitry underlying directional selectivity in the retina. *Nature* **420**, 411–414 (2002).

42. Wei, W., Hamby, A. M., Zhou, K. & Feller, M. B. Development of asymmetric inhibition underlying direction selectivity in the retina. *Nature* **469**, 402–406 (2011).

43. Ding, H., Smith, R. G., Poleg-Polsky, A., Diamond, J. S. & Briggman, K. L. Species-specific wiring for direction selectivity in the mammalian retina. *Nature* **535**, 105–110 (2016).

44. Amthor, F. R., Keyser, K. T. & Dmitrieva, N. A. Effects of the destruction of starburst-cholinergic amacrine cells by the toxin AF64A on rabbit retinal directional selectivity. *Vis.*

Neurosci. **19**, 495–509 (2002).

45. Yonehara, K. *et al.* Spatially asymmetric reorganization of inhibition establishes a motion-sensitive circuit. *Nature* **469**, 407–410 (2011).

46. Yoshida, K. *et al.* A Key Role of Starburst Amacrine Cells in Originating Retinal Directional Selectivity and Optokinetic Eye Movement. *Neuron* **30**, 771–780 (2001).

47. Mauss, A. S., Vlasits, A., Borst, A. & Feller, M. Visual Circuits for Direction Selectivity. *Annu. Rev. Neurosci.* **40**, 211–230 (2017).

48. Vaney, D. I., Sivyer, B. & Taylor, W. R. Direction selectivity in the retina: symmetry and asymmetry in structure and function. *Nat. Rev. Neurosci.* **13**, 194–208 (2012).

49. Mani, A. *et al. A retinal circuit that vetoes optokinetic responses to fast visual motion*. http://biorxiv.org/lookup/doi/10.1101/2021.10.31.466688 (2021) doi:10.1101/2021.10.31.466688.

50. Matsumoto, A., Briggman, K. L. & Yonehara, K. Spatiotemporally Asymmetric Excitation Supports Mammalian Retinal Motion Sensitivity. *Curr. Biol.* **29**, 3277-3288.e5 (2019).

51. Koulen, P., Fletcher, E. L., Craven, S. E., Bredt, D. S. & Wässle, H. Immunocytochemical Localization of the Postsynaptic Density Protein PSD-95 in the Mammalian Retina. *J. Neurosci.* **18**, 10136–10149 (1998).

52. Sassoè-Pognetto, M. *et al.* Colocalization of gephyrin and GABAA-receptor subunits in the rat retina. *J. Comp. Neurol.* **357**, 1–14 (1995).

53. Sassoè-Pognetto, M. & Wässle, H. Synaptogenesis in the rat retina: subcellular localization of glycine receptors, GABAA receptors, and the anchoring protein gephyrin. *J. Comp. Neurol.* **381**, 158–174 (1997).

54. Matsumoto, A. *et al.* Direction selectivity in retinal bipolar cell axon terminals. *Neuron* **0**, (2021).

55. Percival, K. A., Venkataramani, S., Smith, R. G. & Taylor, W. R. Directional excitatory input to direction-selective ganglion cells in the rabbit retina. *J. Comp. Neurol.* **527**, 270–281

(2019).

56. Poleg-Polsky, A. & Diamond, J. S. Imperfect Space Clamp Permits Electrotonic Interactions between Inhibitory and Excitatory Synaptic Conductances, Distorting Voltage Clamp Recordings. *PLOS ONE* **6**, e19463 (2011).

57. Summers, M. T. & Feller, M. B. Distinct inhibitory pathways control velocity and directional tuning in the mouse retina. *Curr. Biol.* **0**, (2022).

58. Yonehara, K. *et al.* The First Stage of Cardinal Direction Selectivity Is Localized to the Dendrites of Retinal Ganglion Cells. *Neuron* **79**, 1078–1085 (2013).

59. Oesch, N., Euler, T. & Taylor, W. R. Direction-Selective Dendritic Action Potentials in Rabbit Retina. *Neuron* **47**, 739–750 (2005).

60. Poleg-Polsky, A. & Diamond, J. S. NMDA Receptors Multiplicatively Scale Visual Signals and Enhance Directional Motion Discrimination in Retinal Ganglion Cells. *Neuron* **89**, 1277–1290 (2016).

61. Poleg-Polsky, A. & Diamond, J. S. Retinal Circuitry Balances Contrast Tuning of Excitation and Inhibition to Enable Reliable Computation of Direction Selectivity. *J. Neurosci.* **36**, 5861–5876 (2016).

62. Sethuramanujam, S. *et al.* A Central Role for Mixed Acetylcholine/GABA Transmission in Direction Coding in the Retina. *Neuron* **90**, 1243–1256 (2016).

63. Sethuramanujam, S. *et al.* "Silent" NMDA Synapses Enhance Motion Sensitivity in a Mature Retinal Circuit. *Neuron* **96**, 1099-1111.e3 (2017).

64. Soodak, R. E. & Simpson, J. I. The accessory optic system of rabbit. I. Basic visual response properties. *J. Neurophysiol.* **60**, 2037–2054 (1988).

65. Giolli, R. A. *et al.* GABAergic neurons comprise a major cell type in rodent visual relay nuclei: an immunocytochemical study of pretectal and accessory optic nuclei. *Exp. Brain Res.* **61**, 194–203 (1985).

66. Grasse, K. L. & Cynader, M. S. Electrophysiology of medial terminal nucleus of accessory

optic system in the cat. *J. Neurophysiol.* **48**, 490–504 (1982).

67. Natal, C. L. & Britto, L. R. G. The rat accessory optic system: effects of cortical lesions on the directional selectivity of units within the medial terminal nucleus. *Neurosci. Lett.* **91**, 154–159 (1988).

68. Simpson, J. I., Soodak, R. E. & Hess, R. The Accessory Optic System and its Relation to the Vestibulocerebellum. in *Progress in Brain Research* (eds. Granit, R. & Pompeiano, O.) vol. 50 715–724 (Elsevier, 1979).

69. Mowrer, O. H. A comparison of the reaction mechanisms mediating optokinetic nystagmus in human beings and in pigeons. *Psychol. Monogr.* **47**, 294–305 (1936).

70. Liu, B., Huberman, A. D. & Scanziani, M. Cortico-fugal output from visual cortex promotes plasticity of innate motor behaviour. *Nature* **538**, 383–387 (2016).

71. Wood, C. C., Spear, P. D. & Braun, J. J. Direction-specific deficits in horizontal optokinetic nystagmus following removal of visual cortex in the cat. *Brain Res.* **60**, 231–237 (1973).

72. Cook, J. E. & Podugolnikova, T. A. Evidence for spatial regularity among retinal ganglion cells that project to the accessory optic system in a frog, a reptile, a bird, and a mammal. *Vis. Neurosci.* **18**, 289–297 (2001).

73. Dann, J. F. & Buhl, E. H. Retinal ganglion cells projecting to the accessory optic system in the rat. *J. Comp. Neurol.* **262**, 141–158 (1987).

74. Ruff, T. *et al.* FLRT3 Marks Direction-Selective Retinal Ganglion Cells That Project to the Medial Terminal Nucleus. *Front. Mol. Neurosci.* **14**, (2021).

75. Tauber, E. S. & Atkin, A. Optomotor Responses to Monocular Stimulation: Relation to Visual System Organization. *Science* **160**, 1365–1367 (1968).

76. Holmgren, C. D. *et al.* Visual pursuit behavior in mice maintains the pursued prey on the retinal region with least optic flow. *eLife* **10**, e70838 (2021).

77. Lee, S. *et al.* An unconventional glutamatergic circuit in the retina formed by vGluT3 amacrine cells. *Neuron* **84**, 708–715 (2014).

78. Sivyer, B., Tomlinson, A. & Taylor, W. R. Simulated Saccadic Stimuli Suppress ON-Type Direction-Selective Retinal Ganglion Cells via Glycinergic Inhibition. *J. Neurosci.* **39**, 4312–4322 (2019).

79. Kittila, C. A. & Massey, S. C. Pharmacology of Directionally Selective Ganglion Cells in the Rabbit Retina. *J. Neurophysiol.* **77**, 675–689 (1997).

80. Carandini, M. & Heeger, D. J. Normalization as a canonical neural computation. *Nat. Rev. Neurosci.* **13**, 51–62 (2012).

81. Barlow, H. B. Summation and inhibition in the frog's retina. *J. Physiol.* **119**, 69–88 (1953).

82. Hartline, H. K., Wagner, H. G. & MacNichol, E. F. The Peripheral Origin of Nervous Activity in the Visual System. *Cold Spring Harb. Symp. Quant. Biol.* **17**, 125–141 (1952).

83. Kuffler, S. W. Discharge patterns and functional organization of mammalian retina. *J. Neurophysiol.* **16**, 37–68 (1953).

84. Dacey, D. M., Crook, J. D. & Packer, O. S. Distinct synaptic mechanisms create parallel S-ON and S-OFF color opponent pathways in the primate retina. *Vis. Neurosci.* **31**, 139–151 (2014).

85. De Monasterio, F. M. & Gouras, P. Functional properties of ganglion cells of the rhesus monkey retina. *J. Physiol.* **251**, 167–195 (1975).

86. Field, G. D. *et al.* High-sensitivity rod photoreceptor input to the blue-yellow color opponent pathway in macaque retina. *Nat. Neurosci.* **12**, 1159–1164 (2009).

87. Suver, M. P. *et al.* Encoding of Wind Direction by Central Neurons in Drosophila. *Neuron* **102**, 828-842.e7 (2019).

88. Martin, H. Osmotropotaxis in the Honey-Bee. *Nature* **208**, 59–63 (1965).

89. Schwenk, K. Why snakes have forked tongues. *Science* **263**, 1573 (1994).

90. Middlebrooks, J. C. & Green, D. M. Sound Localization by Human Listeners. *Annu. Rev. Psychol.* **42**, 135–159 (1991).

91. Stahl, J. S., van Alphen, A. M. & De Zeeuw, C. I. A comparison of video and magnetic

search coil recordings of mouse eye movements. *J. Neurosci. Methods* **99**, 101–110 (2000).

92. Stahl, J. S. Eye Movements of the Murine P/Q Calcium Channel Mutant Rocker, and the Impact of Aging. *J. Neurophysiol.* **91**, 2066–2078 (2004).

93. Zoccolan, D., Graham, B. J. & Cox, D. D. A Self-Calibrating, Camera-Based Eye Tracker for the Recording of Rodent Eye Movements. *Front. Neurosci.* **4**, 193 (2010).

94. Rodieck, R. W. The density recovery profile: A method for the analysis of points in the plane applicable to retinal studies. *Vis. Neurosci.* **6**, 95–111 (1991).

95. Care, R. A. *et al.* Partial Cone Loss Triggers Synapse-Specific Remodeling and Spatial Receptive Field Rearrangements in a Mature Retinal Circuit. *Cell Rep.* **27**, 2171-2183.e5 (2019).

96. Care, R. A. *et al.* Mature Retina Compensates Functionally for Partial Loss of Rod Photoreceptors. *Cell Rep.* **31**, (2020).

97. Remtulla, S. & Hallett, P. E. A schematic eye for the mouse, and comparisons with the rat. *Vision Res.* **25**, 21–31 (1985).

98. Antoine, M. W., Langberg, T., Schnepel, P. & Feldman, D. E. Increased Excitation-Inhibition Ratio Stabilizes Synapse and Circuit Excitability in Four Autism Mouse Models. *Neuron* **101**, 648-661.e4 (2019).