1 **Title: Preclinical assessment of Fibroblast Activation Protein-specific MRI as an alternative**
2 **to PSMA for image-guided focal therapy of prostate cancer**
3

4 Nicole Dmochowska[a*] PhD, Valentina Milanova[a*] PhD, Ramesh Mukkamala[b] PhD, Nguyen T.H.
5 Pham[c] PhD, Madduri Srinivasarao[b] PhD, Lisa M. Ebert PhD[d] Timothy Stait-Gardner[e] PhD, Hien
6 Le[f] MBBS, Anil Shetty[g] MD, Philip S. Low[b] PhD, Benjamin Thierry [a] PhD
7
[a]Future Industries Institute, University of South Australia, Adelaide, South Australia, Australia
[b]Department of Chemistry and Institute for Drug Discovery, Purdue University, West
Lafayette, Indiana, United States
[c]Key Centre for Polymers and Colloids, School of Chemistry, The University of Sydney, Sydney,
NSW, Australia
[d] Centre for Cancer Biology, University of South Australia and SA Pathology, Adelaide, South
Australia, Australia
[e]Nanoscale Organisation and Dynamics Group, Western Sydney University, Sydney, New
South Wales, Australia
[f]Department of Radiation Oncology, Royal Adelaide Hospital, Adelaide, SA, Australia
[g]Ferronova Pty Ltd, Mawson Lakes, SA, Australia
19
20 Corresponding author: Professor Benjamin Thierry, Bioengineering Lab, Future Industries
21 Institute, MM Building, University of South Australia Mawson Lakes Campus, University
22 Boulevard, Mawson Lakes, South Australia 5095, Australia, email:
23 Benjamin.thierry@unisa.edu.au, telephone +618 830 23689.

24
25 **Abstract**
26

Accurate and precise delineation of gross tumour volumes remains a barrier to radiotherapy
dose escalation and boost dosing in the treatment of solid tumours, such as prostate cancer.
Magnetic resonance imaging of tumour molecular targets has the power to enable focal dose
boosting, particularly when combined with technological advances such as MRI-LINAC.
Fibroblast activation protein (FAP) is a transmembrane protein overexpressed in stromal
components of >90% of epithelial carcinomas. Herein we compare targeted MRI of gold
standard PSMA with FAP in the delineation of orthotopic tumours in a mouse model of
prostate cancer. Control (no ligand), FAP and PSMA-targeting iron oxide nanoparticles were
prepared with modification of an MRI agent (FerroTrace). Mice with orthotopic LNCaP
tumours underwent T_2-weighted 3D MRI 24 hours after intravenous injection of contrast
agents. FAP and PSMA nanoparticles produced contrast enhancement on MRI when
compared to control nanoparticles, which was most pronounced on the tumour periphery.
FAP-targeted MRI increased tumour contrast enhancement by ~15% when compared to
PSMA. This study demonstrates preclinical feasibility of PSMA and FAP-targeted MRI which
be used to enable targeted image-guided focal therapy of localized prostate cancer.

Background

Prostate cancer (PCa) is one of the most diagnosed malignancies in men (1) and, owing to its relatively high survival rate and frequent full gland treatment associated morbidity, is the leading cause of treatment–related years lived with disability worldwide. Along with radical prostatectomy, radiotherapy (RT) is a mainstay treatment option for all stages of disease, and is commonly delivered as a uniform dose over the entire prostate gland (2). There is mounting evidence that the risk of biochemical disease recurrence is reduced with dose escalation when greater than 80 Gy, especially in patients with intermediate or high-risk lesions. However, dose intensification of the whole gland is limited by off-target toxicities to adjacent organs at risk including the rectum, urethra, and bladder (3). Following pelvic radiotherapy, up to 30% of patients experience late grade III or IV genitourinary or gastrointestinal toxicities, with persistent long-term negative impacts on quality of life, through both mental and physical wellbeing, largely due to bowel disturbances(4, 5). Although modern RT techniques enable safer and more precise dose delivery to the prostate, treatment-associated morbidity remains a significant concern in the treatment of PCa patients.

PCa is a multifocal disease, often presenting with large primary intraprostatic lesions accompanied by smaller, well defined secondary lesions. Dominant intraprostatic lesions (index lesion) are considered to be the most aggressive, responsible for driving disease progression and prognosis, and are the most common sites of recurrence (6-8). Focal dose escalation RT approaches specifically target index lesions, balancing the risk of recurrence and patient's quality of life. For example, the recently reported FLAME study demonstrated that focal dose escalation under the guidance of multiparametric MRI (mpMRI) could improve biochemical recurrence free survival by 7%, without impacting morbidity and quality of life (9). Early results from a phase II trial investigating MRI-guided focused ultrasound ablation for localized intermediate risk PCa also demonstrated encouraging oncologic and functional outcomes with 93% of patients disease-free at 5 months after treatment (10). In a cohort of low to intermediate risk PCa, gadolinium enhanced MRI-guided focal laser ablation showed promising early oncologic results with 17% of participants requiring additional oncologic treatment after one year of initial treatment, with no significant side effects or impact to quality of life (11).

Target delineation, a key requirement to define gross tumour volumes (GTV) in intraprostatic focal treatment approaches, is commonly referred to as the Achilles' heel of RT. Owing to its superior soft-tissue resolution compared to computed tomography and the ability to acquire non-invasive functional imaging, mpMRI is rapidly becoming a staple tool in guiding PCa RT and more generally focal ablation. Adoption is supported by powerful technological advances such as the integration of MRI with linear accelerators (MRI-LINAC) and MRI-ultrasound (US) fusion set-ups that streamline intraoperative US image guidance with the resolution of MRI. However, mpMRI only has a moderate inter-rater agreement and is prone to a large false-positive rate and underestimation of tumour margins (12, 13). In addition, target delineation with mpMRI is not trivial, which substantially lengthens MRI-guided RT procedures and may

result in greater intrafraction motion. This is further limited by magnet strength and
experience of the imaging team.

A recent study demonstrated increased consensus of prostate specific membrane antigen
(PSMA)-PET volumes with histology compared with mpMRI for the delineation of
intraprostatic GTVs, indicating that imaging with molecular targets increases imaging
specificity (14). Furthermore, GTVs derived from mpMRI significantly underestimated true
tumour volumes compared to PSMA-PET. For intraprostatic radiotherapy boosting, imaging
of molecular targets such as PSMA may therefore advantageously replace or be used in
addition to conventional mpMRI and provides additional biological characterisation of
malignant tissue for dose planning. However, PSMA-PET can intrinsically lead to GTV
underestimation due to various technical aspects such as partial volume effects limiting image
resolution (15, 16). In addition, PET-based target delineation suffers from the absence of
integrated systems such as MRI-LINAC and MRI-US fusion. Conceptually, molecular MRI
should provide improved imaging of tumoural features on a biological level over what is to
date conventionally afforded by PET imaging. However, despite very extensive research in
molecular MRI, especially with nanoparticulate-based contrast agents, clinical translation is
yet to be demonstrated. On the other hand, most of the related research has to date focussed
on diagnostic applications as opposed to the feasibility of accurately imaging and delineating
primary lesions.

Besides the imaging modality itself, a key requirement for molecular MRI is the selection of a
biological target specific to the tumour. PSMA is often significantly overexpressed in PCa and
as such has become the gold-standard target for molecular imaging of PCa (17). Multiple
PSMA ligands are now available for PET imaging with various FDA and EMA-approved PSMA-
targeting probes demonstrating success in the clinic. However, loss of PSMA expression has
been known to occur in neuroendocrine, androgen receptor-negative PCa, reducing the
efficacy of PSMA-targeting probes in these cases. Furthermore, 5-10% of primary PCa or
primary PCa lesions are PSMA negative on PET (18, 19). Therefore, imaging of PSMA is not
suitable for all patients, warranting the validation of additional imaging targets in PCa (20).

Fibroblast activation protein (FAP) is an extracellular serine protease that is upregulated in
stromal components of >90% of epithelial tumours but with low or undetectable expression
in normal tissues. In addition, FAP expression correlates with progression and worse
prognosis (21, 22). Endothelial cells can also express FAP which contributes to the regulation
of tumour angiogenesis, changes in capillary morphology and microvascular reorganization
(23). Specifically, FAP has been found to be expressed by tumour endothelial cells during
capillary formation but is absent in the mature endothelium (24). Additionally, FAP expression
correlates with a significantly increased density of microvessels (25). As such, FAP is
potentially a pan-cancer marker and a promising target for molecular imaging and therapy.
FAP has been shown to be upregulated in PCa, and is predominantly expressed in cancer
associated fibroblasts and reactive stromal cells adjacent to carcinoma cells (26-28). Recent
studies demonstrated that in a population of patients with PSMA negative tumours,
intermediate to high uptake of a [68]Ga-FAP inhibitor was observed in PCa (29, 30).

Furthermore, FAP-specific PET imaging of malignancies with low PSMA expression
demonstrated more precise results when compared to PSMA-PET (31). Imaging of FAP in PCa
has shown promise particularly in castrate resistant prostate cancer (CRPC), with elevated
FAP expression also observed in neuroendocrine prostate cancer (NEPC) and patients
undergoing neoadjuvant androgen deprivation therapy (ADT) (32).

To date, the utility and significance of FAP specific MRI is yet to be assessed. To address this
question, here we report on benchmarking the performance of a FAP-targeted iron oxide
nanoparticulate MRI agent against PSMA in the delineation of PCa in an orthotopic mouse
model. Our experimental approach builds on a previously reported FAP-ligand designed by
ligating fragments of existing FAP ligands that were identified by molecular docking studies
to contribute most prominently to the specificity and affinity of FAP binding. The FAP and
PSMA ligands were substituted from mannose in the FerroTrace maghemite nanoparticulate
MRI agent that is presently undergoing clinical trialling as a lymphotropic contrast agent.
Using this novel imaging platform, we demonstrate that FAP MRI yields superior tumour
contrast and coverage compared to PSMA in a preclinical model of PCa.

Materials

The following chemicals were purchased and used as received without further purification.
Trifluoroacetic acid, 1-Ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride
(EDC.HCl), N-hydroxysuccinimide (NHS), 4-Morpholineethanesulfonic acid, 2-(N-
Morpholino)ethanesulfonic acid hydrate (MES hydrate), 10X Phosphate Buffered Saline were
purchased from Sigma Aldrich. Acetone and dichloromethane (DCM) we purchased from
Chem-Supply. FAP binding ligand was synthesized as previously described (33). PSMA ligand
Clu-CO-Lys-(tBu)$_3$ ester (CAS 1025796-31-9) was synthesized by Advanced Molecular
Technologies (Scoresby, Victoria, Australia).

Analysis of FAP expression in PCa

To evaluate FAP expression in normal prostates and prostate adenocarcinoma, normalized
FAP RNAseq (expected count-deseq2+1) data was obtained from the Therapeutically
Applicable Research to Generate Effect Treatments initiative (TARGET)/the Cancer Genome
Atlas (TCGA)/Genotype Tissue Expression (GTEx) dataset using the UCSC Xena platform
(accessed 26/11/2021) (34). Corresponding patient prostate adenocarcinoma pathological T
staging was obtained from the TCGA prostate adenocarcinoma (PRAD) data set (accessed
26/11/2021). Furthermore, immunohistochemistry staining of FAP in low- and high-grade
prostate adenocarcinoma was obtained from the HPA059739 dataset deposited in the Human
Protein Atlas (https://www.proteinatlas.org/ENSG00000078098-
FAP/pathology/prostate+cancer#) (35).

Magnetic nanoparticle synthesis

The core maghemite iron oxide (γ-Fe$_2$O$_3$) nanoparticles were synthesised via coprecipitation
of iron salts in aqueous solutions and subsequent controlled oxidation using iron nitrate as
previously described (36). To achieve *in vivo* stability, the nanoparticles are colloidally
stabilised by block copolymers synthesised using reversible addition-fragment chain-transfer
(RAFT) polymerisation. The maghemite nanoparticles and RAFT block copolymer coating have
been previously developed for the preparation of a lymphotropic MRI agent (FerroTrace,
Ferronova) currently undergoing clinical trialling (36). Briefly, two types of block copolymer
are attached to the nanoparticle surface: a stabilising-polymer (RAFT-5-MAPC2-15AAM-
3PEO) and a targeting polymer with terminal FAP or PSMA ligands (RAFT-5MAPC2-70AAM-
PSMA/FAP) in a mole ratio of 70% stabilising to 30% targeting.

Preparation of RAFT-5MAPC2-15AAM-3PEO

Polymers were prepared based on a previously published procedure with modifications (37).
Methoxy triethylene glycol modified 2-{[butylsulfanyl)cabonothioyl]sulfanyl}propanoic acid
(1.0 g), acrylamide (2.8 g), 4,4'-azobis(4-cyanovaleric acid) (0.050 g), dioxane (10 g) and water
(10 g) were combined and dissolved before purging with nitrogen gas for 15 minutes and
polymerising at 70°C for 2 hours. The mixture was allowed to cool, opened to air, then
(methacryloyloxy)-ethyl]phosphonic acid (2.5 g) and 4,4'-azobis(4-cyanovaleric acid) (0.050 g)
were added to the reaction mixture. The mixture was purged with nitrogen gas for 15
minutes, then heated at 70°C for 4 hours. The polymer was precipitated in acetone, repurified
and collected by centrifugation.

Preparation of RAFT-5MAPC2-70AAM-PSMA/FAP polymers

RAFT-COOH (2-(((butylthio)carbonothioyl)-thio)-propanoic acid) (0.2 g), 4,4'-azobis(4-
cyanovaleric acid) (0.012 g), acrylamide (4.17 g), dioxane (7.8 g) and water (12.1 g) were
combined and dissolved. The reaction was magnetically stirred at 70°C for 4 hours in an inert
atmosphere. The mixture was allowed to cool, opened to air and [2-(methacryloyloxy)-
ethyl]phosphonic acid (0.81 g) and 4,4'-azobis(4-cyanovaleric acid) (0.012 g) were added to
the reaction mixture. The reaction was purged with nitrogen gas for 15 minutes before
heating to 70°C for 4 hours under magnetic stirring. At this step the RAFT-5MAPC2-70AAM
polymer was precipitated in acetone, repurified, collected by centrifugation and stored prior
to conjugation with the FAP and PSMA ligands.

For preparing RAFT-5MAPC2-70AAM-PSMA, the tert-butyl ester groups of the PSMA ligand
[Clu-CO-Lys-(tBu)$_3$ ester] were first removed using 20% trifluoroacetic acid (TFA) in
dichloromethane (DCM). Briefly, 20 mg/mL PSMA in 20% TFA in DCM was mixed for 3 hours
at room temperature before vacuum drying. The residue was then dissolved in 10-20%
aqueous acetic acid, transferred into a 50 mL falcon tube and a two-fold volume of chloroform
was added. The reaction was mixed thoroughly before the layers were allowed to separate
and the bottom organic layer containing the protective groups and non-volatile by-products
was removed using a needle and syringe. The extraction of the aqueous layer was repeated

twice, and it was dried under high vacuum. The residue was dissolved in glacial acetic acid,
shell-freezed and lyophilized before use. The NH$_2$ derivatized FAP ligand was synthesized as
previously described and used for the preparation of the RAFT-5MAPC2-70AAM-FAP (31).

PSMA and FAP ligands were conjugated to the RAFT-5MAPC2-70AAM polymer using a
carbodiimide coupling reaction. First, 200 mg of poly[[(methacryloyloxy)-ethyl]phosphonic
acid]-*block*-poly(acrylamide), EDC.HCl (48 mg) and NHS (12 mg) were dissolved in 10 mL MES
buffer (pH 5.0 – 5.5). The solution was mixed in a sonic bath for 10 minutes and the activated
polymer was collected by precipitation in acetone followed by centrifugation. Second,
FAP/PSMA ligands (12 mg) were dissolved in 100 µL DMSO and diluted to a total volume of
10 mL with 10X PBS buffer prior to addition to the activated polymer. The reaction mixture
was stirred for 20 hours. The conjugated polymer was purified using centrifugal filters with a
3 kDa molecular weight cut off membrane. The product was diluted to a final concentration
of 50 mg/mL and stored at 4°C until further use.

217 Preparation of PSMA and FAP nanoparticles

17 mg of RAFT-5MAPC2-15AAM-3PEO and 20 mg of RAFT-5MAPC2-70AAM-PSMA/FAP
polymers were dissolved in 2 mL of water and the pH was adjusted to 4 using 0.1M NaOH.
The maghemite nanoparticles (45 mg as dried weight) were added to the polymer solution
under probe sonication. After mixing for 10 minutes, the pH was adjusted to 5.0 with NaOH
(0.1M). Sonication was continued for a total of 30 minutes and the pH was further adjusted
stepwise to 6.0 and 7.0. The nanoparticle suspension was purified using a centrifugal filter
with a 100 kDa molecular weight cut off membrane. Lastly, the nanoparticles were diluted
with 3% saline to obtain a final concentration of 30 mg Fe/mL in 0.9% saline. An illustration
of the PSMA/FAP nanoparticle preparation is shown in figure 1A. For the preparation of non-
targeted nanoparticles the method described above was repeated except RAFT-5MAPC2-
70AAM-PSMA/FAP was replaced by RAFT-5MAPC2-70AAM-COOH.

230 Nanoparticle characterisation

The morphology of the coated nanoparticles was studied using transmission electron
microscope (TEM; JEOL JEM-2100F-HR) equipped with a field emission gun operated at 200
kV. Images were recorded with a CCD Camera (Gatan Orius SC1000). Samples were prepared
by placing a small drop of sample suspension in water onto a 200-mesh carbon-coated copper
grid (ProSciTech) and subsequently evaporating the water in air.

Dynamic Light Scattering (DLS) data was recorded on a Malvern Instruments Zetasizer Nano
ZS at 25°C. The hydrodynamic size of all nanoparticles suspensions was measured at a
concentration of 0.1 mg Fe/mL and isotonic saline (0.9%) was used as suspending medium.
Measurement duration was chosen as automatic.

Surface zeta potential data was acquired using the same instrumentation. The Smoluchowski
method was used for the zeta potential measurements. The surface zeta potential of all

nanoparticle suspensions was measured at a concentration of 1 mg Fe/mL and 10mM saline
was used as the suspending medium.

Cancer cell lines and culture conditions
Cells were routinely cultured in RPMI-1640 medium (Gibco) supplemented with 10% fetal
bovine serum (FBS) (Gibco) and 1% penicillin/streptomycin (Gibco) in a humidified
atmosphere of 5% CO_2 at 37°C. The medium was replaced every 2-3 days and cells were split
once they reached 75-80% confluence using TrypLE Express (Gibco).

Flow cytometric analysis of FAP expression in U87 and C32 cells

Cells were harvested from flasks using TrypLE Select (Thermo Fisher), washed in PBS and
resuspended in FACS buffer (PBS + 1% bovine serum albumin (BSA) + 0.04% sodium azide).
Staining for FAP was performed using either a one- or two-step process. For one-step staining,
90 µl of cells were incubated with 10 µl of anti-FAP-PE (clone 427819; R&D Systems) for 20
minutes at room temperature before washing with FACS buffer. For two-step staining, cells
were incubated with mouse anti-FAP (clone 427819; R&D Systems) at a final concentration of
5µg/ml for 20 mininutes at room temperature and washed with FACS buffer, then incubated
with goat anti-mouse secondary antibody conjugated to AlexaFluor488 (Thermo Fisher #A-
11029) at 10 µg/ml for 20 minutes at room temperature, then washed again in FACS buffer.
Samples were acquired on an LSR Fortessa or Accuri C6 flow cytometer (both BD Biosciences).
Analysis was performed using FCS Express V7 Flow Research Edition (De Novo Software,
Pasadena, CA, USA).

Magnetic nanoparticle *in vitro* binding assay
To test the binding affinity of the PSMA/FAP nanoparticles, cellular uptake in cells expressing
the respective targets was measured *in vitro*. LNCaP cells were seeded in a T25 cell culture
flask at a density of 1 x 10^6 cells per flask. C32 cells were seeded in a 6 well plate at a density
of 3.5 x 10^5 cells per well in the complete cell culture medium as described above. The cells
were placed in a 37°C, 5% CO_2 incubator and allowed to adhere for 24 hours. PSMA, FAP and
control unconjugated nanoparticle suspensions were prepared at a concentration of 0.150
mg Fe/mL in cell culture medium. 6 mL of PSMA or control nanoparticle suspension was added
to each flask containing adherent LNCaP cells or 1 mL of FAP and control nanoparticle
suspension was added to each well containing adherent C32 cells. Three replicates were used
to test each of the nanoparticle formulations. Cells were incubated with the nanoparticle
suspensions for 24 hours in an incubator at 37°C in 5% CO_2 humidified atmosphere. After
incubation, cells were washed 3 times with PBS, detached using TrypLE Express and cell
pellets were collected via centrifugation at 500 g for 5 minutes. The cell pellets were washed
twice with PBS and were dried at 60°C overnight using a heating block. The dried cell pellets
were digested with trace metal grade nitric and hydrochloric acid (1:1 ratio by volume).

Digested samples were diluted with water to a total volume of 10 mL or 3 mL for LNCaP PSMA
nanoparticles and C32 FAP nanoparticle samples, respectively. Iron concentration was
measured by inductively coupled plasma mass spectroscopy (ICP-MS).

Preclinical study in an orthotopic prostate cancer murine model

All animal experiments were approved by the Animal Ethics Committee of the South
Australian Health and Medical Research Institute (SAHMRI) and were conducted in
accordance with Australian National Health and Medical Research Council (NHMRC)
guidelines and the Animal Welfare Act and conformed to the Australian Code for the Care and
Use of Animals for Scientific Purposes. Male NOD scid gamma (NSG) aged 6-8 weeks were
bred and group-housed in a specific pathogen free environment at the SAHMRI. Animals had
ad libitum access to food and water and were clinically assessed daily. Mice were humanely
euthanized via CO_2 inhalation to remove tissues.

Intraprostatic injection of cancer cells was performed as previously described (38). Briefly, a
lower midline incision was made, the bladder and prostate were externalised and 10 μL of
RPMI-1640 containing 1×10^6 LNCaP cells were injected into the anterior prostate using a
precooled Hamilton syringe. To analyse tumour engraftment and growth, mice underwent
weekly *in vivo* bioluminescence imaging (IVIS, Perkin Elmer), approximately 15-20 minutes
post intraperitoneal injection with 150 mg/kg D-luciferin potassium salt.

Magnetic Resonance Imaging

After 5-6 weeks of tumour growth, mice were intravenously injected with 40 mg Fe/kg
nanoparticles in up to 100 μL of saline. After a 24-hour uptake period, mice were
anaesthetised with 2% isoflurane in oxygen and were intraperitoneally injected with an
overdose of pentobarbital prior to transcardial perfusion with 4% paraformaldehyde. Mice
were postfixed in 4% paraformaldehyde prior to storage in PBS with 0.1% sodium azide. Mice
underwent *ex vivo* T2-weighted multiple graded echo sequence acquired using a 16.4T Bruker
Avance scanner (Bruker BioSpin, GmbH), with a 30 mm SAW coil (M2M Imaging, Brisbane,
Australia) (repetition time 40 ms, echo time 10 ms, number of averages 1, slice thickness 0.1
mm) or fast low angle shot (FLASH) sequence using 9.4T Bruker BioSpec scanner (Bruker
BioSpin, GmbH, repetition time 50 ms, echo time 10 ms, number of averages 1, slice thickness
0.1 mm). At the completion of scans, tumours were excised and underwent FLASH scans using
a 11.7T Bruker Avance II scanner (Bruker BioSpin, GmbH, repetition time 100 ms, echo time
5.34 ms, number of averages 10, slice thickness 0.2 mm). All imaging was performed at room
temperature. Horos was used to view scans (Horosproject.org, sponsored by Nimble Co LLC
d/b/a Purview in Annapolis, MD USA).

Calculations of proportions of black pixels were based on protocols previously reported (39,
40). In brief, three regions of interest (ROI) were manually drawn over the prostate tumours
in whole body MR scans using ImageJ software (Bethesda, Maryland, USA) (41). Pixel intensity

histograms were generated for each ROI. A low intensity pixel threshold value was set, based
on the visual analysis of threshold peaks.

Histology and iron staining

After *ex vivo* MR imaging, excised tumours, spleens, kidneys and livers were mounted in
paraffin and 4 μm thick sections were stained with hematoxylin and eosin (H&E) or Prussian
blue using standard protocols. Images were processed using a Nanozoomer (Hamamatsu,
Japan) and were visualized using ImageJ software (Bethesda, Maryland, USA).

Statistical analyses

Data are expressed as mean ± SEM in all cases. The significance of results were determined
by unpaired t testing or one way ANOVA. Differences with a p value of less than 0.05 were
considered to be statistically significant. The relationship between FAP expression and PSA
levels and Gleason scores were assessed using simple linear regression.

Results

Human FAP expression in the normal prostate and prostate adenocarcinoma

Analysis of available transcriptomic data showed a significant increase in FAP expression from
normal prostate to adenocarcinoma ($p<0.0001$) (figure 1a). Further analysis comparing the
expression between pathological T stages demonstrated significant increases from normal
prostate to early stage (I and II) ($p<0.0001$) and to locally advanced (stage III) ($p<0.0001$).
However, there was no additional significant increase to advanced (IV) ($p>0.05$) (figure 1b).
FAP expression was further demonstrated by FAP immunohistochemistry in biopsies of low-
and high-grade prostate adenocarcinoma (figures 1e and f). Linear correlation of Gleason
score with FAP expression showed a weak positive relationship ($R^2 = 0.072$) (figure 1c),
whereas there was no correlation demonstrated with PSA ($R^2 = 0.005$) (figure 1d).

Characterization of magnetic nanoparticles

To assess the characteristics of the prepared PSMA and FAP nanoparticles, a set of physico-
chemical characterisations were carried out. TEM images presented in figure 2b and c, taken
at 60K x magnification, showed near spherical, evenly dispersed FAP and PSMA nanoparticles
even when dried on a carbon film. DLS measurements of the hydrodynamic diameters
confirmed that all of the coated nanoparticles have a narrow size distribution with an average
hydrodynamic size of 60–65 nm when measured in physiological saline (figure 2d). All
nanoparticles were also characterised with similar surface charge ranging from an average of
-17 mV to -20 mV (figure 2e).

***In vitro* binding studies**

The binding affinities of the prepared PSMA/FAP nanoparticles to the PSMA and FAP
receptors were evaluated *in vitro* by comparing their cellular uptake in cancer cell lines which
overexpress the receptors of interest. For nanoparticles used in biomedical applications, it is
important to take into consideration the potential protein adsorption onto the nanoparticle's
surface which can impact the binding activity, also known as the protein corona (42, 43). For
taking the formation of the protein corona into account, cellular uptake studies were
performed in full cell culture media containing 10% serum and incubation time was prolonged
to a 24-hour period.

First, the binding activity of PSMA nanoparticles was tested in the PSMA expressing cell line
LNCaP (44). The results presented in figure 3b indicated that the non-targeted nanoparticles
had a much lower cellular uptake compared to the targeted PSMA nanoparticles. Similarly,
the binding activity of FAP nanoparticles was tested in the human melanoma cell line C32 as
the FAP receptor is expressed in melanoma cell lines (45). Flow cytometric analysis of FAP
expression demonstrated that C32 cells had approximately 25% greater proportion of FAP-
positive cells in contrast to U87 cells, which had been previously demonstrated to have high
levels of FAP expression (figure 3a). The targeted FAP nanoparticles had an uptake almost
twice as high as the non-targeted nanoparticles (figure 3c). Considering the similar
physicochemical features of the PSMA/FAP nanoparticles when compared with the
unconjugated control nanoparticles, these cellular uptake data validate the chemical design
and specific binding affinity of the conjugated nanoparticles to their respective targets.

Magnetic resonance imaging of orthotopic prostate tumours

There was a significant increase in tumour contrast enhancement on T_2-W whole body MRI
after administration of both PSMA nanoparticles and FAP nanoparticles in mice bearing
orthotopic LNCaP tumours compared to the unconjugated control nanoparticle group, as
quantitatively demonstrated by the proportions of intratumoural black pixels (figures 4 and
5, p=0.0358 for PSMA nanoparticles and p<0.0001 for FAP nanoparticles, n=3-4 per group).
FAP nanoparticle administration yielded ~15% improvement of tumour contrast
enhancement compared to PSMA nanoparticles (p=0.0004, n=4 per group). Administration of
non-targeted control nanoparticles resulted in a modest, nonsignificant increase in tumour
contrast relative to the no nanoparticle control group (p=0.2615, n=3 per group).

Distribution of magnetic nanoparticles in tissues

To examine the intratumoural distribution of non-heme iron after the administration of the
PSMA/FAP nanoparticles, Prussian blue staining was performed on sections of LNCaP
orthotopic prostate tumours and normal prostates (figure 6). No positive staining was
observed in normal prostate sections. The absence of Prussian blue staining in sections of
normal prostate for mice administrated with control, FAP and PSMA nanoparticles confirmed
the lack of significant nanoparticle accumulation in the absence of orthotopic tumours. LNCaP
tumours also showed minimal positive basal staining. In agreement with the MRI data,

modest Prussian blue staining was observed after administration of control nanoparticles. On
the other hand, LNCaP tumours showed significant positive staining after the administration
of PSMA and FAP nanoparticles, of which the staining was concentrated on the tumour
periphery. Both PSMA and FAP nanoparticles demonstrated heterogenous intermittent
intratumoural staining, some of which was suggestive of blood vessel staining, of which the
frequency was greater in FAP nanoparticles compared to PSMA nanoparticles.

To further assess the distribution of the iron oxide nanoparticles and their effect on MRI, high
resolution MR imaging of resected tumours treated with either FAP nanoparticles or PSMA
nanoparticles was performed. Preferential contrast enhancement was observed at the
tumour peripheries, in contrast to homogenous contrast enhancement throughout the
tumours (figure 7), suggesting preferential accumulation of the nanoparticles to the highly
vascularised tumour periphery, in qualitative agreement with the Prussian blue staining data.

Additionally, Prussian blue staining was performed to assess the distribution of iron oxide
nanoparticles in the kidney, liver and spleen in orthotopic tumour mice (figure 8). Through
visual assessment there was no increase in Prussian blue staining in the spleen and kidneys in
all nanoparticle groups when compared to the no particle control. As expected, we observed
a marginal increase in Prussian blue staining in the liver compared to the no nanoparticle
control, which was consistent among all nanoparticles. Furthermore, to preliminarily assess
potential acute toxicities, H&E staining was performed on the kidneys, liver and spleen of
LNCaP tumour bearing mice (figure 8). No changes in tissue morphology were observed across
the kidneys, liver and spleen in a control, PSMA and FAP-targeting nanoparticles, when
compared to the no nanoparticle control.

Discussion

Focal boost therapies of PCa aim to selectively eliminate dominant intraprostatic malignant
lesions whilst sparing surrounding off-target tissues, and in doing so achieving oncological
outcomes similar to those afforded by whole gland treatment but with reduced adverse
effects. To enable effective focal treatment with contemporary treatment approaches
including image guided RT, target volumes and safety margins must be precisely defined.
Current MRI techniques such as mpMRI have yielded promising performance in providing
treatment guidance but are susceptible to underestimation of the true tumour volume (46)
and are time and resource demanding. This may be improved through the utilisation of
molecular MRI targeting tumour markers.

Targeting of pan-cancer marker FAP has practise-redefining potential in the treatment of solid
tumours such as PCa, through improved boundary delineation, possibly enabling the use of
focal lesion ablative microboosts. FAP is predominantly expressed on cancer associated
fibroblasts (CAFs) which reside in the tumour stroma and expression is present in virtually all
solid tumours. In addition, unlike cancer cells, CAFs are not undergoing constant mutation,
adding to the promising targeting qualities of FAP (33, 47, 48). Prostate tumours can have
over 50% stromal component, which can be greater in cases of desmoplastic reaction (29). As
such, targeting of the tumour stroma may provide increased imaging sensitivity and contrast

agent retention at the intraprostatic target site compared with targeting of the tumour cells
themselves. PCa metastases have also been demonstrated to be FAP positive, possibly
enabling focal metastatic lesion ablative treatment, with further investigation required. In
support of this train of thought, pilot studies using PSMA-PET have shown efficacy in image-
guided radiotherapy of metastases for the treatment of oligorecurrent PCa (49). In another
study, 1.5T MRI guided stereotactic body RT by LINAC guided by PSMA-PET produced
encouraging oncological and tolerability results in the management of oligometastatic
castration sensitive prostate cancer (50). Recently, FAP-targeted PET has demonstrated fair
to good interobserver agreement rates in the detection of organ and lymph node metastases
across a variety of tumour types (51). There are yet to be clinical studies investigating the use
of molecular MRI contrast agents in image-guided RT. FAP-targeted MRI-guided radiotherapy
may also prove beneficial in palliative radiotherapy, particularly in reducing pain arising from
PCa bone metastases, leading to improved quality of life (52).

FAP-specific imaging may be most beneficial in late stage and clinically aggressive cancers, in
which FAP expression can be more pronounced than PSMA and in cases with doubtful lesions,
therapy failure and/or PSMA downregulation or negative disease (31, 53). Patients
undergoing long-term ADT may have a decreased detection sensitivity on PSMA imaging due
to downregulation of the receptor, although this issue may be mitigated by ceasing ADT prior
to imaging studies (54). Current literature suggests that short-term ADT upregulates PSMA,
suggesting that PSMA-targeted imaging is suitable in these patients.

Although various FAP and PSMA-specific PET tracers are being actively developed, PET
imaging lacks the resolution that is optimal for focal radiotherapy. MR boasts superior soft
tissue contrast, visualization of organ movement and the ability to monitor changes in tumour
and tissue physiology, enabling improved adaptation of target volumes between RT
treatment fractions (55). However, interpretation of PCa mpMRI is susceptible to significant
interpretation variability as demonstrated by discordance in prostate imaging reporting and
data system (PIRADS) score assignment and cancer yield by radiologists (56). MRI provides
high resolution 3D morphological information which can be enhanced with the use of
magnetic core nanoparticles which cause hypointense signals (darkening) on MR imaging by
decreasing the relaxation times of protons in surrounding water molecules. This exogenous
contrast of the tumour may lead to decreased inter-observer variability of GTV delineation.
The utilisation of FAP or PSMA-targeted MRI may also improve the delineation of low-grade
disease, such as Gleason 3+3 cancers which are often missed on MRI (57). Furthermore, it
was found that T2-weighted MRI underestimated GTVs, longest axis and pathological tumour
extent in nearly every case. Furthermore, cellular composition, density and interstitial stromal
space have an effect on diffusion-weighted imaging (DWI) and derived apparent diffusion
coefficient (ADC) maps, which may contribute towards the underestimation of GTVs (58).

In this study we designed, synthesized and characterized FAP and PSMA targeted iron oxide
nanoparticles for molecular MRI and comparatively assessed imaging performance *in vivo* in
an orthotopic model of prostate cancer. To our knowledge, this is the first report describing
the use of FAP specific MRI. The ligand targeted nanoparticles are modified from a

lymphotropic agent that has been validated in a large animal model and presently undergoing
clinical trial (ACTRN12620000831987). This design relies on targeting and stabilising block
copolymers of different lengths that enable efficient targeting using small molecule ligands
as shown in the excellent binding affinity observed here *in vitro* with PSMA and FAP
expressing cells.

Both FAP and PSMA targeted nanoparticles yielded improved tumour accumulation as shown
by increase in MRI contrast in orthotopic tumour bearing mice compared to non-targeted
nanoparticles. It is noteworthy that FAP-targeted MRI outperformed MRI targeting PSMA, the
current clinical imaging benchmark, in contrast enhancement and delineation of prostate
tumours in the orthotopic mouse model used here. The LNCaP tumour model was selected
for this study as it commonly used for preclinical studies of PSMA specific diagnostic and
therapeutic agents. Additionally, the expression of FAP in the stroma of orthotopic LNCaP
tumours has been previously demonstrated as indicated by colocalization of both FAP and a-
smooth muscle actin (59). The excellent imaging performance of the FAP nanoparticles
observed here could be explained by the distribution of FAP expression, which predominantly
occurs in the stroma and vasculature, and might therefore be more accessible to
nanoparticulate imaging agents than markers expressed by tumour cells. Not surprisingly, the
enhanced contrast associated to the presence of the iron oxide nanoparticles was found to
be more pronounced at the periphery of the orthotopic tumours. This can be explained by
the higher density of vascular structures at the periphery of such orthotopic models (60). As
expected, unconjugated nanoparticles provided a modest improvement in the MRI contrast.
Non-specific accumulation of nanoparticles within solid tumours is typically observed in
preclinical mice models, although the precise mechanisms are still controversial (61).
Enhanced tumour contrast on MRI has also been confirmed in several clinical trials for various
dextran-coated iron oxide nanoparticles (62). The ongoing NCT04682847 trial investigates the
use of Ferumoxytol, a non-targeted iron oxide nanoparticle, in MRI-LINAC treatment of
hepatic cancer.

Considering the pan-cancer nature of FAP expression in the stroma of solid tumours, FAP-
specific MRI with rationally designed iron oxide nanoparticles may prove useful in the
delineation of other solid tumours such as gastrointestinal and brain cancer. In the future,
FAP specific MRI may also be useful to guide RT with proton therapies, thereby fully
harnessing the uniquely precise dose deposition afforded by this technique (63).
513

Disclosures
ND, VM, RM, PL and BT hold patents for FAP imaging technologies. BT is a founding member
and shareholder of Ferronova Pty Ltd. ND and VM salaries are funded through Australian
National Health and Medical Research Council grant 1158-755 and each undertake duties
involving research and development that contribute to Ferronova Pty Ltd. AS is an employee
of Ferronova Pty Ltd.
520

521 **Funding**

522 This research was supported by funding from an Australian National Health and Medical
523 Research Council grant number 1158-755 (BT as chief investigator A).

Acknowledgements
We acknowledge the facilities and scientific and technical assistance of the National Imaging
Facility, a National Collaborative Research Infrastructure Strategy (NCRIS) capability, at
SAHMRI PIRL, the University of Western Sydney and the University of Queensland.

CRediT author statement

Nicole Dmochowska: conceptualization, methodology, validation, formal analysis,
investigation, resources, data curation, writing -original draft, writing – review & editing,
visualization, project administration. **Valentina Milanova:** conceptualization, methodology,
validation, formal analysis, investigation, resources, data curation, writing – original draft,
writing – review & editing. **Ramesh Mukkamala:** resources, writing – review & editing.
Nguyen T. H. Pham: methodology, validation, investigation, resources, data curation.
Madduri Srinivasarao: resources, writing – review & editing. **Lisa Ebert:** methodology,
validation, formal analysis, investigation, data curation, writing – review & editing. **Timothy
Stait-Gardner:** resources, data curation, writing – review & editing. **Hien Le:** writing – review
& editing. **Anil Shetty:** methodology, writing – review & editing. **Phil Low:** resources, writing
– review & editing, funding acquisition. **Benjamin Thierry:** conceptualization, methodology,
resources, writing – original draft, writing – review & editing, visualization, supervision,
project administration, funding acquisition.

564 **References**
565

1.Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global Cancer
Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in
185 Countries. *CA Cancer J Clin* 2021;**71**:209-49.

2.Goodman CD, Fakir H, Pautler S, Chin J, Bauman GS. Dosimetric Evaluation of PSMA PET-
Delineated Dominant Intraprostatic Lesion Simultaneous Infield Boosts. *Adv Radiat Oncol*
2020;**5**:212-20.

3.Viani GA, Stefano EJ, Afonso SL. Higher-than-conventional radiation doses in localized
prostate cancer treatment: a meta-analysis of randomized, controlled trials. *Int J Radiat Oncol*
Biol Phys 2009;**74**:1405-18.

4.Mazariego CG, Egger S, King MT, Juraskova I, Woo H, Berry M, et al. Fifteen year quality of
life outcomes in men with localised prostate cancer: population based Australian prospective
study. *BMJ* 2020;**371**:m3503.

5.Matta R, Chapple CR, Fisch M, Heidenreich A, Herschorn S, Kodama RT, et al. Pelvic
Complications After Prostate Cancer Radiation Therapy and Their Management: An
International Collaborative Narrative Review. *Eur Urol* 2019;**75**:464-76.

6.Arrayeh E, Westphalen AC, Kurhanewicz J, Roach M, 3rd, Jung AJ, Carroll PR, et al. Does local
recurrence of prostate cancer after radiation therapy occur at the site of primary tumor?
Results of a longitudinal MRI and MRSI study. *Int J Radiat Oncol Biol Phys* 2012;**82**:e787-93.

7.Pucar D, Hricak H, Shukla-Dave A, Kuroiwa K, Drobnjak M, Eastham J, et al. Clinically
significant prostate cancer local recurrence after radiation therapy occurs at the site of
primary tumor: magnetic resonance imaging and step-section pathology evidence. *Int J*
Radiat Oncol Biol Phys 2007;**69**:62-9.

8.Murray JR, Tree AC, Alexander EJ, Sohaib A, Hazell S, Thomas K, et al. Standard and
Hypofractionated Dose Escalation to Intraprostatic Tumor Nodules in Localized Prostate
Cancer: Efficacy and Toxicity in the DELINEATE Trial. *Int J Radiat Oncol Biol Phys* 2020;**106**:715-
24.

9.Kerkmeijer LGW, Groen VH, Pos FJ, Haustermans K, Monninkhof EM, Smeenk RJ, et al. Focal
Boost to the Intraprostatic Tumor in External Beam Radiotherapy for Patients With Localized
Prostate Cancer: Results From the FLAME Randomized Phase III Trial. *J Clin Oncol*
2021;**39**:787-96.

10.Ghai S, Finelli A, Corr K, Chan R, Jokhu S, Li X, et al. MRI-guided Focused Ultrasound
Ablation for Localized Intermediate-Risk Prostate Cancer: Early Results of a Phase II Trial.
Radiology 2021;**298**:695-703.

11. Walser E, Nance A, Ynalvez L, Yong S, Aoughsten JS, Eyzaguirre EJ, et al. Focal Laser Ablation
of Prostate Cancer: Results in 120 Patients with Low- to Intermediate-Risk Disease. *J Vasc*
Interv Radiol 2019;**30**:401-9 e2.

12. Muller BG, Shih JH, Sankineni S, Marko J, Rais-Bahrami S, George AK, et al. Prostate Cancer:
Interobserver Agreement and Accuracy with the Revised Prostate Imaging Reporting and Data
System at Multiparametric MR Imaging. *Radiology* 2015;**277**:741-50.

13. Ahmed HU, El-Shater Bosaily A, Brown LC, Gabe R, Kaplan R, Parmar MK, et al. Diagnostic
accuracy of multi-parametric MRI and TRUS biopsy in prostate cancer (PROMIS): a paired
validating confirmatory study. *Lancet* 2017;**389**:815-22.

14. Bettermann AS, Zamboglou C, Kiefer S, Jilg CA, Spohn S, Kranz-Rudolph J, et al. [(68)Ga-
]PSMA-11 PET/CT and multiparametric MRI for gross tumor volume delineation in a slice by
slice analysis with whole mount histopathology as a reference standard - Implications for focal
radiotherapy planning in primary prostate cancer. *Radiother Oncol* 2019;**141**:214-9.

15. Piert M, Shankar PR, Montgomery J, Kunju LP, Rogers V, Siddiqui J, et al. Accuracy of tumor
segmentation from multi-parametric prostate MRI and (18)F-choline PET/CT for focal
prostate cancer therapy applications. *EJNMMI Res* 2018;**8**:23.

16. Piert M, Park H, Khan A, Siddiqui J, Hussain H, Chenevert T, et al. Detection of aggressive
primary prostate cancer with 11C-choline PET/CT using multimodality fusion techniques. *J*
Nucl Med 2009;**50**:1585-93.

17. Bouchelouche K, Choyke PL, Capala J. Prostate specific membrane antigen- a target for
imaging and therapy with radionuclides. *Discov Med* 2010;**9**:55-61.

18. Budaus L, Leyh-Bannurah SR, Salomon G, Michl U, Heinzer H, Huland H, et al. Initial
Experience of (68)Ga-PSMA PET/CT Imaging in High-risk Prostate Cancer Patients Prior to
Radical Prostatectomy. *Eur Urol* 2016;**69**:393-6.

19. Maurer T, Gschwend JE, Rauscher I, Souvatzoglou M, Haller B, Weirich G, et al. Diagnostic
Efficacy of (68)Gallium-PSMA Positron Emission Tomography Compared to Conventional
Imaging for Lymph Node Staging of 130 Consecutive Patients with Intermediate to High Risk
Prostate Cancer. *J Urol* 2016;**195**:1436-43.

20. Tosoian JJ, Gorin MA, Rowe SP, Andreas D, Szabo Z, Pienta KJ, et al. Correlation of PSMA-
Targeted (18)F-DCFPyL PET/CT Findings With Immunohistochemical and Genomic Data in a
Patient With Metastatic Neuroendocrine Prostate Cancer. *Clin Genitourin Cancer*
2017;**15**:e65-e8.

21.Zi F, He J, He D, Li Y, Yang L, Cai Z. Fibroblast activation protein alpha in tumor
microenvironment: recent progression and implications (review). *Mol Med Rep*
2015;**11**:3203-11.

22.Liu F, Qi L, Liu B, Liu J, Zhang H, Che D, et al. Fibroblast activation protein overexpression
and clinical implications in solid tumors: a meta-analysis. *PLoS One* 2015;**10**:e0116683.

23.Aimes RT, Zijlstra A, Hooper JD, Ogbourne SM, Sit ML, Fuchs S, et al. Endothelial cell serine
proteases expressed during vascular morphogenesis and angiogenesis. *Thromb Haemost*
2003;**89**:561-72.

24.Xin L, Gao J, Zheng Z, Chen Y, Lv S, Zhao Z, et al. Fibroblast Activation Protein-α as a Target
in the Bench-to-Bedside Diagnosis and Treatment of Tumors: A Narrative Review. *Frontiers in
Oncology* 2021;**11**.

25.Gao LM, Wang F, Zheng Y, Fu ZZ, Zheng L, Chen LL. Roles of Fibroblast Activation Protein
and Hepatocyte Growth Factor Expressions in Angiogenesis and Metastasis of Gastric Cancer.
Pathol Oncol Res 2019;**25**:369-76.

26.Giannoni E, Bianchini F, Masieri L, Serni S, Torre E, Calorini L, et al. Reciprocal activation of
prostate cancer cells and cancer-associated fibroblasts stimulates epithelial-mesenchymal
transition and cancer stemness. *Cancer Res* 2010;**70**:6945-56.

27.LeBeau AM, Brennen WN, Aggarwal S, Denmeade SR. Targeting the cancer stroma with a
fibroblast activation protein-activated promelittin protoxin. *Mol Cancer Ther* 2009;**8**:1378-86.

28.Tuxhorn JA, Ayala GE, Smith MJ, Smith VC, Dang TD, Rowley DR. Reactive stroma in human
prostate cancer: induction of myofibroblast phenotype and extracellular matrix remodeling.
Clin Cancer Res 2002;**8**:2912-23.

29.Hintz HM, Gallant JP, Vander Griend DJ, Coleman IM, Nelson PS, LeBeau AM. Imaging
Fibroblast Activation Protein Alpha Improves Diagnosis of Metastatic Prostate Cancer with
Positron Emission Tomography. *Clin Cancer Res* 2020;**26**:4882-91.

30.Khreish F, Rosar F, Kratochwil C, Giesel FL, Haberkorn U, Ezziddin S. Positive FAPI-PET/CT
in a metastatic castration-resistant prostate cancer patient with PSMA-negative/FDG-positive
disease. *Eur J Nucl Med Mol Imaging* 2020;**47**:2040-1.

31.Kessel K, Seifert R, Weckesser M, Boegemann M, Huss S, Kratochwil C, et al. Prostate-
specific membrane antigen and fibroblast activation protein distribution in prostate cancer:
preliminary data on immunohistochemistry and PET imaging. *Ann Nucl Med* 2022;**36**:293-301.

32.Kesch C, Yirga L, Dendl K, Handke A, Darr C, Krafft U, et al. High fibroblast-activation-
protein expression in castration-resistant prostate cancer supports the use of FAPI-molecular
theranostics. *Eur J Nucl Med Mol Imaging* 2021;**49**:385-9.

33.Roy J, Hettiarachchi SU, Kaake M, Mukkamala R, Low PS. Design and validation of fibroblast
activation protein alpha targeted imaging and therapeutic agents. *Theranostics*
2020;**10**:5778-89.

34.Goldman MJ, Craft B, Hastie M, Repecka K, McDade F, Kamath A, et al. Visualizing and
interpreting cancer genomics data via the Xena platform. *Nat Biotechnol* 2020;**38**:675-8.

35.Uhlen M, Fagerberg L, Hallstrom BM, Lindskog C, Oksvold P, Mardinoglu A, et al.
Proteomics. Tissue-based map of the human proteome. *Science* 2015;**347**:1260419.

36.Krishnan G, Cousins A, Pham N, Milanova V, Nelson M, Krishnan S, et al. Preclinical
evaluation of a mannose-labeled magnetic tracer for enhanced sentinel lymph node retention
in the head and neck. *Nanomedicine* 2022:102546.

37.Bryce NS, Pham BTT, Fong NWS, Jain N, Pan EH, Whan RM, et al. The composition and end-
group functionality of sterically stabilized nanoparticles enhances the effectiveness of co-
administered cytotoxins. *Biomater Sci* 2013;**1**:1260-72.

38.Linxweiler J, Korbel C, Muller A, Hammer M, Veith C, Bohle RM, et al. A novel mouse model
of human prostate cancer to study intraprostatic tumor growth and the development of
lymph node metastases. *Prostate* 2018;**78**:664-75.

39.Ngen EJ, Benham Azad B, Boinapally S, Lisok A, Brummet M, Jacob D, et al. MRI Assessment
of Prostate-Specific Membrane Antigen (PSMA) Targeting by a PSMA-Targeted Magnetic
Nanoparticle: Potential for Image-Guided Therapy. *Mol Pharm* 2019;**16**:2060-8.

40.Ngen EJ, Chen Y, Azad BB, Boinapally S, Jacob D, Lisok A, et al. Prostate-specific membrane
antigen (PSMA)-targeted photodynamic therapy enhances the delivery of PSMA-targeted
magnetic nanoparticles to PSMA-expressing prostate tumors. *Nanotheranostics* 2021;**5**:182-
96.

41.Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis.
Nat Methods 2012;**9**:671-5.

42.Amstad E, Textor M, Reimhult E. Stabilization and functionalization of iron oxide
nanoparticles for biomedical applications. *Nanoscale* 2011;**3**:2819-43.

43.Oh JY, Kim HS, Palanikumar L, Go EM, Jana B, Park SA, et al. Cloaking nanoparticles with
protein corona shield for targeted drug delivery. *Nat Commun* 2018;**9**:4548.

44.Ben Jemaa A, Sallami S, Ceraline J, Oueslati R. A novel regulation of PSMA and PSA
expression by Q640X AR in 22Rv1 and LNCaP prostate cancer cells. *Cell Biol Int* 2013;**37**:464-
70.

45.Tyulkina DV, Pleshkan VV, Alekseenko IV, Kopantseva MR, Sverdlov ED. Expression of the
FAP gene in non-fibroblast human cell lines. Development of cancer-associated fibroblast
models. *Dokl Biochem Biophys* 2016;**470**:319-21.

46.Wysock JS, Lepor H. Multi-parametric MRI imaging of the prostate-implications for focal
therapy. *Transl Androl Urol* 2017;**6**:453-63.

47.Tao L, Huang G, Song H, Chen Y, Chen L. Cancer associated fibroblasts: An essential role in
the tumor microenvironment. *Oncol Lett* 2017;**14**:2611-20.

48.Du H, Che G. Genetic alterations and epigenetic alterations of cancer-associated
fibroblasts. *Oncol Lett* 2017;**13**:3-12.

49.Stefan AK, Katharina S, Clemens K, Erik W, Matthias FH, Sonja K, et al. Clinical outcome of
PSMA-guided radiotherapy for patients with oligorecurrent prostate cancer. *Eur J Nucl Med*
Mol Imaging 2021;**48**:143-51.

50.Mazzola R, Cuccia F, Figlia V, Rigo M, Nicosia L, Giaj-Levra N, et al. Stereotactic body
radiotherapy for oligometastatic castration sensitive prostate cancer using 1.5 T MRI-Linac:
preliminary data on feasibility and acute patient-reported outcomes. *Radiol Med*
2021;**126**:989-97.

51.Serfling SE, Hartrampf PE, Zhi Y, Higuchi T, Rowe SP, Bundschuh L, et al. Interobserver
Agreement Rates on Fibroblast Activation Protein Inhibitor-Directed Molecular Imaging and
Therapy. *Clin Nucl Med* 2022.

52.De Felice F, Piccioli A, Musio D, Tombolini V. The role of radiation therapy in bone
metastases management. *Oncotarget* 2017;**8**:25691-9.

53.Isik EG, Has-Simsek D, Sanli O, Sanli Y, Kuyumcu S. Fibroblast Activation Protein-Targeted
PET Imaging of Metastatic Castration-Resistant Prostate Cancer Compared With 68Ga-PSMA
and 18F-FDG PET/CT. *Clin Nucl Med* 2022;**47**:e54-e5.

54.Vaz S, Hadaschik B, Gabriel M, Herrmann K, Eiber M, Costa D. Influence of androgen
deprivation therapy on PSMA expression and PSMA-ligand PET imaging of prostate cancer
patients. *Eur J Nucl Med Mol Imaging* 2020;**47**:9-15.

55.Otazo R, Lambin P, Pignol JP, Ladd ME, Schlemmer HP, Baumann M, et al. MRI-guided
Radiation Therapy: An Emerging Paradigm in Adaptive Radiation Oncology. *Radiology*
2021;**298**:248-60.

56.Sonn GA, Fan RE, Ghanouni P, Wang NN, Brooks JD, Loening AM, et al. Prostate Magnetic
Resonance Imaging Interpretation Varies Substantially Across Radiologists. *Eur Urol Focus*
2019;**5**:592-9.

57.Priester A, Natarajan S, Khoshnoodi P, Margolis DJ, Raman SS, Reiter RE, et al. Magnetic
Resonance Imaging Underestimation of Prostate Cancer Geometry: Use of Patient Specific
Molds to Correlate Images with Whole Mount Pathology. *J Urol* 2017;**197**:320-6.

58.Hectors SJ, Semaan S, Song C, Lewis S, Haines GK, Tewari A, et al. Advanced Diffusion-
weighted Imaging Modeling for Prostate Cancer Characterization: Correlation with
Quantitative Histopathologic Tumor Tissue Composition-A Hypothesis-generating Study.
Radiology 2018;**286**:918-28.

59.Brennen WN, Rosen DM, Wang H, Isaacs JT, Denmeade SR. Targeting carcinoma-
associated fibroblasts within the tumor stroma with a fibroblast activation protein-activated
prodrug. *J Natl Cancer Inst* 2012;**104**:1320-34.

60.Liu W, Zhu Y, Ye L, Zhu Y, Wang Y. Comparison of tumor angiogenesis in subcutaneous and
orthotopic LNCaP mouse models using contrast-enhanced ultrasound imaging. *Transl Cancer*
Res 2021;**10**:3268-77.

61.Sindhwani S, Syed AM, Ngai J, Kingston BR, Maiorino L, Rothschild J, et al. The entry of
nanoparticles into solid tumours. *Nat Mater* 2020;**19**:566-75.

62.Subhan MA, Yalamarty SSK, Filipczak N, Parveen F, Torchilin VP. Recent Advances in Tumor
Targeting via EPR Effect for Cancer Treatment. *J Pers Med* 2021;**11**.

63.Hoffmann A, Oborn B, Moteabbed M, Yan S, Bortfeld T, Knopf A, et al. MR-guided proton
therapy: a review and a preview. *Radiat Oncol* 2020;**15**:129.

763 **Figures and legends**
764

765
766
767
Figure 1. Transcriptomic analysis of *FAP* expression in clinical samples of prostate cancer. (a)
FAP gene expression values in prostate tumours compared with normal prostate obtained
from GTEX dataset. Red lines represent the medians of each group. (b) *FAP* expression in
relation to pathological T staging, (c) Gleason score and (d) PSA levels. (e) FAP
immunohistochemistry in biopsies of low grade and (f) high grade prostate adenocarcinoma
(images obtained from Human Protein Atlas).

779
780
781
782 **Figure 2**. Preparation and characterization of FAP nanoparticles, PSMA nanoparticles and
783 non-targeted control nanoparticles. (a) Schematic illustration of preparation of FAP/PSMA
784 nanoparticles. (b, c) TEM images of FAP and PSMA nanoparticles at 60K x magnification. (d)
785 Hydrodynamic size distribution by intensity and (e) zeta potentials of FAP and PSMA and
786 unconjugated control nanoparticles measured by dynamic light scattering.
787

789
790
791

Figure 3. Evaluation of *in vitro* binding activity to cells expressing molecular PSMA and FAP
targets. (a) Flow cytometric analysis of proportion of FAP-positive U87 and C32 cells. (b)
Cellular uptake of PSMA nanoparticles and non-targeted nanoparticles in PSMA-positive
LNCaP cells. (c) Cellular uptake of FAP nanoparticles and non-targeted nanoparticles in FAP-
positive C32 cells. Data presented as mean ± SEM from 3-4 replicates per condition. PSMA vs
non-targeted nanoparticles, p<0.0001, t(4)= 16.306, FAP vs non-targeted nanoparticles, p<
0.0001, t(4)= 18.892, independent sample test.

799
800

No nanoparticles

Control nanoparticles

PSMA nanoparticles

FAP nanoparticles

801
802
803
Figure 4. Representative coronal view of the lower abdomen in whole body T_2-W 16.4T MRI
of mice bearing orthotopic prostate tumours 24 hours after intravenous injection with
nanoparticles (40 mg Fe/kg) (representative images). PSMA nanoparticles and FAP
nanoparticles enhance tumour contrast on MRI, relative to the administration of no
nanoparticles and unconjugated control nanoparticles. B indicates bladder, white dotted
border indicates orthotopic prostate tumour.
810
811
812

813
814
815
816 **Figure 5**. (a) Proportion of hypointense (black) pixels on T_2-W 16.4T MRI in murine orthotopic
817 prostate tumours 24 hours after intravenous injection of FAP and PSMA nanoparticles (n=3-
818 4 per group, data presented as mean ± SEM). (b) Representative total pixel distribution of
819 analysed ROIs.

825
826
827
828 **Figure 6**. Representative images of Prussian blue staining of cross sections of normal prostates
829 and orthotopic prostate tumours after intravenous injection with no nanoparticles, control
830 nanoparticles, PSMA nanoparticles and FAP nanoparticles (all nanoparticles 40 mg Fe/kg). All
831 images 5x magnification, scale bar is 500 μm.

PSMA nanoparticles   FAP nanoparticles

Figure 7. High resolution *ex vivo* T$_2$-W 11.7T MRI of murine LNCaP tumours 24 hours after intravenous injection of PSMA and FAP nanoparticles (representative image).

869
870
871
Figure 8. Haematoxylin and eosin (bottom row) and Prussian blue (top row) staining of liver,
kidney and spleen of orthotopic LNCaP tumour mice, 24 hours after intravenous injection with
nanoparticles. No changes in H&E staining were identified in the liver, kidney and spleen in
all nanoparticle groups, compared to the no nanoparticle control. The kidneys and spleen did
not show an increase in Prussian blue staining, relative to the no particle control, however a
slight increase was observed in the liver, which was consistent across all nanoparticles. All
images 20x, scale bar is 100 µm.