1 **Complex multiple introductions drive fall armyworm invasions into Asia and Australia**

2 Rane R[1†], Walsh TK[2†], Lenancker P[3], Gock A[2], Dao TH[4], Nguyen VL[4], Khin TN[5], Amalin D[6], Chittarath K[7], Faheem
3 M[8], Annamalai S[8], Thanarajoo SS[8], Trisyono YA[9], Khay S[10], Kim J[11], Kuniata L[12], Powell K[3], Kalyebi A[13], Otim
4 MH[14], Nam K[15], d'Alençon E[15], Gordon KHJ[2], Tay WT[2†]
5
1. CSIRO, 343 Royal Parade, Parkville, VIC 3052, Australia
2. CSIRO, Black Mountain Laboratories, Clunies Ross Street, ACT 2601, Australia
3. Sugar Research Australia, 71378 Bruce Highway, Gordonvale, QLD 4865, Australia
4. Plant Protection Research Institute, Vietnam
5. Department of Agricultural Research, Myanmar
6. Department of Biology, De La Salle University, Philippines
7. Plant Protection Center, Department of Agriculture, Lao PDR
8. CAB International Southeast Asia, Malaysia
9. Department of Plant Protection, Faculty of Agriculture, Universitas Gadjah Mada, Indonesia
10. Plant Protection Division of CARDI, Ministry of Agriculture, Forestry and Fisheries, Cambodia
11. College of Agriculture and Life Science, Kangwon National University, Republic of Korea
12. New Britain Palm Oil, Ramu Agri Industry Ltd., Papua New Guinea
13. AJSK, P.O Box 16761, Kampala, Uganda.
14. National Crops Resources Research Institute, Namulonge, Kampala, Uganda
15. DGIMI, Université Montpellier, INRAE, Montpellier, France
† Applied BioSciences, Macquarie University, NSW, Australia
22
23 Correspondence: weetek.tay@csiro.au

24
Abstract

The fall armyworm (FAW) *Spodoptera frugiperda* is thought to have undergone a rapid 'west-to-east' spread
since 2016 when it was first identified in western Africa. Between 2018 and 2020, it was also recorded from
South Asia (SA), Southeast Asia (SEA), East Asia (EA), and Pacific/Australia (PA). Population genomic analyses
enabled the understanding of pathways, population sources, and gene flow in this notorious agricultural pest
species. Using neutral single nucleotide polymorphic (SNP) DNA markers, we detected genome introgression
that suggested most populations were overwhelmingly C- and R-strain hybrids. SNP and mitochondrial DNA
markers identified multiple introductions that were most parsimoniously explained by anthropogenic-assisted
spread, i.e., associated with international trade of live/fresh plants and plant products, and involved
'bridgehead populations' in countries to enable successful pest establishment in neighbouring countries.
Distinct population genomic signatures between Myanmar and China do not support the 'African origin
spread' nor the 'Myanmar source population to China' hypotheses. Significant genetic differentiation between
populations from different Australian states supported multiple pathways involving distinct SEA populations.
Our study identified Asia as a biosecurity hotspot and a FAW genetic melting pot, and demonstrated the use of
genome analysis to disentangle preventable human-assisted pest introductions from unpreventable natural
pest spread.

Key words

Bridgehead invasive populations, biosecurity weakness hotspots, secondary source populations, dispersal
pathways, gene flow directionality

Introduction

Movements of exotic arthropods are increasingly being linked with global agricultural and horticultural trade (Bedford 1980, Lopes-da-Silva et al. 2014, Tonnang et al. 2015, Tay et al. 2017, Elfekih et al. 2018, Arnemann et al. 2019, Europhyte 2018), tourism (e.g., Early et al. 2018), and as hitchhiker pests on aircraft (Porter and Hughes 1950) and vessels (e.g., Paini et al. 2018). These exotic arthropods can have significant negative impact on agricultural production (De Barro et al. 2011, Tay et al. 2013, Pozebon et al. 2020, Haile et al. 2021), and the environments (e.g., Bedford 1980, O'Dowd et al. 2003). Good phytosanitary practice together with robust biosecurity preparedness can lower the risk and help mitigate the effects of these invasive species. Invasive arthropods, especially species of agricultural importance, may also introduce novel genotypes (e.g., insecticide resistance genes and alleles; Anderson et al. 2018, Walsh et al. 2018, Tay and Gordon 2019) into related native populations through introgression and hybridisation, or into established populations of invasive species, thereby increasing their genomic diversity and further complicating pest management (e.g., Anderson et al. 2018, Valencia-Montoya et al. 2020).

The fall armyworm (FAW; *Spodoptera frugiperda*), a highly polyphagous noctuid moth native to the tropical and subtropical regions of the Americas, was officially reported in Western Africa in early 2016 (Goergen et al. 2016), followed by rapid detections across Sub-Saharan Africa (e.g., Nagoshi et al. 2018, Otim et al. 2018b, Botha et al. 2019) by February 2018. In Asia, its occurrence was first reported in South Asia (SA) in India in May 2018, and by December 2018, it had been reported in Bangladesh, Sri Lanka, and in Myanmar (i.e., Southeast Asia, SEA), and in China (i.e., East Asia, EA) by January 2019. In SEA, throughout 2019, detection of FAW was reported from Indonesia (Trisyono et al. 2019), Laos, Malaysia, Vietnam (Hang et al. 2020), the Philippines (Navasero et al. 2019), Cambodia (S Khay pers. ob.), and in EA from the Republic of Korea (Seo et al. 2019, Kim et al. 2021) and Japan (FAO 2021) in July 2019. The moth's strong flight ability (e.g., Rose et al. 1975, Sparks 1979, Jones et al. 2019) and potential to contaminate certain agricultural trade commodities (e.g., Early et al. 2018) leads to its success as a pest and its detection in increasing number of countries.

In the native range, FAW is conventionally divided into two host races, the corn (C-strain) and the rice (R-strain) preferred individuals. This distinction is evidenced by phenotypic and genotypic differences between the two. This host separation is thought to be an example of incipient speciation and has been primarily seen in North America. However, even in the native range, particularly in the tropics, this distinction seems to break down. Several studies have shown that, at the genetic level, the invasive populations are hybrids (e.g., Zhang et al. 2020; Tay et al. 2021b, 2022d).

In early 2020, FAW was detected in Pacific/Australia (PA) in Papua New Guinea (Bourke and Sar 2020, Tay et al. 2022a) and in Australia, FAW was first detected in the Erub and Saibai Islands in the Torres Strait in January 2020, and by early February it was confirmed from maize fields in Queensland (QLD), followed by detection in Western Australia (WA), Northern Territory (NT), New South Wales (NSW), Victoria (VIC), and as of April 2021, it was also reported from Tasmania, as well as Norfolk Island in the Pacific, and most recently (March 2022) also in New Zealand. The FAW is now established across northern Australia and has been reported in a number of crops including sorghum and sugarcane. Since reported by Goergen et al. (2016), the rapid detections of FAW across the sub-Saharan African nations followed by Asian (i.e., SA, SEA, EA) detections led to the general acceptance in the scientific and agricultural sectors that the pest spread eastward from an initial incursion into Africa (Day et al. 2017). This perception fits the reported patterns of detection but does it really fit the actual movement of this species across the Old World?

Studies based on nuclear DNA markers involving African (e.g., Schlum et al. 2021, Tay et al. 2022d) and Asian (e.g., Jiang et al. 2022) invasive populations identified gene flow patterns and signatures that suggested the FAW's recent spread was due to multiple introductions, with invasive populations likely to have moved from the east (i.e., EA/SEA) to Africa (e.g., Tay et al. 2022d). Genomic analyses of Chinese and African FAW populations by Gui et al. (2020) also detected signatures that were best explained by independent introduction events in EA, and a likely east-to-west movement of the FAW. The authors' conclusion however, nevertheless maintained the prevailing axiom that FAW arrived in China from western African, with its most recent origin to the Yunnan Province being from the neighbouring country of Myanmar (Sun et al. 2019, Zhang et al. 2019, Gui et al. 2020, Li et al. 2020).

Understanding the spread patterns (e.g., single invasive bridgehead effect (Guillemaud et al. 2011) vs. mass dispersion (Wilson et al. 2009)) and frequencies of introduction (e.g., single introduction (Cock et al. 2017, Nagoshi et al. 2018, Nagoshi et al. 2019b, Gui et al. 2020) vs. multiple introductions (Schlum et al. 2021, Tay et al. 2021a, Tay et al. 2021b, Jiang et al. 2022, Tay et al. 2022a, Tay et al. 2022c)) of the current invasive FAW

populations in Africa, Asia, and Pacific/Australia will have significant implications for the future management
of this pest, especially with respect to the delimitation between human-assisted introductions (i.e.,
preventable via behavioural change) vs. climatic factor-assisted natural migration (i.e., difficult to prevent/not
preventable), insecticide resistance management (Tay et al. 2021a, Tay et al. 2022c), and climate (i.e., hot and
cold) adaptation and tolerance (e.g., Foster and Cherry 1987, Keosentse et al. 2021, Yang et al. 2021, Zhang et
al. 2021), highlighting most likely agents of spread. It also highlights that evaluating possible future incursion
routes makes it necessary to identify weak points in the phytosanitary risk assessments, to enable future
targeted monitoring at the right places, and to critically assess official reporting and announcement dates. The
implications are significant, such as in reverse simulation studies (e.g., Sun et al. 2019, Wu Q et al. 2019, Li et
al. 2020, Qi et al. 2021) that relied on reported detections dates to identify potential wind patterns for
explaining pest dispersal patterns, but have not incorporated appropriate genomic/genetic data to ensure
consistency between multiple datasets, and could result in significant biosecurity concerns relating
to anthropogenic-assisted introductions not being addressed due to the perceived difficulty of preventing pest
spread via natural migration.

In this study, we aim to apply genome-wide single nucleotide polymorphic (SNP) loci to: (i) determine the
genomic diversity of invasive populations of FAW and the likelihood of multiple introduction events, and (ii)
determine the area of origin and introduction pathways of FAW populations in Australia. This information can
then be used to delineate between human-assisted vs. climate-assisted migration patterns of FAW in SEA and
PA regions and inform risk assessments for future incursions of both natural and anthropogenic spread.

Results

Strain assessment

We did not detect any C-strain individual following analysis of 138 fully assembled mitochondrial DNA
genomes (mitogenomes) from Australian samples. Our results are similar to the finding by Piggott et al. (2021)
who detected the C-strain mtCOI haplotype in Australia at very low frequencies. Proportions of C-strain to R-
strain also varied significantly across the different SEA populations (Table S1) in contrast to the patterns
observed in China, India, and African nations (e.g., Otim et al. 2018b, Nagoshi et al. 2020, Zhang et al. 2020,
Jiang et al. 2022, Tay et al. 2022d). All Australian populations analysed for their corn or rice mitochondrial
haplotypes via mitogenome assemblies of whole genome sequencing data therefore contrasted with the
invasive populations from SEA where in some countries (e.g., Myanmar, Vietnam) FAW with the C-strain mtCOI
haplotype made up approximately 50% of the populations examined (Table S1, Figs. 2a, 2b).

Mutations in the Triosephosphate isomerase (*Tpi*) gene used to differentiate the C- and R-strains suggested
that virtually all individuals from the SEA, PA, and South Korean (i.e., EA) carried the C-strain genotype at the
Tpi gene. Of all SEA and PA individuals assessed, none was homozygous for the rice allele at this locus, while a
number of samples (n = 10) were heterozygous C-/R-strains, primarily from the Philippines as well as one
sample each from Malaysia and South Korea. Similar to the conclusions of Zhang et al. (2020), at least in the
invasive range, our finding suggested that the *Tpi* is not a useful marker: not only are the majority of samples
of 'corn' preference, with limited evidence of hybrids and therefore contrasting greatly with whole genome
analysis, but both the *Tpi* and the mitochondrial markers frequently contradicted each other, and do not
provide accurate genealogy in, e.g., hybrid females or their offspring, especially if the female also mated with a
hybrid male.

Mitochondrial Genome analysis

In the invasive populations, we identified for the C-strain at least 12 unique maternal lineages forming a total
of five clades. For the R-strain, conservatively we identified 19 maternal lineages from nine clades (i.e., I - IX;
Fig. 1 'C-strain cladogram'). Combining results from this study and the study of Tay et al. (2022d), we therefore
identified at least 27 unique maternal lineages in the invasive FAW populations from across 14 countries from
Africa, South Asia, East Asia, SEA/Pacific, and Oceania.

The C-strain mitochondrial genomes detected in Southeast Asia (Clade II) differed from the one present in the
Yunnan CY and Indian populations although the four haplotypes clustered with high (87-100%) bootstrap
support. One other C-strain mitogenome was detected in one individual from China (Tay et al. 2022d) and this
remained a singleton haplotype amongst the other eight unique C-strain mitogenomes detected from African
and SE Asian populations. For the R-strain mitogenomes (Fig. 1 'R-strain cladogram'), Clade I consisted of
invasive individuals from all countries sampled to-date, although it did not contain individuals from all

populations (e.g., no individuals from Australia Sf20-4, Sf20-5 or Mackay populations). R-strain Clade I also
contained unique mitogenomes of individuals from the three China Yunnan populations, although other
Chinese R-strain mitogenome haplotypes were also detected in clades II and VI. The second largest clade
(Clade V) contained R-strain mitogenomes shared between individuals from six Australia populations, East
Africa (Tanzania, Uganda), India, South Korea, Laos, PNG, and Malaysia (i.e., blue branches), although excluded
other populations from SEA (i.e., Myanmar, Vietnam, Philippines), China, and Africa (i.e., Malawi, Benin).
Clades II, III, IV, VIII, IX contained mitogenome haplotypes of individuals from either SEA and/or Asia (China,
India) but not individuals from any of the African countries. Clades VII, VIII, and IX also have unique
mitogenome haplotypes of invasive individuals that were more similar to French Guiana (FG code) or the
Caribbean (i.e., Puerto Rico (PR), FG, or Guadeloupe) populations. The overall low bootstrap confidence at
various nodes for both C-strain and R-strain phylogenies reflect the high number of maternal lineages with low
mitogenome nucleotide diversity (i.e., mitogenome haplotypes differed by only very few nucleotide
substitutions) across both invasive and native range populations. These limitations therefore lower their
powers for use to confidently infer invasive population origins.

Fig. 1: Maximum Likelihood cladograms of unique *Spodoptera frugiperda* C-strain and R-strain partial
mitochondrial genomes based on concatenation of the 13 PCGs (11,393 bp) using IQ-Tree with 1,000 UFBoot
replications. Individuals in clades I, II, III, and IV (C-strain) and in Clades I, II, V (R-strain) that are in the same
colour scheme (i.e., green, orange, blue, or pinks) shared 100% nucleotide identity. Mitogenome haplotypes
from native individuals for both C- and R-strains are in khaki green colour. Red and dark grey dots at branch
nodes represent bootstrap values of 87-100% and 74-86%, respectively. Bootstrap values <74% are not shown.
Red arrows indicate invasive individuals' mitogenome haplotypes that are nested within native individuals.
Country and sample codes are listed in Tables S1 and S2.

Nuclear SNP Phylogenetic analysis

Maximum Likelihood (ML) phylogenetic analysis of the global *S. frugiperda* populations based on 870 neutral
single nucleotide polymorphic (SNP) markers provided evidence of complex genetic relationship especially
between individuals representing the invasive populations (Fig. 2). While the New World native populations
(USA, Puerto Rico, Guadeloupe, Mexico, French Guiana, Peru, Brazil) clustered together and shared a basal
(ancestral) lineage, the invasive populations did not identify Africa as the founding invasive population nor
support the axiom of rapid west-to-east spread across the Old World. Selected individuals from Uganda and
Benin appeared to cluster close to China/India populations, but there was no strong bootstrap support for a
shared common ancestor (i.e., bootstrap value <74%). China populations (i.e., CY, YJ, XP from Yunnan; see Fig.
1B in Guan et al. 2020) formed two closely related sister clades with long branch lengths, suggesting these
likely represented separate introduction events. The main Australian FAW population clade consisted of
multiple sister clades and were clustered with varying degree of bootstrap support values. Some of these

Australian populations appeared to shared close relationship with FE (i.e., South Korean) population (>87-
100% support (e.g., selected Burdekin individuals)), while others had largely unknown invasive origin(s).

Southeast Asian countries (i.e., Malaysia, Philippines, Laos PDR, Vietnam, Myanmar) and Pacific/Australia (i.e.,
PNG) have populations that were mixed, and interestingly were ancestral to East African populations (i.e.,
Malawi, Uganda), while some Australian populations (i.e., NSW; NT) were shown to be closely related to these
SE Asian populations with strong bootstrap support (e.g., >87-100% support for NT and NSW populations).
Myanmar and China appeared to not share FAW populations with similar SNP profiles despite being
neighbours. Multiple introductions of FAW with diverse genetic profiles and potentially similarly diverse origins
were therefore likely the main reason for their wide spread detection across the Old World.

Clear evidence of multiple introductions can therefore be seen from the phylogenetic analysis, with unique
introduction events likely underpinning the establishment of FAW populations in EA. (e.g., China) and SEA
(e.g., Malaysia). In Australia, while the NT/NSW populations shared closer evolutionary relationships with SE
Asian populations, the population origin for most of Qld and WA populations was unclear, and could be due to
a sampling effect (e.g., this study could not source FAW populations from e.g., Indonesia, Cambodia, Thailand,
Sri Lanka, Bangladesh, etc.). Some Burdekin and South Korean individuals appeared to share closer
evolutionary lineage, although the factor leading to this detected pattern remained as yet unknown.

Fig. 2: Maximum Likelihood (ML) phylogeny using 879 genome-wide non-coding SNPs with 1000 ultrafast (UF)
bootstrap replications. Node support with <74% are not shown., 74-86% are represented by grey circles, 87-
100% are represented by red circles. Country of origin is colour coded.

Population Statistics and estimates of substructure

The basic population diversity statistics for each population are listed in Table 2. Nucleotide diversity (π) varied
across a narrow range with the lowest (0.237) being from the Malaysian Kedah State population (i.e., reflected
its lab-colony background), and the highest from the Malawian population (0.324), similar to that reported

(Tay et al. 2022d; π = 0.279 – 0.329) in native and invasive populations based on the same set of 870 SNP loci.
The high nucleotide diversity estimates from the SEA, Australia and South Korean populations likely reflected
effects of limited (i.e., 870) highly polymorphic SNPs from non-coding genomic regions being used, and are
comparable to the findings from Tay et al. (2022d) that used the same sets of SNP loci. We did not detect
significant overall differences between the various invasive populations, and for the SE Asian and Australian
populations the nucleotide diversity estimates were generally between 0.258-0.291. All invasive populations
from Africa (Benin, Malawi, Uganda), SA/EA (India, China (CY, XP, XJ), South Korea), SEA (PNG, Malaysia (Johor,
Kedah, Penang), Laos, Vietnam, Myanmar, Philippines), and Australia (WA (Kununurra), NT, QLD (Strathmore,
Walkamin, Burdekin), NSW (Wee Waa)) showed higher average observed heterozygosity (Het_{obs}) than the
average expected heterozygosity (Het_{exp}), with the highest Het_{obs} seen in the Malawian population as also
reported previously (Tay et al. 2022d). Within the Australian populations, highest average observed
heterozygosity was seen in the Strathmore population, while in populations in SEA (i.e., Myanmar, Philippines)
and the EA (i.e., South Korea) all showed similar average Het_{obs}. Negative average F_{IS} values for all populations
were consistent with Het_{obs} being higher than Het_{exp}, and suggested populations were out-breeding (i.e.,
avoidance of matings between related individuals; Wright 1965). Similar to the previous findings (Tay et al.
2022d), the lower Het_{exp} (i.e., Het_{obs} > Het_{exp}; see Luikart and Cornuet 1998) could likewise indicate recent
mixing of distinct populations from SEA that suggest multiple introductions (e.g., Jiang et al. 2022, Tay et al.
2022d *cf*. Cock et al. 2017, FAO 2018, Nagoshi et al. 2019b, Nagoshi et al. 2019a; i.e., due to a recent
bottleneck from a recent western Africa founder event).

The observed heterozygosity excess detected in all invasive range populations could be further explained as
due to population sub-structure and isolation breaking through periodic migration. Significant numbers of loci
(*ca*. 30%) were also shown to not be in Hardy-Weinberg equilibrium (HWE) especially for the Malaysian (i.e.,
Kedah), but also Australian (i.e., Wee Waa, NT, Kununurra), Chinese (e.g., XP), South Korean, and Malawian
populations. Taken as a whole, genetic diversity results from this study therefore suggested that the invasive
Asian (i.e., SA, SEA, EA) FAW populations exhibited signatures of recent mixing of previously separated
populations. Simulated patterns of moth migration of various invasive FAW populations such as between
Myanmar and China (e.g., Sun et al. 2019, Wu Q et al. 2019, Li et al. 2020) and to Australia (Qi et al. 2021) are
incompatible with the population genomic data which suggests these were likely discrete and non-panmictic
FAW populations with the most probable explanation being due to multiple origins of founding populations.

Genetic differentiation analysis

Estimates of pairwise genetic differentiation (F_{ST}) between populations varied significantly (Table 2) and
extended to between populations within a country (e.g., Mackay vs. rest of Australia; Kedah vs. rest of
Malaysia). Of interest are the pairwise estimates between different Australian FAW populations from
Kununurra (Western Australia), Northern Territory, Queensland (Strathmore, Walkamin, Burdekin, Mackay)
and New South Wales (Wee Waa) that represented the most recently reported invasive populations in this
study, and predominantly showed significant differentiation amongst themselves (with the exception of the
two Queensland populations of Mackay and partially for Walkamin) and with other SEA/SA/EA countries. The
majority of non-significant population genetic differentiation estimates were in SEA where the presence of
FAW was reported earlier, i.e., since 2018 (e.g., FAO 2019, 2020) or as early as 2008 (Vu 2008, Nguyen and Vu
2009; see also Tay et al. 2022d), while across Asia (e.g., China) since 2016 but also potentially pre-2014
(Gilligan and Passoa 2014, Tay and Gordon 2019; see also Tay et al. 2022d).

Interestingly, significant genetic differentiation was observed between populations from Yunnan province in
China and populations from Myanmar, Laos, and Vietnam. Penang and Johor (Malaysia) populations were not
significantly differentiated from other SE Asian populations, nor with Ugandan and Malawian populations from
east Africa. Individuals from Benin and Mackay (Queensland, Australia) showed non-significant genetic
differentiation with all populations except with Kedah, and for Mackay also surprisingly with the Wee Waa
population from New South Wales. The South Korean population exhibited significant genetic differentiation
with SE Asian population except with Mackay, India and the Yuanjiang (YJ) population in Yunnan Province.
Finally, the Kedah population, being one of the earliest collected samples from Malaysia and having been
maintained as a laboratory population, showed strong differentiation with all populations (and lowest
nucleotide diversity, π = 0.237; Table 1) further supporting unique, non-African, introduction events in SEA.
Strong genetic differentiation suggested there was limited gene flow to breakdown sub-structure between
populations, and the F_{ST} estimates from these invasive populations therefore failed to support a west-to-east
spread pathway for the FAW. This observation instead suggested the widespread presence of genetically

distinct FAW populations, likely due to independent introductions and therefore also highlighting likely
biosecurity weaknesses especially in East Asia (e.g., China, South Korea) and SEA (e.g., Malaysia).

The genetic diversity of Australian populations identified surprisingly complex sub-structure patterns given the
short time frame of population detections across different northern Australian regions. Significant genetic
differentiation between, e.g., Kununurra (WA), Northern Territory (NT), Queensland (e.g., Strathmore,
Burdekin), and Wee Waa (NSW) populations suggests these populations likely derived from separate
establishment events. The WA Kununurra population was not significantly differentiated from the Johor State
(Malaysia), India and the Cangyuan (CY) China populations, suggesting a potential south-eastern route from
SA/SEA into north-western Australia. Contrasting this, Walkamin and Mackay populations showed non-
significant genetic differentiation with the Madang (PNG) population, suggesting a potential second pathway
for SEA individuals to arrive into the north-eastern region of Australia. Significant genetic differentiation
between WA, NT, and Qld populations suggested that at least during the early stage of pest establishment in
northern Australia, there was limited gene flow to harmonise the unique genetic background carried by these
distinct individuals, some of which exhibited also distinct insecticide resistance profiles (Tay et al. 2022b).

PCA

We selected specific populations to compare using Principal Component Analysis (PCA) as examples to support
evidence of independent introductions, as seen from Fig. 3a between China (CY, YJ, XP) populations vs.
Myanmar, in Fig. 3b (within Malaysian populations between those collected from Penang and Johor States vs.
Kedah State), in Fig. 3c for between China and East Africa (e.g., Uganda, Malawi), and where Benin and India
individuals that grouped with either China or east Africa; and in Fig. 3d between China, Malaysia (Kedah State),
and Australia (NT, NSW)). Genetic variability between Australian populations (e.g., Strathmore (QLD) vs. NT
and NSW) was also evident (Fig. 3d).

PCA also showed that differences existed between FAW populations from the Madang Province in PNG and
with the Strathmore population from Qld (Fig. 3e). The SEA FAW populations from Lao PDR/Vietnam also
exhibited diversity from the South Korean population (Fig. 3f), with the South Korean and Strathmore
populations largely exhibiting similar diversity pattern, while the Madang population shared similarity with
Laos and Vietnam populations. Plotting all SEA populations against China clearly showed that populations from
SEA were distinct from the Chinese FAW populations (Fig. 3g), while in Australia individuals from various
populations shared similarity with both Chinese and SEA FAW. Despite the connectedness of the landscape
between SEA and China, SEA largely appeared to have their own FAW populations, with FAW in SEA and in
China differing in their genome composition overall as shown via PCA.

PCA further enabled visualisation of genetic diversity amongst Australia FAW populations, suggesting that
arrival and establishment of FAW likely involved separate introduction events that followed closely after each
other and over a short timeframe. While it had been anticipated that the southward spread of FAW from SEA
would necessarily lead to Australian FAW and PNG FAW to share similar genetic backgrounds, the Madang
Province FAW population appeared to be different from the Strathmore (Qld) population, with the Madang
population being more similar to Lao PDR/Vietnam populations, and the Strathmore population more similar
to FAW from South Korea.

Pop ID	% HWE	Av. Het (Obs)	s.e.	Av. Hom(Obs)	s.e.	Av. Het(Exp)	s.e.	Av. Hom(Exp)	s.e.	Nt. Div (π)	s.e.	Av. Fis	s.e.
AUS_Kununnura	0.69	0.386	0.011	0.614	0.011	0.272	0.006	0.728	0.006	0.277	0.006	-0.265	0.075
AUS_NT	0.62	0.381	0.015	0.619	0.015	0.236	0.008	0.764	0.008	0.258	0.009	-0.239	0.012
AUS_Strathmore	0.70	0.407	0.011	0.594	0.011	0.286	0.006	0.714	0.006	0.291	0.006	-0.277	0.040
AUS_Walkamin	0.74	0.394	0.010	0.606	0.010	0.281	0.006	0.719	0.006	0.286	0.006	-0.264	0.038
AUS_Burdekin	0.70	0.395	0.011	0.605	0.011	0.281	0.006	0.719	0.006	0.286	0.006	-0.266	0.037
AUS_Mackay	0.72	0.391	0.012	0.609	0.012	0.269	0.007	0.731	0.007	0.290	0.007	-0.212	0.013
AUS_Wee Waa	0.64	0.393	0.013	0.607	0.013	0.257	0.007	0.743	0.007	0.275	0.008	-0.245	0.017
PNG	0.72	0.403	0.011	0.597	0.011	0.284	0.006	0.716	0.006	0.292	0.006	-0.261	0.022
Johore_MYS	0.80	0.400	0.011	0.600	0.011	0.279	0.006	0.721	0.006	0.294	0.006	-0.238	0.017
Kedah_MYS	0.35	0.377	0.016	0.623	0.016	0.224	0.009	0.776	0.009	0.237	0.009	-0.281	0.030
Penang_MYS	0.75	0.396	0.012	0.604	0.012	0.271	0.007	0.729	0.007	0.287	0.007	-0.236	0.015
Laos	0.79	0.390	0.010	0.610	0.010	0.282	0.005	0.718	0.005	0.289	0.006	-0.246	0.034
Vietnam	0.87	0.404	0.011	0.596	0.011	0.286	0.006	0.714	0.006	0.300	0.006	-0.238	0.020
Myanmar	0.73	0.410	0.011	0.590	0.011	0.289	0.006	0.711	0.006	0.298	0.006	-0.264	0.017
India	0.77	0.406	0.010	0.594	0.010	0.289	0.006	0.711	0.006	0.301	0.006	-0.243	0.000
ChinaCY	0.73	0.399	0.011	0.601	0.011	0.279	0.006	0.721	0.006	0.287	0.006	-0.263	0.000
ChinaXP	0.68	0.404	0.011	0.596	0.011	0.280	0.006	0.720	0.006	0.290	0.006	-0.262	0.000
ChinaYJ	0.71	0.403	0.011	0.597	0.011	0.280	0.006	0.720	0.006	0.292	0.006	-0.251	0.000
Philippines	0.74	0.412	0.011	0.588	0.011	0.289	0.006	0.711	0.006	0.297	0.006	-0.271	0.026
Skorea	0.68	0.413	0.012	0.587	0.012	0.278	0.007	0.722	0.007	0.291	0.007	-0.267	0.023
Uganda	0.75	0.434	0.010	0.566	0.010	0.306	0.005	0.694	0.005	0.316	0.005	-0.275	0.000
Malawi	0.69	0.449	0.010	0.551	0.010	0.314	0.005	0.686	0.005	0.324	0.005	-0.291	0.000
Benin	0.79	0.415	0.013	0.586	0.013	0.277	0.007	0.723	0.007	0.317	0.008	-0.184	0.000

315
316
317

Table 2: Population genetic differentiation via pairwise F_{ST} estimates between *Spodoptera frugiperda* populations from the invasive ranges of Africa (Uganda, Malawi,
Benin), South Asia (India), East Asia (China, South Korea), Southeast Asia (Malaysia (Johor, Kedah, Penang States), Laos, Vietnam, Myanmar), and Pacific/Australia (Papua
New Guinea (PNG), AUS; Kununurra (Western Australia, WA), Northern Territory (NT), Strathmore, Walkamin, Burdekin, Mackay (Queensland, Qld), Wee Waa (New South
Wales, NSW). Estimates of significance are: (*) $P \leq 0.05$; $P \leq 0.01$ (**). Values in red are significant differentiation between pairwise comparisons, values in green cells are
not significant. Western African (Benin) population (light grey column) showed non-significant pairwise F_{ST} values with all locations except with the Malaysian Kedah State
population. Refer to Table S1 and Methods section for population details and statistical calculation approach. Refer to (Tay et al. 2022d) for details of African and Asian
FAW populations.

	Kununurra	NT	Strathmore	Walkamin	Burdekin	Mackay	Wee Waa	PNG	Johore	Kedah	Penang	Laos	Vietnam	Myanmar	India	CY	XP	YJ	Philippines	Skorea	Uganda	Malawi	Benin
AUS_Kununurra	-	**	**	**	**		**	**		**	**	**	**	**		**	**		**	**	**	**	
AUS_Northern Territory (NT)	0.044	-	**	**	**	*	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	
AUS_Strathmore	0.013	0.045	-	**	**		**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	
AUS_Walkamin	0.008	0.049	0.008	-	**		**			**	**		**	**		**	**		**	**		**	
AUS_Burdekin	0.015	0.048	0.014	0.010	-		**	**	*	**	**	**	**	**	**	**	**	**	*	**	**	**	
AUS_Mackay	0.012	0.061	0.015	0.008	0.015	-	**			**													
AUS_Wee Waa	0.040	0.090	0.042	0.036	0.043	0.040	-	**	*	**	**	**	**	**	**	**	**	**	**	**	**	**	
PNG	0.014	0.041	0.015	0.009	0.015	0.013	0.037	-	**	**	**				**	**	**	*	*	**	**	**	
Malaysia_Johore	0.014	0.050	0.016	0.009	0.015	0.008	0.034	0.008	-	**						**	**	**		**			
Malaysia_Kedah	0.082	0.145	0.089	0.081	0.083	0.098	0.127	0.080	0.082	-	**	**	**	**	*	**	**	**	*	**	**	**	**
Malaysia_Penang	0.025	0.061	0.027	0.021	0.026	0.023	0.052	0.017	0.017	0.099	-	**	**	**	**	**	**	**	**	**	**	**	
Laos	0.014	0.047	0.016	0.009	0.014	0.011	0.031	0.004	0.003	0.073	0.014	-	**		**	**	**	*		**		**	
Vietnam	0.016	0.048	0.018	0.010	0.015	0.011	0.032	0.003	0.003	0.080	0.014	0.000	-		**	**	**	*		**		**	
Myanmar	0.012	0.045	0.015	0.010	0.015	0.007	0.034	0.004	0.003	0.078	0.014	0.001	0.002	-	**	**	**	*		**		**	
India	0.009	0.049	0.014	0.007	0.013	0.007	0.037	0.014	0.010	0.089	0.025	0.010	0.014	0.009	-	**	**		**	**		**	
China_Cangyuan (CY)	0.018	0.061	0.017	0.015	0.018	0.021	0.049	0.023	0.020	0.093	0.036	0.022	0.024	0.021	0.008	-	**	**	**	**	**	**	
China_Xinping (XP)	0.019	0.061	0.025	0.017	0.022	0.022	0.051	0.026	0.025	0.096	0.036	0.026	0.025	0.020	0.013	0.017	-	**	**	**	**	**	
China_Yuanjiang (YJ)	0.012	0.057	0.013	0.009	0.011	0.018	0.049	0.016	0.019	0.095	0.022	0.015	0.018	0.016	0.007	0.010	0.014	-	**	**	**	**	
Philippines	0.018	0.046	0.019	0.014	0.020	0.014	0.037	0.011	0.010	0.083	0.020	0.009	0.011	0.008	0.017	0.024	0.026	0.021	-	**	**	**	
South Krea	0.027	0.067	0.023	0.022	0.026	0.026	0.054	0.027	0.026	0.100	0.041	0.026	0.025	0.027	0.026	0.030	0.032	0.027	0.030	-	**	**	
Uganda	0.014	0.043	0.012	0.008	0.013	0.008	0.033	0.007	0.004	0.084	0.016	0.005	0.004	0.003	0.009	0.018	0.021	0.013	0.011	0.023	-		
Malawi	0.019	0.047	0.017	0.012	0.019	0.016	0.034	0.010	0.009	0.081	0.017	0.007	0.005	0.007	0.015	0.025	0.028	0.017	0.014	0.027	0.001	-	
Benin	0.015	0.061	0.014	0.008	0.012	0.018	0.049	0.010	0.007	0.106	0.024	0.005	0.008	0.007	0.003	0.014	0.021	0.007	0.012	0.029	0.002	0.006	-

Fig. 3: Principal component analysis (PCA) showing variability between selected FAW populations from their invasive ranges. **(a)** China and Myanmar; **(b)** Kedah and Johor/Penang populations from Malaysia, **(c)** China and east African (Uganda/Malawi) populations, **(d)** Australia (Strathmore, Qld/Northern Territory + New South Wales), China, and Malaysia (Kedah) populations, **(e)** Australia (Strathmore, Qld) and PNG (Madang Province) populations, **(f)** Lao PDR/Vietnam and South Korea populations, **(g)** China and SE Asian (Lao PDR/Vietnam/Myanmar/Philippines/Malaysia) and Pacific/Australia (PNG) populations, and **(h)** Australia, China and Malaysia (Kedah) populations. Note the overall population genomic variability between countries (e.g., **a**, **c-g**) and within countries (e.g., Malaysia **(b)**, Australia **(d)**). Populations with similar genomic variability are also evident, e.g., for Strathmore **(e)** and South Korea **(f)**; and for Madang **(e)** and Lao PDR/Vietnam **(f)**, further supporting potential different population origins of various FAW populations across the current invasive regions. The Southeast Asian and Chinese populations are overall different **(g)**, Australia's FAW populations showed similarity with both Southeast Asia and China **(g, h)**.

DivMigrate analysis

Directionality of gene flow between African, South Asia (Indian), East Asia (China) and SE Asian populations were predominantly from China to east African and Southeast Asian populations (e.g., Figs. 4a, 4b, Fig. S-1), while movements of FAW in Laos and Viet Nam (i.e., the Indochina region) were predominantly with other SEA countries (e.g., with Myanmar and East Africa; Figs. 4c, 4d, Fig. S-2) but with no directional movements to the three Yunnan populations (CY, XP, YJ). Migration directionality with other SE Asian populations (e.g., Johor (JB; Fig. S-3) and Penang (PN, Fig. S-4)) showed that these two populations (but especially the Johor population) were predominantly source populations for Uganda, Malawi, Philippines, Vietnam and PNG (Fig. S-3). Bidirectional migration between Myanmar and Laos PDR populations were also detected with the Johor population from Malaysia (Fig. S-3). When India was selected as the source population, bidirectional migration events were detected with Myanmar and with the Cangyuan (CY) populations (Fig. S-5) while unidirectional migration events from India to Uganda and Malawi and to Laos were detected, and the China Yuanjian (YJ) population showed unidirectional migration to India. Unidirectional migration events from CY and YJ populations to the PNG Madang population were detected, while bidirectional migration events between PNG and Myanmar, Laos PDR, Philippines, Vietnam, and with Uganda and Malawi were also detected (Fig. S-6). No migration events were detected between the West African Benin population and with the South Korean population.

Admixture analysis

Admixture analyses involving all Australian, Southeast Asian and South Korean populations from this study; and native populations from the Americas and Caribbean Islands, and invasive populations from Africa (Benin,

Uganda, Malawi), India, and China (Tay et al. 2022d), provided an overall complex picture of population structure that reflected the species' likely introduction histories across its invasive ranges.

Admixture analysis that excluded New World, African and Indian populations identified four genetic clusters (i.e., K = 4) to best describe these invasive populations from SEA, and EA (i.e., China, South Korea), and Pacific/Australia (Fig. 5a). At K = 4, Australian populations from NT and NSW, YJ population from China, South Korean, and Malaysia's Kedah population, each showed unique admixture patterns (i.e., some individuals from NT and NSW populations lacked cluster 3; most of YJ (but also some CY and XP) individuals lacked clusters 1 and 2; South Korean (e.g., MF individuals) lacked cluster 2; Malaysia's Kedah population lacked evidence of admixture (i.e., reflecting its laboratory culture history) and was made up predominantly by individuals that belonged to cluster 4. Populations from China also differed from most populations from SEA due to the overall absence of genetic cluster 4. Taken as a whole, establishment of the FAW populations in China, Malaysia, vs. other SE Asian populations, and between Australian populations (e.g., NT/NSW *cf.* WA/Qld), likely involved individuals from diverse genetic background (i.e., multiple introductions). At K = 4, the majority of Australian populations appeared to contain genetic clusters similar to China (i.e., cluster 3) and to SEA (i.e., cluster 2).

Overall admixture patterns at best K = 4 in China and SEA remained unchanged when analysed together with African and Indian individuals (Fig. 5b; excluded Australia). Benin individuals were either similar to China or to SEA, while eastern African populations (e.g., Uganda, Malawi) were similar to Southeast Asian populations from e.g., Vietnam, Laos, and is in agreement with the phylogenetic inference (Fig, 3) that identified these African individuals as having loci that were derived from Southeast Asian populations.

Genome-wide SNP loci demonstrated that invasive FAW populations from SEA and Australia exhibited admixed genomic signatures similar to that observed in other invasive populations (Zhang et al. 2020, Tay et al. 2022d). While the current invasive populations in Africa and Asia likely arrived already as hybrids as suggested by Yainna et al. (2020), the Malaysia Kedah State population was potentially established by offspring of a non-admixed female. Distinct admixture patterns in Malaysian FAW populations between Kedah and Johor/Penang states therefore suggested that establishment of these populations was likely as separate introduction events. As reported also in Tay et al. (2022d), the Chinese YJ population appeared to have admixed signature that differed from the XP and CY populations, and suggested that the YJ population could have a different introduction history than the XP and CY populations. Similar multiple genetic signatures based on lesser nuclear markers by Jiang et al. (2022) also supported likely multiple introductions of China Yunnan populations.

Table 3: DivMigrate matrix showing effective migration rates calculated using G_{ST} from source to target invasive populations. Significant unidirectional migration detected from 100 bootstrap replications (at alpha = 0.5) are in red and indicated with *. All three Chinese populations were from the Yunnan province (CY: Cangyuan, XP: Xinping, YJ: Yuanjiang). Malaysian populations are from the states of Johor and Penang as indicated in parentheses.

	Populations	Lao PDR	Malaysia (Johore)	Malaysia (Penang)	Myanmar	Papua New Guinea	Philippines	Vietnam	South Korea	Uganda	India	Benin	China (CY)	Malawi	China (XP)	China (YJ)
																Target
	Lao PDR	-	0.543	0.337	1	0.779	0.707	0.728	0.32	0.865	0.476	0.219	0.313	0.876	0.265	0.314
	Malaysia (Johore)	0.642	-	0.277	0.65	0.548	0.516	0.544	0.283	0.73	0.412	0.198	0.298	0.688	0.225	0.255
	Malaysia (Penang)	0.44	0.313	-	0.426	0.421	0.381	0.357	0.215	0.474	0.274	0.152	0.227	0.553	0.2	0.26
S	Myanmar	0.902	0.553	0.322	-	0.719	0.691	0.648	0.305	0.903	0.49	0.216	0.325	0.836	0.273	0.307
o	Papua New Guinea	0.703	0.447	0.289	0.797	-	0.601	0.613	0.299	0.777	0.42	0.2	0.284	0.723	0.242	0.301
u	Philippines	0.645	0.422	0.268	0.648	0.536	-	0.476	0.297	0.682	0.392	0.189	0.307	0.662	0.261	0.279
r	Vietnam	0.75	0.419	0.281	0.703	0.613	0.506	-	0.281	0.687	0.34	0.181	0.263	0.711	0.235	0.265
c	South Korea	0.351	0.274	0.176	0.354	0.335	0.327	0.295	-	0.394	0.322	0.145	0.291	0.392	0.261	0.27
e	Uganda	0.648	0.401	0.252	0.622	0.542	0.484	0.514	0.273	-	0.412	0.196	0.303	0.905	0.256	0.294
	India	0.545	0.364	0.223	0.544	0.452	0.459	0.412	0.271	0.618	-	0.193	0.488	0.526	0.39	0.449
	Benin	0.315	0.237	0.161	0.3	0.277	0.307	0.26	0.181	0.385	0.311	-	0.254	0.362	0.214	0.249
	China (CY)	0.492	0.379	0.209	0.521	0.459*	0.495*	0.388	0.307	0.622*	0.613	0.191	-	0.522*	0.485	0.547
	Malawi	0.571	0.33	0.233	0.537	0.473	0.456	0.487	0.249	0.792	0.344	0.179	0.258	-	0.219	0.258
	China (XP)	0.397	0.325	0.206	0.507*	0.408*	0.456	0.362	0.278	0.498*	0.452	0.168	0.411	0.485*	-	0.379
	China (YJ)	0.559	0.35	0.261	0.513	0.482	0.488	0.396	0.295	0.635*	0.516	0.213	0.51	0.619*	0.471	-

Note: Australian populations and Malaysian Kedah populations have been excluded (see main text).

Figs. 4a, 4b: DivMigrate analyses with edge weight setting at 0.453 showing unidirectional (yellow arrow lines) and bidirectional (blue arrow lines) gene flow between countries in Africa and South Asia/East Asia/SE Asia.

Significant migration rates (at alpha = 0.5) are in red and as provided in Table 3. Incidences of unidirectional
migration were predominantly detected from China (CHN) Yunnan populations (CY, XP) to SE Asian populations
(e.g., Myanmar (MMR), Laos PDR (Lao), Philippines (PHL)) and to east African populations (e.g., Uganda (UGA),
Malawi (MWI)) (Fig. 4a, 4b). Source populations are CY (Fig. 4a) and XP (Fig. 4b).

Fig. 4a   Fig. 4b

Figs. 4c, 4d: DivMigrate analyses with edge weight setting at 0.453 showing unidirectional (yellow arrow lines)
and bidirectional (blue arrow lines) migration between countries in Africa and South Asia/East Asia/SE Asia.
Migration rates between populations are as provided in Table 3. **Fig. 4c:** Vietnam (VNM) as the source
population identified an incidence of unidirectional migration from Malaysia (MYS) Johor state (JB) to Vietnam,
while bidirectional migration events were detected from Vietnam to other SE Asian (e.g., Philippines (PHL), Lao
PDR (Lao), Myanmar (MMR)), to Pacific/Australia (i.e., Papua New Guinea (PNG)), as well as to east Africa
(Uganda (UGA), Malawi (MWI)). **Fig. 4d:** Lao PDR (LAO) as source population identified bidirectional migration
events between various SEA populations and east African populations, while unidirectional migration events
were identified from India (IND) and China (CHN) Yunnan populations (CY, YJ) to Laos PDR. No migration
events were evident from SE Asian populations to China.

Fig. 4c   Fig. 4d

Fig. 5: Admixture and corresponding CV plots for FAW populations from: **(a)** Australia, China, South Korea, Lao PDR, Myanmar, Malaysia, Philippines, PNG, and Vietnam, and **(b)** Benin, China, India, South Korea, Lao PDR, Myanmar, Malaysia, Philippines, PNG, Tanzania, and Vietnam. Optimal ancestral genetic clusters are K=4 for both admixture plots. Boxed individuals have unique admixture patterns at K=4 when compared with other populations. China FAW lacked Cluster 2 (navy blue colour; present in almost all SEA and Australian FAW), while in NSW and NT some individuals lacked cluster 3. South Korea 'MF' population generally lacked cluster 2, while Kedah (Malaysia) showed distinct (cluster 4) pattern for all individuals. The overall same observations are evident in the admixture plot in (b), with African FAW generally exhibiting admixture patterns similar to SEA populations than to Chinese FAW. With the exception of Kedah (Malaysia) and some Chinese FAW individuals, all FAW in the invasive range showed evidence of genomic admixture (i.e., hybrid signature).

Discussion

In this study, we demonstrated through genomic analysis of genome-wide SNP markers signatures of multiple independent introductions of *S. frugiperda* in Asia (i.e., SEA. EA), the detection of distinct population sub-structures between what had been widely assumed (e.g., Sun et al. 2019, Wu Q et al. 2019, Li et al. 2020) as genetically linked populations between Myanmar and China, of multiple introduction pathways into north western and north eastern Australia, and the identification of potential SEA origins of eastern African (i.e., Uganda, Malawian) FAW populations. The findings highlight the need to infer population movements through genome-wide analyses; while critical assessments are needed of conclusions based on movement studies such as reverse trajectory analyses (Wu Q et al. 2019, Qi et al. 2021) and inference of population genetics relying solely on detection reports and limited (e.g., partial mitochondrial genes; partial TPI gene) DNA markers. Importantly, the study demonstrated the unexpected complexity of FAW populations across its current sub-Saharan, South Asia, Southeast Asia, East Asia, and Pacific/Australia invasive ranges since the species' initial 2016 confirmation in western Africa, and were likely underpinned by human (anthropogenic)-assisted activities, as also cautioned by Early et al. (2018) based on the prevailing opposite wind direction (i.e., east-to-west) in the African continent.

The rapid spread rate of this pest since 2016 to over 70 countries (Czepak et al. 2019) has generally been regarded as a result of its long distance migration ability (e.g., Rose et al. 1975, Jones et al. 2019, Gui et al. 2020, Nagoshi et al. 2020) following a single introduction into west Africa (Goergen et al. 2016, Cock et al. 2017, Nagoshi et al. 2018, Nagoshi et al. 2019b, Nagoshi et al. 2019a) and subsequent wind-assisted dispersal (Day et al. 2017). In contrast, genomic (e.g., Schlum et al. 2021, Tay et al. 2022d) and genetic (Jiang et al. 2022, Tay et al. 2022c) evidence has begun to emerge that the current widespread establishments of FAW populations in Asia were the result of multiple introductions, and likely associated with international agricultural trade. While pre-border interception data from SEA are generally lacking, regular interceptions of FAW in agricultural commodities by Europhyte (e.g., 2018) provided evidence to demonstrate the regular interceptions of FAW from its native countries (and increasingly, also from established invasive populations in Africa) were associated with international trade activities. These 'bridgehead invasive populations' (Guillemaud et al. 2011) established across Asia and Africa enabled subsequent establishments of localised secondary invasive populations via multiple factors including agricultural trade activities and natural migration (e.g., Early et al. 2018, Huesing et al. 2018, Jones et al. 2019).

From the analyses presented here, we suggest that the number of independent introductions of FAW is likely underestimated. For example, while our genomic analyses have identified populations in Malaysia, South Korea, and China (Tay and Gordon 2019, Tay et al. 2022d) with distinct signatures, Jiang et al. (2022) identified six separate genetic signatures in populations from southern and northern China, and Schlum et al. (2021) and Tay et al. (2022d) identified a total of least four separate introductions in African populations (e.g., Kenya, Benin, Uganda, Malawi). Population genetic analyses based on limited partial gene markers have also identified low frequencies of unusual polymorphism patterns in the *Tpi* and mtDNA gene markers in some African and an Indian FAW (e.g., Nagoshi et al. 2019b, Nagoshi et al. 2019a, Jing et al. 2021, Overton et al. 2021). Similarly, Gui et al. (2020) also detected signatures suggestive of independent introductions based on genome wide SNP analyses but have nevertheless maintained the African origin theory. While the conclusions by these authors have reinforced the hypothesis of the west-east movement of FAW, increasing whole genome evidence makes it more reasonable to conclude that these earlier single gene findings were also evidence of multiple unrecognised independent introductions.

Our investigation based on previously described genome-wide neutral SNP markers (Tay et al. 2021b) enabled integration of the published dataset (i.e., of native New World FAW populations and invasive African, Indian, and Chinese FAW populations) with the current SE Asia, East Asia, and Pacific/Australia FAW populations to reveal a complex picture of the pest's invasive history. While multiple introductions especially in the Asian continent (e.g., China, Vietnam, Malaysia; Vu 2008, Nguyen and Vu 2009, Tay and Gordon 2019, Gui et al. 2020, Tay et al. 2021b, Jiang et al. 2022, Tay et al. 2022d; this study) have played an important role leading to the perception of rapid spread, introductions from elsewhere have also contributed to the African establishment and expansion. Further to genomics evidence, resistance allele characterisation and bioassay experiments have also identified invasive populations with non-overlapping unique insecticide resistance

profiles in China (e.g., Lv et al. 2021), Indonesia (Boaventura et al. 2020), Australia (Tay et al. 2021a, Tay et al.
2022c), India (Deshmukh et al. 2020), and Africa (Eriksson 2019, Worku and Ebabuye 2019).

Multiple independent introductions of invasive agricultural pests are well-documented such as for the cryptic
whitefly *Bemisia tabaci* MED and MEAM1 species (e.g., De Barro et al. 2011, de Moraes et al. 2018, Elfekih et
al. 2018), and for the related noctuid pest *Helicoverpa armigera* (e.g., Tay et al. 2017, Arnemann et al. 2019,
see also Jones et al. 2019), and underpinned the perceived rapid spread (e.g., Leite et al. 2014) of these pests
in their introduced habitats. Early confirmation of invasive pests in novel habitats is also often challenging
especially if initial population sizes were small. For example, while the initial detections of *H. armigera* in Brazil
was during the cropping seasons of 2012/2013 (Czepak et al. 2013, Tay et al. 2013), its presence in Brazil was
only subsequently confirmed to be at least as early as in 2008 (Sosa-Gomez et al. 2016) and potentially earlier
elsewhere (e.g., in Peru) in the South American continent (Collins et al. 2007, Tay et al. 2017). Similarly for the
FAW, its presence in Vietnam was reported since 2008 (Vu 2008, Nguyen and Vu 2009, Nguyen et al. 2012),
and pre-border interceptions of FAW from non-native ranges (e.g., China, Indonesia, Israel, Micronesia,
Netherlands, Thailand, and Turkey) back to its native range by the USDA Identification Technology Program
(ITP) were pre-2014 (Gilligan and Passoa 2014, see also Tay et al. 2022d), and pre-dated the 2016 first reported
cases of FAW in east Africa (Goergen et al. 2016). In Uganda, farmer surveys (Kalyebi 2020, Kalyebi et al. 2022)
showed that FAW damage symptoms on maize crops were first identified in the Namutumba (since 2013) and
the Kamuli (since 2014) districts, while (Otim et al. 2018a) reported farmers observed FAW damage symptoms
in eastern and northern Uganda since 2014, although farmers at the time were unaware of this highly invasive
exotic armyworm species. Findings of field surveys by Otim et al. (2018a), Kalyebi (2020), and Kalyebi et al.
(2022) were in agreement with anecdotal observation and with single-gene response on Bt maize in South and
East Africa, that the FAW could have been present in Africa several years prior to its confirmation in west
Africa (Huesing et al. 2018).

While the current FAW populations in Vietnam were shown to have a gene flow directionality from Malaysia,
whether this could be linked to the 2008 report (Vu 2008, Nguyen and Vu 2009) of outbreaks around Hanoi
remained to be answered. Malaysian populations from Penang and Johor States were also shown to be linked
to, and potentially source populations for, Ugandan and Malawian FAW populations (Fig. 2; Fig. S3; Fig. S4), in
addition to the earlier results that also showed Chinese FAW populations to be potential source populations
(Tay et al. 2022d; also Fig. 4a, Fig. 4b). Our study and that of Jiang et al. (2022) therefore identified two
potential biosecurity weakness hotspots in Asia that have contributed to the spread of FAW in recent times.
The study of Jiang et al. (2022) as well as the studies of Tay et al. (2022d) and Gui et al. (2020) therefore
identified at least eight unique genetic/genomic signatures of introduction events across different regions
(e.g., southern and northern regions) in China. While reverse trajectory studies by, e.g., Wu Q et al. (2019)
suggested Yunnan province FAW to have originated from Myanmar, results from the present study showed
that there was very little genetic connectivity between them (e.g., Fig. 3a), while DivMigrate analysis showed
that gene flow directionality was more likely to be from Cangyuan (CY) to Myanmar (Fig. 4a), thereby
highlighting the importance of incorporating genomic analyses to aid interpretations of pest movements.

Southeast Asian and Pacific/Australia countries such as Laos, Myanmar, Vietnam, Philippines, Malaysia, and
PNG, appeared to be the geographic 'genetic melting pot' for the invasive FAW based on our analyses (Figs. 4a-
d; Fig. S1-S6), although their population establishment scenarios were different. For example, our analyses
suggested that Laos, Myanmar, Philippines, and PNG populations were likely linked to 'bridgehead invasive
founder China's CY population', the Vietnam population was more likely associated with individuals from
Malaysia's Johor State 'bridgehead invasive founder population' (Fig. 4C). DivMigrate analyses therefore
suggested that populations in these SE Asian countries have diverse origins, both as potential direct recipients
of New World FAW (e.g., Malaysia, also China) and as recipients of successfully established secondary invasive
populations (e.g., Laos, Vietnam, Myanmar), with subsequent bi-directional movements between these
countries.

Pest spread into Australia originating from Asia and SEA have been shown to be associated with wind patterns
(Eagles et al. 2012), and separate introduction pathways for Western Australia and Queensland insect
populations (e.g., the blue tongue virus vector *Culicoides brevitarsis*) have been reported (Tay et al. 2016).
Similarly, arrival and subsequent detection of *S. frugiperda* populations across the northern region of Australia
also likely represented separate establishments based on nuclear SNP analysis but did not support the reverse

trajectory simulation findings of Qi et al. (2021). Populations between WA, NT, and Qld were shown to be
genetically differentiated, suggesting limited gene flow between these populations at the early stage of
invasions. The Kununurra population was found to be less genetically differentiated with the Johor (Malaysia)
population (but also the population from India), suggesting a route moving south-east from South Asia/SE Asia
into Western Australia, potentially via Indonesia. The Papua New Guinea population was shown to be not
significantly differentiated from the Walkamin (Qld) population, while gene flow directionality analysis linked
Yunnan populations (e.g., CY (Fig. 4a) and XP (Fig. 4b)) with the PNG population.

Boaventura et al. (2020) showed that Indonesian FAW individuals have unique insecticide resistance alleles for
the AChE and VGSC genes. In China, Lv et al. (2021) reported diamide resistance alleles in individuals from
Guangxi and Guizhou Provinces. While this study failed to source Indonesian and Cambodian populations (due
to the difficulty of sharing and sending biological material from Indonesia and Cambodia to Australia),
significant differences in resistance to indoxacarb was nevertheless detected between WA (Kununurra) and
Qld (Strathmore) populations (Tay et al. 2021a, Tay et al. 2022c), and could be due to the different source
populations for the WA and the Qld populations. On-going studies to monitor for potential arrivals of novel
resistance alleles and those already reported in Indonesia and China to Australia will therefore be needed to
inform and impact future FAW management strategies.

Phylogenetic analyses using both mitochondrial genomes and nuclear SNP loci supported multiple
introductions at different scales. At least 12 C-strain mitochondrial genomes based on concatenated PCGs
were identified, with some mitochondrial DNA genomes being detected only in SEA populations (e.g., Lao03 in
Clade II, Lao12 in clade III). In the R-strain, at least 19 unique mitochondrial DNA genomes were detected (Fig.
1). Interestingly, and despite Australia representing the country with the most recent history of FAW's
incursions in this study, unique Australian mitochondrial DNA genomes not reported elsewhere in our other
invasive populations were nevertheless detected (e.g., S3_Sf20-1 (Clade 1), KB1_107_AusRice (Clade III),
03_SF20-5_rice (Clade IV), 52_Sf20-1_Block2_21 (Clade V), and 55_Sf20-1_Block2_24 (Clade IX)). Some of
these unique Australian mitogenomes shared close maternal evolutionary lineage with South Korean
individuals (e.g., Clade III) that was further supported by nuclear SNP phylogeny (Fig. 2). Whether this
represents linked anthropogenic-assisted invasion history remained to be confirmed, although direct flight
between the two countries would be unlikely due to the large distance (*ca.* >7,000 km). Regardless, it
highlights the challenge faced by biosecurity agencies globally in preventing accidental introductions of alien
species.

An important finding from this study has been that the SE Asian populations were largely clustered away from
other Asian (i.e., China, Indian) populations, supporting separate establishment histories of *S. frugiperda*
populations in this region, and demonstrating the role played by independent, potentially anthropogenic-
assisted introductions. The date of *S. frugiperda*'s establishment in China and SEA (e.g., Myanmar, Vietnam),
though unknown, is potentially as early as pre-2014 (Gilligan and Passoa 2014; see also Vu 2008, Nguyen and
Vu 2009, Tay et al. 2022d) and populations from SEA and China remained largely separate at least as recently
as 2019/2020, based on analysis of our 870 SNP loci (see also Jiang et al. (2022) based on microsatellite DNA
markers), contradicting the hypothesis of a Myanmar origin for the Yunnan *S. frugiperda* population (Sun et al.
2019, Wu Q et al. 2019, Li et al. 2020). Taken as a whole and incorporating population substructure (i.e., F_{ST})
analysis, directional migration analysis of invasive *S. frugiperda* populations identified East Asia (e.g., China,
South Korea) and SEA as biosecurity hotspots, while multiple introductions involving genetically distinct *S.*
frugiperda likely occurred in north-western (i.e., Kununurra, WA) and north-eastern (i.e., Burdekin, Qld)
Australia.

Population genomic studies via whole genome sequencing have enabled the spread patterns of this invasive
pest to be re-evaluated to reveal highly complex scenarios that were likely associated with, or at least initiated
by, trade activities. Whole genome sequencing is therefore proposed herein as essential for future biosecurity
monitoring, in order to allow population genomics to directly impact on development of management policies
and strategies (e.g., Guan et al. 2020), and to provide practical information to regulators and to some extent
producers on how best to manage new invasive populations. The use of genome-wide SNP loci obtained via
whole genome sequencing has enabled the elucidation of pathways to disentangle origins and establishment
histories. This, bolsters risk assessment efforts, and provide information to biosecurity agencies and to

international inter-governmental agencies to differentiate between natural and human-assisted spread. While
it would be unlikely to stop natural migration of pests, accidental introductions via human-assisted spread are
preventable. Findings from this study set the precedent that aims to inform biosecurity agencies and policy
makers to effect behavioural change in the case where pest establishment and spread were underpinned by
poor phytosanitary practices.

Material and Methods

FAW specimens for genomic analyses

Spodoptera frugiperda populations from the East Asia (i.e., South Korea), Southeast Asia (SEA; i.e., from:
Myanmar, Laos, Vietnam, Malaysia, Philippines), and Pacific/Australia (i.e., Papua New Guinea, and Australia
(representing new invasive range by *S. frugiperda*)) were surveyed for whole genome sequencing (Table S1;
Fig. 6). Methods of specimen preservation involved collection of larvae/adults from fields and placing directly
into highly concentrated ethanol (95-99.9%) and storing at -20°C until being used in DNA extraction. Samples
sent to CSIRO from Queensland (CSIRO code: Sf20-1) and Western Australia (CSIRO code: SF20-4) were first
used for bioassays purpose (Tay et al. 2021a, Tay et al. 2022c) prior to being stored at -20°C (once laboratory
colonies were established) for DNA extraction purpose. Individuals from these two locations used therefore
represent the original field-collected populations from northern Australia.

DNA extraction and genome library preparation

DNA of individual FAW samples (Table S1) was extracted using the Qiagen DNA extraction kit and eluted in
200μL elution buffer following the manufacturer's protocol. Genomic DNA libraries for individual samples were
prepared, quantified and sent to the Australian Genome Research Facility (AGRF) in Melbourne (Victoria,
Australia) for commercial sequencing.

Fig. 6: Countries where invasive populations of FAW surveyed for genomic analyses in this study were
collected from (see main text, Table S1). Native New World populations (not shown) and invasive populations
from Benin, Uganda, Tanzania, Malawi, India, and China have been reported in Tay et al. (2022d). Geographic
regions of the sampled countries were as defined by World Atlas and are provided in Table S1.

Processing of genome sequences

Genome sequencing data for each individual were trimmed to remove adapter sequences using Trim Galore!
(v 0.6.6; https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) and aligned to the *S. frugiperda*
rice genome (v1.0) (Gouin et al. 2017) using bwa_mem2 (v2) (Vasimuddin et al. 2019). Duplicate alignments

were removed using SAMBLASTER (v 0.1.26) (Faust and Hall 2014) and sorting completed using SAMtools
(v1.9) (Li et al. 2009). Variants were then predicted using BBMap (v38.90) (Bushnell 2014) and normalised
using bcftools (1.9) (Li et al. 2009). These variants were then subsampled using 870 SNP's from Tay et al.
(2022d) for further analyses.

Evolutionary and population genomic analyses

For population and evolutionary genomic analyses of the SEA, South Korean, and Pacific/Australian FAW
populations, we used the same set of the previously reported 870 neutral genome-wide SNPs (Tay et al.
2022d). We also included both native (i.e., Florida, Mississippi, Puerto Rico, Mexico, Guadeloupe, French
Guiana, Peru, Brazil) and invasive (i.e., Benin, Malawi, Uganda, Tanzania, India, China) FAW populations that
were reported in (Tay et al. 2022d) to infer the overall Maximum Likelihood (ML) phylogeny by IQ-Tree (Minh
et al. 2013). Based on the phylogenetic analysis, we undertook separate admixture analyses (Alexander et al.
2009) by: (i) excluding New World, African, and Indian populations (i.e., include East Asia (China + South Korea)
+ Southeast Asia + Pacific/Australia), and (ii) excluding New World and Australian populations (i.e., including
Africa + South Asia (India) + East Asia (Chinese, South Korea) + Southeast Asia + Papua New Guinea).

In the population genomic analysis, native FAW populations were excluded as the implication of multiple
introductions that resulted in the establishment of Old World invasive FAW populations (e.g., Schlum et al.
2021, Tay et al. 2021b, Tay et al. 2022d) meant that inclusion of our limited native populations (i.e., from eight
native localities) vs. the higher number of invasive FAW populations (i.e., from 13 invasive countries) were
unlikely to provide greater certainty to detect potential origins of invasive populations. The study therefore
uses 452 samples with an average sequencing depth of 19.24 (min. 8.71; max. 56.61; median 17.55) for
evolutionary genomic interpretations relating to the gene flow patterns of the global FAW populations.
Detailed description of the methods has been provided (Tay et al. 2021b, Tay et al. 2022d). Admixture was run
using multiple K values (3 to 9), and the cross-validation error (CV) was quantified for each K. The K value with
the least CV error (K=4 in this case) was selected for further interpretation. The 870 SNPs for this present study
have been deposited in CSIRO public Data Access Portal (Rahul et al. 2022a).

Population statistics

We used the populations program in Stacks (Catchen et al. 2013) to estimate the observed (Het_{obs}) and
expected (Het_{exp}) heterozygosities for each population. The number of loci in the global population and
individual populations departing significantly from Hardy-Weinberg equilibrium (HWE) was assessed using
PLINK 2.0 (Chang et al. 2015). The populations program in Stacks was used to calculate nucleotide diversity (π)
and F_{IS} (Wright's (1951) inbreeding coefficient) and used only variant loci with no window size specified. We
used Genepop (v4.7.5) (Raymond and Rousset 1995) to calculate pairwise comparisons of weighted F_{ST} values
between populations, with significance of differentiation between populations tested using the exact G test.

Mitochondrial Genome (mitogenome) analysis

The mitogenomes of all individuals from SEA (n=94), East Asia (South Korea, n=12), and Pacific/Australia
(n=164) were individually assembled following the approach as described in Tay et al. (2022d). Briefly, we used
Geneious 11.1.5 to map raw sequences for each FAW individual against a reference genome (GenBank
accession number MT897262 (for mtCOI C-strain)/MT897275 (for mtCOI R-strain); Tay et al. 2022d), followed
by fine-tuning to resolve regions of poor assembly and ambiguity. Fully assembled mitogenomes were then
annotated using MITOS (Bernt et al. 2013) by selecting invertebrate mitochondrial genetic code 5. Bed files
from MITOS annotations were imported into Geneious 11.1.5 to fine-tune the predicted annotations of gene
and protein coding genes (PCGs).

All FAW mitogenomes from this study and those from Tay et al. (2022d) were combined and aligned using
MAFFT Alignment v7.450 (Katoh et al. 2002, Katoh and Standley 2013) within Geneious 11.1.5, using the
default Auto option for Algorithm, 1.53 for Gap open penalty 0.123 for offset value, and 200PAM / K = 2 for
Scoring matrix. Once aligned, we trimmed and removed all intergenic regions including the A-T rich replication
origin, all tRNAs, and the two rRNA genes, leaving 11,303 bp of the 13 concatenated PCGs. We identified
unique concatenated PCG mitogenomes within individual countries using the DNAcollapser in FaBox (1.5)
(Villesen 2007), and partitioned these unique concatenated PCG mitogenomes prior to inferring ML
phylogenies without outgroups using IQ-tree implementing UFBoot (Minh et al. 2013) with 1,000 replications
to ascertain node confidence. Visualisation of the phylogeny was by Dendrogram (Huson and Scornavacca
2012) to enable a conservative estimate of the number of maternal lineages in the invasive populations. All

assembled and annotated mitogenomes used have been deposited in CSIRO public Data Access Portal (Rahul
et al. 2022b).

Strain assessment

With the non-recombinant nature of the mitogenomes, we analysed the mitogenomes of the FAW according
to their strain classification (i.e., C-strain (corn strain) and the R-strain (rice strain)) separately to identify the
conservative number of maternal founders for both strains across the invasive ranges, based on concatenation
of the 13 PCG regions only, excluding the tRNA and rRNA genes, and the variable intergenic and AT-rich
regions. We follow the naming suggestion of Tay et al. (2022b) and refer the rice-strain and corn-strain instead
as R-strain and C-strain, respectively. Individual FAWs were characterised as the R- or C- strain by comparing
their relevant partial mt*COI* gene regions against previously characterised reference sequences (rice (R-strain):
GenBank MF197867; corn (C-strain): GenBank MF197868; see Otim et al. 2018b). We note that classification of
the FAW host-strain (i.e., R-strain/C-strain) is contentious depending on the markers used (e.g., Dumas et al.
2015 vs. Nagoshi 2010) and may have limited applications in hybrid populations (Tay et al. 2022b, Tay et al.
2022d), we nevertheless provided this information based on both the partial mt*COI* gene identity and the *TPI*
exon 5 gene region as previously described (i.e., for the *TPI* loci, to differentiate between the C-strain and R-
strain *S. frugiperda,* whole genome sequence data were mapped to the SFRU_RICE_002481 contig (Gouin et
al. 2017) at location 12939.

Principle Component Analysis (PCA)

Principal Component Analysis (PCA) was also carried out to increase interpretability of the large and complex
FAW genomic dataset through maximising variance by creating new uncorrelated variables and to aid in
visualisation. PCA based on the 870 SNPs was performed using PLINK v1.9 (Purcell et al. 2007) and visualised
using ggplot2 (Wickham 2016). To enable visualisation of evidence of independent introductions we compared
(i) China (CY, YJ, XP) populations vs. Myanmar, (ii) within Malaysian populations between those collected from
Penang and Johor States vs. Kedah State; (iii) between China and East Africa (e.g., Uganda, Malawi), and (iv)
between China, Malaysia (Kedah State), and Australia (NT, NSW)).

DivMigrate analysis

DivMigrate analysis was carried out using the complete set of 870 SNP loci for all invasive populations from
Africa, Asia, and SEA, with the exception of the Kedah State (Malaysia) population that showed unique
genomic background (and represented a population from the on-going lab-maintained culture first collected
from maize field in 2019). Australia populations representing the 'final arrival' populations, were also excluded
in the DivMigrate analysis to increase interpretability relating to East Africa, South Asia, East Asia and SEA
populations. Relative migration was assessed using the G_{ST} statistics with 100 bootstrap replications and filter
threshold (i.e., visualisation of the number of links in the network plot) was set at 0.453 and plotted using the
graph visualisation and manipulation software Gephi 0.9.2 <www.gephi.org> and countries were separately
drawn using Microsoft PowerPoint for Mac v16.54.

Acknowledgements

We would like to thank Dr Brian Thistleton (Department of Industry, Tourism and Trade, NT), Dr Ian Newton
and Dr Melina Miles (QDAF), Mr Jack Daniel (Northern Australia Crop Research Alliance Pty Ltd, Kununurra
WA), Prof. Myron Zalucki (University of Queensland), Mrs Anna Madden (Wee Waa, NSW), and Dr Simone
Heimoana (CSIRO), for providing Australian FAW specimens for genomic analyses. The Erub Island FAW was
kindly provided by the Department of Agriculture, Water and the Environment (DAWE) and Northern Australia
Quarantine Strategy (NAQS). Van NTH and Huyen NPT (PPRI, Vietnam) helped with FAW specimen and related
metadata collection. DR. Donald Kachigamba provided samples from Malawi. Bill James (CSIRO) helped with
laboratory work, Dr Sharon Downes, Bill James, and Dr Sarina Macfadyen (CSIRO), and Dr Eric Huttner (ACIAR)
provided helpful discussion during the course of this work. The work performed at INRAE was publicly funded
through the ANR (the French National Research Agency, Grant ID 1702-018, given to K.N.) under the
"Investissements d'avenir" programme with the reference ANR-10-LABX-001-01 Labex Agro and coordinated
by Agropolis Fondation under the frame of I-SITE MUSE (ANR-16-IDEX-0006). It was also funded by a grant
from the department of Santé des Plantes et Environnement at Institut national de la recherche agronomique
for K.N. (adaptivesv), and was also financially supported by EUPHRESCO (FAW-spedcom, given to Anne-
Nathalie Volkoff). The project led by CSIRO is a co-investment by the Grains Research and Development
Corporation (GRDC), the Australian Centre for International Agricultural Research (ACIAR), the Cotton Research

and Development Corporation, FMC Australasia and Corteva Agriscience. GRDC manages the project on behalf
of the co-investors.

Data availability

All data used in this study have been deposited in the CSIRO Data Access Portal and are freely accessible,
under the permanent links https://doi.org/10.25919/24gy-g132 and https://doi.org/10.25919/c53m-ts57.

Competing interests

All authors declare no competing interests

References

Alexander, D. H., J. Novembre, and K. Lange. 2009. Fast model-based estimation of ancestry in unrelated
individuals. Genome Res 19: 1655-1664.

Anderson, C. J., J. G. Oakeshott, W. T. Tay, K. H. J. Gordon, A. Zwick, and T. K. Walsh. 2018. Hybridization and
gene flow in the mega-pest lineage of moth, *Helicoverpa*. Proc Natl Acad Sci U S A 115: 5034-5039.

Arnemann, J. A., S. Roxburgh, T. Walsh, J. Guedes, K. Gordon, G. Smagghe, and W. T. Tay. 2019. Multiple
incursion pathways for *Helicoverpa armigera* in Brazil show its genetic diversity spreading in a connected
world. Sci Rep 9: 19380.

Bedford, G. O. 1980. Biology, Ecology, and Control of Palm Rhinoceros Beetles. Annual Review of Entomology
25: 309-339.

Bernt, M., A. Donath, F. Juhling, F. Externbrink, C. Florentz, G. Fritzsch, J. Putz, M. Middendorf, and P. F.
Stadler. 2013. MITOS: improved de novo metazoan mitochondrial genome annotation. Mol Phylogenet
Evol 69: 313-319.

Boaventura, D., M. Martin, A. Pozzebon, D. Mota-Sanchez, and R. Nauen. 2020. Monitoring of Target-Site
Mutations Conferring Insecticide Resistance in *Spodoptera frugiperda*. Insects 11(8), 545.

Botha, A. S., A. Erasmus, H. du Plessis, and J. Van den Berg. 2019. Efficacy of Bt Maize for Control of
Spodoptera frugiperda (Lepidoptera: Noctuidae) in South Africa. Journal of Economic Entomology 112:
1260-1266.

Bourke, M., and S. Sar. 2020. Fall armyworm in Papua New Guinea: how big is the risk?, pp. 5. Australian
National University.

Bushnell, B. 2014. BBMap: A Fast, Accurate, Splice-Aware Aligner. Lawrence Berkeley National Laboratory. .

Catchen, J., P. A. Hohenlohe, S. Bassham, A. Amores, and W. A. Cresko. 2013. Stacks: an analysis tool set for
population genomics. Mol Ecol 22: 3124-3140.

Cock, M. J. W., P. K. Beseh, A. G. Buddie, G. Cafa, and J. Crozier. 2017. Molecular methods to detect
Spodoptera frugiperda in Ghana, and implications for monitoring the spread of invasive species in
developing countries. Sci Rep 7: 4103.

Collins, D., S. Martins, Boonhamm N, K. Walsh, R. Glover, and J. Tomlinson. 2007. Defra SID5 Research
Project Final Report PH0311. Molecular and morphological identification of the eggs and caterpillars of
quarantine listed Noctuidae pp. 21. Department for Environment, Food and Rural Affairs.

Czepak, C., C. Albernaz, L. M. Vivan, H. O. Guimarães, and T. Carvalhais. 2013. First reported occurrence of
Helicoverpa armigera (Hübner) (Lepidoptera: Noctuidae) in Brazil. Pesq. Agropec. Trop., Goiânia 43: 110-
113.

Czepak, C., W. T. Tay, M. H. Otim, S. R. Roy, K. C. A. Godinho, V. Magalhães, and K. F. S. Collier. 2019.
Especial *Spodoptera*: Migração acelerada. Cultivar Grandes Culturas 244: 26-29.

Day, R., P. Abrahams, M. Bateman, T. Beale, V. Clottey, M. Cock, Y. Colmenarez, N. Corniani, R. Early, J.
Godwin, J. Gomez, P. G. Moreno, S. T. \Murphy, B. Oppong-Mensah, N. Phiri, C. Pratt, S. Silvestri, and A.
Witt. 2017. Fall armyworm: impacts and implications for Africa. Outlooks on Pest Management 28: 196-201

De Barro, P. J., S. S. Liu, L. M. Boykin, and A. B. Dinsdale. 2011. *Bemisia tabaci*: a statement of species status.
Annu Rev Entomol 56: 1-19.

de Moraes, L. A., C. Muller, R. C. O. D. Buenos, A. Santos, V. H. Bello, B. R. De Marchi, L. F. M. Watanabe, J.
M. Marubayashi, B. R. Santos, V. A. Yuki, H. M. Takada, D. R. de Barros, C. G. Neves, F. N. da Silva, M. J.
Goncalves, M. Ghanim, L. Boykin, M. A. Pavan, and R. Krause-Sakate. 2018. Distribution and
phylogenetics of whiteflies and their endosymbiont relationships after the Mediterranean species invasion
in Brazil. Sci Rep 8: 14589.

Deshmukh, S., H. B. Pavithra, C. M. Kalleshwaraswamy, B. K. Shivanna, M. S. Maruthi, and D. Mota-Sanchez.
2020. Field efficacy of insecticides for management of invasive fall armyworm, *Spodoptera frugiperda* (J. E.
Smith) (Lepidoptera: Noctuidae) on maize in India. Fla Entomol 103: 221-227.

Dumas, P., J. Barbut, B. Le Ru, J. F. Silvain, A. L. Clamens, E. d'Alencon, and G. J. Kergoat. 2015. Phylogenetic
molecular species delimitations unravel potential new species in the pest genus *Spodoptera* Guenee, 1852
(Lepidoptera, Noctuidae). PLoS One 10: e0122407.

Eagles, D., T. Deveson, P. J. Walker, M. P. Zalucki, and P. Durr. 2012. Evaluation of long-distance dispersal of
Culicoides midges into northern Australia using a migration model. Med Vet Entomol 26: 334-340.

Early, R., P. Gonzalez-Moreno, S. T. Murphy, and R. Day. 2018. Forecasting the global extent of invasion of the
cereal pest *Spodoptera frugiperda*, the fall armyworm. Neobiota 40: 25-50.

Elfekih, S., P. Etter, W. T. Tay, M. Fumagalli, K. Gordon, E. Johnson, and P. De Barro. 2018. Genome-wide
analyses of the *Bemisia tabaci* species complex reveal contrasting patterns of admixture and complex
demographic histories. PLoS One 13: e0190555.

Eriksson, J. 2019. Toxicity bioassays with insecticide formulations used for control of *Spodoptera frugiperda*
(Lepidoptera: Noctuidae). Master of Science in Environmental Science with Integrated Pest Management,
North West University South Africa.

FAO. 2018. Briefing note on FAO actions on fall armyworm in Africa.

FAO. 2019. First detection report of the fall armyworm *Spodoptera frugiperda* (Lepdioptera: Noctuidae) on
maize in Myanmar. Food and Agriculture Organization of the United Nations, International Plant Protection
Convention.

FAO. 2020. Fall armywormm (FAW): impact assessment and the Global Action for Fall Armyworm control, pp.
3.

Faust, G. G., and I. M. Hall. 2014. SAMBLASTER: fast duplicate marking and structural variant read extraction.
Bioinformatics 30: 2503-2505.

Foster, R. E., and R. H. Cherry. 1987. Survival of Fall Armyworm, *Spodoptera frugiperda*, (Lepidoptera,
Noctuidae) Exposed to Cold Temperatures. Fla Entomol 70: 419-422.

Gilligan, T. M., and S. Passoa, C. 2014. LepIntercept, An identification resource for intercepted Lepidoptera
larvae. Identification Technology Program (ITP), USDA-APHIS-PPQ-S&T, Fort Collins, CO.

Goergen, G., P. L. Kumar, S. B. Sankung, A. Togola, and M. Tamo. 2016. First Report of Outbreaks of the Fall
Armyworm *Spodoptera frugiperda* (J E Smith) (Lepidoptera, Noctuidae), a New Alien Invasive Pest in West
and Central Africa. PLoS One 11: e0165632.

Gouin, A., A. Bretaudeau, K. Nam, S. Gimenez, J. M. Aury, B. Duvic, F. Hilliou, N. Durand, N. Montagne, I.
Darboux, S. Kuwar, T. Chertemps, D. Siaussat, A. Bretschneider, Y. Mone, S. J. Ahn, S. Hanniger, A. G.
Grenet, D. Neunemann, F. Maumus, I. Luyten, K. Labadie, W. Xu, F. Koutroumpa, J. M. Escoubas, A.
Llopis, M. Maibeche-Coisne, F. Salasc, A. Tomar, A. R. Anderson, S. A. Khan, P. Dumas, M. Orsucci, J. Guy,
C. Belser, A. Alberti, B. Noel, A. Couloux, J. Mercier, S. Nidelet, E. Dubois, N. Y. Liu, I. Boulogne, O.
Mirabeau, G. Le Goff, K. Gordon, J. Oakeshott, F. L. Consoli, A. N. Volkoff, H. W. Fescemyer, J. H. Marden,
D. S. Luthe, S. Herrero, D. G. Heckel, P. Wincker, G. J. Kergoat, J. Amselem, H. Quesneville, A. T. Groot, E.
Jacquin-Joly, N. Negre, C. Lemaitre, F. Legeai, E. d'Alencon, and P. Fournier. 2017. Two genomes of highly
polyphagous lepidopteran pests (*Spodoptera frugiperda*, Noctuidae) with different host-plant ranges. Sci
Rep 7: 11816.

Guan, F., J. Zhang, H. Shen, X. Wang, A. Padovan, T. K. Walsh, W. T. Tay, K. H. J. Gordon, W. James, C.
Czepak, M. H. Otim, D. Kachigamba, and Y. Wu. 2020. Whole-genome sequencing to detect mutations
associated with resistance to insecticides and Bt proteins in *Spodoptera frugiperda*. Insect Sci 28(3): 627-
638.

Gui, F. R., T. M. Lan, Y. Zhao, W. Guo, Y. Dong, D. M. Fang, H. Liu, H. M. Li, H. L. Wang, R. S. Hao, X. F. Cheng,
Y. H. Li, P. C. Yang, S. K. Sahu, Y. P. Chen, L. Cheng, S. Q. He, P. Liu, G. Y. Fan, H. R. Lu, G. H. Hu, W. Dong,
B. Chen, Y. Jiang, Y. W. Zhang, H. H. Xu, F. Lin, B. Slipper, A. Postma, M. Jackson, B. A. Abate, K. Tesfaye,
A. L. Demie, M. D. Bayeleygne, D. T. Degefu, F. Chen, P. K. Kuria, Z. M. Kinyua, T. X. Liu, H. M. Yang, F. N.
Huang, X. Liu, J. Sheng, and L. Kang. 2020. Genomic and transcriptomic analysis unveils population
evolution and development of pesticide resistance in fall armyworm *Spodoptera frugiperda*. Protein Cell
doi: 10.1007/s13238-020-00795-7.

Guillemaud, T., M. Ciosi, E. Lombaert, and A. Estoup. 2011. Biological invasions in agricultural settings:
Insights from evolutionary biology and population genetics. Cr Biol 334: 237-246.

Haile, F., T. Nowatzki, and N. Storer. 2021. Overview of Pest Status, Potential Risk, and Management
Considerations of *Helicoverpa armigera* (Lepidoptera: Noctuidae) for US Soybean Production. J Integr Pest
Manag 12(1): 3. https://doi.org/10.1093/jipm/pmaa030.

Huesing, J. E., B. M. Prasanna, D. McGrath, P. Chinwada, P. Jepson, and J. L. Capinera. 2018. Chapter 01.
Integrated Pest Management of Fall Armyworm in AFrica: An Introduction., pp. 120. In B. M. Prasanna, J. E.
Huesing, R. Eddy and V. M. Peschke (eds.), Fall armyworm in Africa: A guid or inegrated pest management,
First Edition ed. CDMX: CIMMYT, Mexico.

Huson, D. H., and C. Scornavacca. 2012. Dendroscope 3: an interactive tool for rooted phylogenetic trees and
networks. Syst Biol 61: 1061-1067.

Jiang, Y. Y., Y. Y. Zhang, X. Y. Zhou, X. Y. Hong, and L. Chen. 2022. Population genetics reveal multiple
independent invasions of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in China. Bull Entomol Res: 1-11.

Jing, W., C. Huang, C. Y. Li, H. X. Zhou, Y. L. Ren, Z. Y. Li, L. S. Xing, B. Zhang, X. Qiao, B. Liu, C. H. Liu, Y. Xi, W.
X. Liu, W. K. Wang, W. Q. Qian, S. Mckirdy, and F. H. Wan. 2021. Biology, invasion and management of the
agricultural invader: Fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae). Journal of
Integrative Agriculture 20: 646-663.

Jones, C. M., H. Parry, W. T. Tay, D. R. Reynolds, and J. W. Chapman. 2019. Movement Ecology of Pest
Helicoverpa: Implications for Ongoing Spread. Annu Rev Entomol 64: 277-295.

Kalyebi, A. 2020. Documentation of farmer practices that have been useful in Africa for managing the fall
armyworm (*Spodoptera frugiperda*): A case study from Uganda. pp. 30. GRDC.

Kalyebi, A., T. Walsh, and W. T. Tay. 2022. Farmer perception of impacts of *Spodoptera frugiperda* and
transferability of management practices in Uganda. In Prep.

Katoh, K., and D. M. Standley. 2013. MAFFT multiple sequence alignment software version 7: improvements
in performance and usability. Mol Biol Evol 30: 772-780.

Katoh, K., K. Misawa, K. Kuma, and T. Miyata. 2002. MAFFT: a novel method for rapid multiple sequence
alignment based on fast Fourier transform. Nucleic Acids Res 30: 3059-3066.

Keosentse, O., R. Mutamiswa, H. Du Plessis, and C. Nyamukondiwa. 2021. Developmental stage variation in
Spodoptera frugiperda (Lepidoptera: Noctuidae) low temperature tolerance: implications for
overwintering. Austral Entomol 60: 400-410.

Kim, J., H. Y. Nam, M. Kwon, H. J. Kim, H. J. Yi, S. Haenniger, M. Unbehend, and D. G. Heckel. 2021.
Development of a simple and accurate molecular tool for *Spodoptera frugiperda* species identification
using LAMP. Pest Management Science 77: 3145-3153.

Leite, N. A., A. Alves-Pereira, A. S. Correa, M. I. Zucchi, and C. Omoto. 2014. Demographics and Genetic
Variability of the New World Bollworm (*Helicoverpa zea*) and the Old World Bollworm (*Helicoverpa*
armigera) in Brazil. PLoS ONE 9(11): e113286.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin, and S.
Genome Project Data Processing. 2009. The Sequence Alignment/Map format and SAMtools.
Bioinformatics 25: 2078-2079.

Li, X. J., M. F. Wu, J. Ma, B. Y. Gao, Q. L. Wu, A. D. Chen, J. Liu, Y. Y. Jiang, B. P. Zhai, R. Early, J. W. Chapman,
and G. Hu. 2020. Prediction of migratory routes of the invasive fall armyworm in eastern China using a
trajectory analytical approach. Pest Manag Sci 76: 454-463.

Lopes-da-Silva, M., M. M. Sanches, A. R. Stancioli, G. Alves, and R. Sugayama. 2014. The Role of Natural and
Human-Mediated Pathways for Invasive Agricultural Pests: A Historical Analysis of Cases from Brazil. .
Agricultural Sciences 5: 634-646.

Luikart, G., and J. M. Cornuet. 1998. Empirical evaluation of a test for identifying recently bottlenecked
populations from allele frequency data. Conserv Biol 12: 228-237.

Lv, S. L., Y. Shi, J. C. Zhang, P. Liang, L. Zhang, and X. W. Gao. 2021. Detection of ryanodine receptor target-
site mutations in diamide insecticide-resistant *Spodoptera frugiperda* in China. Insect Science 28: 639-648.

Minh, B. Q., M. A. Nguyen, and A. von Haeseler. 2013. Ultrafast approximation for phylogenetic bootstrap.
Mol Biol Evol 30: 1188-1195.

Nagoshi, R. N. 2010. The Fall Armyworm Triose Phosphate Isomerase (*Tpi*) Gene as a Marker of Strain Identity
and Interstrain Mating. Ann Entomol Soc Am 103: 283-292.

Nagoshi, R. N., G. Goergen, H. D. Plessis, J. van den Berg, and R. Meagher, Jr. 2019a. Genetic comparisons of
fall armyworm populations from 11 countries spanning sub-Saharan Africa provide insights into strain
composition and migratory behaviors. Sci Rep 9: 8311.

Nagoshi, R. N., G. Goergen, K. A. Tounou, K. Agboka, D. Koffi, and R. L. Meagher. 2018. Analysis of strain
distribution, migratory potential, and invasion history of fall armyworm populations in northern Sub-
Saharan Africa. Sci Rep 8: 3710.

Nagoshi, R. N., I. Dhanani, R. Asokan, H. M. Mahadevaswamy, C. M. Kalleshwaraswamy, Sharanabasappa,
and R. L. Meagher. 2019b. Genetic characterization of fall armyworm infesting South Africa and India
indicate recent introduction from a common source population. PLoS ONE 14(5): e0217755.

Nagoshi, R. N., N. N. Htain, D. Boughton, L. Zhang, Y. Xiao, B. Y. Nagoshi, and D. Mota-Sanchez. 2020.
Southeastern Asia fall armyworms are closely related to populations in Africa and India, consistent with
common origin and recent migration. Sci Rep 10: 1421.

Navasero, M. V., M. M. Navasero, G. A. S. Burgonio, K. P. Ardez, M. D. Ebuenga, M. J. B. Beltran, M. B. Bato,
P. G. Gonzales, G. L. Magsino, B. L. Caoili, A. L. A. Barrion-Dupo, and M. F. G. M. Aquino. 2019. Detection
of the fall armyworm, *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) using larval
mrophological characters, and observations on its current local distribution in the Philippines. Philipp Ent
33: 171-184.

Nguyen, T. K. O., and T. P. Vu. Year. Published. Checklist of turfgrass insect pests, morphology, biology and
population fluctuation of *Herpetograma phaeopteralis* (Guenee) (Lepidoptera: Pyralidae) in Ha Noi, in
spring-summer 2008., pp. 1490-1498. *In,* The 3rd National Conference of Ecology and Natural Resources,
22-October, 2009 2009, Hanoi, Vietnam.

Nguyen, V. D., Q. H. Ha, and T. T. C. Nguyen. 2012. Leaves eating worm. In V. L. Pham (ed.), Vietnam insects
and pests.

O'Dowd, D. J., P. T. Green, and P. S. Lake. 2003. Invasional 'meltdown' on an oceanic island. Ecol Lett 6: 812-
817.

Otim, M. H., S. E. Adur, R. M. Ariong, O'bongo, I, G. Aogon, and S. Angudubo. 2018a. The Socio-economic
impact o fall armyworm invasion on the production and income of maize value chain actors in Uganda, pp.
64. National Agricultural Research Organization, Uganda.

Otim, M. H., W. T. Tay, T. K. Walsh, D. Kanyesigye, S. Adumo, J. Abongosi, S. Ochen, J. Sserumaga, S. Alibu,
G. Abalo, G. Asea, and A. Agona. 2018b. Detection of sister-species in invasive populations of the fall
armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae) from Uganda. PLoS ONE 13: e0194571.

Overton, K., J. L. Maino, R. Day, P. A. Umina, B. Bett, D. Carnovale, S. Ekesi, R. Meagher, and O. L. Reynolds.
2021. Global crop impacts, yield losses and action thresholds for fall armyworm (*Spodoptera frugiperda*): A
review. Crop Prot 145: 105641.

Paini, D. R., P. Mwebaze, P. M. Kuhnert, and D. J. Kriticos. 2018. Global establishment threat from a major
forest pest via international shipping: Lymantria dispar. Sci Rep 8: 13723.

Piggott, M. P., F. P. J. Tadle, S. Patel, K. C. Gomez, and B. Thistleton. 2021. Corn-strain or rice-strain?
Detection of fall armyworm, *Spodoptera frugiperda* (JE Smith) (Lepidoptera: Noctuidae), in northern
Australia. Int J Trop Insect Sc 41: 2607-2615.

Porter, J. E., and J. H. Hughes. 1950. Insect Eggs Transported on the Outer Surface of Airplanes. Journal of
Economic Entomology 43: 555-557.

Pozebon, H., R. P. Marques, G. Padilha, O. N. M, I. Valmorbida, J. G. Bevilaqua, W. T. Tay, and J. A.
Arnemann. 2020. Arthropod Invasions Versus Soybean Production in Brazil: A Review. J Econ Entomol
113(4): 1591-1608.

Purcell, S., B. Neale, K. Todd-Brown, L. Thomas, M. A. Ferreira, D. Bender, J. Maller, P. Sklar, P. I. de Bakker,
M. J. Daly, and P. C. Sham. 2007. PLINK: a tool set for whole-genome association and population-based
linkage analyses. Am J Hum Genet 81: 559-575.

Qi, G. J., J. Ma, J. Wan, Y. L. Ren, S. McKirdy, G. Hu, and Z. F. Zhang. 2021. Source Regions of the First
Immigration of Fall Armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae) Invading Australia. Insects
12(12): 1104.

Rane R, Walsh TK, Lenancker P, Gock A, Hang DT, Liem NV, Khin TN, Amalin D, Chittarath K, Faheem M,
Sivapragasam A, Thanarajoo SS, Trisyono YA, Khay S, Kim J, Kuniata L, Powell K, Kalyebi A, Otim MH,
Gordon KHJ, Tay WT. Complex multiple introductions drive fall armyworm invasions into Asia and
Australia: Genome-wide single nucleotide polymorphic (SNP) loci of fall armyworm (FAW) *Spodoptera*
frugiperda from Southeast Asia, East Asia, and Pacifi/Australia. CSIRO Data Access Portal,
https://doi.org/10.25919/24gy-g132 (2022a).

Rane R, Walsh TK, Lenancker P, Gock A, Hang DT, Liem NV, Khin TN, Amalin D, Chittarath K, Faheem M,
Sivapragasam A, Thanarajoo SS, Trisyono YA, Khay S, Kim J, Kuniata L, Powell K, Kalyebi A, Otim MH,
Gordon KHJ, Tay WT. Complex multiple introductions drive fall armyworm invasions into Asia and
Australia: Mitochondrial DNA genomes of fall armyworm (FAW) *Spodoptera frugiperda* from Southeast
Asia, East Asia, and Pacifi/Australia. CSIRO Data Access Portal, https://doi.org/10.25919/c53m-ts57
(2022b).

Raymond, M., and F. Rousset. 1995. Genepop (Version-1.2) - Population-Genetics Software for Exact Tests and
Ecumenicism. J Hered 86: 248-249.

Rose, A. H., R. H. Silversides, and O. H. Lindquist. 1975. Migration Flight by an Aphid, *Rhopalosiphum maidis*
(Hemiptera-Aphididae), and a Noctuid, *Spodoptera frugiperda* (Lepidoptera-Noctuidae). Can Entomol 107:
567-576.

Schlum, K. A., K. Lamour, C. P. de Bortoli, R. Banerjee, R. Meagher, E. Pereira, M. G. Murua, G. A. Sword, A.
E. Tessnow, D. V. Dillon, A. M. L. Ramirez, K. S. Akutse, R. Schmidt-Jeffris, F. N. Huang, D. Reisig, S. J.
Emrich, and J. L. Jurat-Fuentes. 2021. Whole genome comparisons reveal panmixia among fall armyworm
(*Spodoptera frugiperda*) from diverse locations. BMC Genomics 22: 179.

Seo, B. Y., G. S. Lee, J. Park, H. Xi, H. Lee, J. Lee, J. Park, and W. Lee. 2019. The complete mitochondrial
genome of the fall armyworm, *Spodoptera frugiperda* Smith, 1797 (Lepidoptera; Noctuidae), firstly
collected in Korea. Mitochondrial DNA B Resour 4: 3918-3920.

Sosa-Gomez, D. R., A. Specht, S. V. Paula-Moraes, A. Lopes-Lima, S. A. C. Yano, A. Micheli, E. G. F. Morais, P.
Gallo, P. R. V. S. Pereira, J. R. Salvadori, M. Botton, M. M. Zenker, and W. S. Azevedo. 2016. Timeline and
geographical distribution of *Helicoverpa armigera* (Hubner) (Lepidoptera, Noctuidae: Heliothinae) in Brazil.
Rev Bras Entomol 60: 101-104.

Sparks, A. N. 1979. Review of the Biology of the Fall Armyworm (Lepidoptera, Noctuidae). Fla Entomol 62: 82-
87.

Sun, X.-x., C.-x. Hu, H.-r. Jia, Q.-l. Wu, X.-j. Shen, Y.-Y. Jiang, and K.-m. Wu. 2019. Case study on the first
immigration of fall armyworm *Spodoptera frugiperda* invading into China. Journal of Integrative Agriculture
18: 2-10.

Tay, W. T., and K. H. J. Gordon. 2019. Going global - genomic insights into insect invasions. Curr Opin Insect Sci
31: 123-130.

Tay, W. T., P. J. Kerr, and L. S. Jermiin. 2016. Population Genetic Structure and Potential Incursion Pathways of
the Bluetongue Virus Vector *Culicoides brevitarsis* (Diptera: Ceratopogonidae) in Australia. PLoS ONE 11 (1):
e0146699.

Tay, W. T., B. James, and T. Walsh. 2021a. Final Report on Australian Bioassays. Prevention and preparedness
for fall armyworm (*Spodoptera frugiperda*) - Output 2, pp. 31. CSIRO/GRDC.

Tay, W. T., L. Kuniata, W. James, and T. K. Walsh. 2022a. Confirmation of *Spodoptera frugiperda* (Lepidoptera:
Noctuidae) in Papua New Guinea by molecular diagnostics of mitochondrial DNA COI gene. bioRxiv.

Tay, W. T., R. L. J. Meagher, C. Czepak, and A. T. Groot. 2022b. *Spodoptera frugiperda*: ecology, evolution and
management options of an invasive species. Annu Rev Entomol Revised Resubmitted.

Tay, W. T., M. F. Soria, T. Walsh, D. Thomazoni, P. Silvie, G. T. Behere, C. Anderson, and S. Downes. 2013. A
Brave New World for an Old World Pest: *Helicoverpa armigera* (Lepidoptera: Noctuidae) in Brazil. PLoS ONE
8(11): e80134.

Tay, W. T., R. V. Rane, W. James, K. H. J. Gordon, S. Downes, J. Kim, L. Kuniata, and T. K. Walsh. 2022c.
Resistance bioassays and allele characterisation inform *Spodoptera frugiperda* (Lepidoptera: Noctuidae)
management and introduction pathways in Asia and Australia. bioRxiv 2022.02.08.479273.

Tay, W. T., R. Rane, B. James, A. Gock, D. T. Hang, V. L. Nguyen, T. N. Khin, D. Amalin, C. Khonesavanh, F.
Muhammad, S. Annamalai, and T. Walsh. 2021b. Final report on Southeast Asian and Australian FAW
population genomics for biosecurity preparedness. Prevention and preparedness for fall armyworm
(*Spodoptera frugiperda*) - Output 2, pp. 25. CSIRO/GRDC.

Tay, W. T., T. K. Walsh, S. Downes, C. Anderson, L. S. Jermiin, T. K. Wong, M. C. Piper, E. S. Chang, I. B.
Macedo, C. Czepak, G. T. Behere, P. Silvie, M. F. Soria, M. Frayssinet, and K. H. Gordon. 2017.
Mitochondrial DNA and trade data support multiple origins of *Helicoverpa armigera* (Lepidoptera,
Noctuidae) in Brazil. Sci Rep 7: 45302.

Tay, W. T., R. V. Rane, A. Padovan, T. K. Walsh, S. Elfekih, S. Downes, K. Nam, E. D'Alencon, J. P. Zhang, Y. D.
Wu, N. Negre, D. Kunz, D. J. Kriticos, C. Czepak, M. H. Otim, and K. H. J. Gordon. 2022d. Global population
genomic signature of *Spodoptera frugiperda* (fall armyworm) supports complex introduction events across
the Old World. Commun Biol 5: 297.

Tonnang, H. E., S. A. Mohamed, F. Khamis, and S. Ekesi. 2015. Identification and Risk Assessment for
Worldwide Invasion and Spread of *Tuta absoluta* with a Focus on Sub-Saharan Africa: Implications for
Phytosanitary Measures and Management. PLoS ONE 10: e0135283.

Trisyono, Y. A., Suputa, V. E. F. Aryuwandari, M. Hartaman, and J. Jumari. 2019. Occurrence of heavy
infestation by the fall armyworm *Spodoptera frugiperda*, a new alien invasive pest, in corn in Lampung,
Indonesia. J. urnal Perlindungan Tanaman Indonesia 23: 156-160.

Valencia-Montoya, W. A., S. Elfekih, H. L. North, J. I. Meier, I. A. Warren, W. T. Tay, K. H. J. Gordon, A.
Specht, S. V. Paula-Moraes, R. Rane, T. K. Walsh, and C. D. Jiggins. 2020. Adaptive introgression across
semipermeable species boundaries between local *Helicoverpa zea* and invasive *Helicoverpa armigera*
moths. Mol Biol Evol 37(9): 2568-2583.

Vasimuddin, M., S. Misra, H. Li, and S. Aluru. 2019. Efficient Architecture-Aware Acceleration of BWA-MEM
for Multicore Systems. Int Parall Distrib P: 314-324.

Villesen, P. 2007. FaBox: an online toolbox for FASTA sequences. Mol Ecol Notes 7: 965-968.

Vu, T. P. 2008. Insect pests of turf grass, biology, ecology and the control of *Herpetogramma phaeoptralis*
(Guenée) in Hà Nội in Spring Summer 2008. MSc, Hà Nội Agriculture University, Vietnam.

Walsh, T. K., N. Joussen, K. Tian, A. McGaughran, C. J. Anderson, X. Qiu, S. J. Ahn, L. Bird, N. Pavlidi, J.
Vontas, J. Ryu, A. Rasool, I. Barony Macedo, W. T. Tay, Y. Zhang, M. E. A. Whitehouse, P. J. Silvie, S.
Downes, L. Nemec, and D. G. Heckel. 2018. Multiple recombination events between two cytochrome P450
loci contribute to global pyrethroid resistance in *Helicoverpa armigera*. PLoS ONE 13: e0197760.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis, Springer-Verlag New York.

Wilson, J. R. U., E. E. Dormontt, P. J. Prentis, A. J. Lowe, and D. M. Richardson. 2009. Something in the way
you move: dispersal pathways affect invasion success. Trends Ecol Evol 24: 136-144.

Worku, M., and Y. Ebabuye. 2019. Evaluation of efficacy of insecticides against the fall army worm *Spodoptera*
frugiperda. Indian Journal of Entomology 81: 13-15.

Wright, S. 1951. The Genetical Structure of Populations. Ann Eugenic 15: 323-354.

Wright, S. 1965. The Interpretation of Population-Structure by F-Statistics with Special Regard to Systems of
Mating. Evolution 19: 395-420.

Wu Q, Jian Y, and W. K. 2019. Analysis of migration routes of the fall armyworm *Spodoptera frugiperda* (J. E.
Smith) form Myanmar to China. Plant Protection 45: 1-6.

Yainna, S., W. T. Tay, E. Fiteni, F. Legeai, A.-L. Clamens, S. Gimenez, M. Frayssinet, R. Asokan, C. M.
Kalleshwaraswamy, S. Deshmukh, R. L. Meagher, C. A. Blanco, P. Silvie, T. Brévault, A. Dassou, G. J.
Kergoat, T. Walsh, K. Gordon, N. Nègre, E. d'Alençon, and K. Nam. 2020. Genomic balancing selection is
key to the invasive success of the fall armyworm. bioRxiv: 2020.2006.2017.154880.

Yang, X. M., Y. F. Song, X. X. Sun, X. J. Shen, Q. L. Wu, H. W. Zhang, D. D. Zhang, S. Y. Zhao, G. M. Liang, and
K. M. Wu. 2021. Population occurrence of the fall armyworm, *Spodoptera frugiperda* (Lepidoptera:
Noctuidae), in the winter season of China. Journal of Integrative Agriculture 20: 772-782.

Zhang, D. D., S. Y. Zhao, Q. L. Wu, Y. Y. Li, and K. M. Wu. 2021. Cold hardiness of the invasive fall armyworm,
Spodoptera frugiperda in China. Journal of Integrative Agriculture 20: 764-771.

Zhang, L., M. Jin, D. Zhang, Y. Jiang, J. Liu, K. WU, and Y. Xiao. 2019. Molecular identification of invasive fall
armyworm *Spodoptera frugiperda* in Yunnan Province. Plant Protection 45: 19-24.

Zhang, L., B. Liu, W. G. Zheng, C. H. Liu, D. N. Zhang, S. Y. Zhao, Z. Y. Li, P. J. Xu, K. Wilson, A. Withers, C. M.
Jones, J. A. Smith, G. Chipabika, D. L. Kachigamba, K. Nam, E. D'Alencon, B. Liu, X. Y. Liang, M. H. Jin, C.
Wu, S. Chakrabarty, X. M. Yang, Y. Y. Jiang, J. Liu, X. L. Liu, W. P. Quan, G. R. Wang, W. Fan, W. Q. Qian, K.
M. Wu, and Y. T. Xiao. 2020. Genetic structure and insecticide resistance characteristics of fall armyworm
populations invading China. Mol Ecol Resour 20(6): 1682-1696.

1040

1041

1042

1043