CRISPR/Cas9 and FLP-FRT mediated multi-modular engineering of the *cis*-regulatory

landscape of the bithorax complex of *Drosophila melanogaster*

Nikhil Hajirnis[1,+], Shubhanshu Pandey[1], Rakesh K Mishra[1,2,3,*]

1 – CSIR – Centre for Cellular and Molecular Biology, Hyderabad, India

2 – AcSIR – Academy of Scientific and Innovative Research, Ghaziabad, India

3 – Tata Institute for Genetics and Society (TIGS), Bangalore, India

+ - Present address – Department of Anatomy and Neurobiology, University of Maryland,
Baltimore, United States of America

*For correspondence – mishra@ccmb.res.in

Running Title: Engineering of the regulatory landscape of BX-C

Key words: bithorax complex, Hox genes, Regulatory Modules, Genome editing,
CRISPR/Cas9, FLP-FRT

Abstract

The Homeotic genes or *Hox* define the anterior-posterior (AP) body axis formation in

bilaterians and are often present on the chromosome in an order which is collinear to their

function across the AP axis. The expression pattern of *Hox* genes is attributed to the *cis*-

regulatory modules (CRMs) that regulate the genes in a segment-specific manner. In the

bithorax complex (BX-C), one of the two *Hox* complexes in *Drosophila melanogaster*, even

the CRMs consisting of enhancers, initiators, insulators, and Polycomb/trithorax response

elements are organized in order that is collinear to their function in the thoracic and abdominal

region. Much of these findings are based on the analysis of hundreds of mutations in the BX-

C. However, targeted genomic rearrangements comprising of duplications, inversions, etc., that

can reveal the basis of collinearity and the number of regulatory modules with respect to body

segments have not been reported. In the present study, we generated a series of transgenic lines

with the insertion of FRT near the regulatory domain boundaries, to shuffle the CRMs

associated with the posterior *Hox*, *Abd-B*, of the BX-C. Using these FRT lines, we created

several alterations such as deletion, duplication, or inversion of multiple CRMs to comprehend

their peculiar genomic arrangement and numbers in the BX-C.

Introduction

Eukaryotic gene regulation is a complex process facilitated by a combination of *cis*-regulatory elements (CREs) and *trans*-acting factors (TAFs). The fine-tuning of expression of a gene is performed by several *cis*-regulatory elements that include initiator elements, enhancers, repressors, Polycomb/trithorax response elements and promoter targeting sequences (Narlikar and Ovcharenko 2009; Crocker et al. 2016; Iampietro et al. 2010; Gaston and Jayaraman 2003; Zhou and Levine 1999; Ringrose and Paro 2004). In *Drosophila melanogaster* genome, these elements are often populated in 15-20 Kb regions that regulate the activity of associated genes. Such regions where several CREs populate together and are responsible for the expression of a gene in a cell, tissue, or segment-specific manner are called *cis*-regulatory modules (CRMs) (Ho et al. 2009; Starr et al. 2011; Bekiaris et al. 2018). Plentiful studies have been dedicated towards understanding the role of each of these elements individually (Polychronidou and Lohmann 2013; Kyrchanova et al. 2011; Cléard et al. 2006; Akbari et al. 2008; Narlikar and Ovcharenko 2009). However, only a handful of them addressed the collective functioning of the CRMs (Lelli et al. 2012; Zinzen et al. 2009; Starr et al. 2011; Hajirnis and Mishra 2021). One such region in the *Drosophila* genome that heavily relies on the collective functioning of the CRMs, includes a set of developmentally important genes, called *Hox*. These genes code for transcription factors that bind to the DNA in a sequence-specific manner. The expression pattern of these genes defines the anterior-posterior body axis of a developing bilaterian embryo (Foronda et al. 2009; Hajirnis and Mishra 2021).

Homeotic genes or *Hox* were discovered in *Drosophila melanogaster* wherein they are present in a spatially colinear manner on the chromosomes (Lewis 1978). That is, the genes are arranged in a complex and, the genes that are present at one end of the complex are responsible for the development of the anterior part of an embryo while the genes present at the other end are

responsible for the development of the posterior segments. The *Drosophila Hox* genes are

arranged in two clusters, the Antennapedia complex (ANT-C) and the bithorax complex (BX-

C) (Kaufman et al. 1990; Peifer et al. 1987).

The posterior-most *Hox* gene, *Abd-B* is regulated by four distinct CRMs, *infradbominal-5*

(*iab5*), *iab6*, *iab7,* and *iab8/9*. Notably, in addition to the arrangement of *Hox*, even the CRMs

of the BX-C genes are spatially arranged in a manner colinear to the A-P body axis that they

determine (Karch et al. 1985; Maeda 2006). Thus, *iab5, iab6, iab7* and *iab8/9* are responsible for

the development of the four abdominal segments, A5, A6, A7 and A8/9.

In addition to the spatially collinear arrangement of the modules, the numbers of the modules

also corroborate with the number of the segments that they provide identity; the four CRMs

drive the identity of four abdominal segments.

Each of the *Abd-B* modules is separated by chromatin domain boundaries in the order *MCP*

that separates *iab4* and *iab5* (Mihaly et al. 1998), *Fab6* separating *iab5* and *iab6* (Pérez-Lluch et

al. 2008), *Fab7* separating *iab6* and *iab7* (Gyurkovics et al. 1990a), and, *Fab8* that separates *iab7*

and *iab8* (Barges et al. 2000). A deletion of several regions of the CRM causes a loss of function

of the associated gene that leads to anteriorization of the respective segment (Celniker et al.

1990). For example, the deletion of *iab6* elements leads to a transformation of A6 to A5 (Galloni

et al. 1993). The module that becomes activated in a segment, remains activated even in the

posterior segment (Bowman et al. 2014; Maeda and Karch 2015). Thus, when a posterior module

is not available in case of a deletion, the adjacent anterior module defines the segment identity.

Such flies have two copies of A5 followed by A7 (Mihaly et al. 2006). On the contrary, the

deletion of boundaries, such as *Fab7*, would lead to the posteriorization of A6 to A7. This is

due to ectopic derepression of posterior module *iab7* that prevails over the anterior module

iab6, and drives the identity of A6. The fly, thus, has A5 followed by two copies of A7

(Gyurkovics et al. 1990b; Mihaly et al. 1998; Iampietro et al. 2010).

Although there are several inversions and a translocation reported for different regions of the

BX-C (Celniker and Lewis 1993; Ho et al. 2009; Hendrickson and Sakonju 1995; Stuttgart and

Herberth 1988; Gligorov et al. 2018), there are no known cases of targeted duplications or

inversions of CRMs concerning their effect on adult segments. Further, many questions

concerning the regulation of the BX-C remain unanswered (Bender 2020). For instance, how

well do the CRMs rely upon each other for the accessibility of the chromatin to the transcription

factors? Is the relative positioning and order of the CRMs important to respond to the upstream

spatial cues? What is the relevance of the defined number of modules in the system and can

the system accommodate additional modules? The sequential opening of the chromatin

represented by the loss of repressive H3K4me3 marks is observed across the BX-C CRMs

through anterior to posterior parasegments. If the order or number of CRMs are altered, would

the sequential opening model maintain its status-quo?

In order to answer such questions, we generated transgenic lines with FRT insertions in targeted

regions across the *Abd-B* CRMs. these lines could be used to create multiple deletions,

duplications, and inversions for the desired set of CRMs (Fig. 1A). We modified the *cis-*

regulatory landscape of the posterior-most BX-C gene, *Abd-B*, to generate useful resources that

will help us advance our understanding of the role of *cis*-regulatory modules in regulating Hox.

Further, our study also features a set of *in-vivo* systems to comprehend the arrangement and

number of *cis*-regulatory modules in the BX-C. This system can be repurposed to shuffle the

arrangement and number of regulatory domains in order to understand the functioning of CRMs

in addition to our existing knowledge about the individual regulatory elements.

Results

Generation of FRT knock-in transgenes

The BX-C is heavily decorated with binding sites for transcription factors and chromatin remodelers (Négre et al. 2011). Further, a single nucleotide transition of a repressor binding site was shown to have an effect on the ectopic derepression of *iab5* in A3 leading to a prominent A3 to A5 homeotic transformation of the segment (Ho et al. 2009). Thus, the BX-C CRMs need to be tightly regulated and leave narrow margins to make amendments in the genome without affecting *cis-* motifs. Therefore, to insert FRT sites without any further perturbation, we utilized the available data for experimental validations of the BX-C and ChIP data on modENCODE to connote the *Abd-B cis*-regulatory landscape into distinct *cis*-regulatory modules, *iab5* through *iab8* (Fig 1A). The boundaries separating these domains were also annotated based on experimental results from previous studies (Iampietro et al. 2008; Mishra and Karch 1999; Postika et al. 2018; Kyrchanova et al. 2015). Further, regions marking P/TREs distal to the boundaries were annotated based on earlier experimental validations and previously reported PRE mapping tool, PRE mapper (Pérez-Lluch et al. 2008; Mishra et al. 2001; Singh and Mishra 2015; Postika et al. 2021a; Srinivasan and Mishra 2020). The target sites for FRT insertions were selected such that the FRT lands between a boundary and a CRM without perturbing the boundary-PRE combination (Fig. 1A, Supplemental Table S1).

The FRT sequences were cloned in different orientations with required homology arms (Supplemental Table S2) for CRISPR-Cas9 mediated homology-directed repair (Port et al. 2015). The FRTs are inserted between *MCP-iab5*, *Fab6-iab6*, *iab7-Fab8,* and *Fab8-iab8* in

different orientations as shown in Fig 1A. For example, the line *MP_F_i5* has FRT inserted between *MCP* and *iab5* with the direction of FRT towards *iab5*. Similarly, the *i6_F_F6* line has an insertion of FRT between *Fab6* and *iab6* with the direction of FRT towards *Fab6*.

The FRT insertion is validated by amplifying the region using primers eccentric to the FRT site. Further, the FRT contains an *Xba-I* restriction digestion site. Thus, the amplified products for positive transgenes are digested with the endonuclease and render two distinct bands on agarose gel electrophoresis (Supplemental Fig. S1, Supplemental Table S6). The FRT insertions are further confirmed by sequencing.

Other than the flies that have FRTs inserted in a single locus, we also generated lines that have a couple of FRTs inserted in the same as well as opposite orientations. The single FRT lines can be used as landing platforms for various other transgenic assays and can also be repurposed to generate targeted deletions and duplications of the CRM-boundary combinations in the locus (Horn and Handler 2005; Phan et al. 2017; Golic and Lindquist 1989). The double FRT with the FRTs in the same orientation can be used to generate deletion in *-cis* or duplication in *-trans*(Golic et al. 1997). The duplicated line can further be used to generate subsequent modules of the *cis*-regulatory landscape by repurposing along with single or double FRTs. On the other hand, the double-FRT line with the two FRTs in opposite orientation can be used to generate inversion of the locus to re-arrange the order of the CRMs and boundary (Golic et al. 1997). Some of the combinations using single FRTs and possibilities of using double FRTs are mentioned in the following sections. A summary of all the FRT lines and genomic rearrangements generated in the study is presented in Fig. 1.

Deletion of *iab5-Fab6* reveals the non-autonomous function of CRMs

The number of *Abd-B* CRMs corroborates the number of abdominal segments. Thus, we

rationalized that the deletion of information about the development of one of the segments can

cause an impact on the number of segments formed. Although larger deletions in the region

were shown to affect only the identity of segments but not the number, the exact effect in adults

is not clear. This is especially because many of the larger deletions including *abd-A* and *Abd-*

B CDS lead to embryonic lethality in homozygous conditions. Thus, removing a combination

of CRM along with boundary will leave the locus with altered modularity characterized by

lesser numbers of CRM-boundary combinations. Towards this, we generated a line that lacks

a ~15.14 kb region spanning *iab5* and *Fab6* (R6.0 – 3R: 16,872,062..16,887,207). Thus, the

fly is left with three CRMs instead of four in wild-type animals (Fig. 2A).

To obtain the line lacking *iab5* and *Fab6*, we crossed *hsFlp*; *MP_F_i5* (FRT inserted between

MCP and *iab5* with the direction of *FRT* towards *iab5*) with *hsFlp*; *F6_F_i6* (FRT inserted

between *Fab6* and *iab6* with the direction of FRT towards *iab6*), see Fig 2A. Both lines express

Flippase recombinase (FLP) under the influence of heat-shock inducible promoter (*hs*). The

trans-heterozygous progenies with one allele with an FRT between *MCP* and *iab5* while the

other with an FRT between *Fab6* and *iab6* were given heat shock at 37°C for 90 min at an

interval of 24 hrs. Heat shocks were given from the late second instar larval stages until the

late pupal stages of *Drosophila* development. FLP activity was confirmed as described

previously (Pignoni et al. 1997); see methods and supplementary information (Supplemental

Fig. S2) for details.

Once the heat-shocked animals eclosed (G1), they were mated with third chromosome

balancers to prevent genetic recombination of target loci. The progenies from these crosses

(G2) were pooled in batches of 10 and were again crossed with third chromosome balancers.

Once enough activity was observed in the vial, the flies were screened via PCR for the desired

deletion. The PCR products of flies showing amplification of deletion specific regions were

confirmed via Sanger sequencing (Supplemental Fig. S3).

We observed a striking anteriorization of A5 and A6 into copies of A4 in the deletion mutants

Del_i5-F6. The A5 and A6 in wild-type males are characterized by complete melanization

whereas, the A4 has melanization in the posterior end. The homozygous fly for the deletion of

iab5-Fab6 shows a transformation of A5 to A4 characterized by the loss of melanization.

Moreover, even the A6 was transformed into a copy of A4 characterized by the loss of

melanization and appearance of ventral bristles which are otherwise absent in a wild-type male

(Fig. 2B-C).

Previous studies showed the functioning of CRMs in an autonomous manner wherein the

autonomy is largely maintained by the boundary elements demarcating the modules (Hagstrom

et al. 1996; Maeda and Karch 2015). Deletions in *iab6* or *iab7* do not affect the functioning of

each other in the associated segments A6 or A7. For instance, the deletion of *iab6* regions

would cause A6 to transform into a copy of A5 but does not have an impact on A7 or A8

(Iampietro et al. 2010). On the other hand, the deletion of a boundary would ectopically activate

the posterior CRM. For example, the deletion of *Fab6* would lead to the transformation of A5

into a copy of A6. Akin to that, the deletion of *Fab7* will transform A6 into A7 (Mihaly et al.

1998). Therefore, we predicted that the deletion of *iab5-Fab6* will lead to the transformation of

A5 into a copy of A6, a phenotype largely dominated by the deletion of *Fab6*.

In contrast to our expectations, we observed both A5 and A6 transforming into copies of A4 in

males and females (Fig 2B-E). Since the deletion of *Fab6* does not impact *iab6* activity, we

reasoned that the regions in *iab5* are responsible for *iab6* activation, thereby defying the

autonomy of *iab6*. Interestingly, Postika et al. reported a similar effect observed due to

deletions of several regions within *iab5*. Since the deletion we generated encompassed all the

regions mentioned by Postika et al., we see a similar phenotype with a clean transformation of

A5 and A6 into copies of A4 (Postika et al. 2021b, 2021a). Thus, although the system is left with

modules responsible for the formation of A6 through A8; the activators in anterior CRM, *iab5*,

are required for the execution of the spatial code in A6 but not in A7 or A8.

Deletion of *iab6-iab7* transforms A6 and A7 into copies of A5

The transgenic assays and genetic interactions from previous studies have revealed that

boundaries interact with each other (Maeda and Karch 2007; Kyrchanova et al. 2011; Singh and

Mishra 2015). Therefore, we deleted a region spanning *iab6* through *iab7* thereby juxta

positioning *Fab6* next to *Fab8* (Fig. 3A). Towards this, we generated a line with two FRTs

inserted in the same orientation in two different loci within *Abd-B cis*-regulatory landscape.

One of the FRTs was inserted between *Fab6* and *iab6* with the direction of FRT towards *Fab6*

(*i6_F_F6*). The other FRT was knocked-in between *iab7* and *Fab8* with the orientation of FRT

towards *iab7* (*F8_F_i7*) as shown in Fig. 3A. Although the FRTs were inserted in *cis*- and a

homozygous fly could have rendered us the required deletion, we are also open to the

possibility of the recombination to have occurred in *-trans* as indicated in Fig. 3A.

The FLP mediated recombination under the influence of a heat-shock inducible promoter was

carried out as described in the previous section. To molecularly validate the deletion, the

genomic DNA of the putative mutants was isolated and amplified using a combination of

primers specific for the deletion locus upon recombination; an expected juxta positioning of

Fab6 and *Fab8*. The repositioning would amplify a specific product with the forward primer

for screening FRT between *Fab6* and *iab6* (F6i6_ScrF) and the reverse primer for screening

FRT between *iab7* and *Fab8* (i7F8_ScrR). A resultant product of 974 bp confirms the deletion

event as shown in Supplemental Fig. S4. The endogenous locus at the approximate junction of

Fab6 and *iab6* was chosen as a control locus. This region was amplified in the genomic DNA

of a wild-type CS fly but not the deletion mutant (Supplemental Fig. S4C). The amplified

product for deletion mutants was confirmed by sequencing (Supplemental Fig. S4D). The total

length of the deleted region spans 31.374 kb from 3R: 16,887,208 to 3R: 16,918,581 (Genome

assembly R6.0).

We observed a phenotype as was expected with the deletion of the CRMs *iab6* and *iab7* that

are responsible for determining the identities of A6 and A7. There is a complete loss of function

in the homozygous flies carrying the desired deletion as characterized by the complete

transformation of A6 and A7 into copies of A5 (Fig. 3B-E). Additionally, the males also show

genital rotation of varying degrees and are sterile. The deletion of intermittent boundary *Fab7*,

and, juxtapositioning of *Fab6* and *Fa8* do not cause an effect on the functioning of *iab5*. The

phenotype is also apparent in females wherein a wild-type female has 8-10 narrow groups of

drooping bristles in A7. In the mutant females, A7 is broader, and a complete set of sternites

appear on the ventral surface of the segment (Fig. 3D-E). The juxta positioning of *Fab6* and

Fab8 does not seem to have an effect on regulation of *Abd-B* via *iab5*.

Duplication of *iab6-iab7* renders phenotype dominated by the posterior module *iab7*

The number of CRMs in the BX-C is equivalent to the number of segments that they provide

identity. Hence, it is important to understand the functional correlation of the number of CRMs

and their role in providing identity to respective segments. Towards this, we generated a mutant

line with duplication in *iab6* through *iab7* such that the modularity is lost. The duplication

spans 31.374 kb (3R: 16,887,208..16,918,581). This arrangement juxtaposes an additional copy

of *iab6* distal to *iab7* (Fig. 4A) without a boundary. Using the double FRT line used for

obtaining the deletion of *iab6-iab7*, we generated a recombinant line with the duplication of

the same region as indicated in Fig. 4A.

To obtain the duplication line, we screened the recombinants from the progenies of flies set for

obtaining the deletion of *iab6* through *iab7* in the previous section (Supplemental Table S9-

S10). The expected recombination of the two loci in *-trans* is depicted in Fig. 4A with

dichromatic representations of the parent and prospective recombinant alleles. Note that the

duplication allele has three FRTs. One of the FRTs is derived from recombining FRTs between

Fab6-iab6 and *iab7-Fab8*. The other FRTs are present from the parent allele. This line is a

novel playground to recombine various transgenic animals having FRTs at different loci in the

Abd-B cis-regulatory landscape. The three FRTs can be repurposed differently to alter the

modularity of the locus in varied manners.

For the molecular validation, the genomic DNA of putative recombinants was isolated and

amplified by primers specific to the duplication locus. The forward primer to screen FRT

between *iab7* and *Fab8* (i7F8_ScrF) and the reverse primer to screen FRT between *Fab6* and

iab6 (F6i6_ScrR) amplified a specific 534 bp product indicating the presence of duplication

allele. The endogenous *Fab6-iab6* and *iab7-Fab8* loci were used as controls. The amplified

products were confirmed using sequencing (Supplemental Fig. S5).

The fly with the duplication of *iab6* through *iab7* (*Dp_i6-i7*) has multiple changes in the

arrangement of modules. The fly has two *iab6* modules, one flanked by *Fab6* and *Fab7* as in

wild-type conditions, and the other is juxtaposed next to *iab7* without intermittent boundary.

We observe that there is a partial posteriorization of A5 in adult males as suggested by the

decrease in the number of bristles in sternites and partial morphological changes in the dorsal

A5 as indicated by dotted shape in Fig. 4C. Further, the A6 identity is completely transformed

into a copy of A7 towards the dorsal side. However, on the ventral end, A6 shows a partial

anteriorization as suggested by the presence of a few bristles as well as loss of sclera formation.

There are no apparent changes in the male A7 of the transgenic animals despite two copies of

iab7. A similar effect of A6 to A7 transformation is also evident in females (Fig. 4B-E).

The presence of extra modules was particularly interesting to understand the effect of relative

positioning of *iab6* and i*ab7*. We thus expected a mixed gain and loss of function of A6 as well

as A7. We were also open to the possibility of observing a segment with novel features since

the duplicated *iab7-iab6* formed a larger module flanked by *Fab7*. Moreover, the presence of

extra modules would also change the relative positioning of *iab5* with respect to the *Abd-B*

promoter and thus an effect in A5 was also expected. However, a strong A6 to A7 and further

posteriorization of A5 indicate the dual copies of *iab7* were dominant over the two copies of

anterior module *iab6* as well as the wild-type *iab5* irrespective of relative positioning.

Duplication of *iab6-Fab8* alters modularity and renders variable phenotype

In the previous duplication, the modularity was perturbed due to the absence of a boundary

element between *iab7* and *iab6*. Therefore, to preserve modularity, we duplicated the 34.776

kb (3R: 16,887,208..16,921,983) region from *iab6* through *Fab8*, keeping the modularity

intact. Thus, each CRM is still flanked by a combination of boundary-PRE as shown in Fig. 5.

Further, to keep one of the CRMs as an internal control, we left *iab5* intact and duplicated *iab6*

through *Fab8*.

To obtain a fly with proposed duplication, we genetically crossed the flies with FRT insertions

between *Fab6-iab6* and *Fab8-ia8* to obtain a trans-heterozygous line. These flies were in the

background of *Flp* recombinase expressed under the control of a heat-shock inducible promoter

as described earlier. A brief schematic of flies used and the duplication upon recombination is

depicted in Fig. 5A. For details, see methods and Supplemental Table S9-S10.

Next, to validate the duplication event molecularly, the genomic DNA of prospective

recombinants was isolated and amplified by primers specific to the duplication locus

(F8i8_ScrF and F6i6_ScrR). The intact loci at *Fab6-iab6* and *Fab8-iab8* were amplified as

controls. Only the flies possessing the desired duplication showed an amplified product

(Supplemental Fig. S6C). Genomic DNA from CS flies was used as a control. Unlike the

duplication locus that was amplified only in the positive recombinants, the *Fab6-iab6* and

Fab8-iab8 loci were amplified in both the duplication and CS genomic DNA. The products

were later confirmed by sequencing (Supplemental Fig. S6C-D).

In the *Dp_i6-F8* flies, the combination of CRM-boundary is intact. Thus, we expected a mixed

gain and loss of functions in A6 and A7. Since *iab6* and *iab7* are regulators of *Abd-B*, an effect

could also have been obtained in A5 and A8. The extra copies of *iab6* and *iab7* were modularly

flanked by *Fab6* and *Fab7* akin to the wild-type scenario. Surprisingly, we observed a variety

of phenotypes with perturbations ranging from disturbances in A1 to a normal-looking fly

similar to wild-type (Fig. 5B-H).

We classified the phenotypes into six different categories as follows: Category A (Cat-A)

mutants show the most extreme phenotypes with all abdominal segments from A1 to A8

severely perturbed. Several flies have disturbances in left-right or dorsal-ventral patterning.

These flies are often sterile (Fig 5C). Cat-B animals showed disturbances in segments A5 to

A8 (Fig. 5D). The rationale for this category is the role of *iab6* and *iab7* to regulate *Abd-B* that

defines A5-A8. Any segment perturbed anterior to A5 was not counted in this category, and

instead, was noted in the previous category, A. The Cat-C mutants displayed disturbances in

A6 as well as A7 (Fig. 5E). Since the duplicated *iab*s are responsible for providing identities

to A6 and A7, flies having perturbations in only these segments were considered a separate
category. Cat-D mutants had a perturbation in only one of the abdominal segments. Often these
perturbations were restricted to A6 or A7 (Fig. 5F). In the next category, Cat-E mutants display
a slight perturbation of dorsal melanization and often rendered "wavy" segmental boundaries
(Fig. 5G). The Cat-F mutants show a wild-type phenotype with no observable change in the
cuticle or ventral bristle patterning (Fig. 5H).

Both males and females showed the mentioned pattern. However, females were better to
observe via imaging owing to clear demarcations of seven abdominal segments followed by
genitalia. Also, the cuticles of these animals were extremely brittle, and therefore, the abdomen
was directly imaged under the stereomicroscope (Fig. 5B-H; see methods for details).

All the classes of mutants were confirmed to have the same genotype molecularly. Further, the
inbreeding of mutants from all the phenotypes, including Cat-F (normal segments), rendered
progenies with varying phenotypes again. The development of progenies with inconsistent
phenotypes is true for heterozygous and homozygous flies from these categories. Hence, we
selected only category F adults and set up a breeding experiment to check the distribution of
progeny with different phenotypes.

We observed almost half the population for homozygous (45.24%) and heterozygous (41.35%)
progenies were normal (Cat-F). In contrast, the remaining categories had a smaller number of
flies in the population (Fig. 5I). Interestingly, ~25% heterozygous (24.67%) and ~17%
homozygous (16.52%) flies showed phenotypes grouped under category A. The heterozygous
flies showed 1.5 times a greater number of progenies belonging to Cat-A than homozygous
(Fig 5I). Our results suggest that more heterozygous flies tend to be severely affected by the
duplication of CRMs (Supplementary Data).

Thus although the duplication of *iab6-Fab7* perturbs the genomic modularity of the locus

(previous section), the segmental boundaries are intact. However, duplication of modules

keeping genomic modularity intact (*Dp_iab6-Fab8*) disrupts the segmental boundary in a

manner that requires further investigations to understand the BX-C better.

The inversion of *iab5* through *iab7* provides mixed and altered identities of the associated

segments, A5, A6 and A7

The CRMs of the BX-C are present in a spatially colinear manner. The three CRMs, *iab5*, *iab6*

and *iab7* regulate *Abd-B* to provide identities to A5, A6 and A7 respectively. The specific

ordering of the CRMs can be consequential towards co-regulation of the associated gene (Mateo

et al. 2019; Maeda and Karch 2015; Hajirnis and Mishra 2021). One way to test this hypothesis is

to invert the order of these regions. Towards this, we generated a 46.520 kb

(3R:16,872,062..16,918,581) inversion of *iab5* through *iab7* (Fig 6).

We generated a fly having two FRTs in the opposite orientations inserted between two loci:

MCP-iab5 and *iab7-Fab8*. The FRT inserted between *MCP-iab5* was oriented towards *iab5*,

while the FRT inserted between *iab7* and *Fab8* was oriented towards *iab7*. These flies were

crossed with a third chromosome balancer, and the heterozygous larvae were given heat shock

as described earlier (Supplemental Table S11-S12). The eclosing adults post-heat shock (G0)

were crossed with balancers in anticipation that their gametes had undergone the desired

recombination (Fig. 6A). The progenies (G1) emerging from these crosses were further crossed

with third chromosome balancers. The G1 flies, after three to four days of mating, were

sacrificed for molecular screening.

To molecularly screen for the inversion event, the forward primer for *MCP-iab5* locus

(Mi5_ScrF) and the forward primer for *iab7-Fab8* locus (i7F8_ScrF) were repurposed. The

forward primer for the *iab7-Fab8* locus is now used as a reverse primer for *MCP-iab7*

recombined locus. Similarly, the recombined locus of *iab5-Fab8* was screened by repurposing

the reverse primer to screen FRT at *MCP-iab5* (Mi5_ScrR) locus with the reverse primer to

screen FRT at the *iab7-Fab8* region (i7F8_ScrR) as explained in Supplemental Fig. S7. A

specific product upon repurposing the primers is obtained only for the flies that had an

inversion event. The endogenous loci of *MCP-iab5* and *iab7-Fab8* were amplified only in the

CS fly and not in the inversion line. The recombined loci were further confirmed by sequencing

(Supplemental Fig. S7).

An expected outcome of this inversion was a wild-type phenotype for the mutant animal if all

the elements are perfectly exchangeable. The other direct possibility was that the "code" for

segmental identity resides in the order of CRMs, and therefore reversing their order might

reverse the identities of abdominal segments. In this case, we expect a fly that forms A4,

followed by A7, A6 and A5 in the anteroposterior order. However, we noted flies with different

phenotypes for different segments (Fig 6B-E).

We must also consider that the order of CRMs was reversed, and their relative positioning was

also exchanged. For instance, the native position of *iab5* now has the presence of *iab7* while

iab5 replaces *iab7* as it is juxtaposed next to *Fab8* (Fig. 6A). On the contrary, the *iab6* is in the

same position as before but is present in an opposite orientation. The re-orientation of *iab6* also

includes the change in directionality of flanking boundaries *Fab6* and *Fab7*. Notedly the

boundaries function in an orientation-dependent manner (Zhou et al. 1996; Hogga and Karch 2002;

Postika et al. 2018). Therefore, altering the relative positioning of boundaries can also contribute

to the phenotypes obtained.

The A5 segment of the flies shows partial gain and loss of function. The numbers and density

of ventral bristles decrease, indicating a gain of function to A6 (Fig 6C-D). Occasionally,

several males also show partial loss of melanization in A5, indicating partial transformation

into A4. The mixed gain and loss of function phenotypes indicate a cell-type-specific behavior

of the modules within a particular segment of the mutant animal. The ventral A5 also develops

a sclera similar to A6, indicating slight posteriorization (Fig. 6D). The PREs associated with

the boundaries in the BX-C provide temporal and cell-type-specific regulation of the adjacent

CRMs (Ringrose and Paro 2004). The absence of any major PRE associated with a boundary

near *iab5* partly explains the incomplete anterior- and posteriorization upon inversion.

Next, even though *iab6* is in the same position relative to wild-type condition, it shows a partial

gain of function transformation into A7 as indicated by narrowing of the dorsal melanization

and sternite in the males (Fig. 6C). The effect is also clearly observed in A6 of the females

wherein the ventral bristles attain the identity of the A7 (Fig. 6D-E). Notedly, although the *iab6*

remains in the same relative order concerning adjoining boundaries and CRMs, the direction

of the module is opposite. It has been previously shown that the direction of the BX-C

boundaries including *Fab6* and *Fab7* is crucial to performing the insulator bypass activity

(Zhou et al. 1996; Postika et al. 2018). In the inversion mutant, since both the boundaries are

inverted, *iab6* is misregulated. The studies on orientation were done individually on different

elements. The rearrangement in the inversion locus is a complex interplay between many of

them. Therefore, the mutants should be probed deeper, especially for the chromatin

conformations of the CRMs to understand the interaction of different elements upon inversion.

Further, the A7 shows an opposite transformation when compared to A6. A distinct dorsal

melanized segment is present in males in the position of A7, thus indicating anteriorization of

the segment into a copy of A6 (Fig. 6C). However, the segment is not completely transformed

into A6. A similar effect is evident in the A7 of adult females by the appearance of extra bristles

and broadening of the sclera (Fig. 6E). The male flies also have a genital rotation of varying

degrees and is not consistent. Additionally, homozygous flies with the inversion of *iab5*

through *iab7* are sterile.

As mentioned earlier, an inversion such as this includes multiple disruptions of the regulatory

landscape of the associated gene and although individual elements have been dissected earlier,

the complex interplay of multiple elements is interesting to probe in such scenarios.

Discussion

Homeotic genes or *Hox* code for sequence-specific transcription factors that define the anterior-

posterior body axis of a developing bilaterian embryo (Hajirnis and Mishra 2021). The *Hox* genes

were discovered in *Drosophila melanogaster* wherein they display a striking property of spatial

collinearity (Lewis 1978). That is, the genes are arranged in a complex and, the genes that are

present at one end of the complex are responsible for the development of the anterior part of

an embryo while the genes present at the other end are responsible for the development of the

posterior regions. The posterior *Hox* complex in *Drosophila melanogaster*, the bithorax

complex presents us with a unique opportunity to study the relevance of the higher-order

arrangement of the *cis*-regulatory modules (CRMs) of the homeotic genes (Maeda 2006). The

three genes of the BX-C: *Ubx*, *abd-A* and *Abd-B* are regulated by nine *cis*-regulatory modules

abx/bx and bxd/pbx regulating *Ubx* in posterior T2 through A1, *iab2* through *iab4* regulating

abd-A in A2 through A4, and, *iab5* through *iab8* regulating *Abd-B* in A5 through A8 (Maeda

and Karch 2015). The *iab*s are arranged in a spatially collinear manner with respect to the

segments that are affected upon mutating them. Also, the numbers of the *iab*s corroborate with

the number of abdominal segments formed in the fly. In the present study, we generated a set

of FRT transgenes to alter the *cis*-regulatory landscape of the posterior-most *Hox* gene *Abd-B*.

These lines can be used independently to insert test elements in the fly or can be repurposed to

alter the regulatory landscape in *Drosophila*.

The role of many of the *cis*-regulatory elements or modules is known in their native structure

(Iampietro et al. 2010, 2008; Zhou and Levine 1999; Cléard et al. 2006). Notedly, many of the

existing studies regarding genome manipulation of a similar kind are limited by the status-quo

of the targeted manipulations, that is, the alterations lack the flexibility to incorporate novel

features or re-shuffle the existing modules (Li et al. 2015; Guo et al. 2015; Fabre et al. 2017). In

the current study, the use of FLP-FRT at modular junctions makes the system more dynamic

and robust. The alterations in the genome are quickly interchangeable. In fact, in the case of

duplication of *iab6-iab7*, the fly has three FRTs and thus it can be repurposed to develop a

plethora of downstream combinations of modules. For instance, if the *iab7-iab6* fused domain

is behaving like a "super-module", an additional such module can be introduced in the system

using FLP mediated recombination using the lines we generated. Furthermore, each of the lines

can be followed individually in different directions. For example, it is interesting to probe the

chromatin landscape and chromatin interaction in the fly bearing an inversion of *iab5* through

iab7. Previous studies have shown sequential derepression of the BX-C *cis*-regulatory modules

from anterior to posterior segments; a feature called open for business model of the BX-C

regulation (Bowman et al. 2014; Maeda and Karch 2015). However, one key aspect to probe in the

sequential opening model is the ability of these modules to sense the spatial cue of accessibility

(Kyrchanova et al. 2015). Therefore, in the case of inversion, segment-specific methylation

marks would reveal novel aspects of sequential opening. Moreover, understanding segment-

specific interactions of enhancers and promoters via techniques like ORCA and Hi-C will shed

light upon the functioning of these modules in the altered scenario (Mateo et al. 2019).

In cases such as *Dp_iab6-Fab8*, where the modularity is intact with respect to the positioning

of CRMs and boundaries but the number of modules has been altered; the variability in

phenotype is interesting to probe. For instance, the variability can arise from the change in

dosages of the non-coding RNAs (ncRNA) generated from the *cis*-regulatory locus of *Abd-B*

(Garaulet and Lai 2015; Gummalla et al. 2014). Or, the variability could be caused by ectopic

transvection happening in different cell types in a stochastic manner (Vazquez et al. 2006;

Hendrickson and Sakonju 1995). One way to probe the latter is to investigate the role of epigenetic

modifiers such as PcG and trxG proteins to modulate the phenotypic variability (Singh and

Mishra 2015). Genes like *Zeste* are known players of transvection (Sipos et al. 1998; Birve et al.

2001). The effects of mutations in such genes can be probed to understand the nature of

variability present in the system. The other possibility pertaining to the differential dosages of

ncRNAs can be probed by *in-situ* hybridization (Arib et al. 2015). However, the definitive

changes in the ncRNAs corresponding with the adult phenotypes require further investigation.

With the advancement of genomics and gene-editing techniques, many such mutations as

presented in the current study can be generated to understand the evolutionary significance of

the CRMs and *Hox* arrangement in different organisms. Additionally, comprehending the

epigenomic landscape and DNA looping becomes important to grasp the depths of CRM

functioning in such conditions.

In summary, we generated a set of transgenic flies that can alter the modularity of *cis*-regulatory

domains in the BX-C of *Drosophila melanogaster* and open avenues to explore the

fundamental basis of body axis formation. The novel and unanticipated phenotypes obtained

in the study clearly demonstrate that our current understanding of the mechanisms of regulation

of bithorax complex is substantially inadequate. This study also shows the potential of targeted

re-arrangement of the modules to elucidate the role of the genomic landscape of the bithorax

complex. These resources will shed light on important aspects of *Hox* regulation including but

not limited to finer details of the order of the CRMs, chromatin conformations, enhancer-promoter interactions, *modus operandi* of regulatory boundaries, or causal relation between histone modifications and expression of associated Hox. Genetic interactions of the novel modules will provide crucial insights underlying their concerted functioning and open a new paradigm for the business of the bithorax complex.

Materials and Methods

Primer designing for FRT insertion

The primers for targeting Cas9 were obtained after submitting query sequences for different regions of the BX-C at http://crispor.tefor.net/. *Drosophila melanogaster* BDGP release 6 (R6.0) was selected as the reference genome, and 20bp crRNA with *S. pyogenes* Cas9 5'-NGG3' PAM was curated for obtaining targets. From the list of suitable gRNAs, sequences with an MIT specificity score of 97 or more (preferably 100), CFD specificity score of 97 or more (preferably 100) and a Lindel score of 80 or more were selected as preferred guides. Due care was taken to select primers with the least number of off-targets as indicated in the relevant column.

Upon successful selection of guides, 1000-1200 bp upstream and downstream region of the cut site was selected as a query to design primers for amplification and cloning of homology arm for donor constructs in https://www.ncbi.nlm.nih.gov/tools/primer-blast/. For all the FRT insertions, the centromeric proximal homology arm to the cut site was denoted as the left homology arm (LH), while the distal homology arm was denoted as the right homology arm

(RH). The reverse primer for LH was selected as 20-24 bp immediately centromeric proximal

to the Cas9 cut site (3 bases upstream of PAM). Similarly, the forward primer for RH was

selected as 20-24 bp immediately distal to the cut site. Keeping the LH reverse primer as

constant (constraint), a forward primer was curated on the online tool.

Similarly, keeping forward primer of RH the constraint, a reverse primer for amplifying RH

was curated with the same parameters as stated earlier. Further, to clone the arms with FRTs

in the desired orientation and, in pBSKS via Takara® In-fusion (Cat# 638915), the forward

primer was added with an overhang complementary to 15 bp immediately upstream of EcoRV

digested pBSKS. The reverse primer was also incorporated with a 15 bp o/h homologous to 15

bp immediately downstream of EcoRV digested pBSKS. The reverse primer of LH was

provided with an overhang for 2/3rd the size of the 34 bp minimal FRT sequence:

5′-GAAGTTCCTATTCtctagaaaGtATAGGAACTTC-3′

Note that the nucleotides indicated by uppercase alphabets represent binding sites for subunits

of the flippase enzyme (FLP). The lowercase alphabets indicate the asymmetric core of the

FRT site, which gives the site a particular orientation. A similar o/h was also provided on the

forward primer of RH, ensuring that it also has a 15 bps homology to the FRT overhang of LH

reverse primer. The complete list of primers is provided in Supplemental Table S4-S6.

Fly stocks and culture

Flies were grown in standard cornmeal yeast extract medium at 25°C unless otherwise

specified.

Generation of transgenic lines with FRT insertions at various regions within the Abd-B cis-regulatory landscape

All the donor constructs to target FRT were cloned in pBSKS vector with ~1 Kb homology arms from the region flanking the site of FRT insertion. The arms were amplified with overhangs for FRT sequences with 15-20 bp homology. The homology arms with FRT sequences were cloned in EcoRV digested pBSKS vector using standard protocol for Takara® in-fusion assembly (Cat#102518). All the constructs to express the guideRNA were cloned in pCFD3 vectors as described earlier.

The donor and guide expressing constructs were co-injected in *nos-Cas9 Drosophila* embryos (BDSC#54591) as previously described or outsourced to Centre for Cellular and Molecular Platforms (C-CAMP), Bangalore, India. The injected G0s and G1 flies were crossed with third chromosome balancer as single fly crosses. The G1s were screened for the desired transgenes by using primers that eccentrically flanked the FRT site. The amplified product was digested using *Xba-I* to validate the presence of FRT prior to sequencing based confirmation. A detailed schematics for obtaining the FRT transgenes is shown in Supplemental Table S2-S3 and S7-S8.

Generation of recombinants by Flp mediated recombination

The transgenic lines with desired FRTs were crossed and obtained with a cassette containing Flippase recombinase coding region under the influence of heat-shock inducible promoter (hsFLP; BDSC#7). The FLP activity was validated using a fly expressing Gal4 under the

influence of Act5C promoter ubiquitously upon FRT excision (BDSC#4779) combined with

UAS-GFP and hsFlp expressing flies (see supplementary information, Supplemental Fig. S2).

To obtain deletion and duplication, the flies with desired FRTs were crossed to obtain trans-

heterozygous fly with respect to the position of FRT. The trans-heterozygous late second instar

larvae are given heat shock at 37°C for 90 mins followed by a daily heat-shock periodically

after every 24 hrs till eclosion. The detailed schematics of the crosses for obtaining deletion

and duplication is shown in Supplemental Table S9-S10. For obtaining inversion, a fly with

two FRTs in -cis is given heat shock as standardized. The schematics of genetic crosses are

provided in Supplemental Table S11-S12.

Population assay of variable phenotype for *Dp_i6-F8*

25 females and 15 males of either *Dp_i6-F8* or 25 females of *Dp_i6-F8* and 15 males of CS

lines were mated and the number of progenies eclosing counted with different phenotypes. The

distribution of heterozygous and homozygous progenies was recorded. The flies with affected

abdominal segments A1-A8 were counted as category A. The flies with a distortion in A5-A8

were counted in category B. Category C included flies with aberrations in A6-A7. The flies

having a defect in only one segment (often A6 or A7) were grouped as Category D. Category

E mutants had minor defects in segments without perturbing the segmental boundaries, often

characterized as wavy segmental borders. The category F flies appeared normal when

compared to a wild-type fly. The distribution was normalized as the percentage of the total

population for a given category. The detailed counting is presented in Supplementary Data in

form of a workbook.

Cuticle preparations of adult abdominal segments

Adult abdominal cuticles were prepared as described earlier. Briefly, 2-3 days old adult flies

were dehydrated in 70% ethanol for 24 hrs. The flies are then boiled at 70°C in 10% potassium

hydroxide for 90 mins followed by three washes of autoclaved distilled water. Next, the flies

are washed in autoclaved water for 90 mins at 70°C and transferred to room temperature 70%

ethanol.

Mounting and imaging of cuticle preparations

The cuticles are transferred to a watch glass containing 70% ethanol and head-thorax region is

separated from the abdominal segments by a dissection needle. The abdominal segments are

given a sharp cut along the dorsal midline and transferred to a clean glass slide containing P700

halocarbon oil (Sigma #H8898). The cuticle is spread in desired orientation using dissection

needles and covered with cover slip. The cuticles were imaged at 1.0X objective, 35X zoom

on Zeiss AxioZoom V16 stereo microscope.

Competing Interest Statement

We declare no competing interest.

Acknowledgments

Authors would like to acknowledge the members of RKM group for constant discussion during

the conceptualization and execution of the project. NH is a fellow of Department of

Biotechnology, Govt of India and thanks the agency for timely financial support. RKM is

recipient of JC Bose National fellowship, India. The authors also thank CSIR, India, JC Bose

National Fellowship, India, and Tata Institute for Genetics and Society, India for financial

support at various levels. Further, the authors are thankful to Sonal Nagarkar Jaiswal lab for

providing hsFlp and Act5C>STOP>Gal4; UAS-GFP flies, and, Centre for Cellular and

Molecular Platforms (C-CAMP), Bangalore for injecting some of the constructs used in the

study. The authors would especially like to acknowledge the arduous efforts of Mandar Naik,

Binita Ghosh and Padmasini Chary in assisting NH with initial screenings of FRT transgenes

and standardizations of heat shock experiments for the FLP expressing flies.

Author Contributions

RKM conceptualized the project. RKM and NH designed the experiments. NH executed the

project. SP assisted NH in a molecular screening of the recombinants, cuticle preparations, and

imaging. NH wrote the manuscript with inputs from RKM. RKM supervised the project. NH

and RKM edited the manuscript.

References

Akbari OS, Bae E, Johnsen H, Villaluz A, Wong D, Drewell RA. 2008. A novel promoter-
    tethering element regulates enhancer-driven gene expression at the bithorax complex in
    the Drosophila embryo. *Development* **135**: 123–131.

Arib G, Cléard F, Maeda RK. 2015. Following the intracellular localization of the iab-
    8ncRNA of the bithorax complex using the MS2-MCP-GFP system. *Mechanisms of
    Development* **138**: 133–140.

Barges S, Mihaly J, Galloni M, Hagstrom K, Müller M, Shanower G, Schedl P, Gyurkovics
    H, Karch F. 2000. The Fab-8 boundary defines the distal limit of the bithorax complex
    iab-7 domain and insulates iab-7 from initiation elements and a PRE in the adjacent iab-
    8 domain. *Development* **127**: 779–790.

Bekiaris PS, Tekath T, Staiger D, Danisman S. 2018. Computational exploration of cis -
regulatory modules in rhythmic expression data using the " Exploration of Distinctive
CREs and CRMs " (EDCC) and " CRM Network Generator " (CNG) programs. *PLoS*
ONE **13**: e0190421.
https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0190421.

Bender WW. 2020. Molecular Lessons from the Drosophila Bithorax Complex. *Genetics*
216: 613–617.

Birve A, Sengupta AK, Beuchle D, Larsson J, Kennison JA, Rasmuson-Lestander A, Muller
J. 2001. Su(z)12, a novel Drosophila Polycomb group gene that is conserved in
vertebrates and plants. *Development* **128**: 3371–3379.

Bowman SK, Deaton AM, Domingues H, Wang PI, Sadreyev RI, Kingston RE, Bender W.
2014. H3K27 modifications define segmental regulatory domains in the Drosophila
bithorax complex. *Elife* **3**: 1–13.

Celniker SE, Lewis EB. 1993. Molecular basis of Transabdominal-a sexually dimorphic
mutant of the bithorax complex of Drosophila. *Developmental Biology* **90**: 1566–1570.

Celniker SE, Sharma S, Keelan DJ, Lewis EB. 1990. The molecular genetics of the bithorax
complex of Drosophila: Cis-regulation in the Abdominal-B domain. *EMBO Journal* **9**:
4277–4286.

Cléard F, Moshkin Y, Karch F, Maeda RK. 2006. Probing long-distance regulatory
interactions in the Drosophila melanogaster bithorax complex using Dam identification.
Nature Genetics **38**: 931–935.

Crocker J, Preger-Ben Noon E, Stern DL. 2016. *The Soft Touch: Low-Affinity Transcription*
Factor Binding Sites in Development and Evolution. 1st ed. Elsevier Inc.

Fabre PJ, Leleu M, Mormann BH, Lopez-Delisle L, Noordermeer D, Beccari L, Duboule D.
2017. Large scale genomic reorganization of topological domains at the HoxD locus.
Genome Biology **18**: 1–15.

Foronda D, Navas LFDE, Garaulet DL, Sánchez-herrero E. 2009. Function and specificity of
Hox genes. *International Journal of Developmental Biology* **1419**: 1409–1419.

Galloni M, Gyurkovics H, Schedl P, Karch F. 1993. The bluetail transposon: evidence for
independent cis-regulatory domains and domain boundaries in the bithorax complex.
The EMBO Journal **12**: 1087–1097.

Garaulet DL, Lai EC. 2015. Hox miRNA regulation within the Drosophila Bithorax complex:
Patterning behavior. *Mechanisms of Development* **138**: 151–159.

Gaston K, Jayaraman PS. 2003. Transcriptional repression in eukaryotes: Repressors and
repression mechanisms. *Cellular and Molecular Life Sciences* **60**: 721–741.

Gligorov D, Karch F, Maeda RK, Wolfner MF, Frei Y, Sitnik JL, Prince E. 2018. The
lncRNA male-specific abdominal plays a critical role in Drosophila accessory gland
development and male fertility. *PLOS Genetics* **14**: e1007519.

Golic KG, Lindquist S. 1989. The FLP recombinase of yeast catalyzes site-specific
recombination in the drosophila genome. *Cell* **59**: 499–509.

Golic MM, Rong YS, Petersen RB, Lindquist SL, Golic KG. 1997. FLP-mediated DNA
mobilization to specific target sites in Drosophila chromosomes. *Nucleic Acids Research*
25: 3665–3671. https://academic.oup.com/nar/article/25/18/3665/1046549 (Accessed
May 18, 2022).

Gummalla M, Galetti S, Maeda RK, Karch F. 2014. Hox gene regulation in the central
nervous system of Drosophila. *Frontiers in Cellular Neuroscience* **8**: 1–12.

Guo Y, Xu Q, Canzio D, Shou J, Li J, Gorkin DU, Jung I, Wu H, Zhai Y, Tang Y, et al. 2015.
CRISPR Inversion of CTCF Sites Alters Genome Topology and Enhancer/Promoter
Function. *Cell* **162**: 900–910.

Gyurkovics H, Gausz J, Kummer J, Karch F. 1990a. A new homeotic mutation in the
Drosophila bithorax complex removes a boundary separating two domains of regulation.
EMBO J **9**: 2579–2585.

Gyurkovics H, Gausz J, Kummer J, Karch F. 1990b. A new homeotic mutation in the
Drosophila bithorax complex removes a boundary separating two domains of regulation.
The EMBO Journal **9**: 2579–2585.

Hagstrom K, Muller M, Schedl P. 1996. Fab-7 functions as a chromatin domain boundary to
ensure proper segment specification by the Drosophila bithorax complex. *Genes and
Development* **10**: 3202–3215.

Hajirnis N, Mishra RK. 2021. Homeotic Genes: Clustering, Modularity, and Diversity.
Frontiers in Cell and Developmental Biology **9**: 1–20.

Hendrickson JE, Sakonju S. 1995. Cis and trans interactions between the iab regulatory
regions and abdominal-A and abdominal-B in Drosophila melanogaster. *Genetics* **139**:
835–848.

Ho MCW, Johnsen H, Goetz SE, Schiller BJ, Bae E, Tran DA, Shur AS, Allen JM, Rau C,
Bender W, et al. 2009. Functional evolution of cis-regulatory modules at a homeotic
gene in Drosophila. *PLoS Genetics* **5**. https://doi.org/10.1371/journal.pgen.1000709.

Hogga I, Karch F. 2002. Transcription through the iab-7 cis-regulatory domain of the
bithorax complex interferes with maintenance of Polycomb-mediated silencing.
Development **129**: 4915–4922.

Horn C, Handler AM. 2005. Site-specific genomic targeting in Drosophila. *Proc Natl Acad
Sci U S A* **102**: 12483–12488. www.pnas.orgcgidoi10.1073pnas.0504305102 (Accessed
May 18, 2022).

Iampietro C, Cléard F, Gyurkovics H, Maeda RK, Karch F. 2008. Boundary swapping in the
Drosophila Bithorax complex. *Development* **135**: 3983–7.

Iampietro C, Gummalla M, Mutero A, Maeda RK. 2010. Initiator Elements Function to
Determine the Activity State of BX-C Enhancers. *PLoS Genetics* **6**.

Karch F, Weiffenbach B, Peifer M, Bender W, Duncan I, Celniker S, Crosby M, Lewis EB.
1985. The abdominal region of the bithorax complex. *Genes, Development, and Cancer:*
The Life and Work of Edward B Lewis **43**: 327–357.

Kaufman TC, Seeger MA, Olsen G. 1990. Molecular and Genetic Organization of the
Antennapedia Gene Complex of Drosophila melanogaster. *Advances in Genetics* **27**:
309–362.

Kyrchanova O, Ivlieva T, Toshchakov S, Parshikov A, Maksimenko O, Georgiev P. 2011.
Selective interactions of boundaries with upstream region of Abd-B promoter in
Drosophila bithorax complex and role of dCTCF in this process. *Nucleic Acids Research*
39: 3042–3052.

Kyrchanova O, Mogila V, Wolle D, Paolo J, White R, Georgiev P, Schedl P. 2015. The
boundary paradox in the Bithorax complex. *MOD* **138**: 122–132.

Lelli KM, Slattery M, Mann RS. 2012. Disentangling the many layers of eukaryotic
transcriptional regulation. *Annu Rev Genet* **46**: 43–68.

Lewis EB. 1978. A gene complex controlling segmentation in Drosophila. *Nature* **276**: 565–
570.

Li J, Shou J, Guo Y, Tang Y, Wu Y, Jia Z, Zhai Y, Chen Z, Xu Q, Wu Q. 2015. Efficient
inversions and duplications of mammalian regulatory DNA elements and gene clusters
by CRISPR/Cas9. *Journal of Molecular Cell Biology* **7**: 284–298.

Maeda RK. 2006. The ABC of the BX-C: the bithorax complex explained. *Development* **133**:
1413–1422. http://dev.biologists.org/cgi/doi/10.1242/dev.02323.

Maeda RK, Karch F. 2007. Making connections: boundaries and insulators in Drosophila.
Current Opinion in Genetics and Development **17**: 394–399.

Maeda RK, Karch F. 2015. The open for business model of the bithorax complex in
Drosophila. *Chromosoma* **124**: 293–307.
http://www.ncbi.nlm.nih.gov/pubmed/26067031 (Accessed July 12, 2016).

Mateo LJ, Murphy SE, Hafner A, Cinquini IS, Walker CA, Boettiger AN. 2019. Visualizing
DNA folding and RNA in embryos at single-cell resolution. *Nature*.
http://dx.doi.org/10.1038/s41586-019-1035-4.

Mihaly J, Barges S, Sipos L, Maeda R, Cléard F, Hogga I, Bender W, Gyurkovics H, Karch
F. 2006. Dissecting the regulatory landscape of the Abd-B gene of the bithorax complex.
Development **133**: 2983–93.

Mihaly J, Hogga I, Barges S, Galloni M, Mishra RK, Hagstrom K, M??ller M, Schedl P,
Sipos L, Gausz J, et al. 1998. Chromatin domain boundaries in the Bithorax complex.
Cellular and Molecular Life Sciences **54**: 60–70.

Mishra RK, Karch F. 1999. Boundaries that demarcate structural and functional domains of
chromatin. *Journal of Biosciences* **24**: 377–399.

Mishra RK, Mihaly J, Barges S, Spierer a, Karch F, Hagstrom K, Schweinsberg SE, Schedl
P. 2001. The iab-7 polycomb response element maps to a nucleosome-free region of

chromatin and requires both GAGA and pleiohomeotic for silencing activity. *Mol Cell Biol* **21**: 1311–1318.

Narlikar L, Ovcharenko I. 2009. Identifying regulatory elements in eukaryotic genomes. *Briefings in Functional Genomics and Proteomics* **8**: 215–230.

Négre N, Brown CD, Ma L, Bristow CA, Miller SW, Wagner U, Kheradpour P, Eaton ML, Loriaux P, Sealfon R, et al. 2011. A cis-regulatory map of the Drosophila genome. *Nature* **471**: 527–531.

Peifer M, Karch FF, Bender W. 1987. The bithorax complex: control of segment identity. *Genes Dev* **1**: 891–898.

Pérez-Lluch S, Cuartero S, Azorín F, Espinàs ML. 2008. Characterization of new regulatory elements within the Drosophila bithorax complex. *Nucleic Acids Research* **36**: 6926–6933.

Phan QV, Contzen J, Seemann P, Gossen M. 2017. Site-specific chromosomal gene insertion : Flp recombinase versus Cas9 nuclease. *Nature Scientific Reports* **7**: 1–12.

Pignoni F, Hu B, Zavitz KH, Xiao J, Garrity PA, Zipursky SL. 1997. The Eye-Specification Proteins So and Eya Form a Complex and Regulate Multiple Steps in Drosophila Eye Development. *Cell* **91**: 881–891.

Polychronidou M, Lohmann I. 2013. Cell-type specific cis-regulatory networks: insights from Hox transcription factors. *Fly (Austin)* **7**: 13–17.

Port F, Muschalik N, Bullock SL. 2015. CRISPR with independent transgenes is a safe and robust alternative to autonomous gene drives in basic research. *bioRxiv* 017384. http://biorxiv.org/content/early/2015/04/01/017384.abstract.

Postika N, Metzler M, Affolter M, Müller M, Schedl P, Georgiev P, Kyrchanova O. 2018. Boundaries mediate long-distance interactions between enhancers and promoters in the Drosophila Bithorax complex. *PLoS Genetics* **14**: 1–21.

Postika N, Schedl P, Georgiev P, Kyrchanova O. 2021a. Mapping of functional elements of the Fab-6 boundary involved in the regulation of the Abd-B hox gene in Drosophila melanogaster. *Scientific Reports* **11**: 1–11. https://doi.org/10.1038/s41598-021-83734-8.

Postika N, Schedl P, Georgiev P, Kyrchanova O. 2021b. Redundant enhancers in the iab-5 domain cooperatively activate Abd-B in the A5 and A6 abdominal segments of Drosophila. *Development* **148**: 1–20.

Ringrose L, Paro R. 2004. Epigenetic Regulation of Cellular Memory by the Polycomb and Trithorax Group Proteins. *Annual Review of Genetics* **38**: 413–443.

Singh NP, Mishra RK. 2015. Specific combinations of boundary element and Polycomb response element are required for the regulation of the Hox genes in Drosophila melanogaster. *Mechanisms of Development* **138**: 141–150.

Sipos L, Mihály J, Karch F, Schedl P, Gausz J, Gyurkovics H. 1998. Transvection in the Drosophila Abd-B domain: Extensive upstream sequences are involved in anchoring distant cis-regulatory regions to the promoter. *Genetics* **149**: 1031–1050.

Srinivasan A, Mishra RK. 2020. Genomic organization of Polycomb Response Elements and
its functional implication in Drosophila and other insects. *Journal of Biosciences* **45**.

Starr MO, Ho MCW, Gunther EJM, Tu YK, Shur AS, Goetz SE, Borok MJ, Kang V, Drewell
RA. 2011. Molecular dissection of cis-regulatory modules at the Drosophila bithorax
complex reveals critical transcription factor signature motifs. *Developmental Biology*
359: 176–189. http://dx.doi.org/10.1016/j.ydbio.2011.07.028.

Stuttgart L, Herberth F. 1988. Paternal Imprinting of Iversion of Uab[1] Causes Homeotic
Transformations in Drosophila. *Genetics* **118**: 103–107.

Vazquez J, Muller M, Pirrotta V, Sedat JW. 2006. The Mcp Element Mediates Stable Long-
Range Chromosome–Chromosome Interactions in Drosophila. *Molecular Biology of the*
Cell **17**: 2158–2165.

Zhou J, Barolo S, Szymanski P, Levine M. 1996. The Fab-7 element of the bithorax complex
attenuates enhancer-promoter interactions in the Drosophila embryo. *Genes and*
Development **10**: 3195–3201.

Zhou J, Levine M. 1999. A novel cis-regulatory element, the PTS, mediates an anti-insulator
activity in the Drosophila embryo. *Cell* **99**: 567–75.

Zinzen RP, Girardot C, Gagneur J, Braun M, Furlong EEM. 2009. Combinatorial binding
predicts spatio-temporal cis-regulatory activity. *Nature* **462**: 65–70.

  Figure legends

Figure 1.Targeting the cis-regulatory modules of the bithorax complex regulating *Abd-B*:
(A) The proposed experiment for the current study is to generate modular deletion, duplication
and inversion within the *cis*-regulatory landscape of *Abd-B*. Top row shows the normal
arrangement of the CRMs (green solid arrows) of *Abd-B*, iab5, iab6, iab7 and iab8 demarcated
by the boundaries MCP, Fab6, Fab7 and Fab8 (brown rectangle and pentagon arrows). The
blue arrow depicts iab4 which is a regulator of *abd-A*. The re-arrangement of the CRMs of
different nature are shown below the wild-type representation. The concomitant expression of
Abd-B driven by the four CRMs in the four abdominal segments of the fly is shown on right.
(B) Top row: Representation of the wild-type locus of *Abd-B*. Green arrows are regulators of
Abd-B, blue arrow is regulator of abd-A, brown rectangles represent the boundaries and yellow
triangles represent the PREs. Red arrows pointing down indicates the sites proposed for FRT
insertion for which constructs were generated. Bottom rows indicate the nature of transgenes
generated with FRTs (red chevron arrow) in different locations and orientations. (C) The lines
with altered CRMs generated in this study.

Figure 2. Deletion of *iab5* and *Fab6*: (A) Representation of the wild-type Abd-B cis-
regulatory landscape and dichromatic representation of flies with endogenous FRT (red
chevron arrow) between *MCP-iab5* (*MP_F_i5*), and, *Fab6-iab6* (*F6_F_i6*). The bottom row
indicates the deletion product formed as a result of recombination indicated by red-dotted line
across the middle rows. (B) Abdominal cuticle preparation of adult male flies two days post
eclosion. The curved segments on the left form the dorsal side of the abdomen, while the
straight lane of bristles towards the right is the ventral sternites. Each dorsal segment has a
distinct pattern of melanisation, as indicated by blue and green arrows. The blue arrows indicate
melanisation in the posterior ends of segments A2-A4 under the influence of *abd-A* driven by
iab2 through *iab4*. The green arrows indicate complete melanisation of A5 and A6 influenced
by the levels of *Abd-B* regulated by *iab5* and *iab6* in respective segments. Each ventral segment
also has a distinct arrangement of bristles. The A6 does not have any bristles on sternite in the
wild-type fly, as shown by the red arrowhead and dotted shape. (C) The abdominal cuticle of
adult males with deletion of *iab5* and *Fab6*. The males show a strong homeotic transformation
of A5 to A4 characterized by loss of dorsal melanization (curved segments on the right). The
flies also show A6 to A4 transformation characterized by loss of melanization of dorsal
segment and appearance of bristles on ventral sternites (bold red arrow). (D) Abdominal cuticle
preparation adult female flies. The dorsal melanization and the pattern of tergites and sternites
are largely indistinct in females. However, the A6 of the females have a hardened sclera in
wild-type females (dotted shape). Note that A7 of the wild-type flies have fewer bristles that
droop towards the genitalia. The A7 of the fly is also observably smaller than the other
segments of the fly. (E) Abdominal cuticle prep of adult *Del_i5-F6* female fly. Note that the
sternites in A5 and A6 are strikingly similar to A4 as indicated by bold arrowheads. The tergites
and melanisation pattern also appears strikingly similar to A4 in A5 and A6. Thus, both males
and females had similar cuticular homeotic transformations in the flies lacking known *iab5* and
Fab6 regions.

Figure 3. Deletion of *iab6-iab7*: (A) Representation of the wild-type *Abd-B* cis-regulatory
landscape and dichromatic representation of flies with endogenous FRT (red chevron arrow)
between *Fab6-iab6* (*i6_F_F6*), and, *iab7-Fab8* (*F8_F_i7*). The bottom row indicates the
deletion product formed as a result of recombination indicated by red-dotted line across the

middle rows. (B-E) Abdominal cuticle preparation of adult flies two days post eclosion. (B) Abdominal cuticle preparation of wild-type male. (C) The abdominal cuticle of adult males with deletion of *iab*6 through *iab7*. The flies show a strong homeotic transformation of A6 and A7 to copies of A5. The A6 to A5 transformation is characterised by broadened A6 and sternal bristles' appearance (bold red arrow). The A7 to A5 transformation is characterised by a distinct segment with dorsal melanisation similar to A5. The transformed segment also has sternal bristles similar to A5. (D) Abdominal cuticle preparation of wild type female. (E) Abdominal cuticle prep of adult *Del_i6-i7* female fly. Note that the sternites in A6 and A7 are strikingly similar to A5 (bold red arrows). The tergites and melanisation pattern also appears strikingly similar to A5 in A6 and A7. Thus, both males and females had similar cuticular homeotic transformations in the flies lacking known *iab6* through *iab7* regions.

Figure 4. Duplication of *iab6* through *iab7*: (A) Representation of the wild-type *Abd-B* cis-regulatory landscape and dichromatic representation of flies with endogenous FRT (red chevron arrow) between *Fab6-iab6* (*i6_F_F6*), and, *iab7-Fab8* (*F8_F_i7*). The bottom row indicates the duplication product formed as a result of recombination indicated by red-dotted line across the middle rows. (B-E) Abdominal cuticle preparation of adult male flies two days post eclosion. (B) Abdominal cuticle preparation of wild-type male. (C) The abdominal cuticle of adult males with duplication of *iab*6 through *iab7*. The flies show a strong homeotic transformation of A6 to a copy of A7. The transformation is characterised by loss of segment with dorsal melanization. The A6 also has partial signatures of anteriorization indication by the appearance of sternal bristles and loss of sclera. (bold red arrow). The A5 of the fly also shows partial posteriorization indicated by loss of sternal bristles. The partial posteriorization is also suggested by slight narrowing of dorsal abdomen akin to A6 of a wild-type fly (shown by the red dotted peripheral structure). (D) Abdominal cuticle preparation of adult female flies. (E) Abdominal cuticle prep of adult *Dp_i6-i7* female fly. The A5 of the fly appears normal. The A6 shows a clear homeotic transformation into a copy of A7 by the appearance of A7-specific sternal bristles. Note that the sternites in A6 and A7 are strikingly similar (bold red arrows). The tergites and melanisation patterns of A6 and A7 also appears strikingly similar. Thus, both males and females had similar cuticular homeotic transformations in the flies lacking known *iab6* through *iab7* regions. The A7 of the flies show loss of melanization, indicating either transformation into A8 or uncharacterised structures.

Figure 5. Duplication of *iab6* through *iab7*: (A) Representation of the wild-type *Abd-B cis*-regulatory landscape and dichromatic representation of flies with endogenous FRT (red chevron arrow) between *Fab6-iab6* (*i6_F_F6*), and, *Fab8-iab8* (*F8_F_i8*). The bottom row indicates the duplication product formed as a result of recombination indicated by red-dotted line across the middle rows. (B-H) Bright-field images of the dorsal abdomen of two-days post eclosion adult females from different classes of mutants. The images are represented in grayscale for easier visualization of the segmental pattern. (B) A wild-type CS fly is shown as a control for patterning defects in the mutants. (C-H) Flies with variable phenotype classified into different categories depending upon the extent of segmental aberration. See text for details. (I) A standard histogram plotted for the distribution of 827 heterozygous and 557 homozygous flies for the duplication of *iab6* through *iab8* with different phenotypes grouped under categories A-F (see text for details). Blue bars indicate the percentage distribution of heterozygous flies, and the orange bars represent the percentage distribution of the homozygous flies.

Figure 6. Inversion of *iab5* through *iab7*: (A) Simple representation of flies with endogenous
FRT (red chevron arrow) between *MCP* and *iab5* with the direction of FRT towards *iab5*
(*MP_F_i5*). The flies also have another FRT inserted between *iab7* and *Fab8* with the
direction of FRT towards *iab7* (*F8B_F_i7*). The activity of FLP causes an inversion of iab5
through iab7 as shown in the bottom row. The FRT mediated recombination happens across
the two FRTs in *cis-* as indicated by curved dotted arrows in the middle rows. (B) Cuticle
preparation of adult male abdomen. (C) The abdominal cuticle of adult males with inversion
of *iab5* through *iab7*. The flies show a strong homeotic transformation of A6 to a copy of A7.
The transformation is characterised by loss of segment with dorsal melanization. The A6 also
has partial signatures of anteriorization indicated by the appearance of sternal bristles and loss
of sclera (bold red arrow). The A5 of the fly also shows partial posteriorization indicated by
loss of sternal bristles. The partial posteriorization is also suggested by slight narrowing of
dorsal abdomen akin to A6 of a wild-type fly (shown by the red dotted peripheral structure).
(D) Abdominal cuticle preparation and mounting of 2 days post eclosion adult female flies.
(E) Abdominal cuticle prep of adult *Dp_i6-i7* female fly. The A5 of the fly appears normal.
The A6 shows a clear homeotic transformation into a copy of A7 by the appearance of A7-
specific sternal bristles. Note that the sternites in A6 and A7 are strikingly similar (bold red
arrows). The tergites and melanisation patterns of A6 and A7 also appear strikingly similar.

925   Figure 1

931 Figure 2

935   Figure 3

939   Figure 4

943   Figure 5

947   Figure 6