Hotspot of *de novo* telomere addition stabilizes linear amplicons

in yeast grown in sulfate-limiting conditions

Remington E. Hoerr[1], Alex Eng[2], Celia Payen[2,3], Sara C. DiRienzi[2,4], M. K. Raghuraman[2], Maitreya J.

Dunham[2], Bonita J. Brewer[2], Katherine L. Friedman[1]

[1] Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA, Email:

remington.hoerr@vanderbilt.edu; katherine.friedman@vanderbilt.edu

[2] Department of Genome Sciences, University of Washington, Seattle, WA, USA, Email: engal@uw.edu;

raghu@uw.edu; bbrewer@uw.edu; maitreya@uw.edu;

[3] Current affiliation: IFF, Wilmington, Delaware, USA, Email: celia.e.payen@iff.com

[4] Current affiliation: Department of Molecular Virology and Microbiology, Baylor College of Medicine,

Houston TX, USA, Email: sara.dirienzi@bcm.edu

Keywords: Copy number variation; gene amplification; evolutionary genomics; *de novo* telomere

addition; Cdc13; DNA repair; *Saccharomyces cerevisiae*

Abstract:

Evolution is driven by the accumulation of competing mutations that influence survival. A broad form of genetic variation is the amplification or deletion of DNA (≥50 bp) referred to as copy number variation. In humans, copy number variation may be inconsequential, contribute to minor phenotypic differences, or cause conditions such as birth defects, neurodevelopmental disorders, and cancers. To identify mechanisms that drive copy number variation, we monitored the experimental evolution of *Saccharomyces cerevisiae* populations grown under sulfate-limiting conditions. Cells with increased copy number of the gene *SUL1*, which encodes a primary sulfate transporter, exhibit a fitness advantage. Previously, we reported interstitial inverted triplications of *SUL1* as the dominant rearrangement in a haploid population. Here, in a diploid population, we find instead that small linear fragments containing *SUL1* form and are sustained over several generations. Many of the linear fragments are stabilized by *de novo* telomere addition within a telomere-like sequence near *SUL1* (within the *SNF5* gene). Using an assay that monitors telomerase action following an induced chromosome break, we show that this region acts as a hotspot of *de novo* telomere addition and that required sequences map to a region of <250 base pairs. Consistent with previous work showing that association of the telomere-binding protein Cdc13 with internal sequences stimulates telomerase recruitment, mutation of a four-nucleotide motif predicted to associate with Cdc13 abolishes *de novo* telomere addition. Our study suggests that internal telomere-like sequences that stimulate *de novo* telomere addition can contribute to adaptation by promoting genomic plasticity.

Introduction:

Genomic variation is the basis of evolution. Mutations, including both single-nucleotide polymorphisms and larger genomic rearrangements, can rise in prevalence within a population if they confer a fitness advantage. Copy number variation (CNV) refers to a genomic rearrangement that results in loss or gain of DNA (Alkan et al. 2011; MacDonald et al. 2014). CNV is most likely to have a phenotypic impact when the DNA that is duplicated or lost contains one or more genes that impact cellular fitness. In humans, CNV is associated with a variety of disorders primarily (but not exclusively) related to intellectual disabilities, developmental delays, and predisposition to cancers (reviewed in Zhang et al. 2009; Shaikh 2017; Doran and Pennington 2022; R. H. Song et al. 2022).

CNV can arise through a variety of chromosomal rearrangements stemming from errors in replication fork progression, recombination, or in DNA damage repair pathways (Hastings et al. 2009). The frequency of CNVs is often increased in cancer cells due to dysregulation of DNA repair or replication pathways (Doran and Pennington 2022). Gene amplification can present in a variety of configurations including both direct and inverted tandem duplication (additional copies at the original location), dispersed duplication (integration of additional copies elsewhere in the genome), or double minute chromosomes that are small circular extrachromosomal fragments of DNA that lack a centromere (Biedler et al. 1983; Chakraborty and Ay 2018; Doran and Pennington 2022).

The origin of CNVs is not well understood, in part because the initiating event may be replaced over time by more beneficial secondary and tertiary events that confer increased growth advantages (Ilić et al. 2022). For example, double minute chromosomes may in some cases serve as recombinational intermediates in the formation of tandem or dispersed duplications (Turner et al. 2017; Rosswog et al. 2021; K. Song et al. 2022). Since most studies of CNV in humans rely on samples from a single point in time, they provide only a snapshot of the most dominant rearrangement (Chakraborty

and Ay 2018). This limitation highlights the need for experimental systems in which the temporal

relationship between events can be observed.

The budding yeast *Saccharomyces cerevisiae* utilizes DNA repair pathways similar to those of

human cells but is more easily manipulated and monitored for genomic rearrangements (Krogh and

Symington 2004; Wellinger and Zakian 2012). For example, growth under specific environmental

conditions selects for amplification of one of several genes (e.g., *HXT6/7*, *CUP1, GAP1,* or *SUL1*) (Brown

et al. 1998; Gresham et al. 2008; Kao and Sherlock 2008; Gresham et al. 2010; Zhang et al. 2013).

Growth of continuous yeast cultures in a chemostat allows the genetic composition of the population to

be monitored over time, providing an attractive system to monitor CNV formation and persistence in a

heterogenous population. Previously reported evolution experiments with *S. cerevisiae* describe several

different types of chromosomal rearrangements that increase gene copy number. Extrachromosomal

circular DNAs, similar to the double minute chromosomes described above, have been reported

(Gresham et al. 2010). Additionally, persistent linear fragments have also been described (Dorsey et al.

1992; Narayanan et al. 2006). Circular molecules lack DNA ends, but linear fragments must protect their

ends from nucleolytic degradation. In most cases, protection of the ends is afforded by the telomere, a

repetitive, protein-bound DNA sequence that resists nuclease action and engagement with the DNA

repair machinery (Blackburn 1991). Where such linear fragments have been described, they arise from

palindromic duplication, resulting in an isochromosomal fragment capped at both ends by copies of the

same pre-existing telomere (Dorsey et al. 1992; Narayanan et al. 2006).

In haploid laboratory strains grown in sulfate-limiting chemostats, the fitter variants almost

always contain an inverted interstitial triplication of the primary sulfate transporter gene (*SUL1*) and

variable amounts of the flanking regions in an otherwise unrearranged chromosome (Araya et al. 2010;

Payen et al. 2014; Brewer et al. 2015). Here, we report that diploids of the same laboratory strain

appear to follow a different path to *SUL1* amplification. Instead of chromosomally stabilized amplicons,

we find extrachromosomal linear fragments that contain *SUL1* and the nearby endogenous telomere at

one end and a new telomere added to a sequence centromere proximal to *SUL1* at the other end.

Similar fragments arose independently multiple times, suggesting that one or more sequences

centromere proximal to *SUL1* have a propensity to stimulate new telomere formation. Indeed, we show

that a sequence within the *SNF5* gene is sufficient to stimulate *de novo* telomere addition when moved

to an ectopic location. The ability of this sequence to stimulate *de novo* telomere addition requires at

least two short sequence motifs that are predicted to associate with the telomere binding protein Cdc13

(Eldridge et al. 2006) and can be functionally replaced by artificial recruitment of Cdc13.

Methods and Materials:

<u>Strains and Plasmids</u>

All strains for SiRTA analyses were derived from S288c and are listed in File S1. Plasmid pAB180

(pRS414-ADHpromoter-GBD-*CDC13*) and pAB220 (pRS414-ADHpromoter-GBD) were gifts of A. Bianchi.

Sulfate-limited chemostat experiments were performed on strains of the S288c background (FY3 and

FY5) (Gresham et al. 2008; Miller et al. 2013). Creation of the *ars228* deletion has been described

(Brewer et al. 2015). Strains for the HO cleavage assay were generated as described in Ngo et al. 2020.

In brief, the HO endonuclease recognition sequence was integrated on chromosome VII by one-step

gene replacement using plasmid pHO invC (based on JH2017; gift of J. Haber). The *URA3* gene was

integrated by one-step gene replacement using pRS306 (Sikorski and Hieter 1989) as template. Putative

SiRTAs were inserted on chromosome VII using CRISPR/Cas9 as described (Anand et al. 2017) using gRNA

sequence TGCGGCAAGTTCATCTTCCA. Sequences of different length within *SNF5* were amplified from

genomic yeast DNA with appropriate primers (File S1) and additional sequences were added that lie

upstream (5'-TTTCTTTGGAAAACGTTGAAAATGAGGTTCTATGATCTAC) and downstream (5'-

AGAACATAGAATAAATTTGGTACTGGAACGTTGATTAACT) of the gRNA recognition site. *RAD52* was

replaced by one-step gene replacement using pFA6a-TRP1 (Longtine et al. 1998) as template.

<u>Contour-clamped homogeneous electric field (CHEF) gels</u>

Agarose plugs for CHEF gel electrophoresis were generated using the method by Lucas Argueso (Kwan et

al. 2016). For separation of whole chromosomes, run conditions in the BioRad CHEF-DRII were 1% LE

agarose in 0.5XTBE in 2.3 L 0.5% TBE running buffer at 14˚C. Switch times were either 47" to 170" at

165V for 39-62 hours or 300" to 900" at 100V for 68 hr. Linear fragments were separated from

chromosomes by CHEF gels with switch times 1.7" to 2.6" at 198 V for 20 hr. Restriction analysis of

linear fragments were performed by conventional agarose gel electrophoresis in 0.4% ME agarose gels

in 1XTBE at 1V/cm for 19 hr at room temperature. Standard protocols for restriction digestion,

Southern blotting and hybridization using ^{32}P-labeled PCR probes were used. Linear fragments were

concentrated from plug supernatants using Centricon Filter devices and then ethanol precipitated.

<u>Sequence analysis of linear fragments</u>

Whole genome sequencing of fragments purified from plug supernatants and split read analysis were as

described (Payen et al. 2014).

<u>HO cleavage assay</u>

The HO cleavage assay was performed as described in Ngo et al. 2020. In brief, cells were grown in SD-

Ura medium (+2% raffinose) overnight to an optical density at 600 nm (OD600) of 0.6-0.9. Cells were

plated on yeast extract peptone medium with 2% galactose (YEPGal). Simultaneously, cells were serially

diluted in water and plated on yeast extract peptone medium with 2% dextrose (YEPD) to determine

total cell number. When plasmid selection was required (pRS414, pAB220, and pAB180), cells were

plated on synthetic medium lacking tryptophan and containing either 2% glucose or 2% galactose. After

3 days, colonies were counted and at least 100 individual colonies were transferred to plates containing

5-fluoroorotic acid (5-FOA) to determine the frequency of *URA3* loss. For each experiment, at least 30 5-

FOA resistant clones were grown in YEPD for DNA extraction by MasterPure™ Yeast DNA Purification Kit

(Lucigen). The relative location of each chromosome truncation event was determined using multiplex

PCR with primers that anneal centromere or telomere proximal to the SiRTA (File S1). *De novo* telomere

addition events were amplified using CHECKCHR7GUIDE1F and TelomereRev as primers. The purified

PCR product was sequenced using Sanger sequencing to determine the site of *de novo* telomere

addition. Full data sets corresponding to Figures 3 and 4 are found in File S2.

Results:

<u>Linear fragments containing *SUL1* accumulate in cells grown in sulfate-limiting conditions</u>

Under sulfate-limiting conditions, *S. cerevisiae* cells with increased capacity for sulfate transport

exhibit a selective growth advantage (Gresham et al. 2008). The most common genomic rearrangement

observed after prolonged competitive growth of haploid lab strains in low-sulfate conditions is an

interstitial triplication of a chromosomal region containing the primary sulfate-transporter gene *SUL1*

and its adjacent origin of replication *ARS228*, located ~25 kb from the right telomere of chromosome II

(Payen et al. 2014; Brewer et al. 2015)*.* From the structure of these amplicons—the central copy of the

triplication is in inverted orientation—Brewer et al. (2011) proposed a model for their generation that

relies on the presence of the adjacent origin of replication and novel junctions that occur at pre-existing

flanking short interrupted inverted repeats (Origin Dependent Inverted Repeat Amplification; ODIRA).

That the process is a byproduct of replication was inferred from the observation that deletion of *ARS228*

results in larger amplicons that encompass the next nearest replication origin (Brewer et al. 2015).

A key feature of the ODIRA model is the generation of an extrachromosomal intermediate that

later recombines with the chromosomal locus (Brewer et al. 2011). In an attempt to gather evidence for

the unique extrachromosomal intermediate, we constructed a diploid heterozygous for the *ARS228*

deletion (*ARS228/ars228Δ*; Brewer et al. 2015) and two diploid control strains (*ARS228/ARS228* and

ars228Δ/ars228Δ) and carried out independent evolution experiments in sulfate-limited chemostats (n =

10, 4, and 4, respectively). We reasoned that if the intermediate arose from the *ARS228* chromosome

and subsequently recombined with the *ars228Δ* chromosome, the resulting amplicon would contain

copies of both *ARS228 and ars228Δ*.

Contrary to expectations, these diploids did not produce the same type of inverted interstitial

amplicons of *SUL1*. Array comparative genome hybridization (aCGH) of each of the diploid populations

after ~180 generations (33-36 days) under sulfate-limitation revealed consistent amplification of the

SUL1 region (e.g., Figure 1A, Figure S1). However, in each case the amplicon was not interstitial but

continued through the right telomere. The amplified sequences were limited to this region of

chromosome II and the new junctions were narrowly restricted to a small region of chromosome II near

the *SNF5* gene.

To determine the genomic location of the amplicons, we analyzed samples taken every few days

through the 33 days of chemostat growth and analyzed the yeast karyotype by contour-clamped

homogeneous electric field (CHEF) gels. Southern blotting and hybridization with probes specific to

SUL1 detected chromosome II as expected, but also revealed the generation of small (~30 kilobases)

molecules in all samples from all three genotypes (e.g., Figure 1B, C; Figures S1 and S2). Because they

were generated in diploid cells with both copies of *ARS228* deleted (Figure S1C) we propose that the

adjacent origin of replication *ARS229* can support the creation and maintenance of these fragments.

The sizes of these molecules estimated from their migration on CHEF gels were consistent with the sizes

of the amplicon detected by aCGH suggesting that they were acentric linear fragments.

The linear fragments usually appeared early in the course of the selection (~by 50 generations)

and often persisted through the end of the experiment. In half of the cases, chromosomally integrated

SUL1 amplicons were the predominant form by the end of the chemostat run. In all but one of these

cases, the transient linear fragments disappeared concurrently with the emergence of new, slower-

migrating *SUL1*-hybridizing bands (Figures S1 and S2). The size of the major amplicon from the final day

in the chemostat culture matched the size of the transient linear fragments [in all but one of the

cultures (S702)], suggesting a precursor-product relationship between the linear fragments and the

chromosomally integrated form. One of the most intriguing aspects of these free linear fragments is

their consistent small size—ranging between ~26 and 42 kb (Figure 1C). Since all of the linear fragments

extend through the right telomere of chromosome II, the newly formed left ends of the linear fragments

appear to arise near or within *SNF5*, a gene centromere proximal to *SUL1*.

Linear fragments are stabilized by addition of a new telomere

While selective pressure can partially compensate for the lack of a centromere, stability of a

linear fragment requires protection at both ends from exonucleases. Normal chromosome ends

(telomeres) in *S. cerevisiae* consist of a thymine and guanine (TG)-rich strand that extends 10-15 bases

past the complementary adenine and cytosine (AC)-rich strand to create a 3' overhang (Blackburn 1991).

This TG-rich single-stranded overhang interacts with the enzyme telomerase, which utilizes its reverse

transcriptase catalytic subunit and intrinsic AC-rich RNA template to add telomeric repeats to the

chromosome end (Singer and Gottschling 1994; Lingner et al. 1997). In yeast, the 3' terminating strand

consists of a heterogeneous sequence pattern with a single thymine followed by one to three guanines

(abbreviated TG_{1-3}) (Wellinger and Zakian 2012).

The aCGH analyses (Figures 1A and S1) suggested that the right end of the linear fragment

contained the endogenous right telomere, but the structure of the left end was initially unclear. More

detailed analysis of the linear fragments was facilitated by the development of a method to purify these

small molecules through passive diffusion from agarose CHEF gel plugs. Over the course of several

weeks, the small fragments diffuse out of the agarose plug and into the supernatant where they can be

recovered, while the intact chromosomes remain trapped in the plug (Figure S3). Purified linear

fragments were digested with *Eco*NI, which cleaves three times in the terminal region of chromosome II

(Figure 2A). Following separation by CHEF gel, the samples were Southern blotted and probed

sequentially with sequences that hybridize to either *SUL1* or telomeric DNA (Figure 2B). *Eco*NI digestion

separated *SUL1* from the right telomere and allowed us to query both ends of the linear fragments. The

restriction fragments containing the right telomere hybridized to telomeric DNA ($C_{1-3}A/G_{1-3}T$) and not to

SUL1 (Figure 2B, gold rectangles). However, fragments of multiple sizes (ranging from ~3-25 kb)

hybridized to both the *SUL1* and the telomeric probe (Figure 2B), suggesting that the left end of each

fragment was also capped by telomeric DNA.

To determine the sites of telomeric repeat addition, we determined the sequence of fragments

purified from the plug supernatants by Illumina sequencing. The results revealed that nine of the 15

fragments contained telomeric repeats clustered within a ~100 bp region of the gene *SNF5*,

approximately ~7 kb from *SUL1* (Figure S4). With one exception, there was no evidence of subtelomeric

sequences, but rather the telomeric DNA was appended directly to the chromosome II sequence,

consistent with addition of a new telomere to internal sequences by telomerase.

Sequences in the *SNF5* gene stimulate unusually high levels of *de novo* telomere addition

The observation that many of the telomere addition events associated with extrachromosomal

SUL1 fragments cluster within a small (~100 bp) region is reminiscent of sequences in the yeast genome

that are hotspots of *de novo* telomere addition (termed SiRTAs or Sites of Repair-associated Telomere

Addition) (Stellwagen et al. 2003; Obodo et al. 2016; Hoerr et al. 2021). SiRTAs contain strand-specific,

TG-rich sequence motifs (Obodo et al. 2016; Ngo et al. 2020). These motifs can provide a substrate for

the acquisition of telomeric sequences if they become exposed in single-stranded DNA either as a result

of 5' end resection following a DSB or at nicks in single-stranded gaps. In either case, the resulting TG-

rich 3' overhang can invade a similar sequence and acquire a telomere through break-induced

replication or can undergo *de novo* telomere addition by direct action of telomerase (Putnam and

Kolodner 2017). Such events are most easily assayed in cases where the new telomere is added to the

centromere-proximal side of the break, causing a terminal truncation. In contrast, the putative SiRTA

within *SNF5* is oriented in the opposite orientation on the right arm of the chromosome. Rather than

stabilizing the centromere-containing region of the chromosome, *de novo* telomere addition stabilizes a

terminal acentric fragment containing *SUL1*.

Like telomeres, SiRTAs have an asymmetric sequence pattern in which one strand is relatively

rich in T and G nucleotides (Obodo et al. 2016; Ngo et al. 2020). For telomerase to act upon a SiRTA, this

TG-rich strand must be exposed in single-stranded DNA. SiRTAs have a bipartite structure consisting of

one TG-rich region that is most often the site of telomere addition (the "Core") and a second TG-rich

"Stim" sequence located 5' to the Core on the TG-rich strand (Obodo et al. 2016). The Stim sequence

itself is rarely targeted for telomere addition but is required to stimulate *de novo* telomere addition in

the Core (Obodo et al. 2016). Based on this known structure, we speculated that the ~100 bp region of

SNF5 most frequently observed to undergo telomere addition during generation of the linear fragments

represents the Core sequence and that an adjacent sequence would provide Stim function. For a

telomere to be added to the left end of the acentric fragment of chromosome II, the TG-rich sequence

must be on the bottom or 3' to 5' strand (Fig. 3A). Analysis of this strand showed that a 718 nucleotide

(nt) region including and extending 5' to the putative Core sequence (chromosome II coordinates

779784 to 780501) is unusually TG-rich. Sequences of 300 nt from the 3' to 5' strand generated as 10

base sliding windows extending across this region contain, on average, 66% T or G nucleotides

[(T+G)/300 = 0.66 ±0.05 n= 42; Figure S5A). In contrast, 300 nt sequences randomly selected from the

yeast genome (n=493) contain, on average, 50% T or G nucleotides [(T+G)/300 = 0.50±0.04], as expected

from an unbiased distribution relative to each other and the two strands (Figure S5A). To ensure that

our analysis included any potential Stim sequence, we initially tested this entire 718 nt sequence for its

ability to stimulate *de novo* telomere addition.

To determine the ability of this 718 nt sequence to stimulate telomere addition, we took

advantage of an assay in which the sequence of interest is integrated on the left arm of chromosome VII,

replacing sequences between chromosome VII coordinates 18086 and 18108 (Ngo et al. 2020). This

strain (Lydeard et al. 2010; Obodo et al. 2016) contains a single recognition site for the yeast

homothallic switching (HO) endonuclease that allows regulated generation of a double-strand break

approximately 2 kb distal to the sequence being tested (Figure 3A). Of note, the *SNF5* sequence is

inserted on chromosome VII in the inverted orientation relative to the centromere compared with its

orientation on chromosome II. However, because the endogenous location is on the right arm of

chromosome II and the insertion site is on the left arm of chromosome VII, the TG-rich sequence that

will stimulate *de novo* telomere addition is, in both cases, located on the bottom (3' to 5') strand. In

both configurations, exposure of the TG-rich strand in single-stranded DNA (for example, by resection of

a DSB occurring to the left of the sequence of interest) followed by telomerase action on that 3'

overhang will stabilize the chromosome fragment located to the right of the lesion. On chromosome II,

de novo telomere addition will create an acentric terminal fragment. On chromosome VII, *de novo*

telomere addition will truncate the chromosome. The site at which the putative SiRTA is integrated is

distal to the last essential gene on the chromosome arm (*BRR6*), ensuring that cells adding a *de novo*

telomere within the sequence of interest remain viable. Because the sequence being tested is present

both at its endogenous location and at the integrated site on chromosome VII, we deleted *RAD52* to

prevent repair by ectopic recombination.

Growth of this strain on media containing galactose induces expression of the HO endonuclease,

generating a double-strand break (DSB) distal to the putative SiRTA (Lydeard et al. 2010; Obodo et al.

2016). Correct repair restores the cleavage site, leading to a cycle of cleavage and repair that culminates

in cell death or mutation/loss of the HO site. Among the cells that survive HO cleavage, those that have

lost a *URA3* marker distal to the HO site are identified through selection for growth on media containing

5-fluoroorotic acid (5-FOA) and are subsequently screened through a combination of PCR and Sanger

sequencing to identify telomere addition events (Obodo et al. 2016). The fraction of 30 analyzed clones

that have undergone *de novo* telomere addition within the sequence of interest represents the

"relative" frequency of *de novo* telomere addition and is used as a measure of SiRTA function (Obodo et

al. 2016). Previously described SiRTAs display relative frequencies of *de novo* telomere addition ranging

from 10-35% (Obodo et al. 2016; Ngo et al. 2020).

In this case, analysis of 5-FOA resistant clones surviving HO cleavage from four independent

assays (30 clones each) showed that 30.8% ±5% had incurred truncation events mapping to the 718 bp

sequence (Figure 3B). Each of the 51 5-FOA resistant clones was analyzed by PCR using one primer that

anneals to the AC-rich telomeric strand and a second primer that anneals centromere-proximal to the

718 bp sequence (Mangahas et al. 2001). In every case, a product of the expected size was produced

which, when sequenced, revealed the direct addition of telomeric DNA to the chromosome II sequence

(File S2). Taken together, these results indicate that this region functions as a SiRTA (subsequently

referred to as the "*SNF5*" SiRTA due to its location within the *SNF5* open reading frame). 50 of the 51

telomere addition events mapped to ~180 bases at the 3' end of the 718 base region, relative to the TG-

rich strand (Figure 3 and Figure S6). This ~180 base region includes the ~100 base region in which

telomere addition was observed in the linear fragments and 69% of the HO-induced events map within 3

bases of a telomere-addition event observed on the linear molecules. However, six of nine telomere

additions in the linear fragments lie 3' to the vast majority of telomere additions observed after HO

induction (Figure S6). It is unclear why events observed in the linear fragments from the sulfate limited

chemostats are distributed differently than in the HO assay.

***De novo* telomere addition requires specific motifs associated with binding of Cdc13**

To refine the location of sequences required for *de novo* telomere addition, we tested a series of

fragments that included the region within which telomere addition occurred but were progressively

truncated at the opposite (5') end of the TG-rich strand. Fragments of 456 and 226 bp stimulated *de*

novo telomere addition as strongly as the full 718 bp sequence (Figure 3B). In contrast, no telomere

addition events were observed in a fragment of 175 bp, despite retention of virtually all sequences that

were targeted by telomerase in the context of the larger fragment (Figure 3A, B). We conclude that the

SNF5 SiRTA has a structure similar to other characterized SiRTAs, with stimulatory sequences that map

to the ~50 bp region delimited by the 226 and 175 bp fragments. We refer to this 50 bp region as the

SNF5 SiRTA Stim. Although the fraction of T and G nucleotides is elevated throughout the 718 bp region

(Figure S5B, left axis), the ratio of G to T is notably higher in the centromere proximal 226 nucleotide

region that is sufficient for *de novo* telomere addition than in the remainder of the region (Figure S5B,

right axis). Because telomeres conform to a pattern of TG_{1-3}, the higher G to T ratio is consistent with a

sequence that is more "telomere-like."

In *S. cerevisiae*, telomerase is primarily recruited to the telomere by the single-stranded binding

protein Cdc13 (Pennock et al. 2001). Binding of Cdc13 to TG-rich Stim sequences rendered single-

stranded as a result of 5' end resection following a DSB has been implicated in the high frequency of *de*

novo telomere addition within a SiRTA (Obodo et al. 2016; Epum et al. 2020). Therefore, we

hypothesized that the 50 bp *SNF5* SiRTA Stim functions through recruitment of Cdc13. The optimal

Cdc13 binding site within telomeric repeats has been characterized (5'-GTGTGGGTGTG) (Anderson et al.

2003; Eldridge et al. 2006), but single nucleotide substitutions within most positions are well tolerated.

The largest decrease in Cdc13 binding affinity is observed when G1, G3, or T4 nucleotide positions are

substituted (<u>GTGT</u>GGGTGTG), suggesting that a 5'-GxGT-3' motif may be necessary (but not sufficient)

for Cdc13 binding (Anderson et al. 2003; Eldridge et al. 2006). The *SNF5* SiRTA Stim sequence defined

above contains two 5'-GxGT-3' motifs on the TG-rich (bottom) strand (Figure 4A). To determine if these

motifs contribute to the high frequency of *de novo* telomere addition within the SiRTA, we mutated

each 5'-GxGT-3' motif to its complementary CxCA in the context of the 718 bp fragment. Remarkably,

mutation of one or both motifs nearly eliminated *de novo* telomere addition (Figure 4A, B), suggesting

that both are required for SiRTA function. This result is consistent with previous reports that SiRTA

activity likely requires more than one Cdc13 binding site (Strecker et al. 2017). We hypothesize that the

SiRTA sequence must reach a threshold of Cdc13 binding to stimulate *de novo* telomere addition and

this threshold is not met in the absence of one or both 5'-GxGT-3' motifs.

To further interrogate the requirement of Cdc13 binding to stimulate *de novo* telomere

addition, we replaced the 50 bp region of the *SNF5* SiRTA Stim sequence with two copies of the Gal4

upstream activating sequence (Stim::2xUAS) to allow specific recruitment of the Gal4 DNA binding

domain (GBD) (Zhang and Durocher 2010; Obodo et al. 2016). We introduced an empty vector, a vector

expressing GBD alone, or a vector expressing a fusion of GBD to full-length Cdc13 into cells with the

SNF5 SiRTA Stim::2xUAS fragment integrated at the chromosome VII test site. As expected, *SNF5* SiRTA

Stim::2xUAS did not support high frequency *de novo* telomere addition with empty vector or GBD alone.

Consistent with previous reports that artificial recruitment of Cdc13 to the SiRTA supports *de novo*

telomere addition (Zhang and Durocher 2010; Obodo et al. 2016), expression of the GBD-Cdc13 fusion

protein supported *de novo* telomere addition in 24.4% ±8.3% of clones tested from three independent

assays and stimulation was dependent on the presence of the UAS sequences (Figure 4C). These results

further support our hypothesis that a threshold of Cdc13 binding must be met to stimulate *de novo*

telomere addition and that without the 5'-GxGT-3' motif-containing Stim sequence, this critical

threshold is not satisfied.

Discussion:

In this work, we demonstrate that diploid laboratory yeast strains grown in sulfate-limiting

conditions generate small, extrachromosomal linear DNA fragments containing the primary sulfate

transporter *SUL1* that persist because they provide a selective growth advantage. The linear fragments

are capped at one end by sequences from the endogenous right telomere and at the other end by *de*

novo telomere addition events clustered in a region centromere proximal to *SUL1*. Fragments of similar,

but not identical, size arise in multiple independent cultures, suggesting that some sequence feature or

features in this region are responsible for the creation and/or stabilization of these linear fragments.

These results raise three intertwined questions: 1) What features of the region make it a hotspot for

telomere addition? 2) What process generates the substrate for telomere addition in the sulfate

limiting chemostats (in the absence of HO cleavage)? And 3) Why do the diploids produce a different

form of *SUL1* amplicon than their isogenic haploids?

Telomere addition: We find multiple sequence features within the *SNF5* gene that contribute to

the stabilization of linear fragments through *de novo* telomere addition. We demonstrate that a DNA

fragment from the *SNF5* gene that contains the common sites of telomere addition is sufficient to

stimulate *de novo* telomere addition at an ectopic location, similar to other characterized hotspots of *de*

novo telomere addition (SiRTAs) (Obodo et al. 2016; Ngo et al. 2020). Previous work demonstrated that

association of Cdc13 with a SiRTA is critical for the stimulation of telomere addition (Obodo et al. 2016;

Epum et al. 2020). However, the number and/or affinity of Cdc13 binding sites sufficient to support *de*

novo telomere formation are unclear. Strecker et al. (2017) placed differing lengths of telomeric repeats

adjacent to a double-strand break to determine a threshold length above which telomere addition is

stimulated. Telomere addition is strongly inhibited at sequences below that threshold length (Strecker

et al. 2017). Hypomorphic mutations in Cdc13 increase this threshold value, suggesting that the net

number and/or affinity of Cdc13 binding sites likely determines the threshold at which a sequence acts

to stimulate telomere addition (Strecker et al. 2017). Our results are consistent with this hypothesis.

Remarkably, mutation of just four nucleotides corresponding to a predicted binding site for Cdc13

(Anderson et al. 2003; Eldridge et al. 2006) nearly eliminates detectable *de novo* telomere addition at

the *SNF5* SiRTA in the ectopic assay. Our results suggest that there are likely two Cdc13 binding sites in

the ~50 bp fragment that together constitute the *SNF5* SiRTA Stim. Mutation of either site lowers the

total amount of Cdc13 binding below the threshold required for this sequence to function as a hotspot

of *de novo* telomere addition. Consistent with this interpretation, replacement of the 50 bp *SNF5* SiRTA

Stim sequence with two copies of the Gal4 upstream activating sequence supports *de novo* telomere

addition only when cells express the Gal4 DNA binding domain fused to Cdc13.

Substrate for telomere addition: We showed that *SNF5* SiRTA activity is supported by a minimal

fragment of 226 bp. This small region comprises approximately ~1% (226 bp/20.2 kb) of the total region

of chromosome VII in which a viable chromosome truncation could occur (between the HO site and the

most distal essential gene on this chromosome arm), and yet nearly one third of all truncation events

were mapped to the *SNF5* SiRTA. This clustering of *de novo* telomere additions in a small region of *SNF5*

could result from chromosome fragility in this region; however, observations from the ectopic assay

argue against this model. The *SNF5* SiRTA undergoes unusually high levels of *de novo* telomere addition

even when the initiating chromosome break is intentionally induced two kilobases distal to the eventual

site at which telomerase acts, suggesting that it is a site at which DNA ends are preferentially repaired.

We conclude that this region of *SNF5* is not necessarily more prone to double stranded breaks, but

rather has the fortuitous combination of features that permit telomere addition.

While we have not measured double stranded breaks in chemostat cultures, we have evidence

that replication errors, instead of double stranded breaks, may generate aberrant intermediates that are

subsequently targeted for repair (Brewer et al. 2015). The discontinuous nature of DNA replication forks

provides an opportunity for nascent leading strands to anneal to the lagging strand template during fork

reversal at sites of interrupted inverted repeats (described as ODIRA in Brewer et al. 2011). In a scan of

the region at the 5' end of the *SNF5* gene we find a ~5 fold increase in the density of short interrupted

inverted repeats (Figure S7A) resulting primarily from a sequence in the promoter region (5'-TGTTGTTG-

3') that is complementary to the several stretches of CAA triplet repeats within the N-terminal coding

region of *SNF5*. When compared to >10,000 regions across the genome, the *SNF5* region is enriched for

short interrupted inverted repeats (Fig S7B). The regression of a replication fork proceeding toward the

centromere from *ARS228* or another distal origin (e.g. *ARS229*) could provide multiple opportunities for

this sequence on the leading strand to find a match with a complementary sequence in the single

stranded gaps on the lagging strand and thereby create a closed loop. After completion of replication by

a fork established centromere proximal to the *SNF5* region, the hairpin capped linear could be released

(Brewer et al. 2011) and suffer one of several fates. Replication in the next cell cycle would generate an

isochromosomal palindromic linear fragment. Alternatively, recognition of the single stranded loop by

nucleases and repair enzymes could provide the substrate for telomere addition (Figure S7C,D).

Ploidy specific amplification pathways: In previous experiments with haploid laboratory strains,

we have not observed the generation of linear fragments—essentially all events were interstitial

inverted triplications on chromosome II (Payen et al. 2014). In some cases, rearrangements that would

be lethal in haploids are tolerated in diploids by the presence of a second, unrearranged chromosome

(Zhang et al. 2013). It is unlikely that a similar phenomenon is affecting the nature of *SUL1* amplification

in diploid and haploid cells as there are no essential genes beyond *SNF5* on chromosome II, so loss of

this terminal fragment should be tolerable in haploids. In point of fact, we do not detect loss of the

distal markers on chromosome II in the diploids. The aCGH profiles indicated that there were no regions

at the right end of chromosome II that were present at fewer than two copies per diploid genome and

CHEF gels did not reveal any truncated variants of chromosome II (Figures 1, S1, and S2). While these

results are consistent with repair of a broken chromosome II by homology directed repair off the

homolog or sister chromatid (perhaps by break-induced replication), it is also possible that a double

stranded break was not the initiating event. We would like to propose that the same type of ODIRA

event that can explain the events that occur in haploid cells is also occurring in the diploids, but that the

resolution of the intermediate differs between haploids and diploids. Differences in the regulation of

repair pathways between haploid and diploid cells may contribute to the different outcomes (Li and Tye

2011). There may be a race between replication of the hairpin intermediate and the processing of the

single stranded loop that is influenced by the levels of repair enzymes. In support of this notion, we

have recently observed that in some haploid wild strains with varied genetic backgrounds, linear

amplicons similar to those described here, also arising from the vicinity of the *SNF5* gene, are the

preferred amplicon (manuscript in preparation). Analysis of the transcriptomes and proteomes of these

strains could identify co-varying RNA/proteins that influence the amplicon outcome. Regardless of

mechanism, our observations in yeast may provide a useful model for the types of rearrangements that

occur in humans and other organisms.

Finally, results presented here suggest that interstitial TG-rich hotspots of *de novo* telomere

addition that are oppositely oriented relative to the chromosome arm can stabilize an acentric fragment

to provide, in some circumstances, a selective advantage. These observations highlight a possible

evolutionary consequence of SiRTA sequences in *S. cerevisiae*, regardless of orientation, by increasing

adaptation and genome plasticity.

Data Availability

Strains and plasmids are available upon request. Source data for Figures 3, 4, and S6 are provided in File

S2. Sequencing data will be available from NIH Sequence Read Archive (SRA). aCGH data will be

available from NIH Gene Expression Omnibus (GEO).

Acknowledgements

We thank James Haber and Alessandro Bianchi for generously providing us with strains and plasmids

and Natalie Guzman for initial exploratory studies.

Funding

This work was supported by National Institutes of Health awards R01GM123292 to KLF, R35GM122497

to BJB, and R01GM094306 to MJD and by National Science Foundation award 1516330 to MJD. REH was

supported by National Institutes of Health award T32GM137793. The funders had no role in study

design, data collection and analysis, decision to publish, or preparation of the manuscript.

Figure legends:

Figure 1. Amplification of *SUL1* chromosomal fragments in sulfate-limited chemostats. A) Array

Comparative Genomic Hybridization (aCGH) of the population of a diploid yeast strain 33 days after

continuous growth in a sulfate limited chemostat. The right end of chromosome II, containing the gene

for the primary sulfur transporter *SUL1*, is present in 2-3 copies in excess of the genomic copies

(normalized to 1). The presence of two discontinuities at 782 kb and 788 kb suggests two discrete

amplicons within the cell population. B) CHEF gel electrophoresis of population samples across the 33

days of continuous growth in sulfate limited medium. The ethidium bromide stained gel displays the

total yeast karyotype. The CHEF gel was Southern blotted and probed sequentially for *SUL1* and *CEN2* as

indicated. Two extrachromosomal bands of low molecular weight are detected by the *SUL1* probe from

days 17 onward. C) CHEF gel electrophoresis, under conditions to determine the sizes of

extrachromosomal amplicons, from specific days of 15 individual chemostat runs. The ethidium

bromide stained gel and Southern blot hybridized with *SUL1* reveal multiple small extrachromosomal

fragments. The sizes of extrachromosomal amplicons vary from 26 kb to 40 kb with a recurring event

producing a fragment of approximately 32 kb (red arrow).

Figure 2. Linear fragments are capped by new telomeric sequences. A) Positions of the *Eco*NI restriction

sites at the right end of chromosome II (coordinates 770 to 813 kb) and of two probes [$C_{(1-3)}A/G_{(1-3)}T$ and

SUL1 (indicated by the horizontal bars)]. Chromosome II-R telomeric sequences are indicated with gold

triangles. Proposed telomeric sequences at variable positions at the left margin of the amplicons are

indicated by the orange triangles. B) Linear fragments purified from the plug supernatant (see Figure

S3) for five cultures (S708-S712) were digested with *Eco*NI and separated on a CHEF gel. Southern blots

were sequentially hybridized with a *SUL1* probe and a telomere specific probe ($C_{(1-3)}A/G_{(1-3)}T$). Cultures

S710 and S712 contain two species of extrachromosomal molecules. Orange triangles indicate the

fragments with *de novo* telomeres. Fragments boxed in gold correspond to the native right telomere.

Figure 3. The *SNF5* SiRTA is sufficient to stimulate *de novo* telomere addition at an ectopic location. A)

Top schematic: the endogenous right arm of chromosome II; distance between the endogenous *SNF5*

sequence and *SUL1* is shown. Triangles represent the endogenous telomere. Middle schematic: lines

representing the relative size of each fragment from *SNF5* that was integrated and tested (actual size in

bp shown). Black arrows above the 718 bp fragment depict approximate locations of *de novo* telomere

addition events stimulated by HO cleavage. Blue arrows below the 718 bp fragment depict approximate

locations of spontaneous *de novo* telomere addition events from the chemostat experiments. Arrow

size is proportional to the number of independent events at that location (see Figure S6 for exact

numbers and locations). Bottom schematic: modified left arm of chromosome VII. Distances between

the integrated *SNF5* sequence and the HO cleavage site or the most distal essential gene (*BRR6*) on this

chromosome arm are shown. B) Chromosome truncation events following HO cleavage preferentially

involve *de novo* telomere addition in the SiRTA. At least 30 clones that survived HO cleavage and lost

the *URA3* marker were assayed for each experiment. The percentage of clones that incurred telomere

addition within the sequence of interest is shown. Bar height is the average of three to four

independent experiments; error bars represent standard deviations. Significant difference is indicated

(* p<0.05) by unpaired Student's *t-test* comparing 226 bp to 175 bp. See File S2 for data used to

generate graph.

Figure 4. The *SNF5* SiRTA requires Cdc13 binding to stimulate *de novo* telomere addition. A) Top: lines

representing 718, 175 bp, and *SNF5* SiRTA Stim::2XUAS fragments integrated and tested on

chromosome VII. Sites of *de novo* telomere addition are depicted as in Figure 3. Bottom: Twenty-nine

nucleotides of the TG-rich strand of the *SNF5* SiRTA Stim sequence are shown from 3' to 5'. Note that a

double-strand break that occurs to the left of this sequence will expose the TG-rich sequence in single-

stranded DNA following 5' end resection. 5'-GxGT-3' motifs and corresponding mutations are

highlighted in gray. B) Percentage of 5-FOA resistant clones that incurred *de novo* telomere addition

within the 718 bp *SNF5* SiRTA in the presence of the indicated mutation. Wild-type (WT) samples are

repeated from Figure 3 for comparison. Samples with statistically significant values by one-way ANOVA

with *post hoc* Dunnett's are indicated (**, p< 0.01). C) Percentage of 5-FOA resistant clones that

incurred *de novo* telomere addition within the 175 bp *SNF5* fragment when 2 copies of the Gal4 UAS are

added (Stim::2xUAS) or when the 175 bp fragment is present alone. Cells were transformed with

pRS414 (empty vector) or pRS414 expressing either Gal4 DNA binding domain (GBD) or GBD fused to the

N-terminus of Cdc13 as indicated. For B and C, at least 30 colonies were assayed for each experiment.

Bar height is the average of three to four independent experiments; error bars represent standard

deviations. Samples with statistically significant values by one-way ANOVA with *post hoc* Tukey's are

indicated (*, $p < 0.01$; ***, $p < 0.001$). See File S2 for data used to generate graph.

Literature Cited

Alkan C, Coe BP, Eichler EE. 2011. Genome structural variation discovery and genotyping. Nat Rev Genet.

12(5):363–376. doi:10.1038/nrg2958.

Anand R, Memisoglu G, Haber J. 2017. Cas9-mediated gene editing in *Saccharomyces cerevisiae*. Protoc

Exch. doi:10.1038/protex.2017.021a.

Anderson EM, Halsey WA, Wuttke DS. 2003. Site-directed mutagenesis reveals the thermodynamic

requirements for single-stranded DNA recognition by the telomere-binding protein Cdc13. Biochemistry.

42(13):3751–3758. doi:10.1021/bi027047c.

Araya CL, Payen C, Dunham MJ, Fields S. 2010. Whole-genome sequencing of a laboratory-evolved yeast

strain. BMC Genomics. 11(1):88. doi:10.1186/1471-2164-11-88.

Biedler JL, Meyers MB, Spengler BA. 1983. Homogeneously staining regions and double minute

chromosomes, prevalent cytogenetic abnormalities of human neuroblastoma cells. Adv Cell Neurobiol.

4:267–307. doi:10.1016/B978-0-12-008304-6.50015-4.

Blackburn EH. 1991. Structure and function of telomeres. Nature. 350(6319):569–573.

doi:10.1038/350569A0.

Brewer BJ, Payen C, DiRienzi SC, Higgins MM, Ong G, Dunham MJ, Raghuraman MK. 2015. Origin-

dependent inverted-repeat amplification: tests of a model for inverted DNA amplification. PLoS Genet.

11(12):e1005699. doi:10.1371/journal.pgen.1005699.

Brewer BJ, Payen C, Raghuraman MK, Dunham MJ. 2011. Origin-dependent inverted-repeat

amplification: A replication-based model for generating palindromic amplicons. PLoS Genet.

7(3):e1002016. doi:10.1371/journal.pgen.1002016.

Brown CJ, Todd KM, Rosenzweig RF. 1998. Multiple duplications of yeast hexose transport genes in

response to selection in a glucose-limited environment. Mol Biol Evol. 15(8):931–942.

doi:10.1093/OXFORDJOURNALS.MOLBEV.A026009.

Chakraborty A, Ay F. 2018. Identification of copy number variations and translocations in cancer cells

from Hi-C data. Bioinformatics. 34(2):338–345. doi:10.1093/BIOINFORMATICS/BTX664.

Doran CG, Pennington SR. 2022. Copy number alteration signatures as biomarkers in cancer: a review.

Biomark Med. 16(5):371–386. doi:10.2217/BMM-2021-0476.

Dorsey M, Peterson C, Bray K, Paquin CE. 1992. Spontaneous Amplification of the ADH4 gene in

Saccharomyces cerevisiae. Genetics. 132(4):943–50. doi:10.1093/genetics/132.4.943.

Eldridge AM, Halsey WA, Wuttke DS. 2006. Identification of the determinants for the specific recognition

of single-strand telomeric DNA by Cdc13. Biochemistry. 45(3):871–879. doi:10.1021/bi0512703.

Epum EA, Mohan MJ, Ruppe NP, Friedman KL. 2020. Interaction of yeast Rad51 and Rad52 relieves

Rad52-mediated inhibition of *de novo* telomere addition. PLoS Genet. 16(2):e1008608.

doi:10.1371/journal.pgen.1008608.

Gresham D, Desai MM, Tucker CM, Jenq HT, Pai DA, Ward A, DeSevo CG, Botstein D, Dunham MJ. 2008.

The repertoire and dynamics of evolutionary adaptations to controlled nutrient-limited environments in

yeast. PLoS Genet. 4(12). doi:10.1371/JOURNAL.PGEN.1000303.

Gresham D, Usaite R, Germann SM, Lisby M, Botstein D, Regenberg B. 2010. Adaptation to diverse

nitrogen-limited environments by deletion or extrachromosomal element formation of the *GAP1* locus.

Proc Natl Acad Sci U S A. 107(43):18551. doi:10.1073/PNAS.1014023107.

Hastings PJ, Lupski JR, Rosenberg SM, Ira G. 2009. Mechanisms of change in gene copy number. Nat Rev

Genet. 10(8):551. doi:10.1038/NRG2593.

Hoerr RE, Ngo K, Friedman KL. 2021. When the ends justify the means: regulation of telomere addition

at double-strand breaks in yeast. Front Cell Dev Biol. 9(18):655377. doi:10.3389/fcell.2021.655377.

Ilić M, Zaalberg IC, Raaijmakers JA, Medema RH. 2022. Life of double minutes: generation, maintenance,

and elimination. Chromosoma. epub ahead. doi:10.1007/S00412-022-00773-4.

Kao KC, Sherlock G. 2008. Molecular characterization of clonal interference during adaptive evolution in

asexual populations of *Saccharomyces cerevisiae*. Nat Genet. 40(12):1499–1504. doi:10.1038/NG.280.

Krogh BO, Symington LS. 2004. Recombination proteins in yeast. Annu Rev Genet. 38(1):233–271.

doi:10.1146/annurev.genet.38.072902.091500.

Kwan EX, Wang XS, Amemiya HM, Brewer BJ, Raghuraman MK. 2016. rDNA copy number variants are

frequent passenger mutations in *Saccharomyces cerevisiae* deletion collections and *de novo*

transformants. G3 Genes, Genomes, Genet. 6(9):2829–2838. doi:10.1534/g3.116.030296.

Li XC, Tye BK. 2011. Ploidy dictates repair pathway choice under DNA replication stress. Genetics.

187(4):1031. doi:10.1534/GENETICS.110.125450.

Lingner J, Cech TR, Hughes TR, Lundblad V. 1997. Three ever shorter telomere (*EST*) genes are

dispensable for *in vitro* yeast telomerase activity. Proc Natl Acad Sci U S A. 94(21):11190–11195.

doi:10.1073/pnas.94.21.11190.

Longtine MS, McKenzie A, Demarini DJ, Shah NG, Wach A, Brachat A, Philippsen P, Pringle JR. 1998.

Additional modules for versatile and economical PCR-based gene deletion and modification in

Saccharomyces cerevisiae. Yeast. 14(10):953–961. doi:10.1002/(SICI)1097-0061(199807)14:10<953::AID-

YEA293>3.0.CO;2-U.

Lydeard JR, Lipkin-Moore Z, Jain S, Eapen V V., Haber JE. 2010. Sgs1 and Exo1 redundantly inhibit break-

induced replication and *de novo* telomere addition at broken chromosome ends. PLoS Genet. 6(5):25.

doi:10.1371/journal.pgen.1000973.

MacDonald JR, Ziman R, Yuen RKC, Feuk L, Scherer SW. 2014. The database of genomic variants: a

curated collection of structural variation in the human genome. Nucleic Acids Res. 42(D1):D986.

doi:10.1093/nar/gkt958.

Mangahas JL, Alexander MK, Sandell LL, Zakian VA. 2001. Repair of chromosome ends after telomere

loss in *Saccharomyces*. Mol Biol Cell. 12(12):4078–4089. doi:10.1091/mbc.12.12.4078.

Miller AW, Befort C, Kerr EO, Dunham MJ. 2013. Design and use of multiplexed chemostat arrays. J Vis

Exp.(72):1–6. doi:10.3791/50262.

Narayanan V, Mieczkowski PA, Kim HM, Petes TD, Lobachev KS. 2006. The pattern of gene amplification

is determined by the chromosomal location of hairpin-capped breaks. Cell. 125(7):1283–1296.

doi:10.1016/J.CELL.2006.04.042.

Ngo K, Epum EA, Friedman KL. 2020. Emerging non-canonical roles for the Rad51–Rad52 interaction in

response to double-strand breaks in yeast. Curr Genet. 66(5):917–926. doi:10.1007/s00294-020-01081-

z.

Obodo UC, Epum EA, Platts MH, Seloff J, Dahlson NA, Velkovsky SM, Paul SR, Friedman KL. 2016.

Endogenous hot spots of *de novo* telomere addition in the yeast genome contain proximal enhancers

that bind Cdc13. Mol Cell Biol. 36(12):1750–1763. doi:10.1128/mcb.00095-16.

Payen C, Di Rienzi SC, Ong GT, Pogachar JL, Sanchez JC, Sunshine AB, Raghuraman MK, Brewer BJ,

Dunham MJ. 2014. The dynamics of diverse segmental amplifications in populations of *Saccharomyces*

cerevisiae adapting to strong selection. G3 Genes, Genomes, Genet. 4(3):399–409.

doi:10.1534/g3.113.009365.

Pennock E, Buckley K, Lundblad V. 2001. Cdc13 delivers separate complexes to the telomere for end

protection and replication. Cell. 104(3):387–396. doi:10.1016/S0092-8674(01)00226-4.

Putnam CD, Kolodner RD. 2017. Pathways and mechanisms that prevent genome instability in

Saccharomyces cerevisiae. Genetics. 206(3):1187–1225. doi:10.1534/genetics.112.145805.

Rosswog C, Bartenhagen C, Welte A, Kahlert Y, Hemstedt N, Lorenz W, Cartolano M, Ackermann S,

Perner S, Vogel W, et al. 2021. Chromothripsis followed by circular recombination drives oncogene

amplification in human cancer. Nat Genet. 53(12):1673–1685. doi:10.1038/S41588-021-00951-7.

Shaikh TH. 2017. Copy number variation disorders. Curr Genet Med Rep. 5(4):183. doi:10.1007/S40142-

017-0129-2.

Sikorski RS, Hieter P. 1989. A system of shuttle vectors and yeast host strains designed for efficient

manipulation of DNA in *Saccharomyces cerevisiae*. Genetics. 122(1):19–27.

doi:10.1080/00362178385380431.

Singer MS, Gottschling DE. 1994. *TLC1*: template RNA component of *Saccharomyces cerevisiae*

telomerase. Science. 266(5184):404-9. doi:10.1126/science.7545955.

Song K, Minami JK, Huang A, Dehkordi SR, Lomeli SH, Luebeck J, Goodman MH, Moriceau G, Krijgsman

O, Dharanipragada P, et al. 2022. Plasticity of extrachromosomal and intrachromosomal BRAF

amplifications in overcoming targeted therapy dosage challenges. Cancer Discov. 12(4):1046–1069.

doi:10.1158/2159-8290.CD-20-0936.

Song RH, Gao CQ, Zhao J, Zhang JA. 2022. An update evolving view of copy number variations in

autoimmune diseases. Front Genet. 12:794348. doi:10.3389/FGENE.2021.794348.

Stellwagen AE, Haimberger ZW, Veatch JR, Gottschling DE. 2003. Ku interacts with telomerase RNA to

promote telomere addition at native and broken chromosome ends. Genes Dev. 17(19):2384–2395.

doi:10.1101/gad.1125903.

Strecker J, Stinus S, Pliego Caballero M, Szilard RK, Chang M, Durocher D. 2017. A sharp Pif1-dependent

threshold separates DNA double-strand breaks from critically short telomeres. Elife. 6:e23783.

doi:10.7554/eLife.23783.001. https://doi.org/10.7554/eLife.23783.001.

Turner KM, Deshpande V, Beyter D, Koga T, Rusert J, Lee C, Li B, Arden K, Ren B, Nathanson DA, et al.

2017. Extrachromosomal oncogene amplification drives tumour evolution and genetic heterogeneity.

Nature. 543(7643):122–125. doi:10.1038/NATURE21356.

Wellinger RJ, Zakian VA. 2012. Everything you ever wanted to know about *Saccharomyces cerevisiae*

telomeres: beginning to end. Genetics. 191(4):1073–1105. doi:10.1534/genetics.111.137851.

Zhang F, Gu W, Hurles ME, Lupski JR. 2009. Copy number variation in human health, disease, and

evolution. Annu Rev Genomics Hum Genet. 10:451. doi:10.1146/ANNUREV.GENOM.9.081307.164217.

Zhang H, Zeidler AFB, Song W, Puccia CM, Malc E, Greenwell PW, Mieczkowski PA, Petes TD, Argueso JL.

2013. Gene copy-number variation in haploid and diploid strains of the yeast *Saccharomyces cerevisiae*.

Genetics. 193(3):785–801. doi:10.1534/GENETICS.112.146522.

Zhang W, Durocher D. 2010. *De novo* telomere formation is suppressed by the Mec1-dependent

inhibition of Cdc13 accumulation at DNA breaks. Genes Dev. 24(5):502–515. doi:10.1101/gad.1869110.