Title: Comparative immune responses to *Mycobacterium tuberculosis* in people with latent

infection or sterilizing protection

Authors: Emilie Jalbert[1], Cuining Liu[2], Vidya Mave[3,4,5], Nancy Lang[1], Anju Kagal[6], Chhaya Valvi[7],

Mandar Paradkar[3,4,5], Nikhil Gupte[3,5], Rahul Lokhande[8], Renu Bharadwaj[6], Vandana Kulkarni[3],

Amita Gupta[5], Adriana Weinberg[9]*

Affiliations:

[1]Department of Pediatrics, University of Colorado-Denver Anschutz Medical Campus, Aurora,

CO.

[2]Department of Biostatistics and Informatics, Colorado School of Public Health, University of

Colorado-Denver Anschutz Medical Campus, Aurora, CO.

[3]Byramjee Jeejeebhoy Government Medical College- Johns Hopkins University Clinical Research

Site (BJGMC-JHU CRS), Pune, Maharashtra India.

[4]Johns Hopkins India Pvt Limited (JHIPL), Pune, Maharashtra, India.

[5]School of Medicine, Center for Clinical Global Health Education (CCGHE), Johns Hopkins

University, Baltimore, Maryland, USA.

[6]Department of Microbiology, Byramjee Jeejeebhoy Government Medical College and Sassoon

General Hospital, Pune, Maharashtra, India.

[7]Department of Pediatrics, Byramjee Jeejeebhoy Government Medical College and Sassoon

General Hospital, Pune, Maharashtra, India

[8]Department of Pulmonary Medicine, Byramjee Jeejeebhoy Government Medical College and

Sassoon General Hospital, Pune, Maharashtra, India

[9]Departments of Pediatrics, Medicine and Pathology, University of Colorado-Denver Anschutz

Medical Campus, Aurora, CO

*Corresponding author: adriana.weinberg@cuanschutz.edu

Conflicts of interest: AW receives research grants from GlaxoSmithKline and Merck (moneys to

CU) and personal fees from Merck and Sequirus. No other conflicts exist.

Abstract

Tuberculosis (TB) affects 2 billion people worldwide and causes 1.5 million deaths every year. There is great need for new TB vaccines that are more efficacious than the currently licensed BCG vaccine, which provides only limited protection. Our goal was to identify potential targets for new TB vaccines by characterizing the immune responses that distinguish individuals with sterilizing protection against TB (TB-resisters), defined by presence of TB-specific immune responses and absence of latent infection, from individuals with latent TB infection (LTBI-participants).

Cryopreserved peripheral blood mononuclear cells (PBMC) from 13 TB-resisters and 10 LTBI-participants were analyzed by high dimensional spectral flow cytometry after overnight *M. tuberculosis (Mtb)* antigenic stimulation or unstimulated control. Activation of conventional and nonconventional T cells, NK cells, and antigen presenting cells (APC) was compared between the two groups. Compared with LTBI-participants, TB-resisters had significantly higher proportions of conventional and nonconventional T cells expressing granzyme B (GranzB) and PD-1, and of polyfunctional cells in unstimulated and *Mtb*-stimulated conditions. Conversely LTBI-participants had higher expression of CD25, CD69, CD107a, IL10 and IFNγ. An unbiased cluster analysis revealed higher frequency of recently described CD8+GMM+GranzB+ T cells in unstimulated PBMC from TB-resisters than LTBI-participants. APC activation revealed very few differences between TB-resisters and LTBI-participants. An exploratory analysis of responses in 14 BCG-recipients with minimal exposure to TB showed multiple differences with TB-resisters and LTBI-participants in PBMC activation; lower polyfunctionality of T cells and APC in *Mtb*-stimulated PBMC; and absence of CD8+GMM+GranzB+ T cells.

We conclude that combined increased T cell expression of GranzB and checkpoint inhibitors may contribute to immune protection against TB and may be targeted by new vaccines.

Summary

To identify immunologic targets for tuberculosis vaccines more efficacious than BCG, we

investigated the immune responses that differentiate individuals with sterile protection against *M.*

tuberculosis (Mtb) infection (TB-resisters) from those who develop latent tuberculosis infection

(LTBI-participants). TB-resisters showed higher proportions of conventional and unconventional

T cells with markers of cytotoxicity, immunologic checkpoint inhibitors, and polyfunctionality in

unstimulated and ex vivo *Mtb*-stimulated mononuclear cells. Antigen presenting cells and NK cells

did not significantly differentiate TB-resisters from LTBI-participants. A complementary analysis

of immune responses in BCG recipients with minimal exposure to tuberculosis by virtue of having

lived most of their lives in areas of low TB endemicity revealed very different unstimulated and

Mtb-stimulated immunologic profiles and lower T cell and antigen presenting cell polyfunctionality

compared to both TB-resisters and LTBI-participants. A combination of increased cytotoxic and

regulatory responses may contribute to protection against tuberculosis and may be targeted by

new vaccines.

Introduction

Tuberculosis (TB) is a major global health problem, since approximately 2 billion people are infected with *M. tuberculosis* (*Mtb*). Vaccines are the most powerful tools for limiting the morbidity and mortality of many infectious diseases, but Bacillus Calmette-Guérin (BCG)—the only licensed TB vaccine—confers limited protection[1-7]. BCG administered to infants at birth is ~80% effective in preventing disseminated and central nervous system TB. In contrast, BCG has variable efficacy against pulmonary TB when administered to infants or adults, and many studies could not demonstrate any efficacy[1-8]. BCG generates TB-specific Th1 cell-mediated immunity (CMI) that has been considered essential for protection against TB, although this concept has been disputed by recent studies[9-11]. For example, recent vaccine candidates that generate more robust *Mtb*-specific Th1 CMI than BCG did not have greater efficacy against pulmonary TB than BCG[9, 12-14]. Additional evidence that *Mtb* Th1 responses may not predict protection against *Mtb* infection is that 10 to 20% of the individuals with latent TB infection, as defined by the presence of *Mtb* CMI measured by tuberculin skin test (TST) or IFNγ release assays (IGRA), a measure of Th1 immunity, develop active TB disease over time. In contrast, 20% of individuals with household contacts with highly contagious active pulmonary TB infection do not develop clinical disease and have negative TST or IGRA[15]. These individuals, designated as TB-resisters, have antibodies and limited CMI responses to *Mtb*, confirming prior *Mtb* infection, but their T cells do not produce IFNγ in response to *Mtb* ex vivo restimulation[16, 17]. They also have genetic traits that differentiate them from people with LTBI [18-21]. TB-resisters are deemed to have sterilizing immunity against *Mtb*[22, 23]. However, the mechanisms of protection against *Mtb* infection in TB-resisters are incompletely understood[17, 24]. Identifying these mechanisms would benefit the development of new vaccines that may confer sterilizing immune protection against *Mtb*[25].

In addition to adaptive CMI, innate immunity is a critical mechanism of protection against

infections. Innate immunity is rapidly deployed and constitutes the first line of immune-mediated

defense. Moreover, innate immune responses critically contribute to the development of antigen-

specific adaptive CMI and establish a feedback mechanism that allows them to be also boosted

by adaptive CMI[26-32]. Persistent memory-like innate immune responses have been described

against viruses, tumors, and other antigens, including *Mtb*[28, 32-43]. Natural killer (NK) cells, γδ

T cells, NKT cells, invariant NKT (iNKT) cells, macrophages, monocytes (mono) and dendritic

cells (DC) have been shown to undergo clonal expansions and/or epigenetic modifications after

exposure to immunogens that can confer antigen specificity and/or allow them to activate

transcription programs that improve their functionality[43, 44]. Recent studies described a

germline encoded, mycolyl-reactive iNKT cell subset (GEMT) that recognizes the lipid glucose

monomycolate in the context of CD1b and participates in the elimination of *Mtb* from the host[40,

42, 45]. Compared to other iNKT, CD1b-restricted GEMT cells have a more diverse T-cell receptor

repertoire, which permitted the identification of GEMT cell clonal expansions in patients

recovering from TB and their persistence in the host for several years after *Mtb* elimination[40,

42]. Another nonconventional T cell subset of interest is the mucosal-associated invariant T cells

(MAIT) that developmentally share some characteristics both with iNKT and γδ T cells. MAIT can

recognize microbial riboflavin derivatives presented by the major histocompatibility-related

receptor 1 (MR1) and play a critical role in antibacterial, including *Mtb*, pulmonary defenses via

cytokine production and cytotoxicity[46-48]. MR1+ MAIT were shown to account for most IFNγ

production in response to BCG[49]. MR1- MAIT lack the receptor for antigen recognition and are

deemed to respond to secreted cytokines[50, 51]. The role of innate immune responses in

sterilizing immunity against *Mtb* infection has not been studied.

The overarching goal of our study was to identify innate and adaptive immune responses that

differentiate TB-resisters from people with LTBI (LTBI-participants) among household contacts of

active pulmonary TB cases. We leveraged a longitudinal cohort study of household contacts of

individuals with active pulmonary TB disease [Cohort For TB Research By The Indo-US Medical

Partnership Multicentric Prospective Observational Study (C-TRIUMPH)] in Pune, India, by using

advanced spectral flow cytometry technology and high-dimensional analytic tools on peripheral

blood mononuclear cells (PBMC) archived from the study participants[23, 52, 53]. To further

understand how the responses to *Mtb* in LTBI-participants and TB-resisters may differ from those

induced by BCG, we also included in our analysis a group with documented BCG vaccination and

with minimal exposure to TB.

Results

Characteristics of the study population

The study used PBMC from 13 TB-resisters and 11 LTBI-participants in C-TRIUMPH, who had

intense exposure to active pulmonary TB household contacts (**Table 1**). Importantly, TB-resisters

and LTBI-participants had similarly high exposure scores to TB (median=7), a measure of the

likelihood of acquiring *Mtb* infection from a household index case[54]. Also recruited were 14

BCG-recipients with limited exposure to TB by virtue of having spent most of their lives in the

United States or other countries with low incidence of TB. In addition to these differences in

upbringing and environmental exposures, we noted that BCG-recipients significantly differed from

C-TRIUMPH participants in race/ethnicity, but not in age or BMI.

146 **Table 1. Characteristics of the study population**

	TB-Resisters	LTBI+ participants	BCG	P-Value* (TB-Resisters vs. LTBI+)	P-Value* (three-group)
N	13	11	14		
Median years of age (range)	24 (9, 65)	29 (7, 62)	29 (20, 41)	0.81	0.71
N Female sex (%)	6 (46)	4 (36)	10 (71.4)	0.70	0.19
N Race and Ethnicity (%)					
Asian (Indian subcontinent)	13 (100)	11 (100)	0 (0)		
Asian (not Indian subcontinent)	0 (0)	0 (0)	3 (21.4)	na	<0.0001
Black	0 (0)	0 (0)	2 (14.3)		
Hispanic Non-Black	0 (0)	0 (0)	5 (35.7)		
Non-Hispanic White	0 (0)	0 (0)	4 (28.6)		
Median BMI (range)	23.3 (13.1, 30.5)	20.4 (13.5, 29.8)	22.8 (20, 30.6)	0.49	0.49
Median exposure score (range)	7 (4, 9)	7 (4, 9)	na	0.84	na
N with BCG scar (%)	9 (64.3)	5 (35.7)	na	0.24	na

148
149 * ANOVA for continuous variables and Fisher's exact for categorical variables.
150 **Abbreviations**: BCG=Bacille de Calmette-Guerin; BMI=Body mass index (kg/m^2); LTBI=latent
151 tuberculosis infection; N=number; na=not applicable
152
153

Phenotypic and functional characteristics of innate and adaptive immune responses in TB-

resisters and LTBI-participants

We used two high dimensional flow cytometry panels [T-cell and antigen-presenting-cell (APC)]

on PBMC stimulated with *Mtb* antigen or unstimulated controls (**Table S1**). In the T-cell panel, we

characterized CD4+ and CD8+ conventional T cells (Tconv), NKT, γδ T, GEMT, iNKT, MR1+ and

MR1- MAIT, and NK cells. Functionality was assessed by expression of CD25 and CD69

activation markers; PD1 immunologic checkpoint receptor; CD107a (lysosome degranulation)

and granzyme B (GranzB) cytotoxicity markers; GMCSF, IL2, IL10, IL17, IFNγ, and TNFα

cytokines; and Ki67 proliferation marker. In the APC panel, we characterized functionality of

CD14+ monocytes (Mono), CD123+ plasmacytoid DC (pDC), CD11c+CD14- total conventional

DC (cDC total), CD141+ cDC1 and CD1c+ cDC2 subsets using the following functional readouts:

CD40, CD80 and CD83 activation markers; PDL1 immunologic checkpoint ligand; and GMCSF,

IL1β, IL8, IL10, IL12p40, IL27 and TNFα cytokines. Gating trees are shown in **Fig S1**.

In *unstimulated* conditions, 10 functional cell subsets had significantly higher frequencies in TB-

resisters compared with LTBI-participants, consisting of GranzB-expressing CD4+ Tconv, CD8+

Tconv, CD8+ iNKT and CD8+ NKT; PD1-expressing CD4+ Tconv, CD8+ Tconv, CD8+ iNKT and

MR1- MAIT; CD40+ pDC; and IL1β+ Monos (**Fig 1**). In contrast, we only detected a single subset

(Ki67+MR1+ MAIT) with significantly higher frequencies in LTBI-participants compared with TB-

resisters.

Many of the functional differences between TB-resisters and LTBI-participants observed in

unstimulated PBMC persisted after *Mtb* ex vivo stimulation (**Fig 2**). Differences between TB-

resisters and LTBI-participants in *Mtb-stimulated* PBMC included higher proportions of (1)

GranzB-expressing CD4+ Tconv, CD8+ Tconv, CD8+ iNKT. CD8+ NKT and MR1- MAIT cells;

and (2) PD1-expressing CD4+ Tconv, CD8+ Tconv and MR1- MAIT in TB-resisters; and higher

proportions of (1) CD25-expressing CD4+ and CD8+ Tconv and NKT, CD8+ iNKT, and MAIT

MR1-; (2) CD107+CD8+ iNKT and NKT; (3) IFNγ-expressing CD4+ and CD8+ iNKT; (4)

CD8+IL10+ iNKT; and (5) CD69+ γδ T cells in LTBI-participants.

Mtb-memory responses were characterized by subtracting frequencies of unstimulated from *Mtb-*

stimulated corresponding PBMC. Using this definition, we identified 6 differentially expressed

functional subsets with significantly higher frequencies in LTBI-participants compared with TB-

resisters: CD25+ and CD69+ γδ T cells; CD4+CD25+ Tconv; CD8+IFNγ+ iNKT; and

CD8+CD107a+ NKT (**Fig 3**). There were no *Mtb-memory* subsets with higher frequency in TB-

resisters than in LTBI-participants.

Polyfunctionality of the immune cell subsets in TB-resisters and LTBI-participants

Polyfunctionality analyses based on Boolean gating of CD25, CD107a, GranzB, IFNγ, IL10 and

PD1 in the T cell panel showed higher marker co-expression in TB-resisters than in LTBI-

participants in *unstimulated* conditions on CD8+ iNKT, MR1- MAIT, CD8+ NKT and CD4+ and

CD8+ Tconv (**Fig 4A**). In *Mtb-stimulated* conditions, CD8+ iNKT, MR1- MAIT, CD8+ NKT, and

CD4+ and CD8+ Tconv maintained higher polyfunctionality in TB-resisters than LTBI-participants.

LTBI-participants did not have any subsets with higher polyfunctionality than TB-resisters. There

were no differences in polyfunctionality of NK and γδ T cells in Mtb-stimulated or unstimulated

conditions between TB-resisters and LTBI-participants (**Fig S2**).

In the APC analysis, we evaluated the expression of CD40, CD80, IL1β, IL27, PDL1 and TNFα

on Mono, cDC1, cDC2, cDC total, and pDC. **Figure 4B** shows that TB-resisters had higher

proportions of polyfunctional Mono than LTBI-participants in *unstimulated* conditions. No other

differences were observed between TB-resisters and LTBI-participants.

Coordination of the immune responses in TB-resisters and LTBI-participants

207
We complemented the analysis of manually gated PBMC subsets with Phenograph unbiased

analysis, which identified 8 T-cell and 1 APC clusters that significantly differed between TB-

resisters and LTBI-participants in *unstimulated* PBMC; 3 T-cell and 1 APC clusters in *Mtb-*

stimulated conditions; and 5 T-cell and 3 APC *Mtb-memory* clusters (**Fig S3**). Most clusters

confirmed findings already observed in the manually gated analysis, with the following exceptions:

(1) Two T-cell and one APC *Mtb-memory* clusters (**Fig S3**) with higher relative frequencies in TB-

resisters than LTBI-participants, in contrast to the manual gating analysis that did not identify

excess *Mtb-memory* subsets in TB-resisters compared with LTBI-participants; and (2) A

CD8+GMM+GranzB+ Tconv subset that had not been included in the prespecified manual gating

analysis (**Fig 5 and S3**). This subset had higher frequency in *unstimulated* PBMC from TB-

resisters compared with LTBI-participants (FDR-corrected p=0.008).

Immune responses in BCG-recipients

The analysis of functional phenotypes in BCG-recipients showed significant differences with TB-

resisters in 54 *unstimulated,* 41 *Mtb-stimulated*, and 41 *Mtb-memory* PBMC subset responses

(**Fig S4**). Compared to LTBI-participants, BCG-recipients showed differences in 56 *unstimulated*,

32 *Mtb-stimulated*, and 24 *Mtb-memory* responses (**Fig S5**).

The comparison of polyfunctional NK and T cells in BCG-recipients and TB-resisters showed

significantly higher frequencies of polyfunctional MR1+ and MR1- MAIT and CD4+ and CD8+

NKT and Tconv in TB-resisters both in *unstimulated* and *Mtb-stimulated* PBMC, while BCG-

recipients had higher frequencies of polyfunctional CD4+ iNKT cells (**Fig 6A**). Compared with

BCG-recipients, LTBI-participants had higher frequencies of polyfunctional MR1+ MAIT cells in

unstimulated PBMC, while BCG-recipients had higher frequencies of CD4+ iNKT (**Fig 6B**). In *Mtb-*

stimulated conditions, LTBI-participants showed higher frequencies of polyfunctional MR1+ MAIT

and CD4+ Tconv, while BCG-recipients had higher frequencies of polyfunctional CD4+ and CD8+

iNKT.

The polyfunctionality analysis of APC showed that compared with BCG-recipients, TB-resisters

had higher proportions of polyfunctional Monos, cDC1, cDC2 and pDC in *unstimulated* PBMC

(**Fig 6C**). After *Mtb-stimulation* the cDC1 and cDC2 also had higher polyfunctionality in TB-

resisters and none of the APC subsets had higher polyfunctionality in BCG-recipients. Compared

to BCG-recipients LTBI-participants had higher frequencies of polyfunctional cDC1 in

unstimulated PBMC, while BCG-recipients had higher frequencies of polyfunctional cDC1 and

cDC2 after *Mtb stimulation* (**Fig 6D**).

One of the two recent descriptions of GMM+ Tconv showed their presence in response to BCG

administration in infants, but not in adults[55]. On comparing the frequency of

CD8+GMM+GranzB+ T cells in BCG-recipients with TB-resisters and LTBI-participants in our

study, we also found very small frequencies of this cell subset (≤0.005% out of total lymphocytes)

in BCG-recipients, significantly lower compared with either of the *Mtb*-exposed groups (FDR-

corrected p≤0.00013; **Fig S7**).

Discussion

The goal of this study was to characterize responses unique to TB-resisters compared to LTBI-

participants that may be targeted by new TB vaccines. A consistent finding was increased

GranzB-expressing conventional and nonconventional T cells in *Mtb-stimulated* and *unstimulated*

PBMC of TB-resisters. This hallmark was observed in the prespecified manual gating and in the

unsupervised cluster analyses and included expression of GranzB by Tconv, NKT and iNKT cells.

The *Mtb-memory* analysis reveal higher cytotoxic responses in TB-resisters using the unbiased

cluster analysis but not the manual gating. Collectively, these findings suggest that TB-resisters

may mount higher and/or faster cytotoxic responses upon exposure to *Mtb* than LTBI-participants,

which may contribute to the clearance of the infectious agent before establishing latency.

Of special interest was a population of CD8+GMM+GranzB+ Tconv that was not part of the

prespecified manual gating but was identified by the Phenograph cluster analysis. This cell

subset, which has only recently been described, was found in people with active TB, in whom

CD8+GMM+ T cells displayed upregulated cytotoxic transcription profiles[56]. The authors

concluded that this cell subset may contribute to the clearance of *Mtb*-infected host cells. Notably,

we found higher proportions of CD8+GMM+GranzB+ T cells in TB-resisters compared with LTBI-

participants, suggesting that this cell subset may contribute to the sterilizing immune protection

of TB-resisters against *Mtb* infection.

TB-resisters also had higher proportions of PD1-expressing Tconv, iNKT and MR1- MAIT than

LTBI-participants in *unstimulated* PBMC and after ex vivo *Mtb stimulation*. PD1 is an immunologic

checkpoint inhibitor commonly expressed on activated T cells[57]. PD1 expression contributes to

quenching the immune response after removal of the stimulating agent, limiting the inflammatory

response and tissue destruction. Recent reports showed an association of increased T cell

activation with TB morbidity and with increased risk of active *Mtb* infection in BCG-recipients[58-

60]. Moreover, PD1 deficient mice as well as people treated with PD1/PDL1 blocking agents have

increased risk of *Mtb* infection, morbidity, and reactivation[61-63]. Collectively, these observations

suggest that the increased expression of PD1 in TB-resisters may limit their susceptibility to *Mtb*

tissue destruction and, perhaps, propagation of infection.

A distinguishing characteristic of the immune response in TB-resisters from LTBI-participants was

polyfunctionality. TB-resisters had higher proportions of polyfunctional Tconv and

nonconventional T cells both in *Mtb-stimulated* and *unstimulated* conditions. Polyfunctional CD4+

Tconv, characterized by expression of IL2, IFNγ and/or TNFα have been previously associated

with protection against active TB infection in animal models[64]. However, the role of

polyfunctional CD4+ T cell responses in protection against *Mtb* infection in humans remains

uncertain, particularly since a recent study showed that IL2, IFNγ and/or TNFα polyfunctional

CD4+ T cell responses generated by BCG did not correlate with protection against TB disease in

vaccinated infants[11]. In our study, cytokine production made a minor contribution to the

polyfunctional T cell responses, which predominantly expressed GranzB, CD107a, CD25 and

PD1, and involved mostly CD8+ T cells, including NKT, iNKT and Tconv. In addition, CD4+ Tconv

and MR1- MAIT cells also showed higher polyfunctionality in TB-resisters than LTBI-participants

expressing the same markers as the CD8+ Tconv. These findings suggest that high proportions

of cytotoxic Tconv and nonconventional T cells may contribute to the immune protection of TB-

resisters against *Mtb* infection.

Mtb-memory T cell responses identified by the expression of single activation markers were

higher in LTBI-participants than in TB-resisters, although two *Mtb-memory* CD4+ Tconv clusters

revealed by the Phenograph analysis had higher frequencies in TB-resisters than in LTBI-

participants. These data confirm that TB-resisters develop *Mtb*-specific T cell memory, but in

general *Mtb-memory* responses are higher in LTBI-participants. This finding may be related to

the observation that 25% of people with LTBI have evidence of active *Mtb* replication, which

continuously exposes the immune system to *Mtb* antigens and maintains high levels of memory

cells[65-68].

APC displayed few differentiating features between TB-resisters and LTBI-participants. In

unstimulated PBMC, TB-resisters had higher frequencies of activated and polyfunctional Mono

by manual gating and Boolean analyses, respectively, and cDC2 by cluster analysis. However,

no differences were found in *Mtb-stimulated* responses between TB-resisters and LTBI-

participants. It is important to note that *Mtb* contains multiple toll-like receptor (TLR) agonists,

which hinders the discrimination of *Mtb-stimulated* responses from nonspecific stimulation TLR-

mediated activation of APC. Nevertheless, the Phenograph analysis identified an activated Mono

cluster with higher frequency in TB-resisters than LTBI-participants.

The analysis of *Mtb-memory* responses in BCG-recipients using single markers of activation

showed both higher and lower responses in multiple T-cell and APC subsets compared with TB-

resisters or LTBI-participants. These differences may have been related to the difference in

environmental exposures, including *Mtb*, in addition to genetic backgrounds of BCG-recipients

compared with TB-resisters or LTBI-participants. The Boolean analysis of polyfunctional

responses narrowed down the differences between BCG-recipients and LTBI-participants or TB-

resisters. Overall, BCG-recipients had higher frequencies of *unstimulated* and *Mtb-stimulated*

iNKT polyfunctional cells than both LTBI-participants and TB-resisters. BCG-recipients also had

lower frequencies of *unstimulated* and/or *Mtb-stimulated* Tconv, $\gamma\delta$, NKT, and/or MAIT

polyfunctional cells compared to both LTBI-participants and TB-resisters, but a larger number of

polyfunctional subsets differentiated BCG-recipients from TB-resisters than LTBI-participants.

Compared to TB-resisters, BCG-recipients also had universally lower frequencies of *stimulated*

and *Mtb-stimulated* polyfunctional APC subsets. Compared to LTBI-participants, BCG-recipients

had lower frequencies only of *stimulated* and *Mtb-stimulated* polyfunctional cDC1 and higher

frequencies of polyfunctional cDC2. Collectively, these findings suggest superior polyfunctionality

of T cells and APC in TB-resisters compared with BCG-recipients.

Our study has both strengths and limitations. The main limitations consisted of the small number

of participants and different demographic features of BCG-recipients compared with C-TRIUMPH

participants (e.g., race/ethnicity). Although our study did not measure antibody responses, which

have been recently shown to differentiate TB-resisters from LTBI-participants[17, 69], the CMI

analysis was comprehensive and revealed new findings unique in TB-resisters.

We conclude that both innate and adaptive responses distinguish TB-resisters from LTBI-

participants. Unique elements of the cell-mediated immune responses to *Mtb* in TB-resisters are

high cytotoxic potential, expression of immunologic checkpoint inhibitors and polyfunctionality of

Tconv and nonconventional T cells, suggesting that some or all may be key factors in an effective

immune response against *Mtb*. Although TB-resisters displayed these characteristics even

without ex vivo *Mtb stimulation*, it is reasonable to propose that a vaccine that increases some or

all these responses in the form of *Mtb-memory* may be more effective than BCG.

Materials and Methods

<u>Participants.</u> Cryopreserved PBMC were obtained from 13 TB-resisters and 11 LTBI-participants

in C-TRIUMPH [23, 52, 53]. LTBI-participants were defined by positive IGRA and/or TST at entry.

TB-resisters had negative IGRA and TST at entry and over the 2 years of follow up. BCG-

recipients whose immunization status was documented by medical history, who lived most of their

lives outside of TB endemic areas, and who had negative IGRA were enrolled at the University of

Colorado Anschutz Medical Campus (CU-AMC).

<u>IGRA</u> testing used Quantiferon Gold in Tube kit (Qiagen) performed as per manufacturer's

instructions.

<u>Flow Cytometry.</u> Testing conditions were optimized using a convenience sample of PBMC from

LTBI-participants and donors without previous exposure to TB or BCG submitted to stimulation

for several time intervals with different concentrations of irradiated *Mtb*, whole cell lysate, cell

membrane, PPD and BCG. *Mtb* cell membrane provided the best discrimination between LTBI-

participants and TB-unexposed and was selected for this study. PBMC were thawed in serum-

free media (AIM V, Gibco) supplemented with benzonase (50 units/mL, Millipore). PBMC were

resuspended at $2*10^6$ cells/ml and stimulated with cell membrane (BEI Resources Strain

CDC1551, Cell Membrane Fraction, NR-14832) at 20ug/ml in AIM V or unstimulated overnight.

Brefeldin-A and Monensin (SIGMA) at 5ug/ml each were added for the last 4h of culture. For the

T-cell panel, CD107a-BV785 (Biolegend) was added for the last 4h of culture. Cells were washed

with PBS, stained with Zombie NIR Fixable Viability dye (Biolegend), and washed with PBS+1%

albumin. For the APC panel, cells were treated with Human TruStain FcX (Biolegend), surface-

stained with CD16-eFluor450, CD83-PerCP-eFluor710 (Thermo Fisher/eBioscience), CD40-

BV480, CD141-BV750, CD11c-BB515 (BD Biosciences), CD80-BV510, CD1c-BV605, CD123-

BV650, PDL1-BV785, HLA-DR-AlexaFluor488, CD3-, CD19-, CD20- and CD56-PerCP-Cy5.5,

CD14-APC-Fire750 (Biolegend), brilliant stain buffer plus (BD Biosciences) and True-stain

monocyte blocker (Biolegend). Cells were washed and treated with eBioscience Foxp3 Staining

Buffer Set (Thermo Fisher/eBioscience), then stained with IL8-BV421, IL10-BV711, IL1β-PE (BD

Biosciences), GMCSF-PE-Dazzle594, IL12p40-AlexaFluor647 (Biolegend), IL27-APC, TNFα-

AlexaFluor700 (Thermo Fisher/eBioscience) and brilliant stain buffer plus. For the T-cell panel,

cells were treated with 50% human AB serum (Gemini) in PBS+1% albumin, washed and surface-

stained with tetramers: CD1b-unloaded-FITC, CD1b-GMM-APC, and MR1-BV421. Cells were

washed, stained with TCR Vα7.2-BV605, TCR Vα24-Jα18-PE-Cy7 (Biolegend) and TCR γδ-PE

(BD Biosciences) followed by CD16-eFluor450, CD3-AlexaFluor532 (Thermo

Fisher/eBioscience), CD56-BV480, CD25-BB515 (BD Biosciences), CD8-BV510, PD1-BV711,

CD69-PE-Cy5, CD161-AlexaFluor700 (Biolegend) and brilliant stain buffer plus. Cells were

washed, treated with Foxp3 Staining Buffer Set, then stained with IL17-BV570, IL2-BV650, IL10-

PE-Dazzle594, GranzymeB-APC-Fire750 (Biolegend), IFNγ-BV750, GMCSF-AlexaFluor647 (BD

Biosciences), Ki67-PerCP-eFluor710, TNFα-PerCP-Cy5.5 (Thermo Fisher/eBioscience) and

brilliant stain buffer plus. Cells from both panels were washed and fixed with PBS+1%

paraformaldehyde, acquired on the Cytek Aurora™ (Cytek Biosciences), and analyzed with

FlowJo (Becton Dickinson). Two leucopack controls were used in each run to ensure inter-run

reproducibility and to examine technical effects.

Boolean analysis. Boolean combinatorial gates were created using the 6 cytokines/activation

markers, generating 64 distinct activation phenotypes. Graphical representation was performed

using SPICE 6.0 software[70]. The data were analyzed using permutations (10,000 iterations)

included in the software.

UMAP analysis. For T cell clustering, we used 67,000 down-sampled events from 13 TB-resisters

and 9 LTBI-participants for a total of 2,948,000 events in the concatenated file. For APC

clustering, we used data from 11 TB-resisters and 8 LTBI-participants for a total of 502,479 events

in the concatenated file. Parameters were rescaled to ArcSinh prior to running UMAP with

phenograph in FlowJo on the concatenated file using parameters, Euclidean distance, 15 nearest

neighbors and minimum distance of 0.5[71].

Statistical Testing of Cell Lineage and Function Between TB-Resisters and LTBI-participants. To

compare the average cell lineage frequencies obtained from manual and data-driven gating

between groups, we used beta regression (*betareg* R package, v3.1-3) with a logit-link to model

cell lineage percent (regression outcome; between 0 and 100% of events) on group (primary

explanatory variable of interest; LTBI-participant versus TB-resisters), adjusting for participant

age in years as a covariate [72]. Similarly, to test for functional differences among *unstimulated*

conditions, we modeled percent positivity (outcome; between 0 and 100% of cells in given lineage

expressing functional marker) on group, correcting for age. "Rare" outcomes (i.e., outcomes of

0% in over 1/3rd of participants) were excluded from testing due to their low variability among our

participants. For each set of statistical tests, we defined significant between-group differences as

a non-zero group regression coefficient (Wald test, multiple testing adjustment using the

Benjamini-Hochberg false discovery rate < 0.05)[73].

To examine differences in functional markers under *Mtb-stimulated* conditions, we

repeated this modeling procedure but with the stimulated percent positivity values as the

regression outcome variable, adjusting for age. We also examined differences in *Mtb-memory* by

additionally adjusting for each participant's corresponding baseline values (unstimulated percent

positivity values in their paired control sample) as a covariate. The *Mtb-memory* analysis thus

emphasizes markers that were not already differentially abundant between groups prior to

stimulation, or stimulation effects potentially masked by unstimulated differences. Adjusting for

baseline levels are conceptually similar to "subtracting" the unstimulated values from the Mtb-

stimulated values; as a consequence, we visualize Mtb-memory with Δ-values, defined by

stimulated frequencies minus unstimulated frequencies. Functional markers with limited changes

following Mtb-stimulation (i.e., Δ-values of magnitude ≤ 0.1% in over 1/3rd of participants) were

excluded from testing due to their low variability among our participants and tendency to

recapitulate unstimulated tests. Significant between-group differences in Mtb-memory were

based on FDR < 0.05.

These modeling procedures was also repeated for the TB-resister versus BCG recipient

comparisons and the LTBI-participants versus BCG recipient comparisons. All analyses were

performed in the R statistical package v4.0.2 unless otherwise specified. Additional

methodological details and analysis code are publicly available at

https://github.com/chooliu/Sterile_vs_Latent_TB_Immunity.

Declaration of approval. The study was approved by the Colorado Multiple Institutional Review

Board (COMIRB) and by the IRBs of the Byramjee Jeejeebhoy Govt. Medical College, India and

Johns Hopkins Medicine. All participants signed informed consent.

Author contributions: AW designed the study, oversaw the laboratory assays, analyzed the

data and prepared the manuscript. EJ developed the flow cytometry assays, analyzed the data

and contributed to manuscript preparation. CL performed the statistical analysis and contributed

to manuscript preparation. NL conducted regulatory and clinical activities at CU-AMC. For the

clinical research cohort activities in Pune, India: VM, MP, RL, CV oversaw the clinical enrollment

and follow-up activities; VK, RB, AK conducted the laboratory activities and specimen

management; NG conducted the data management activities; MP oversaw all the day-to-day

study coordination activities; and AG, VM obtained the funding for the clinical cohort protocol,

and AG, VM, and MP developed the cohort protocol and monitored the overall study conduct at

BJGMC JHU CRS. All authors reviewed and approved the manuscript.

Acknowledgments: The authors thank Ms. Sarah Lazarus for technical support; Drs. Seshadri

and Moody for technological and material support on GEMT tetramers[74] and GMM,

respectively; and the NIH tetramer core for MR1 tetramers and for CD1b monomers used for

generation of GEMT tetramers. The MR1 tetramer technology was developed jointly by Dr.

James McCluskey, Jamie Rossjohn, and David Fairlie[75], and the material was produced by

the NIH Tetramer Core Facility as permitted to be distributed by the University of Melbourne.

The following reagent was obtained through BEI Resources, NIAID, NIH: *Mycobacterium*

tuberculosis, Strain CDC1551, Cell Membrane Fraction, NR-14832.

Funding: The study was supported by NIAID grant 1R21AI134129 (AW). CTRIUMPH is part of

the Regional Prospective Observational Research for Tuberculosis consortium for India funded

with Federal funds from the Government of India's (GOI) Department of Biotechnology (DBT), the

Indian Council of Medical Research (ICMR), the USA National Institutes of Health (NIH), the

National Institute of Allergy and Infectious Diseases (NIAID), the Office of AIDS Research (OAR),

and distributed in part by CRDF Global (USB1-31147-XX-13 to AG). This work was also

supported by the NIH funded Johns Hopkins Baltimore-Washington-India Clinical Trials Unit for

NIAID Networks (UM1AI069465 to AG). Rahul Lokhande and Chhaya Valvi were supported by

the BJGMC JHU HIV TB Program funded by the Fogarty International Center, NIH

(D43TW009574). The contents of this publication are solely the responsibility of the authors and

do not represent the official views of the DBT, the ICMR, the NIH, or CRDF Global. Any mention

of trade names, commercial projects or organizations does not imply endorsement by any of the

sponsoring organizations. The funders had no role in study design, data collection and analysis,

decision to publish, or preparation of the manuscript.

Data and materials availability: The flow cytometry was partially uploaded to ImmPort

(https://www.immport.org) under study accession SDY1694. Information about C-TRIUMPH can

be found at

https://main.ccghe.net/content/report-c-triumph-cohort-tuberculosis-research-indo-us-medical-

partnership.

References

1. Fine PE. Variation in protection by BCG: implications of and for heterologous immunity.
Lancet. 1995;346(8986):1339-45. Epub 1995/11/18. PubMed PMID: 7475776.
2. Barreto ML, Pereira SM, Pilger D, Cruz AA, Cunha SS, Sant'Anna C, et al. Evidence of
an effect of BCG revaccination on incidence of tuberculosis in school-aged children in Brazil:
second report of the BCG-REVAC cluster-randomised trial. Vaccine. 2011;29(31):4875-7. Epub
2011/05/28. doi: 10.1016/j.vaccine.2011.05.023. PubMed PMID: 21616115.
3. Basu Roy R, Sotgiu G, Altet-Gomez N, Tsolia M, Ruga E, Velizarova S, et al. Identifying
predictors of interferon-gamma release assay results in pediatric latent tuberculosis: a protective
role of bacillus Calmette-Guerin?: a pTB-NET collaborative study. American journal of
respiratory and critical care medicine. 2012;186(4):378-84. Epub 2012/06/16. doi:
10.1164/rccm.201201-0026OC. PubMed PMID: 22700862; PubMed Central PMCID:
PMC3443812.
4. Chan PC, Yang CH, Chang LY, Wang KF, Kuo YC, Lin CJ, et al. Lower prevalence of
tuberculosis infection in BCG vaccinees: a cross-sectional study in adult prison inmates. Thorax.
2013;68(3):263-8. Epub 2012/09/29. doi: 10.1136/thoraxjnl-2012-202208. PubMed PMID:
23019256.
5. Eisenhut M, Paranjothy S, Abubakar I, Bracebridge S, Lilley M, Mulla R, et al. BCG
vaccination reduces risk of infection with Mycobacterium tuberculosis as detected by gamma
interferon release assay. Vaccine. 2009;27(44):6116-20. Epub 2009/09/01. doi:
10.1016/j.vaccine.2009.08.031. PubMed PMID: 19715782.
6. Roy A, Eisenhut M, Harris RJ, Rodrigues LC, Sridhar S, Habermann S, et al. Effect of
BCG vaccination against Mycobacterium tuberculosis infection in children: systematic review
and meta-analysis. BMJ. 2014;349:g4643. Epub 2014/08/07. doi: 10.1136/bmj.g4643. PubMed
PMID: 25097193; PubMed Central PMCID: PMC4122754.
7. Soysal A, Millington KA, Bakir M, Dosanjh D, Aslan Y, Deeks JJ, et al. Effect of BCG
vaccination on risk of Mycobacterium tuberculosis infection in children with household
tuberculosis contact: a prospective community-based study. Lancet. 2005;366(9495):1443-51.
Epub 2005/10/26. doi: 10.1016/S0140-6736(05)67534-4. PubMed PMID: 16243089.
8. Dockrell HM, Smith SG. What Have We Learnt about BCG Vaccination in the Last 20
Years? Front Immunol. 2017;8:1134. Epub 2017/09/29. doi: 10.3389/fimmu.2017.01134.
PubMed PMID: 28955344; PubMed Central PMCID: PMCPMC5601272.
9. Karp CL, Wilson CB, Stuart LM. Tuberculosis vaccines: barriers and prospects on the
quest for a transformative tool. Immunol Rev. 2015;264(1):363-81. Epub 2015/02/24. doi:
10.1111/imr.12270. PubMed PMID: 25703572; PubMed Central PMCID: PMC4368410.
10. Matsumiya M, Satti I, Chomka A, Harris SA, Stockdale L, Meyer J, et al. Gene
expression and cytokine profile correlate with mycobacterial growth in a human BCG challenge
model. J Infect Dis. 2015;211(9):1499-509. Epub 2014/11/09. doi: 10.1093/infdis/jiu615.
PubMed PMID: 25381367; PubMed Central PMCID: PMC4392868.
11. Kagina BM, Abel B, Scriba TJ, Hughes EJ, Keyser A, Soares A, et al. Specific T cell
frequency and cytokine expression profile do not correlate with protection against tuberculosis
after bacillus Calmette-Guerin vaccination of newborns. Am J Respir Crit Care Med.
2010;182(8):1073-9. Epub 2010/06/19. doi: 10.1164/rccm.201003-0334OC. PubMed PMID:
20558627; PubMed Central PMCID: PMCPMC2970848.
12. Bhatt K, Verma S, Ellner JJ, Salgame P. Quest for correlates of protection against
tuberculosis. Clin Vaccine Immunol. 2015;22(3):258-66. Epub 2015/01/16. doi:
10.1128/CVI.00721-14. PubMed PMID: 25589549; PubMed Central PMCID: PMC4340894.
13. Tameris M, Geldenhuys H, Luabeya AK, Smit E, Hughes JE, Vermaak S, et al. The
candidate TB vaccine, MVA85A, induces highly durable Th1 responses. PLoS One.

2014;9(2):e87340. doi: 10.1371/journal.pone.0087340. PubMed PMID: 24498312; PubMed
Central PMCID: PMC3911992.

14.   Tameris MD, Hatherill M, Landry BS, Scriba TJ, Snowden MA, Lockhart S, et al. Safety
and efficacy of MVA85A, a new tuberculosis vaccine, in infants previously vaccinated with BCG:
a randomised, placebo-controlled phase 2b trial. Lancet. 2013;381(9871):1021-8. doi:
10.1016/S0140-6736(13)60177-4. PubMed PMID: 23391465.

15.   Tait DR, Hatherill M, Van Der Meeren O, Ginsberg AM, Van Brakel E, Salaun B, et al.
Final Analysis of a Trial of M72/AS01E Vaccine to Prevent Tuberculosis. N Engl J Med.
2019;381(25):2429-39. Epub 2019/10/30. doi: 10.1056/NEJMoa1909953. PubMed PMID:
31661198.

16.   Lu LL, Smith MT, Yu KKQ, Luedemann C, Suscovich TJ, Grace PS, et al. IFN-gamma-
independent immune markers of Mycobacterium tuberculosis exposure. Nat Med.
2019;25(6):977-87. Epub 2019/05/22. doi: 10.1038/s41591-019-0441-3. PubMed PMID:
31110348; PubMed Central PMCID: PMCPMC6559862.

17.   Simmons JD, Stein CM, Seshadri C, Campo M, Alter G, Fortune S, et al. Immunological
mechanisms of human resistance to persistent Mycobacterium tuberculosis infection. Nat Rev
Immunol. 2018;18(9):575-89. Epub 2018/06/14. doi: 10.1038/s41577-018-0025-3. PubMed
PMID: 29895826; PubMed Central PMCID: PMCPMC6278832.

18.   Stein CM, Zalwango S, Malone LL, Won S, Mayanja-Kizza H, Mugerwa RD, et al.
Genome scan of M. tuberculosis infection and disease in Ugandans. PLoS One.
2008;3(12):e4094. Epub 2009/01/01. doi: 10.1371/journal.pone.0004094. PubMed PMID:
19116662; PubMed Central PMCID: PMCPMC2605555.

19.   Seshadri C, Sedaghat N, Campo M, Peterson G, Wells RD, Olson GS, et al.
Transcriptional networks are associated with resistance to Mycobacterium tuberculosis
infection. PLoS One. 2017;12(4):e0175844. Epub 2017/04/18. doi:
10.1371/journal.pone.0175844. PubMed PMID: 28414762; PubMed Central PMCID:
PMCPMC5393882.

20.   Moller M, Kinnear CJ, Orlova M, Kroon EE, van Helden PD, Schurr E, et al. Genetic
Resistance to Mycobacterium tuberculosis Infection and Disease. Front Immunol. 2018;9:2219.
Epub 2018/10/16. doi: 10.3389/fimmu.2018.02219. PubMed PMID: 30319657; PubMed Central
PMCID: PMCPMC6170664.

21.   Cobat A, Poirier C, Hoal E, Boland-Auge A, de La Rocque F, Corrard F, et al. Tuberculin
skin test negativity is under tight genetic control of chromosomal region 11p14-15 in settings
with different tuberculosis endemicities. J Infect Dis. 2015;211(2):317-21. Epub 2014/08/22. doi:
10.1093/infdis/jiu446. PubMed PMID: 25143445; PubMed Central PMCID: PMCPMC4279780.

22.   Stein CM, Zalwango S, Malone LL, Thiel B, Mupere E, Nsereko M, et al. Resistance and
Susceptibility to Mycobacterium tuberculosis Infection and Disease in Tuberculosis Households
in Kampala, Uganda. Am J Epidemiol. 2018;187(7):1477-89. Epub 2018/01/06. doi:
10.1093/aje/kwx380. PubMed PMID: 29304247; PubMed Central PMCID: PMCPMC6031055.

23.   Mave V, Chandrasekaran P, Chavan A, Shivakumar S, Danasekaran K, Paradkar M, et
al. Infection free "resisters" among household contacts of adult pulmonary tuberculosis. PLoS
One. 2019;14(7):e0218034. Epub 2019/07/19. doi: 10.1371/journal.pone.0218034. PubMed
PMID: 31318864; PubMed Central PMCID: PMCPMC6638997.

24.   Meermeier E, Lewinsohn D. Early clearance versus control: what is the meaning of a
negative tuberculin skin test or interferon-gamma release assay following exposure to
Mycobacterium tuberculosis? [version 1; peer review: 2 approved]. F1000Research.
2018;7(664). doi: 10.12688/f1000research.13224.1.

25.   Jeyanathan M, Yao Y, Afkhami S, Smaill F, Xing Z. New Tuberculosis Vaccine
Strategies: Taking Aim at Un-Natural Immunity. Trends Immunol. 2018;39(5):419-33. Epub
2018/02/13. doi: 10.1016/j.it.2018.01.006. PubMed PMID: 29429859.

26.   Adams NM, O'Sullivan TE, Geary CD, Karo JM, Amezquita RA, Joshi NS, et al. NK Cell
Responses Redefine Immunological Memory. J Immunol. 2016;197(8):2963-70. doi:
10.4049/jimmunol.1600973. PubMed PMID: 27824591; PubMed Central PMCID:
PMCPMC5108457.
27.   Soudja SM, Chandrabos C, Yakob E, Veenstra M, Palliser D, Lauvau G. Memory-T-cell-
derived interferon-gamma instructs potent innate cell activation for protective immunity.
Immunity. 2014;40(6):974-88. Epub 2014/06/17. doi: 10.1016/j.immuni.2014.05.005. PubMed
PMID: 24931122; PubMed Central PMCID: PMC4105986.
28.   Berrien-Elliott MM, Wagner JA, Fehniger TA. Human Cytokine-Induced Memory-Like
Natural Killer Cells. J Innate Immun. 2015;7(6):563-71. doi: 10.1159/000382019. PubMed
PMID: 25924651.
29.   Gottschalk C, Mettke E, Kurts C. The Role of Invariant Natural Killer T Cells in Dendritic
Cell Licensing, Cross-Priming, and Memory CD8(+) T Cell Generation. Front Immunol.
2015;6:379. doi: 10.3389/fimmu.2015.00379. PubMed PMID: 26284065; PubMed Central
PMCID: PMC4517377.
30.   Slauenwhite D, Johnston B. Regulation of NKT Cell Localization in Homeostasis and
Infection. Front Immunol. 2015;6:255. doi: 10.3389/fimmu.2015.00255. PubMed PMID:
26074921; PubMed Central PMCID: PMC4445310.
31.   Waggoner SN, Cornberg M, Selin LK, Welsh RM. Natural killer cells act as rheostats
modulating antiviral T cells. Nature. 2012;481(7381):394-8. doi: 10.1038/nature10624. PubMed
PMID: 22101430; PubMed Central PMCID: PMC3539796.
32.   Zheng J, Liu Y, Lau YL, Tu W. gammadelta-T cells: an unpolished sword in human anti-
infection immunity. Cell Mol Immunol. 2013;10(1):50-7. Epub 2012/10/16. doi:
10.1038/cmi.2012.43. PubMed PMID: 23064104; PubMed Central PMCID: PMC4003172.
33.   Firth MA, Madera S, Beaulieu AM, Gasteiger G, Castillo EF, Schluns KS, et al. Nfil3-
independent lineage maintenance and antiviral response of natural killer cells. J Exp Med.
2013;210(13):2981-90. doi: 10.1084/jem.20130417. PubMed PMID: 24277151; PubMed Central
PMCID: PMC3865482.
34.   Gasteiger G, Hemmers S, Bos PD, Sun JC, Rudensky AY. IL-2-dependent adaptive
control of NK cell homeostasis. J Exp Med. 2013;210(6):1179-87. doi: 10.1084/jem.20122571.
PubMed PMID: 23650439; PubMed Central PMCID: PMC3674698.
35.   Sun JC, Beilke JN, Lanier LL. Adaptive immune features of natural killer cells. Nature.
2009;457(7229):557-61. doi: 10.1038/nature07665. PubMed PMID: 19136945; PubMed Central
PMCID: PMC2674434.
36.   Sun JC, Lanier LL. Natural killer cells remember: an evolutionary bridge between innate
and adaptive immunity? Eur J Immunol. 2009;39(8):2059-64. doi: 10.1002/eji.200939435.
PubMed PMID: 19637199; PubMed Central PMCID: PMC2819266.
37.   Shimizu K, Sato Y, Shinga J, Watanabe T, Endo T, Asakura M, et al. KLRG+ invariant
natural killer T cells are long-lived effectors. Proc Natl Acad Sci U S A. 2014;111(34):12474-9.
Epub 2014/08/15. doi: 10.1073/pnas.1406240111. PubMed PMID: 25118276; PubMed Central
PMCID: PMC4151776.
38.   Reeves RK, Li H, Jost S, Blass E, Li H, Schafer JL, et al. Antigen-specific NK cell
memory in rhesus macaques. Nat Immunol. 2015;16(9):927-32. doi: 10.1038/ni.3227. PubMed
PMID: 26193080; PubMed Central PMCID: PMC4545390.
39.   Hoft DF, Brown RM, Roodman ST. Bacille Calmette-Guerin vaccination enhances
human gamma delta T cell responsiveness to mycobacteria suggestive of a memory-like
phenotype. J Immunol. 1998;161(2):1045-54. Epub 1998/07/22. PubMed PMID: 9670986.
40.   Montamat-Sicotte DJ, Millington KA, Willcox CR, Hingley-Wilson S, Hackforth S, Innes J,
et al. A mycolic acid-specific CD1-restricted T cell population contributes to acute and memory
immune responses in human tuberculosis infection. J Clin Invest. 2011;121(6):2493-503. Epub

2011/05/18. doi: 10.1172/JCI46216. PubMed PMID: 21576820; PubMed Central PMCID:
PMC3104771.
41.   Shen Y, Zhou D, Qiu L, Lai X, Simon M, Shen L, et al. Adaptive immune response of
Vgamma2Vdelta2+ T cells during mycobacterial infections. Science. 2002;295(5563):2255-8.
Epub 2002/03/23. doi: 10.1126/science.1068819. PubMed PMID: 11910108; PubMed Central
PMCID: PMC2872146.
42.   Van Rhijn I, Kasmar A, de Jong A, Gras S, Bhati M, Doorenspleet ME, et al. A
conserved human T cell population targets mycobacterial antigens presented by CD1b. Nat
Immunol. 2013;14(7):706-13. Epub 2013/06/04. doi: 10.1038/ni.2630. PubMed PMID:
23727893; PubMed Central PMCID: PMC3723453.
43.   Netea MG, Dominguez-Andres J, Barreiro LB, Chavakis T, Divangahi M, Fuchs E, et al.
Defining trained immunity and its role in health and disease. Nat Rev Immunol. 2020;20(6):375-
88. Epub 2020/03/07. doi: 10.1038/s41577-020-0285-6. PubMed PMID: 32132681; PubMed
Central PMCID: PMCPMC7186935.
44.   Netea MG, Joosten LA. Master and commander: epigenetic regulation of macrophages.
Cell Res. 2016;26(2):145-6. Epub 2016/01/16. doi: 10.1038/cr.2016.5. PubMed PMID:
26768770; PubMed Central PMCID: PMCPMC4746612.
45.   James CA, Seshadri C. T Cell Responses to Mycobacterial Glycolipids: On the
Spectrum of "Innateness". Front Immunol. 2020;11:170. Epub 2020/03/03. doi:
10.3389/fimmu.2020.00170. PubMed PMID: 32117300; PubMed Central PMCID:
PMCPMC7026021.
46.   Martin E, Treiner E, Duban L, Guerri L, Laude H, Toly C, et al. Stepwise development of
MAIT cells in mouse and human. PLoS Biol. 2009;7(3):e54. doi: 10.1371/journal.pbio.1000054.
PubMed PMID: 19278296; PubMed Central PMCID: PMCPMC2653554.
47.   Li J, Yu S, Zhang Y, Shen J, Lao S, Li B, et al. Tissue-infiltrating mucosal-associated
invariant T cells play an important role against Mycobacterium tuberculosis infection in
tuberculosis pleurisy. FASEB J. 2016. doi: 10.1096/fj.201600441R. PubMed PMID: 27507648.
48.   Le Bourhis L, Martin E, Peguillet I, Guihot A, Froux N, Core M, et al. Antimicrobial
activity of mucosal-associated invariant T cells. Nat Immunol. 2010;11(8):701-8. doi:
10.1038/ni.1890. PubMed PMID: 20581831.
49.   Suliman S, Murphy M, Musvosvi M, Gela A, Meermeier EW, Geldenhuys H, et al. MR1-
Independent Activation of Human Mucosal-Associated Invariant T Cells by Mycobacteria. J
Immunol. 2019;203(11):2917-27. Epub 2019/10/16. doi: 10.4049/jimmunol.1900674. PubMed
PMID: 31611259; PubMed Central PMCID: PMCPMC6859375.
50.   Godfrey DI, Koay HF, McCluskey J, Gherardin NA. The biology and functional
importance of MAIT cells. Nat Immunol. 2019;20(9):1110-28. Epub 2019/08/14. doi:
10.1038/s41590-019-0444-8. PubMed PMID: 31406380.
51.   Greene JM, Dash P, Roy S, McMurtrey C, Awad W, Reed JS, et al. MR1-restricted
mucosal-associated invariant T (MAIT) cells respond to mycobacterial vaccination and infection
in nonhuman primates. Mucosal Immunol. 2017;10(3):802-13. Epub 2016/10/21. doi:
10.1038/mi.2016.91. PubMed PMID: 27759023; PubMed Central PMCID: PMCPMC5397382.
52.   Gupte A, Padmapriyadarsini C, Mave V, Kadam D, Suryavanshi N, Shivakumar SV, et
al. Cohort for Tuberculosis Research by the Indo-US Medical Partnership (CTRIUMPH):
protocol for a multicentric prospective observational study. BMJ Open. 2016;6(2):e010542.
Epub 2016/02/27. doi: 10.1136/bmjopen-2015-010542. PubMed PMID: 26916698; PubMed
Central PMCID: PMCPMC4769396.
53.   Chandrasekaran P, Mave V, Thiruvengadam K, Gupte N, Shivakumar S, Hanna LE, et
al. Tuberculin skin test and QuantiFERON-Gold In Tube assay for diagnosis of latent TB
infection among household contacts of pulmonary TB patients in high TB burden setting. PLoS
One. 2018;13(8):e0199360. Epub 2018/08/02. doi: 10.1371/journal.pone.0199360. PubMed
PMID: 30067752; PubMed Central PMCID: PMCPMC6070176.

54.   Saunders MJ, Wingfield T, Datta S, Montoya R, Ramos E, Baldwin MR, et al. A
household-level score to predict the risk of tuberculosis among contacts of patients with
tuberculosis: a derivation and external validation prospective cohort study. Lancet Infect Dis.
2020;20(1):110-22. Epub 2019/11/05. doi: 10.1016/S1473-3099(19)30423-2. PubMed PMID:
31678031; PubMed Central PMCID: PMCPMC6928575.
55.   Gela A, Murphy M, Rodo M, Hadley K, Hanekom WA, Boom WH, et al. Effects of BCG
vaccination on donor unrestricted T cells in two prospective cohort studies. EBioMedicine.
2022;76:103839. Epub 20220208. doi: 10.1016/j.ebiom.2022.103839. PubMed PMID:
35149285; PubMed Central PMCID: PMCPMC8842032.
56.   James CA, Xu Y, Aguilar MS, Jing L, Layton ED, Gilleron M, et al. CD4 and CD8 co-
receptors modulate functional avidity of CD1b-restricted T cells. Nat Commun. 2022;13(1):78.
Epub 20220110. doi: 10.1038/s41467-021-27764-w. PubMed PMID: 35013257; PubMed
Central PMCID: PMCPMC8748927.
57.   Francisco LM, Sage PT, Sharpe AH. The PD-1 pathway in tolerance and autoimmunity.
Immunological Reviews. 2010;236(1):219-42. doi: https://doi.org/10.1111/j.1600-
065X.2010.00923.x.
58.   Fletcher HA, Snowden MA, Landry B, Rida W, Satti I, Harris SA, et al. T-cell activation is
an immune correlate of risk in BCG vaccinated infants. Nat Commun. 2016;7:11290. Epub
2016/04/14. doi: 10.1038/ncomms11290. PubMed PMID: 27068708; PubMed Central PMCID:
PMCPMC4832066.
59.   Sakai S, Kauffman KD, Sallin MA, Sharpe AH, Young HA, Ganusov VV, et al. CD4 T
Cell-Derived IFN-gamma Plays a Minimal Role in Control of Pulmonary Mycobacterium
tuberculosis Infection and Must Be Actively Repressed by PD-1 to Prevent Lethal Disease.
PLoS Pathog. 2016;12(5):e1005667. Epub 2016/06/01. doi: 10.1371/journal.ppat.1005667.
PubMed PMID: 27244558; PubMed Central PMCID: PMCPMC4887085.
60.   Wang F, Hou H, Xu L, Jane M, Peng J, Lu Y, et al. Mycobacterium tuberculosis-specific
TNF-alpha is a potential biomarker for the rapid diagnosis of active tuberculosis disease in
Chinese population. PLoS One. 2013;8(11):e79431. Epub 2013/11/19. doi:
10.1371/journal.pone.0079431. PubMed PMID: 24244502; PubMed Central PMCID:
PMCPMC3823617.
61.   Barber DL, Sakai S, Kudchadkar RR, Fling SP, Day TA, Vergara JA, et al. Tuberculosis
following PD-1 blockade for cancer immunotherapy. Sci Transl Med. 2019;11(475). Epub
2019/01/18. doi: 10.1126/scitranslmed.aat2702. PubMed PMID: 30651320.
62.   Lazar-Molnar E, Chen B, Sweeney KA, Wang EJ, Liu W, Lin J, et al. Programmed
death-1 (PD-1)-deficient mice are extraordinarily sensitive to tuberculosis. Proc Natl Acad Sci U
S A. 2010;107(30):13402-7. Epub 2010/07/14. doi: 10.1073/pnas.1007394107. PubMed PMID:
20624978; PubMed Central PMCID: PMCPMC2922129.
63.   Fujita K, Yamamoto Y, Kanai O, Okamura M, Hashimoto M, Nakatani K, et al. Incidence
of Active Tuberculosis in Lung Cancer Patients Receiving Immune Checkpoint Inhibitors. Open
Forum Infect Dis. 2020;7(5):ofaa126. Epub 2020/05/12. doi: 10.1093/ofid/ofaa126. PubMed
PMID: 32391404; PubMed Central PMCID: PMCPMC7200083.
64.   Lewinsohn DA, Lewinsohn DM, Scriba TJ. Polyfunctional CD4(+) T Cells As Targets for
Tuberculosis Vaccination. Front Immunol. 2017;8:1262. Epub 2017/10/21. doi:
10.3389/fimmu.2017.01262. PubMed PMID: 29051764; PubMed Central PMCID:
PMCPMC5633696.
65.   Barry CE, 3rd, Boshoff HI, Dartois V, Dick T, Ehrt S, Flynn J, et al. The spectrum of
latent tuberculosis: rethinking the biology and intervention strategies. Nat Rev Microbiol.
2009;7(12):845-55. Epub 2009/10/27. doi: 10.1038/nrmicro2236. PubMed PMID: 19855401;
PubMed Central PMCID: PMCPMC4144869.

66.   Walzl G, Ronacher K, Hanekom W, Scriba TJ, Zumla A. Immunological biomarkers of
tuberculosis. Nat Rev Immunol. 2011;11(5):343-54. Epub 2011/04/09. doi: 10.1038/nri2960.
PubMed PMID: 21475309.
67.   Achkar JM, Jenny-Avital ER. Incipient and subclinical tuberculosis: defining early
disease states in the context of host immune response. J Infect Dis. 2011;204 Suppl 4:S1179-
86. Epub 2011/10/26. doi: 10.1093/infdis/jir451. PubMed PMID: 21996700; PubMed Central
PMCID: PMCPMC3192549.
68.   Cardona PJ. Revisiting the natural history of tuberculosis. The inclusion of constant
reinfection, host tolerance, and damage-response frameworks leads to a better understanding
of latent infection and its evolution towards active disease. Arch Immunol Ther Exp (Warsz).
2010;58(1):7-14. Epub 2010/01/06. doi: 10.1007/s00005-009-0062-5. PubMed PMID:
20049645.
69.   Lu LL, Smith MT, Yu KKQ, Luedemann C, Suscovich TJ, Grace PS, et al. IFN-γ-
independent immune markers of Mycobacterium tuberculosis exposure. Nature Medicine.
2019;25(6):977-87. doi: 10.1038/s41591-019-0441-3.
70.   Roederer M, Nozzi JL, Nason MC. SPICE: exploration and analysis of post-cytometric
complex multivariate datasets. Cytometry A. 2011;79(2):167-74. Epub 2011/01/26. doi:
10.1002/cyto.a.21015. PubMed PMID: 21265010; PubMed Central PMCID: PMCPMC3072288.
71.   McInnes LH, J. UMAP: uniform manifold approximation and projection for dimension
reduction. https://arxivorg/abs/180203426. 2018.
72.   Cribari-Neto F, Zeileis A. Beta regression in R. 2009.
73.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society Series B
(Methodological). 1995;57(1):289-300.
74.   Layton ED, Yu KKQ, Smith MT, Scriba TJ, De Rosa SC, Seshadri C. Validation of a
CD1b tetramer assay for studies of human mycobacterial infection or vaccination. J Immunol
Methods. 2018. Epub 2018/04/24. doi: 10.1016/j.jim.2018.04.004. PubMed PMID: 29684428.
75.   Corbett AJ, Eckle SB, Birkinshaw RW, Liu L, Patel O, Mahony J, et al. T-cell activation
by transitory neo-antigens derived from distinct microbial pathways. Nature.
2014;509(7500):361-5. Epub 2014/04/04. doi: 10.1038/nature13160. PubMed PMID: 24695216.
758
759

Supporting Information

Table S1. Flow cytometry panels

Figure S1. Gating strategy. **Top**: T cells including NK cells. **Bottom**: APC.

Figure S2. Polyfunctionality of the immune responses to Mtb in TB-resisters (LTBIneg) compared with LTBI-participants (LTBIpos). Data were derived from 13 TB-resisters and 11 LTBI+ participants. Pie charts 1-4 show proportions of cells expressing no markers (dark blue) or ≥1 marker (other colors) in response to Mtb stimulation or control. Arches show the markers expressed by the responding cells. The permutation tables present p values for comparisons across TB-resisters unstimulated (1), TB-resisters Mtb-stimulated (2), LTBI+ participants unstimulated (3) and LTBI+ participants Mtb-stimulated (4) conditions. Tables show unadjusted p-values for each comparison. Pink boxes indicate significant differences defined by FDR p < 0.05.

Fig S3. Cluster analysis of adaptive and innate immune responses in TB-resisters and LTBI-participants. Data were derived from 13 TB-resisters and 9 LTBI-participants for the T cell clustering and 12 TB-resisters and 8 LTBI-participants for the APC clustering. **A**: Forest trees of lnOR and 95% CI, means, and FDR-corrected p values of clusters significantly different between the two groups. **B**: Heat map of T-cell clusters identified by Phenograph. **C**: Heat map of APC clusters.

Figure S4. **Comparison of immune cell subset frequencies in BCG recipients and TB-Resisters.** Data were derived from 13 TB-Resisters and 14 BCG recipients. Forest plot displays ln-transformed odds ratios (lnOR) and 95% confidence intervals (CI). The dotted line shows no

average effect (lnOR = 0, corresponding to OR = 1). Features on the right side of this line are

more highly expressed among TB-Resisters (OR > 1), and features on the left side are more

highly expressed among BCG recipients (OR < 1). The table shows the absolute OR and the

means of each parameter in TB-resisters and LTBI-participants. **Top**: unstimulated PBMC;

Middle: Mtb-stimulated; **Bottom**: Mtb-memory

Figure S5. **Comparison of immune cell subset frequencies in BCG recipients and LTBI-**

articipants. Data were derived from 11 LTBI-participants and 14 BCG-recipients. Forest plot

displays ln-transformed odds ratios (lnOR) and 95% confidence intervals (CI). The dotted line

shows no average effect (lnOR = 0, corresponding to OR = 1). Features on the right side of this

line are more highly expressed among LTBI-participants (OR > 1), and features on the left side

are more highly expressed among BCG recipients (OR < 1). The table shows the absolute OR

and the means of each parameter in TB-resisters and LTBI-participants. **Top**: unstimulated

PBMC; **Middle**: Mtb-stimulated; **Bottom**: Mtb-memory.

Figure S6. Polyfunctionality of select T cell responses in BCG-recipients by comparison

with TB-resisters (LTBIneg) and LTBI-participants (LTBIpos). Data were derived from 14

BCG-recipients, 13 TB-resisters, and 11 LTBI-participants. **A:** BCG-recipients and TB-resisters;

B: BCG-recipients and LTBI-participants. Pie charts 1-4 show proportions of cells expressing no

markers (dark blue) or ≥1 marker (other colors) in response to Mtb stimulation or control. Arches

show the markers expressed by the responding cells. The permutation tables present p values

for comparisons across all groups and conditions. Tables show unadjusted p-values for each

comparison. Pink boxes indicate significant differences defined by FDR p < 0.05.

Figure 1. Functional phenotypes differentially expressed in unstimulated PBMC from TB-resisters and LTBI-participants. Data were derived from 13 TB-resisters and 11 LTBI-participants. PBMC were analyzed after overnight rest in culture medium. **A**. Forest plot displays ln-transformed odds ratios (lnOR) and 95% confidence intervals (CI). The dotted line shows no average effect (lnOR = 0, corresponding to OR = 1). Features on the right side of this line are more highly expressed among LTBI-participants (OR > 1), and features on the left side are more highly expressed among TB-resisters (OR < 1). The table shows the absolute OR and the means of each parameter in TB-resisters and LTBI-participants. **B.** Typical scatter plots exemplifying differences in GranzB+ CD4+ and CD8+ Tconv between TB-resisters (blue dots) and LTBI-participants (red dots). **C**. Dot plots showing frequencies of GranzB+ CD4+ and CD8+ Tconv in all participants.

Figure 2. Functional phenotypes differentially expressed in *Mtb*-stimulated PBMC from TB-resisters and LTBI-participants. Data were derived from 13 TB-resisters and 9 LTBI-participants that contributed PBMC to the T cell analysis. PBMC were analyzed after pre-optimized overnight stimulation with *Mtb* cell membrane (see methods). **A**. Forest plot displays natural log (ln)-transformed odds ratios (lnOR) and 95% confidence intervals (CI). The dotted line shows no average effect (lnOR = 0, corresponding to OR = 1). Features on the right side of this line are more highly expressed among LTBI-participants (OR > 1), and features on the left side are more highly expressed among TB-resisters (OR < 1). The table shows the absolute OR and the means of each parameter in TB-resisters and LTBI-participants. **B.** Typical scatter plots exemplifying differences in PD1+ CD4+ and CD8+ Tconv between TB-resisters and LTBI-participants. **C**. Frequencies of GranzB+ CD4+ and CD8+ Tconv in all participants, including medians, upper and lower quartiles marked on the graph.

843
844
Figure 3. *Mtb*-**specific memory responses in TB-resisters and LTBI-participants.** Data were

derived from 13 TB-resisters and 9 LTBI-participants that contributed PBMC to the T cell analysis.

Mtb-memory was assessed by subtracting the frequency of activated cell subsets inn

unstimulated PBMC from *Mtb*-stimulated PBMC. **A**. Forest plot displays ln-transformed odds

ratios (lnOR) and 95% confidence intervals (CI) of the differences mentioned above. The dotted

line shows no average effect (lnOR = 0, corresponding to OR = 1). Features on the right side of

this line are more highly expressed among LTBI-participants (OR > 1), and features on the left

side are more highly expressed among TB-resisters (OR < 1). The table shows the absolute OR

and the means of each parameter in TB-resisters and LTBI-participants. **B.** Typical scatter plots

exemplifying differences in *Mtb*-memory CD8+IFNγ+ iNKT and between TB-resisters and LTBI-

participants. **C**. Frequencies of CD8+IFNγ+ iNKT in all participants, including medians, upper and

lower quartiles marked on the graph.

Figure 4. Polyfunctionality of immune responses in TB-resisters and LTBI-participants.

Data were derived from 13 TB-resisters and 11 LTBI-participants. Pie charts 1-4 show proportions

of cells expressing no markers (dark blue) or ≥1 marker (other colors) in response to *Mtb*

stimulation or unstimulated control in the T cell panel (**A**) and APC panel (**B**). Arches show the

markers expressed by the responding cells. The permutation tables present p-values for

comparisons across TB-resisters unstimulated (1), TB-resisters *Mtb*-stimulated (2), LTBI-

participants unstimulated (3) and LTBI-participants *Mtb*-stimulated (4) conditions. Tables show

unadjusted p-values for each comparison. Pink boxes indicate significant differences defined by

FDR p < 0.05.

870
871
Figure 5. Comparison of CD8+GMM+GranzB+ Tconv subset in TB-resisters and LTBI-

participants. Data were derived from 13 TB-resisters and 9 LTBI-participants that contributed

2,948,000 events down sampled to 67,000 events in the cleanup gate. **A**: T-cell UMAPs showing

concatenated stimulated and unstimulated cells (upper left corner), distribution of major cell

subsets (upper right corner); Distribution of unstimulated T cells in Resisters and LTBI-participants

(middle row); CD8+GMM+GranzB+ cluster in Resister and LTBI-participants (bottom row).

Representation of typical scatter plot. **B**: Individual results from each participant with medians,

upper and lower quartiles are identified on the graph. **C:** Typical representation of the frequency

of CD8+GMM+GranzB+ T cells in LTBI-participants and Resisters.

Figure 6. Polyfunctionality of immune responses in BCG-recipients by comparison with TB-resisters and LTBI-participants. Data were derived from 14 BCG-recipients, 13 TB-resisters, and 11 LTBI-participants. Pie charts 1-4 show proportions of cells expressing no markers (dark blue) or ≥1 marker (other colors) in response to *Mtb* stimulation or unstimulated control. Arches show the markers expressed by the responding cells. The permutation tables show unadjusted p-values for each comparison. Pink boxes indicate significant differences defined by FDR $p < 0.05$). **A:** T cell panel in BCG-recipients and TB-resisters; **B**: APC panel in BCG-recipients and TB-resisters; C: T cell panel in BCG-recipients and LTBI-participants; **D**: APC

891   panel in BCG-recipients and LTBI-participants. The T cell panels show selected subsets with

892   significant differences of BCG-recipients with TB-resisters and/or LTBI-participants defined by

893   FDR p<0.05. Remaining subsets are shown in **Fig S6**.