1

Lymphatic dysfunction in lupus contributes to cutaneous photosensitivity and lymph node B cell responses

Mir J. Howlader[1,2,*], William G. Ambler[1,3,†,*], Madhavi Latha S. Chalasani[1,4,‡], Aahna Rathod [1], Ethan S. Seltzer[1,¶] Ji Hyun Sim[1,4], Jinyeon Shin[5], Noa Schwartz[1,6,§], William D Shipman III[1,7,8,§§], Dragos C. Dasoveanu[1,9,¶¶], Camila B. Carballo[10] , Ecem Sevim[6,†], Salma Siddique[2,6, ‡‡], Yurii Chinenov[11], Scott A. Rodeo[10,12], Doruk Erkan[6], Raghu P. Kataru[5], Babak Mehrara[5], Theresa T. Lu[1,2,4,6]

[1]Autoimmunity and Inflammation Program, Hospital for Special Surgery Research Institute; New York, NY, USA
[2]Biochemistry, Structural Biology, Cell Biology, Developmental Biology and Molecular Biology Graduate Program, Weill Cornell Medicine, New York, NY, USA
[3]Pediatric Rheumatology, Department of Medicine, Hospital for Special Surgery; New York, NY, USA
[4]Department of Microbiology and Immunology, Weill Cornell Medicine; New York, NY, USA
[5]Division of Plastic and Reconstructive Surgery, Department of Surgery, Memorial Sloan Kettering Cancer Center; New York, NY, USA
[6]Rheumatology, Department of Medicine, Hospital for Special Surgery; New York, USA
[7]Weill Cornell/Rockefeller/Sloan-Kettering Tri-Institutional MD-PhD Program, New York, NY USA
[8]Immunology and Microbial Pathogenesis Graduate Program, Weill Cornell Medicine, New York, NY, USA
[9]Physiology, Biophysics, and Systems Biology Graduate Program, Weill Cornell Medicine, New York, NY USA
[10]Orthopedic Soft Tissue Research Program, Hospital for Special Surgery Research Institute; New York, NY, USA
[11]David Z. Rosenweig Genomics Research Center, Hospital for Special Surgery Research Institute; New York, NY, USA
[12]Department of Orthopedics, Hospital for Special Surgery; New York, NY, USA
[†]Current address: Systemic Autoimmunity Branch, NIAMS/NIH; Bethesda, Maryland, USA
[‡]Current address: NJ Bio; Princeton, NJ, USA
[¶] Current address: Graduate Program in Bioscience, Rockefeller University; New York, NY, USA
[§]Current address: Department of Medicine (Rheumatology), Albert Einstein College of Medicine/Montefiore Medical Center; Bronx, NY, USA
[§§]Current address: Department of Dermatology, Yale University; New Haven, CT, USA
[¶¶]Current address: Department of Cancer Immunology, Genentech; South San Francisco, CA, USA
[‡‡]Current address: Nemours Hospital for Children, Wilmington, DE, USA

*Denotes shared first authorship

Corresponding author:

Theresa T. Lu
Hospital for Special Surgery
535 East 70th Street
New York, NY 10021
Tel: 212-774-2532
Email: lut@hss.edu
52

53   **Conflict of Interests statement:** The authors have declared that no conflict of interest

54   exists.

Abstract

Patients with systemic lupus erythematosus (SLE) are photosensitive, developing skin inflammation with even ambient ultraviolet radiation (UVR), and this cutaneous photosensitivity can be associated with UVR-induced flares of systemic disease, which can involve increased autoantibodies and further end organ injury. Mechanistic insight into the link between the skin responses and autoimmunity is limited. Signals from skin are transmitted directly to the immune system via lymphatic vessels, and here we show evidence for potentiation of UVR-induced lymphatic flow dysfunction in SLE patients and murine models. Improving lymphatic flow by manual lymphatic drainage (MLD) or with a transgenic model with increased lymphatic vessels reduces both cutaneous inflammation and lymph node B and T cell responses, and long term MLD reduces splenomegaly and titers of a number of autoantibodies. Mechanistically, improved flow restrains B cell responsesS in part by stimulating a lymph node fibroblastic reticular cell-monocyte axis. Our results point to lymphatic modulation of lymph node stromal function as a link between photosensitive skin responses and autoimmunity and as a therapeutic target in lupus, provide insight into mechanisms by which the skin state regulates draining lymph node function, and suggest the possibility of MLD as an accessible and cost-effective adjunct to add to ongoing medical therapies for lupus and related diseases.

Introduction

Photosensitivity, a cutaneous sensitivity to ultraviolet radiation (UVR), affects the majority of systemic lupus erythematosus (SLE) patients, but, in addition to inflammatory skin lesions, UVR exposure can trigger systemic disease flares in both patients and murine models that can include increased autoantibody levels and further end organ injury (1-5). Currently, medications used for photosensitive skin responses include topical steroids, topical calcineurin inhibitors, and antimalarials such as hydroxychloroquine. More importantly, lupus patients are advised to reduce UVR exposure by avoiding the sun, wearing protective clothing, and wearing sunscreen to prevent photosensitive skin responses and their sequelae (6-8). Photosensitive skin inflammation, the accompanying risk of systemic disease flares, and the lifestyle modifications needed to prevent these all contribute to disrupting patients' quality of life (9-11). Advances have begun to delineate the mechanisms that contribute to photosensitivity (12-14). However, the link between UVR-induced skin inflammation and increased autoantibody titers in these photosensitive patients remains poorly understood.

Skin communicates with lymphoid tissues via lymphatic vessels which transport cells and interstitial fluid to skin-draining lymph nodes where immune responses occur and can be regulated. This lymphatic transport serves both to clear fluid and inflammation from the skin and to deliver antigens, antigen-presenting cells, and mediators that impact lymph node function (15, 16). Reduced lymphatic flow results in exacerbated skin inflammation and, over time, can lead to the development of autoantibodies (17, 18). This latter scenario is not well understood mechanistically but the combination of skin inflammation and autoimmunity is reminiscent of SLE, and raises

the possibility of lymphatic dysfunction in SLE, which has been described anecdotally

(19-22).

Lymphatic flow brings both dendritic cells and lymph fluid to the subcapsular sinus

of the draining lymph node, at which point they take divergent paths. Dendritic cells

migrating from the skin actively leave the sinus to enter the nodal parenchyma where T

and B cells are located (23). Lymph fluid containing soluble molecules, on the other

hand, flows into the conduit system (24). The conduits consist of a central collagen core

ensheathed by fibroblastic reticular cells (FRCs), and the potential space between the

collagen core and the FRCs allows for flow of lymph fluid throughout the lymph node

until the lymph leaves the node mainly via efferent lymphatic flow. Comprising the wall of

the conduit system, then, FRCs are among the major first sensors of signals flowing

from the skin. FRCs, in turn, play critical roles regulating T and B cell responses (25-

27), and we recently showed that immunization of healthy mice upregulated FRC-

derived CCL2 which promoted CCR2+ Ly6C+ monocyte expression of reactive oxygen

species (ROS) to limit plasmablast survival (28). Lymphatic input from skin to lymph

nodes that alters FRC phenotype, then, could potentially impact lymph node B cell

responses.

In this study, we examined for lymphatic dysfunction in both human SLE and

murine SLE models and sought to understand the consequences on both skin and

lymph node function. We show evidence for potentiation of UVR-induced flow

dysfunction in human SLE and in multiple photosensitive SLE models. Improving

lymphatic flow by manual lymphatic drainage (MLD) or with a transgenic model with

increased lymphatic flow reduced both UVR-induced skin inflammation, draining lymph

node B and T cell responses, and MLD over a prolonged period of time reduced

splenomegaly and titers of a number of autoantibodies. We further show that improving

lymphatic flow upregulates FRC CCL2, and that depleting monocytes limits the flow-

induced reduction in plasmablasts. These results suggest that SLE skin is primed for

UVR-induced lymphatic flow dysfunction, and the dysfunction contributes to both

cutaneous photosensitive responses and, by modulating a FRC-monocyte axis, lymph

node B cell activity in disease. This scenario suggests that improving lymphatic flow and

its consequences on the lymph node stromal microenvironment may be therapeutically

useful in SLE.

Results

Evidence of dysfunctional dermal lymph flow in SLE patients and mouse models

We assessed for evidence of lymphatic flow alterations in the skin of SLE patients and SLE mouse models. In human skin, greater lumenal area of lymphatic vessels can be reflective of distention from reduced lymphatic flow (29-31). We examined biopsies that we had previously obtained from the sun-exposed forearm skin of patients with SLE and with persistently positive antiphospholipid antibodies (APL), a condition that can overlap with SLE. These biopsies were taken from sites of livedo reticularis, a lacy pattern of prominent veins on otherwise normal-appearing skin that is not considered to be related to photosensitivity and that affects both SLE and APL patients (32). We compared the samples from SLE patients (some of whom also had APL) to that of APL patients without SLE. SLE skin showed dilated lymphatic vessels when compared to APL-only skin **(Figure 1A-B**). There was no change in the density of lymphatic vessels **(Figure 1C).** While these results are in the context of livedo skin, the differences between SLE and non-SLE patients suggested that there is dysfunctional lymphatic flow in human SLE skin that is exposed to UVR.

We examined lymphatic flow in lupus model mice by assaying for retention in the skin of intradermally injected Evans blue dye (33) after UVR treatment. We focused on ear skin as it has little fur, allowing the skin to be directly exposed to UVR. We used two photosensitive lupus models, the MRL-Fas^{lpr} (LPR) spontaneous lupus model (34-36) and the imiquimod (IMQ) lupus model that is induced by chronic epicutaneous IMQ application (37). The LPR mice and their MLR-MpJ (MRL) controls were treated with 1000J UVB/m2/day of UVR for 4 days prior to Evans blue dye injection in the ear skin (**Figure 1D**). LPR mice showed increased Evans blue dye retention in the ear when compared to MRL mice **(Figure 1E)**, suggesting reduced flow out of the ear skin. As

expected for the photosensitive LPR mice, there was greater UVR-induced skin swelling

in LPR mice when compared to control MRL mice **(Figure 1F).** The greater Evans blue

dye retention in LPR mice was specific to UVR exposure, as LPR mice showed no

increases in Evans blue dye retention or ear thickness at baseline without UVR (**Figure**

1G-H).

The UVR-induced lymphatic alterations and ear swelling were accompanied by the

accumulation of inflammatory cells. We had previously shown that inflammatory Ly6C^{hi}

monocytes accumulated after one day of UVR in LPR mice (35). After 4 days of UVR,

monocytes numbers accumulated in greater numbers in both MRL and LPR mice treated

with UVR at comparable levels **(Figure 1I)**. However, TCR$\alpha\beta$+ CD4 T cells and

TCR$\alpha\beta$+CD3+CD4-CD8- "double negative" (DN) T cells that are characteristic of LPR

mice (38) both accumulated in greater numbers in LPR mice compared to MRL mice at

this time point **(Figure 1J-K)**. Neutrophil, CD8 T cells, and TCR$\gamma\delta$ T cells showed no

changes between LPR and MRL mice **(Supplemental Figure 1A-C)**. Together with the

Evans blue dye experiments, these results indicated that lymphatic flow dysfunction

accompanies the increased inflammation in the skin (as indicated by the greater edema

and T cell infiltrate) in UVR-treated LPR mice. These results suggested the possibility

that lymphatic flow dysfunction, by failing to remove fluid and inflammatory mediators

from the skin, was contributing to the increased skin inflammation of the lupus model

mice.

We examined the IMQ model (37), inducing this model in B6 mice by applying

IMQ, a TLR7 agonist, to the skin for 4-5 weeks, yielding B6-IMQ mice **(Figure 1L)**. IMQ

was applied on the right ear only, and the skin on the rest of the body, including the left

ear, was considered "non-lesional" skin and reflective of systemic disease. We have

shown previously that the "non-lesional" left ear in IMQ mice is similar to non-lesional

skin in human lupus in expressing an interferon signature whereas the right ear has a

less robust interferon signature (39). Additionally, the right ear, even without UVR

exposure, showed upregulation of apoptotic pathways (**Supplemental Figure 2A**),

suggesting that there was tissue damage from repeated local treatment of IMQ. To

assess effects of UVR exposure without confounding results from direct IMQ treatment,

we focused on the left, non-lesional ear. The left ear showed increased Evans blue dye

retention with increased skin swelling upon UVR exposure when compared to vehicle-

painted controls **(Figure 1M-N)**. Similarly, the footpad of B6-IMQ mice showed increased

Evans blue dye retention when compared to B6 mice (**Figure 1O**), suggesting that the

dermal lymphatic dysfunction in B6-IMQ mice affected non-lesional skin throughout the

body and that the left ear results were not reflective of direct exposure to IMQ that had

transferred from the right ear. The right ear also showed increased Evans blue dye

retention and increased ear thickness (**Supplemental Figure 2B-C**).

Characterization of the inflammatory infiltrate in B6-IMQ mice showed a higher

baseline monocyte number in B6-IMQ compared to B6 mice, and UVR exposure for 1

day showed greater monocyte accumulation in B6-IMQ mice (**Figure 1P**). Monocyte

accumulation continued to increase with additional days of UVR exposure, and were

equally high in B6-IMQ and B6 ears after 4 days of UVR (**Figure 1P**). In contrast, CD4 T

cell number accumulation was limited until after 4 days of UVR, when B6-IMQ showed

higher numbers of CD4 T cells than B6 mice (**Figure 1Q**). Neutrophils, TCR$\alpha\beta$ CD8 T

cells, and TCR$\gamma\delta$ T cells showed no changes between IMQ and control mice after 4 days

of UVR exposure (**Supplemental Figure 1D-F**). Similar to LPR mice, then, B6-IMQ mice

showed that UVR-induced skin inflammation after 4 days of treatment included lymphatic

dysfunction along with tissue swelling and T cell accumulation. Together, our findings in

two distinct SLE models suggested that photosensitive skin is characterized in part by

UVR-induced lymphatic dysfunction, supporting the evidence of lymphatic dysfunction in

sun-exposed skin in human SLE.

Improving lymphatic flow reduces photosensitivity of skin in SLE models

We asked about the contribution of reduced lymphatic flow to cutaneous

photosensitive responses in the SLE models by using two different approaches to

improve lymphatic flow. One approach was through manual lymphatic drainage (MLD), a

technique used by physical therapists to reduce swelling in patients with acquired or

congenital lymphedema (40, 41). We administered MLD targeting the left ear once a day

during the course of UVR exposure **(Figure 2A-B).** This treatment decreased Evans

blue retention in LPR mice relative to handling controls **(Figure 2C and Supplemental**

Figure 3A), suggesting that MLD was successful in improving lymphatic flow. MLD also

ameliorated UVR-induced skin inflammation, with reductions in ear swelling **(Figure 2D-**

E), CD4 T cell numbers and interferon gamma (IFNg) expression **(Figure 2F-H)**, and DN

T cell numbers and IFNg expression **(Figure 2I-J)**. The proportion of regulatory T cells

(Treg) and T helper 17 cells (Th17) within the CD4 T cell population, the proportion of IL-

17-expressing DN cells, and the number of CD8 T cells did not change with MLD

(**Supplemental Figure 3B-E**). Monocyte and neutrophil numbers also remained

unchanged (**Supplemental Figure 3F-G).** These results suggested that improving

lymphatic flow with MLD was able to both reduce ear swelling and T cell accumulation

and activity in UVR-treated LPR mice.

MLD had similar effects in B6-IMQ mice **(Figure 2K-P)**. Evans blue dye retention

was reduced **(Figure 2L)**, suggesting improved lymphatic flow, and this was

accompanied by reduced ear swelling **(Figure 2M)**. Both total CD4 T cell numbers and

IFNg-expressing Th1 proportion were also decreased with MLD **(Figure 2N-P)**. Similar

to LPR mice, MLD had no effect on Treg, Th17, CD8 T cell, monocyte, or neutrophil

accumulation **(Supplemental Figure 3H-L)**. Our results together suggested that

improving lymphatic flow with MLD ameliorated UVR-induced cutaneous inflammation in

both the LPR and IMQ SLE models.

Our second approach to improve lymph flow was by using transgenic

Flt4Cre^{ERT2}PTEN$^{fl/fl}$ mice (42). When treated with tamoxifen, these mice have lymphatic

endothelial cell (LEC)-specific deletion of PTEN, an antagonist of critical

VEGFC/VEGRF3 signaling. This results in expansion of functional lymphatic vessels,

improved lymphatic flow from skin, and reduced UVR-induced skin inflammation in

healthy (ie non-lupus) mice (42). We confirmed that tamoxifen treatment of

Flt4Cre^{ERT2}PTEN$^{fl/fl}$ mice specifically deleted PTEN from LECs (**Supplemental Figure**

4A). We designated the tamoxifen-treated mice as "LEC^{PTEN} mice" and non-tamoxifen-

treated mice as "LEC^{WT} mice", and induced the IMQ model in them to generate LEC^{PTEN}-

IMQ and LEC^{WT}-IMQ mice (**Figure 2Q**). Reduced Evans blue dye retention in the left ear

of LEC^{PTEN}-IMQ mice compared to LEC^{WT}-IMQ controls confirmed improved lymphatic

drainage (**Figure 2R**), and this was associated with reduced UVR-induced ear swelling

(Figure 2S), suggesting that improving lymphatic flow reduced the ear swelling. Similar

to MLD, the genetic approach to improving lymphatic flow reduced CD4 T cell numbers

in LEC^{PTEN}-IMQ mice compared to LEC^{WT}-IMQ mice (**Figure 2T-U**). Notably, tamoxifen

treatment of non-transgenic B6 mice did not reduce Evans blue dye retention or ear

swelling **(Figure 2V-W),** suggesting that effects in LEC^{PTEN}-IMQ mice were attributable to

PTEN deletion rather than to tamoxifen treatment. Treg, CD8 T cell, monocyte, and

neutrophil numbers remained unchanged (**Supplemental Figure 3M-P**). The right ear of

LEC^{PTEN}-IMQ mice also showed reduced Evans blue dye retention and ear swelling

when compared to LEC^{WT}-IMQ mice and no changes in monocyte and neutrophil

numbers (**Supplemental Figure 4B-E**). As with the left ear, B6-IMQ mice showed no

reduction in Evans blue dye retention or ear thickness with tamoxifen (**Supplemental**

Figure 4F-G). Consistent with prior reports that improved lymphatic drainage can

reduce cutaneous inflammation (33, 42, 43), our results from using both an acute

physical approach in two lupus models and a long-term genetic approach showed that

improving lymphatic flow from skin reduces cutaneous UVR-induced inflammation in

SLE model mice, suggesting that the lymphatic dysfunction contributes to photosensitive

skin responses in SLE.

Improving lymphatic flow reduces draining lymph node B cell responses in SLE

models*.*

We asked whether lymphatic flow alterations in SLE models contributed to

modulating immune activity in downstream lymph nodes. Improving lymphatic flow with

MLD over 4 days in LPR mice during UVR exposure did not affect overall lymph node

cellularity or overall B cell numbers in draining auricular nodes (**Figure 3A-B**), but did

reduce germinal center B cell and plasmablast numbers (**Figure 3C-D**). CD4 T cell

numbers were also reduced (**Figure 3E-F**), as was the frequency of TFH cells

(**Supplemental Figure 5A**), which could have contributed to reduced germinal center B

cell responses. Th1 (**Figure 3G**), Tregs, and Th17 frequencies were unchanged

(**Supplemental Figure 5B-C**). CD8 (**Figure 3H**), and DN T cell (**Figure 3I**) numbers

were unchanged, as was the expression of IL-17 by DN cells (**Supplemental Figure**

5D). These data suggested that improving lymphatic flow with MLD in LPR mice reduces

draining lymph node germinal center and plasmablast responses along with overall CD4

T cell numbers and the proportion of TFH cells.

Similar to LPR mice, MLD in B6-IMQ mice showed no effects on total LN cellularity

or on overall B cells (**Fig 3J-K)** but did reduce germinal center B cell, plasmablast, and

CD4 T cell numbers (**Fig 3L-O)**. Note that the reduction in germinal center (2.2 fold) and

plasmablast (2.4 fold) numbers reflects a partial reduction rather than normalization to a

non-lupus state, as UVR-treated B6 mice have very few germinal center cells and

plasmablasts (1002+/-859 and 312+/-220 per lymph node, respectively). Unlike in LPR

mice, MLD in B6-IMQ mice did not reduce TFH cells **(Supplemental Figure 5E)** but did

reduce the TH1 frequency **(Figure 3P).** Tregs and Th17 cell frequencies

(Supplemental Figure 5F-G) and CD8 T cell numbers **(Figure 3Q)** were unchanged.

These data together showed that MLD in LPR and IMQ models reduced B cell

responses and CD4 T cell numbers and differentiation, thus suggesting that even a short

period of improved lymphatic flow from skin can reduce B and CD4 T cell responses.

In contrast to LPR and B6-IMQ mice treated with MLD, LEC^PTEN-IMQ mice showed

decreases in lymph node cellularity and overall B cells when compared to LEC^WT-IMQ

mice **(Figure 3R-S).** As with LPR and B6-IMQ mice, germinal center B cell, plasmablast,

and CD4 T cell numbers were reduced (**Figure 3T-W).** Th1 **(Figure 3X)**, TFH, and Treg

frequencies **(Supplemental Figure 5H-I)** and CD8 T cell numbers **(Figure 3Y)** were

unchanged. These data suggest that improving lymph flow genetically and before

induction of lupus disease activity reduced overall lymph node cellularity, and, similar to

short term MLD, preferentially reduced B cell and CD4 responses.

Longer duration of improved lymphatic flow reduces systemic disease activity in

SLE mice

While our MLD was targeted to the left ear and draining left auricular lymph node,

reduced inflammation at the site of greatest UVR exposure (left ear versus the fur-

covered back, for example) and its draining lymph node could potentially have a

systemic effect, as soluble mediators including antigen and inflammatory cytokines can

travel from the skin to draining lymph node and then, via efferent flow from the lymph

node, out to the systemic circulation to the spleen and end organs (12). Also, reduced B

and T cell responses in the draining lymph node would lead to fewer cells and

autoantibodies leaving the lymph node to reach the circulation and spleen and other

tissues to carry out inflammatory, effector, or memory functions. Furthermore, some

inflammatory cells can also leave the skin directly into the blood circulation by reverse

transmigration from tissue into blood vessels, as has been shown for neutrophils in

UVR-treated skin (44).

We thus asked if improving lymphatic flow in the LEC^{PTEN}-IMQ mice or by local

MLD can affect parameters of systemic disease activity. We first looked at the

splenomegaly that characterizes both the IMQ and LPR models (37, 45). Spleens do not

have afferent lymphatics (46) and so splenic changes are considered to reflect systemic

changes. LEC^{PTEN}-IMQ mice had reduced splenic weight compared to LEC^{WT}-IMQ mice

(**Figure 4A**), suggesting reduction in systemic disease activity with improved lymphatic

flow. To determine if MLD could also affect systemic disease activity, we treated LPR

mice with MLD for 4-5 weeks, a duration that would allow indirect splenic changes to

occur and for turnover of pre-MLD autoantibodies (47). MLD-treated LPR mice showed

reduced splenic weight compared to mice that did not receive MLD, suggesting that

longer term MLD can reduce systemic disease (**Figure 4B**). We further measured levels

of serum autoantibodies after 4-5 weeks of UVR with and without MLD. While high titer

anti-DNA and ribonucleoprotein (RNP) antibodies remained unchanged, antibodies

against complement pathway components (C5, C9, Factor B, Factor I, Factor P), histone

2B, and beta2 glycoprotein were reduced (**Fig 4C,D**). Interestingly, LPR mice also

expressed autoantibodies found in dermatomyositis patients (TIF1gamma, Jo1, PL-7,

SAE1/SAE2, MDA5, NXP2, mi-2, PM/Scl100) who also demonstrate photosensitivity

(48), and some of these (Jo-1,NXP2, PM/Scl100) were also reduced with MLD (**Fig**

4C,D). The reduced splenomegaly and autoantibody levels suggest that longer term

MLD can reduce systemic disease activity.

Improving lymphatic flow increases lymph node FRC CCL2, monocyte ROS

generation, and restrains plasmablast responses in a monocyte-dependent

manner

To understand how improving lymphatic flow limits lymph node B cell responses,

we considered that FRCs are among the initial sensors of lymph fluid within lymph nodes

and asked if improving lymphatic flow could drive an FRC-monocyte axis that we have

previously shown to regulate plasmablast accumulation in healthy mice (28). In this axis,

stromal CCL2 in the T zone and the medulla induced local CCR2+ Ly6^{hi} monocytes to

upregulate ROS expression, which then limited survival of plasmablasts that were

colocalized with the CCL2-expressing FRCs and monocytes. Upon MLD of IMQ-treated

CCL2-GFP reporter mice, FRCs in the draining auricular lymph nodes showed

upregulated GFP, suggestive of upregulated CCL2 expression (**Figure 5A**). By anti-

CCL2 staining, an increase in FRC CCL2 was also detectable in LEC^{PTEN}-IMQ mice

compared with control LEC^{WT}-IMQ mice (**Figure 5B**). Additionally, MLD also increased

FRC numbers **(Figure 5C),** which likely reflected the increased proliferation rate, as

indicated by increased Ki-67 levels **(Figure 5D-E).** This expansion of FRC numbers

likely added further to the level of CCL2 sensed by CCR2+ cells in the lymph node.

Together, our results suggested that improving lymphatic flow increased the level of

FRC CCL2 in draining lymph nodes.

In association with the stromal CCL2 upregulation, Ly6C^{hi} monocyte numbers were

unchanged (**Figure 5F**), but monocyte ROS expression was increased (**Figure 5G**). The

monocyte ROS increase was specific to this population, as B cells did not show a similar

increase (**Figure 5G**). This finding of unchanged monocyte numbers but upregulated

ROS expression was consistent with the FRC-monocyte axis in healthy mice (28).

Together, these results were consistent with a model whereby improving lymphatic flow

increased stromal CCL2 which then increased CCR2+ Ly6C^{hi} monocyte expression of

ROS to control B cell responses in draining lymph nodes.

To test this model, we took two approaches to ask the extent to which monocytes

were required for the reduced B cell responses seen with improved lymphatic flow. We

used anti-Gr-1 (28) that depleted both monocytes (**Figure 5H-J**) and neutrophils

(**Supplemental Figure 6A**) in B6-IMQ mice during UVR and MLD. This depletion was

associated with restoration of germinal center B cell and plasmablast numbers to the

higher levels seen in mice without MLD (**Figure 5K-L; compare with Figure 3L-M),**

suggesting that MLD-driven reduction in B cell responses was dependent on myeloid

cells. To further assess the role of monocytes without depleting neutrophils, we induced

the IMQ model in CCR2-DTR mice, injected DT to deplete monocytes (**Figure 5M-O)** but

not neutrophils (**Supplemental Figure 6B),** and then treated with UVR and MLD **(Figure**

5M). Germinal center B cell numbers were not altered but plasmablast numbers were

restored to the higher levels seen in mice without MLD (**Figure 5P-Q; compare with**

Figure 3L-M). The results of the Gr-1 depletion and the CCR2-DTR model together

suggested that improved lymphatic flow limits plasmablast accumulation in a monocyte-

dependent manner, while the reduction in germinal center B cells is mediated by other

mechanisms. All together, our data suggested a model whereby restoring lymphatic flow

in UVR-treated SLE mice reduces draining lymph node B cell responses at least in part

by upregulating stromal CCL2 and increasing monocyte ROS to limit plasmablast

accumulation.

Discussion

Lymphatic flow is critical for clearing inflammatory mediators from peripheral

tissues and communicating with draining lymph nodes, and our results suggested that

the link between UVR-induced photosensitive skin responses and increased

autoimmunity in SLE is at least in part due to a propensity for SLE skin to develop UVR-

induced lymphatic dysfunction. By correcting this dysfunction using either MLD or a

transgenic model with increased lymphatic flow, we showed that this lymphatic

dysfunction contributed to both upstream cutaneous photosensitive responses and

downstream draining lymph node B and T cell responses **(see Figure 6 graphical**

summary). While the contribution of lymphatic dysfunction to increasing skin

inflammation is consistent with findings in non-lupus models (42, 49), we establish here

that lymphatic dysfunction also contributes to B and T cell responses in lymph nodes.

Our results suggest that lymphatic dysfunction prevents optimal function of the

regulatory FRC-monocyte axis in lymph nodes that normally limits plasmablast

responses, and that improving lymphatic function and reducing B cell responses, over

the long term, can reduce autoantibody titers. Reduced antibody titers, in turn, has the

potential to lead to reduced deposition of immune complexes and limiting inflammation

and damage in skin, kidneys, and other end organs. Together, our results suggested

that UVR-induced lymphatic flow dysfunction is a contributing factor to lupus

pathophysiology and points to lymphatic modulation of lymph node stromal function as a

therapeutic target for UVR-induced disease flares.

Our finding that lymphatic flow modulates lymph node stromal phenotype in SLE

models underscores the importance of the communication between tissue and lymph

nodes and highlights FRC regulation as an outcome of this communication. In this

setting, FRCs act as a rheostat that senses and converts peripheral tissue signals into

regulators of lymph node activity. We showed that improving flow was connected

functionally to the FRC CCL2-monocyte ROS axis that we had previously delineated in

the setting of immune responses in non-lupus models. Consistent with the findings in

healthy mice, this axis contributed to limiting plasmablast responses and not germinal

center responses. Anatomically, CCL2 FRCs are positioned with plasmablasts within the

T cell zone and medulla, where they are able to modulate local monocytes, and

consequently the plasmasblasts (28, 50). FRCs, similar to fibroblasts in the synovium

and other tissues (51, 52), are comprised of multiple subsets that have distinct functions

related to their anatomic positioning within lymph nodes (25, 26); it will be interesting to

further understand how different FRC subsets are modulated phenotypically and

functionally by changes in lymphatic flow.

The ways by which lymphatic flow modulates FRC phenotype remains to be

determined. FRCs are excellent mechanosensors and respond to environmental

alterations to modulate immune function (53-55), and our results may reflect FRC

sensing of changes in parameters such as shear stress as lymphatic flow changes. It is

also possible that our results reflect FRC sensing of different soluble mediators

originating in the skin as lymphatic flow and consequent skin inflammation is modulated

(56). For example, with the MLD-induced reduction of T cell interferon gamma

expression in skin, there could be reduced levels of interferon that reach the lymph node

FRCs , thus altering FRC phenotype. The reduced skin inflammation could also include

reduced type I IFN expression in skin, with reduced type I IFN flowing to draining lymph

nodes. Here, it is interesting to note that IFNAR deletion from FRCs has been shown to

upregulate FRC CCL2 expression (57), raising the possibility that reduced type I IFN

coming from the skin could cause FRC CCL2 upregulation and activate the FRC-

monocyte axis to limit B cell responses. Yet another potential way by which improving

lymphatic flow can modulation FRC function is that while dendritic cell mobilization from

skin to lymph nodes is relatively well preserved even in the face of changes in lymphatic

fluid flow (58), dendritic cells can modulate FRC phenotype (59) and skin dendritic cell

alterations induced by the reduced skin inflammation upon improving lymphatic flow may

potentially alter FRC function upon dendritic cell migration to draining nodes. Further

elucidation of how lymphatic manipulation impacts the lymph node stromal compartment

will be an important future direction.

The reduced lymphatic flow and relationship to B cell responses in lupus models

echoes the findings of Swartz and colleagues who showed in otherwise healthy mice

that a dearth of dermal lymphatics leads over time to autoantibodies generation (17).

Our data showing that lymphatic function is connected to a lymph node FRC-monocyte

axis that we have previously shown in healthy mice (28) to limit B cell responses

provides a potential mechanism to explain these findings. This would suggest that the

lymphatic regulation of the FRC-monocyte axis is a physiologic mechanism for immune

regulation, and that the effects of UVR exposure on autoimmunity in lupus is, in part, a

disruption of this physiologic lymphatic-FRC-monocyte axis.

Our study leads to many more questions. It will be interesting to understand the

other mechanisms by which lymphatic flow impacted lymph node immune function. For

example, the lymphatic-modulated FRC-monocyte axis in lupus model mice contributed

to limiting plasma cell responses and not germinal center responses. In LPR mice,

reduced TFH numbers by improved lymphatic flow may have contributed to reduced

germinal center responses; how improving lymphatic flow reduced TFH numbers or the

T cell interferon gamma expression in both LPR and IMQ models in lymph nodes and

the extent to which these changes are the consequence of FRC changes remains to be

examined. It will also be interesting to understand the mechanisms by which lupus leads

to lymphatic dysfunction. Type I IFN has been shown to inhibit dermal lymphatic fluid

transport in the setting of a vaccinia skin infection model (60), suggesting the possibility

that the high IFN-I tone in non-lesional human and murine lupus skin (39, 61-63)

combined with UVR-induced IFN-I upregulation (64) could be contributing to the

lymphatic dysfunction that we observe. The high interferon tone and other features of

lupus skin may also contribute to lymphatic flow changes even without UVR exposure

that was not captured with our Evans Blue dye assay but that may have immune

consequences. Additionally, lymphatic endothelial cells play important roles in

regulating the immune cells that enter and migrate within the lymphatics (65-67); the

ways in which lymphatic endothelial cell phenotype is altered in lupus skin and the

impact on autoimmunity will be interesting to understand going forward.

Our results have clinical implications. Potentially, MLD could be used in addition to

current medical therapies to reduce local cutaneous inflammation or, over a larger area

over time, to contribute to reducing systemic disease. Examination will be needed to

assess the utility of MLD as an accessible and relatively inexpensive adjunct approach

to ameliorate disease in SLE.

Materials and Methods

Study Design

The purpose of this study was to examine lymphatic function in the skin of

photosensitive SLE mouse models and human SLE skin and to understand the

contributions of lymphatic function on cutaneous photosensitive and lymph node

responses. Laboratory mice were used as subjects for experiments. Human skin

sections were also analyzed. Evans blue lymphangiography was used to assess

lymphatic flow and flow cytometry was used to identify and quantify cell numbers. For

experiments, sample sizes of n=3-21 animals per condition were evaluated in 1 to 11

independent experiments.

Human Samples Staining and Imaging

FFPE sections from skin biopsies of 8 patients with SLE and/or persistently positive APL

examined in (32) were used to stain for lymphatic vessels. Patients in the study had

active livedo reticularis and staining for lymphatic vessels was done on biopsies taken

from the purple areas of the livedo. Five patients had SLE, three of which had

concomitant persistent positive APL, and three patients had persistent positive APL

without SLE. Two archived samples from healthy donors were also used. Ethical

approval for this study was obtained from Hospital for Special Surgery Institutional

Review Board (IRB) (IRB number: 2015-256), where participants had signed written

informed consents for both the initial study and for the subsequent study of skin

biopsies.

Five micrometer paraffin sections were deparaffinized in xylene and rehydrated in a

graded alcohol series. After a final wash with distilled water, specimen slides were

placed in boiling Antigen Unmasking Solution, Tris-Based (Vector Laboratories) for 15

minutes. Slides were briefly washed in water and thereafter in phosphate buffered saline

(PBS)-0.025% triton-X-100 prior to blocking non-specific binding sites with PBS-3%

bovine serum albumin (BSA) for 30 minutes at room temperature. Double

immunostaining was performed with 1:40 dilution in PBS/0.5% BSA of anti-PDPN (D2-

40, Biolegend, San Diego, CA) and 1:100 dilution of anti-CD31/PECAM-1 (Novus

Biologicals, Centennial, CO) overnight at 4°C. Slides were washed with PBS-0.025%

triton-X-100 and endogenous peroxidase was inhibited by incubating slides in 3% H_2O_2

for 15 minutes in the dark. The sections were then washed in distilled water followed by

wash buffer and incubated with alkaline phosphatase-conjugated donkey anti-mouse

and horse radish peroxidase-conjugated donkey anti-rabbit secondary antibodies (both

Jackson ImmunoResearch, West Grove, PA) at 1:100 dilution (prepared in tris-buffered

saline (TBS) with 0.5% BSA) for 1 hour at room temperature. The reaction was revealed

with a 3,3′-diaminobenzidine Substrate Kit for peroxidase and fast blue substrate (both

Sigma-Aldrich, Burlington, MA) for alkaline phosphatase. At the end of incubation, slides

were washed in TBS-0.025% triton-X-100, mounted with Clear-Mount (Electron

Microscopy Sciences) and baked at 56°C for 20 minutes. Imaging was performed with

Leica Aperio CS2 slide scanner at 40X magnification and ImageJ (NIH) analysis

software used to measure the lumenal area of PDPN+CD31+ lymphatic vessels. Image

analysis was performed blinded to the patient diagnosis.

Mice

Mice between 6-15 weeks of age were used unless otherwise specified. Both male and

female mice were used for experiments. All experiments were performed with age and

sex matched controls. C57Bl/6, CCL2$^{-/-}$ (68), CCL2-GFP (69), MRL^{MpJ} (MRL) and MRL-

Fas[lpr] (LPR) mice were originally from Jackson Laboratory (JAX, Bar Harbor, Maine) and

bred at our facility. PTEN[fl/fl] CreERT2 (42) were as described. CCR2-DTR (70) mice

were bred at our facility. All animal procedures were performed in accordance with the

regulations of the Institutional Animal Care and Use Committee at Weill Cornell Medicine

(New York, NY).

Manual lymphatic drainage

MLD was adapted to the mouse by a licensed physical therapist (CBC) with experience

in MLD. MLD targeting the left ear was performed daily as specified. ProDerma smooth,

powder-free latex gloves (Uniseal, Alhambra, CA) were used to minimize friction and

cutaneous trauma and all movements were done at a speed of 1-2 seconds per

movement. Mice were anesthetized throughout procedure. The first step was the

clearing step, and this started with passive motion of the left forelimb, where the mouse

was placed in a supine position, held by the paw and clockwise rotations were

performed (20 repetitions) to clear the supra-clavicular and axillary area. Subsequently,

stationary circular movements with light pressure of the index finger were applied on the

left submandibular area followed by the auricular area (20 reps each). The mouse was

then placed in the right lateral decubitus position to proceed with the reabsorption step

which was performed in a specific sequence: 1. The left ear was held gently with a

pincer grip (thumb and index finger) and light sweeping movements with the index finger

were done on the ventral surface of the ear from distal to proximal (50 reps); 2. Placed in

a prone position, the ear was gently held with a pincer grip and sweeping movements

were done from distal to proximal on the dorsum of the base of the ear (where the large

collecting lymphatic vessels are and can be visualized by Evan's blue

lymphangiography) (200 reps); 3. Placed back to the right lateral decubitus position,

sweeping movements at the base of the ear toward the auricular lymph node were

performed (20 reps); 4. Finally placed in supine position again, the same clockwise

rotations of the left forelimb was performed (20 reps). Handling controls were

anesthetized in the same manner as for MLD, and the mouse was placed prone and left

ear was held with a pincer grip similarly to MLD for 5 minutes.

Mouse treatments

For UVR treatments, mice were exposed to 1000 - 1500 J UVB/m^2/day for four

consecutive days using a bank of four FS40T12 sunlamps, as previously described (35).

For long term treatment, mice were exposed to UVR for four consecutive days the first

week and then for 3 consecutive days/week in the following weeks. To measure ear

swelling after UVR exposure, a caliper (Mitutoyo, Industry, CA) was used. Each ear was

measured in the anterior half of the ear three times and the average was taken.

For the IMQ-induced lupus mouse model, mice were painted on the dorsal and ventral

sides of the right ear with 5% imiquimod cream (Taro Pharmaceutical) 3x/week for 4-5

weeks (~50 mg/mouse total cumulative dose) (37).

For mice receiving tamoxifen, tamoxifen (Sigma-Aldrich) in corn oil was injected

intraperitoneally (IP) at a dose of 300mg/kg/day every other day for three doses.

For monocyte depletion studies in the B6-IMQ mice, anti-Gr1 (RB6-8C5) or isotype

control IgG (LTF-2) (both BioXCell, Lebanon, NH) were injected IP at a dose of 250μg in

200μl PBS on indicated days. To deplete monocytes in the CCR2-DTR mice, diphtheria

toxin (Enzo Life Sciences, Farmingdale, NY) was injected IP at a dose of 250ng in 200 ul

PBS.

Evans blue lymphatic function assay

Evans blue retention assay was performed as previously described (71). Mouse ears

were injected intradermally at the tip using a Hamilton syringe (#1701, 10 ul syringe) with

a 30-gauge needle. 1µl of 2% Evans blue (Sigma-Aldrich) was injected and mice were

euthanized 22-24 hours later. The harvested ear was placed in 300µl formamide at 58ºC

overnight to extract Evans blue, which was quantified by absorbance with a Multiskan

Ascent plate reader (Titertek, Pforzheim, Germany) at 620nm using a titration curve.

ROS Staining

Intracellular ROS was measured using 5-(and6)chloromethyl-2',7'-

dichlorodihydrofluorescein diacetate, acetyl ester (CM-H2DCFDA, Thermo Fisher

Scientific), as in (28). The dye was reconstituted at 5 mmol/L in DMSO and stored at

−20°C. Cells were prepared in RPMI and stained with 1/500 dilution of the stock in PBS

for 30 minutes at 37°C prior to flow cytometric analysis.

Flow cytometry staining and quantification

For flow cytometric staining of skin, single cell suspensions were generated as

previously described (72). In brief, ears were finely minced, digested in type II

collagenase (616 U/mL; Worthington Biochemical Corporation, Lakewood, NY), dispase

(2.42 U/mL; Life Technologies, Carlsbad, CA), and DNAse1 (80 µg/mL; Sigma-Aldrich),

incubated at 37°C while shaking at 100 rpm, triturated with glass pipettes, and filtered.

For staining of lymph node cells, hematopoietic cells from lymph nodes were obtained by

mashing the lymph nodes and extruding the cells through a 70µm strainer. Stromal cells

were obtained as previously described (73); lymph nodes were minced, digested with

type II collagenase (616 U/mL) and DNAase1 (40μg/mL) at 37°C while shaking at 50

rpm, triturated with glass pipettes, and filtered.

To count cells, the single cell suspension from the whole lymph node or ear was washed

and resuspended in 300 ul of buffer. Ten ul was taken to be counted on the Multisizer 4e

Coulter Counter (Beckman Coulter) and this count was used to calculate total number of

cells per lymph node. One to two million cells per sample were stained, and most of the

sample was run on the flow cytometer. To obtain absolute number of a particular

population of cells per ear or lymph node, the frequency of these cells (of the total) in the

facs analysis was multiplied by the total number of cells per tissue (as calculated by the

Counter Counter data).

For flow cytometry analysis, gating of specific populations was performed after excluding

debris, doublets, and dead cells using 4',6-diamidino-2-phenylindole dihydrochloride

(DAPI, Invitrogen) for non-fixed cells. Antibodies are from Biolegend unless otherwise

specified. Samples were treated with anti-mouse CD16/32 (Fc block, clone 93) prior to

staining with additional antibodies. Gating strategies and antibodies used-- B cells:

CD45+(30-F11) B220+(RA3-6132); monocytes: CD45+, B220-, CD3-(145-2C11),

CD11b+(M1/70), Ly6C^{hi}(HK1.4), Ly6G-(1A8); neutrophils: CD45+, CD11b+, Ly6C^{med},

Ly6G+; germinal center B cells: CD3-, B220+, GL7+(GL7), PNA+(Vector Laboratories);

plasmablasts: CD3-, B220med-lo, CD138+(281-2). In digested tissues, plasmablasts

were identified by either intracellular IgG (IgG1-A85-1, IgG2a/b-R2-40, IgG3-R40-82, all

BD Biosciences) or intracellular Ig kappa+(187.1) (Southern Biotech) using BD

Cytofix/Cytoperm kit (BD Biosciences) in lieu of CD138. Plasmablasts were confirmed by

Ki67(16A8) staining. Lymphatic endothelial cells: CD45-, CD31+ (390), PDPN+ (8.1.1);

blood endothelial cells: CD45-CD31-, PDPN-; fibroblastic reticular cells: CD45-, CD31-,

PDPN+. CD4 T cells: CD45+CD4+TCRab+ (H57-597); regulatory T cells: TCRab+,

CD4+, CD25+ (PC61), intracellular Foxp3+ (FJK-16s); follicular helper T cells: TCRab+,

CD4+, CXCR5+ (L138D7), PD1+ (29F.1A12); Th1 cells: TCRab+, CD4+, Foxp3-,

CXCR5 -, intracellular IFNg+ (XMG1.2, eBioscience); Th17 cells: TCRab+, CD4+,

Foxp3-, CXCR5-, intracellular IL17+ (TC11-18H10, BD Biosciences). The intracellular

stains for CD4+ T cell subset was done after fixing cells with the eBioscience™

Intracellular Fixation & Permeabilization Buffer Set (Cat: 88-8824-00). For Th1 and Th17

cells, staining was done after cells were stimulated with Cell Activation Cocktail with

Brefeldin A (Biolegend, Cat no. 423304) in 37C incubator for 4 hours. CCL2 was

identified by either using CCL2-GFP reporter mice or by staining. Staining for CCL2 was

performed using CCL2-FITC (245, Invitrogen) and signal was amplified by staining with

anti-FITC-biotin (1F8-1E4, Jackson ImmunoResearch) and subsequently streptavidin-

FITC (Invitrogen). A negative control for GFP signal was done by using non-transgenic

mice in CCL2-GFP mice experiments. CCL2$^{-/-}$ mice were used as negative control for

staining experiments.

For flow cytometry analysis, cells were analyzed using a FACSCanto or

FACSSymphony (BD Biosciences) and FlowJo Software (Tree Star).

Cell Sorting:

For qPCR of skin cell populations, cells from ear skin were pooled and then sorted using

a BD Influx (BD Biosciences). LECs were selected as DAPI-CD45-PDPN+CD31+ cells.

BECs were selected as DAPI-CD45-PDPN-CD31+ cells. Fibroblasts were selected as

DAPI-CD45-PDPN+CD31- cells. Macrophages were selected as DAPI-

CD45+CD11b+F4/80+ cells.

RNA Extraction

RNA was extracted from sorted cells using an RNAEasy Plus Kit (Qiagen, Germantown,

MD) and quality confirmed on a BioAnalyzer 2100 (Agilent Technologies, Santa Clara,

CA).

Real time PCR

cDNA was synthesized (iScript kit, Bio-Rad, Hercules, CA) from extracted RNA and real-

time PCR (iQ SYBR-Green Supermix kit, Bio-Rad)) was performed using primers for

PTEN (Mm_Pten_1_SG QuantiTect Primer Assay, Qiagen) and GAPDH (R:

TTGAAGTCGCAGGAGACAACCT, F: ATGTGTCCGTCGTGGATCTGA).

Autoantigen Microarray:

Blood was collected from mice, left at room temperature for 1 hour, and then centrifuged

at 3000rpm for 3 minutes. The supernatant containing the serum was then collected and

frozen in -80C until ready to be shipped for autoantigen microarray profiling at the

Genomics and Microarray Core Facitlity, UT Southwestern Medical Center. At the core,

the samples are treated with DNAse I, diluted 1:50, and incubated with autoantigen

array. The autoantibodies binding to the antigens on the array are detected with Cy3

labeled anti-IgG and cy5 labeled anti-IgM and the arrays are scanned with GenePix®

4400A Microarray Scanner. The images are analyzed using GenePix 7.0 software to

generate GPR files. The averaged net fluorescent intensity (NFI) of each autoantigen is

normalized to internal controls (IgG or IgM).

Statistics

For figures showing normalized values, each individual replicate experiment was

normalized for that experiment. For experiments that contained more than one control

sample, the mean was obtained for the control samples, and the individual control and

experimental samples were divided by this mean value to normalize to the control mean.

The Shapiro-Wilk test was used to test for normality. Unpaired t test was used for normal

data and Mann-Whitney U test was used otherwise.

Author Contributions

WGA, NS, and TTL conceived the study; MJH, WGA, MLC, AR, ESS, JHS, JS, NS, WDS, DCD, CBC, RPK, YC designed and performed the experiments and analyzed the data; ES, DE, SS, RPK, and BM provided key reagents and intellectual input. CBC and SAR designed experiments and provided intellectual input. TTL designed, supervised, and interpreted experiments. MJH, WGA, and TTL wrote the manuscript; all authors contributed to critical reading and editing.

Acknowledgments

The authors thank the Lu Lab for critical reading of the manuscript. This work was

supported by R01AI079178-121S1 (MJH), NIH T32AR071302-01 to the Hospital for

Special Surgery Research Institute Rheumatology Training Program (NS and WDS),

NIH MSTP T32GM007739 to the Weill Cornell/Rockefeller/Sloan-Kettering Tri-

Institutional MD-PhD Program (WDS), Emerson Collective Cancer Research Fund

691032 (RPK), NIH R01AI079178 (TTL), R01AR084694 (TTL), R21 AR081493 (TTL),

DOD W81XWH-22-1-0627(TTL), Lupus Research Alliance (TTL), the St. Giles

Foundation (TTL); and NIH Office of the Director grant S10OD019986 to Hospital for

Special Surgery.

References

1.   Tsokos GC, Lo MS, Costa Reis P, and Sullivan KE. New insights into the
immunopathogenesis of systemic lupus erythematosus. *Nat Rev Rheumatol.*
2016;12(12):716-30.
2.   Stannard JN, and Kahlenberg JM. Cutaneous lupus erythematosus: updates on
pathogenesis and associations with systemic lupus. *Curr Opin Rheumatol.*
2016;28(5):453-9.
3.   Schmidt E, Tony HP, Bröcker EB, and Kneitz C. Sun-induced life-threatening
lupus nephritis. *Annals of the New York Academy of Sciences.* 2007;1108:35-40.
4.   Foering K, Chang AY, Piette EW, Cucchiara A, Okawa J, and Werth VP.
Characterization of clinical photosensitivity in cutaneous lupus erythematosus.
Journal of the American Academy of Dermatology. 2013;69(2):205-13.
5.   Ansel JC, Mountz J, Steinberg AD, DeFabo E, and Green I. Effects of UV
radiation on autoimmune strains of mice: increased mortality and accelerated
autoimmunity in BXSB male mice. *J Invest Dermatol.* 1985;85(3):181-6.
6.   Ahluwalia J, and Marsch A. Photosensitivity and photoprotection in patients with
lupus erythematosus. *Lupus.* 2019;28(6):697-702.
7.   Kuhn A, Gensch K, Haust M, Meuth AM, Boyer F, Dupuy P, et al.
Photoprotective effects of a broad-spectrum sunscreen in ultraviolet-induced
cutaneous lupus erythematosus: a randomized, vehicle-controlled, double-blind
study. *J Am Acad Dermatol.* 2011;64(1):37-48.
8.   Vilá LM, Mayor AM, Valentín AH, Rodríguez SI, Reyes ML, Acosta E, et al.
Association of sunlight exposure and photoprotection measures with clinical
outcome in systemic lupus erythematosus. *P R Health Sci J.* 1999;18(2):89-94.
9.   Mikita N, Ikeda T, Ishiguro M, and Furukawa F. Recent advances in cytokines in
cutaneous and systemic lupus erythematosus. *J Dermatol.* 2011;38(9):839-49.
10. Ogunsanya ME, Kalb SJ, Kabaria A, and Chen S. A systematic review of patient-
reported outcomes in patients with cutaneous lupus erythematosus. *Br J*
Dermatol. 2017;176(1):52-61.
11. Tebbe B, and Orfanos CE. Epidemiology and socioeconomic impact of skin
disease in lupus erythematosus. *Lupus.* 1997;6(2):96-104.
12. Sim JH, Ambler WG, Sollohub IF, Howlader MJ, Li TM, Lee HJ, et al. Immune
Cell–Stromal Circuitry in Lupus Photosensitivity. *The Journal of Immunology.*
2021;206(2):302-9.
13. Kahlenberg JM. Rethinking the Pathogenesis of Cutaneous Lupus. *J Invest*
Dermatol. 2021;141(1):32-5.
14. Patel J, Borucki R, and Werth VP. An Update on the Pathogenesis of Cutaneous
Lupus Erythematosus and Its Role in Clinical Practice. *Curr Rheumatol Rep.*
2020;22(10):69.
15. Oliver G, Kipnis J, Randolph GJ, and Harvey NL. The Lymphatic Vasculature in
the 21(st) Century: Novel Functional Roles in Homeostasis and Disease. *Cell.*
2020;182(2):270-96.
16. Fonseca DM, Hand TW, Han SJ, Gerner MY, Glatman Zaretsky A, Byrd AL, et al.
Microbiota-Dependent Sequelae of Acute Infection Compromise Tissue-Specific
Immunity. *Cell.* 2015;163(2):354-66.

17. Thomas SN, Rutkowski JM, Pasquier M, Kuan EL, Alitalo K, Randolph GJ, et al.
Impaired Humoral Immunity and Tolerance in K14-VEGFR-3-Ig Mice That Lack
Dermal Lymphatic Drainage. *The Journal of Immunology.* 2012;189(5):2181-90.
18. Kajiya K, and Detmar M. An Important Role of Lymphatic Vessels in the Control
of UVB-Induced Edema Formation and Inflammation. *J Invest Dermatol.*
2006;126(4):920-2.
19. Rajasekhar L, Habibi S, Sudhakar P, and Gumdal N. Lymphatic obstruction as a
cause of extremity edema in systemic lupus erythematosus. *Clin Rheumatol.*
2013;32 Suppl 1:S11-3.
20. Daniel A, Bagnato G, Vital E, and Del Galdo F. Chylous ascites in a patient with
an overlap syndrome: a surprising response to rituximab. *BMJ case reports.*
2017;2017.
21. Dalvi SR, Yildirim R, Santoriello D, and Belmont HM. Pseudo-pseudo Meigs'
syndrome in a patient with systemic lupus erythematosus. *Lupus.*
2012;21(13):1463-6.
22. Lee CK, Han JM, Lee KN, Lee EY, Shin JH, Cho YS, et al. Concurrent
occurrence of chylothorax, chylous ascites, and protein-losing enteropathy in
systemic lupus erythematosus. *J Rheumatol.* 2002;29(6):1330-3.
23. Ulvmar MH, Werth K, Braun A, Kelay P, Hub E, Eller K, et al. The atypical
chemokine receptor CCRL1 shapes functional CCL21 gradients in lymph nodes.
Nat Immunol. 2014;15(7):623-30.
24. Gretz JE, Norbury CC, Anderson AO, Proudfoot AE, and Shaw S. Lymph-borne
chemokines and other low molecular weight molecules reach high endothelial
venules via specialized conduits while a functional barrier limits access to the
lymphocyte microenvironments in lymph node cortex. *J Exp Med.*
2000;192(10):1425-40.
25. Li L, Wu J, Abdi R, Jewell CM, and Bromberg JS. Lymph node fibroblastic
reticular cells steer immune responses. *Trends Immunol.* 2021;42(8):723-34.
26. Pikor NB, Cheng H-W, Onder L, and Ludewig B. Development and
Immunological Function of Lymph Node Stromal Cells. *The Journal of*
Immunology. 2021;206(2):257-63.
27. Krishnamurty AT, and Turley SJ. Lymph node stromal cells: cartographers of the
immune system. *Nat Immunol.* 2020;21(4):369-80.
28. Dasoveanu DC, Park HJ, Ly CL, Shipman WD, Chyou S, Kumar V, et al. Lymph
node stromal CCL2 limits antibody responses. *Sci Immunol.* 2020;5(45).
29. Yazici Y, Erkan D, Levine DM, Parker TS, and Lockshin MD. Protein-losing
enteropathy in systemic lupus erythematosus: report of a severe, persistent case
and review of pathophysiology. *Lupus.* 2002;11(2):119-23.
30. von der Weid PY, Rehal S, and Ferraz JG. Role of the lymphatic system in the
pathogenesis of Crohn's disease. *Curr Opin Gastroenterol.* 2011;27(4):335-41.
31. Rossi A, Sozio F, Sestini P, Renzoni EA, Khan K, Denton CP, et al. Lymphatic
and blood vessels in scleroderma skin, a morphometric analysis. *Human*
Pathology. 2010;41(3):366-74.
32. Sevim E, Siddique S, Chalasani MLS, Chyou S, Shipman W, O'Shea O, et al.
Mammalian Target of Rapamycin Pathway Assessment in Antiphospholipid
Antibody Positive Patients with Livedo. *J Rheumatol.* 2022.
33. Huggenberger R, Siddiqui SS, Brander D, Ullmann S, Zimmermann K,
Antsiferova M, et al. An important role of lymphatic vessel activation in limiting
acute inflammation. *Blood.* 2011;117(17):4667-78.

34. Furukawa F. Photosensitivity in cutaneous lupus erythematosus: lessons from
mice and men. *J Dermatol Sci.* 2003;33(2):81-9.
35. Shipman WD, Chyou S, Ramanathan A, Izmirly PM, Sharma S, Pannellini T, et
al. A protective Langerhans cell–keratinocyte axis that is dysfunctional in
photosensitivity. *Science Translational Medicine.* 2018;10(454):eaap9527.
36. Menke J, Hsu MY, Byrne KT, Lucas JA, Rabacal WA, Croker BP, et al. Sunlight
Triggers Cutaneous Lupus through a CSF-1-Dependent Mechanism in MRL-
Faslpr Mice. *J Immunol.* 2008;181(10):7367-79.
37. Yokogawa M, Takaishi M, Nakajima K, Kamijima R, Fujimoto C, Kataoka S, et al.
Epicutaneous application of toll-like receptor 7 agonists leads to systemic
autoimmunity in wild-type mice: a new model of systemic Lupus erythematosus.
Arthritis Rheumatol. 2014;66(3):694-706.
38. Mizui M, Koga T, Lieberman LA, Beltran J, Yoshida N, Johnson MC, et al. IL-2
protects lupus-prone mice from multiple end-organ damage by limiting CD4-CD8-
IL-17-producing T cells. *J Immunol.* 2014;193(5):2168-77.
39. Li TM, Zyulina V, Seltzer ES, Dacic M, Chinenov Y, Daamen AR, et al. The
interferon-rich skin environment regulates Langerhans cell ADAM17 to promote
photosensitivity in lupus. *eLife.* 2024;13:e85914.
40. Lurie F, Malgor RD, Carman T, Dean SM, Iafrati MD, Khilnani NM, et al. The
American Venous Forum, American Vein and Lymphatic Society and the Society
for Vascular Medicine expert opinion consensus on lymphedema diagnosis and
treatment. *Phlebology.* 2022;37(4):252-66.
41. Lasinski BB, McKillip Thrift K, Squire D, Austin MK, Smith KM, Wanchai A, et al.
A systematic review of the evidence for complete decongestive therapy in the
treatment of lymphedema from 2004 to 2011. *Pm r.* 2012;4(8):580-601.
42. Kataru RP, Baik JE, Park HJ, Ly CL, Shin J, Schwartz N, et al. Lymphatic-
specific intracellular modulation of receptor tyrosine kinase signaling improves
lymphatic growth and function. *Sci Signal.* 2021;14(695):eabc0836.
43. Schwager S, Renner S, Hemmerle T, Karaman S, Proulx ST, Fetz R, et al.
Antibody-mediated delivery of VEGF-C potently reduces chronic skin
inflammation. *JCI Insight.* 2018;3(23).
44. Skopelja-Gardner S, Tai J, Sun X, Tanaka L, Kuchenbecker JA, Snyder JM, et al.
Acute skin exposure to ultraviolet light triggers neutrophil-mediated kidney
inflammation. *Proc Natl Acad Sci U S A.* 2021;118(3).
45. Liu Y, Seto NL, Carmona-Rivera C, and Kaplan MJ. Accelerated model of lupus
autoimmunity and vasculopathy driven by toll-like receptor 7/9 imbalance. *Lupus
Science & Medicine.* 2018;5(1):e000259.
46. Lewis SM, Williams A, and Eisenbarth SC. Structure and function of the immune
system in the spleen. *Sci Immunol.* 2019;4(33).
47. Vieira P, and Rajewsky K. The half-lives of serum immunoglobulins in adult mice.
Eur J Immunol. 1988;18(2):313-6.
48. Hodgkinson LM, Wu TT, and Fiorentino DF. Dermatomyositis autoantibodies:
how can we maximize utility? *Annals of translational medicine.* 2021;9(5):433.
49. Kajiya K, Sawane M, Huggenberger R, and Detmar M. Activation of the VEGFR-
3 pathway by VEGF-C attenuates UVB-induced edema formation and skin
inflammation by promoting lymphangiogenesis. *J Invest Dermatol.*
2009;129(5):1292-8.
50. Mohr E, Serre K, Manz RA, Cunningham AF, Khan M, Hardie DL, et al. Dendritic
cells and monocyte/macrophages that create the IL-6/APRIL-rich lymph node

microenvironments where plasmablasts mature. *J Immunol.* 2009;182(4):2113-
23.
51. Smith MH, Gao VR, Periyakoil PK, Kochen A, DiCarlo EF, Goodman SM, et al.
Drivers of heterogeneity in synovial fibroblasts in rheumatoid arthritis. *Nature*
Immunology. 2023;24(7):1200-10.
52. Wei K, Korsunsky I, Marshall JL, Gao A, Watts GFM, Major T, et al. Notch
signalling drives synovial fibroblast identity and arthritis pathology. *Nature.*
2020;582(7811):259-64.
53. Tomei AA, Siegert S, Britschgi MR, Luther SA, and Swartz MA. Fluid flow
regulates stromal cell organization and CCL21 expression in a tissue-engineered
lymph node microenvironment. *J Immunol.* 2009;183(7):4273-83.
54. Horsnell HL, Tetley RJ, De Belly H, Makris S, Millward LJ, Benjamin AC, et al.
Lymph node homeostasis and adaptation to immune challenge resolved by
fibroblast network mechanics. *Nat Immunol.* 2022;23(8):1169-82.
55. Assen FP, Abe J, Hons M, Hauschild R, Shamipour S, Kaufmann WA, et al.
Multitier mechanics control stromal adaptations in the swelling lymph node. *Nat*
Immunol. 2022;23(8):1246-55.
56. Howlader MJ, Rashighi M, Santambrogio L, and Lu TT. Lymphatic messengers:
Non-antigen soluble mediators from diseased tissues to draining lymph nodes.
Curr Opin Immunol. 2025;93:102536.
57. Perez-Shibayama C, Islander U, Lütge M, Cheng HW, Onder L, Ring SS, et al.
Type I interferon signaling in fibroblastic reticular cells prevents exhaustive
activation of antiviral CD8(+) T cells. *Sci Immunol.* 2020;5(51).
58. Platt AM, Rutkowski JM, Martel C, Kuan EL, Ivanov S, Swartz MA, et al. Normal
dendritic cell mobilization to lymph nodes under conditions of severe lymphatic
hypoplasia. *J Immunol.* 2013;190(9):4608-20.
59. Kumar V, Dasoveanu DC, Chyou S, Tzeng T-C, Rozo C, Liang Y, et al. A
dendritic-cell-stromal axis maintains immune responses in lymph nodes.
Immunity. 2015;42:719-30.
60. Loo CP, Nelson NA, Lane RS, Booth JL, Loprinzi Hardin SC, Thomas A, et al.
Lymphatic Vessels Balance Viral Dissemination and Immune Activation following
Cutaneous Viral Infection. *Cell Reports.* 2017;20(13):3176-87.
61. Billi AC, Ma F, Plazyo O, Gharaee-Kermani M, Wasikowski R, Hile GA, et al.
Nonlesional lupus skin contributes to inflammatory education of myeloid cells and
primes for cutaneous inflammation. *Science Translational Medicine.*
2022;14(642):eabn2263.
62. Der E, Suryawanshi H, Morozov P, Kustagi M, Goilav B, Ranabothu S, et al.
Tubular cell and keratinocyte single-cell transcriptomics applied to lupus nephritis
reveal type I IFN and fibrosis relevant pathways. *Nat Immunol.* 2019;20(7):915-
27.
63. Psarras A, Alase A, Antanaviciute A, Carr IM, Md Yusof MY, Wittmann M, et al.
Functionally impaired plasmacytoid dendritic cells and non-haematopoietic
sources of type I interferon characterize human autoimmunity. *Nature*
Communications. 2020;11(1):6149.
64. Skopelja-Gardner S, An J, Tai J, Tanaka L, Sun X, Hermanson P, et al. The early
local and systemic Type I interferon responses to ultraviolet B light exposure are
cGAS dependent. *Sci Rep.* 2020;10(1):7908.

65. Clement CC, D'Alessandro A, Thangaswamy S, Chalmers S, Furtado R, Spada
S, et al. 3-hydroxy-L-kynurenamine is an immunomodulatory biogenic amine. *Nat*
Commun. 2021;12(1):4447.
66. Russo E, Teijeira A, Vaahtomeri K, Willrodt AH, Bloch JS, Nitschke M, et al.
Intralymphatic CCL21 Promotes Tissue Egress of Dendritic Cells through
Afferent Lymphatic Vessels. *Cell Rep.* 2016;14(7):1723-34.
67. Saxena V, Piao W, Li L, Paluskievicz C, Xiong Y, Simon T, et al. Treg tissue
stability depends on lymphotoxin beta-receptor- and adenosine-receptor-driven
lymphatic endothelial cell responses. *Cell Rep.* 2022;39(3):110727.
68. Lu B, Rutledge BJ, Gu L, Fiorillo J, Lukacs NW, Kunkel SL, et al. Abnormalities in
monocyte recruitment and cytokine expression in monocyte chemoattractant
protein 1-deficient mice. *J Exp Med.* 1998;187(4):601-8.
69. Shi C, Jia T, Mendez-Ferrer S, Hohl TM, Serbina NV, Lipuma L, et al. Bone
marrow mesenchymal stem and progenitor cells induce monocyte emigration in
response to circulating toll-like receptor ligands. *Immunity.* 2011;34(4):590-601.
70. Hohl TM, Rivera A, Lipuma L, Gallegos A, Shi C, Mack M, et al. Inflammatory
monocytes facilitate adaptive CD4 T cell responses during respiratory fungal
infection. *Cell Host Microbe.* 2009;6(5):470-81.
71. Escobedo N, Proulx ST, Karaman S, Dillard ME, Johnson N, Detmar M, et al.
Restoration of lymphatic function rescues obesity in Prox1-haploinsufficient mice.
JCI Insight. 2016;1(2).
72. Chia JJ, Zhu T, Chyou S, Dasoveanu DC, Carballo C, Tian S, et al. Dendritic
cells maintain dermal adipose-derived stromal cells in skin fibrosis. *J Clin Invest.*
2016;126(11):4331-45.
73. Chyou S, Benahmed F, Chen J, Kumar V, Tian S, Lipp M, et al. Coordinated
regulation of lymph node vascular-stromal growth first by CD11c+ cells and then
by T and B cells. *J Immunol.* 2011;187(11):5558-67.

Figure 1. Patients with SLE and murine SLE models show evidence of cutaneous

lymphatic dysfunction

(A-C) Punch biopsies of forearm skin from healthy control (HC), positive antiphospholipid

antibodies (APL) without SLE, and SLE subjects with or without APL were stained for

CD31+PDPN+ lymphatic vessels. (A) Representative photomicrographs. Arrowheads

point to lymphatic vessels. (B) Lumenal area per vessel. Each symbol is an individual

subject. (C) Number of lymphatic vessels per tissue area. (D-H) LPR mice and MRL

controls were treated with ultraviolet radiation (UVR) for 4 consecutive days, injected in

ear pinna with Evans blue dye (EB) 1 day after final UVR dose, and ear harvested to

assess EB content 1 day later, as in (D). (E) EB retention and (F) ear swelling was

quantified after UVR or (G,H) with no UVR. (I-K) Ears were examined by flow cytometry

at 24 hours after final dose of UVR. (I) Monocyte, (J) TCR$\alpha\beta$ CD4+ T cell, and (K)

TCRab+ CD3+ CD4-CD8- "double negative" (DN) T cell numbers. (L-O) B6 mice

received IMQ on the right ear and exposed to UVR, and EB retention after intradermal

injection of left ear or footpad was compared to responses in vehicle-treated control

mice, as in (L). (M) EB retention in ear. (N) Ear swelling. (O) EB retention in footpad. (P-

Q) Left ear was collected 24 hours after 1 or 4 doses of UVR. (P) Monocyte and (Q)

TCR$\alpha\beta$ CD4 T cell numbers. Each symbol represents one mouse; n = 2 to 21 per

condition; data are from 2 (I, J, K), 3 (M, O), 4 (G, H), 6 (E, F), 10 (N), and 11 (P, Q)

independent experiments. Normality was assessed using the Shapiro-Wilk test. If

normal, unpaired t test was used. If data were not normal, Mann-Whitney U test was

used. ***$P<0.001$, **$P<0.01$, *$P<0.05$, NS=not significant. Error bars represent SD.

Figure 2. Improving lymphatic flow reduces cutaneous photosensitive responses in SLE models

(A-J) LPR mice and (K-P) B6-IMQ mice were treated with UVR and manual lymphatic drainage (MLD) targeting the left ear or were control handled. Left ear was then examined. (A, K) Experimental design. (B) Illustration of MLD technique. Please see

Methods for details. (C, L) EB retention and (D, E, M) ear thickness, absolute (D,M) or

normalized (E). (F, N) TCR$\alpha\beta$ CD4 T cell numbers, (G,O) normalized to controls. (H, P)

Percentage of CD4 T cells that express IFNg. (I) DN T cell numbers and (J) percentage

that express IFNg. (Q-U) Flt4Cre[ERT2] PTEN[fl/fl] mice treated with tamoxifen (LEC[PTEN]) or

without (LEC[WT]) were treated with IMQ on right ear and UVR before left ear skin

assessment, as in (Q). (R) EB retention, (S) skin thickness, (T) TCR$\alpha\beta$ CD4 T cell

numbers, and (U) IFNg+ percentage. (V, W) Non-transgenic B6 mice were treated as

described in (Q). (V) EB retention and (W) ear thickness. Each symbol represents one

mouse; n= 3 to 13 per condition; data are from 2 (H, J, L, V, W), 3 (N, O, P, R, T, U), 4 (F,

G, I, S), 6 (C), 7 (D, E), and 9 (M) independent experiments. Normality was assessed

using the Shapiro-Wilk test. If normal, unpaired t test was used. If data were not normal,

Mann-Whitney U test was used. ***P<0.001, **P<0.01, *P<0.05, NS=not significant.

Error bars represent SD.

Figure 3. Improving lymphatic flow reduces draining lymph node B and T cell

responses in SLE models

(A-Q) Left auricular lymph nodes of (A-I) LPR, (J-Q) B6-IMQ, and (R-X) LEC^PTEN-IMQ

and LEC^{WT}- IMQ mice that were treated as in Figure 2A, K, and Q were examined. (A, J,

R) Lymph node cellularity. (B, K, S) B cell, (C, L, T) germinal center B cell, (D, M, U)

plasmablast, and (E, N, V) TCR$\alpha\beta$ CD4 T cell numbers. (F, O, W) CD4 T cell numbers

normalized to control. (G, P, X) Percentage of CD4 T cells that express IFNg. (H, Q, Y)

CD8 and (I) DN T cell numbers. Each symbol represents one mouse; n= 4 to 17 per

condition; data are from 2 (X), 3 (G, P, Q), 4 (A-C, H-K, M, V, W, Y), 5 (D, N, O), 6 (E, F,

L, R, U), and 8 (S, T) independent experiments. Normality was assessed using the

Shapiro-Wilk test. If normal, unpaired t test was used. If data were not normal, Mann-

Whitney U test was used. ***P<0.001, **P<0.01, *P<0.05, NS=not significant. Error bars

represent SD.

Figure 4. Improving lymphatic flow long term reduces systemic disease activity

(A) Splenic weight of LEC^PTEN-IMQ and LEC^WT-IMQ mice treated with UVR for 4 days.

(B) Splenic weight of LPR mice treated with UVR and MLD concurrently or control

handled for 5 weeks. (C,D) Heatmap of normalized signal intensity (NSI) from

autoantigen microarray panel for IgG of MRL mice, LPR mice treated with UVR, and

LPR mice treated with both UVR and MLD for 4-5 weeks. (C) Each column represents

one mouse and autoantibodies with significant differences between LPR UV and LPR

UV+MLD are labelled in red. (D) Each column represents average NSI of all mice serum

samples. Each symbol represents one mouse; n= 1 to 12 per condition; data are from 1

(B), 2 (C, D), and 6 (A) independent experiments. Normality was assessed using the

Shapiro-Wilk test. If normal, unpaired t test was used. If data were not normal, Mann-

Whitney U test was used. ***$P<0.001$, **$P<0.01$, *$P<0.05$, NS=not significant. Error bars

represent SD.

1002

1003

1004

Figure 5. Improving lymphatic flow increases lymph node FRC proliferation, FRC

CCL2, monocyte ROS generation, and limits plasmablast numbers in a monocyte-

dependent manner

(A-Q) Left auricular lymph nodes of indicated mice exposed to 4 days of UVR were

examined. (A) FRC expression of GFP in CCL2-GFP reporter mice treated with IMQ that

received MLD or control handling with UVR. (B) CCL2 expression by FRCs in LEC^{PTEN}-

IMQ and LEC^{WT}- IMQ mice. (A, B) Representative histogram (left) and graph (right).

MFI=geometric mean fluorescence intensity. (C-G) B6-IMQ mice received 4 days of +/-

MLD with UVR. (C) FRC numbers. (D) Percentage of FRCs that express ki67, (E)

normalized to control handled mice. (F) Ly6C^{hi} monocyte numbers, flow cytometry gating

(left) and numbers (right). (G) Monocyte ROS measured using CM-H2DCFDA,

representative histograms (left) and relative MFI of CM-H2DCFDA (right). (H-L) B6-IMQ

mice were treated with anti-Gr-1 or isotype control at days -1, 0, and +2 of UVR and

MLD treatments as shown in (H). (I) Monocyte numbers and (J) normalized to isotype

control. (K) Germinal center B cell and (I) plasmablast numbers. (M-Q) CCR2-DTR mice

were treated with DT at days 0 and 2 of UVR and MLD treatments as shown in (M). (N)

Monocyte numbers and (O) normalized to control. (P) Germinal center B cell and (Q)

plasmablast numbers. Each symbol represents one mouse; n= 6 to 19 per condition;

data are from 2 (B), 3 (A, I-L), 5 (N, O), 6 (D-G, P, Q), and 7 (C) independent

experiments. Normality was assessed using the Shapiro-Wilk test. If normal, unpaired t

test was used. If data were not normal, Mann-Whitney U test was used. ***$P<0.001$,

**$P<0.01$, *$P<0.05$, NS=not significant. Error bars represent SD.

1027

1028

1029

1030