1 **Title: Maturational networks of fetal brain activity**
2

3 **Authors:** Karolis VR[1,2,*], Fitzgibbon SP[2], Cordero-Grande L[3], Farahibozorg SR[2], Price AN[1],
4 Hughes EJ[1], Fetit AE[4,5], Kyriakopoulou V[1], Pietsch, M[1], Rutherford MA[1], Rueckert D[4,6], Hajnal
5 JV[1], Edwards AD[1,7], O'Muircheartaigh J[1,7,8], Duff EP[2,9,#] & Arichi T[1,7,10,#]
6

    1. Centre for the Developing Brain, School of Biomedical Engineering and Imaging
       Sciences, King's College London, UK.
    2. Wellcome Centre for Integrative Neuroimaging, FMRIB, Nuffield Department of
       Clinical Neurosciences, University of Oxford, UK.
    3. Biomedical Image Technologies, ETSI Telecomunicación, Universidad Politécnica de
       Madrid & CIBER-BBN, Spain
    4. Biomedical Image Analysis Group, Department of Computing, Imperial College
       London, UK.
    5. UKRI CDT in Artificial Intelligence for Healthcare, Department of Computing,
       Imperial College London, UK
    6. Department of Medicine and Informatics, Technical University of Munich, Germany.
    7. MRC Centre for Neurodevelopmental Disorders, King's College London, UK.
    8. Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry,
       Psychology & Neuroscience, King's College London, UK
    9. Department of Brain Sciences, Faculty of Medicine, Imperial College London, UK.
    10. Paediatric Neurosciences, Evelina London Children's Hospital, Guy's and St Thomas'
       NHS Foundation Trust, UK.
       * - corresponding author, slava.karolis@kcl.ac.uk
       # - equally contributing last authors

ABSTRACT
A key feature of the fetal period is the rapid emergence of organised patterns of
spontaneous brain activity. However, characterising this process in utero using functional
MRI is inherently challenging and requires analytical methods which can capture the
constituent developmental transformations. Here, we introduce a novel analytical
framework, termed "maturational networks", that achieves this by conceptualising
functional networks as an emerging property of the developing brain. Compared to standard
network analysis methods that assume consistent patterns of connectivity across
development, our method incorporates age-related changes in connectivity directly into
network estimation. This enables us to reveal the richness of emerging functional
connections and the hierarchy of their maturational relationships in the in-utero brain with
unprecedented anatomical specificity. We show that the associative areas play a central role
within prenatal functional architecture, therefore indicating that functional connections
supporting high-level human cognition start emerging even prior to exposure to the extra-
utero environment.

47
48
49 **INTRODUCTION**
50
Does a 'thing' possess invariant properties that define its 'being', or does its essence reveal
itself in the process of a perpetual change, i.e., in its 'becoming'? This ancient intellectual
dilemma, conceived by an early Greek philosopher Heraclitus, has been entwined in the
centuries-long evolution of human knowledge (*1, 2*). At its core, it reflects a fundamental
problem of selecting an appropriate representational framework for studying a
phenomenon while offering a choice between two extreme alternatives. On the one hand, a
description of invariant (canonical, typical) characteristics serves a purpose of giving a
phenomenon a concrete definition and thus differentiating it from other things. On the
other hand, representations that characterise a phenomenon as a process are more fitting if
the phenomenon constitutes a sequence of superseding transient states with ill-defined
invariant characteristics.
62
The notion of functional networks in the fetal brain is a case in point for the latter. Evidence
from animal models suggests that intrinsically generated neural activity in the prenatal brain
first begins with local direct propagation before progressing to larger bursts of spontaneous
activity which help to establish local circuitry (*3*). At around 26 weeks of gestation, ex-utero
functional MRI (fMRI) studies of very preterm infants (*4*) show that spatially distinct resting-
state networks can be identified, initially consisting of local patterns of connectivity with a
lack of long range interhemispheric or dorsocaudal connections. Towards term equivalent
age, these networks evolve into a set of spatially distributed (multi-nodal) co-activation
patterns resembling those seen in adults (*5, 6*), reflecting a generic drift of organic functions
towards forming increasingly complex systems (*7*). Such rapid developmental changes mean
that functional networks in the prenatal period possess the attributes of an intrinsically non-
static entity, a characteristic example of Heraclitian "becoming".
75
Previous research has demonstrated that, despite enormous technological challenges,
functional connectivity in utero can also be studied using resting-state fMRI (*8-10*). This
opens up an opportunity for the use of standard approaches to group-level fMRI network
analyses (*11*) such as group independent component analysis (group-ICA) (*12-14*). The latter
describes functional networks as a collection of spatial maps (*15*), each of them charting
areas linked together by the strength of covariation between the timecourses of their
fluctuating intrinsic activity. However, utility of this method for application with fetal data
remains an open question, both conceptually and when considering the unique signal
properties of the data acquired in utero. Conceptually, an assumption embedded into this
method is that a group-level spatial map characterises a canonical form of a functional
network with respect to its individual manifestations, thereby downgrading developmental
changes in its spatial layout to the status of non-systematic, and likely underestimated (*16*),
inter-subject variability. On a practical level, application of group-ICA to fetal data typically
renders maps of poorly localised and segregated regions, lacking network-like features, such
as the presence of spatially non-contingent brain areas (*11*). This may be explained by the
weakness of long-distance connectivity in the fetal brain but may also be a consequence of
inherently high levels of motion and low signal-to-noise ratio in this data, which adversely
affects the detection of long-distance connections (*17, 18*). As a result, coherent

developmental features that are fundamental to both a definition and understanding of the
neuroscientific basis of functional networks in utero are likely lost using this standard
approach.
97
In this paper, in order to capture the maturational transiency of connectivity states, we
introduce an alternative perspective on resting-state functional networks in utero, which we
call "maturational networks", or matnets for conciseness. The key feature of this framework
is that it incorporates age-related changes in connectivity into network estimation, thereby
characterising functional networks as an emerging property of the brain. At its core, it builds
on Flechsig's idea (*19*), that functionally related areas mature together. In contrast to the
standard analytical approach of ICA, which utilises correlational structure to factorize
networks, our approach leverages age-related changes in correlations in order to
characterise maturational modes of variation in the data. The utility of this approach is
demonstrated in in-utero fMRI data acquired as part of the developing Human Connectome
Project (dHCP) (*20, 21*), an open science initiative aiming to map brain connectivity across
the perinatal period, that were reconstructed and preprocessed using specially developed
methodologies (*22-24*). We show that our approach overcomes inherent limitations of fMRI
data acquired in-utero for characterising mid- and long-distance connectivity, and for
inference about the developmental trajectory of the fetal functional connectome. Moreover,
it enables factorisation of spatial patterns that fit better the concept of resting-state
network as we understand it from the studies of more mature brains, that is, as spatially
distributed configurations encompassing non-adjacent brain areas (*25, 26*). Finally, we show
that maturational networks lead to a new perspective on the macro-scale developmental
relationships in the human brain, the "maturational connectome".
118
119
120 **RESULTS**
121
Resting state fMRI data from 144 healthy fetuses with an age range between 25 and 38
weeks gestation (Supplementary Figure 1) were acquired over 12.8 mins on a 3T Philips
Achieva system (Best, NL) (*27*) as part of the developing Human Connectome Project (dHCP).
All of the fetal brain images were clinically reported and showed appropriate appearances
for their gestational age with no acquired lesions or congenital malformations. The data
underwent dynamic geometric correction for distortions, slice-to-volume motion correction
(*22, 23*) and temporal denoising (*24*), followed by their registration to a common space to
enable group-level analyses (*6*).
130
131
132
133
134 *The framework*
135
In order to demonstrate the utility of our approach, we note that developmental changes in
a spatial layout of functional networks can be modelled retrospectively within the standard
group-ICA approach using several post-processing steps (*14*), as shown in Figure 1A. The
results of this modelling can therefore serve as a reference for comparison with the results
of matnets modelling. In brief, the conventional modelling approach involves the estimation

of group-level ("canonical") spatial maps, followed by the two steps of dual regression (DR)
(*14*), i.e., a sequence of spatial and temporal regressions performed against individual data,
in order to obtain subject-specific variants of the group maps, followed by a mass-univariate
(i.e., voxelwise) modelling of the latter using age as a covariate. The key step is the dual
regression step, that "permits the identification of between-subject differences in resting
functional connectivity based on between-subject similarities"(*14*), where a subject-specific
map represents the individualised manifestation of a group map.

148
In contrast, our matnets approach, shown in Figure 1B, attempts to derive maps of
maturational modes of variation in a direct manner, in essence by reversing the order of
operations while omitting the intermediate steps of dual regression. It runs as follows. At the
first step, a dense N voxels by N voxels connectome is computed for each subject separately.
Each element of the dense connectome is then fitted across subjects with age as covariate
and converted using t-statistics into a maturational dense connectome, i.e., a matrix in
which elements contain the estimates of the age effect. An ICA factorisation of the
maturational dense connectome is then performed to obtain spatial maps that comprise
voxels with similar maturational profiles across the study period, or matnets.

158
Figure 1. Two approaches to maturational analysis of the functional networks. (A) Group-ICA + dual regression pipeline and
its outputs. The pipeline allows modelling maturational changes in the spatial layout of the networks using mass-univariate
analysis of the subject-specific variants of the group maps. The latter are derived using dual regression. (B) Pipeline for
derivation of maturational networks. It directly leverages age-related changes to derive networks instead of estimating
subject-specific variants of the group-level maps. se – standard error

Univariate spatial properties of group-average correlations and age-related differences in
correlations

166
The efficiency of either method for network analysis, for instance in terms of their ability to
discover meaningful spatial relationships, is contingent on the relevant signal properties of
the data, which remain poorly understood for the in-utero fMRI. A brief description of these

properties would assist subsequent interpretations and inform analytical choices.
Consequently, we provide a short summary of the univariate spatial properties of the two
metrics that are expected to shape the results of the group-ICA and matnets analyses:
respectively, group-average correlations and the effect of age (t-value) on the strength of
correlations.

The generic spatial structure of the two metrics can be easily appreciated by considering
connectivity maps from seed regions to the whole brain ("seed-to-brain" maps). The maps of
the group-average correlation for six cortical seeds (3 for each hemisphere; estimated from
the correlation between the mean timecourse of voxels within a seed mask and timecourses
of all voxels in the brain, and then averaged across subjects) are shown in Figure 2A (left
panel). The conspicuous feature of these maps is a presence of a strong distance dependent
gradient, indicating signal smearing over the immediate neighbourhood of the seed. This
effect transgresses anatomical boundaries, as demonstrated in a context where the
anatomical and purely spatial distances can be disentangled (Supplementary Figure 2) and
shows a spatially indiscriminate character as it could equally be replicated for seeds located
in the white matter (Supplementary Figure 3).

In comparison, the configuration of the spatial maps for the age-related effect on correlation
(that is, instead of being averaged across subjects, the seed-to-brain correlation maps were
fitted were fitted voxel-wise with age as a covariate) for the same set of seeds reveals two
components of relevance: a negative local component and a positive mid- and long-distance
component (Figure 2A, right panel). The negative local component is revealed by a
distribution of high negative values in the proximity of the seed. This local component, which
implies that the strength of distance-dependent gradients in connectivity structure is
negatively associated with age at a short distance, occurs in a spatially indiscriminate
manner (Supplementary Figure 4), though less obviously in white matter, possibly due to a
greater signal blurring within this tissue. Otherwise, the positive mid- and long-distance
component is characterised by an age-related increase in correlation strength between seed
and other grey matter regions.

Furthermore, the spatial distance also determines a similarity (i.e., spatial correlation)
between pairs of maps (Figure 2B) for group-average correlation, which suggests that spatial
distance may become a dominant factor for the fusion of the voxels into networks in
analyses based on the correlational structure of the data, such as group-ICA. Conversely, the
same figure (Figure 2B) also shows that the similarity between age-effect maps was
unaffected by the spatial distance between seeds used to produce these maps. This suggests
that leveraging positive age-related associations for the network construction can
potentially reveal a rich set of spatially distributed patterns with improved specificity. In this
view, matnets were derived using a factorization of the positively thresholded maturational
dense connectome.

Group ICA maps and estimated age-related differences in their layout

The results of the conventional group-ICA factorisation are shown in Figure 3A. The
appearance of the spatial maps suggest that they inherit certain signal properties that had
previously been revealed in the univariate analysis. Thus, their "blurry" appearance is
reminiscent of the increased local signal correlations observed in univariate maps of seed-to-
brain group-average correlations. In addition, the location of the peaks in many group-ICA
maps tended to be biased away from the cortex towards the white matter and a local low-

to-high ramp of the component values could often be traced along the boundary between grey and white matter tissues (Supplementary Figure 5). Despite the above characteristics, most components have anatomically plausible layouts, encompassing a diverse range of functionally relevant areas. The components where peaks were most firmly located within cortical ribbon, were found in sensorimotor and pre-motor areas (e.g., components #16,17, 23, 24).

Meanwhile, the analysis of age-related changes in the spatial layout of the networks using the dual regression approach appear to be affected by a specific bias, as shown using the examples of the spatial maps of the first 3 components and the corresponding maps of the age effect in Figure 3B, demonstrating a negative effect of age (i.e., a decrease of connectivity with age) in the most representative component voxels. This somewhat counter-intuitive pattern was observed for all group-ICA components. Furthermore, as Figure 3C shows, there was a high negative spatial correlation between component group-ICA component spatial maps and corresponding t-maps of the age effect. As we will discuss later, this pattern appears to be a direct consequence of the signal properties earlier highlighted in the context of the univariate analyses, showing that there is a negative association between age and strength of correlations for voxels surrounding a seed.

Figure 3. Results of group-ICA analysis. All spatial maps are shown in radiological orientation. (A) Group-level spatial maps, thresholded at abs(z) >3. (B). Spatial maps of the first 3 components (upper row) and the corresponding maps of the age

Maturational networks (matnets)

The above analysis demonstrates an inability to reconstruct coherent maturational
relationships in the fetal fMRI data using tools that are widely used in standard network
analysis in pediatric and adult populations. In contrast, results from the maturational
network factorization, presented in Figure 4A, reveal spatial configurations of a high
anatomical validity, including locality within the grey matter (Supplementary Figure 6). In
order to ascertain the robustness of the method, we repeated the analysis in split-half
samples and found a good replicability of the component spatial properties (Supplementary
Figures 7-10). Compared to the group-ICA, the characterized matnets showed a greater
spatial complexity, encompassing non-adjacent areas. For instance, the main node of matnet
#11 spatially overlapped with that of group-ICA #24 but in addition encompassed areas in
lateral central and pre-motor cortices. Another example is the bilateral matnet component
#7, in which the left-hemisphere sub-division overlapped with a spatially compact group-ICA
component #16. Furthermore, the matnet maps tended to demonstrate more anatomically
specific local variations of intensity compared to the group-ICA maps (Figure 4B) which are
reminiscent of the spatial specificity in the age-effect seed-to-brain maps from the
univariate analyses. For instance, the matnet map #7 in Figure 4B has multiple poles,
distributed across the somatosensory, motor and premotor cortices, which suggests an early
integration of local circuits supporting different functions. In contrast, group-ICA
components were typically characterised by a tendency to have only one centre-of-gravity.

Whole-brain maturational relationships

Within the standard group-ICA+DR approach, a compact description of the functional relationships between brain networks can be provided using "netmats"(*28*), a matrix of correlations between component timecourses estimated in step 1 of DR. Here we demonstrate that our framework can be used to present a maturational analogue to the netmats approach. In other words, to provide a whole-brain view on the functional architecture of the in-utero brain via a description of maturational relationships between matnets, which we call "maturational connectome" for conciseness. As shown in Figure 5A, this involves regression of the maturational networks against columns of the thresholded maturational dense connectome in order to obtain a N components by N voxels matrix of estimated regression coefficients (an analogue of component timecourses of DR1), followed by calculation of N components by N components matrix, in which elements represent correlations between each pair of rows in the regression coefficient matrix.

A three-dimensional embedding of the maturational connectome (Figure 5B), allows one to appreciate a generic structure of the maturational relationships. Here a point in space indicates a relative location of a network with respect to other networks, with a shorter distance between networks being indicative of stronger maturational ties (see below for a concrete definition of the latter). 5 groups of networks can be identified using hierarchical clustering, based on the networks' location in the embedded space (Figure 5C). The first group (coded brown) consisted of networks that combined the posterior and anterior peri-insular areas with occipital, auditory and ventral sensorimotor areas. The second group (coded green) consisted of two smaller sub-groups: one comprising dorsolateral pre-motor, dorsolateral prefrontal and medial pre- and supplementary motor areas; the other combining frontal anterior cingulate with inferior parietal and superior lateral occipital cortices, extending into medial posterior areas (precuneus). Adjacent to this group, there was a two-network group (coded blue), comprising dorsal sensorimotor areas. The fourth group (coded violet) comprised ventral frontal and orbitofrontal areas. Finally, the last group (coded purple) combined ventral occipito-temporal areas with dorsal parietal and sensorimotor areas.

311
312   *Figure 5. Maturational connectome. (A) Pipeline for derivation and analysis of maturational connectome (B) Maturational*
313   *connectome embedding and their split into groups, based on hierarchical clustering (C) Hierarchical clustering tree.*
314

We note a possibility of a dual view on the above results, assisting their interpretation. In the
above, the results of the regression coefficient matrix were treated as pseudo-timeseries, in
analogy to DR1 outputs. An alternative view is made possible by the fact that one dimension
of the estimated regression coefficient matrix is equal to the number of voxels and therefore
it can also be viewed as a collection of spatial maps, one per each matnet. These -
complementary – maps have a relatively straightforward interpretation: they depict areas to
which corresponding matnets tends to develop connections to in an age-related manner.
From this perspective, the clustering into 5 groups is determined by a spatial similarity of
their complementary maps. This fact permits a straightforward identification of
"maturational hubs" for each group as maps that characterise shared connectivity profiles
within each group of matnets, for instance, by means of principle component analysis.
326
Figure 6A shows the first principal component maps of complementary maps in each groups.
Thus, one can observe a likely vascular contribution in group 1 and 2, evident by the fact that
the maps contain areas overlapping with the circle of Willis. In parallel, both maps also
contain brain areas which are spatially distinct from the areas reflecting vascular
development. For group 1, these areas are bilateral dorsal somatosensory and adjacent
parietal cortices and bilateral cerebellum. For the group 2, prefrontal group, the hubs are
located in bilateral IFG and superior bank of anterior STG, bilateral insula, bilateral STS. For
the sensorimotor (3rd) group, the hubs are not expressed well but some preferential
connectivity to right insula and (predominantly right) striatum and thalamus can be
observed. For the 4th (ventral frontal and orbitofrontal) group, the hubs were located in the
bilateral SFG and MFG. Finally, for group 5 (ventral visual stream areas), the hubs were
located in (predominantly right) lateral parietal and dorsal parieto-occipital cortices and
right posterior perisylvian cortices.
340
Furthermore, as a proof-of-principle that maturational relationships are determined by an
age-related *increase* of connectivity and as well as a demonstration of the potential
application of the method to the study individual variability, two results can be presented.
One is derived by repeating the above regression procedure for each subject's dense
connectome, thereby obtaining subject-specific complementary maps (analogous to DR2
outputs), which were subsequently fitted across subject with age as covariate
(Supplementary Figure 11). The other is derived by estimating the temporal coupling
between each group of maturational networks and a corresponding maturational hub as a
function of age. Figure 6B shows the results for two maturational groups, whereas age-
related trajectories for all 25 components are shown in Supplementary Figure 12.

Figure 6. Matnets and their complementary maps. A. Spatial maps of matnets (top), their complementary maps (bottom), and the 1st PC of the complementary maps (right) in each matnet group. B. Examples of temporal correlations between time courses of matnets and their complementary map for two matnet groups (see Supplementary Figure 12 for all maturational components). Both matnets and complementary maps were thresholded at z >5 in order to reduce a degree of potential spatial overlap between the two. The time courses were computed as weighted averages of the above-threshold voxels.

DISCUSSION

In this paper, we present a description of emerging functional network architecture in in-utero brain with an unprecedented anatomical specificity, providing insights on the organisational principles of the fetal functional brain. This was made possible by developing a novel modeling framework that characterizes functional networks as an emerging property of the brain. Within this framework, the fusion of voxels into a network is determined by the similarity of their maturational profiles with respect to the rest of the brain. In effect, this

represents a computational implementation of Flechsig's principle (*19*) that states that
concordant maturation characterises functionally related areas. In an implicit form,
Flechsig's principle has been previously utilized in the studies of structural covariance (*29*) in
developmental cohorts (*30, 31*), including fetuses (*32*). Here we apply the principle explicitly
to the study of brain's functional organisation.

We have showed maturational networks (Figure 4) unveil spatially distributed patterns of
connections with a remarkable anatomical specificity for the in-utero data, owing to their
reliance on the benign signal properties that reveal an age-dependent increase of mid- and
long-distance connectivity in a spatially selective manner. We have also showed that
maturational networks represent a coherent way of characterising maturational patterns in
the context of fetal fMRI, compared to inference using the standard approach (Figure 3), in
which results appear to be affected by specific biases (we will discuss this below).

Taken together, the results indicate that fundamental features of neonatal and even adult-
like functional architecture occur prior to the exposure to extrauterine environmental
influences. This is reflected in a range of motifs characteristic of the neonatal brain
connectivity, which can be viewed as the eventual target for maturational processes in
utero. Thus, several networks revealed a non-negligible bilateral component, that agrees
with the studies of pre-term and term born babies (*4-6*), as well as in-utero seed-based
connectivity fMRI studies (*8*), suggesting that interhemispheric coupling becomes
established during this period. The maturational networks also characterised a range of non-
trivial functional relationships that are similarly observed in neonatal data (*6*), such as
functional associations between the inferior parietal regions and precuneus; between the
anterior cingulate cortex and lateral orbito-frontal cortex, between the medial and lateral
(pre)-motor cortices; between the central sulcus and posterior insular cortex; or between
the dorsal and ventral stream regions. This demonstrates that these emerging functional
relationships across spatially distinct regions are an intrinsic property of the brain and
provides crucial validation of the findings of neonatal studies where the complementary role
of environmental influences had been unclear.

An additional level of insight into the developmental sequelae of the fetal functional brain is
provided by a low-dimensional embedding of the whole-brain maturational connectome,
allowing an appreciation of mutual relationships between networks (Figure 5). A
conspicuous generic feature of this connectome is the tendency for homologous
contralateral networks to cluster together. Overall, the clustering analysis identifies two
larger groups that occupy the central location in the embedded sapce and three smaller,
more peripheral, groups. Based on the areas that dominate their anatomical layout, the
three smaller clusters of networks can be labelled as orbitofrontal, ventral visual and
sensorimotor groups. Of the larger groups, one was dominated by the cortical nodes of
perception and bodily sensation (occipital, auditory and somatosensory limbic areas) but
also included nodes in the motor and motor limbic (*33*) (anterior cingulate and anterior
insular) cortices. The other larger group was dominated by the functional nodes responsible
for an environmental interaction through action (dorsolateral and medial pre-motor cortex
and pre-frontal areas), but also included a sub-group of networks which spatially overlap
with nodes of the future default mode networks (*25, 34*), such as precuneus, anterior
cingulate and angular gyrus. Notably, the location of the latter within the embedding space

was midway between the notional perception group and the remaining networks of the notional action group, hinting both towards hub connectivity patterns and their apparent role in modulating internal and external inputs whilst mind-wandering or performing cognitively demanding tasks later in life (*35*).

The framework also allowed us to describe maturational hubs of the in-utero connectome, characterised as regions in which networks within a matnet group form preferential connectivity in an age-related manner. Areas in the dorsal somatosensory and adjacent parietal cortex which process sensation and spatial information; as well as the cerebellum were identified as hubs for the first group of matnets, combining perception, bodily sensation and motor limbic areas located outside sensorimotor cortices, suggesting integration of information across different perceptual and limbic domains towards a central cortical processing unit. The analysis also reveals an important role of associative areas implicated in high-level cognition within the brain connectome from the onset of the brain functional development. Thus, the hub for ventral occipital and temporal areas, which in the adult brain encodes representations of abstract visual information, is found in the posterior parietal cortex, including the right IPS and the posterior node of ventral attentional network (VAN) (*36*). This supports evidence, similarly observed at the level of individual matnets (e.g., matnet #6), of an ongoing integration of the ventral and dorsal stream representations. We also observed an emergence of links between ventral action-related limbic areas, representing internal motives and drives in the adult brain (*37*), and areas associated with encoding representations of abstract rules for goal-directed behavior and with executive control. This is made evident by the fact that the major hubs for the action group, which among others included networks in lateral prefrontal cortex, were located in the anterior perisylvian and insular cortices, posterior ventral orbitofrontal and anterior temporal cortices, overlapping with limbic and motor limbic cortices and the prefrontal hub of the adult VAN. Reciprocally, the dorsal prefrontal areas implemented in the adult dorsal attentional network (including frontal eye field), working memory and executive control were identified as a hub for ventral orbitofrontal matnet group, which in the adult brain are known to project feedback pathways to the dorsolateral prefrontal cortex, providing the latter with information on internal environment (*38*). These findings indicate that the neural machinery for linking decisions and actions to internal wishes and motives start emerging as early as the fetal period in human life.

These results challenge the view that the transition from fetal to a more mature functional architecture is manifested by the shift of functional hubs from primary to associative areas (*39, 40*) and aligns with earlier studies of structural connectivity in full-term and preterm neonates showing that adult-like features of the structural connectome can already be observed at this early period (*41*). The distinction between more mature and in-utero functional connectome features appear to be signified by a relative disconnection along posterior-anterior axis, such as between nodes of fronto-parietal networks which are considered responsible for the integration of information across behavioral domains in the adult brain. Similarly, consistent with previous findings in preterm neonates (*4*), no strong evidence of the links between medial posterior and anterior nodes of the default mode network were observed, which at this period appear to be integrated, respectively, within parietal and frontal networks and lack a prominent role in the observed whole brain functional architecture.

Maturation entails competing physiological processes which may potentially leave a footprint on the properties of the fMRI signal (*42, 43*), thereby raising a question about the biological underpinnings of the age-related signal changes implicated in the derivation of maturational networks. For instance, one cannot exclude the possibility that changes in the long-distance connectivity, in the absence of a mature structural connectome, are in part due to the coordinated development of the brain's vasculature (*44*). De-confounding the latter from the estimates of neural connectivity is a contentious issue even in the context of adult resting-state imaging (*45, 46*). In the fetal brain, the problem may be further exacerbated as the development of brain neural systems goes hand in hand with the development of other organ functions including the vasculature and thus are likely collinear to the degree that the two are indistinguishable at a level visible to fMRI.

Compared to maturational networks, group-ICA components identified with a standard group-ICA approach had diminished spatial complexity and anatomical specificity and were biased towards the white matter. Notably, the results of dual regression modeling showed that local connectivity within group-ICA networks diminishes with age. Such characteristics fit well those of the functional nodes described in the fetal animal studies, which center on the cortical subplate and act as local amplifiers of the thalamic activity with spread that does not conform to anatomical boundaries (*3, 47*). This may suggest that group-ICA and maturational networks truthfully reflect two different states of the fetal functional brain: a truly "fetal" subplate-centered (*48*) and locally active state depicted by the group-ICA, that gives way to the adult-like cortex-centered and spatially distributed state of maturational networks. Against this intriguing interpretation, though not necessary incompatible with it, are the results of the univariate analysis of the connectivity metrics. The latter demonstrates that the correlational structure of the data, that underlies the derivation of the group-ICA components, is dominated by a spatially smooth and non-linear distance-dependent gradient, which scales negatively with age. The factors that make biologically-motivated interpretation of this gradient unlikely is the spatially indiscriminate character of these phenomena combined with a violation of anatomical boundaries, including the large connectivity distance between the two brain hemispheres which in reality are separated by a CSF filled inter-hemispheric fissure.

An initial hypothesis to explain the origins of distance-dependent gradients and its interaction with age can be based on the potential contribution of two factors: motion and effective resolution. The role of motion on connectivity estimates has been demonstrated in adult imaging, where it has been shown to decrease long-range connectivity and overestimate local connectivity (*17, 18*). Although we used a comprehensive image processing pipeline to account for head motion during data acquisition, fetal imaging data is still especially susceptible to this effect as fetuses have virtually no motion-free periods. Even if the fetus stays still, maternal breathing cycles and endogenous motion in the non-rigid tissues surrounding the fetal head continue to cause a constant change of position. Under these circumstances, effective resolution naturally leads to age-related differences in the effect, which likely explains the dual regression result showing a decrease in connectivity with age within the most representative component voxels. The brain undergoes a 3-fold growth in size over the studied period, which implies that real-world separation between pairs of voxels in a standard space is smaller for younger subjects than for older ones and

514   thus a greater effect of distance as measured in the common space. In light of the
515   differences in signal properties between the grey and white matter and their modulation by
516   age, the possible contribution of other factors such as modulation of the BOLD signal itself
517   and/or the role of age-related changes in tissue content also should not be disregarded.
518
Below we outline several limitations of the study. First, the current study has the well-known
limitations of cross-sectional analyses whereby between-subject variability can be
confounded with aging effects. Nevertheless, cross-sectional data are expected to dominate
fetal research for a foreseeable future, as scanning mothers during pregnancy on multiple
occasions presents both ethical and practical challenges. In the meantime, one can strive for
better estimates of cross-sectional trajectories, using improved modelling and larger data
samples. Our results are based on one of the largest fetal fMRI data sets both in terms of the
number of subjects and the number of volumes per subject. However, further improvements
in modelling can be achieved when data for the full fetal dHCP cohort will be made openly
available to the neuroscientific community in the coming year. This would increase the
current data sample by a factor of nearly 2.
530
The second limitation concerns generalization of our conclusions to other data samples,
especially in the context of fetal fMRI as a novel field, where norms of data acquisition are
yet to be established. Unfortunately, fetal fMRI has not as yet stepped in into the age of
normative open-access big data (*49*) which has enabled recent progress in the study of ex-
utero connectivity, (e.g.,(*50*)). However, the qualitative comparison of our results with the
results drawn from other studies gives us a certain confidence that our results are not
specific to our sample. For instance, there was a remarkable similarity between our group-
ICA results and the group-ICA results reported in a recent paper (*11*), despite considerable
differences in the acquisition sequence (multi- vs single-band), spatial image corrections
(dynamic distortion and slice-to-volume corrections vs volumetric alignment only) and de-
noising pipelines (predominantly motion parameter-based vs. ICA-based). Furthermore, the
qualitative characteristics of group-ICA components as well as the dominance of distance-
dependent gradient over the correlational structure also appear to be reproducible across
the studies (*11*).
545
In conclusion, we describe a novel framework that delineates the emergence of resting state
networks in the fetal human brain with remarkable spatial specificity and provides a
comprehensive model of inter-areal maturational relationships, assigning a central role to
the brain regions associated with active environmental interaction through perceptual and
motor-planning mechanisms. A discerning feature of this maturational network framework is
a prospective incorporation of the variable-of-interest (here, age) into network estimation.
This can potentially make the method adaptable to other applications, such as studying early
human development through childhood, network maturation in neurodevelopmental
disorders such as autism, ageing and exploring the connectivity underpinnings of changing
patterns of behavior across the lifespan.
556
557
558   **METHODS**
559
560   *Data*

Resting-state fMRI data were acquired in 151 fetuses (62 females, 77 males, 5 unknown),
median age = 29.5w, range = [25 38], with Philips Achieva 3T system (Best, NL) and a 32-
channel cardiac coil using a single-shot EPI (TR/TE = 2200/60) sequence consisting of 350
volumes of 48 slices each, slice grid 144 x 144, isotropic resolution = 2.2 mm, multi-band
(MB) factor = 3 and SENSE factor = 1.4 (*27*). All fetal brain images were reported by a
neuroradiologist as showing appropriate appearances for their gestational age with no
acquired lesions or congenital malformations of clinical significance. Data from 7 fetuses did
not pass visual quality assessment due to excessive motion and failure in image
reconstruction.

The data of the remaining 144 fetuses were preprocessed using a dedicated pipeline (*22-24*).
In brief, the data underwent MB-SENSE image reconstruction, dynamic shot-by-shot B0 field
correction by phase unwrapping and slice-to-volume (S2V) motion correction (*22*). The data
were then temporally denoised using several sets of confound regressors, aiming to address
various types of artefacts. The denoising model combined volume censoring regressors,
aiming to reject volumes (at a heuristically selected threshold) (Supplementary Figure 13),
highpass (1/150 Hz) filtering regressors of direct cosine transform matrix in order to remove
slow frequency drift in the data, 6 white matter and cerebrospinal fluid component
timecourses (obtained using subject-level ICA within a combined white matter + CSF mask,
(e.g.,(*51*), and 3 variants of voxelwise 4d denoising maps in order to account for the local
artefacts in the data: 1) folding maps (N=2) which aggregate time courses of voxels linked in
multiband acquisition to voxels in the original data, aiming at filtering out leakage artefacts;
2) density maps, representing temporal evolution of an operator that compensates for the
volume alterations a result of distortion in phase encoding direction, and aiming to filter out
residual effects of distortion correction on the voxel timecourses; and 3) motion-parameter-
based regressors, expanded to include first and second order volume-to-volume and slice-
to-slice differentials as well as their square terms, aiming to remove motion-related artefacts
(*52, 53*).

Registration to the group space

A 4D atlas of the developing brain (available at https://brain-development.org/brain-
atlases/fetal-brain-atlases/)(*54*) was used as a template space for data registration. A
schematic depiction of the registration to a common template space is shown in
Supplementary Figure 14. The mapping between a functional native space and the common
template space is constructed as the concatenation of several intermediate transformations,
which ascertain a gradual alignment between spaces to minimise risks of gross misalignment
as a result of the substantial differences in the brain topology across the range of gestation
ages (*6*): 1) rigid alignment between mean functional and anatomical scans calculated using
FLIRT (*55*); 2) a non-linear transformation between an anatomical T2 scan and an age-
matched template calculated using ANTs (*56*) ; 3) a sequence of non-linear transformations
between templates of adjacent ages (e.g., 24 and 25, 25 and 26, etc.), also calculated using
ANTs. These transformations were concatenated to create a one-step mapping between
functional and group template space, avoiding multiple interpolations. The template
corresponding to GA=37 weeks was selected as a common space for group analysis based on
the considerations that it has a greatest effective resolution and topological complexity. An

608 additional group space was created by symmetrizing the GA=37 week template with respect
609 to the brain midline, with appropriate adjustment of the mapping from the functional native
610 spaces, that included an additional non-linear transform from non-symmetrical-to-
611 symmetrical template spaces. After registering the functional MRI data to the template
612 space, they were smoothed using 3mm Gaussian kernel.
613
614 ***Data analysis***
615
Univariate analyses. The seeds for the seed-to-brain analysis were determined empirically
using the results of modelling age-related changes in interhemispheric connectivity between
pairs of homologous voxels (Supplementary Figure 15), performed in the symmetrical
template space (*27*). The subject-specific maps of homologous voxel connectivity were
obtained by calculating the correlation between timecourses of homologous voxels in the
two hemispheres. The age-effect map was obtained via a voxel-wise regression with age as a
covariate. The seeds for grey matter were created by thresholding the age-effect map from
the above analysis at z > 3, which rendered 3 sizable clusters of voxels (14, 32, and 45
voxels). Given the absence of positive age-related increase in connectivity between
homologous voxels for white matter areas, the white matter seeds were created by
thresholding the age-effect map of interhemispheric connectivity negatively at z < -3, and
then manually adjusting clusters to fit the size of the grey matter clusters. Because the seeds
were defined in the symmetrical template space, the seed-to-brain connectivity analysis was
also performed in this space. The seed-to-brain group-average correlation map was
calculated by first calculating individual maps of correlations between time course of a seed
and time courses of all voxels in the brain and then averaging these maps across subjects.
The age-effect map was obtained by fitting individual maps voxelwise using age as a
covariate.
634
635 ***Group-ICA***. The derivation of group-average modes-of-variation and their subject-specific
636 variants was performed using the protocol of FSL MELODIC for group-ICA analyses(*15*),
637 including FSL MELODIC's Incremental Group Principal component analysis (MIGP step (*12*)),
638 and the standard procedure of dual regression, implemented in FSL (*57*). The number of
639 derived components was set to 25.
640
Maturational modes of variation. The pipeline for derivation of maturational modes of
variation is shown in Figure 1B. First, a symmetrical matrix of correlations between each pair
of voxels in the brain mask was calculated, aka "dense connectome", for each subject
separately. Each element of the dense connectome was fitted across subjects with age as
covariate, rendering a voxel-by-voxel matrix of age-effect beta coefficients. The matrix was
then converted into t-values, rendering maturational dense connectome, subsequently
thresholded at 0 in order to leverage the age-dependent increases in correlations in network
estimation. The rationale for positive thresholding is described in the Results section. In
order to perform connectome factorisation, an intermediate step of dimensionality
reduction, analogous to MIGP, was applied. For this, the maturational dense connectome
(size: N voxels by N voxels) was split column-wise into 200 blocks (size: N voxels by N
voxels/200. At the initial step, a matrix consisting of the first two blocks was formed and
subsequently reduced to 500 components using singular value decomposition. An iterative
procedure was then run that consisted of concatenating the current matrix of 500

components with a following block and subsequent reduction to 500 components by SVD, until all blocks were exhausted. The output of this procedure was used to obtain the final factorisation of 25 components using FSL MELODIC.

Maturational connectome analysis. The pipeline for derivation of the maturational connectome is shown in Figure 5A. It consists of the regression of the maturational networks against the maturational dense connectome in order to obtain #networks by #voxels matrix of regression coefficients. Correlations between each pair of rows of the matrix were then estimated, collected into a matrix which constitutes the maturational connectome. In order to reveal a structure of the whole-brain maturational relationships, the maturational connectome matrix was embedded into 3-dimensional space using an eigendecomposition of a graph normalised Laplacian. A point in the embedding space indicates a relative location of a network with respect to other networks (i.e., a shorter distance means closer maturational ties). A partition of networks into groups of networks was performed using the Ward method of hierarchical clustering (*58*), based on the network coordinates in the embedding 3D space.

Funding

The Developing Human Connectome Project was funded by the European Research Council under the European Union Seventh Framework Programme (FP/20072013)/ERC Grant Agreement no. 319456. The authors also acknowledge support in part from the Wellcome Engineering and Physical Sciences Research Council (EPSRC) Centre for Medical Engineering at Kings College London [WT 203148/Z/16/Z], the Medical Research Council (MRC) Centre for Neurodevelopmental Disorders [MR/N026063/1], and the Department of Health through an NIHR Comprehensive Biomedical Research Centre Award (to Guy's and St. Thomas' National Health Service (NHS) Foundation Trust in partnership with King's College London and King's College Hospital NHS Foundation Trust). VK and TA were supported by a MRC Clinician Scientist Fellowship [MR/P008712/1] and MRC translation support award [MR/V036874/1]. JOM is supported by a Sir Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal Society [Grant Number 206675/Z/17/Z]. LCG is supported by the Comunidad de Madrid-Spain Support for R&D Projects [BGP18/00178]. AEF acknowledges additional funding from the UKRI CDT in Artificial Intelligence for Healthcare in his role as a Senior Teaching Fellow [grant number EP/S023283/1].

REFERENCES:

1.   I. Kant, *Critique of Pure Reason*. (Penguin Classics, 2007), pp. 784.
2.   F. Nietzsche, *The Birth of Tragedy*. (Penguin Classics, 1993), pp. 160.
3.   R. Khazipov, H. J. Luhmann, Early patterns of electrical activity in the developing cerebral cortex of humans and rodents. *Trends Neurosci* **29**, 414-418 (2006).
4.   V. Doria *et al.*, Emergence of resting state networks in the preterm human brain. *Proc Natl Acad Sci U S A* **107**, 20015-20020 (2010).
5.   M. Eyre *et al.*, The Developing Human Connectome Project: typical and disrupted perinatal functional connectivity. *Brain* **144**, 2199-2213 (2021).

6.   S. P. Fitzgibbon *et al.*, The developing Human Connectome Project (dHCP) automated
resting-state functional processing framework for newborn infants. *Neuroimage* **223**,
117303 (2020).
7.   S. J. Gould, *Ontogeny and Phylogeny*. (Belknap Press of Harvard University Press,
1985), pp. 518.
8.   M. E. Thomason *et al.*, Cross-hemispheric functional connectivity in the human fetal
brain. *Sci Transl Med* **5**, 173ra124 (2013).
9.   M. E. Thomason *et al.*, Age-related increases in long-range connectivity in fetal
functional neural connectivity networks in utero. *Dev Cogn Neurosci* **11**, 96-104
(2015).
10. G. Ferrazzi *et al.*, Resting State fMRI in the moving fetus: A robust framework for
motion, bias field and spin history correction. *Neuroimage* **101**, 555-568 (2014).
11. L. Ji, C. L. Hendrix, M. E. Thomason, Empirical evaluation of human fetal fMRI
preprocessing steps. *Network Neuroscience*, 1-37 (2022).
12. S. M. Smith, A. Hyvarinen, G. Varoquaux, K. L. Miller, C. F. Beckmann, Group-PCA for
very large fMRI datasets. *Neuroimage* **101**, 738-749 (2014).
13. V. D. Calhoun, T. Adali, G. D. Pearlson, J. J. Pekar, A method for making group
inferences from functional MRI data using independent component analysis. *Hum
Brain Mapp* **14**, 140-151 (2001).
14. C. F. Beckmann, C. E. Mackay, S. M. Smith, Group comparison of resting-state FMRI
data using multi-subject ICA and dual regression. *NeuroImage* **47**, S148 (2009).
15. C. F. Beckmann, S. M. Smith, Probabilistic independent component analysis for
functional magnetic resonance imaging. *IEEE Trans Med Imaging* **23**, 137-152 (2004).
16. J. D. Bijsterbosch *et al.*, The relationship between spatial configuration and functional
connectivity of brain regions. *Elife* **7**, (2018).
17. J. D. Power, K. A. Barnes, A. Z. Snyder, B. L. Schlaggar, S. E. Petersen, Spurious but
systematic correlations in functional connectivity MRI networks arise from subject
motion. *Neuroimage* **59**, 2142-2154 (2012).
18. K. R. A. Van Dijk, M. R. Sabuncu, R. L. Buckner, The influence of head motion on
intrinsic functional connectivity MRI. *Neuroimage* **59**, 431-438 (2012).
19. P. Flechsig, Developmental (myelogenetic) localisation of the cerebral cortex in the
human subject. *Lancet* **158**, 1027-1030 (1901).
20. A. D. Edwards *et al.*, The Developing Human Connectome Project Neonatal Data
Release. *Frontiers in Neuroscience* **16**, (2022).
21. http://www.developingconnectome.org.
22. L. Cordero Grande, A. N. Price, D. Christiaens, J. Hutter, J. V. Hajnal, in *Proceedings of
the 26th annual meeting of the ISMRM*. (Paris, France, 2018), pp. 208.
23. L. Cordero-Grande, E. J. Hughes, J. Hutter, A. N. Price, J. V. Hajnal, Three-dimensional
motion corrected sensitivity encoding reconstruction for multi-shot multi-slice MRI:
Application to neonatal brain imaging. *Magn Reson Med* **79**, 1365-1376 (2018).
24. V. Karolis *et al.*, in *Organisation for Human Brain Mapping*. (2021), pp. 2112.
25. S. M. Smith *et al.*, Correspondence of the brain's functional architecture during
activation and rest. *Proc Natl Acad Sci U S A* **106**, 13040-13045 (2009).
26. J. S. Damoiseaux *et al.*, Consistent resting-state networks across healthy subjects.
Proc Natl Acad Sci U S A **103**, 13848-13853 (2006).
27. A. N. Price *et al.*, in *Proceedings of the annual meeting of the International Society of
Magnetic Resonance in Medicine (ISMRM)*. (2019), pp. 244.

28.   S. M. Smith *et al.*, A positive-negative mode of population covariation links brain
connectivity, demographics and behavior. *Nat Neurosci* **18**, 1565-1567 (2015).

29.   A. Mechelli, K. J. Friston, R. S. Frackowiak, C. J. Price, Structural covariance in the
human cortex. *J Neurosci* **25**, 8303-8310 (2005).

30.   D. Fenchel *et al.*, Development of Microstructural and Morphological Cortical Profiles
in the Neonatal Brain. *Cereb Cortex* **30**, 5767-5779 (2020).

31.   V. R. Karolis *et al.*, Volumetric grey matter alterations in adolescents and adults born
very preterm suggest accelerated brain maturation. *Neuroimage* **163**, 379-389
(2017).

32.   J. Xia *et al.*, Fetal cortical surface atlas parcellation based on growth patterns. *Hum*
Brain Mapp **40**, 3881-3899 (2019).

33.   A. D. Craig, How do you feel? Interoception: the sense of the physiological condition
of the body. *Nat Rev Neurosci* **3**, 655-666 (2002).

34.   M. E. Raichle *et al.*, A default mode of brain function. *P Natl Acad Sci USA* **98**, 676-682
(2001).

35.   M. E. Raichle, The Brain's Default Mode Network. *Annu Rev Neurosci* **38**, 433-447
(2015).

36.   M. Corbetta, G. L. Shulman, Control of goal-directed and stimulus-driven attention in
the brain. *Nat Rev Neurosci* **3**, 201-215 (2002).

37.   H. Barbas, General cortical and special prefrontal connections: principles from
structure to function. *Annu Rev Neurosci* **38**, 269-289 (2015).

38.   H. Barbas, M. A. Garcia-Cabezas, How the prefrontal executive got its stripes. *Curr*
Opin Neurobiol **40**, 125-134 (2016).

39.   S. Oldham, A. Fornito, The development of brain network hubs. *Dev Cogn Neurosci*
36, 100607 (2019).

40.   M. I. van den Heuvel *et al.*, Hubs in the human fetal brain network. *Dev Cogn*
Neurosci **30**, 108-115 (2018).

41.   G. Ball *et al.*, Rich-club organization of the newborn human brain. *Proc Natl Acad Sci*
U S A **111**, 7456-7461 (2014).

42.   J. J. Harris, C. Reynell, D. Attwell, The physiology of developmental changes in BOLD
functional imaging signals. *Dev Cogn Neuros-Neth* **1**, 199-216 (2011).

43.   M. Pietsch *et al.*, A framework for multi-component analysis of diffusion MRI data
over the neonatal period. *Neuroimage* **186**, 321-337 (2019).

44.   B. Lacoste, C. H. Gu, Control of cerebrovascular patterning by neural activity during
postnatal development. *Mech Develop* **138**, 43-49 (2015).

45.   M. G. Bright, J. R. Whittaker, I. D. Driver, K. Murphy, Vascular physiology drives
functional brain networks. *Neuroimage* **217**, (2020).

46.   K. Murphy, R. M. Birn, P. A. Bandettini, Resting-state fMRI confounds and cleanup.
Neuroimage **80**, 349-359 (2013).

47.   M. T. Colonnese, M. A. Phillips, Thalamocortical function in developing sensory
circuits. *Curr Opin Neurobiol* **52**, 72-79 (2018).

48.   I. Kostovic, The enigmatic fetal subplate compartment forms an early tangential
cortical nexus and provides the framework for construction of cortical connectivity.
Prog Neurobiol **194**, 101883 (2020).

49.   V. Rajagopalan, S. Deoni, A. Panigrahy, M. E. Thomason, Is fetal MRI ready for
neuroimaging prime time? An examination of progress and remaining areas for
development. *Dev Cogn Neuros-Neth* **51**, 100999 (2021).

50. S. M. Smith *et al.*, Resting-state fMRI in the Human Connectome Project. *Neuroimage*
80, 144-168 (2013).
51. Y. Behzadi, K. Restom, J. Liau, T. T. Liu, A component based noise correction method
(CompCor) for BOLD and perfusion based fMRI. *Neuroimage* **37**, 90-101 (2007).
52. R. H. R. Pruim, M. Mennes, J. K. Buitelaar, C. F. Beckmann, Evaluation of ICA-AROMA
and alternative strategies for motion artifact removal in resting state fMRI.
Neuroimage **112**, 278-287 (2015).
53. R. H. R. Pruim *et al.*, ICA-AROMA: A robust ICA-based strategy for removing motion
artifacts from fMRI data. *Neuroimage* **112**, 267-277 (2015).
54. A. Serag *et al.*, Construction of a consistent high-definition spatio-temporal atlas of
the developing brain using adaptive kernel regression. *Neuroimage* **59**, 2255-2265
(2012).
55. M. Jenkinson, S. Smith, A global optimisation method for robust affine registration of
brain images. *Med Image Anal* **5**, 143-156 (2001).
56. B. B. Avants *et al.*, A reproducible evaluation of ANTs similarity metric performance in
brain image registration. *Neuroimage* **54**, 2033-2044 (2011).
57. S. M. Smith *et al.*, Advances in functional and structural MR image analysis and
implementation as FSL. *Neuroimage* **23**, S208-S219 (2004).
58. J. H. Ward, Hierarchical Grouping to Optimize an Objective Function. *Journal of the*
American Statistical Association **58**, 236-244 (1963).
815
816