Structural basis for the unique multifaceted interaction of DPPA3 with the UHRF1 PHD

finger

Keiichi Hata[1,#], Naohiro Kobayashi[2,#], Keita Sugimura[3], Weihua Qin[4], Deis Haxholli[4], Yoshie

Chiba[3], Sae Yoshimi[1], Gosuke Hayashi[5], Hiroki Onoda[1], Takahisa Ikegami[6], Christopher B.

Mulholland[4], Atsuya Nishiyama[3], Makoto Nakanishi[3], Heinrich Leonhardt[4], Tsuyoshi

Konuma[6,*], Kyohei Arita[1,*]

[1]Structural Biology Laboratory and [6]Structural Epigenetics Laboratory, Graduate School of

Medical Life Science, Yokohama City University, 1-7-29, Suehiro-cho, Tsurumi-ku, Yokohama,

Kanagawa 230-0045, Japan.

[2]RIKEN Center for Biosystems Dynamics Research, Suehiro-cho 1-7-22, Tsurumi, Yokohama,

Kanagawa 230-0045, Japan.

[3]Division of Cancer Cell Biology, The Institute of Medical Science, The University of Tokyo,

4-6-1 Shirokanedai, Minato-ku, Tokyo 108-8639, Japan.

[4]Faculty of Biology, Ludwig-Maximilians-Universität München, Munich, Germany.

[5]Department of Biomolecular Engineering, Nagoya University, Furo-cho, Chikusa-ku, Nagoya

464-8603, Japan.

* To whom correspondence should be addressed. Tel: +81 45 508 7225; Fax: +81 45 508 7365;

Email: aritak@yokohama-cu.ac.jp

Correspondence may also be addressed to Tsuyoshi Konuma. Tel: +81 45 508 7218; Fax: +81

45 508 7362; Email: konumax@yokohama-cu.ac.jp

[#] The authors wish it to be known that, in their opinion, the first two authors should be regarded

as Joint First Authors.

Present Address: Hiroki Onoda, Synchrotron Radiation Research Center, Nagoya University,

Furo-cho, Chikusa-ku, Nagoya, 464-8603, JAPAN.

ABSTRACT

Ubiquitin-like with PHD and RING finger domain-containing protein 1 (UHRF1)-dependent DNA methylation is essential for maintaining cell fate during cell proliferation. Developmental pluripotency-associated 3 (DPPA3) is an intrinsically disordered protein that specifically interacts with UHRF1 and promotes passive DNA demethylation by inhibiting UHRF1 chromatin localization. However, the molecular basis of how DPPA3 interacts with and inhibits UHRF1 remains unclear. We aimed to determine the structure of the mouse UHRF1 plant homeodomain (PHD) complexed with DPPA3 using nuclear magnetic resonance. Induced α-helices in DPPA3 upon binding of UHRF1 PHD contribute to stable complex formation with multifaceted interactions, unlike canonical ligand proteins of the PHD domain. Mutations in the binding interface and unfolding of the DPPA3 helical structure inhibited binding to UHRF1 and its chromatin localization. Our results provide structural insights into the mechanism and specificity underlying the inhibition of UHRF1 by DPPA3.

INTRODUCTION

Cytosine DNA methylation of the CpG sequence in mammals plays a pivotal role in embryogenesis, retrotransposon silencing, X-chromosome inactivation, genome imprinting, and carcinogenesis(1). Mammalian cells undergo two waves of methylation changes: DNA methylation and demethylation(2). After fertilization, DNA methylation patterns derived from gametes are erased during early embryogenesis and re-established during cellular development(3). DNA methylation patterns in primordial germ cells (PGCs) are erased globally, and sex-specific DNA methylation patterns are established during germ cell development(2, 4). After establishment, cell-type-specific DNA methylation patterns are faithfully propagated after each cycle of replication to maintain cellular identity. DNMT1, a maintenance DNA methyltransferase, and UHRF1 (ubiquitin-like PHD and RING finger domain-containing protein 1, also known as Np95/ICBP90), which is a ubiquitin E3-ligase and recruiter of DNMT1, are essential for DNA methylation maintenance(5–8). UHRF1 specifically binds to hemi-methylated DNA and ubiquitinates histone H3 and PCNA-associated factor 15 (PAF15) to recruit DNMT1 to chromatin and replication sites(9–13). The two distinct ubiquitin signals are involved in replication-coupled and uncoupled DNA methylation maintenance(9, 14, 15).

UHRF1 functions as a reader of epigenetic marks, hemi-methylated DNA, and the H3K9me2/3 modification(16–19) to regulate its ubiquitination activity(20). Further, it serves as a binding platform for DNA replication factors and epigenetic modifiers, which are involved in DNA methylation maintenance, gene expression, DNA damage repair, and tumorigenesis(9, 21–28). Additionally, mouse DPPA3 (developmental pluripotency associated 3, also known as Stella/PGC7; hereafter mDPPA3) is a novel ligand that regulates the binding of UHRF1 to chromatin(29, 30). mDPPA3, an intrinsically disordered protein (IDP), is specifically expressed in PGCs, oocytes, and preimplantation embryos, and plays an important role in the formation of oocyte-specific DNA methylation patterns by preventing excessive *de novo* DNA methylation mediated by UHRF1(29, 31–33). Overexpression of mDPPA3 in somatic cells

such as NIH3T3, HEK293, and mouse embryonic stem cells, and in non-mammalian species,

results in genome-wide DNA demethylation, indicating that mDPPA3 is a passive DNA

demethylation factor that inhibits the cellular functions of UHRF1(29, 34, 35).

UHRF1 has five functional domains: a ubiquitin-like domain (UBL), tandem tudor

domain (TTD), plant homeodomain (PHD) finger, SET and RING associated domain (SRA),

and really interesting new gene (RING) (Figure 1A). mDPPA3 interacts with the UHRF1 PHD

finger, resulting in the inhibition of chromatin binding of UHRF1(35, 36). The UHRF1 PHD

finger also interacts with the N-terminal $^{1}ARTK^{4}$ in histone H3 (H3) and $^{1}VRTK^{4}$ in PAF15, in

which strict recognition of the main chain amino group at the first residue of H3 and PAF15 is

critical for their ubiquitination(9, 19). The PHD finger is one of the largest families of

chromatin-reader domains. They have been found in more than 100 human proteins, most of

which recognize K4 methylation state in the H3 N-terminal tail(37). Notably, with a few

exceptions, the recognition mode of the first amino acid residue amino group in ligands is

conserved among almost all PHD fingers(38). Indeed, acetylation of the N-terminus of the H3

tail abolished binding to the PHD finger of UHRF1(19). Given that the ARTK/VRTK-like

sequence is not present at the N-terminus of mDPPA3, the molecular mechanisms by which

the UHRF1 PHD finger recognizes DPPA3 is unknown.

Here, we aimed to determine the solution structure of mouse UHRF1 PHD (mPHD)

in complex with the C-terminal fragment of mDPPA3 using nuclear magnetic resonance

(NMR), and identified the unique multifaceted interaction of mDPPA3 with mPHD. Although

the $^{88}VRT^{90}$ cassette of mDPPA3 is not located at its N-terminus, unlike H3 or PAF15, we found

that it was recognized by a shallow acidic groove on the mPHD in a manner similar to the N-

terminus of H3 and PAF15. Structural induction of the two α-helices of mDPPA3 provided

several additional binding sites for mPHD, which plays an important role in stable complex

formation. Structure-guided mutagenesis and functional assays using *Xenopus* egg extracts and

mouse embryonic stem cells (mESCs) helped evaluate the key amino acid residues of mDPPA3

that negatively regulate the binding of mUHRF1 to chromatin, and shed light on the

mechanisms underlying chromatin delocalization of mUHRF1 by mDPPA3. Our data provide

insight into the diversity of recognition of ligand proteins by the PHD finger and contribute to

the understanding of its key role in epigenetic maintenance.

MATERIALS AND METHODS

Protein expression and purification

For NMR and ITC experiments, cDNA of mouse UHRF1 PHD (residues 304–372)

was sub-cloned into a pGEX6P-1 plasmid (Cytiva) at 5'-BamHI and 3'-XhoI sites for protein

expression with N-terminal glutathione S-transferase (GST). The protein was expressed in

E.coli BL21 (DE3) in Luria–Bertani medium (LB) containing 50 µg/mL ampicillin. When the

optical density at 660 nm (O.D.$_{660}$) of the cells reached 0.7, 0.2 mM isopropyl β-d-

thiogalactoside (IPTG) was added to the medium and the cells were further harvested for 3 h

at 37°C. The cells were suspended with lysis buffer (40 mM Tris-HCl [pH 8.0], 300 mM NaCl,

10% glycerol, 30 µM Zn-acetate, 0.5 mM tris(2-carboxyethyl)phosphine (TCEP)) and

disrupted by sonication. After the cell debris were removed by centrifugation, the supernatant

was loaded onto a GST-affinity column, GS4B (Cytive). After the protein was eluted from the

column using reduced glutathione, the GST-tag was cleaved by HRV-3C protease. The sample

was further purified using HiTrap Q HP anion-exchange chromatography (Cytiva). Final

purification was performed using HiLoad 26/600 Superdex 75 size-exclusion chromatography

(Cytiva) equilibrated with buffer 1×phosphate-buffered saline (adjusted to pH 7.0; PBS)

containing 1 mM DTT.

mDPPA3, residues 76–128W, for NMR and ITC experiments was expressed as a six

histidine-tagged ubiquitin fusion protein. The procedures for cell culture were the same as that

for mPHD. The cells were suspended in lysis buffer (30 mM HEPES [pH7.5], 400 mM NaCl,

0.1% Nonidet P-40 (NP-40), 40 mM imidazole). After cell lysis by sonication and removal of

cell debris by centrifugation, the supernatant was loaded to histidine-tag affinity column Ni

Sepharose 6 Fast Flow (Cytiva), and the sample was eluted from the column using an elution

buffer containing 500 mM imidazole. Next, the histidine-tag was removed by *Saccharomyces*

cerevisiae ubiquitin carboxyl-terminal hydrolase YUH1. The sample was further purified using

HiTrap SP HP cation-exchange chromatography (Cytiva) and finally purified by HiLoad

26/600 Superdex 75 size-exclusion chromatography equilibrated with 1×PBS buffer containing

1 mM DTT.

For structure determination using NMR, cDNA of mDPPA3 (residues 76–127) was

inserted into the 3' end of the mPHD (residues 304–372) in pGEX6P-1 plasmid for expression

as fusion protein. The procedures for cell culture of the mPHD-mDPPA3 were same as for

mPHD. The cells were suspended with lysis buffer (40 mM Tris-HCl [pH 8.0], 300 mM NaCl,

10% glycerol, 30 μM Zn-acetate, 0.5 mM TCEP) and then disrupted by sonication. After the

cell debris was removed by centrifugation, the supernatant was loaded onto GS4B. The protein

was eluted from the column using reduced glutathione, and then GST-tag was cleaved by HRV-

3C protease. The sample was further purified using HiLoad 26/600 Superdex 75 size-exclusion

chromatography equilibrated with 1×PBS buffer containing 1 mM DTT.

For preparation of ^{15}N-labeled or ^{13}C, ^{15}N-double labeled mPHD, mDPPA3 and

mPHD-mDPPA3, M9 minimal media containing 0.5 g/L ^{15}NH$_4$Cl or 0.5 g/L ^{15}NH$_4$Cl and 1 g/L

^{13}C-glucose was used instead of LB media. Site-directed mutagenesis was performed by

designing two primers containing the mutations. The mutants of mPHD and mDPPA3 were

purified using the same protocol.

GST pull-down assay

MagneGST^{TM} Glutathione Particles (Promega) were used for the assay. The truncated

GST-mUHRF1 (10 μg) were immobilized on the beads (10 μL) equilibrated with the binding

buffer (20 mM Tris-HCl [pH 7.5], 150 mM NaCl, 1 mM DTT, 10% glycerol, and 0.5 % NP-

40). After the unbound proteins were washed-out thrice by the 200 μL of the binding buffer, 10 μg of C-terminal fragment of mDPPA3 (residues 61–150 or 76–128W) was incubated with the beads for 2 h at 4°C. After incubation, the unbound protein was washed thrice using 200 μL of the binding buffer. The bound proteins were boiled for 2 min at 95°C in an SDS sample buffer and analyzed by SDS-PAGE.

ITC

Microcal PEAQ-ITC (Malvern) was used for the ITC measurements. Wild-type and mutants of mPHD and mDPPA3 were dissolved in 10 mM HEPES (pH 7.5) buffer containing 150 mM NaCl and 0.25 mM TCEP. All measurements were carried out at 293K. The data were analyzed with Microcal PEAQ-ITC analysis software using a one-site model. For each interaction, at least three independent titration experiments were performed to show the dissociation constants with the mean standard deviations.

NMR

All NMR experiments were performed on Bruker BioSpin Avance III HD spectrometers with a TCI triple-resonance cryogenic probe-head with basic ^{1}H resonance frequencies of 500.13, 800.23 and 950.15 MHz. Three-dimensional (3D) spectra for main-chain signal assignments: HNCACB, HN(CO)CACB, HNCA, HN(CO)CA, HNCO and HN(CA)CO, and for side-chain signal assignments: HBHA(CBCACO)NH, H(CCO)NH, CC(CO)NH, HCCH-TOCSY and (H)CCH-TOCSY, for structure analysis: ^{1}H-^{13}C NOESY-HSQC, and ^{1}H-^{15}N NOESY-HSQC spectra were acquired at 293 K for 660 μM [^{13}C, ^{15}N]-mPHD-mDPPA3 fusion protein dissolved in PBS buffer (pH 7.0) containing 1 mM DTT and 5% D_2O. For the main-chain signal assignments of isolated proteins, 500 μM [^{13}C, ^{15}N]-mPHD in the free, 350 μM [^{13}C, ^{15}N]-mPHD in the complex with mDPPA3 at molar ratio of 1:2, 500 μM [^{13}C, ^{15}N]-mDPPA3 in the free state, and 350 μM [^{13}C, ^{15}N]-mDPPA3 in the complex with

mPHD at molar ratio of 1:2 were used in the buffer same as the fusion protein. The spectral

widths (the number of total data points) of each spectrum were 24 ppm (2,048) for the [1]H

dimension, and 30 ppm (256) for the [15]N dimension. All 3D spectra except for [1]H-[13]C NOESY-

HSQC were acquired by means of non-uniform sampling (NUS) to randomly reduce t1 and t2

time-domain data points typically around 25%. The uniformly sampled data were reconstructed

from the raw NMR data according to the sparse sampling schedules using several techniques

such as IST, SMILE, MDD and IRLS(39–41). Chemical shift perturbation experiments were

performed by recording 2D [1]H-[15]N HSQC spectra of 30 µM [[15]N]-mPHD dissolved in the same

buffer. All NMR spectra were processed using NMRPipe(42). For the NMR analysis, an

integrated package of NMR tools named MagRO-NMRViewJ, version 2.01.39 [the upgraded

version of Kujira(43)], on NMRView(44) was used for automated signal identification and

noise filtration using convolutional neural networks [CNNfilter(45)]. The filtered signal lists

were applied to calculations for automated signal assignments by FLYA(46) and then the signal

assignments were used for prediction of dihedral angles by TALOS+(47), and automated NOE

assignments and structure calculation by CYANA(48) (Table 1). Finally, water refinement

calculations by AMBER12 were performed for the lowest energy structures (20 models) (Table

2).

Furthermore, we ran fully automated peak picking and noise filtration on MagRO for

all spectra required for FLYA calculation using structure calculation mode with peak tables for

the spectra: [1]H-[15]N HSQC, HNCO, HN(CA)CO, HNCA, HN(CO)CA, CBCA(CO)NH,

HNCACB, HCCH-TOCSY,[15]N-edited NOESY, and [13]C-edited NOESY. After the 1st FLYA

calculation, ~80% of backbone and ~60% side-chain signals were automatically assigned.

These assigned chemical shifts were imported into MagRO from the output flya.tab file,

according to the following criteria: among 20 assigned chemical shift tables described in

flya.tab, a cut-off value of 80% was set to eliminate poorly populated assigned chemical shifts

after the final consolidation stage, and several proton signals such as Thr-OH, Ser-OH, His-

H$_{e2}$, His-H$_{d1}$, and Phe-H$_z$ were also neglected. Following visual inspection of the assigned data using 3D spectra such as H(CCCO)NH, CC(CO)NH and HBHA(CO)NH to confirm and correct the assignments, 2nd FLYA calculation was performed with the confirmed assigned chemical shifts to assign the remaining signals. TALOS+ calculation was performed using chemical shifts for ^{1}HN, ^{15}N, ^{13}C$_a$, ^{13}C$_b$, and ^{13}CO to predict the dihedral angles (phi and psi) of backbone. MagRO automatically converted the predicted dihedral angles to talosp.aco restrain file in the CYANA format except for the data which may not be trustful: low predicted order parameter (less than 0.8) and worse class annotated as "Warn" or "Dyn", as well as with setting a minimum angle (±20-deg). In the CYANA calculation, we applied 6 upper- and lower-limit distance constraints (total 82) to form tetrahedral coordinates for each Zn^{2+} atom, which was topologically linked with pseudo residues "PL" and "LL2". We assumed that the members of residues involved in the three Zn-finger coordinates from the typical fashion found in PHD domains, namely site1: C-C-C-C, site2: C-C-C-C, and site3: C-C-H-C.

For the obtained CYANA structures (20 models) with the lowest target function, implicit water refinement calculations were performed by AMBER12. Dihedral angle constraints (derived from TALOS+ prediction) and distance constraints including additional chirality and backbone omega angle constraints were converted for AMBER format using the SANDER tool(49). The distance constraints exported by CYANA (final.upl) were exclusively used (purely derived from experimental NOE data). Metal coordinate parameters, ZAFF.prep and ZAFF.frcmod, were appended to the standard force field, ff99SB, for the three zinc-finger cites according to the AMBER tutorial [http://ambermd.org/tutorials/advanced/tutorial20/ZAFF.htm]. In each initial stage of the refinement, 500 steps of energy minimization (250 steps of steepest gradient, followed by 250 steps of conjugate gradient decent) without electro-static energy and NMR constrain terms. The calculation was followed by a short molecular dynamics calculation (total 100 psec, time step 1.0 fsec, using SHAKE algorism for bond and angle stabilization) using electro-static

energy based on generalized born model (salt concentration: 100 mM, disabled Surface Accessibility (SA) function, electro-static potential radius cut-off: 18 Å) and NMR constrain terms. After the temperature was gradually increased from 0 to 300 K for 1,500 steps, dynamic calculation at 300 K was ran for remaining steps. In the final stage of the refinement, 2,000 steps of energy minimization were performed with the same energy terms.

^{1}H-^{15}N heteronuclear NOE (het-NOE) spectra of mPHD-mDPPA3 were measured with and without ^{1}H saturation applied before the start of the measurement[50]. The het-NOE values were calculated as ratios of the signal intensities of the two spectra. For signal intensity of each residue, the experimental error was estimated from the target signal around noise amplitude.

For the competition experiments, ^{1}H-^{15}N HSQC spectra were measured at 293K for 80 µM [^{15}N]-mPHD in the presence of mDPPA3$_{76\text{-}128W}$ and/or the H3 1-37W peptide at molar ratios (mPHD:mPDDA3:H3) of 1:0:0, 1:2:0, 1:0:2 and 1:2:8.

CD

Far-UV circular dichroism (CD) spectra were obtained using a JASCO J-720W model spectrometer. All samples were prepared to a concentration of 20 µM. Measurements were performed at 293K with a path length of 1 mm.

In vitro **ubiquitination assay**

Protein expression in *E. coli* and purification of mouse UBA1 (E1), human UBE2D3 (E2), mouse UHRF1 (E3), C-terminal FLAG tagged-H3$_{1\text{-}37W}$ and ubiquitin were performed according to the methodology in previous reports[9]. Ubiquitination reaction mixtures contained 100 µM ubiquitin, 200 nM E1, 8 µM E2, 3 µM E3, 5 mM ATP, and 10 µM C-terminal FLAG tagged-H3$_{1\text{-}37W}$ in the presence and absence of 10 µM mDPPA3$_{76\text{-}128W}$ in ubiquitination reaction buffer (50 mM Tris-HCl [pH 8.0], 50 mM NaCl, 5 mM MgCl$_2$, 0.1% Triton X-100, 2

mM DTT). The mixture was incubated at 30°C for 3 h, and the reaction was stopped by adding 3×SDS loading buffer. The reaction was analyzed by SDS-PAGE, followed by Western blotting using 1/5,000 diluted anti-FLAG antibody (Cell Signaling Technology, #2368).

Xenopus interphase egg extracts and purification of chromatin

Xenopus laevis was purchased from Kato-S Kagaku and handled according to the animal care regulations at the University of Tokyo. Interphase egg extracts were prepared as described previously(11). Unfertilized *Xenopus* eggs were dejellied in 2.5% thioglycolic acid-NaOH (pH 8.2) and were washed three times in 0.2×MMR buffer (5 mM HEPES-KOH [pH 7.6], 100 mM NaCl, 2 mM KCl, 0.1 mM EDTA, 1 mM MgCl$_2$, 2 mM CaCl$_2$). After activation in 1×MMR supplemented with 0.3 µg/mL calcium ionophore, eggs were washed four times with EB buffer (10 mM HEPES-KOH [pH 7.7], 100 mM KCl, 0.1 mM CaCl$_2$, 1 mM MgCl$_2$, 50 mM sucrose). Packed eggs were crushed by centrifugation (BECKMAN, Avanti J-E, JS13.1 swinging rotor) for 20 min at 18,973×g. Egg extracts were supplemented with 50 µg/mL cycloheximide, 20 µg/mL cytochalasin B, 1 mM DTT, 2 µg/mL aprotinin, and 50 µg/mL leupeptin and clarified for 20 min at 48,400×g (Hitachi, CP100NX, P55ST2 swinging rotor). The cytoplasmic extracts were aliquoted and stored at -80°C. Chromatin purification after incubation in egg extracts was performed as previously described with modifications. Sperm nuclei were incubated in egg extracts supplemented with an ATP regeneration system (20 mM phosphocreatine, 4 mM ATP, 5 µg/mL creatine phosphokinase) at 3000–4000 nuclei/µL at 22°C. Aliquots (15 µL) were diluted with 150-200 µL chromatin purification buffer (CPB; 50 mM KCl, 5 mM MgCl$_2$, 20 mM HEPES-KOH [pH 7.7]) containing 0.1% NP-40, 2% sucrose and 2 mM NEM. After incubating on ice for 5 min, extracts were layered over 1.5 mL of CPB containing 30% sucrose and centrifuged at 15,000×g for 10 min at 4°C. Chromatin pellets were resuspended in 1×Laemmli sample buffer, heated for 5 min and analyzed by SDS-PAGE. Recombinant Flag-tagged mDPPA3 was added to egg extracts at 400 nM.

DNA methylation was monitored by the incorporation of S-[methyl-3^H]-adenosyl-L-methionine. Extracts supplemented with S-[methyl-3^H]-adenosyl-L-methionine and sperm nuclei were incubated at 22°C in the presence or absence of recombinant Flag-tagged mDPPA3 protein. After 90 min incubation, the reaction was stopped by the addition of CPB containing 2% sucrose up to 300 μL. Genomic DNA was purified using a Wizard Genomic DNA Purification Kit (Promega) according to the manufacturer's instructions. The incorporation of radioactivity was quantified with a liquid scintillation counter.

GST pull-down assay using *Xenopus* egg extracts

For GST pull-down experiments using *Xenopus* egg extracts, mouse DPPA3 cDNA was sub-cloned into a pGEX4T-3 plasmid using In-Fusion (Clontech) according to the manufacturer's instructions. GST or GST-mDPPA3 proteins were expressed in *E.coli* BL21 (BL21-CodonPlus) by the addition of 0.1 mM IPTG to media followed by incubation for 12 h at 20°C. Bacteria cells were harvested and resuspended in lysis buffer (20 mM HEPES-KOH [pH 7.6], 500 mM NaCl, 0.5 mM EDTA, 10% glycerol, 1 mM DTT) supplemented with 0.5% NP-40 and protease inhibitors and were then disrupted by sonication on ice. After the cell debris was removed by centrifugation, the recombinant proteins were immobilized on GS4B resin by incubation for 2 h at 4°C. After the unbound proteins was washed-out by the lysis buffer, interphase egg extracts were incubated with the beads for 2 h at 4°C. After incubation, the beads were washed four times with CPB containing 2% sucrose, 500 mM KCl and 0.1% Triton X-100. The washed beads were resuspended in 20 μL of 2×Laemmli sample buffer and 20 μL of 1×Laemmli sample buffer, boiled for 5 min at 100°C, and analyzed by immunoblotting.

For FLAG-tagged protein expression in insect cells, 3×FLAG-tagged mDppa3 WT or mutants were sub-cloned into the pVL1392 vector. Baculoviruses were produced using a BD BaculoGold Transfection Kit and a BestBac Transfection Kit (BD Biosciences), following the manufacturer's protocol. Proteins were expressed in Sf9 insect cells by infection with viruses

expressing 3×FLAG-tagged mDPPA3 WT or its mutants for 72 h at 27°C. Sf9 cells from a 750 mL culture were collected and lysed by resuspending them in 30 mL lysis buffer, followed by incubation on ice for 10 min. A soluble fraction was obtained after centrifugation of the lysate at 15,000×g for 15 min at 4°C. The soluble fraction was incubated with 250 µL anti-FLAG M2 affinity resin equilibrated with lysis buffer for 4 h at 4°C. The beads were collected and washed with 10 mL of wash buffer and then with 5 mL of EB (20 mM HEPES-KOH [pH 7.5], 100 mM KCl, 5 mM MgCl$_2$) containing 1 mM DTT. Each recombinant protein was eluted twice in 250 µL EB containing 1 mM DTT and 250 µg/mL 3×FLAG peptide (Sigma-Aldrich). Eluates were pooled and concentrated using a Vivaspin 500 (GE Healthcare).

Cell culture and cell line generation

Dppa3_KO/UHRF1-GFP mESCs were described previously(35, 51) and were cultivated in Dulbecco's Modified Eagle's Medium-High Glucose supplemented with homemade recombinant LIF(52), 16% Fetal Bovine Serum (FBS, Sigma) 0.1 mM β-mercaptoethanol ready to use, (Life Technologies), 2 mM L-glutamine, respectively 100 U/mL and 100 µg/mL of Pen/Strep (Sigma), 1×non-essential amino acids (Sigma) and with 2i inhibitors (1 µM PD32591 and 3 µM CHIR99021, Axon Medchem). Cells were cultured in gelatine (0.2%) coated Corning dishes at 37°C in a 5% CO$_2$ incubator and gently dissociated for passaging at 1:8 ratio every second day with Stem Pro Accutase (Gibco). Cells were washed with Dulbecco's Phosphate Buffer Saline cell culture grade.

To generate stable doxycycline inducible Dppa3 mESC lines, 400,000 Dppa3_KO/UHRF1-GFP mESCs were seeded into one well of a 6-well plate 4 h before transfection. Then cells were transfected with equimolar amounts of pSBtet-3×FLAG-Dppa3_wt-mScarlet-PuroR or pSBtet-3×FLAG-Dppa3_mutants-mScarlet-PuroR or pSBtet-3×FLAG-B.Taurus Dppa3-mScarlet-PuroR, and the Sleeping Beauty transposase, pCMV(CAT)T7-SB100(53) (Addgene plasmid #34879) vector using Lipofectamine 3000

(Thermo Fisher Scientific) according to manufacturer's instructions. After 48 h transfection, cells were seeded onto p100 plates into clonal density under puromycin selection with a concentration of 1 µg/mL for 7 days. After the seventh day selection, colonies with mScarlet signal under induction of doxycycline (1 µg/mL) were picked and mixed for further analyses. The stable cell lines were kept in puromycin selection of 1 µg/mL starting two weeks after expansion.

Sleeping beauty constructs

To generate inducible mouse Dppa3 mutants and Bos Taurus Dppa3 expression constructs, the sequences coding for mDppa3 mutants, including R104A, R89A/V90A, L91A/V94G, M102P/E109P, and Bos Taurus Dppa3 were synthesized as gBlocks (IDT, Coralville, IA, USA) and inserted into the pSB_Avi_3×FLAG_insert_mScarlet_Puro vector (linearized by AsiSI and NotI) using Gibson assembly based on protocol instructions of 2×HiFi master mix (New England Biolabs). The correct constructs were selected based on Sanger Sequencing results (Eurofins).

Cellular fractionation and western blotting

Cell fractionation was performed as described previously with minor modifications(54). Approximately 1×10^7 mESCs were resuspended in 100 µL of buffer A (10 mM HEPES [pH 7.9], 10 mM KCl, 1.5 mM $MgCl_2$, 340 mM sucrose, 10% glycerol, 0.1% Triton X-100, 1 mM DTT, 1 mM phenylmethylsulfonyl fluoride (PMSF), 1×mammalian protease inhibitor cocktail (PI; Roche)) and incubated for 5 min on ice. Nuclei were collected by centrifugation (5 min, 1300×g, 4°C) and the cytoplasmic fraction (supernatant) was cleared again by centrifugation (15 min, 20,000×g, 4°C). Nuclei were washed once with buffer A and resuspended in 100 µL of buffer A. The cytoplasmic and nuclear fractions were supplemented with 4×Laemmli buffer and boiled for10 min at 95°C.

Western blots were performed using the following antibodies: anti-alpha-Tubulin (monoclonal; 1:2000; Sigma, T9026), rabbit anti-H3 (polyclonal; 1:5000; Abcam, ab1791), mouse anti-GFP (monoclonal; 1:5000; Roche), goat anti-rabbit HRP (polyclonal; 1:5000; BioRad), rabbit anti-mouse HRP (1:5000; Invitrogen, Cat # A27025).

Live cell imaging

Live cell experiments were conducted with a Nikon TiE microscope equipped with a Yokogawa CSU-W1 spinning disk confocal unit (pinhole size 50 μm), together with a Andor Borealis illumination unit, [Andor ALC600 laser combiner (405 nm/ 488 nm/ 561 nm/ 640 nm)]. The images were acquired with an Andor IXON 888 Ultra EMCCD camera, with 100×/1,45NA oil immersion objective through the interface of the software NIS Elements (Version 5.02.00) in Perfect Focus System with lasers at 488 nm for GFP, 561 nm for mScarlet and 640 nm for Sir-DNA. For live cell imaging, 80,000 cells were seeded into one well of an 8 well glass bottom chamber slide (Ibidi) coated one day before with Geltrex Ready-to-Use (Thermo Fisher Scientific). The next day, 1 μM Sir-DNA (Spirochrome) for DNA staining was added into medium 1 h before live cell imaging. Images were acquired before and after 50 min of doxycycline induction with the same laser power, acquisition time and gain.

Targeted bisulfite amplicon sequencing (TaBAseq)

TaBAseq was performed as described previously(35). According to the manufacturer's instructions, there were 1×10^6 ESCs culturing in 2iLIF condition for genomic isolation with QIAamp DNA Mini Kit (Qiagen). 500 ng of gDNA was used for bisulfite conversion followed by the instructions of EZ DNA Methylation-Gold Kit (Zymo), which was eluted in a 2×20 μL Elution Buffer.

TaBAseq is based on two sequential PCRs. The first one amplifies locus-specific LINE-1 elements and the second one indexes the sample-specific amplicon with Ilumina's

Truseq and Nextera compatible overhangs. The amplicon-specific PCR was performed in a total volume of 25 µL containing 0.4 µM of forward and reverse primers, 2 mM betaiinitialne (Sigma-Aldrich), 10mM tetramethylammonium chloride solution (Sigma-Aldrich), 1×MyTaq Reaction Buffer, 0.5 units of MyTaq HS polymerase (Bioline) and 1 µL of bisulfite converted DNA (12.5 ng). The cycling parameters were as follows: 5 min at 95°C for initial denaturation, 40 cycles (95°C for 20 s, 58°C for 30 s, 72°C for 25 s) and with the final elongation 72°C for 3 min. To check for the quality and yield of the PCR reaction, it was run on a 2% agarose gel and to purify the PCR amplicons, it was used CleanPCR beads with 1.8 × the volume of the remaining PCR reaction. The magnetic beads were immobilized with DynaMag-96 Side Magnet (Thermo Fisher) for 5 min. After the supernatant is removed, the beads are washed 2 × with 150 µL of fresh 70 % ethanol. After leaving the beads to air-dry for 5 min, DNA was eluted in 15 µL of elution buffer (10 mM Tris-HCl [pH 8.0]). DNA concentration was determined using Nanodrop (Thermo Fisher Scientific).

To index the amplicons, a second PCR was performed in a 25 µL total volume, containing 0.08 µM (1 µL of a 2 µM stock) of the Indexing Primers, respectively i5 and i7, 1×MyTaq Reaction Buffer, 0.5 units of MyTaq HS Polymerase (Bioline) and 1 µL of the purified PCR purified amplicon. The cycling parameters are as follows: 5 min at 95°C for initial denaturation, 18 cycles (95°C for 20 s, 55°C for 30 s, 72°C for 40 s) and with the final elongation at 72°C for 5 min. Again 2% agarose gel was used to determine the yield and purity of the PCRs. The PCR-indexed amplicons were purified as described above using the CleanPCR magnetic beads. The DNA concentration for each sample was determined with Quant-iT PicoGreen dsDNA Assay Kit (Thermo Fisher Scientific). The final library was created by pooling together an equimolar ratio of all PCR products in a final concentration of 1 ng/µL of the total library. The final library concentration was once again estimated via Quant-iT PicoGreen, while the size distribution and quality of the library were assessed using Bioanalyzer. Ilumina Miseq was used to sequence the dual-indexed TaBAseq library with a

2×300 bp paired-end, with 5% sequencing coverage. The sequencing data was analyzed with a

TABSAT package(55).

Protein sequence alignments

Protein sequence alignments for UHRF1 and DPPA3 were performed using Jalview software

(www.jalview.org) with sequences retrieved from Uniprot for UHRF1: (Q8VDF2_Mus

musculus; Q7TPK1_Rattus norvegicus; G3RVG5_Gorilla gorilla gorilla; Q96T88_Homo

sapiens; H2QF26_Pan troglodytes; A0A3Q7TCY9_Vulpes vulpes; A0A3Q7V1Z9_Ursus

arctos horribilis; A0A2U3VKN8_Odobenus rosmarus divergence; A0A2Y9KPL3_Enhydra

lutris kenyoni; A7E320_Bos Taurus) and DPPA3 (Q8QZY3_Mus musculus; Q6IMK0_Rattus

norvegicus; G3RB81_Gorilla gorilla gorilla; Q6W0C5_Homo sapiens; H2Q5C8_Pan

troglodytes; A0A3Q7U513_Vulpes vulpes; A0A3Q7W0Q7_Ursus arctos horribilis;

A0A2U3ZR98_Odobenus rosmarus divergence; A0A2Y9IX83_Enhydra lutris kenyoni;

A9Q1J7_ Bos Taurus). Sequences are aligned with the multiple sequence alignment algorithm

MAFFT (http://mafft.cbrc.jp/alignment/software/). The percentage of protein identity

conserved for both proteins is calculated through Pairwise alignment algorithm between Mus

musculus and different species. The phylogenetic tree was generated with online tools,

https://www.genome.jp (PhyML) and iTOL.

19
20

RESULTS

Biochemical assay for determining essential regions for complex formation

The C-terminal fragment of mDPPA3 is required for binding to mUHRF1 (Figure 1A,

B)(36). GST pull-down assays using truncated mouse UHRF1 showed that the PHD finger

(residues 304–372: mPHD) was sufficient for binding to the C-terminal fragment of mDPPA3

(residues 61–150) (Supplementary Figure S1A, B). Further optimization revealed that

mDPPA3 residues 76–127, followed by an additional Trp (mDPPA3$_{76-128W}$), were sufficient to

bind to the mPHD (Figure 1C). This interaction was validated by isothermal titration

calorimetry (ITC) and NMR spectroscopy. mDPPA3$_{76-128W}$ bound to mPHD with a K_D of 27.7

nM, which is stronger than the binding of the N-terminal tail of H3 (K_D: 1590 nM) and PAF15

(K_D: 3523 nM) to mPHD, which are well-known ligands of the UHRF1 PHD finger (Figure

1D, Supplementary Figure S2). The ^{1}H-^{15}N heteronuclear single quantum coherence (HSQC)

spectrum of [^{15}N]-mPHD titrated with non-labeled mDPPA3$_{76-128W}$ was markedly different

from that of apo-mPHD, in which nearly all HSQC signals were shifted in the slow-exchange

regime on the chemical shift timescale (Supplementary Figure S3A, B). The weighted averages

of the ^{1}H and ^{15}N chemical shift differences ($\Delta\delta$) between the apo-mPHD and mPHD bound to

mDPPA3 showed relatively large values for Glu333, Leu337, Met343, and Glu362 in mPHD

(Supplementary Figure S3A–C). ITC data indicated that the L337A mutation in mPHD

severely weakened the binding to mDPPA3$_{76-128W}$ with a K_D > 43,000 nM (Figure 1D,

Supplementary Figure S2) and also abolished the binding to the N-terminal tail of H3 and

PAF15 (Supplementary Figure S2). The ^{1}H-^{15}N HSQC spectrum of mPHD L337A showed that

the mutation had a modest effect on the native conformation (Supplementary Figure S3D).

Collectively, Leu337 of mPHD is commonly used as the interface for binding to mDPPA3, H3,

and PAF15.

Overall structure of mPHD in complex with mDPPA3

To uncover the molecular mechanism by which mDPPA3 binds to mPHD with high affinity, we determined the structure of the complex using solution NMR analysis. As a target for structural analysis, we designed a chimeric protein, mDPPA3$_{76-127}$ was fused to the C-terminal end of mPHD (mPHD-mDPPA3). The ^{1}H-^{15}N HSQC spectrum of the mPHD moiety in [^{15}N]-mPHD-mDPPA3 was superimposed on that of [^{15}N]-mPHD mixed with non-labeled mDPPA3$_{76-128W}$, validating that the chimeric protein has a binding mode equal to that of the isolated proteins (Supplementary Figure S4A).

The ensemble of the mPHD-mDPPA3 structures was well converged and showed a low average root mean square deviation (rmsd) of 0.40 Å for Cα atom coordinates (Figure 2A, B, Table 1 and 2). The mPHD moiety comprised of pre-PHD (residues 304–321) and core-PHD (residues 322–372) encompassing three zinc finger motifs (Zn1-3) and an anti-parallel β-sheet (Figure 2A). The residues 307–369 of the mPHD moiety were superimposed on the solution structure of apo-mPHD (PDB: 6VFO) with a Cα rmsd of 3.1 Å (Supplementary Figure S5), indicating that the overall structure of mPHD did not change substantially upon binding of mDPPA3, except for a loop region, as mentioned later.

The ^{1}H-^{15}N HSQC spectrum of [^{15}N]-mDPPA3 in the free state showed sharp signals in a very narrow range of ^{1}H chemical shifts corresponding to a randomly coiled state, indicating the unstructured conformation of apo-mDPPA3 (Supplementary Figure S4B). Binding of mPHD induced two α-helices in mDPPA3: a short α-helix, αS1 (residues 90–95) and a long α-helix αL2 (residues 97–118), which are connected via a short turn, resulting in an 'L' shaped motif in mDPPA3 (Figure 2A, B). The helical structure induced upon binding of mPHD was further experimentally confirmed by NMR and circular dichroism (CD) spectrum analysis. Secondary structure prediction by TALOS-N using the backbone ^{1}HN, ^{15}N, ^{13}C$_\alpha$, ^{13}C$_\beta$, and ^{13}CO chemical shifts showed the induction of helical structure of mDPPA3 upon binding of mPHD (Supplementary Figure S4C, D)(56). The CD spectrum of mPHD mixed with

mDPPA3 showed a negative peak at 222 nm, whereas this property was not observed in the spectra of mPHD and mDPPA3 alone (Figure 2C). ^{1}H-^{15}N heteronuclear nuclear Overhauser effect (het-NOE) of [^{15}N]-mPHD-mDPPA3, which is sensitive to local conformational flexibility on the picosecond to nanosecond timescale(57), was consistently high except for the residues on the N-terminal region of the mPHD moiety, residues 76–84, and the C-terminus of mDPPA3 (Figure 2B, D). The NMR structure showed that residues 85–118 of mDPPA3 in the complex were structurally well converged (Figure 2B) and ITC data demonstrated that mDPPA3$_{85–118}$ bound to mPHD with a $K_D = 45$ nM, which is comparable to the binding affinity of mDPPA3$_{76–128W}$, indicating that residues 85–118 of mDPPA3 were sufficient for binding to mPHD (Supplementary Figure S2). mDPPA3 provides a wide interface for contact with mPHD; the estimated contact area between the two proteins was ~1360 Å^2, which comprises three parts: a ^{88}VRT^{90} cassette, and αS1 and αL2 of mDPPA3 (Figure 2E). In contrast, H3 and PAF15 bind to the human UHRF1 PHD finger with a contact area of ~400 Å^2 (PDB: 3ASL) and ~360 Å^2 (PDB: 6IIW), respectively. The contact areas of the complex structures are much smaller than those of the mPHD:mDPPA3 complex, supporting the stronger binding affinity between mPHD and mDPPA3. NMR titration experiments demonstrated that mDPPA3 preferentially bound to mPHD even in the presence of excess H3 peptide (Supplementary Figure S4E), indicating that mDPPA3-binding to mPHD competes with the H3-binding.

Recognition of the conserved VRT cassette in mDPPA3 by mPHD

The ^{88}VRT^{90} cassette of mDPPA3 adopted an extended conformation and fit into the shallow acidic groove on mPHD via an intermolecular anti-parallel β sheet (Figure 2A). The side chain of Val88 in the ^{88}VRT^{90} cassette was surrounded by Leu336, Val357, Pro358, Glu360, and Trp363 of mPHD within van der Waals contacts (Figure 3A). The guanidino group of Arg89 in the cassette was located within a distance that enabled hydrogen bonding with the side-chain carboxyl groups of Asp339 and Glu362 of mPHD (Figure 3A). The side-chain

hydroxyl and methyl groups of Thr90 in mDPPA3 interacted with mPHD differently: the methyl group formed hydrophobic interactions with Leu336, Leu337, and Val357 of mPHD, whereas the hydroxyl group was positioned within the hydrogen bond distance with the main chain amide of Ser93, indicating its function as a helix-cap for the N-terminus of the following αS1 (Figure 3B). ITC data showed that the R89A/T90A mutations in the [88]VRT[90] cassette of mDPPA3 abolished its interaction with mPHD (Figure 3C). This interaction is supported by GST pull-down experiments with *in vitro* translated full-length mUHRF1 and GST-mDPPA3, which showed that the R89A/T90A mutations are sufficient to block the binding of mDPPA3 to mUHRF1 (Figure 3D), indicating that the [88]VRT[90] cassette in mDPPA3 plays a critical role in binding to mPHD.

The [88]VRT[90] cassette of mDPPA3 was well-conserved in the N-terminal sequences of PAF15 ([1]VRT[3]) and H3 ([1]ART[3]). Although the [88]VRT[90] cassette of mDPPA3 is not located at its N-terminal end, the cassette was recognized by the shallow acidic groove of mPHD, which is used for PAF15/H3-binding (Figure 3E–G). Notably, compared with the PHD moiety structure in the complexes with PAF15/H3 and mDPPA3, the conformation of loop residues 356–364 (hereafter loop[PHD]) in mPHD was markedly different (Figure 3H). In the complex with PAF15/H3, loop[PHD] functions as a wall to recognize the N-terminal amino group of PAF15/H3 by hydrogen bonding using the main chain carbonyl oxygen of Glu355 (Figure 3F, G). In contrast, in the complex with mDPPA3, loop[PHD] is shifted outward away from the amide nitrogen of Val88 of mDPPA3, resulting in disruption of the hydrogen bond between the amide nitrogen in mDPPA3 and the main-chain carbonyl oxygen of Glu360 (corresponding to residue Glu355 in humans) in the loop[PHD] of mPHD (Figure 3H). The results indicate that loop[PHD] has an intrinsically flexible capability to accommodate the VRT cassette of mDPPA3. The structural rearrangement of loop[PHD] permits the peptide bond moiety between Arg87–Val88 to enter the groove (Figure 3E, H). The side-chain conformation of Arg89 of mDPPA3 in the complex with mPHD was different from that of Arg2 of H3/PAF15 (Figure 3E–G). Given that

ITC data for the R89A mutation of mDPPA3 showed an ~18-fold decrease in binding affinity

to mPHD (K_D = 361 nM; Figure 3C, Supplementary Figure S2), the Arg89 guanidino group of

mDPPA3 interacts with the acidic surface of mPHD.

Binding mode of two-helices in mDPPA3 to mPHD

In addition to the [88]VRT[90] cassette, mDPPA3 utilizes two α-helices, αS1 and αL2, to

bind to mPHD. The αS1 of mDPPA3 forms a small hydrophobic cluster composed of the side

chain methyl groups of Leu91, Val94, and Leu95, of which Leu91 and Val94 interact with the

hydrophobic patch of mPHD comprising Leu337 and Ala344 (Figure 4A). As mentioned before,

the L337A mutation in mPHD largely impaired the interaction with mDPPA3; similarly, the

L91A/V94A mutations in mDPPA3 also reduced the binding affinity to mPHD with a K_D of

1084 nM (Figure 4B), indicating that αS1 contributes to stable complex formation.

αL2 of mDPPA3, comprising residues 99–118, was embedded into a concave surface

between the pre- and core-PHD domains (Figure 2E). Notably, the introduction of the helix

breaking residue M102P/E109P mutations, or truncation of αL2 residues 76–106 in mDPPA3

resulted in a significant reduction in binding to mPHD, indicating that the formation of αL2 is

critical for interaction with mPHD (Figure 4B, Supplementary Figure S2). Interestingly, the

Arg104 guanidino group on αL2 in mDPPA3 was deeply buried in the binding interface of the

complex, in which the side chain formed hydrogen bonds with the main chain carbonyl oxygens

of Lys320 and Cys321 in the pre-PHD domain (Figure 4A). Ile108 and Arg111 of mDPPA3

also interacted with Cys341 and Asp342 of the core-PHD domain, indicating that αL2 bridges

the pre- and core-PHD domains (Figure 4A). Although mutations of Ile108 and Arg111 of

mDPPA3 had a limited effect on binding to mPHD (Figure 4B, Supplementary Figure S2), the

R104A mutation significantly reduced the binding affinity to mPHD (Figure 4B). GST pull-

down experiments demonstrated that the L91A/V94A mutation in αS1 reduced the binding of

GST-mDPPA3 to full-length mUHRF1 modestly (Figure 4C), showing a limited effect on full-

length proteins. In contrast, the M102P/E109P and R104A mutations in αL2 of mDPPA3 markedly reduced binding to mUHRF1 (Figure 4C), indicating that αL2 of mDPPA3 plays a pivotal role in complex formation with full-length proteins.

Mutational analyses of interaction interface of DPPA3 with UHRF1 in *Xenopus* egg extracts and mESCs

To test the function of the DPPA3 mutants, we measured their ability to inhibit UHRF1-dependent maintenance of DNA methylation in *Xenopus* egg extracts (Figure 5A). We compared the inhibitory activity of GST-mDPPA3 (wild-type) with that of the R89A/T90A, L91A/V94A, M102P/E109P, and R104A mutants. Our results showed that wild-type mDPPA3 inhibited the chromatin recruitment of UHRF1 and DNA methylation; in contrast, the R89A/T90A mutant showed a severely impaired inhibitory effect (Figure 5B, C). Concordant with the ITC data and GST-pull down assay, the L91A/V94A mutant was ineffective in inhibiting UHRF1 (Figure 5B). Mutants linked to the αL2 helix structure (M102P/E109P and R104A) were unable to inhibit UHRF1 chromatin loading and DNA methylation (Figure 5B, C). In addition, *in vitro* ubiquitination assay using C-terminal FLAG tagged H3 1-37W as a substrate demonstrated that WT mDPPA3 inhibited the ubiquitination of H3 catalyzed by UHRF1, however, the mutants mDPPA3 failed to do so (Supplementary Figure S6), suggesting that DPPA3 represses the E3-ligase activity of UHRF1.

To further examine the importance of the interaction interface of DPPA3 and UHRF1, we analyzed chromatin displacement and nucleocytoplasmic translocation of UHRF1 by inducing the expression of DPPA3 mutants in mouse embryonic stem cells (mESCs). We generated inducible mDPPA3-mScarlet expression cassettes harboring mutations R89A/T90A, L91A/V94G, R104A, and M102P/E109P. After introducing these expression cassettes into DPPA3 knock out/UHRF1-GFP (D3KO/U1GFP) mESCs, we used live-cell imaging to observe the localization of UHRF1-GFP (Figure 6A). Wild-type mDPPA3 caused chromatin

displacement and nucleocytoplasmic translocation of UHRF1-GFP. In contrast, the mDPPA3

mutants failed to efficiently re-localize UHRF1-GFP (Figure 6A). Furthermore, biochemical

fractionation experiments showed that UHRF1-GFP was detected in the cytoplasm after

induction of wild-type mDPPA3, whereas the mDPPA3 mutants showed low activity for

UHRF1-GFP displacement and export to the cytoplasm. (Figure 6B). The L91A/V94G(A)

mutation of mDPPA3 had a more severe effect on mESCs than on *Xenopus* egg extracts (Figure

5B and 6B). We examined the ability of wild-type DPPA3 and the mutants to rescue the

hypermethylation of LINE-1 (long interspersed nuclear element) elements in DPPA3 knock-

out mESCs (Figure 6C). The mDPPA3 mutants failed to restore low methylation levels

(although the M102P/E109P mutant was somewhat less effective) (Figure 6C), indicating that

binding of mDPPA3 to UHRF1 PHD finger is required for DNA demethylation.

Next, we examined the evolutionary conservation of the DPPA3-UHRF1 interaction

interface. Although UHRF1 is highly conserved, phylogenetic analysis of DPPA3 showed the

prevalence of two divergent groups: group 1, with more than 24.6% similarity (*Mus musculus*,

Rattus norvegicus, *Gorilla gorilla*, *Homo sapiens*, and *Pan troglodytes*) and group 2 which

showed less than 18.6% identity (*Vulpes vulpes*, *Ursus arctos horribilis*, *Odobenus rosmarus*,

Enhydra lutris kenyoni, and *Bos taurus*) (Supplementary Figure S7), suggesting rapid evolution

of DPPA3. This is consistent with the observation that DPPA3 has recently evolved in

mammals and does not appear to have catalytic activity(35). ~~Notably, the VRT cassette of~~

~~DPPA3, a central element in the interaction with UHRF1, was highly conserved except in~~ *~~Bos~~*

~~taurus~~ ~~(Figure 1B), which could not displace UHRF1 in mESCs from chromatin (Figure 6A).~~

Collectively, the results of the functional assays indicate that the key residues of

mDPPA3 identified in our NMR structure of the complex are important for UHRF1 regulation.

DISCUSSION

The PHD finger is a well-known reader domain for post-translational modifications

of histone H3(58). Several structural studies of PHD fingers in complex with their ligands have revealed an N-terminal recognition rule for ligand recognition that can be applied to the vast majority of PHD fingers. PHD fingers have a shallow acidic groove for recognition of the N-terminus of ligands, in which the amino group of the first amino acid residue in the ligand forms hydrogen bond (s) with the PHD finger, resulting in a binding affinity with K_D in the μM range. Our NMR solution structure showed that mDPPA3 binds to mPHD via multifaceted interactions, utilizing the VRT cassette and two α-helices. The ^{88}VRT^{90} cassette of mDPPA3 is not located at the N-terminal end. However, the cassette was recognized by the shallow acidic groove of mPHD in a similar manner to the N-terminus of ^{1}ART^3 in H3 and ^{1}VRT^3 in PAF15. This binding is reinforced by the two α-helices unique to mDPPA3, resulting in high-affinity binding to mPHD (K_D = 27.7 nM). Although the N-terminus recognition rule does not apply to mDPPA3 binding to mPHD, stable complex formation is ensured by the multifaceted interaction provided by mDPPA3. The VRT cassette of mDPPA3 is well conserved among various species (Figure 1B), suggesting that binding of the VRT cassette to the UHRF1 PHD finger while disregarding the N-terminus recognition rule is a common molecular mechanism for the complex formation of DPPA3 and UHRF1. Intriguingly, phosphorylation of H3T3 negatively regulates the chromatin binding of UHRF1(59). NetPhos3.1 (https://services.healthtech.dtu.dk/service.php?NetPhos-3.1) predicts the high probability that T90 of mDPPA3 is target for protein kinase A (^{86}R$\underline{R}$VR$\underline{T}^{90}$), suggesting that post-translational modifications of mDPPA3 regulate its binding to UHRF1.

The VRT cassette and αS1 helix of DPPA3 identified in the NMR structure were relatively conserved between species, whereas the αL2 helix was less conserved and had a short insertion in the group 2 species of the DPPA3 family (Figure 1B, Supplementary Figure S7). AlphaFold2 predicted that the corresponding regions in humans (UniProt: Q6W0C5) and rats (UniProt: Q6IMK0) show a long helical structure, suggesting that the formation of the αL2 in the DPPA3 family is structurally conserved.

In general, a conventional PHD finger consists of one core-PHD domain, including two zinc fingers(58). In contrast, the UHRF1 PHD finger contains a pre-PHD domain, which provides an additional binding surface for mDPPA3. Protein sequence analysis of UHRF1 comparing 10 different species showed that UHRF1 was highly conserved during mammalian evolution (>40% similarity) (Supplementary Figure S7). In particular, the UHRF1 pre-PHD and PHD domains were very similar (61.1%) (Figure 1B), suggesting the conservation of the domain in the UHRF1 family. The αL2 of mDPPA3 is embedded in the concave surface formed between the pre- and core-PHD domains in mPHD. Breaking the folding of the αL2 helix severely impaired the binding affinity to mPHD, indicating that αL2 formation by mDPPA3 is crucial for its interaction with mPHD. Thus, the unique structural features of mPHD, consisting of pre- and core-PHDs, and of mDPPA3 with the two helices for enforcing VRT cassette binding ensure highly specific structural complementarity for complex formation. As a consequence, DPPA3 does not bind to the conventional PHD finger that lacks the pre-PHD domain.

The binding of the UHRF1 PHD domain to H3 is involved in its chromatin localization(29, 35, 36). The shallow acidic groove on mPHD is a common binding platform for mDPPA3 and H3. In addition to the shallow acidic groove, mDPPA3 utilizes two helices for binding to mPHD, which increases its binding affinity. DPPA3-binding to mPHD totally competes with the H3-binding (Supplementary Figure S4E). Thus, the molecular mechanism by which mDPPA3 inhibits chromatin binding of UHRF1 includes the preferential binding of mDPPA3 to the UHRF1 PHD finger. Indeed, mDPPA3 inhibited the ubiquitination of H3 catalyzed by UHRF1 *in vitro* (Supplementary Figure S6), suggesting that DPPA3 inhibits chromatin binding of UHRF1 and represses the E3-ligase activity of UHRF1 in cells. The binding mode of DPPA3 to UHRF1 is unique and DPPA3-binding explicitly inhibits chromatin loading of UHRF1. UHRF1 is overexpressed in many cancer cells and downregulation of UHRF1 in these cells leads to reactivation of tumor suppressor gene expression(60).

Overexpression of DPPA3 leads to tumor differentiation in hepatocellular carcinoma in which UHRF1 nuclear translocation is impeded(61). Thus, our structural analysis of the UHRF1 PHD in complex with DPPA3 may provide a framework for the design of new anticancer drugs. Peptide-like inhibitors that mimic the αL2 of DPPA3 and specifically bind to the concave surface between pre- and core-PHD domains inhibit excessive UHRF1 in cancer cells.

ACCESSION NUMBERS

NMR data and refined coordinate was deposited in the Protein Data Bank with

accession code 7XGA and Biological Magnetic Resonance Bank with accession code

bmr36483. Supplementary Information is linked to the online version of the paper at

www.nature.com/nature. Correspondence should be addressed to aritak@yokohama-cu.ac.jp.

Source data are provided with this paper.

Supplementary Data statement

Supplementary Data are available at NAR Online.

FUNDING

This study was supported by MEXT/JSPS KAKENHI (JP18H02392, JP19H05294 and

JP19H05741 to K.A., and 19H05285 and 21H00272 to A.N.), PRESTO (14530337) from JST

to K.A., Takeda Science Foundation (1871140003) to K.A., the grant for 2021–2023 Strategic

Research Promotion (No. SK201904) of Yokohama City University to K.A., Research

Development Fund of Yokohama City University to T.K. and the Deutsche

Forschungsgemeinschaft (DFG grant SFB 1064/A17 # 213249687 to H.L.).

CONFLICT OF INTEREST

The authors declare no competing interests.

REFERENCES

1. Schübeler,D. (2015) Function and information content of DNA methylation. *Nature*, **517**, 321–6.

2. Zeng,Y. and Chen,T. (2019) DNA Methylation Reprogramming during Mammalian Development. *Genes, Vol. 10, Page 257*, **10**, 257.

3. Greenberg,M.V.C. and Bourc'his,D. (2019) The diverse roles of DNA methylation in mammalian development and disease. *Nature Reviews Molecular Cell Biology, 20:10*, **20**, 590–607.

4. Seisenberger,S., Peat,J.R., Hore,T.A., Santos,F., Dean,W. and Reik,W. (2013) Reprogramming DNA methylation in the mammalian life cycle: building and breaking epigenetic barriers. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, **368**.

5. Bostick,M., Kim,J.K., Estève,P.-O., Clark,A., Pradhan,S. and Jacobsen,S.E. (2007) UHRF1 plays a role in maintaining DNA methylation in mammalian cells. *Science (New York, N.Y.)*, **317**, 1760–1764.

6. Sharif,J., Muto,M., Takebayashi,S., Suetake,I., Iwamatsu,A., Endo,T. a, Shinga,J., Mizutani-Koseki,Y., Toyoda,T., Okamura,K., *et al.* (2007) The SRA protein Np95 mediates epigenetic inheritance by recruiting Dnmt1 to methylated DNA. *Nature*, **450**, 908–912.

7. Smets,M., Link,S., Wolf,P., Schneider,K., Solis,V., Ryan,J., Meilinger,D., Qin,W. and Leonhardt,H. (2017) DNMT1 mutations found in HSANIE patients affect interaction with UHRF1 and neuronal differentiation. *Human Molecular Genetics*, **26**, 1522–1534.

8. von Meyenn,F., Iurlaro,M., Habibi,E., Liu,N.Q., Salehzadeh-Yazdi,A., Santos,F., Petrini,E., Milagre,I., Yu,M., Xie,Z., *et al.* (2016) Impairment of DNA Methylation Maintenance Is the Main Cause of Global Demethylation in Naive Embryonic Stem Cells. *Molecular Cell*, **62**, 848–861.

9. Nishiyama,A., Mulholland,C.B., Bultmann,S., Kori,S., Endo,A., Saeki,Y., Qin,W., Trummer,C., Chiba,Y., Yokoyama,H., *et al.* (2020) Two distinct modes of DNMT1 recruitment ensure stable maintenance DNA methylation. *Nature Communications*, **11**, 1222.

10. Ishiyama,S., Nishiyama,A., Saeki,Y., Moritsugu,K., Morimoto,D., Yamaguchi,L., Arai,N., Matsumura,R., Kawakami,T., Mishima,Y., *et al.* (2017) Structure of the Dnmt1 Reader Module Complexed with a Unique Two-Mono-Ubiquitin Mark on Histone H3 Reveals the Basis for DNA Methylation Maintenance. *Molecular Cell*, **68**, 350-360.e7.

11. Nishiyama,A., Yamaguchi,L., Sharif,J., Johmura,Y., Kawamura,T., Nakanishi,K., Shimamura,S., Arita,K., Kodama,T., Ishikawa,F., *et al.* (2013) Uhrf1-dependent H3K23 ubiquitylation couples maintenance DNA methylation and replication. *Nature*, **502**, 249–53.

12. Qin,W., Wolf,P., Liu,N., Link,S., Smets,M., Mastra,F. La, Forné,I., Pichler,G., Hörl,D., Fellinger,K., *et al.* (2015) DNA methylation requires a DNMT1 ubiquitin interacting motif (UIM) and histone ubiquitination. *Cell research*, **25**, 911–29.

13. Karg,E., Smets,M., Ryan,J., Forné,I., Qin,W., Mulholland,C.B., Kalideris,G., Imhof,A., Bultmann,S. and Leonhardt,H. (2017) Ubiquitome Analysis Reveals PCNA-Associated Factor 15 (PAF15) as a Specific Ubiquitination Target of UHRF1 in Embryonic Stem Cells. *Journal of molecular biology*, **429**, 3814–3824.

14. Ming, X., Zhang, Z., Zou, Z., Lv, C., Dong, Q., He, Q., Yi, Y., Li, Y., Wang, H. and Zhu, B. (2020) Kinetics and mechanisms of mitotic inheritance of DNA methylation and their roles in aging-associated methylome deterioration. *Cell research*, **30**, 980–996.

15. Nishiyama,A. and Nakanishi,M. (2021) Navigating the DNA methylation landscape of cancer. *Trends in Genetics*, **37**, 1012–1027.

16. Arita,K., Ariyoshi,M., Tochio,H., Nakamura,Y. and Shirakawa,M. (2008) Recognition of hemi-methylated DNA by the SRA protein UHRF1 by a base-flipping mechanism.

Nature, **455**, 818–821.

17. Avvakumov,G. V, Walker,J.R., Xue,S., Li,Y., Duan,S., Bronner,C., Arrowsmith,C.H. and Dhe-Paganon,S. (2008) Structural basis for recognition of hemi-methylated DNA by the SRA domain of human UHRF1. *Nature*, **455**, 822–825.

18. Hashimoto,H., Horton,J.R., Zhang,X., Bostick,M., Jacobsen,S.E. and Cheng,X. (2008) The SRA domain of UHRF1 flips 5-methylcytosine out of the DNA helix. *Nature*, **455**, 826–829.

19. Arita,K., Isogai,S., Oda,T., Unoki,M., Sugita,K., Sekiyama,N., Kuwata,K., Hamamoto,R., Tochio,H., Sato,M., *et al.* (2012) Recognition of modification status on a histone H3 tail by linked histone reader modules of the epigenetic regulator UHRF1. *Proceedings of the National Academy of Sciences*, **109**, 12950–12955.

20. Harrison,J.S., Cornett,E.M., Goldfarb,D., Darosa,P.A., Li,Z.M., Yan,F., Dickson,B.M., Guo,A.H., Cantu,D. V, Kaustov,L., *et al.* (2016) Hemi-methylated DNA regulates DNA methylation inheritance through allosteric activation of H3 ubiquitylation by UHRF1. *eLife*, **5**.

21. Unoki,M., Nishidate,T. and Nakamura,Y. (2004) ICBP90, an E2F-1 target, recruits HDAC1 and binds to methyl-CpG through its SRA domain. *Oncogene*, **23**, 7601–7610.

22. Achour,M., Fuhrmann,G., Alhosin,M., Rondé,P., Chataigneau,T., Mousli,M., Schini-Kerth,V.B. and Bronner,C. (2009) UHRF1 recruits the histone acetyltransferase Tip60 and controls its expression and activity. *Biochemical and Biophysical Research Communications*, **390**, 523–528.

23. Kim,J.K., Estève,P.O., Jacobsen,S.E. and Pradhan,S. (2009) UHRF1 binds G9a and participates in p21 transcriptional regulation in mammalian cells. *Nucleic Acids Research*, **37**, 493–505.

24. Hervouet,E., Lalier,L., Debien,E., Cheray,M., Geairon,A., Rogniaux,H., Loussouarn,D., Martin,S. a., Vallette,F.M. and Cartron,P.F. (2010) Disruption of Dnmt1/PCNA/UHRF1

interactions promotes tumorigenesis from human and mice glial cells. *PLoS ONE*, **5**.

25. Zhang,Z.-M., Rothbart,S.B., Allison,D.F., Cai,Q., Harrison,J.S., Li,L., Wang,Y.,
Strahl,B.D., Wang,G.G. and Song,J. (2015) An Allosteric Interaction Links USP7 to
Deubiquitination and Chromatin Targeting of UHRF1. *Cell reports*,
10.1016/j.celrep.2015.07.046.

26. Kori,S., Ferry,L., Matano,S., Jimenji,T., Kodera,N., Tsusaka,T., Matsumura,R., Oda,T.,
Sato,M., Dohmae,N., *et al.* (2019) Structure of the UHRF1 Tandem Tudor Domain
Bound to a Methylated Non-histone Protein, LIG1, Reveals Rules for Binding and
Regulation. *Structure*, **27**, 485-496.e7.

27. Ferry,L., Fournier,A., Tsusaka,T., Adelmant,G., Shimazu,T., Matano,S., Kirsh,O.,
Amouroux,R., Dohmae,N., Suzuki,T., *et al.* (2017) Methylation of DNA Ligase 1 by
G9a/GLP Recruits UHRF1 to Replicating DNA and Regulates DNA Methylation.
Molecular Cell, **67**, 550-565.e5.

28. Hahm,J.Y., Kim,J.-Y., Park,J.W., Kang,J.-Y., Kim,K.-B., Kim,S.-R., Cho,H. and Seo,S.-
B. (2019) Methylation of UHRF1 by SET7 is essential for DNA double-strand break
repair. *Nucleic Acids Research*, **47**, 184–196.

29. Li,Y., Zhang,Z., Chen,J., Liu,W., Lai,W., Liu,B., Li,X., Liu,L., Xu,S., Dong,Q., *et al.*
(2018) Stella safeguards the oocyte methylome by preventing de novo methylation
mediated by DNMT1. *Nature*, **564**, 136–140.

30. Zhao,S., Xu,J., Liu,S., Cui,K., Li,Z. and Liu,N. (2019) Dppa3 in pluripotency
maintenance of ES cells and early embryogenesis. *Journal of Cellular Biochemistry*,
120, 4794–4799.

31. Saitou,M., Barton,S.C. and Surani,M.A. (2002) A molecular programme for the
specification of germ cell fate in mice. *Nature, 418:6895*, **418**, 293–300.

32. Sato,M., Kimura,T., Kurokawa,K., Fujita,Y., Abe,K., Masuhara,M., Yasunaga,T.,
Ryo,A., Yamamoto,M. and Nakano,T. (2002) Identification of PGC7, a new gene

expressed specifically in preimplantation embryos and germ cells. *Mechanisms of development*, **113**, 91–94.

33. Payer,B., De Sousa Lopes,S.M.C., Barton,S.C., Lee,C., Saitou,M. and Surani,M.A. (2006) Generation of stella-GFP transgenic mice: a novel tool to study germ cell development. *Genesis*, **44**, 75–83.

34. Funaki,S., Nakamura,T., Nakatani,T., Umehara,H., Nakashima,H. and Nakano,T. (2014) Inhibition of maintenance DNA methylation by Stella. *Biochemical and Biophysical Research Communications*, **453**, 455–460.

35. Mulholland,C.B., Nishiyama,A., Ryan,J., Nakamura,R., Yiğit,M., Glück,I.M., Trummer,C., Qin,W., Bartoschek,M.D., Traube,F.R., *et al.* (2020) Recent evolution of a TET-controlled and DPPA3/STELLA-driven pathway of passive DNA demethylation in mammals. *Nature Communications*, **11**.

36. Du,W., Dong,Q., Zhang,Z., Liu,B., Zhou,T., Xu,R. ming, Wang,H., Zhu,B. and Li,Y. (2019) Stella protein facilitates DNA demethylation by disrupting the chromatin association of the RING finger–type E3 ubiquitin ligase UHRF1. *Journal of Biological Chemistry*, **294**, 8907–8917.

37. Jain,K., Fraser,C.S., Marunde,M.R., Parker,M.M., Sagum,C., Burg,J.M., Hall,N., Popova,I.K., Rodriguez,K.L., Vaidya,A., *et al.* (2020) Characterization of the plant homeodomain (PHD) reader family for their histone tail interactions. *Epigenetics & Chromatin, 13:1*, **13**, 1–11.

38. Musselman,C.A. and Kutateladze,T.G. (2011) Handpicking epigenetic marks with PHD fingers. *Nucleic acids research*, **39**, 9061–9071.

39. Orekhov,V.Y. and Jaravine,V.A. (2011) Analysis of non-uniformly sampled spectra with multi-dimensional decomposition. *Progress in Nuclear Magnetic Resonance Spectroscopy*, **59**, 271–292.

40. Ying,J., Delaglio,F., Torchia,D.A. and Bax,A. (2017) Sparse multidimensional iterative

lineshape-enhanced (SMILE) reconstruction of both non-uniformly sampled and conventional NMR data. *Journal of Biomolecular NMR*, **68**, 101–118.

41. Hyberts,S.G., Milbradt,A.G., Wagner,A.B., Arthanari,H. and Wagner,G. (2012) Application of iterative soft thresholding for fast reconstruction of NMR data non-uniformly sampled with multidimensional Poisson Gap scheduling. *Journal of Biomolecular NMR*, **52**, 315–327.

42. Delaglio,F., Grzesiek,S., Vuister,G.W., Zhu,G., Pfeifer,J. and Bax,A. (1995) NMRPipe: A multidimensional spectral processing system based on UNIX pipes. *Journal of Biomolecular NMR, 6:3*, **6**, 277–293.

43. Kobayashi,N., Iwahara,J., Koshiba,S., Tomizawa,T., Tochio,N., Güntert,P., Kigawa,T. and Yokoyama,S. (2007) KUJIRA, a package of integrated modules for systematic and interactive analysis of NMR data directed to high-throughput NMR structure studies. *Journal of Biomolecular NMR*, **39**, 31–52.

44. Johnson,B.A. and Blevins,R.A. (1994) NMR View: A computer program for the visualization and analysis of NMR data. *Journal of Biomolecular NMR, 4:5*, **4**, 603–614.

45. Kobayashi,N., Hattori,Y., Nagata,T., Shinya,S., Güntert,P., Kojima,C. and Fujiwara,T. (2018) Noise peak filtering in multi-dimensional NMR spectra using convolutional neural networks. *Bioinformatics*, **34**, 4300–4301.

46. Schmidt,E. and Güntert,P. (2012) A new algorithm for reliable and general NMR resonance assignment. *Journal of the American Chemical Society*, **134**, 12817–12829.

47. Shen,Y., Delaglio,F., Cornilescu,G. and Bax,A. (2009) TALOS+: A hybrid method for predicting protein backbone torsion angles from NMR chemical shifts. *Journal of Biomolecular NMR*, **44**, 213–223.

48. Güntert,P. and Buchner,L. (2015) Combined automated NOE assignment and structure calculation with CYANA. *Journal of Biomolecular NMR*, **62**, 453–471.

49. Case,D.A., Cheatham,T.E., Darden,T., Gohlke,H., Luo,R., Merz,K.M., Onufriev,A.,

Simmerling,C., Wang,B. and Woods,R.J. (2005) The Amber biomolecular simulation programs. *Journal of Computational Chemistry*, **26**, 1668–1688.

50. Clore,G.M., Driscoll,P.C., Gronenborn,A.M. and Wingfield,P.T. (1990) Analysis of the Backbone Dynamics of Interleukin-1β Using Two-Dimensional Inverse Detected Heteronuclear 15N-1H NMR Spectroscopy. *Biochemistry*, **29**, 7387–7401.

51. Mulholland,C.B., Smets,M., Schmidtmann,E., Leidescher,S., Markaki,Y., Hofweber,M., Qin,W., Manzo,M., Kremmer,E., Thanisch,K., *et al.* (2015) A modular open platform for systematic functional studies under physiological conditions. *Nucleic Acids Research*, **43**, e112.

52. Laible,G., Wolf,A., Dorn,R., Reuter,G., Nislow,C., Lebersorger,A., Popkin,D., Pillus,L. and Jenuwein,T. (1997) Mammalian homologues of the Polycomb-group gene Enhancer of zeste mediate gene silencing in Drosophila heterochromatin and at S. cerevisiae telomeres. *The EMBO journal*, **16**, 3219–3232.

53. Mátés,L., Chuah,M.K.L., Belay,E., Jerchow,B., Manoj,N., Acosta-Sanchez,A., Grzela,D.P., Schmitt,A., Becker,K., Matrai,J., *et al.* (2009) Molecular evolution of a novel hyperactive Sleeping Beauty transposase enables robust stable gene transfer in vertebrates. *Nature genetics*, **41**, 753–761.

54. Méndez,J. and Stillman,B. (2000) Chromatin association of human origin recognition complex, cdc6, and minichromosome maintenance proteins during the cell cycle: assembly of prereplication complexes in late mitosis. *Molecular and cellular biology*, **20**, 8602–8612.

55. Pabinger,S., Ernst,K., Pulverer,W., Kallmeyer,R., Valdes,A.M., Metrustry,S., Katic,D., Nuzzo,A., Kriegner,A., Vierlinger,K., *et al.* (2016) Analysis and Visualization Tool for Targeted Amplicon Bisulfite Sequencing on Ion Torrent Sequencers. *PLOS ONE*, **11**, e0160227.

56. Shen,Y. and Bax,A. (2013) Protein backbone and sidechain torsion angles predicted from

NMR chemical shifts using artificial neural networks. *Journal of Biomolecular NMR*, **56**, 227–241.

57. Clore,G.M., Driscoll,P.C., Gronenborn,A.M. and Wingfield,P.T. (1990) Analysis of the Backbone Dynamics of Interleukin-1β Using Two-Dimensional Inverse Detected Heteronuclear 15N-1H NMR Spectroscopy. *Biochemistry*, **29**, 7387–7401.

58. Sanchez,R. and Zhou,M.M. (2011) The PHD Finger: A Versatile Epigenome Reader. *Trends in biochemical sciences*, **36**, 364.

59. Rajakumara,E., Wang,Z., Ma,H., Hu,L., Chen,H., Lin,Y., Guo,R., Wu,F., Li,H., Lan,F., *et al.* (2011) PHD Finger Recognition of Unmodified Histone H3R2 Links UHRF1 to Regulation of Euchromatic Gene Expression. *Molecular Cell*, **43**, 275–284.

60. Sidhu,H. and Capalash,N. (2017) UHRF1: The key regulator of epigenetics and molecular target for cancer therapeutics. *Tumour biology : the journal of the International Society for Oncodevelopmental Biology and Medicine*, **39**, 1–11.

61. Yan,Q., Zhang,Y., Fang,X., Liu,B., Wong,T.L., Gong,L., Liu,S., Yu,D., Liu,M., Jiang,L., *et al.* (2021) PGC7 promotes tumor oncogenic dedifferentiation through remodeling DNA methylation pattern for key developmental transcription factors. *Cell Death and Differentiation*, **28**, 1955–1970.

2   **Figure 1: Characterization of the interaction between mUHRF1 and mDPPA3.**

3   **(A)** Schematic of the domain organization of mouse UHRF1 and DPPA3. L1-L4 indicate the

4   connecting linkers. **(B)** Protein sequence alignment of the UHRF1 pre-PHD/core-PHD

5   domains and DPPA3 residues 85-127 between different species with Jalview software. The

6   cyan and yellow dotted lines indicate the interaction interface identified in this study. **(C)** GST

1   pull-down experiments using mPHD and mDPPA3$_{76-128W}$. Bait proteins mean GST-fused

2   mUHRF1 that is immobilized on GST-beads. Proteins are stained by Coomassie Brilliant Blue.

3   **(D)** ITC measurements for the mPHD wild-type (WT)/mutant and mDPPA3$_{76-128W}$.

4   Superimposition of enthalpy change plots with standard deviations.

Figure 2: Structural analysis of mPHD-mDPPA3.

(A) Overall NMR structure of mPHD-mDPPA3. Pre-PHD, core-PHD, and mDPPA3 are shown

as orange, pink, and green cartoons, respectively. Zinc ions are depicted as gray-sphere models.

(B) Overlay of 20 NMR structures in which the structurally unconverged regions of mDPPA3 are indicated. The color scheme is the same as that in Figure 2A. **(C)** CD spectra of mPHD alone (pink), mDPPA3$_{76-128W}$ alone (green), and the mPHD in complex with mDPPA3$_{76-128W}$ (black). The sum of CD spectra of mPHD alone and mPDDA3$_{76-128W}$ alone is shown as gray. **(D)** Backbone $\{^1H\}$-^{15}N heteronuclear NOE of mPHD-mDPPA3. The het-NOE values were color-coded orange and pink for mPHD and green for mDPPA3$_{76-127}$ regions. **(E)** Electrostatic surface potential of mPHD. The surface colors red and blue represent negative and positive charges, respectively. mDPPA3 is depicted as a green cartoon with a stick model of the key residues that interact with mPHD.

Figure 3: Recognition of the [88]VRT[90] cassette of mDPPA3 by mPHD.

(A) Recognition of Val88 and Arg89 in mDPPA3. The mDPPA3 residues are shown as a green

stick model and transparent sphere model of Val88 methyl groups overlaid on the stick model.

mPHD residues that are involved in the recognition of Val88 of mDPPA3 are depicted as a pink

stick model superimposed on a transparent sphere model, and Asp339/Glu362 are shown as

pink stick models. The black dashed lines indicate the hydrogen bonds. **(B)** Recognition of

Thr90 of mDPPA3 showing a green stick model for hydrophobic residues of mPHD and a pink

stick model with a transparent sphere model. The hydrogen bond is indicated by the black

dashed line. **(C)** ITC measurements using mutants in the VRT cassette of mDPPA3 and mPHD

WT. Superimposition of enthalpy change plots with standard deviations. **(D)** GST pull-down assay to detect the interaction between full-length mUHRF1 and full-length GST-mDPPA3 wild-type (WT) and mutant proteins. **(E)** The upper panel shows the overall structure of mPHD (pre-PHD: orange surface, core-PHD: pink surface) bound to mDPPA3 (green stick model). The lower panel shows recognition of the ^{88}VRT^{90} cassette of mDPPA3. The red dashed lines indicate the hydrogen bonds. **(F)** The upper panel shows the overall structure of hPHD (pre-PHD: orange surface, core-PHD: gray surface) bound to H3 (cyan stick model) (PDB: 3ASL). The lower panel shows recognition of ^{1}ARTK^{4} of H3 by human PHD. **(G)** The upper panel shows the overall structure of hPHD (pre-PHD: orange surface, core-PHD: light-purple surface) bound to PAF15 (yellow stick model) (PDB: 6IIW). The lower panel shows recognition of ^{1}VRTK^{4} of PAF15 by hPHD. **(H)** Overlay of the N-terminus of H3, PAF15, and V88 of mDPPA3. The double arrow indicates the structural difference between the loops in the PHDs.

Figure 4: Contribution of the two helices of mDPPA3 for binding to mPHD.

(A) Interaction between the two helices of mDPPA3 and mPHD. The top-left panel displays

the overall structure of the mPHD-mDPPA3 complex in the cartoon mode. The top-right panel

shows the interaction between the αS1 of mDPPA3 and the hydrophobic patch of mPHD. The

bottom-left panel depicts the binding of Arg104 in mDPPA3 to the pre-PHD domain. The black

dashed lines indicate the hydrogen bonds. The bottom-right panel shows the interaction

between Cys341/Asp342 in core-PHD and Ile108/Arg111 on the αL2 of mDPPA3. The color

scheme is the same as that shown in Figure 3A. Transparent sphere models of the side chains

of hydrophobic residues were superimposed on the corresponding stick models. **(B)** ITC

measurements using mutants of α-helices of mDPPA3 and WT mPHD. Superimposition of

enthalpy change plots with standard deviations. **(C)** GST pull-down assay to detect the

interaction between full-length mUHRF1 and GST-mDPPA3 WT and mutants in αS1 and αL2.

Figure 5: The interaction interface between mDPPA3 and UHRF1 is important for the

inhibition of DNA methylation and UHRF1 chromatin binding in *Xenopus* egg extracts.

(A) Experimental design for functional analysis of mDPPA3 mutants using *Xenopus* egg

extracts. **(B)** Sperm chromatin was incubated with interphase *Xenopus* egg extracts

supplemented with buffer (+buffer), GST-mDPPA3 (+mDPPA3), or each mDPPA3 mutant.

Chromatin fractions were isolated and immunoblotted using the indicated antibodies.

Representative data from n = 3 independent experiments. **(C)** Sperm chromatin was added to

interphase egg extracts supplemented with radiolabeled S-[methyl-^{3}H]-adenosyl-L-methionine

and WT-GST-mDPPA3 or each mDPPA3 mutants. The efficiency of maintenance DNA

1   methylation was assessed by the incorporation of radio-labeled methyl groups from S-[methyl-

2   ^{3}H]-adenosyl-L-methionine (^{3}H-SAM) into DNA purified from egg extracts.

Figure 6: Effect of nuclear localization of mUHRF1 by mDPPA3 mutants in mouse ESCs

(A) Representative images illustrating the localization of UHRF1-GFP, and mouse and *B. taurus* DPPA3-mScarlet fusions in live U1G/D3KO + pSB-D3-mSC ESCs after doxycycline induction. DNA counterstain: SiR-DNA. Scale bar: 5 μm. **(B)** Subcellular distribution of UHRF1-GFP before and after DPPA3-mScarlet induction as determined using cell fractionation and western blot analyses. Cells were fractionated into cytoplasmic (C) and nuclear fractions (N). Anti-tubulin and anti-H3 blots were performed to identify the two

fractions. Ponceau S stained blots were used as a loading control. UHRF1-GFP was detected

using an anti-GFP antibody to determine its distribution in the C and N fractions. **(C)** DNA

methylation of LINE-1 repetitive elements were measured by targeted amplicon bisulfite

sequencing (TaBAseq) for each cell line (n=3). In the boxplot, the horizontal lines represent

the median values, box limits indicate the 25th and 75th percentiles as determined by R

software; whiskers extend to minimum and maximum values. A one-tailed t-test was performed

and p values are indicated.

Table 1.   NMR and refinement statistics for protein structures

	mPHD-mDPPA3 fusion
NMR distance and dihedral constraints	
Distance constraints	
Total NOE	1897
Intra-residue	364
Inter-residue	
Sequential ($\lvert i - j \rvert = 1$)	587
Medium-range ($\lvert i - j \rvert < 4$)	419
Long-range ($\lvert i - j \rvert > 5$)	527
Intermolecular	
Hydrogen bonds	15
Total dihedral angle restraints	100
φ	50
ψ	50
Structure statistics	
Violations (mean and s.d.)	
Distance constraints (Å)	0.057 +/- 0.006
Dihedral angle constraints (º)	1.577 +/- 0.078
Max. dihedral angle violation (º)	11.6
Max. distance constraint violation (Å)	0.38

Table 2. AMBER refinement statistics for complexes

	mPHD-mDPPA3
Structure statistics	
Energy (mean and s.d.)	
E-AMBER (kcal/mol)	-5314.54 +/- 19.75
NMR NOE	36.31 +/- 2.59
NMR Dihedral	6.92 +/- 0.87
Violations	
Number of distance constraints > 0.1 Å, > 0.5 Å	24.6 +/- 3.3, 0+/-0
Number of dihedral angle constraints > 2.5º, > 5º	15.9 +/- 2.1, 12.1+/-1.5
Max. dihedral angle violation (º)	31.5
Max. distance constraint violation (Å)	0.45
Ramachandran plot statistics (%)	
Residues in favored regions	88 +/- 1
Residues in allowed regions	11 +/- 2
Residues in outlier regions	1 +/- 0
Average pairwise r.m.s. deviation** (Å)	
Heavy	0.850
Backbone (Ca, N, CO)	0.396

[1]H-[1]H distance (exclusively derived from the NOE peaks) and dihedral (predicted by TALOS+) constraints for the AMBER refinements were identical to those applied to the CYANA calculations.

Ordered residues were automatically identified by Fit_Robot, residues on PHD domain: 9-15,18-60,62-68, and peptide: 86-112,127-129.

Supplementary Figure S1: GST pull-down assay for detecting the interaction between truncate version of GST fused mUHRF1 and mDPPA3.

(A) Schematic figure of domain structures of mUHRF1 that used for the pull-down assay **(B)** GST pull-down experiments using truncated forms of GST-mUHRF1 and C-terminal fragment of mDPPA3 (residues 61–150). Proteins are stained by Coomassie Brilliant Blue.

Supplementary Figure S2: ITC thermograms (upper) and plots of corrected heat values (lower) for the indicated binding experiments.

Supplementary Figure S3: NMR analysis.

(A) Overlay of ^{1}H-^{15}N HSQC spectra of 30 μM mPHD showing chemical shift changes upon

titration with mDPPA3$_{76–128W}$ of 0 μM (black), 15 μM (blue), 30 μM (green) and 60 μM (red).

Square regions inside the HSQC spectra were expanded (right panels). **(B)** Signal assignments

in ^{1}H-^{15}N HSQC spectra of mPHD in the free state (left) and in the complex state with

mDPPA3$_{76–128W}$ (right). **(C)** Weighted average chemical shift differences of ^{1}H and ^{15}N

resonances between free mPHD and mPHD in complex withmDPPA3$_{76–128W}$. Dashed line

represents mean plus 2 standard deviations. **(D)** Overlay of ^{1}H-^{15}N HSQC spectra of mPHD

wild type (Black) and L337A mutant (red).

Supplementary Figure S4: NMR analysis.

(A) Overlay of ^{1}H-^{15}N HSQC spectra of [^{15}N]-mPHD-mDPPA3 (Black) and [^{15}N]-mPHD

mixed with mDPPA3$_{76-128W}$ (red). **(B)** Overlay of ^{1}H-^{15}N HSQC spectra of [^{15}N]-mDPPA3$_{76-}$

$_{128W}$ in the free state (black) and in the complex with mPHD (red). **(C)** Weighted average

chemical shift differences of ^{1}H and ^{15}N resonances between free mDPPA3$_{76-128W}$ and

mDPPA3$_{76-128W}$ complex with mPHD. **(D)** Secondary structure prediction using the TALOS-

N program for mDPPA3 in the free state and in the complex with mPHD. **(E)** Overlay of ^{1}H-

^{15}N HSQC spectra of [^{15}N]-mPHD in the presence of mDPPA3$_{76-128W}$ and/or the H3 1–37W

peptide at molar ratios (mPHD:mDPPA3:H3) of 1:0:0 (black), 1:2:0 (red), 1:0:2 (blue) and

1:2:8 (green).

mPHD in the complex with mDPPA3
apo-mPHD (PDB: 6VFO)

Supplementary Figure S5: Structural comparison of mPHD and mDPPA3.

Overlay of mPHD moiety (colored pink, residues 304–372) in the complex with mDPPA3 on
apo-mPHD (PDB: 6VFO, colored yellow, residues 303–380) solved by NMR. Blue dotted line
indicates the loop that recognizes N-terminus of H3/PAF15.

2 **Supplementary Figure S6: *In vitro* ubiquitination assay of H3 using UHRF1 as a E3-ligase.**

3 C-terminal FLAG tagged-H3$_{1-37W}$ was ubiquitinated using in house purified E1, E2, and

4 UHRF1 (E3). The ubiquitinated H3 was detected using anti-FLAG antibody. Equimolar excess

5 of mDPPA3 WT or mutant to H3 was added.

Supplementary Figure S7 Phylogenetic tree of the UHRF1 and DPPA3.

Phylogenetic tree of UHRF1 and DPPA3 in different species rooting by midpoint. The pairwise
identities of UHRF1 and DPPA3 between mouse and different species were indicated, light
pink (low) and light blue (high). UHRF1 is conserved between species, but DPPA3 is not and
forms two clusters, relative conserved (>24.6%) and not conserved (<18.6%).