In situ visualization of opioid and cannabinoid drug effects using phosphosite-specific

GPCR antibodies

Sebastian Fritzwanker[1], Falko Nagel[2], Andrea Kliewer[1], Viviane Stammer[1], Stefan Schulz[1,2,*]

[1]Institut für Pharmakologie und Toxikologie, Universitätsklinikum Jena, Friedrich-Schiller-

Universität Jena, Drackendorfer Straße 1, D-07747 Jena, Germany

[2]7TM Antibodies GmbH, Hans-Knöll-Straße 6, D-07745 Jena, Germany

[*]**Corresponding author**

Emails: stefan.schulz@med.uni-jena.de

Abstract

G protein-coupled receptors (GPCRs) are important signal transducers that are

phosphorylated upon activation at intracellular serine and threonine residues. Although

antibodies that specifically recognize the phosphorylation state of GPCRs have been available

for many years, efficient immunolocalization of phosphorylated receptors in their tissues of

origin has not been possible. Here, we show that phosphorylation of receptors is highly

unstable during routine immunohistochemical procedures, requiring the use of appropriate

phosphatase inhibitors particular during tissue perfusion, post-fixation, and cryoprotection but

not during immunostaining of tissue sections. We provide proof of concept using

phosphorylation state-specific μ-opioid receptor (MOP) and cannabinoid receptor 1 (CB1)

antibodies. Indeed, three of four well-characterized phosphosite-specific MOP antibodies,

including pS375-MOP, pT376-MOP, and pT379-MOP, showed robust neuronal

immunostaining in brain and spinal cord sections of opioid-treated mice only after inclusion

of phosphatase inhibitors. We then extended this approach to the CB1 receptor and

demonstrated that one of three newly-generated phosphosite-specific CB1 antibodies, namely

pS425-CB1, showed striking staining of fibers and varicosities in brain slices from

cannabinoid-treated mice. Although subsequent experiments showed that phospho-CB1

immunostaining was less sensitive to phosphatases, we conclude that the use of phosphatase

inhibitors should always be considered in the development of immunohistochemical

procedures for new phosphosite-specific GPCR antibodies. In summary, we anticipate that

this improved protocol will facilitate the widespread use of phosphorylation state-specific

antibodies to monitor the activation of endogenous GPCRs under physiological and

pharmacological conditions. Our approach may also prove useful to confirm target

engagement of GPCR drug candidates in native tissues.

Introduction

Agonist-driven serine/threonine phosphorylation is a biologically and pharmacologically important process that primarily initiates desensitization and internalization of G protein-coupled receptors (GPCRs) [1-5]. Phosphorylation also increases the interaction of GPCRs with intracellular adapter proteins such as β-arrestins, which can trigger a second wave of signaling [1-5]. Thus, analysis of agonist-driven phosphorylation can provide valuable insights into the receptor activation state and ligand pharmacology. A widely used approach to analyze GPCR phosphorylation is the use of phosphosite-specific antibodies [6-9]. When available, such antibodies are valuable tools to elucidate the temporal dynamics of receptor phosphorylation, identify relevant kinases and phosphatases, and detect receptor activation using immunoblotting techniques [7, 10-15]. However, previous attempts using routine immunohistochemical approaches to unequivocally reveal agonist-induced phosphorylation of endogenous GPCRs in native tissues largely failed.

In intact cells, GPCR dephosphorylation is regulated in time and space, beginning immediately after receptor activation at the plasma membrane [16]. Indeed, phosphosite-specific antibodies in combination with siRNAs have led to the identification of distinct protein phosphatase 1 (PP1) and PP2 catalytic subunits as bona fide GPCR phosphatases [17-22]. For many receptors, dephosphorylation is complete within 10 to 30 minutes after agonist washout [17, 18, 21]. A notable exception is the agonist-induced phosphorylation of S341/S343 at the SST2 somatostatin receptor, which persists for considerably longer periods [21]. Consequently, activated SST2 receptors could be successfully localized using pS341/pS343-SST2 antibodies under routine immunohistochemical conditions [23]. However, this approach could not easily be reproduced for other SST2 sites or other GPCRs.

The ability to visualize activated and phosphorylated GPCRs in their tissues of origin

would provide important clues to the physiological and pharmacological regulation of

receptor activation. In particular, it would allow for distinction between currently-activated

and resting GPCR populations in the context of particular physiological or behavioral

conditions. We were therefore motivated to develop and validate immunohistochemical

fixation and staining procedures that can be universally applied to prototypical GPCRs. Our

improved protocol will facilitate the widespread application of phosphosite-specific

antibodies as GPCR activation sensors in academic and pharmaceutical research.

Results

Dephosphorylation of GPCRs occurs rapidly in intact cells and is mediated by the protein

phosphatases PP1 and PP2 [16]. Serine/threonine phosphatases are known to be activated during

cell lysis and tissue fixation procedures. We therefore tested the inclusion of appropriate

protein phosphatase inhibitors (PPIs) during tissue fixation and immunohistochemical

staining. For this purpose, mice were treated with either methadone or saline, perfusion fixed,

and slices of brain and spinal cord were stained with phosphosite-specific μ-opioid receptor

(MOP) or phosphorylation-independent np-MOP antibodies. As expected, staining using the

np-MOP antibody was present in the superficial layers of the spinal cord, regardless of drug

treatment or the presence of phosphatase inhibitors (**Fig. 1**). In contrast, immunostaining for

pT379-MOP was only detected in methadone-treated animals when a cocktail of appropriate

phosphatase inhibitors was present during both fixation and staining procedures (**Fig. 1**). Four

unique serine and threonine residues are phosphorylated in MOP in response to full agonists [8,

15]. Phosphosite-specific antibodies have been generated for all of these sites, which work

equally well in immunoblot applications [7, 24, 25]. We found that three out of four, namely

pS375-MOP, pT376-MOP and pT379-MOP, but not pT370-MOP, are also well suited for

immunohistochemical staining of phosphorylated MOP under these conditions (**Fig. 2 upper**

panel). Neither p-MOP nor np-MOP staining was apparent in MOP knockout mice (**Fig. 2**

lower panel). To further substantiate the specific binding of phospho-MOP antibodies, we

performed peptide neutralization controls. When pS375-MOP or pT379-MOP antibodies were

incubated with an excess of their respective immunizing peptides containing S375 or T379 in

phosphorylated form, immunostaining was completely absent in the superficial layers of the

spinal cord (**Fig. 3**). In contrast, immunostaining was virtually unaffected by the addition of

the corresponding non-phosphorylated peptides, indicating unequivocal detection of agonist-

dependent S375 and T379 phosphorylation of MOP in tissues (**Fig. 3**).

The antagonist naloxone drives MOP into an inactive state, preventing agonist-

induced phosphorylation. When mice were treated with the selective MOP agonist fentanyl,

we found the typical staining pattern for pS375-MOP, pT376-MOP and pT379-MOP as

evidence of MOP activation. In contrast, this staining was not detected in animals injected

with naloxone immediately prior to fentanyl challenge, indicating that MOP activation was

blocked by the antagonist (**Fig. 4**). Collectively, our results support the notion that agonist-

induced MOP phosphorylation is specifically detected in brain sections under these

conditions. Notably, agonist-driven changes in phospho-immunofluorescence staining of

MOP were robust enough to allow for quantification even in relatively small groups of

animals (Supplemental Figure S1). Given the considerable costs involved in using PPIs, we

then tested whether their presence is necessary during both fixation and staining procedures.

The results depicted in **Figure 5**, clearly revealed that inclusion of PPIs during tissues fixation

is an absolute requirement. They also show that after fixation, it is not obligatory to add PPIs

to each solution for successful immunostaining (**Fig. 5**).

In an effort to perform dual immunofluorescence labeling, we combined the rabbit

pT376-MOP antibody (depicted in green, **Fig. 6**) with guinea pig np-MOP antibody (depicted

in magenta, **Fig. 6**). This procedure allows to discern populations of non-phosphorylated

MOP from activated and phosphorylated MOP, which dramatically increases in the

superficial layer of the spinal cord after administration of fentanyl or methadone (depicted in

white in overlay images, **Fig. 6**). Next, we compared the immunostaining patterns obtained

for phospho-MOP with total MOP. While there was good overlap in some brain regions,

including habenula, fasciculus retroflex (**Fig. 7**), unexpected low or even absent pS375-MOP,

pT376-MOP, nor pT379-MOP immunostaining was observed in the nucleus accumbens and

caudate putamen, despite high levels of MOP expression (**Fig. 7**). These results suggest that

region-specific MOP activation and phosphorylation patterns exist, which can now be

visualized for the first time.

The final set of experiments was designed to extend this new approach to another

prototypical GPCR namely the CB1 cannabinoid receptor. To this end, three newly-generated

phosphosite-specific CB1 antibodies including pS425-CB1, pS429-CB1 and pS441-CB1 were

tested under the new assay conditions. In fact, the pS425-CB1 antibody produced staining in

hippocampus and cortex closely resembling that of the np-CB1 antibody only when mice

were treated with CP-55940 [26] (**Fig. 8**). Notably, pS425-CB1 immunostaining was much less

prominent in untreated animals but not absent (**Fig. 8**). Staining in both untreated and

cannabinoid-treated animals was completely abolished when the p425-CB1 antibody was

neutralized with its immunizing peptide (**Fig. 8**). Prior administration of the antagonist

AM251 reduced CP-55940-induced phospho-CB1 immunostaining to that seen in untreated

mice (**Fig. 8**). To corroborate these findings using an independent biochemical approach,

groups of three animals each were treated with either vehicle or CP55940. CB1 receptors

were then immunoprecipitated using a goat np-CB1 antibody and CB1 phosphorylation was

determined using a rabbit pS425-CB1 antibody (**Fig. 9**). To confirm equal loading, the blot

was stripped and total CB1 was detected using a rabbit np-CB1 antibody (**Fig. 9**). The results

depicted in Figure 9, reveal the existence of basal pS425 phosphorylation of CB1 that is

strongly increased after administration of the high-efficacy agonist CP55940 (**Fig. 9**). When

CB1 immunoprecipitates were subjected to λ-phosphatase digestion, both basal and agonist-

driven CB1 phosphorylation were completely diminished, whereas total CB1 was still

detectable. Collectively, these findings support the conclusion that CB1 phosphorylation can

be specifically detected. We then tested whether the presence of PPIs during tissues fixation is

an absolute requirement for successful pS425-CB1 immunostaining. Surprisingly, we found

that CB1 phosphorylation appears to be much less sensitive to in situ phosphatase digestion

than MOP phosphorylation (Supplemental Fig. S2). Finally, we used a combination of goat

np-CB1 and rabbit pS425-CB1 antibodies to perform dual immunofluorescence labeling. At

higher magnification it became apparent that the p425-CB1 antibody exclusively labelled

fibers and terminals similar to that seen with the np-CB1 antibody (**Fig. 10**). Interestingly,

overlay images revealed that in CP-55940-treated mice activated and phosphorylated CB1

receptors were densely concentrated in varicosities (depicted in white, **Fig. 10**), whereas CB1

was detected in fibers appears to reside in the non-phosphorylated state (depicted in magenta,

Fig. 10).

Discussion

GPCRs are privileged targets for small molecule drugs in almost all therapeutic areas [27].

However, to date, the visualization of therapeutic drug effects on endogenously expressed

GPCRs in native tissues has remained a major challenge. Activation of GPCRs by their

endogenous ligands or exogenous agonists results in conformational changes that are

recognized by a family of kinases termed G protein-coupled receptor kinases (GRKs) [1-5]. The

unique ability of GRKs to recognize activated receptors results in agonist-dependent

phosphorylation at intracellular serine and threonine residues [1-5]. Thus, analysis of agonist-

driven phosphorylation of GPCRs can provide valuable insights into receptor activation.

Antibodies that specifically recognize the GPCR phosphorylation state have been available in

some cases over the last 20 years. Indeed, phosphosite-specific GPCR antibodies have proven

useful to detect receptor activation and to profile the pharmacological properties of new

ligands [12-15]. To date, however, their utility has been largely limited to immunoblotting

approaches using heterologous receptor expression systems, while detection of

phosphorylated GPCRs from tissue lysates has been mostly unsuccessful, due to their low

abundance *in vivo*.

Here, we have systematically developed an improved immunohistochemical staining

procedure that allows for visualization of opioid drug effects in the mouse brain *in vivo*. We

also show that this approach can be easily extended to other prototypical GPCRs. Key to this

new protocol was the observation that receptor phosphorylation is highly unstable during

routine immunohistochemistry and that for many GPCRs inclusion of protein phosphatase

inhibitors may be an absolute requirement during perfusion, post-fixation and cryoprotection

of tissues. Several lines of evidence suggest that our approach facilitates the localization of

bona fide phosphorylated GPCRs. First, phospho-GPCR immunostaining strongly increased

when animals had been treated with agonist. Conversely, immunostaining was not detected

when agonist action was blocked by antagonist. Second, robust phospho-GPCR

immunostaining was only apparent when appropriate protein phosphatase inhibitors were

included in all steps of fixation and staining procedures. Third, phospho-GPCR

immunostaining was completely neutralized by preincubation with cognate phosphorylated

peptides but not with the corresponding non-phosphorylated peptides.

In the present study, we had access to four different phospho-MOP and three different

phospho-CB1 antibodies which had previously been validated by immunoblotting techniques.

We observed that three out of four antibodies for MOP and one out of three antibodies for

CB1 produced specific phospho-immunostaining in tissues suggesting that each phosphosite-

specific GPCR antibody has to be validated individually for immunohistochemical

applications. Moreover, we observed that phospho-MOP staining was critically dependent on

the presence of PPIs, whereas CB1 phosphorylation appeared to be more stable in tissues.

Nevertheless, we conclude that use of PPIs should always be considered in the development

of immunohistochemical procedures for new phosphosite-specific GPCR antibodies.

It is well known that endogenous GPCRs are notoriously difficult to detect in tissues

by immunohistochemical staining. Here, we not only establish a protocol for in situ

visualization of activated and phosphorylated GPCRs. We also demonstrate that phospho-

immunostaining can disclose previously unappreciated functional information not seen with

phosphorylation-independent GPCR antibodies. One example is the striking mismatch

between total and phospho-immunostaining for MOP in caudate putamen. This mismatch

cannot be explained be low detection level of phospho-MOP staining as many other brain

regions with comparable total MOP revealed prominent phospho-MOP. The most likely

explanation for the relative lack of phospho-MOP staining in caudate putamen is the low or

absent expression of relevant GRK family members [32]. Notably, lack of MOP

phosphorylation in some brain regions relative to others could have profound consequences

on MOP desensitization and may explain well-known differences in tolerance development to

different opioid effects. Another example is the different subcellular localization of

phosphorylated and non-phosphorylated CB1 receptors. While total CB1 appears to be evenly

distributed along fibers and varicosities, phosphorylated CB1 is particular enriched in

terminals indicating compartmentalization of GRK-mediated phosphorylation.

In summary, we provide a proof of concept for using phosphorylation state-specific

antibodies as sensors to elucidate GPCR activation in native tissues. Our approach not only

facilitates the visualization of the effects of agonists and antagonists in preclinical animal

studies. It also allows for elucidation of different cellular and subcellular activation patterns in

response to external stimuli or drug administration. Therefore, phosphorylation state-specific

antibodies are a new class of markers for GPCR activation that are likely to find widespread

application in academic and pharmaceutical research.

Figure legends

Fig. 1. Comparison of phospho-MOP immunohistochemistry in the presence or absence of

protein phosphatase inhibitors. Animals were either treated with methadone or saline for 30

min, transcardially perfused, fixed and stained in the presence (+) or absence (-) of protein

phosphatase inhibitors (PPI). Shown are confocal images of coronal sections of the spinal

cord stained with pT379-MOP or np-MOP antibody. Note that PPIs need to be present during

both fixation and staining procedures to obtain agonist-induced phospho-MOP

immunostaining. Scale bar = 250 µm.

Fig. 2. Immunohistochemical staining of agonist-induced MOP phosphorylation in the mouse

spinal cord. Animals were treated with methadone or saline for 30 min, transcardially

perfused, fixed and stained in the presence of PPIs. Shown are confocal images of coronal

sections of the spinal cord stained with phosphosite-specific antibodies pT370-MOP, pS375-

MOP, pT376-MOP and pT379-MOP, or phosphorylation independent np-MOP antibody.

Note that pS375-MOP, pT376-MOP and pT379-MOP, but not pT370-MOP, revealed an

agonist-dependent MOP phosphorylation immunostaining in a pattern closely resembling that

of np-MOP immunostaining. Scale bar = 250 µm.

Fig. 3. Peptide neutralization of agonist-induced MOP phosphorylation staining in the mouse

spinal cord. Animals were treated with methadone for 30 min, transcardially perfused, fixed

and stained in the presence of PPIs. Shown are confocal images of coronal sections of the

spinal cord stained with phosphosite-specific antibodies pS375-MOP or pT379-MOP in the

presence or absence of their immunizing peptides containing S375 or T379 in phosphorylated

form (p-Peptide) or the corresponding non-phosphorylated peptide (np-Peptide). Note that

phospho-MOP immunostaining was completely neutralized by excess of phosphorylated

peptide but not of non-phosphorylated peptide. Scale bar = 250 µm.

Fig. 4. Antagonist block of phospho-MOP immunostaining. Animals were either treated with

saline or fentanyl for 15 min. Where indicated, animals were pretreated with naloxone for 10

min followed by 15-min fentanyl treatment. Animals were then transcardially perfused, fixed

and stained in the presence of PPIs. Shown are confocal images of coronal sections of the

spinal cord stained with phosphosite-specific antibodies pS375-MOP, pT376-MOP or pT379-

MOP or phosphorylation independent np-MOP antibody. Note that agonist-induced phospho-

MOP immunostaining is diminished by antagonist treatment. Scale bar = 250 µm.

Fig. 5. Comparison of phospho-MOP detection after staining in the presence or absence

of PPIs. Animals were treated with methadone for 30 min, transcardially perfused and fixed

in the presence (+) or absence (-) of PPIs and also stained in the presence (+) or absence (-) of

PPIs. Shown are confocal images of coronal sections of the spinal cord stained with pS375-

MOP, pT379-MOP or np-MOP antibody. Note that PPIs need to be present during the whole

fixation procedures but not during the staining protocol to obtain agonist-induced phospho-

MOP immunostaining. Scale bar = 250 µm.

Fig. 6. Double staining of agonist-induced p-MOP and np-MOP in mouse brain. Animals

were either treated with NaCl (30 min), Fentanyl (15 min) or Methadone (30 min). Animals

were then transcardially perfused, fixed and stained. Shown are confocal images of coronal

sections of the spinal cord stained with phosphosite-specific antibody pT376-MOP (green)

and phosphorylation independent np-MOP antibody (magenta). Note that there is a strong

overlap of np-MOP and pT376-MOP in the merged panels (white) in agonist treated animals

compared to NaCl treated mice. Scale bar = 250 µm.

Fig. 7. Agonist-induced phospho-MOP immunostaining in mouse brain. Animals were treated

with methadone for 30 min, transcardially perfused, fixed and stained in the presence of PPIs.

Shown are confocal images of coronal brain sections stained with phosphosite-specific

antibodies pS375-MOP, pT376-MOP or pT379-MOP or phosphorylation independent np-

MOP antibody. Note that all three phosphosite-specific antibodies detected agonist-induced

MOP phosphorylation in the medial habenula and fasciculus retroflexus but not in nucleus

accumbens or caudate putamen, brain regions known to be rich in MOP receptors as shown

by np-MOP antibody staining. Scale bar = 250 µm.

Fig. 8. Agonist-induced phospho-CB1 immunostaining in mouse brain. Animals were

either treated with Vehicle or the CB1-agonist CP-55940 for 30 min. Where indicated,

animals were pretreated with the CB1-antagonistAM251 for 45 min followed by 30 min CP-

55940 treatment. Animals were then transcardially perfused, fixed and stained in the presence

of PPIs. Shown are confocal images of coronal brain sections stained with phosphosite-

specific antibody pS425-CB1, in the presence or absence of the p-Peptide containing S425 in

phosphorylated form, or phosphorylation independent np-CB1 antibody. Note that agonist-

induced phopho-CB1 immunostaining is diminished by antagonist treatment. Note that

phospho-CB1 immunostaining was completely neutralized by excess of phosphorylated

peptide. Scale bar = 1000 µm and 250 µm.

Fig. 9. Immunoblotting of phosphorylated CB1 in agonist treated mice. Six different

wildtype animals were used. Number 1 to 3 were treated with vehicle and Number 4 to 6

with CP-55940 for 30 min. Organs were removed rapidly and frozen in liquid nitrogen After

tissue preparations, CB1 receptors were extracted using goat np-CB1 bound to protein A

beads. After blotting, the membranes were treated with the phosphosite-specific rabbit

antibody pS425-CB1, stripped (dashed line) and reprobed with rabbit phosphorylation-

independent anti-CB1. Note that there is some CB1-phosphorylation constitutive signal in

vehicle-treated mice. Molecular mass on the left in kDa.

Fig. 10. Double staining of agonist-induced p-CB1 and np-CB1 in mouse brain. Animals

were either treated with Vehicle (A) or CP-55940 (B, C) for 30 min. Animals were then

transcardially perfused, fixed and stained. Shown are confocal images of coronal brain

sections of cortex (CTX) and hippocampal formation (HPF) stained with phosphosite-specific

antibody pS425-CB1 (green) and phosphorylation independent np-CB1 antibody (magenta).

Note that there is a strong overlap of np-CB1 and pS425-CB1 in the merged panels (white) in

CP-55940 treated animals (B) compared to vehicle treated mice. Note that in cortical neurons,

only CB1 located in varicosities undergoes agonist-induced phosphorylation. Scale bar =

1000 µm, 100 µm and 2 µm.

Methods

Antibodies

<u>Primary Antibodies:</u>

The phosphorylation state-specific rabbit MOP antibodies pT370-MOP (7TM0319B), pS375-

MOP (7TM0319C), pT376-MOP (7TM0319D), pT379-MOP (7TM0319E), the

phosphorylation state-specific rabbit CB1 antibody pS425-CB1 (7TM0056A) as well as the

phosphorylation-independent rabbit antibody np-MOP (7TM0319N) and phosphorylation-

independent guinea pig antibody np-MOP (7TM0319N-GP) were provided by 7TM

Antibodies (www.7tmantibodies.com). In addition, we used rabbit np-MOP from abcam

(ab134054), rabbit np-CB1 from Cayman (1000659) and goat np-CB1 from Frontier Institute

(MFSR100610).

<u>Secondary Antibodies:</u>

We used a biotinylated AB from Jackson Immuno Research (711-065-152, anti-rabbit) and

the respective streptavidine conjugates from ThermoFisher (S21381, Alexa Fluor^{TM} 555

conjugate) and (S11223, Alexa Fluor^{TM} 488 conjugate) for the amplified staining procedure.

For the direct stainings we used Cy^{TM}3 (706-165-148, anti-guinea pig) and Cy^{TM}3 (705-165-

147, anti-goat), both obtained from Jackson ImmunoResearch.

Animals

Mice (C57BL/6J obtained from JAX^{TM}) were housed 2–5 per cage under a 12-hr light-dark

cycle with ad libitum access to food and water. All animal experiments were approved by

Thuringian state authorities and performed in accordance with European Commission

regulations for the care and use of laboratory animals. Our study is reported in accordance

with ARRIVE guidelines [28]. In all experiments, male and female mice aged 8–30 weeks

between 25 and 35 g body weight were used.

Drugs and routes of administration

All drugs were freshly prepared prior to use and were injected subcutaneously (s.c.) or

intraperitoneally (i.p.) in unanaesthetized mice at a volume of 10 μl/g bodyweight. Opioid

drugs were diluted in 0.9% (w/v) saline and cannabinoid drugs in vehicle solution, containing

13.2% ethanol and 0.25% Tween80 in 0.9% saline, for injections. Drugs were obtained and

used as follows: fentanyl citrate (0.3 mg/kg for 15 min; s.c.) (B. Braun 06900650),

levomethadone hydrochloride (15 mg/kg for 30 min; s.c.) (Sanofi-Aventis 07480196),

naloxone (2 mg/kg for 25 min; s.c.) (Ratiopharm 04788930), CP-55940 (0.75 mg/kg for 30

min; i.p.) (Sigma-Aldrich C1112) and AM251 (3 mg/kg for 45 min; i.p.) (MedChemExpress

183232-66-8). A total of 35 mice (n=4-6 per treatment condition) were used.

Immunohistochemistry

Mice were deeply anaesthetized with isoflurane (CP-Pharma 4001404) and then subjected to a

transcardial perfusion with calcium-free Tyrode's solution containing protein phosphatase-

inhibitors (+PPIs) (1 tablet PhosSTOP per 10 ml) (Roche 04906845001) followed by

Zamboni's fixative containing 4% paraformaldehyde and 0.2% picric acid in 0.1 M phosphate

buffer pH 7.4 +PPIs. Brains and spinal cords were rapidly dissected and postfixed in the same

fixative for 4 h at room temperature. The tissue was cryoprotected by immersion in 10%

sucrose +PPIs followed by 30% sucrose +PPIs for 48 h at 4 °C before sectioning using a

freezing microtome. Tissue was cut into 40 μm sections. Free-floating sections were washed

in PBS with or without (+/-) PPIs (depending on the experiment and investigated GPCR) and

incubated in methanol containing 0.3% H_2O_2 for 30 min. After washing in PBS-T (PBS +

0,3% Tween®20), the sections were blocked in PBS containing 0.3% Triton X-100, +/- PPIs

and 10% NGS for 2 h. Subsequently, the sections were incubated with primary antibody in

PBS containing 0.3% Triton X-100, 2% NGS +/- PPI overnight at 4°C. Where indicated,

primary antibodies were preincubated with 1 µg/ml of their cognate phospho-peptide or the

corresponding non-phospho-peptide for 1 h at room temperature. When necessary, staining of

primary antibody was detected using the biotin amplification procedure as described [29, 30].

Briefly, tissue sections were transferred to biotinylated donkey anti-rabbit IgG (1:300 in PBS

containing 0.3% Triton X-100, +/-PPIs and 10% NGS) for 2 h, washed in PBS-T +/-s PPIs

and then incubated in AB solution (reagents from the Vector ABC kit; 25 µl A and 25 µl B in

10 ml PBS+0.3% Triton X-100 +/- PPIs) for 60 min, washed again in PBS-T +/- PPIs and

transferred to biotinylated tyramine (BT) solution (BT was prepared as described by Adams

[31], 5 µl BT+0.01% H_2O_2 in 1 ml PBS+0.3% Triton X-100 and +/- PPIs) for 20 min, followed

by a final incubation step in streptavidin-AlexaFluor555 conjugate 1:400 in PBS+0.3% Triton

X-100 containing 10% NDS +/- PPI) overnight at 4°C. For double-staining procedures, the

primary np-AB was incubated overnight as indicated and after several washing steps on the

following day conjugated to the secondary AB, again overnight. Sections were then mounted

onto SuperFrost Plus glass slides (ThermoFisher 15438060) and cover slipped with Eukitt

(ORSAtec). Specimens were examined using a Zeiss LSM 900 laser scanning confocal

microscope equipped with ZEN software for image analysis.

Western Blot and immunoprecipitation

Wildtyp mice were treated with agonist or vehicle. Mice were anesthetized with isoflurane,

killed by cervical dislocation, and brains were quickly dissected, excluding the cerebellum.

Brain samples were immediately frozen in liquid nitrogen. Brains were transferred to ice-cold

detergent buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 5 mM EDTA, 1% Nonidet P-40,

0.5% sodium deoxycholate, 0.1% sodium dodecyl sulfate (SDS), containing protease and

phosphatase inhibitors), homogenized, and centrifuged at 14,000 × g for 30 min at 4°C. The

supernatant was then immunoprecipitated with the polyclonal goat np-CB1 antibody bound to

protein A-agarose beads (20333, ThermoScientific, Germany) for 90 min at 4°C. Proteins

were eluted from the beads with SDS sample buffer for 25 min at 43°C and then resolved on

8% SDS-polyacrylamide gels. After electroblotting, membranes were incubated with pS425-

MOP antibody, followed by detection using a chemiluminescence detection system. Blots

were subsequently stripped and incubated again with the rabbit np-CB1 antibody to confirm

equal loading of the gels.

Data availability

All data supporting the findings of this study are available within the article and its

supplementary information files. Additional information, relevant data and unique biological

materials will be available from the corresponding author upon reasonable request. Source

data are provided with this paper.

References

1. Drube, J. et al. GPCR kinase knockout cells reveal the impact of individual GRKs on arrestin binding and GPCR regulation. *Nat Commun* **13**, 540 (2022).

2. Gurevich, E.V., Tesmer, J.J., Mushegian, A. & Gurevich, V.V. G protein-coupled receptor kinases: more than just kinases and not only for GPCRs. *Pharmacol Ther* **133**, 40-69 (2012).

3. Liggett, S.B. Phosphorylation barcoding as a mechanism of directing GPCR signaling. *Sci Signal* **4**, pe36 (2011).

4. Premont, R.T. & Gainetdinov, R.R. Physiological roles of G protein-coupled receptor kinases and arrestins. *Annu Rev Physiol* **69**, 511-534 (2007).

5. Tobin, A.B., Butcher, A.J. & Kong, K.C. Location, location, location...site-specific GPCR phosphorylation offers a mechanism for cell-type-specific signalling. *Trends Pharmacol Sci* **29**, 413-420 (2008).

6. Doll, C. et al. Agonist-selective patterns of micro-opioid receptor phosphorylation revealed by phosphosite-specific antibodies. *Br J Pharmacol* **164**, 298-307 (2011).

7. Grecksch, G. et al. Analgesic tolerance to high-efficacy agonists but not to morphine is diminished in phosphorylation-deficient S375A mu-opioid receptor knock-in mice. *J Neurosci* **31**, 13890-13896 (2011).

8. Just, S. et al. Differentiation of opioid drug effects by hierarchical multi-site phosphorylation. *Mol Pharmacol* **83**, 633-639 (2013).

9. Prihandoko, R., Bradley, S.J., Tobin, A.B. & Butcher, A.J. Determination of GPCR Phosphorylation Status: Establishing a Phosphorylation Barcode. *Curr Protoc Pharmacol* **69**, 2 13 11-12 13 26 (2015).

430   10.   Divorty, N. et al. Agonist-induced phosphorylation of orthologues of the orphan

431   receptor GPR35 functions as an activation sensor. *J Biol Chem*, 101655 (2022).

432   11.   Illing, S., Mann, A. & Schulz, S. Heterologous regulation of agonist-independent mu-

433   opioid receptor phosphorylation by protein kinase C. *Br J Pharmacol* **171**, 1330-1340

434   (2014).

435   12.   Mann, A. et al. New phosphosite-specific antibodies to unravel the role of GRK

436   phosphorylation in dopamine D2 receptor regulation and signaling. *Sci Rep* **11**, 8288

437   (2021).

438   13.   Mann, A. et al. Agonist-induced phosphorylation bar code and differential post-

439   activation signaling of the delta opioid receptor revealed by phosphosite-specific

440   antibodies. *Sci Rep* **10**, 8585 (2020).

441   14.   Mann, A. et al. Agonist-selective NOP receptor phosphorylation correlates in vitro and

442   in vivo and reveals differential post-activation signaling by chemically diverse

443   agonists. *Sci Signal* **12** (2019).

444   15.   Miess, E. et al. Multisite phosphorylation is required for sustained interaction with

445   GRKs and arrestins during rapid mu-opioid receptor desensitization. *Sci Signal* **11**

446   (2018).

447   16.   Kliewer, A., Reinscheid, R.K. & Schulz, S. Emerging Paradigms of G Protein-

448   Coupled Receptor Dephosphorylation. *Trends Pharmacol Sci* **38**, 621-636 (2017).

449   17.   Doll, C. et al. Deciphering micro-opioid receptor phosphorylation and

450   dephosphorylation in HEK293 cells. *Br J Pharmacol* **167**, 1259-1270 (2012).

451   18.   Lehmann, A., Kliewer, A., Gunther, T., Nagel, F. & Schulz, S. Identification of

452   Phosphorylation Sites Regulating sst3 Somatostatin Receptor Trafficking. *Mol*

453   *Endocrinol* **30**, 645-659 (2016).

454   19.   Lehmann, A., Kliewer, A., Martens, J.C., Nagel, F. & Schulz, S. Carboxyl-terminal

455   receptor domains control the differential dephosphorylation of somatostatin receptors

456   by protein phosphatase 1 isoforms. *PLoS One* **9**, e91526 (2014).

457   20.   Petrich, A. et al. Phosphorylation of threonine 333 regulates trafficking of the human

458   sst5 somatostatin receptor. *Mol Endocrinol* **27**, 671-682 (2013).

459   21.   Poll, F., Doll, C. & Schulz, S. Rapid dephosphorylation of G protein-coupled

460   receptors by protein phosphatase 1beta is required for termination of beta-arrestin-

461   dependent signaling. *J Biol Chem* **286**, 32931-32936 (2011).

462   22.   Tran, T.M. et al. Characterization of beta2-adrenergic receptor dephosphorylation:

463   Comparison with the rate of resensitization. *Mol Pharmacol* **71**, 47-60 (2007).

464   23.   Waser, B. et al. Phosphorylation of sst2 receptors in neuroendocrine tumors after

465   octreotide treatment of patients. *Am J Pathol* **180**, 1942-1949 (2012).

466   24.   Fritzwanker, S. et al. HA-MOP knockin mice express the canonical micro-opioid

467   receptor but lack detectable splice variants. *Commun Biol* **4**, 1070 (2021).

468   25.   Kliewer, A. et al. Phosphorylation-deficient G-protein-biased mu-opioid receptors

469   improve analgesia and diminish tolerance but worsen opioid side effects. *Nat Commun*

470   **10**, 367 (2019).

471   26.   Davis, M.I. et al. The cannabinoid-1 receptor is abundantly expressed in striatal

472   striosomes and striosome-dendron bouquets of the substantia nigra. *PLoS One* **13**,

473   e0191436 (2018).

474   27.   Santos, R. et al. A comprehensive map of molecular drug targets. *Nat Rev Drug*

475   *Discov* **16**, 19-34 (2017).

476   28.   Percie du Sert, N. et al. The ARRIVE guidelines 2.0: Updated guidelines for reporting

477   animal research. *PLoS Biol* **18**, e3000410 (2020).

29.   Schreff, M. et al. Distribution, targeting, and internalization of the sst4 somatostatin

receptor in rat brain. *J Neurosci* **20**, 3785-3797 (2000).

30.   Schulz, S. et al. Immunolocalization of two mu-opioid receptor isoforms (MOR1 and

MOR1B) in the rat central nervous system. *Neuroscience* **82**, 613-622 (1998).

31.   Adams, J.C. Biotin amplification of biotin and horseradish peroxidase signals in

histochemical stains. *J Histochem Cytochem* **40**, 1457-1463 (1992).

32.   Allen Brain Atlas, https://atlas.brain-map.org

Acknowledgements: We want to thank Svetlana Würl and Monique Brendel for expert

technical assistance. This work was supported by the European Regional Development Fund

(EFRE) and the Free State of Thuringia (grant: 2020 FE 0146) to 7TM Antibodies GmbH.

Author contributions: S.S. conceived and initiated the project and designed all experiments

with S.F. . S.F. and A.K. performed and analyzed immunohistochemistry. F.N. affinity-

purified all phosphosite-specific antibodies used in this study. The manuscript was written

and revised by S.S. and S.F. with input from other authors.

Competing interests: S.S. is the founder and scientific advisor of 7TM Antibodies GmbH,

Jena, Germany. F.N. is an employee of 7TM Antibodies. All other authors declare no

competing interests.

In situ visualization of opioid and cannabinoid drug effects using phosphosite-specific

GPCR antibodies

Sebastian Fritzwanker[1], Falko Nagel[2], Andrea Kliewer[1], Viviane Stammer[1], Stefan Schulz[1,2,*]

[1]Institut für Pharmakologie und Toxikologie, Universitätsklinikum Jena, Friedrich-Schiller-

Universität Jena, Drackendorfer Straße 1, D-07747 Jena, Germany

[2]7TM Antibodies GmbH, Hans-Knöll-Straße 6, D-07745 Jena, Germany

[*]**Corresponding author**

Emails: stefan.schulz@med.uni-jena.de

Abstract

G protein-coupled receptors (GPCRs) are important signal transducers that are

phosphorylated upon activation at intracellular serine and threonine residues. Although

antibodies that specifically recognize the phosphorylation state of GPCRs have been available

for many years, efficient immunolocalization of phosphorylated receptors in their tissues of

origin has not been possible. Here, we show that phosphorylation of receptors is highly

unstable during routine immunohistochemical procedures, requiring the use of appropriate

phosphatase inhibitors particular during tissue perfusion, post-fixation, and cryoprotection but

not during immunostaining of tissue sections. We provide proof of concept using

phosphorylation state-specific μ-opioid receptor (MOP) and cannabinoid receptor 1 (CB1)

antibodies. Indeed, three of four well-characterized phosphosite-specific MOP antibodies,

including pS375-MOP, pT376-MOP, and pT379-MOP, showed robust neuronal

immunostaining in brain and spinal cord sections of opioid-treated mice only after inclusion

of phosphatase inhibitors. We then extended this approach to the CB1 receptor and

demonstrated that one of three newly-generated phosphosite-specific CB1 antibodies, namely

pS425-CB1, showed striking staining of fibers and varicosities in brain slices from

cannabinoid-treated mice. Although subsequent experiments showed that phospho-CB1

immunostaining was less sensitive to phosphatases, we conclude that the use of phosphatase

inhibitors should always be considered in the development of immunohistochemical

procedures for new phosphosite-specific GPCR antibodies. In summary, we anticipate that

this improved protocol will facilitate the widespread use of phosphorylation state-specific

antibodies to monitor the activation of endogenous GPCRs under physiological and

pharmacological conditions. Our approach may also prove useful to confirm target

engagement of GPCR drug candidates in native tissues.

Introduction

Agonist-driven serine/threonine phosphorylation is a biologically and pharmacologically important process that primarily initiates desensitization and internalization of G protein-coupled receptors (GPCRs) [1-5]. Phosphorylation also increases the interaction of GPCRs with intracellular adapter proteins such as β-arrestins, which can trigger a second wave of signaling [1-5]. Thus, analysis of agonist-driven phosphorylation can provide valuable insights into the receptor activation state and ligand pharmacology. A widely used approach to analyze GPCR phosphorylation is the use of phosphosite-specific antibodies [6-9]. When available, such antibodies are valuable tools to elucidate the temporal dynamics of receptor phosphorylation, identify relevant kinases and phosphatases, and detect receptor activation using immunoblotting techniques [7, 10-15]. However, previous attempts using routine immunohistochemical approaches to unequivocally reveal agonist-induced phosphorylation of endogenous GPCRs in native tissues largely failed.

In intact cells, GPCR dephosphorylation is regulated in time and space, beginning immediately after receptor activation at the plasma membrane [16]. Indeed, phosphosite-specific antibodies in combination with siRNAs have led to the identification of distinct protein phosphatase 1 (PP1) and PP2 catalytic subunits as bona fide GPCR phosphatases [17-22]. For many receptors, dephosphorylation is complete within 10 to 30 minutes after agonist washout [17, 18, 21]. A notable exception is the agonist-induced phosphorylation of S341/S343 at the SST2 somatostatin receptor, which persists for considerably longer periods [21]. Consequently, activated SST2 receptors could be successfully localized using pS341/pS343-SST2 antibodies under routine immunohistochemical conditions [23]. However, this approach could not easily be reproduced for other SST2 sites or other GPCRs.

The ability to visualize activated and phosphorylated GPCRs in their tissues of origin

would provide important clues to the physiological and pharmacological regulation of

receptor activation. In particular, it would allow for distinction between currently-activated

and resting GPCR populations in the context of particular physiological or behavioral

conditions. We were therefore motivated to develop and validate immunohistochemical

fixation and staining procedures that can be universally applied to prototypical GPCRs. Our

improved protocol will facilitate the widespread application of phosphosite-specific

antibodies as GPCR activation sensors in academic and pharmaceutical research.

Results

Dephosphorylation of GPCRs occurs rapidly in intact cells and is mediated by the protein

phosphatases PP1 and PP2 [16]. Serine/threonine phosphatases are known to be activated during

cell lysis and tissue fixation procedures. We therefore tested the inclusion of appropriate

protein phosphatase inhibitors (PPIs) during tissue fixation and immunohistochemical

staining. For this purpose, mice were treated with either methadone or saline, perfusion fixed,

and slices of brain and spinal cord were stained with phosphosite-specific μ-opioid receptor

(MOP) or phosphorylation-independent np-MOP antibodies. As expected, staining using the

np-MOP antibody was present in the superficial layers of the spinal cord, regardless of drug

treatment or the presence of phosphatase inhibitors (**Fig. 1**). In contrast, immunostaining for

pT379-MOP was only detected in methadone-treated animals when a cocktail of appropriate

phosphatase inhibitors was present during both fixation and staining procedures (**Fig. 1**). Four

unique serine and threonine residues are phosphorylated in MOP in response to full agonists [8,

15]. Phosphosite-specific antibodies have been generated for all of these sites, which work

equally well in immunoblot applications [7, 24, 25]. We found that three out of four, namely

pS375-MOP, pT376-MOP and pT379-MOP, but not pT370-MOP, are also well suited for

immunohistochemical staining of phosphorylated MOP under these conditions (**Fig. 2 upper**

panel). Neither p-MOP nor np-MOP staining was apparent in MOP knockout mice (**Fig. 2**

lower panel). To further substantiate the specific binding of phospho-MOP antibodies, we

performed peptide neutralization controls. When pS375-MOP or pT379-MOP antibodies were

incubated with an excess of their respective immunizing peptides containing S375 or T379 in

phosphorylated form, immunostaining was completely absent in the superficial layers of the

spinal cord (**Fig. 3**). In contrast, immunostaining was virtually unaffected by the addition of

the corresponding non-phosphorylated peptides, indicating unequivocal detection of agonist-

dependent S375 and T379 phosphorylation of MOP in tissues (**Fig. 3**).

The antagonist naloxone drives MOP into an inactive state, preventing agonist-

induced phosphorylation. When mice were treated with the selective MOP agonist fentanyl,

we found the typical staining pattern for pS375-MOP, pT376-MOP and pT379-MOP as

evidence of MOP activation. In contrast, this staining was not detected in animals injected

with naloxone immediately prior to fentanyl challenge, indicating that MOP activation was

blocked by the antagonist (**Fig. 4**). Collectively, our results support the notion that agonist-

induced MOP phosphorylation is specifically detected in brain sections under these

conditions. Notably, agonist-driven changes in phospho-immunofluorescence staining of

MOP were robust enough to allow for quantification even in relatively small groups of

animals (Supplemental Figure S1). Given the considerable costs involved in using PPIs, we

then tested whether their presence is necessary during both fixation and staining procedures.

The results depicted in **Figure 5**, clearly revealed that inclusion of PPIs during tissues fixation

is an absolute requirement. They also show that after fixation, it is not obligatory to add PPIs

to each solution for successful immunostaining (**Fig. 5**).

In an effort to perform dual immunofluorescence labeling, we combined the rabbit

pT376-MOP antibody (depicted in green, **Fig. 6**) with guinea pig np-MOP antibody (depicted

in magenta, **Fig. 6**). This procedure allows to discern populations of non-phosphorylated

MOP from activated and phosphorylated MOP, which dramatically increases in the

superficial layer of the spinal cord after administration of fentanyl or methadone (depicted in

white in overlay images, **Fig. 6**). Next, we compared the immunostaining patterns obtained

for phospho-MOP with total MOP. While there was good overlap in some brain regions,

including habenula, fasciculus retroflex (**Fig. 7**), unexpected low or even absent pS375-MOP,

pT376-MOP, nor pT379-MOP immunostaining was observed in the nucleus accumbens and

caudate putamen, despite high levels of MOP expression (**Fig. 7**). These results suggest that

region-specific MOP activation and phosphorylation patterns exist, which can now be

visualized for the first time.

The final set of experiments was designed to extend this new approach to another

prototypical GPCR namely the CB1 cannabinoid receptor. To this end, three newly-generated

phosphosite-specific CB1 antibodies including pS425-CB1, pS429-CB1 and pS441-CB1 were

tested under the new assay conditions. In fact, the pS425-CB1 antibody produced staining in

hippocampus and cortex closely resembling that of the np-CB1 antibody only when mice

were treated with CP-55940 [26] (**Fig. 8**). Notably, pS425-CB1 immunostaining was much less

prominent in untreated animals but not absent (**Fig. 8**). Staining in both untreated and

cannabinoid-treated animals was completely abolished when the p425-CB1 antibody was

neutralized with its immunizing peptide (**Fig. 8**). Prior administration of the antagonist

AM251 reduced CP-55940-induced phospho-CB1 immunostaining to that seen in untreated

mice (**Fig. 8**). To corroborate these findings using an independent biochemical approach,

groups of three animals each were treated with either vehicle or CP55940. CB1 receptors

were then immunoprecipitated using a goat np-CB1 antibody and CB1 phosphorylation was

determined using a rabbit pS425-CB1 antibody (**Fig. 9**). To confirm equal loading, the blot

was stripped and total CB1 was detected using a rabbit np-CB1 antibody (**Fig. 9**). The results

depicted in Figure 9, reveal the existence of basal pS425 phosphorylation of CB1 that is

strongly increased after administration of the high-efficacy agonist CP55940 (**Fig. 9**). When

CB1 immunoprecipitates were subjected to λ-phosphatase digestion, both basal and agonist-

driven CB1 phosphorylation were completely diminished, whereas total CB1 was still

detectable. Collectively, these findings support the conclusion that CB1 phosphorylation can

be specifically detected. We then tested whether the presence of PPIs during tissues fixation is

an absolute requirement for successful pS425-CB1 immunostaining. Surprisingly, we found

that CB1 phosphorylation appears to be much less sensitive to in situ phosphatase digestion

than MOP phosphorylation (Supplemental Fig. S2). Finally, we used a combination of goat

np-CB1 and rabbit pS425-CB1 antibodies to perform dual immunofluorescence labeling. At

higher magnification it became apparent that the p425-CB1 antibody exclusively labelled

fibers and terminals similar to that seen with the np-CB1 antibody (**Fig. 10**). Interestingly,

overlay images revealed that in CP-55940-treated mice activated and phosphorylated CB1

receptors were densely concentrated in varicosities (depicted in white, **Fig. 10**), whereas CB1

was detected in fibers appears to reside in the non-phosphorylated state (depicted in magenta,

Fig. 10).

Discussion

GPCRs are privileged targets for small molecule drugs in almost all therapeutic areas [27].

However, to date, the visualization of therapeutic drug effects on endogenously expressed

GPCRs in native tissues has remained a major challenge. Activation of GPCRs by their

endogenous ligands or exogenous agonists results in conformational changes that are

recognized by a family of kinases termed G protein-coupled receptor kinases (GRKs) [1-5]. The

unique ability of GRKs to recognize activated receptors results in agonist-dependent

phosphorylation at intracellular serine and threonine residues [1-5]. Thus, analysis of agonist-

driven phosphorylation of GPCRs can provide valuable insights into receptor activation.

Antibodies that specifically recognize the GPCR phosphorylation state have been available in

some cases over the last 20 years. Indeed, phosphosite-specific GPCR antibodies have proven

useful to detect receptor activation and to profile the pharmacological properties of new

ligands [12-15]. To date, however, their utility has been largely limited to immunoblotting

approaches using heterologous receptor expression systems, while detection of

phosphorylated GPCRs from tissue lysates has been mostly unsuccessful, due to their low

abundance *in vivo*.

Here, we have systematically developed an improved immunohistochemical staining

procedure that allows for visualization of opioid drug effects in the mouse brain *in vivo*. We

also show that this approach can be easily extended to other prototypical GPCRs. Key to this

new protocol was the observation that receptor phosphorylation is highly unstable during

routine immunohistochemistry and that for many GPCRs inclusion of protein phosphatase

inhibitors may be an absolute requirement during perfusion, post-fixation and cryoprotection

of tissues. Several lines of evidence suggest that our approach facilitates the localization of

bona fide phosphorylated GPCRs. First, phospho-GPCR immunostaining strongly increased

when animals had been treated with agonist. Conversely, immunostaining was not detected

when agonist action was blocked by antagonist. Second, robust phospho-GPCR

immunostaining was only apparent when appropriate protein phosphatase inhibitors were

included in all steps of fixation and staining procedures. Third, phospho-GPCR

immunostaining was completely neutralized by preincubation with cognate phosphorylated

peptides but not with the corresponding non-phosphorylated peptides.

In the present study, we had access to four different phospho-MOP and three different

phospho-CB1 antibodies which had previously been validated by immunoblotting techniques.

We observed that three out of four antibodies for MOP and one out of three antibodies for

CB1 produced specific phospho-immunostaining in tissues suggesting that each phosphosite-

specific GPCR antibody has to be validated individually for immunohistochemical

applications. Moreover, we observed that phospho-MOP staining was critically dependent on

the presence of PPIs, whereas CB1 phosphorylation appeared to be more stable in tissues.

Nevertheless, we conclude that use of PPIs should always be considered in the development

of immunohistochemical procedures for new phosphosite-specific GPCR antibodies.

It is well known that endogenous GPCRs are notoriously difficult to detect in tissues

by immunohistochemical staining. Here, we not only establish a protocol for in situ

visualization of activated and phosphorylated GPCRs. We also demonstrate that phospho-

immunostaining can disclose previously unappreciated functional information not seen with

phosphorylation-independent GPCR antibodies. One example is the striking mismatch

between total and phospho-immunostaining for MOP in caudate putamen. This mismatch

cannot be explained be low detection level of phospho-MOP staining as many other brain

regions with comparable total MOP revealed prominent phospho-MOP. The most likely

explanation for the relative lack of phospho-MOP staining in caudate putamen is the low or

absent expression of relevant GRK family members [32]. Notably, lack of MOP

phosphorylation in some brain regions relative to others could have profound consequences

on MOP desensitization and may explain well-known differences in tolerance development to

different opioid effects. Another example is the different subcellular localization of

phosphorylated and non-phosphorylated CB1 receptors. While total CB1 appears to be evenly

distributed along fibers and varicosities, phosphorylated CB1 is particular enriched in

terminals indicating compartmentalization of GRK-mediated phosphorylation.

216   In summary, we provide a proof of concept for using phosphorylation state-specific

217 antibodies as sensors to elucidate GPCR activation in native tissues. Our approach not only

218 facilitates the visualization of the effects of agonists and antagonists in preclinical animal

219 studies. It also allows for elucidation of different cellular and subcellular activation patterns in

220 response to external stimuli or drug administration. Therefore, phosphorylation state-specific

221 antibodies are a new class of markers for GPCR activation that are likely to find widespread

222 application in academic and pharmaceutical research.

 Figure legends

 Fig. 1. Comparison of phospho-MOP immunohistochemistry in the presence or absence of

 protein phosphatase inhibitors. Animals were either treated with methadone or saline for 30

 min, transcardially perfused, fixed and stained in the presence (+) or absence (-) of protein

 phosphatase inhibitors (PPI). Shown are confocal images of coronal sections of the spinal

 cord stained with pT379-MOP or np-MOP antibody. Note that PPIs need to be present during

 both fixation and staining procedures to obtain agonist-induced phospho-MOP

 immunostaining. Scale bar = 250 µm.

 Fig. 2. Immunohistochemical staining of agonist-induced MOP phosphorylation in the mouse

 spinal cord. Animals were treated with methadone or saline for 30 min, transcardially

 perfused, fixed and stained in the presence of PPIs. Shown are confocal images of coronal

 sections of the spinal cord stained with phosphosite-specific antibodies pT370-MOP, pS375-

 MOP, pT376-MOP and pT379-MOP, or phosphorylation independent np-MOP antibody.

 Note that pS375-MOP, pT376-MOP and pT379-MOP, but not pT370-MOP, revealed an

 agonist-dependent MOP phosphorylation immunostaining in a pattern closely resembling that

 of np-MOP immunostaining. Scale bar = 250 µm.

 Fig. 3. Peptide neutralization of agonist-induced MOP phosphorylation staining in the mouse

 spinal cord. Animals were treated with methadone for 30 min, transcardially perfused, fixed

 and stained in the presence of PPIs. Shown are confocal images of coronal sections of the

 spinal cord stained with phosphosite-specific antibodies pS375-MOP or pT379-MOP in the

 presence or absence of their immunizing peptides containing S375 or T379 in phosphorylated

 form (p-Peptide) or the corresponding non-phosphorylated peptide (np-Peptide). Note that

phospho-MOP immunostaining was completely neutralized by excess of phosphorylated

peptide but not of non-phosphorylated peptide. Scale bar = 250 µm.

Fig. 4. Antagonist block of phospho-MOP immunostaining. Animals were either treated with

saline or fentanyl for 15 min. Where indicated, animals were pretreated with naloxone for 10

min followed by 15-min fentanyl treatment. Animals were then transcardially perfused, fixed

and stained in the presence of PPIs. Shown are confocal images of coronal sections of the

spinal cord stained with phosphosite-specific antibodies pS375-MOP, pT376-MOP or pT379-

MOP or phosphorylation independent np-MOP antibody. Note that agonist-induced phospho-

MOP immunostaining is diminished by antagonist treatment. Scale bar = 250 µm.

Fig. 5. Comparison of phospho-MOP detection after staining in the presence or absence

of PPIs. Animals were treated with methadone for 30 min, transcardially perfused and fixed

in the presence (+) or absence (-) of PPIs and also stained in the presence (+) or absence (-) of

PPIs. Shown are confocal images of coronal sections of the spinal cord stained with pS375-

MOP, pT379-MOP or np-MOP antibody. Note that PPIs need to be present during the whole

fixation procedures but not during the staining protocol to obtain agonist-induced phospho-

MOP immunostaining. Scale bar = 250 µm.

Fig. 6. Double staining of agonist-induced p-MOP and np-MOP in mouse brain. Animals

were either treated with NaCl (30 min), Fentanyl (15 min) or Methadone (30 min). Animals

were then transcardially perfused, fixed and stained. Shown are confocal images of coronal

sections of the spinal cord stained with phosphosite-specific antibody pT376-MOP (green)

and phosphorylation independent np-MOP antibody (magenta). Note that there is a strong

overlap of np-MOP and pT376-MOP in the merged panels (white) in agonist treated animals

compared to NaCl treated mice. Scale bar = 250 μm.

Fig. 7. Agonist-induced phospho-MOP immunostaining in mouse brain. Animals were treated

with methadone for 30 min, transcardially perfused, fixed and stained in the presence of PPIs.

Shown are confocal images of coronal brain sections stained with phosphosite-specific

antibodies pS375-MOP, pT376-MOP or pT379-MOP or phosphorylation independent np-

MOP antibody. Note that all three phosphosite-specific antibodies detected agonist-induced

MOP phosphorylation in the medial habenula and fasciculus retroflexus but not in nucleus

accumbens or caudate putamen, brain regions known to be rich in MOP receptors as shown

by np-MOP antibody staining. Scale bar = 250 μm.

Fig. 8. Agonist-induced phospho-CB1 immunostaining in mouse brain. Animals were

either treated with Vehicle or the CB1-agonist CP-55940 for 30 min. Where indicated,

animals were pretreated with the CB1-antagonistAM251 for 45 min followed by 30 min CP-

55940 treatment. Animals were then transcardially perfused, fixed and stained in the presence

of PPIs. Shown are confocal images of coronal brain sections stained with phosphosite-

specific antibody pS425-CB1, in the presence or absence of the p-Peptide containing S425 in

phosphorylated form, or phosphorylation independent np-CB1 antibody. Note that agonist-

induced phopho-CB1 immunostaining is diminished by antagonist treatment. Note that

phospho-CB1 immunostaining was completely neutralized by excess of phosphorylated

peptide. Scale bar = 1000 μm and 250 μm.

Fig. 9. Immunoblotting of phosphorylated CB1 in agonist treated mice. Six different

wildtype animals were used. Number 1 to 3 were treated with vehicle and Number 4 to 6

with CP-55940 for 30 min. Organs were removed rapidly and frozen in liquid nitrogen After

tissue preparations, CB1 receptors were extracted using goat np-CB1 bound to protein A

beads. After blotting, the membranes were treated with the phosphosite-specific rabbit

antibody pS425-CB1, stripped (dashed line) and reprobed with rabbit phosphorylation-

independent anti-CB1. Note that there is some CB1-phosphorylation constitutive signal in

vehicle-treated mice. Molecular mass on the left in kDa.

Fig. 10. Double staining of agonist-induced p-CB1 and np-CB1 in mouse brain. Animals

were either treated with Vehicle (A) or CP-55940 (B, C) for 30 min. Animals were then

transcardially perfused, fixed and stained. Shown are confocal images of coronal brain

sections of cortex (CTX) and hippocampal formation (HPF) stained with phosphosite-specific

antibody pS425-CB1 (green) and phosphorylation independent np-CB1 antibody (magenta).

Note that there is a strong overlap of np-CB1 and pS425-CB1 in the merged panels (white) in

CP-55940 treated animals (B) compared to vehicle treated mice. Note that in cortical neurons,

only CB1 located in varicosities undergoes agonist-induced phosphorylation. Scale bar =

1000 µm, 100 µm and 2 µm.

Methods

Antibodies

Primary Antibodies:

The phosphorylation state-specific rabbit MOP antibodies pT370-MOP (7TM0319B), pS375-MOP (7TM0319C), pT376-MOP (7TM0319D), pT379-MOP (7TM0319E), the phosphorylation state-specific rabbit CB1 antibody pS425-CB1 (7TM0056A) as well as the phosphorylation-independent rabbit antibody np-MOP (7TM0319N) and phosphorylation-independent guinea pig antibody np-MOP (7TM0319N-GP) were provided by 7TM Antibodies (www.7tmantibodies.com). In addition, we used rabbit np-MOP from abcam (ab134054), rabbit np-CB1 from Cayman (1000659) and goat np-CB1 from Frontier Institute (MFSR100610).

Secondary Antibodies:

We used a biotinylated AB from Jackson Immuno Research (711-065-152, anti-rabbit) and the respective streptavidine conjugates from ThermoFisher (S21381, Alexa Fluor™ 555 conjugate) and (S11223, Alexa Fluor™ 488 conjugate) for the amplified staining procedure. For the direct stainings we used Cy™3 (706-165-148, anti-guinea pig) and Cy™3 (705-165-147, anti-goat), both obtained from Jackson ImmunoResearch.

Animals

Mice (C57BL/6J obtained from JAX™) were housed 2–5 per cage under a 12-hr light-dark cycle with ad libitum access to food and water. All animal experiments were approved by Thuringian state authorities and performed in accordance with European Commission regulations for the care and use of laboratory animals. Our study is reported in accordance with

ARRIVE guidelines [28]. In all experiments, male and female mice aged 8–30 weeks between 25

and 35 g body weight were used.

Drugs and routes of administration

All drugs were freshly prepared prior to use and were injected subcutaneously (s.c.) or

intraperitoneally (i.p.) in unanaesthetized mice at a volume of 10 µl/g bodyweight. Opioid drugs

were diluted in 0.9% (w/v) saline and cannabinoid drugs in vehicle solution, containing 13.2%

ethanol and 0.25% Tween80 in 0.9% saline, for injections. Drugs were obtained and used as

follows: fentanyl citrate (0.3 mg/kg for 15 min; s.c.) (B. Braun 06900650), levomethadone

hydrochloride (15 mg/kg for 30 min; s.c.) (Sanofi-Aventis 07480196), naloxone (2 mg/kg for

25 min; s.c.) (Ratiopharm 04788930), CP-55940 (0.75 mg/kg for 30 min; i.p.) (Sigma-Aldrich

C1112) and AM251 (3 mg/kg for 45 min; i.p.) (MedChemExpress 183232-66-8). A total of 35

mice (n=4-6 per treatment condition) were used.

Immunohistochemistry

Mice were deeply anaesthetized with isoflurane (CP-Pharma 4001404) and then subjected to a

transcardial perfusion with calcium-free Tyrode's solution containing protein phosphatase-

inhibitors (+PPIs) (1 tablet PhosSTOP per 10 ml) (Roche 04906845001) followed by

Zamboni's fixative containing 4% paraformaldehyde and 0.2% picric acid in 0.1 M phosphate

buffer pH 7.4 +PPIs. Brains and spinal cords were rapidly dissected and postfixed in the same

fixative for 4 h at room temperature. The tissue was cryoprotected by immersion in 10% sucrose

+PPIs followed by 30% sucrose +PPIs for 48 h at 4 °C before sectioning using a freezing

microtome. Tissue was cut into 40 µm sections. Free-floating sections were washed in PBS

with or without (+/-) PPIs (depending on the experiment and investigated GPCR) and incubated

in methanol containing 0.3% H_2O_2 for 30 min. After washing in PBS-T (PBS + 0,3%

Tween®20), the sections were blocked in PBS containing 0.3% Triton X-100, +/- PPIs and

10% NGS for 2 h. Subsequently, the sections were incubated with primary antibody in PBS

containing 0.3% Triton X-100, 2% NGS +/- PPI overnight at 4°C. Where indicated, primary

antibodies were preincubated with 1 µg/ml of their cognate phospho-peptide or the

corresponding non-phospho-peptide for 1 h at room temperature. When necessary, staining of

primary antibody was detected using the biotin amplification procedure as described [29, 30].

Briefly, tissue sections were transferred to biotinylated donkey anti-rabbit IgG (1:300 in PBS

containing 0.3% Triton X-100, +/-PPIs and 10% NGS) for 2 h, washed in PBS-T +/-s PPIs and

then incubated in AB solution (reagents from the Vector ABC kit; 25 µl A and 25 µl B in 10

ml PBS+0.3% Triton X-100 +/- PPIs) for 60 min, washed again in PBS-T +/- PPIs and

transferred to biotinylated tyramine (BT) solution (BT was prepared as described by Adams [31],

5 µl BT+0.01% H_2O_2 in 1 ml PBS+0.3% Triton X-100 and +/- PPIs) for 20 min, followed by a

final incubation step in streptavidin-AlexaFluor555 conjugate 1:400 in PBS+0.3% Triton X-

100 containing 10% NDS +/- PPI) overnight at 4°C. For double-staining procedures, the

primary np-AB was incubated overnight as indicated and after several washing steps on the

following day conjugated to the secondary AB, again overnight. Sections were then mounted

onto SuperFrost Plus glass slides (ThermoFisher 15438060) and cover slipped with Eukitt

(ORSAtec). Specimens were examined using a Zeiss LSM 900 laser scanning confocal

microscope equipped with ZEN software for image analysis.

Western Blot and immunoprecipitation

Wildtyp mice were treated with agonist or vehicle. Mice were anesthetized with isoflurane,

killed by cervical dislocation, and brains were quickly dissected, excluding the cerebellum.

Brain samples were immediately frozen in liquid nitrogen. Brains were transferred to ice-cold

detergent buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 5 mM EDTA, 1% Nonidet P-40,

0.5% sodium deoxycholate, 0.1% sodium dodecyl sulfate (SDS), containing protease and

phosphatase inhibitors), homogenized, and centrifuged at 14,000 × g for 30 min at 4°C. The

supernatant was then immunoprecipitated with the polyclonal goat np-CB1 antibody bound to

protein A-agarose beads (20333, ThermoScientific, Germany) for 90 min at 4°C. Proteins were

eluted from the beads with SDS sample buffer for 25 min at 43°C and then resolved on 8%

SDS-polyacrylamide gels. After electroblotting, membranes were incubated with pS425-MOP

antibody, followed by detection using a chemiluminescence detection system. Blots were

subsequently stripped and incubated again with the rabbit np-CB1 antibody to confirm equal

loading of the gels.

Data availability

All data supporting the findings of this study are available within the article and its

supplementary information files. Additional information, relevant data and unique biological

materials will be available from the corresponding author upon reasonable request. Source data

are provided with this paper.

References

1. Drube, J. et al. GPCR kinase knockout cells reveal the impact of individual GRKs on arrestin binding and GPCR regulation. *Nat Commun* **13**, 540 (2022).

2. Gurevich, E.V., Tesmer, J.J., Mushegian, A. & Gurevich, V.V. G protein-coupled receptor kinases: more than just kinases and not only for GPCRs. *Pharmacol Ther* **133**, 40-69 (2012).

3. Liggett, S.B. Phosphorylation barcoding as a mechanism of directing GPCR signaling. *Sci Signal* **4**, pe36 (2011).

4. Premont, R.T. & Gainetdinov, R.R. Physiological roles of G protein-coupled receptor kinases and arrestins. *Annu Rev Physiol* **69**, 511-534 (2007).

5. Tobin, A.B., Butcher, A.J. & Kong, K.C. Location, location, location...site-specific GPCR phosphorylation offers a mechanism for cell-type-specific signalling. *Trends Pharmacol Sci* **29**, 413-420 (2008).

6. Doll, C. et al. Agonist-selective patterns of micro-opioid receptor phosphorylation revealed by phosphosite-specific antibodies. *Br J Pharmacol* **164**, 298-307 (2011).

7. Grecksch, G. et al. Analgesic tolerance to high-efficacy agonists but not to morphine is diminished in phosphorylation-deficient S375A mu-opioid receptor knock-in mice. *J Neurosci* **31**, 13890-13896 (2011).

8. Just, S. et al. Differentiation of opioid drug effects by hierarchical multi-site phosphorylation. *Mol Pharmacol* **83**, 633-639 (2013).

9. Prihandoko, R., Bradley, S.J., Tobin, A.B. & Butcher, A.J. Determination of GPCR Phosphorylation Status: Establishing a Phosphorylation Barcode. *Curr Protoc Pharmacol* **69**, 2 13 11-12 13 26 (2015).

10.   Divorty, N. et al. Agonist-induced phosphorylation of orthologues of the orphan

receptor GPR35 functions as an activation sensor. *J Biol Chem*, 101655 (2022).

11.   Illing, S., Mann, A. & Schulz, S. Heterologous regulation of agonist-independent mu-

opioid receptor phosphorylation by protein kinase C. *Br J Pharmacol* **171**, 1330-1340

(2014).

12.   Mann, A. et al. New phosphosite-specific antibodies to unravel the role of GRK

phosphorylation in dopamine D2 receptor regulation and signaling. *Sci Rep* **11**, 8288

(2021).

13.   Mann, A. et al. Agonist-induced phosphorylation bar code and differential post-

activation signaling of the delta opioid receptor revealed by phosphosite-specific

antibodies. *Sci Rep* **10**, 8585 (2020).

14.   Mann, A. et al. Agonist-selective NOP receptor phosphorylation correlates in vitro and

in vivo and reveals differential post-activation signaling by chemically diverse

agonists. *Sci Signal* **12** (2019).

15.   Miess, E. et al. Multisite phosphorylation is required for sustained interaction with

GRKs and arrestins during rapid mu-opioid receptor desensitization. *Sci Signal* **11**

(2018).

16.   Kliewer, A., Reinscheid, R.K. & Schulz, S. Emerging Paradigms of G Protein-

Coupled Receptor Dephosphorylation. *Trends Pharmacol Sci* **38**, 621-636 (2017).

17.   Doll, C. et al. Deciphering micro-opioid receptor phosphorylation and

dephosphorylation in HEK293 cells. *Br J Pharmacol* **167**, 1259-1270 (2012).

18.   Lehmann, A., Kliewer, A., Gunther, T., Nagel, F. & Schulz, S. Identification of

Phosphorylation Sites Regulating sst3 Somatostatin Receptor Trafficking. *Mol*

Endocrinol **30**, 645-659 (2016).

19.   Lehmann, A., Kliewer, A., Martens, J.C., Nagel, F. & Schulz, S. Carboxyl-terminal

455   receptor domains control the differential dephosphorylation of somatostatin receptors

456   by protein phosphatase 1 isoforms. *PLoS One* **9**, e91526 (2014).

457   20.   Petrich, A. et al. Phosphorylation of threonine 333 regulates trafficking of the human

458   sst5 somatostatin receptor. *Mol Endocrinol* **27**, 671-682 (2013).

459   21.   Poll, F., Doll, C. & Schulz, S. Rapid dephosphorylation of G protein-coupled

460   receptors by protein phosphatase 1beta is required for termination of beta-arrestin-

461   dependent signaling. *J Biol Chem* **286**, 32931-32936 (2011).

462   22.   Tran, T.M. et al. Characterization of beta2-adrenergic receptor dephosphorylation:

463   Comparison with the rate of resensitization. *Mol Pharmacol* **71**, 47-60 (2007).

464   23.   Waser, B. et al. Phosphorylation of sst2 receptors in neuroendocrine tumors after

465   octreotide treatment of patients. *Am J Pathol* **180**, 1942-1949 (2012).

466   24.   Fritzwanker, S. et al. HA-MOP knockin mice express the canonical micro-opioid

467   receptor but lack detectable splice variants. *Commun Biol* **4**, 1070 (2021).

468   25.   Kliewer, A. et al. Phosphorylation-deficient G-protein-biased mu-opioid receptors

469   improve analgesia and diminish tolerance but worsen opioid side effects. *Nat Commun*

470   **10**, 367 (2019).

471   26.   Davis, M.I. et al. The cannabinoid-1 receptor is abundantly expressed in striatal

472   striosomes and striosome-dendron bouquets of the substantia nigra. *PLoS One* **13**,

473   e0191436 (2018).

474   27.   Santos, R. et al. A comprehensive map of molecular drug targets. *Nat Rev Drug*

475   *Discov* **16**, 19-34 (2017).

476   28.   Percie du Sert, N. et al. The ARRIVE guidelines 2.0: Updated guidelines for reporting

477   animal research. *PLoS Biol* **18**, e3000410 (2020).

478   29.   Schreff, M. et al. Distribution, targeting, and internalization of the sst4 somatostatin

479   receptor in rat brain. *J Neurosci* **20**, 3785-3797 (2000).

480   30.   Schulz, S. et al. Immunolocalization of two mu-opioid receptor isoforms (MOR1 and

481   MOR1B) in the rat central nervous system. *Neuroscience* **82**, 613-622 (1998).

482   31.   Adams, J.C. Biotin amplification of biotin and horseradish peroxidase signals in

483   histochemical stains. *J Histochem Cytochem* **40**, 1457-1463 (1992).

32.   Allen Brain Atlas, https://atlas.brain-map.org

Acknowledgements: We want to thank Svetlana Würl and Monique Brendel for expert

technical assistance. This work was supported by the European Regional Development Fund

(EFRE) and the Free State of Thuringia (grant: 2020 FE 0146) to 7TM Antibodies GmbH.

Author contributions: S.S. conceived and initiated the project and designed all experiments

with S.F. . S.F. and A.K. performed and analyzed immunohistochemistry. F.N. affinity-

purified all phosphosite-specific antibodies used in this study. The manuscript was written

and revised by S.S. and S.F. with input from other authors.

Competing interests: S.S. is the founder and scientific advisor of 7TM Antibodies GmbH,

Jena, Germany. F.N. is an employee of 7TM Antibodies. All other authors declare no

competing interests.

	Medial habenula		fasciculus retroflexus		Nucleus accumbens		Caudoputamen	
	NaCl	Methadone	NaCl	Methadone	NaCl	Methadone	NaCl	Methadone
pS375-MOP								
pT376-MOP								
pT379-MOP								
np-MOP								