Nutrient limitation mimics artemisinin tolerance in malaria

Audrey C. Brown, Michelle D. Warthan, Anush Aryal, Shiwei Liu, Jennifer L. Guler*
Department of Biology, University of Virginia, Charlottesville, VA.

Abstract

Mounting evidence demonstrates that nutritional environment can alter pathogen drug sensitivity. While the nutrient-rich media used for standard *in vitro* culture contains supra-physiological nutrient concentrations, pathogens encounter a relatively restrictive environment *in vivo*. We assessed the effect of nutrient limitation on the protozoan parasite that causes malaria and demonstrated that short-term growth under physiologically-relevant mild nutrient stress (or "metabolic priming") triggers increased tolerance of the potent antimalarial drug dihydroartemisinin (DHA). We observed beneficial effects using both short-term survival assays and longer-term proliferation studies, where metabolic priming increases parasite survival to a level previously defined as DHA resistant (>1% survival). We performed these assessments by either decreasing single nutrients that have distinct roles in parasite metabolism or using a media formulation with reductions in many nutrients that simulates the human plasma environment. We determined that priming-induced DHA tolerance was restricted to parasites that had newly invaded the host red blood cell but the effect was not dependent on genetic background. The molecular mechanisms of this intrinsic effect mimic aspects of genetic artemisinin tolerance, including translational repression, autophagy, and protein export. This finding suggests regardless of the impact on survival rates, environmental stress could stimulate changes that ultimately directly contribute to drug tolerance. Because metabolic stress is likely to occur more frequently *in vivo* compared to the stable *in vitro* environment, priming-induced drug tolerance has ramifications for how *in vitro* results translate to *in vivo* studies. Improving our understanding of how pathogens adjust their metabolism to impact survival of current and future drugs is an important avenue of research to prevent and slow the spread of resistance.

Introduction

In vitro culture provides microbes with a stable environment for growth with relatively little challenge to organismal homeostasis. For example, the supra-physiological concentrations of nutrients *in vitro* largely protect cultured organisms from the stress of starvation conditions. Yet, nutrient limitation occurs frequently *in vivo*[1-4]. Therefore, the consequences of dynamic adaptation to these stressors, such as variations in nutrient levels, are largely missed by standard *in vitro* experimental designs.

Interestingly, data from across the tree of life has shown that the relationship between stress and negative fitness consequences is not necessarily linear. The adaptive benefits of low doses of mild environmental stress overwhelm the adverse effects from the stressor in a biphasic dose response phenomenon termed hormesis ([5,6]; **Fig. 1A**). Given this effect, adjustment of metabolism in response to homeostatic challenge from mild nutrient stress may increase pathogen fitness and survival of subsequent stressors, such as drug pressure.

Indeed, metabolic adjustability to varied nutritional environments has been shown to alter drug sensitivity across diverse pathogens[7,8]. For example, in *Trypanasoma cruzi* reduction of exogenous glutamine decreases flux through the sterol synthesis pathway rendering parasites sensitive to azole drugs[9]. Meylan, *et al*. found *Pseudomonas aeruginosa* supplementation with specific carbon sources increased susceptibility to tobramycin[10]. Increasing proline concentration confers halofuginone resistance in *P. falciparum*[11]. The nutritional environment *in vitro* is highly divergent from conditions pathogens are exposed to in the context of human infection, with nutrient concentrations frequently found at considerably higher levels in culture[12-14]. Discrepancies in metabolic activity between *in vitro* and *in vivo* environments brought about by non-physiological conditions can lead to reduced translatability of findings from culture-based experiments when metabolic flux through pathways is altered[15,16]. For example, glycerol metabolism sensitizes *Mycobacterium tuberculosis* to pyrimidine-imidazoles through accumulation of toxic metabolites[15]. However, these drugs are ineffective *in vivo* where changes in nutrient availability lead to differences in carbon metabolism.

These previously characterized changes in pathogen drug sensitivity as a consequence of varied nutritional environment typically depend on either 1) changes in flux through a specific metabolic pathway relevant to a given drug target or 2) a general slowing of growth leading to decreased uptake and activation of drug. Here, we present a case where nutrient stress induces a change in drug sensitivity of the human malaria parasite, *Plasmodium falciparum*. We demonstrate that short-term growth under mild, physiologically relevant nutrient limitation leads to superior recovery from drug treatment. We observe this effect under distinct conditions that impact the parasites differently; therefore, it is not dependent on a change in flux within a specific metabolic

pathway nor an overall decrease in growth rate (as in above examples). Importantly, the increased survival rates achieve a level that reaches the field's definition of "resistance," yet genetic mutations that confer resistance are not necessary. We determined that the induction of stress response pathways within the parasite occurs during nutrient limitation, which prime the parasite to better withstand drug treatment. This study emphasizes the link between nutrient limitation, stress response pathways, and survival of drug treatment and has important implications for the evolution of drug resistant pathogens.

Results

Metabolic priming increases parasite recovery from DHA

We sought to determine the effect of mild nutrient stress on the survival of *P. falciparum*. Asynchronous parasite cultures were subjected to short-term (36-72h) incubation in media restricted of either hypoxanthine or thiamine (**Table 1**, termed "metabolic priming"). *Plasmodium* lack the ability to *de novo* synthesize purines, instead requiring scavenging from the environment. In serum-free malaria cultures, the purine derivative, hypoxanthine, is a standard additive to fulfill this requirement. In contrast, *Plasmodium* can both *de novo* synthesize or scavenge thiamine[17]. In addition to different modes of acquisition for these nutrients, hypoxanthine and thiamine represent distinct metabolite classes: a nucleoside important for nucleic acid synthesis and a cofactor for apicoplast and mitochondrial metabolism, respectively. The duration of incubation and level of nutrient limitation for these metabolites were optimized to induce stress while keeping the majority of the parasite population viable (**Supplemental Fig. 1**).

We treated parasites grown in either metabolic priming conditions or standard media (formulations detailed in *Methods*) with a rapid-acting, highly potent antimalarial drug (200nM dihydroartemisinin (DHA) for 6h, **Fig. 1B**). DHA is an artemisinin derivative that causes widespread damage within parasites by promiscuously alkylating different classes of biomolecules, including heme and numerous proteins, thereby disrupting multiple biological pathways[18-20]. Following removal of drug and continued culture for 10 days in standard media, primed parasites showed a significant increase in cumulative proliferation compared to non-primed controls (hypoxanthine primed *N*= 5, *p* = 0.0126; thiamine primed *N=4*, *p* = 0.0074) (**Fig 1C-D**). We detected this effect in both priming conditions despite the restricted nutrients in each condition being of unrelated metabolite classes, irrespective of the parasite line (**Supplemental Table 1**, *Dd2* originally from Southeast Asia and *NF54* from Africa). The increased cumulative proliferation is driven by altered response to DHA treatment; we only observed this effect in primed parasites that were subjected to DHA treatment but not in those treated with vehicle alone (**Supplemental Fig. 2**).

During metabolic priming experiments, the level of stress applied to the parasites impacted whether we observed increased cumulative proliferation. For example, a >10% growth reduction over the course of low hypoxanthine metabolic priming was found to facilitate the phenotype (**Fig. 1C**), while failure to detect a growth phenotype (**Supplemental Fig. 3A**) or overly stressful conditions (**Supplemental Fig. 3B**) were detrimental to post-DHA cumulative proliferation. These data support a biphasic dose response to metabolic priming characteristic of hormesis. However, this pattern is dependent on the limiting nutrient; in other conditions growth during priming did not serve as a proxy for estimating dose response. For example, thiamine-free metabolic priming did not lead to growth reduction but still led to increased cumulative proliferation following DHA treatment (**Fig. 1D**). The differential effect of priming treatments on growth is likely due to parasite ability to scavenge or *de novo* synthesize thiamine, whereas *P. falciparum* are strictly purine auxotrophs.

Metabolic priming is stage-specific and facilitates increased DHA survival rates
Because our priming conditions alter the parasites' response to DHA (**Fig. 1C-D**), and DHA survival patterns are known to be dependent on the parasite life cycle stage (with highest survival at 0-3hr post-invasion[21], **Fig. 2A**), we evaluated 1) stage effects of metabolic priming and 2) stage-specificity of the proliferation effect.

First, we compared parasite stage distribution in primed and non-primed populations to assess if subtle changes in lifecycle stage occur during priming conditions (i.e. stage effects). We did not detect any significant differences in the percentage of ring-stage parasites in non-primed compared to primed samples (Non-primed: 61.6% ± 2.1%, Primed: 61.3% ± 2.3% (mean ± S.E.M); p=0.67), (**Supplemental Table 1**). Therefore, an alteration of cell cycle progression is not contributing to differences in proliferation following DHA treatment.

Second, we performed shorter term DHA survival assays on highly synchronous parasite populations following priming to determine if changes in parasite proliferation are stage-specific. To assess the behavior of early stage parasites (**Fig. 2A**), we performed "ring stage survival assays" (RSAs) on highly synchronous populations of 0-3h rings[21]. Metabolic priming prior to DHA treatment increased ring stage survival (**Fig. 2B**); we observed a mean 2.1-fold increase for both 200nM and 700nM DHA, which is the concentration used for our study (**Fig. 1C-D**) and the standard concentration for DHA RSAs, respectively. Notably, metabolic priming increased survival percentages to a level that is considered resistant (>1% for 700nM DHA for 6h[21]). Additionally, we tested later stage parasites (24-27h) in a modified RSA that assesses trophozoite survival (TSA[21], **Fig. 2C**). We detected no significant increase in DHA survival following metabolic priming, indicating the benefit of metabolic priming is ablated in trophozoites (**Fig. 2D**). In addition to showing stage-specificity of the effect, these experiments also showed that higher cumulative

proliferation in primed samples post-DHA is a result of an increased percentage of parasites surviving DHA treatment, as opposed to, increased proliferation per parasite.

Metabolic priming increases survival in parasites with genetic DHA resistance

We sought to assess if the survival benefits of metabolic priming also apply to DHA resistant parasites (RSA >1%[21]). Using our long term growth assay, we found that metabolic-priming does improve post-drug recovery of a moderately DHA resistant parasite strain that harbors a mutation in the *kelch13* gene (**Fig. 3A**; MRA-1238, RSA: 6.2%, *kelch13* I543T mutant[22]). We confirmed this effect using RSAs; metabolic priming increases survival of a high DHA concentration (700mM) by a mean of 2.1-fold (**Fig. 3B**). This parasite line was not tested at the lower 200nM concentration as for DHA sensitive lines due to their known resistance status; other DHA resistant lines (i.e. MRA 1240, RDA: 88.2%, *kelch 13* R539T) could not be assessed because of their high tolerance to 700nM DHA levels. These data in combination with data generated using DHA sensitive lines (**Figs. 1** and **2, Supplemental Table 1;** *Dd2* and *NF54*), indicate that increased DHA survival following metabolic priming is not dependent on genetic background.

Autophagy contributes to increased DHA survival

Phosphorylation of the translation initiation factor eIF2α occurs in *Plasmodium* in response to nutritional stress[23,24]. Additionally, increased levels of eIF2α phosphorylation are protective for DHA treatment[25]. In higher eukaryotes, phosphorylation of eIF2α can promote activation of the PI3K complex and increase transcription of autophagy-related genes[26,27]. In *Plasmodium*, increased PI3K activity mediates artemisinin resistance [28]. Further, polymorphisms in autophagy-related genes, such as ATG18 and ATG7, are associated with decreased artemisinin sensitivity in Southeast Asian parasite populations[29,30]. Recent evidence indicates the role of eIF2α phosphorylation in promoting autophagy is conserved in *Plasmodium*[31]. Thus, autophagy may be activated by our metabolic priming in a manner associated with DHA survival.

To explore the role of autophagy in priming, we used a parasite strain with conditional regulation of ATG8 expression; removal of the small molecule, anhydrotetracycline (aTc), leads to knockdown of ATG8, which is the essential for autophagosome formation and therefore essential for successful macroautophagy[32]. When we compared cumulative proliferation post-DHA between parasites with low versus normal levels of ATG8 expression (**Supplemental Fig. 4 A-B**), we detected a 3.6-fold reduction in the effect of hypoxanthine metabolic priming on survival when ATG8 is repressed (304% vs 85% proliferation difference 10d post-DHA; **Supplemental Fig. 4C-D**, **Supplemental Table 1**). Along with translational repression, this result suggests that autophagy contributes to priming-induced pro-survival phenotype. However, the effect is difficult to fully ascertain using this approach, given that autophagy plays a role in the recovery of DHA treatment[29,31]; this role is confirmed in our studies by the difference in post-drug recovery

time between non-primed parasites with and without ATG8 expression. Parasites expressing ATG8 surpass 5% parasitemia about 4.2 days earlier than parasites with reduced ATG8 expression following a 200nM DHA pulse (ATG8 On: 7.3d ± 0.7d (S.E.M.), ATG8 Off: 11.5d ± 0.5d, **Supplemental Fig. 4E**)

Physiological growth conditions promote increased DHA survival
Nutrient limitation is a physiologically relevant stress that parasites must adapt to in the context of human infection. Our low hypoxanthine and thiamine-free metabolic priming conditions mimic nutrient stress from a single nutrient deficiency. While this type of sole micronutrient deficiency occurs *in vivo*[33,34], this is not the only relevant nutritional difference between in vitro cultures and clinical infections. For example, even when human hosts are not suffering micronutrient deficiencies, *in vivo* levels of not just one, but many, nutrients are dramatically lower compared to the supra-physiological levels in standard RPMI-media based formulations[12-14].

We hypothesized that the nutritional differences between RPMI and human plasma is sufficient to induce a metabolic priming-like effect similar to that seen with our established conditions. To test this, we utilized Human Plasma-Like Media (HPLM) as a base for *P. falciparum* culture[35]. Under routine propagation conditions, parasites maintained in HPLM grew equivalent to those in RPMI based media (**Fig. 4A**). HPLM and RPMI based cultures increased an average of 18 ± 1.9 and 19 ± 2.5 fold (mean ± S.E.M) per replication cycle, respectively.

We adapted parasites to HPLM based culture for at least 7 days prior to treating with DHA and kept parasites in HPLM for 10 days of recovery. Propagation in HPLM prior to DHA pulse led to increased cumulative proliferation during post-drug recovery compared parasites in RPMI based media (**Fig. 4B**, mean day 10 difference of 459%, *p* = 0.0037). As with low hypoxanthine and thiamine-free priming conditions, growth in HPLM also increased DHA survival in RSAs (**Fig. 4C**, mean difference of 1.4-fold including both 200 and 700nM DHA samples). These data provide evidence that related pro-survival changes are occurring in response to both single nutrient deficiency and broader reductions of environmental nutrients.

Metabolic priming shifts transcriptome toward artemisinin tolerance
To more fully explore pathways that are altered in response to nutritional environment, we assessed gene expression in synchronous ring stage non-primed parasites and parasites primed with low hypoxanthine, thiamine-free, or HPLM conditions. Principal component analysis showed that each group had a distinct transcriptional profile, with hypoxanthine primed parasites clustering separate from the other three groups on PC1 and the remaining groups separating on PC2 (**Fig. 5A**). This pattern parallels the number of significantly deferentially expressed (DE) genes

detected in experimental groups compared to non-primed RPMI-based control samples with hypoxanthine primed parasites yielding the largest number of differences (**Fig. 5B**).

To identify pathways contributing to the increases in survival, we performed Gene Set Enrichment Analysis (GSEA) across priming conditions (**Fig. 5C**). Some metabolic pathways not previously implicated in stress responses, but important for parasite biology, were altered in a condition-dependent manner. For example, pyrimidine biosynthesis and apicoplast metabolism were significantly down-regulated in in thiamine-free and HPLM groups, respectively (**Fig. 5C**). Given the possible impact of ATG8 knockdown on our phenotype (**Supplemental Fig. 4C-D**), we also assessed if the autophagy pathway was altered in primed samples. GSEA indicated autophagy was significantly upregulated only in a condition-specific manner in hypoxanthine samples but not across all groups (**Fig. 5C**), indicating that autophagy is unlikely to be a primary driver of the observed phenotype.

Overall, GSEA showed that 25 of 463 total pathways were significantly enriched across all groups (**Supplemental Table 2**, including pathways heavy in exported proteins (*e.g.* protein-protein interactions between *P. falciparum* and the red blood cell). Upregulation of exported proteins has been previously observed in both the artemisinin resistance transcriptome and as a response to nutrient limitation[36-38]. Translational regulation was also significantly upregulated in all three conditions in concordance with previous knowledge of the eIF2α-mediated response under nutrient stress (**Fig. 5C**). This pathway includes genes with roles in processes known to impact DHA survival including the unfolded protein response (UPR) and proteasomal degradation (**Supplemental Table 2**).

Due to the large number of DE genes following hypoxanthine priming, and relatively small number in thiamine-free samples, it was necessary to limit the complexity of our results to further assess individual transcripts contributing to our phenotype. We specifically focused on the top 15 up- and down-regulated genes in each group as determined by lowest adjusted p-values. This yielded 37 up- and 42 down-regulated unique genes across the three treatment conditions as there was some overlap between groups. From this list of genes, we identified those that had an adjusted p-value of ≤0.05 in all groups as a curated list of 26 up- and 20 down-regulated genes of interest (**Table 2**). This curated list contained many genes implicated in artemisinin genetic tolerance, such as *coronin* (PF3D7_1251200), a gene known to be sufficient to induce artemisinin tolerance when mutated[39]. When compared to a list of 151 genes known to be basally altered (without active DHA treatment; **Supplemental Table 3**) in genetically artemisinin tolerant parasites[39-43], 8 of our curated genes were represented (**Table 2**). Further, changes in this curated set of 151 genes (designated as "ART Tolerance" in **Fig. 5C**) were highly enriched by GSEA in all groups (in the top 1% of 463 pathways by adjusted p-value). These data

indicate that nutrient limitation remodels metabolism in parasites to induce an artemisinin tolerance-like state.

Discussion

In the current study, we have demonstrated that physiologically relevant nutrient limitation stimulates intrinsic pathways that increase *P. falciparum* survival of drug treatment (**Figs. 1-4**). Our analyses indicate genes within these altered pathways directly contribute to genetic artemisinin tolerance (**Fig. 5C**), thus we speculate that nutrient limitation offers a preamble for resistance evolution. Our findings connect to other discoveries in the field on cellular responses to both nutrient limitation and drug treatment and provide a framework for how the nutritional environment impacts pathogen homeostasis. We explore these connections and their relevance below.

Limiting nutrient levels for malaria parasites in vitro

The supra-physiological concentrations of nutrients found in common media formulations, such as RPMI-1640, can potentiate genetic, transcriptional, and metabolic changes in parasite biology (reviewed in [12,14]). Here, we observed improved parasite survival following limitation of single nutrients in *in vitro* growth media; hypoxanthine and thiamine were chosen in part due to contrasting metabolic attributes outlined in the *Results* section, but also due to potential relevance to conditions *P. falciparum* may experience *in vivo* (see further discussion below). This approach is similar to other studies that induced nutrient stress by reduction of a single amino acid[44,45] or select metabolite classes[24,31,46]. We also observed the survival boost when parasites were grown in HPLM, a physiological media formulation that has reduced concentrations of multiple nutrients[13]. To our knowledge, this is the first use of HPLM as a media base for the propagation of *P. falciparum*; to induce starvation in previous studies, nutrient-rich RPMI 1640 base media is diluted[46], producing a blanket reduction in nutrient levels. Regardless of the approach, the *P. falciparum* response to artemisinin treatment, when not protected by this supra-physiological glut of nutrients, had not been previously evaluated *in vitro*. This study begins to connect many disparate observations to improve our understanding of how environment impacts cellular responses.

Molecular connections between nutrient limitation and DHA survival

We demonstrate that metabolic priming triggers broad alterations of translational regulation (**Fig. 5C**), including genes associated with the UPR and proteasomal degradation. Previous studies in *Plasmodium* connected nutrient limitation with translational regulation (via eIF2α phosphorylation[23,24]) and separately, eIF2α phosphorylation with artemisinin-induced latency[25]. Our study establishes a direct link between nutrient limitation and DHA survival. Additionally, our data support a role for autophagy in priming-induced DHA survival (**Supplementary Fig. 4C-D** and

Fig. 5C). Given that the activation of stress-induced autophagy in *P. falciparum* and higher eukaryotes is partially dependent on translational regulation by eIF2α phosphorylation[31,47], the connection between nutrient limitation, eIF2a-autophagy, and DHA survival is growing stronger. However, the extent to which eIF2α phosphorylation-dependent latency is mediating this phenotype remains to be explored.

A surprising outcome of our study was the discovery that nutrient limitation stimulated an artemisinin tolerance-like transcriptional state. This adaptive response, which was stimulated by all priming conditions (**Fig. 5**), triggered a change in expression for genes that had previously been linked to genetic artemisinin tolerance (**Table 2**). Future investigations should determine the longevity of metabolic priming changes for DHA survival and whether these adaptive changes facilitate development of genetic tolerance. This phenomenon has been observed in the development of antibiotic resistant bacteria[48] and recently proposed for *Plasmodium* artemisinin tolerance by Zhu, *et al*.[38]. Of note, more than half of the top transcriptional markers for resistance identified by Zhu, *et al*. were significantly altered (by adjusted p-value) in the same direction in at least one of our priming conditions (32/58 genes).

Translation from in vitro to in vivo results
Analyses of how the *in vitro* environment affects experiments performed on organisms outside of the native environment remains limited. Metabolic driven changes in pathogen drug sensitivity due to nutritional environment can lead to discordance between *in vitro* results and *in vivo* responses. Our data provide evidence of such an effect from nutritional environment on *P. falciparum* DHA survival between standard *in vitro* culture conditions and more *in vivo*-like environments (**Fig. 6**). Moving forward, use of physiological HPLM based media may be appropriate for selected assessments on the translatability of findings from culture conditions to clinical cases. Future studies should also investigate if environmentally-driven changes in survival are relevant to other antimalarial drugs and how any parasite survival benefits may be targeted to enhance the killing potential of malaria drug treatments.

Relevance for the in vivo host environment
Increasing resistance to artemisinin and its partner drugs necessitates development of new antimalarial drugs. Nucleotide metabolism is a potential future drug target in *P. falciparum*; antimalarials are currently being developed to inhibit both purine scavenging (including hypoxanthine) and pyrimidine biosynthesis[49-52]. As some of these candidates move closer to being approved in the clinic, it is imperative to understand how parasites respond to the stress of nucleotide deprivation. Similarly, thiamine limitation has potential physiological relevance to *P. falciparum* infection as clinical thiamine deficiency is common in malaria endemic regions, such as Southeast Asia[33,34], and evidence suggests malaria infection exacerbates this deficiency[53].

Additionally, metabolic network reconstruction data implies that drug resistance status can alter metabolism, making parasites more reliant on thiamine import for survival[54]. This is corroborated by increased expression in artemisinin tolerant isolates of thiamine pyrophosphokinase, a gene necessary for conversion of imported thiamine to the active thiamine pyrophosphate form[55].

Utilizing artemisinin with partner drugs of distinct mechanisms of action has been one key to the effectiveness of antimalarial drugs over recent decades. Partner drugs minimize the development of tolerance or resistance. Yet, the potential for partner drugs that target metabolism (as described above) to induce an artemisinin tolerant-like transcriptional state poses an unexpected challenge that should be considered in the development of new antimalarial combinations.

Combating anti-malarial drug resistance is especially critical as the spread of genetic artemisinin tolerance threatens to undo progress in reducing malaria mortality over recent decades[56]. However, the challenges of fighting drug resistance are not unique to malaria. The interplay between nutrient limitation, stress response pathways, and survival of drug treatment has important implications for the evolution of drug resistance across diverse pathogens. How microbes adjust their metabolism in ways that impact survival of current and future drugs is an important avenue of research to prevent and slow the spread of drug resistant pathogens.

Materials and Methods

Parasites and Growth

Plasmodium falciparum lines MRA 150 (Dd2), MRA 1000 (NF54), and MRA 1238 were obtained from the Malaria Research and Reference Reagent Resource Center (MR4, BEI Resources). The ATG8 conditional TetR-Dozi knockdown line was a generous gift from Ellen Yeh (Stanford University, Stanford, CA). *Plasmodium* cultures were maintained in A+ human erythrocytes (Valley Biomedical, Winchester, VA) at 3% hematocrit in RPMI 1640 HEPES (Sigma Aldrich, St Louis, MO) supplemented with 0.5% Albumax II Lipid-Rich BSA (Sigma Aldrich, St Louis, MO) and 50 mg/L hypoxanthine (Thermo Fisher Scientific, Waltham, MA). This formulation is referred to as "standard media" throughout the manuscript. ATG8 TetR-DOZI parasites were maintained with 0.5μM anhydrotetracycline (Sigma Aldrich, St Louis, MO) or supplemented with isopentyl pyrophosphate (Isoprenoids LC, Tampa, FL) as previously described[32]. Cultures were grown at 37°C and individually flushed with 5% oxygen, 5% carbon dioxide, 90% nitrogen gas. Dilution of cultures with uninfected erythrocytes and changing of culture medium was performed every other day. Parasitemia was determined by flow cytometry using SYBR Green I staining. Cultures were confirmed negative for mycoplasma approximately monthly using a LookOut Mycoplasma PCR detection kit (Sigma Aldrich, St Louis, MO).

Metabolic Priming

Low hypoxanthine media was made using RPMI 1640 HEPES base with reduced exogenous hypoxanthine addition to a final concentration of 0.5µM. Thiamine-free media was custom ordered to be identical to RPMI 1640 HEPES without the addition of thiamine hydrochloride. Hypoxanthine was added to thiamine-free media to 50 mg/L. Human Plasma-Like Media (HPLM, Thermo Fisher Scientific, Waltham, MA) was used without added HEPES or exogenous hypoxanthine. All media formulations were supplemented with 0.5% Albumax II Lipid-Rich BSA.

Uninfected erythrocytes were pre-incubated prior to use in priming experiments in the appropriate low nutrient medium for 48h at 37°C, 3-6% hematocrit, and 5% oxygen, 5% carbon dioxide, 90% nitrogen gas. Pre-incubated uninfected erythrocytes were seeded with infected erythrocyte culture to a starting parasitemia of <0.5% and resuspended in control medium for non-primed samples or low nutrient medium for primed samples. Cultures were allowed to incubate for the prescribed period of time dependent on the nutrient deprived (**Table 1**). For 72h incubations, media was refreshed at 48h.

Aliquots were taken for flow cytometry measurement with SYBR Green I and MitoProbe DiIC1(5) kit (both Thermo Fisher Scientific, Waltham, MA). Primed samples were compared to non-primed counterparts for reduction in growth, mitochondrial membrane potential (MMP), and percentage of ring stage parasites. Priming was considered successful when the following conditions were met for low hypoxanthine primed samples compared to non-primed: 1) an approx. >10% growth decrease, 2) little to no decrease in MMP (approx. <10% drop), and 3) ring stage percentages within approx. 10% of non-primed. Priming in thiamine-free and HPLM conditions was considered successful if both conditions "2" and "3" were met. Following successful priming, both primed and non-primed samples were treated with either 200nM dihydroartemisinin (DHA, Sigma Aldrich, St Louis, MO) or DMSO for 6h. To remove drug, cultures were washed three times in media and all samples were resuspended in normal, full-nutrient RPMI + Albumax media and returned to the incubator. Cultures were measured every other day with SYBR Green I and MitoProbe for 10-14 days post drug treatment.

Biological replicates were performed in technical duplicate. For statistical analysis, growth data 10 days post-DHA was log transformed and tested for normality by Shapiro-Wilks test prior to running a repeated measures two-way ANOVA followed by Sidak's multiple comparisons test. RSA and TSA data was tested by ordinary two-way ANOVA followed by Dunnett's multiple comparisons test. Analysis and visualizations were made using GraphPad Prism version 7.04.

Ring and Trophozoite Stage Survival Assays
Survival assays were performed as previously described with minor modifications[57]. Briefly, all metabolically primed samples were suspended in the relevant media at 36h, 72h, or ≥7d prior to

drug treatment for thiamine-, hypoxanthine-, and HPLM-primed conditions, respectively. Approximately 30h prior to 0-3h ring generation, 35mL cultures were synchronized with 5% D-sorbitol and allowed to progress until the culture was predominantly schizonts. Schizonts were isolated by layering 4 mL of culture over 4 mL of 75% Percoll, centrifugation, and collection of the intermediate band. Isolated schizonts were washed then added to uninfected erythrocytes in the appropriate media formulation. Exactly three hours later, a rapid D-sorbitol synchronization was performed (10m at 37 °C followed by 5s vortex) to remove any uninvaded late-stage parasites. Parasites for TSAs were returned to the incubator for exactly 24h before drug treatment, whereas parasites for RSAs were immediately pulsed with either DMSO or DHA (200nM or 700nM) for 6h. Following drug treatment, all cells were washed multiple times and returned to RPMI + ALB media for 66h prior to assessment of survival by flow cytometry measurement with SYBR Green I and MitoProbe DiIC1(5) kit.

Immunoblotting
Anti-ATG8 antibody was a gift from Ellen Yeh (Stanford University, Stanford, CA). Anti-ATG8 was used in conjunction goat Anti-guinea pig IgG H&L (Alexa Fluor® 488) (Abcam, Waltham, MA). Bound antibodies were detected on a Bio-Rad Imager and quantified using Image Lab Software (Bio-Rad, Hercules, CA).

RNAseq
Ring stage samples were generated by synchronization with 5% sorbitol 88h and 44h prior to harvest to accommodate the ~44h erythrocytic cycle duration of Dd2[58]. Primed sample groups were grown as described above in their designated media for 72h, 36h, and 7d prior to extraction for hypoxanthine, thiamine, and HPLM groups, respectively. Starting 24h prior to RNA extraction, blood smears were assessed every 3 hours until reinvasion was observed to approximate the age of rings in final samples. In all RNAseq samples, reinvasion was first observed 18h prior to harvest. At sample collection, erythrocytes were lysed in 0.15% saponin before RNA was extracted from parasite pellets using a Direct-zol RNA Miniprep Kit (Zymo Research) according to the manufacturer's instructions. An aliquot of each sample was taken for assessment with a 2100 Bioanalyzer using the RNA 6000 Pico assay (Agilent Technologies, USA) before the remainder of sample was snap frozen in liquid nitrogen and stored at -80C.

Samples were shipped to Genewiz, Inc. for standard RNAseq with polyA selection. Raw reads obtained from Genewiz, Inc. were checked for overall quality and trimmed to remove adapters and low-quality bases from 3' ends (with parameter -q 25) using TRIMGALORE (v. 0.6.7). STAR (v. 2.7.9a) was used to align trimmed reads with parameters – genomeSAindexNbases 11 and – alignIntronMax 500. Featurecounts (v. 2.0.1) was used for gene counting with parameter -t 'exon'. Differential expression analysis was done using DESeq2 (v. 1.34.0). Genes were considered

significantly differentially expressed if the following criteria were met 1) Benjamini-Hochberg adjusted p-value of ≤0.05 and 2) log2-fold change of ≥1.0 or ≤ -1.0. Gene set enrichment analysis (GSEA) was performed using fgsea (v. 1.20.0). 3D7 genome sequence for alignment and annotation, gene product descriptions, and metabolic pathway information were pulled from PlasmoDB (Release 55). Visualizations and analyses were generated in R version 4.1.2 using Tidyverse (v. 1.3.1) packages.

Acknowledgements

Special thanks to Ellen Yeh (Stanford University) for kindly providing us with the ATG8 Tet-R DOZI parasite line and ATG8 antibody and to Manoj Duraisingh (Harvard T.H. Chan School of Public Health) for facilitating initial trials of parasite growth in HPLM. This project was supported with funding through NSF CAREER Award No. 2046472. ACB was supported by an institutional training grant (5T32GM008136-33) and a fellowship from the ARCS Metro-Washington Chapter.

Tables and Figure Legends

Table 1. **Length and concentration of nutrient restrictions for metabolic priming.** *Priming was considered successful for this nutrient when an approximately 10-50% growth decrease occurred, there was little to no decrease in mitochondrial membrane potential (MMP) (approx. <10% drop), and ring stage percentages were within <10% ± non-primed. **Priming was considered successful for these nutrients when the latter two above conditions (MMP and staging) were met but no growth reduction criteria were required.

Nutrient Restricted	Class of Restricted Nutrient	Conc. of Restricted Nutrient	Length of Restriction
Hypoxanthine*	Purine	0.5uM	48-72h
Thiamine**	Cofactor	0.0uM	36h
Multiple (HPLM)**	Multiple	Variable	≥ 7d

Table 2. Genes of Interest Curated from Differentially Expressed Gene Lists. Asterisk indicates genes found to be basally altered in artemisinin mutants (from **Supplemental Table 3**).

Gene ID	Product Description
Upregulated	
*PF3D7_0214600**	serine/threonine protein kinase STK2, putative
PF3D7_0402200	surface-associated interspersed protein 4.1 (SURFIN 4.1), pseudogene
PF3D7_0418600	regulator of chromosome condensation, putative
PF3D7_0613900	myosin E, putative
PF3D7_0618000	conserved Plasmodium membrane protein, unknown function
PF3D7_0707300	rhoptry-associated membrane antigen
PF3D7_0724900	kinesin-20, putative
*PF3D7_0817600**	conserved protein, unknown function
*PF3D7_0822900**	PhIL1-interacting candidate PIC2
PF3D7_0903600	conserved protein, unknown function
PF3D7_0911100	START domain-containing protein, putative
PF3D7_0919900	regulator of chromosome condensation-PP1-interacting protein

PF3D7_1026600	conserved Plasmodium protein, unknown function
*PF3D7_1035300**	glutamate-rich protein GLURP
PF3D7_1126700	conserved Plasmodium protein, unknown function
PF3D7_1143100	AP2 domain transcription factor AP2-O
*PF3D7_1145200**	serine/threonine protein kinase, putative
*PF3D7_1206300**	conserved Plasmodium protein, unknown function
*PF3D7_1223100**	cAMP-dependent protein kinase regulatory subunit
*PF3D7_1251200**	coronin
PF3D7_1252400	reticulocyte binding protein homologue 3, pseudogene
PF3D7_1327300	conserved Plasmodium protein, unknown function
PF3D7_1335400	reticulocyte binding protein 2 homologue a
PF3D7_1356800	serine/threonine protein kinase ARK3, putative
PF3D7_1371600	erythrocyte binding like protein 1, pseudogene
PF3D7_1423300	serine/threonine protein phosphatase 7
Downregulated	
PF3D7_0308000	DNA polymerase delta small subunit, putative
PF3D7_0510100	KH domain-containing protein, putative
PF3D7_0512800	conserved Plasmodium protein, unknown function
PF3D7_0711000	AAA family ATPase, CDC48 subfamily
PF3D7_0715900	cation diffusion facilitator family protein, putative
PF3D7_0801100	28S ribosomal RNA
PF3D7_0802200	1-cys peroxiredoxin
PF3D7_0923800	thioredoxin reductase
PF3D7_1130900	conserved Plasmodium protein, unknown function
PF3D7_1239200	AP2 domain transcription factor, putative
PF3D7_1303800	conserved Plasmodium protein, unknown function
PF3D7_1322300	BRCT domain-containing protein, putative
PF3D7_1330600	elongation factor Tu, putative
PF3D7_1334500	MSP7-like protein
PF3D7_1359600	conserved Plasmodium protein, unknown function
PF3D7_1371800	Plasmodium exported protein, unknown function
PF3D7_1419200	thioredoxin-like protein, putative
PF3D7_1440200	stromal-processing peptidase, putative
PF3D7_1443900	heat shock protein 90, putative
PF3D7_MIT04100	large subunit ribosomal RNA fragment D

Figure 1. Mild nutrient stress prior to DHA treatment increases post-drug recovery. A) The biphasic dose response to stress characteristic of hormesis. B) Paradigm used for metabolic priming followed by DHA challenge and recovery. C-D) Post-DHA recovery in standard media of *P. falciparum* primed in either low hypoxanthine or thiamine-free media. Bars represent S.E.M. of technical replicates within one representative experiment. Results from all independent assays are detailed in Supplemental Table 1 (hypoxanthine deprivation; *N*=5, thiamine deprivation; *N*=4).

Figure 2. Increases in post-DHA survival are mediated by rings. A) *P. falciparum* asexual blood stage replication cycle. DHA tolerance is highest in 0-3h rings. B-C) Percent survival for early rings (B, 0-3h, *N*=3-4) or trophozoites (C, 24-27h, *N*=1-2) measured 66h after treatment with 200 or 700nM DHA for 6h. Bars represent S.E.M. Parasite line: *Dd2*. RSA p-values are as follows: Non-primed vs. low hypoxanthine: 200nM = 0.031, 700nM = 0.056. Non-primed vs. thiamine free:

200nM = 0.038, 700nM = 0.073. TSA p-values are as follows. Non-primed vs. low hypoxanthine: 200nM = 0.407, 700nM = 0.930. Non-primed vs. thiamine free: 200nM = 0.796, 700nM = 0.999. P-values <0.1 are summarized with one asterisk (*). P-values <0.05 are summarized with two asterisks (**).

Figure 3. Metabolic priming effects are conserved in parasite lines carrying genetic DHA resistance. A) Post-DHA recovery of MRA-1238 line in standard media and primed in low hypoxanthine or thiamine free media. Bars represent S.E.M. of technical replicates within one representative experiment. B) Percent survival for early rings measured 66h after treatment with 700nM DHA for 6h (N=1-3). Bars represent S.E.M. Parasite line: *MRA-1238*. RSA p-values are as follows: Non-primed vs. low hypoxanthine: 700nM = 0.012. Non-primed vs. thiamine free: 700nM = 0.026. P-values <0.1 are summarized with one asterisk (*). P-values <0.05 are summarized with two asterisks (**).

Figure 4. Physiologic nutrient media increases post-DHA recovery. A) Average growth rate of *Dd2* parasites propagated in RPMI 1640 versus HPLM based media. N=10 representing 5 replication cycles of two independent growth assays. B) Post-DHA recovery in standard media of *P. falciparum* primed in HPLM media. Bars represent S.E.M. of technical replicates within one representative experiment. Results from independent assays are detailed in Supplemental Table 1 (N=2). C) Percent survival for early rings measured 66h after treatment with 200 or 700nM DHA for 6h (N=1). Parasites were adapted to HPLM based media at least 7 days prior to drug treatment and maintained in HPLM through the RSA reading. Bars represent S.E.M. Parasite line: *Dd2*. RPMI = Roswell Park Memorial Institute Media. HPLM = Human Plasma-Like Media.

Figure 5. Metabolic Priming induces artemisinin tolerant-like transcriptional changes. A) PCA plot of gene expression from each condition. B) Volcano plots of differentially expressed genes for each condition. Log2 fold change thresholds set at -1 and +1. Adjusted p-value set to ≤0.05. C) GSEA for selected pathways. TRUE and FALSE = adjusted p-value of <0.05 and >0.05, respectively. RPMI = Roswell Park Memorial Institute Media. HPLM = Human Plasma-Like Media. GSEA = Gene Set Enrichment Analysis. PCA = Principal Component Analysis.

Figure 6. Schematic summary of priming induced artemisinin tolerance. Nutrient limitation directly impacts parasite survival following dihydroartemisinin (DHA) treatment. These conditions are more relevant in the human host than in standard in vitro conditions. Transcriptional changes induced by nutrient limitation mimic those that contribute to artemisinin tolerance.

References

Berney, M. & Berney-Meyer, L. Mycobacterium tuberculosis in the Face of Host-Imposed Nutrient Limitation. *Microbiology spectrum* **5**, doi:10.1128/microbiolspec.TBTB2-0030-2016 (2017).
Mancio-Silva, L. *et al.* Nutrient sensing modulates malaria parasite virulence. *Nature* **547**, 213-216, doi:10.1038/nature23009 (2017).

Mansour, S. C. *et al.* Bacterial Abscess Formation Is Controlled by the Stringent Stress Response and Can Be Targeted Therapeutically. *EBioMedicine* **12**, 219-226, doi:10.1016/j.ebiom.2016.09.015 (2016).

Tattoli, I. *et al.* Amino acid starvation induced by invasive bacterial pathogens triggers an innate host defense program. *Cell host & microbe* **11**, 563-575, doi:10.1016/j.chom.2012.04.012 (2012).

Calabrese, E. J. Hormesis: Path and Progression to Significance. *Int J Mol Sci* **19**, doi:10.3390/ijms19102871 (2018).

Rossnerova, A. *et al.* The Molecular Mechanisms of Adaptive Response Related to Environmental Stress. *Int J Mol Sci* **21**, 7053, doi:10.3390/ijms21197053 (2020).

Martínez, J. L. & Rojo, F. Metabolic regulation of antibiotic resistance. *FEMS Microbiology Reviews* **35**, 768-789, doi:10.1111/j.1574-6976.2011.00282.x (2011).

Yang, J. H., Bening, S. C. & Collins, J. J. Antibiotic efficacy-context matters. *Current opinion in microbiology* **39**, 73-80, doi:10.1016/j.mib.2017.09.002 (2017).

Dumoulin, P. C., Vollrath, J., Tomko, S. S., Wang, J. X. & Burleigh, B. Glutamine metabolism modulates azole susceptibility in Trypanosoma cruzi amastigotes. *eLife* **9**, doi:10.7554/eLife.60226 (2020).

Meylan, S. *et al.* Carbon Sources Tune Antibiotic Susceptibility in Pseudomonas aeruginosa via Tricarboxylic Acid Cycle Control. *Cell chemical biology* **24**, 195-206, doi:10.1016/j.chembiol.2016.12.015 (2017).

Herman, J. D. *et al.* A genomic and evolutionary approach reveals non-genetic drug resistance in malaria. *Genome biology* **15**, 511, doi:10.1186/preaccept-1067113631444973 (2014).

Brown, A. C. & Guler, J. L. From Circulation to Cultivation: Plasmodium In Vivo versus In Vitro. *Trends in parasitology* **36**, 914-926, doi:https://doi.org/10.1016/j.pt.2020.08.008 (2020).

Cantor, J. R. The Rise of Physiologic Media. *Trends Cell Biol* **29**, 854-861, doi:10.1016/j.tcb.2019.08.009 (2019).

LeRoux, M., Lakshmanan, V. & Daily, J. P. Plasmodium falciparum biology: analysis of in vitro versus in vivo growth conditions. *Trends in parasitology* **25**, 474-481, doi:10.1016/j.pt.2009.07.005 (2009).

Pethe, K. *et al.* A chemical genetic screen in Mycobacterium tuberculosis identifies carbon-source-dependent growth inhibitors devoid of in vivo efficacy. *Nat Commun* **1**, 57, doi:10.1038/ncomms1060 (2010).

Pillai, A. D. *et al.* Solute restriction reveals an essential role for clag3-associated channels in malaria parasite nutrient acquisition. *Molecular pharmacology* **82**, 1104-1114, doi:10.1124/mol.112.081224 (2012).

Müller, S. & Kappes, B. Vitamin and cofactor biosynthesis pathways in Plasmodium and other apicomplexan parasites. *Trends in parasitology* **23**, 112-121, doi:10.1016/j.pt.2007.01.009 (2007).

Ismail, H. M. *et al.* Artemisinin activity-based probes identify multiple molecular targets within the asexual stage of the malaria parasites Plasmodium falciparum 3D7. *Proceedings of the National Academy of Sciences of the United States of America* **113**, 2080-2085, doi:10.1073/pnas.1600459113 (2016).

Meunier, B. & Robert, A. Heme as Trigger and Target for Trioxane-Containing Antimalarial Drugs. *Accounts of Chemical Research* **43**, 1444-1451, doi:10.1021/ar100070k (2010).

Wang, J. *et al.* Haem-activated promiscuous targeting of artemisinin in Plasmodium falciparum. *Nature communications* **6**, 10111-10111, doi:10.1038/ncomms10111 (2015).

Witkowski, B. *et al.* Novel phenotypic assays for the detection of artemisinin-resistant Plasmodium falciparum malaria in Cambodia: in-vitro and ex-vivo drug-response studies. *The Lancet. Infectious diseases* **13**, 1043-1049, doi:10.1016/s1473-3099(13)70252-4 (2013).

Pholwat, S. *et al.* The Malaria TaqMan Array Card Includes 87 Assays for Plasmodium falciparum Drug Resistance, Identification of Species, and Genotyping in a Single Reaction. *Antimicrobial agents and chemotherapy* **61**, doi:10.1128/aac.00110-17 (2017).

Babbitt, S. E. *et al.* ≪em>Plasmodium falciparum≪/em> responds to amino acid starvation by entering into a hibernatory state. *Proceedings of the National Academy of Sciences* **109**, E3278, doi:10.1073/pnas.1209823109 (2012).

Fennell, C. *et al.* PfeIK1, a eukaryotic initiation factor 2α kinase of the human malaria parasite Plasmodium falciparum, regulates stress-response to amino-acid starvation. *Malaria Journal* **8**, 99, doi:10.1186/1475-2875-8-99 (2009).

Zhang, M. *et al.* Inhibiting the Plasmodium eIF2α Kinase PK4 Prevents Artemisinin-Induced Latency. *Cell host & microbe* **22**, 766-776.e764, doi:10.1016/j.chom.2017.11.005 (2017).

Choi, H. J. *et al.* A novel PI3K/mTOR dual inhibitor, CMG002, overcomes the chemoresistance in ovarian cancer. *Gynecologic Oncology* **153**, 135-148, doi:https://doi.org/10.1016/j.ygyno.2019.01.012 (2019).

Deegan, S., Saveljeva, S., Gorman, A. M. & Samali, A. Stress-induced self-cannibalism: on the regulation of autophagy by endoplasmic reticulum stress. *Cellular and molecular life sciences : CMLS* **70**, 2425-2441, doi:10.1007/s00018-012-1173-4 (2013).

Mbengue, A. *et al.* A molecular mechanism of artemisinin resistance in Plasmodium falciparum malaria. *Nature* **520**, 683-687, doi:10.1038/nature14412 (2015).

Breglio, K. F. *et al.* A single nucleotide polymorphism in the Plasmodium falciparum atg18 gene associates with artemisinin resistance and confers enhanced parasite survival under nutrient deprivation. *Malar J* **17**, 391, doi:10.1186/s12936-018-2532-x (2018).

Dwivedi, A. *et al.* Functional analysis of Plasmodium falciparum subpopulations associated with artemisinin resistance in Cambodia. *Malar J* **16**, 493, doi:10.1186/s12936-017-2140-1 (2017).

Ray, A. *et al.* Autophagy Underlies the Proteostasis Mechanisms of Artemisinin Resistance in P. falciparum Malaria. *mBio* **0**, e00630-00622, doi:10.1128/mbio.00630-22 (2022).

Walczak, M., Ganesan, S. M., Niles, J. C. & Yeh, E. ATG8 Is Essential Specifically for an Autophagy-Independent Function in Apicoplast Biogenesis in Blood-Stage Malaria Parasites. *mBio* **9**, e02021-02017, doi:10.1128/mBio.02021-17 (2018).

Johnson, C. R. *et al.* Thiamin deficiency in low- and middle-income countries: Disorders, prevalences, previous interventions and current recommendations. *Nutrition and health* **25**, 127-151, doi:10.1177/0260106019830847 (2019).

Whitfield, K. C. *et al.* High prevalence of thiamine (vitamin B1) deficiency in early childhood among a nationally representative sample of Cambodian women of

childbearing age and their children. *PLoS neglected tropical diseases* **11**, e0005814, doi:10.1371/journal.pntd.0005814 (2017).

Cantor, J. R. *et al.* Physiologic Medium Rewires Cellular Metabolism and Reveals Uric Acid as an Endogenous Inhibitor of UMP Synthase. *Cell* **169**, 258-272.e217, doi:10.1016/j.cell.2017.03.023 (2017).

Rawat, M. *et al.* Identification of Co-Existing Mutations and Gene Expression Trends Associated With K13-Mediated Artemisinin Resistance in Plasmodium falciparum. *Frontiers in Genetics* **13** (2022).

Tewari, S. G. *et al.* Short-term metabolic adjustments in Plasmodium falciparum counter hypoxanthine deprivation at the expense of long-term viability. *Malaria Journal* **18**, 86, doi:10.1186/s12936-019-2720-3 (2019).

Zhu, L. *et al.* Artemisinin resistance in the malaria parasite, Plasmodium falciparum, originates from its initial transcriptional response. *Communications Biology* **5**, 274, doi:10.1038/s42003-022-03215-0 (2022).

Demas, A. R. *et al.* Mutations in Plasmodium falciparum actin-binding protein coronin confer reduced artemisinin susceptibility. *Proceedings of the National Academy of Sciences of the United States of America* **115**, 12799-12804, doi:10.1073/pnas.1812317115 (2018).

Mok, S. *et al.* Artemisinin-resistant K13 mutations rewire Plasmodium falciparum's intra-erythrocytic metabolic program to enhance survival. *Nature Communications* **12**, 530, doi:10.1038/s41467-020-20805-w (2021).

Rocamora, F. *et al.* Oxidative stress and protein damage responses mediate artemisinin resistance in malaria parasites. *PLoS pathogens* **14**, e1006930, doi:10.1371/journal.ppat.1006930 (2018).

Siddiqui, G., Srivastava, A., Russell, A. S. & Creek, D. J. Multi-omics Based Identification of Specific Biochemical Changes Associated With PfKelch13-Mutant Artemisinin-Resistant Plasmodium falciparum. *The Journal of infectious diseases* **215**, 1435-1444, doi:10.1093/infdis/jix156 (2017).

Witkowski, B. *et al.* Increased tolerance to artemisinin in Plasmodium falciparum is mediated by a quiescence mechanism. *Antimicrobial agents and chemotherapy* **54**, 1872-1877, doi:10.1128/aac.01636-09 (2010).

Babbitt, S. E. *et al.* Plasmodium falciparum responds to amino acid starvation by entering into a hibernatory state. *Proc Natl Acad Sci U S A* **109**, E3278-3287, doi:10.1073/pnas.1209823109 (2012).

McLean Kyle, J., Jacobs-Lorena, M. & Miller Louis, H. Plasmodium falciparum Maf1 Confers Survival upon Amino Acid Starvation. *mBio* **8**, e02317-02316, doi:10.1128/mBio.02317-16 (2017).

Nair, S. *et al.* Nutrient Limitation Magnifies Fitness Costs of Antimalarial Drug Resistance Mutations. *Antimicrobial agents and chemotherapy* **66**, e01529-01521, doi:10.1128/aac.01529-21 (2022).

Humeau, J. *et al.* Phosphorylation of eukaryotic initiation factor-2α (eIF2α) in autophagy. *Cell Death Dis* **11**, 433-433, doi:10.1038/s41419-020-2642-6 (2020).

Fridman, O., Goldberg, A., Ronin, I., Shoresh, N. & Balaban, N. Q. Optimization of lag time underlies antibiotic tolerance in evolved bacterial populations. *Nature* **513**, 418-421, doi:10.1038/nature13469 (2014).

Cassera, M. B., Zhang, Y., Hazleton, K. Z. & Schramm, V. L. Purine and pyrimidine pathways as targets in Plasmodium falciparum. *Curr Top Med Chem* **11**, 2103-2115, doi:10.2174/156802611796575948 (2011).

Evans, G. B., Tyler, P. C. & Schramm, V. L. Immucillins in Infectious Diseases. *ACS infectious diseases* **4**, 107-117, doi:10.1021/acsinfecdis.7b00172 (2018).

Phillips, M. A. *et al.* A long-duration dihydroorotate dehydrogenase inhibitor (DSM265) for prevention and treatment of malaria. *Science translational medicine* **7**, 296ra111-296ra111, doi:10.1126/scitranslmed.aaa6645 (2015).

Llanos-Cuentas, A. *et al.* Antimalarial activity of single-dose DSM265, a novel plasmodium dihydroorotate dehydrogenase inhibitor, in patients with uncomplicated Plasmodium falciparum or Plasmodium vivax malaria infection: a proof-of-concept, open-label, phase 2a study. *The Lancet Infectious Diseases* **18**, 874-883, doi:https://doi.org/10.1016/S1473-3099(18)30309-8 (2018).

Mayxay, M. *et al.* Thiamin deficiency and uncomplicated falciparum malaria in Laos. *Tropical medicine & international health : TM & IH* **12**, 363-369, doi:10.1111/j.1365-3156.2006.01804.x (2007).

Carey, M. A., Papin, J. A. & Guler, J. L. Novel Plasmodium falciparum metabolic network reconstruction identifies shifts associated with clinical antimalarial resistance. *BMC genomics* **18**, 543, doi:10.1186/s12864-017-3905-1 (2017).

Mok, S. *et al.* Artemisinin resistance in Plasmodium falciparum is associated with an altered temporal pattern of transcription. *BMC genomics* **12**, 391, doi:10.1186/1471-2164-12-391 (2011).

WHO. World Malaria Report. (World Health Organization, Geneva, Switzerland, 2021).

Witkowski B, M. D., Amaratunga C, Fairhurst RM. Ring-stage Survival Assays (RSA) to evaluate the in-vitro and ex-vivo susceptibility of Plasmodium falciparum to artemisinins. (Institute Pasteur du Cambodge – National Institutes of Health Procedure RSAv1, 2013).

Reilly, H. B., Wang, H., Steuter, J. A., Marx, A. M. & Ferdig, M. T. Quantitative dissection of clone-specific growth rates in cultured malaria parasites. *International journal for parasitology* **37**, 1599-1607, doi:10.1016/j.ijpara.2007.05.003 (2007).

A

Survival / Death — Drug Tolerance

Stress →

B

Mild nutrient starvation stress

Antimalarial drug stress

Low Nutrient	200nM DHA	Monitor Proliferation & Viability Daily
36-72h	6h	10-14 days

C

Hypoxanthine Deprivation

Normalized Proliferation

— Primed
● Non-Primed

K13 WT

Days post-DHA

D

Thiamine Deprivation

Normalized Proliferation

— Primed
● Non-Primed

K13 WT

Days post-DHA

A

Hypoxanthine Deprivation

Primed

Non-Primed

K13 MT

Normalized Proliferation

Days post-DHA

B

RSA 0-3 h.p.i.

Non-Primed

Hypoxanthine

Thiamine

Percent Survival

**

**

700nM DHA