The development of intracytoplasmic membranes in alphaproteobacteria involves the conserved mitochondrial crista-developing Mic60 protein

Sergio A. Muñoz-Gómez[1,*,+], Lawrence Rudy Cadena[2,3,*], Alastair T. Gardiner[4], Michelle M. Leger[5], Shaghayegh Sheikh[2,3], Louise Connell[6], Tomáš Bilý[2], Karel Kopejtka[4], J. Thomas Beatty[7], Michal Koblížek[4], Andrew J. Roger[8], Claudio H. Slamovits[8], Julius Lukeš[2,3], Hassan Hashimi[2,3,+]

Affiliations

[1]Ecologie Systématique Evolution, Université Paris-Saclay, AgroParisTech, Orsay, France.
[2]Institute of Parasitology, Biology Center, Czech Academy of Sciences, České Budějovice (Budweis), Czech Republic.
[3]Faculty of Science, University of South Bohemia, České Budějovice (Budweis), Czech Republic.
4Center Algatech, Institute of Microbiology, Czech Academy of Sciences, Třeboň, Czech Republic.
[5]Institute of Evolutionary Biology (CSIC-Universitat Pompeu Fabra), Barcelona, Catalonia, Spain.
[6]Department of Chemistry and Biomolecular Sciences, University of Ottawa, Ottawa, Canada.
[7]Department of Microbiology and Immunology, University of British Columbia, Vancouver, Canada.
[8]Centre for Comparative Genomics and Evolutionary Bioinformatics, Department of Biochemistry and Molecular Biology, Dalhousie University, Halifax, Nova Scotia, Canada.

*Equal contribution
+Correspondence to: sergio.munoz@universite-paris-saclay.fr; hassan@paru.cas.cz

Abstract

Mitochondrial cristae expand the surface area of respiratory membranes and ultimately allow for the evolutionary scaling of respiration with cell volume across eukaryotes. The discovery of Mic60 homologs among alphaproteobacteria, the closest extant relatives of mitochondria, suggested that cristae might have evolved from bacterial intracytoplasmic membranes (ICMs). Here, we investigated the predicted structure and function of alphaproteobacterial Mic60, and a protein encoded by an adjacent gene Orf52, in two distantly related purple alphaproteobacteria, *Rhodobacter sphaeroides* and *Rhodopseudomonas palustris*. In addition, we assessed the potential physical interactors of Mic60 and Orf52 in *R. sphaeroides*. We show that the three α-helices of mitochondrial Mic60's mitofilin domain, as well as its adjacent membrane-binding amphipathic helix, are present in alphaproteobacterial Mic60. The disruption of Mic60 and Orf52 caused photoheterotrophic growth defects, which are most severe under low light conditions, and both their disruption and overexpression lead to enlarged ICMs in both studied alphaproteobacteria. We also found that alphaproteobacterial Mic60 physically interacts with BamA, the homolog of Sam50, one of the main physical interactors of eukaryotic Mic60. This interaction, responsible for making contact sites at mitochondrial envelopes, has been conserved in modern alphaproteobacteria despite more than a billion years of evolutionary divergence. Our results suggest a role for Mic60 in photosynthetic ICM development and contact site formation at alphaproteobacterial envelopes. Overall, we provide support for the hypothesis that mitochondrial cristae evolved from alphaproteobacterial ICMs, and therefore have improved our understanding of the nature of the mitochondrial ancestor.

Keywords: *Cereibacter*, *Rhodobacter*, *Rhodopseudomonas*, chromatophores, eukaryote, endosymbosis, MICOS.

Introduction

Mitochondria are organelles inferred to have been present in the last common ancestor of all eukaryotes (reviewed in Roger et al. 2017). Unlike most other organelles (e.g., the endoplasmic reticulum, nucleus, cytoskeleton, etc.), the mitochondrion evolved from an endosymbiont most closely related to extant alphaproteobacteria (Muñoz-Gómez et al. 2022; Martijn et al. 2018; Fan et al. 2020). In aerobic eukaryotes, mitochondria produce most of the ATP of the cell through aerobic respiration, i.e., the harnessing of energy through the coupling of electron transport to chemiosmosis with oxygen as a terminal electron acceptor, in addition to compartmentalizing other metabolic pathways (Roger et al. 2017; Hammond et al. 2022). Because aerobic respiration occurs at internalized membranes that

can expand greatly, mitochondria allow for the proportional increase (or linear scaling) of respiration
with cell volume across eukaryotes (Schavemaker and Muñoz-Gómez 2022). Mitochondria are thus
one of the innovations that likely allowed many eukaryotes to achieve larger cell volumes coupled to
relatively fast growth rates, and ultimately opened new evolutionary trajectories. Elucidating the origin
of mitochondria and their respiratory membranes may shed light on the origin of eukaryotic cells.

The specialization of mitochondria as respiratory organelles is most clearly reflected in their internal
structure. The mitochondrial inner membrane (MIM) invaginates into specialized sub-compartments
called cristae, the structural hallmarks of the organelle (Daems and Wisse 1966; Mannella et al. 1994;
Perkins et al. 1997). The MICOS (Mitochondrial Contact Site and Cristae Organizing System)
complex and oligomers of the F_1F_O-ATP synthase are two of the most evolutionarily conserved factors
responsible for the development and shape of cristae (Zick et al. 2009; Wideman and Muñoz-Gómez
2016). Whereas ATP synthase oligomers bend crista membranes at their rims to produce diverse
crista shapes (Kühlbrandt 2019; Pánek et al. 2020), the MICOS complex creates both crista junctions
(CJs) and contact sites (CSs) that compartmentalize, stably anchor, and maintain cristae at the
mitochondrial envelope (Rampelt et al. 2017). These functions of the MICOS complex appear to be
largely conserved across phylogenetically disparate eukaryotes, such as in the animal *Homo sapiens*,
the fungus *Saccharomyces cerevisiae*, the land plant *Arabidopsis thaliana*, and the parasitic protist
Trypanosoma brucei (Kozjak-Pavlovic 2017; Wollweber et al. 2017; Michaud et al. 2016; Kaurov et al.
2018).

Studies on the evolutionary history of MICOS revealed that this multi-protein complex is ancestrally
present in all eukaryotes and predates the origin of mitochondria (Muñoz-Gómez et al. 2015; Huynen
et al. 2016). The central and scaffolding subunit of the MICOS complex, the Mic60 protein, traces
back to the *Alphaproteobacteria*, the group from which mitochondria descended. Indeed, Mic60
serves as a phylogenetic marker that is uniquely present in the *Alphaproteobacteria* and the
mitochondrial lineage (Muñoz-Gómez et al. 2022). While only the C-terminal signature mitofilin
domain of Mic60 is sufficiently conserved at the sequence level, the overall predicted secondary
structure of Mic60 has also been conserved in both mitochondrial and alphaproteobacterial homologs
(Muñoz-Gómez et al. 2015). In addition to having homologues of Mic60, many alphaproteobacteria
also develop either lamellar or vesicular intracytoplasmic membranes (ICMs) that house diverse
electron transport chains involved in methanotrophy, nitrification, and anoxygenic photosynthesis
(Drews 1991; Niederman 2006; Pinevich 1997; LaSarre et al. 2018; Sener et al. 2016). This raises the
possibility that Mic60 is involved in the development and/or stability of ICMs in alphaproteobacteria,
and that mitochondrial cristae evolved from ancestral alphaproteobacterial ICMs (Muñoz-Gómez et al.
2017). Support for the functional conservation of alphaproteobacterial Mic60 comes from its capacity
to bind and bend membranes *in vitro* and heterologously in the gammaproteobacterium *Escherichia*
coli (Tarasenko et al. 2017). However, the precise role of Mic60 has not yet been directly studied in
alphaproteobacteria, and thus the evolutionary relationship between cristae and ICMs remains
unknown.

Though early ideas linked cristae to ICMs based simply on morphological resemblance, the
iconography of the field (i.e., the aggregate of scientific diagrams) has mostly depicted cristae as
post-endosymbiotic adaptations of mitochondria (Muñoz-Gómez et al. 2017). To better understand
the function of Mic60 in alphaproteobacteria and the origin of mitochondrial cristae, we investigated
the role of *mic60*, and its adjacent gene *orf52*, in two distantly related purple alphaproteobacteria: the
vesicular ICM-developing *Rhodobacter* (*Cereibacter*) *sphaeroides* (*Rhodobacterales*) and lamellar
ICM-developing *Rhodopseudomonas palustris* (*Rhizobiales*). We first explored the sequence and
structural conservation of the alphaproteobacterial Mic60 homologue. We then experimentally
investigated the effects of the disruption and overexpression of Mic60 and Orf52 in photoheterotrophic
growth and ICM development. Finally, we assessed the higher-order assembly and physical
interactors of alphaproteobacterial Mic60 and Orf52.

Results

<u>The predicted secondary and tertiary structure of alphaproteobacterial Mic60 is conserved</u>

Previous studies have shown that the overall predicted secondary structure of Mic60 is conserved
both in eukaryotes and alphaproteobacteria (Muñoz-Gómez et al. 2015; Huynen et al. 2016). In the
former, Mic60 consists of an N-terminal pre-sequence (a targeting signal to the mitochondrion),
followed by a transmembrane segment, a central region with coiled-coils, and a C-terminal signature
mitofilin domain. The same motifs and domains are found in the same order in the
alphaproteobacterial Mic60 homolog, except for the N-terminal pre-sequence which is missing, as
expected. Recently, mitochondrial Mic60 has been shown to deform membranes, and thus likely to
introduce curvature at CJs (Hessenberger et al. 2017; Tarasenko et al. 2017). This membrane-
deforming capability depends on a lipid-binding site (LBS) that is found in between the central coiled-
coils and the C-terminal mitofilin domain. This LBS comprises two α-helices (LBS1 and LBS2), the
first of which is amphipathic and presumably inserts itself into the MIM (Hessenberger et al. 2017).
LBS1 is extremely important for the function of Mic60 as its removal or mutation leads to the loss of
membrane binding and deformation, and also to phenotypes quite similar to those obtained when the
entire *MIC60* gene is deleted in *S. cerevisiae* (Hessenberger et al. 2017).

To investigate whether the membrane-bending α-helices of eukaryotic Mic60 are present in its
alphaproteobacterial homologs, we first performed profile Hidden Markov Model (pHMM)-sequence
searches against the UniProtKB database. We also predicted the secondary structure of both
eukaryotic and alphaproteobacterial Mic60 homologs with JPred4 (Drozdetskiy et al. 2015). A detailed
inspection of both the alignments and predicted secondary structure revealed that
alphaproteobacterial Mic60 has retained the two α-helices that comprise the LBS of eukaryotic Mic60
(Fig. 1A). Helical wheel projections further show that LBS1 is amphipathic in both *R. sphaeroides* and
R. palustris (Fig.1B, C). In addition, the functionally critical amino acid position Phe573 in the yeast
Chaetomium thermophilum (Hessenberger et al. 2017) is largely conserved among
alphaproteobacteria (e.g., Phe327 in *R. sphaeroides*; see Fig. 1A). Next, we predicted the tertiary
structure of the Mic60 homologs of the yeast *Lachancea thermotolerans*, whose crystal structure was
recently partially solved (Bock-Bierbaum et al. 2022), and the alphaproteobacteria *R. sphaeroides* and
R. palustris using AlphaFold2 (Fig. 1D-G). The predicted structures confirm the presence of the two α-
helices (LBS) in the linker region between the central coiled-coils and the C-terminal mitofilin domain
(Fig. 1D, E). Furthermore, the predicted structures show that the three conserved α-helices that
comprise the mitofilin domain of eukaryotic Mic60 (α1-3M), as well as the last two small α-helices of
the central coiled-coil region (α2-3C), are also present in alphaproteobacterial Mic60 (Fig. 1D-G).
Structural alignments reveal that the C-terminal region of alphaproteobacterial Mic60 (i.e.,
LBS+mitofilin) largely overlap with that of its eukaryotic homolog (Fig. 1F, G). The major structural
differences between eukaryotic and alphaproteobacterial Mic60 homologs are that the former is, on
average, a longer protein with a larger segment of central coiled coils, and has a transmembrane
segment much closer to the N-terminus of the protein (Fig. 1D, E). These observations indicate that
alphaproteobacteria Mic60 has a largely conserved secondary and tertiary structure, and that it may
bind and bend membranes through a mechanism similar to that of its eukaryotic ortholog.

Figure 1. Evolutionary conservation of the secondary and tertiary structure of mitochondrial and alphaproteobacterial Mic60. A. Alignment of C-terminal signature mitofilin domain and its adjacent N-terminal region of representative alphaproteobacteria and eukaryotes. This alignment has the Mic60 amino acid sequence of *R. sphaeroides* as a reference and was obtained from JPred4 (Drozdetskiy et al. 2015). The amino acid colors follow a coloring scheme based on physicochemical properties, and the intensity of the color reflect evolutionary conservation of the site in the alignment. The red arrow points to the conserved Phe327 in the *R. sphaeroides* Mic60 homolog. Common gaps are hidden from the alignment as per JPred4 output. **B.** Helical wheel projection of the amphipathic helix that comprises LBS1 in *R. sphaeroides* as predicted by the HELIQUEST web server (Gautier et al. 2008). **C.** Helical wheel projection of the amphipathic helix that comprises LBS1 in *R. palustris* predicted as in B. **D.** Predicted tertiary structures of the Mic60 homolog of the alphaproteobacterium *R. sphaeroides*. **E.** Predicted tertiary structures of the Mic60 homolog of the yeast *L. thermotolerans*. α-helices are colored according to domain (mitofilin in red, LBS1 and LBS2 in blue, middle coiled coils in green, and transmembrane segment in orange) and follow the coordinates predicted by JPred4. **F.** Structural alignment of the C-terminal region of Mic60 homologs from the eukaryote *L. thermotolerans* (yellow) and the bacterium *R. sphaeroides* (red). **G.** Structural alignment of the C-terminal region of Mic60 homologs from the eukaryote *L. thermotolerans* (yellow) and the bacterium *R. palustris* (pink).

Alphaproteobacterial *mic60* is clustered with a syntenic neighboring gene, *orf52*

A survey of the genomic context of *mic60* in several alphaproteobacterial species revealed that *mic60* is genetically linked to genes involved in the heme biosynthesis pathway (Huynen et al. 2016). In most alphaproteobacterial genomes, *mic60* is downstream of *hemC* (hydroxymethylbilane synthase; HMBS) and *hemD* (uroporphyrinogen-III synthase; UROS), and upstream of a hypothetical protein-coding gene sometimes (mis)annotated as *hemY*. All four genes have the same orientation and are usually tightly clustered with little intergenic space in between them, which may suggest that they are functionally related or co-transcribed as part of the same operon. This is consistent with the regulatory requirements of Mic60 according to its hypothesized function in ICM development, which requires heme biosynthesis for the proper assembly of cytochromes (Muñoz-Gómez et al. 2017). Indeed, HemD is occasionally fused to Mic60 in members of the *Rhodospirilales, Kiloniellales*, and *Rhizobiales* (Fig. 2A) (Huynen et al. 2016). In the yeast *S. cerevisiae*, the MICOS complex has been reported to interact with the enzyme ferrochelatase (HemH in alphaproteobacteria) which catalyzes

the insertion of ferrous iron into protoporphyrin IX, the eighth and final step in heme biosynthesis
(Dietz et al. 2021).

Although, *hemC* and *hemD* are genuine enzymes of the heme biosynthesis pathway, the gene
downstream of *mic60* clearly does not encode a heme biosynthetic enzyme. This protein is usually
misannotated as HemY because it contains a conserved hemY_N domain (PF07219) at its N-
terminus. Confusingly, this domain is unrelated to genuine *hemY* (protoporphyrinogen IX oxidase),
which is rather uncommon among alphaproteobacteria. Instead of *hemY*, most alphaproteobacteria
use the product of *hemJ* to synthesize protoporphyrin IX (very few alphaproteobacteria have a
genuine *hemY* gene homologous to that of *E. coli* and *Bacillus subtilis*) (Kobayashi et al. 2014). Here,
the protein encoded by the gene downstream of *mic60* will be referred to as Orf52 based on its
predicted size of 52 kDa in *R. sphaeroides* 2.4.1. Like alphaproteobacterial Mic60, Orf52 is an integral
membrane protein, but it possesses two transmembrane segments instead of one at its N-terminus,
and seems to expose its bulk to the periplasmic space. Moreover, Orf52 contains several
tetratricopeptide repeat motifs, which are usually involved in protein-protein interactions and found in
proteins that are part of multiprotein complexes. In *R. sphaeroides*, the *mic60*□*orf52* gene pair is co-
transcribed as indicated by a transcription start site (TSS) in the intergenic region between *hemD* and
mic60 (Myers et al. 2020) (Fig 2A). In the magnetosome gene island (MAI) of *Magnetospirillum
gryphiswaldense*, *mic60* and *orf52* are also co-transcribed separately from their neighboring genes
(Dziuba et al. 2021). The conserved motif order and composition, as well as transcriptional coupling,
of Mic60 and Orf52 indicate that these proteins have structural roles and may physically interact with
each other at alphaproteobacterial envelopes.

<u>Mic60 and Orf52 have broad phylogenetic distributions that overlap with the distribution of ICMs in the</u>
<u>*Alphaproteobacteria*</u>

Prokaryotic homologs of Mic60 have been previously shown to be restricted to alphaproteobacterial
species, thus serving as a synapomorphy for the clade that comprises both mitochondria and the
Alphaproteobacteria (Muñoz-Gómez et al. 2015; Huynen et al. 2016; Muñoz-Gómez et al. 2022).
Moreover, phylogenetically diverse alphaproteobacteria are known to develop extensive bioenergetic
ICMs that house electron transport chains associated with physiological processes such as
methanotrophy, nitrification (nitrite oxidation), and (aerobic or anaerobic) anoxygenic photosynthesis
(Drews 1991; Niederman 2006; Pinevich 1997; Muñoz-Gómez et al. 2017; LaSarre et al. 2018).
However, it is unclear to what extent the presence of Mic60, which has been hypothesized to be
involved in ICM development (Muñoz-Gómez et al. 2017), correlates with the occurrence of ICMs. To
better understand the evolutionary relationship between Mic60 and ICMs, we investigated their large-
scale phylogenetic distributions in the *Alphaproteobacteria*. To do this, we searched the GTDB R207
database that currently comprises more than 7,684 alphaproteobacterial genomes, with pHMMs for
Mic60, Orf52, and markers for ICM-associated physiologies (Fig. 2B).

These analyses show that both Mic60 and Orf52 have a broad and dense phylogenetic distribution
that encompasses the much more sporadic distribution of reported ICMs in the *Alphaproteobacteria*
(Fig. 2B). Methanotrophy and nitrite oxidation are restricted to a very few genera (e.g., *Methylocella*,
Methylosinus, and *Methylocystis*, and *Nitrobacter*), whereas photoautotrophy (or the capacity to
harvest light's energy to fix carbon dioxide) is phylogenetically widespread (Fig. 2B). The presence of
these ICM-associated physiologies largely overlaps with those species reported to develop extensive
ICMs (Fig. 2B). Furthermore, the prediction of a photoautotrophic physiology in several species not
yet studied ultrastructurally suggests that these may also develop extensive ICMs. On the other hand,
phototrophs (represented by aerobic anoxygenic photoheterotrophs) have a much broader
phylogenetic distribution (Imhoff et al. 2018) (Fig. 2B), though they are not often associated with the
presence of extensive ICMs. It is known, however, that aerobic anoxygenic phototrophs can develop
less conspicuous ICMs under some environmental conditions (Iba et al. 1988). Altogether, these
phylogenetic patterns suggests that both Mic60 and Orf52 are ancestrally present in the
Alphaproteobacteria and are required by extant species that either have or lack the capacity to
develop ICMs.

Figure 2. Genomic context of *mic60* and phylogenetic distribution of Mic60, Orf52, and ICMs across the
Alphaproteobacteria. A. The genomic context for *mic60* in representative ICM-developing alphaproteobacteria
and their relatives. Black circles to right of species identifiers denote alphaproteobacteria that develop extensive
ICMs. The bent arrow denotes the TSS in *R. sphaeroides.* Representative taxa were selected manually and a
supermatrix of 117 single-copy marker genes was assembled with GToTree (Lee 2019). The compositional

heterogeneity of the supermatrix was decreased with ⌐trimmer and a phylogenetic tree was inferred with IQ-
TREE and the LG+C60+F+G4 model (Nguyen et al. 2015). **B.** A comprehensive phylogenetic tree of the
Alphaproteobacteria that displays the distributions of Mic60, Orf52, markers for ICM-associated physiologies, and
reported ICMs across a maximally diverse set of taxa down sampled from the GTDB R207 database (Parks et al.
2022). A supermatrix of single-copy marker genes was assembled with GTDB-Tk, a phylogenetic tree was
inferred with IQ-TREE (-fast mode), and the longest-branching taxa were identified and removed with TreeShrink
(Mai and Mirarab 2018). The down sampling was performed with Treemmer (Menardo et al. 2018) and
constrained in such a way that all taxa with ICMs and ICM-associated physiologies were retained. TreeViewer
was used to display the phylogenetic distributions of traits on the tree. Protein searches were performed with
hmmsearch of the HMMER suite using both Pfam and custom pHMMs (Mistry et al. 2021).

<u>Knock out of *mic60* and *orf52* significantly affects photoheterotrophic growth in *R. sphaeroides* and *R.</u>
<u>palustris</u>

Purple alphaproteobacteria develop extensive intracellular membranes (ICM) in the presence of light
and the absence of oxygen (Drews 1991; Drews and Golecki 2004; Niederman 2006). These ICMs
house the photosynthetic apparatus and electron transport chain, which is generally composed of
light-harvesting complexes 1 and 2 (LH1 and LH2), a type II reaction center (RC), a cytochrome bc_1, a
periplasmic cytochrome c_2, and an ATP synthase (Greening and Lithgow 2020). By means of
cytochrome bc_1, which is also shared with the respiratory chain, the photosynthetic chain creates a
proton motive force across the ICM that is harvested by the ATP synthase to produce ATP
(Niederman 2006). ICMs are often continuous with, but sometimes detached from (Noble et al. 2018),
the cytoplasmic membrane (CM) (Scheuring et al. 2014), just as cristae are continuous with the MIM.
In *S. cerevisiae* mitochondria, the disruption of Mic60 leads to functional defects such as decreased
growth rate under respiratory conditions (i.e., non-fermentable media) and increased production of
oxygen radicals (John et al. 2005; Warnsmann et al. 2022). If Mic60 is involved in the development of
photosynthetic ICMs, then its disruption should lead to growth defects in the absence of oxygen and
the presence of light (i.e., photosynthetic conditions).

To assess whether Mic60 and Orf52 have an impact on photoheterotrophic growth and ICM
development, we first knocked out these genes in two phylogenetically distant purple
alphaproteobacteria amenable to reverse genetics, *R. sphaeroides* and *R. palustris* (Jaschke et al.
2011; Jiao and Newman 2007). To create knock-out strains, suicide plasmids containing the knock-
out gene construct flanked by homologous stretches were transferred into both species via
conjugation using a suitable *E. coli* strain. After selection and counter-selection for a first and second
recombination to insert the suicide plasmid into the host genome, and excise it, respectively, these
knock-out strains (Δ*mic60* and Δ*orf52*) were confirmed by PCR assays using several sets of internal
and external primers that flank the knock-out gene construct junctions (Fig. S1A-D).

Purple alphaproteobacteria develop a moderate amount of ICMs under anoxia and high light intensity.
The Δ*mic60* strains had a slower photoheterotrophic growth rate relative to *mic60*[+]*orf52*[+] wild type
(WT) in both *R. sphaeroides* and *R. palustris* (Fig. 3A, 3C). Under these conditions, the Δ*orf52*
decreased photoheterotrophic growth significantly in *R. palustris* but not in *R. sphaeroides*.
Furthermore, purple alphaproteobacteria develop larger amounts of ICMs and upregulate the LH2
complex to increase light capture under anoxia and low light intensity. As expected, both Δ*mic60* and
Δ*orf52* strains displayed even slower photoheterotrophic growth rates in both *R. sphaeroides* and *R.*
palustris in low light (Fig. 3B, 3D. Interestingly, Δ*orf52* displayed much more severe decreases in
growth rate, and even lower yield at stationary phase, than Δ*mic60* in *R. palustris* (Fig. 3B). This is
opposite from what was observed in *R. sphaeroides* (Fig. 3D) and indicates that these proteins might
contribute differently to photoheterotrophic growth in these two distantly related species. In summary,
the slower photoheterotrophic growth rate of the knock-out strains, especially under low light when
larger amounts of ICMs are required, suggests that both Mic60 and Orf52 affect the development of
photosynthetic ICMs.

Figure 3. The disruption of the *mic60* and *orf52* genes causes defects in photoheterotrophic growth and
ICM development. A-D. Growth curves of *R. palustris* WT and Δ*mic60* and Δ*orf52* strains under high (**A**) and
low (**B**) light and *R. sphaeroides* under high (**C**) and low (**D**) light. Turbidity was measured at 565 nm and
expressed in arbitrary units given on the *y*-axis. Time points are expressed in hours on the *x*-axis. Growth stages
are color-coded according to the labels at the top. Colored asterisks show statistical significance of differences at
each time point between WT and either Δ*mic60* (blue) or Δ*orf52* (red). **E.** Exemplar TEM micrographs of each *R.*
palustris strain grown in low light. **F.** Column scatter plot of the ratio of *R. palustris* ICM area/whole cell area (*y*-
axis) measured from 20 cells imaged using TEM in *R. palustris* WT, knock-out (Δ*mic60* and Δ*orf52*) and
overexpression (Mic60↑ and Orf52↑) strains. Mean and standard deviation shown by middle bar and whiskers,
respectively. Values and asterisks above each column represent statistical significance of the difference of each
population in comparison to WT. **G.** Exemplar electron tomograms of each *R. sphaeroides* strain grown in low
light. Blue arrows point at aberrant ICMs. **H.** Column scatter plot of tubules/cell per section (*y*-axis) for *R.*
sphaeroides WT, knock-out (Δ*mic60* and Δ*orf52*) and overexpression (Mic60↑ and Orf52↑) strains. The number
of sections, cells and tubules for each cell lines are given in the table below the *x*-axis. Mean and standard
deviation shown by middle bar and whiskers, respectively. Asterisks above each column represent statistical
significance of the difference of each mutant population in comparison to WT. Related to Movie S1-S4. Scale
bars in (**D**) and (**F**), 100 nm. Statistical significance: *, P<0.05; **, P<0.01; ***, P<0.001; ****, P<0.0001. See also
Fig. S1.

Mic60 and Orf52 are involved in the formation of lamellar and vesicular ICMs

The disruption of mitochondrial Mic60 leads to structural defects such as the loss of CJs, the
detachment of cristae from the mitochondrial envelope, and elongated crista membranes (Rabl et al.
2009; Kaurov et al. 2018). Hypotheses about the role of Mic60 in ICM development postulate that this
protein might be responsible for creating ICM junctions (ICMJs) and CSs that compartmentalize ICMs
and anchor them to the bacterial envelope (Muñoz-Gómez et al. 2017). The loss of Mic60 and Orf52
from ICM-developing alphaproteobacteria might thus lead to the loss of ICMJs or detached and
elongated ICMs. Their overexpression, in contrast, might lead to a larger number of ICMJs or
branching and larger ICMs.

To directly address whether the gene products of *mic60* and *orf52* are involved in ICM formation, we
generated *R. palustris* and *R. sphaeroides* strains capable of overexpressing Mic60 and Orf52
(Mic60↑ and Orf52↑, respectively). These strains were verified by reverse transcription quantitative
PCR (RT-qPCR) upon induction for six hours with 1 mM isopropyl-β-D-thiogalactoside (IPTG). Both
Mic60↑ and Orf52↑ strains had an increase in gene expression of ~7⎕9-fold in *R. sphaeroides*, and of
about two-fold in *R. palustris*, relative to ex-conjugants grown in the absence of IPTG (Fig. S1C). The
knock-out and WT strains were grown under photoheterotrophic conditions and low light and IPTG-
induced for six hours (in case of overexpression ex-conjugants) to maximize ICM development. After
harvesting, the strains were cryopreserved, contrasted, and imaged using transmission electron
microscopy (TEM). Randomized TEM micrographs were blindly scored to quantitatively evaluate
phenotypes in accord with the cytoplasmic membrane-adjacent stacked lamellar ICMs of *R. palustris*
and the uniformly distributed vesicular ICMs of *R. sphaeroides* (LaSarre et al. 2018) (Fig. 3E-H, Movie
S1-4).

In *R. palustris*, the area occupied by the lamellar ICMs increased relative to the whole cell area in
both knock-out (Δ*mic60* and Δ*orf52*) and overexpression (Mic60↑ and Orf52↑) strains, as compared to
the WT (ratio ICM area/cell area=0.18±0.03) (Fig. 3E-F). This increase was statistically significant in
Mic60↑ (0.29±0.03) but not in Δ*mic60* (0.24±0.03). Both Δ*orf52* (0.32±0.03) and Orf52↑ (0.40±0.04)
demonstrated higher and statistically significant increases in ICM:cell area ratios when compared to
their Mic60 counterparts. Wavy ICM membranes and/or wide gaps between lamellae were often
observed in the knock-out strains (Fig 3E). In *R. sphaeroides*, we scored for the appearance of
elongated ICMs (i.e., tubules) per cell as these were rarely observed in the WT strain
(tubules/cell/section=0.04±0.02) (Fig 3G-H). Both Δ*mic60* (0.46±0.07) and Mic60↑ (0.82±0.13)
showed a statistically highly significant increase in the number of tubules, whereas these structures
were observed to a lesser extent in Δ*orf52* (0.54±0.19) and Orf52↑ (0.74±0.20); the most conspicuous
outliers (≥3 tubules/cell/section) were observed in the latter cells. Electron tomograms were used to
render representative images for *R. sphaeroides* cells whose ICM membranes were not well
contrasted (Fig 3G, Movie S1-4). This revealed the presence of elongated ICMs of various lengths
and volumes in all the mutant strains assayed, indicating an increase in ICM membrane area.
Furthermore, the periplasm of the mutants appears to be expanded compared to the WT.

The ultrastructural defects of the Δ*mic60* and Δ*orf52* strains of *R. sphaeroides* and *R. palustris* are
consistent with their lower photoheterotrophic growth dynamics. Namely, the Δ*orf52* strain exhibits
more pronounced phenotypes than the Δ*mic60* strains in *R. palustris*, whereas the opposite is true in
R. sphaeroides. We also observed that both the knock-out and overexpression strains led to ICM area
expansion. It is possible that both the disruption and overproduction of Mic60 lead to unregulated
membrane growth that, under the image analyses employed here, produce seemingly similar
phenotypes. The sporadic wide gaps observed between the ICM lamellae in *R. palustris*, and the
expanded periplasm in *R. sphaeroides*, may be a consequence of the loss of ICMJs between lamellae
and the loss of CSs with the outer membrane, respectively. However, improved sample preservation
and larger-scale volumetric electron microscopy will be required to quantitatively assess changes in
the shape and number of ICMJs, which potentially differ between knock-out and overexpression
strains. The expansion of photosynthetic ICMs is similar to the enlarged respiratory cristae observed
in *S. cerevisiae* and *T. brucei* mitochondria defective of Mic60, and to the enlarged magnetosomes
reported in the magnetotactic alphaproteobacterium *M. gryphiswaldense* upon deletion of the *mic60*
paralog found within the MAI (see Discussion). Together, the ultrastructural defects displayed by both
Δ*mic60* and Δ*orf52* suggest that these genes are involved in the development of photosynthetic ICMs
in purple alphaproteobacteria.

<u>Mic60 and Orf52 are assembled into a 250 kDa complex in the ICMs of *R. sphaeroides*</u>

Eukaryotic MICOS comprises 6-9 subunits in *H. sapiens*, *S. cerevisiae*, and *T. brucei*. Four of them,
namely Mic60, Mic10, Mic19 and Mic12, are inferred to have been ancestral to eukaryotes (Muñoz-
Gómez et al. 2015; Huynen et al. 2016). Furthermore, the MICOS complex has been suggested to
engage in interactions with a myriad of other proteins, such as the outer membrane β-barrel insertase
Sam50, the outer membrane translocase Tom40, and the mitochondrial intermembrane space
assembly protein Mia40 (Pánek et al. 2020; Muñoz-Gómez et al. 2015b). It is thus possible that

alphaproteobacterial Mic60 and Orf52 are part of a larger multi-protein complex at
alphaproteobacterial envelopes.

To investigate whether Mic60 and Orf52 form part of a multi-protein complex, *R. sphaeroides* was
conjugated with a suitable *E. coli* strain to transfer plasmids for the IPTG-inducible expression of
either Mic60 or Orf52 bearing a C-terminal hexa-histidine (6xHis) tag. The expression of Mic60-6xHis
and Orf52-6xHis was verified by Western blot analysis using anti-6xHis antibodies (Fig. 4A). Mic60-
6xHis was found to migrate as a ~62 kDa protein, in contrast to its theoretical 43.8 kDa molecular
weight; this is likely explained by the anomalous mobility of acidic proteins in SDS-PAGE (Guan et al.
2015). The two overexpression strains, i.e., Mic60-6xHis↑ and Orf52-6xHis↑, alongside the
Mic60⁺Orf52⁺ (WT) control strain, were disrupted by high pressure homogenization to allow for ICM
isolation (i.e., chromatophores) by differential centrifugation. The absorption spectra of the isolated
ICMs from each strain were essentially identical, which indicates that the overexpression of Mic60-
6xHis and Orf52-6xHis does not affect the RC-LH1:LH2 protein composition ratio of ICMs (Fig. 4B).

The isolated ICMs were resolved in blue native PAGE (BN-PAGE) gels and transferred onto a
membrane for probing with an anti-His antibody (Fig. 4C). Both Mic60-6xHis and Orf52-6xHis
incorporate into a ~250 kDa multi-protein complex, with the antibody signal absent from the WT lane.
Notably, the immunoreactive ~250 kDa-size band seems to correspond to a faint and similarly sized
Coomassie-stained band in the BN-PAGE gel. The proteins from four excised ~250 kDa bands were
eluted and analyzed by liquid chromatography-tandem mass spectroscopy (LC-MS/MS) to identify
proteins migrating in this region of the BN-PAGE gel (Fig. S2A). Mic60 and Orf52 were among the
175 top proteins identified with a mean intensity score >23 in all four independent biological replicates
and thus considered high-confidence interactors (Dataset 1). The top-three hits in the ~250 kDa band
corresponded to the RC complex subunits H, M, and L, which were previously shown to co-migrate
with Mic60 in the BN-PAGE gels (D'Amici et al. 2010). This is consistent with both Mic60 and Orf52
previously being detected in isolated ICM and/or in its developmental precursor, i.e., 'upper
pigmented bands' (UPB) (D'Amici et al. 2010; Jackson et al. 2012). These experiments thus show
that both Mic60 and Orf52 are part of a higher-order assembly complex of ~250 kDa present in
photosynthetic ICMs.

Figure 4. Mic60 and Orf52 assemble into a 250 kDa protein complex in *R. sphaeroides*. A. Immunoblot
verifying the expression of Mic60-6xHis and Orf52-6xHis. **B.** Absorption spectra (~280-950 nm scan, *x*-axis) of
ICMs isolated from Mic60-6xHis, Orf52-6xHis, and WT control strains. Relative absorbance given on *y*-axis and
spectra are normalized to the bacteriochlorophyll *a* Qx peak at 590 nm. Spectra are stacked on top of each other
and color-coded according to the legend on the upper right-hand corner. **C.** Protein complexes from detergent-
solubilized isolated ICMs as resolved by BN-PAGE. Scissor symbol indicates the ~250 kDa band excised for MS
analysis on left gel (see also Fig. S3A and Dataset 1). Right gel shows immunoblot demonstrating that Mic60-

6xHis and Orf52-6xHis assemble into a ~250 kDa band. **D.** Immunoblot of Mic60-6xHis and Orf52-6xHis AP
eluates showing that the 6xHis-tagged bait proteins were successfully purified from detergent-solubilized isolated
ICMs. For **A**, **C**, and **D**, molecular weight markers shown on left. All immunoblots use arrows to point at specific
antibody signals from Mic60-6xHis and Orf52-6xHis (absent for WT) controls whereas asterisks denoted non-
specific band used as loading control.

<u>Mic60 and Orf52 interact with the outer membrane β-barrel insertase BamA</u>

Among several interactions reported for the eukaryotic MICOS complex (e.g., with Tom40, Sam50,
Mia40), it was suggested that the Mic60-Sam50 interaction may have predated the origin of
mitochondria (Muñoz-Gómez et al. 2015b). This interaction is required for making CSs at the
mitochondrial envelope of phylogenetically disparate eukaryotes such as *H. sapiens*, *S. cerevisiae*,
and *T. brucei* (Kozjak-Pavlovic 2017; Wollweber et al. 2017; Kaurov et al. 2018), and the bacterial
homolog of Sam50, BamA, is a ubiquitous protein of bacteria surrounded by two membranes
(diderms). In these bacteria (e.g., *Proteobacteria*), BamA required for the assembly of β-barrels in the
outer membrane (Höhr et al. 2018; Doyle and Bernstein 2019; Voulhoux et al. 2003). Furthermore,
mic60 is genetically linked to *orf52* and co-transcribed as part of the same operon (Myers et al. 2020).
These observations and inferences prompted us to next investigate potential physical interactors of
alphaproteobacterial Mic60 and Orf52.

We isolated ICMs from *R. sphaeroides* and performed affinity purifications (AP) of Mic60-6xHis and
Orf52-6xHis. The ICMs were solubilized with a mixture of digitonin (1.5% w/v) and Triton X-100 (0.1%
v/v), and the cleared lysate was incubated with cobalt-coated Dynabeads to capture the 6xHis-tagged
baits. After extensive washing, the successful capture of Mic60-6xHis and Orf52-6xHis was confirmed
on a portion of beads taken for Western blot analysis with an anti-His antibody (Fig. 4D)—a mock AP
was done on *R. sphaeroides* WT as a control for non-specific binding. The remaining beads from
triplicate APs of Mic60-6xHis, Orf52-6xHis and the mock controls were trypsinized and analyzed by
LC-MS/MS. Protein enrichment in bait-APs in comparison to the mock control was quantified using
label-free quantification as previously described (Cadena et al. 2021). In each AP, 497 high-
confidence proteins were found (Dataset 2) based on mean Andromeda confidence scores >100 in
three biological replicates (Cox et al. 2011). Among these, proteins were considered to have true
physical interaction if they (1) had Log_2-transformed fold-enrichment and p-values larger than three
(Fig. 4A-B), (2) were part of the higher-order ~250 kDa complex (see above; Dataset 1), (3) were
previously reported in ICM and/or upper pigmented bands (or UPBs, which are ICM precursors; Fig.
5C-D), and (4) were predicted to have transmembrane segments or a β-barrel structure (Fig. 5C-D).

The data show that both Mic60-6xHis and Orf52-6xHis interact with BamA (Fig. 5, S2), the β-barrel
insertase subunit of the β-barrel assembly machine (BAM) complex (Voulhoux et al. 2003). Somewhat
unexpectedly, Mic60-6xHis and Orf52-6xHis do not interact with each other, as neither protein was
found to co-purify with either bait protein (Fig. 5A-B). We cannot rule out that the C-terminal 6x-His
tag may have interfered with their interaction, although this is unlikely due to its small size. Mic60 was
also not found to interact with the enzyme HemD, although these proteins have fused in some
alphaproteobacteria (Fig 2A) (Huynen et al. 2016). This was not unexpected as the HemD-Mic60
fusion protein is predicted to be membrane-anchored at its midpoint, having the HemD fragment
exposed to the cytoplasm and Mic60 fragment on the periplasm. Moreover, the high degree of
enrichment of Mic60 relative to BamA (Fig 5B, 5D) is consistent with the idea that this protein
engages in homotypic interactions and forms homooligomers as its eukaryotic counterparts (Bock-
Bierbaum et al. 2022; Hessenberger et al. 2017; John et al. 2005).

In contrast to Mic60, which exhibits a strong interaction solely with BamA, Orf52 also interacts with a
β-barrel protein annotated as Outer Membrane Efflux (OM Efflux) and a very small 4.8 kDa protein
with an N-terminal TMD that is encoded by the RSP_7517 locus (Fig. 5A, 5C; Fig. S3D). The former
belongs to the TolC protein family and was found to be more abundant in ICM than the UBP precursor
(Jackson et al. 2012), whereas the latter is among the most enriched proteins in the Orf52 AP. The
enrichment of RSP_7515 is surprising as it was not previously detected in ICMs or UPBs (Jackson et
al., 2012). It was however, detected in the ~250 kDa BN-PAGE band, albeit below our defined
threshold by a single peptide (Dataset 1). This may be explained by the fact that this protein is highly
hydrophobic and would only yield two peptides by trypsinization, making it less amenable to detection

by shotgun LC-MS/MS, as employed for analyzing the protein composition of the ~250 kDa BN-PAGE
band (Barrera and Robinson 2011) (Fig. S3D). In conclusion, the Sam50 homolog BamA appears to
be the only physical interactor of alphaproteobacterial Mic60 (Fig. 5E), and this interaction has been
conserved despite ~2 billion years of evolutionary divergence of modern alphaproteobacteria from
mitochondria.

Figure 5. The interactomes of Mic60 and Orf52 demonstrate that both proteins interact with the outer
membrane β-barrel protein BamA in *R. sphaeroides*. A-B. Volcano plots of proteins identified by MS in AP
eluates of Mic60-6xHis (**A**) and Orf52-6xHis (**B**), with -Log P-values for each protein (*y*-axis) plotted against Log$_2$-
tranformed fold-enrichment over WT negative control (*x*-axis); thresholds for each value for a protein considered
enriched interactors are indicated by dotted lines, demarking the shaded area of each plot. Baits and enriched
interactors are indicated by labelled and colored large dots, whereas other proteins are demarked by small, grey
dots. **C-D.** Summary of enriched interactors found in Mic60-6xHis (**C**) and Orf52-6xHis (**D**) APs showing the
same color scheme as in **A** and **B**. From the left of protein names: schemas depicting the domain architecture of
each interacting protein; its fold change in APs compared to WT negative controls; presence in upper pigmented
band (UPB) and ICM proteomes (Jackson et al., 2012); presence in ~250 kDa band proteome (Dataset 1);
theoretical molecular (Mol.) weight. Black dots indicate presence in the proteome; grey dot indicates presence
below our defined threshold; *, presence in ICM according to MS analysis of D'Amici *et al.* (2010). **E.** Schema of

Mic60 and Orf52 interactions in *R. sphaeroides*. Boxed contains a key to depicted proteins, with N- and C-termini
shown in red. Related to Fig. S2 and Dataset 2.

Discussion

The overarching aim of this study was to investigate the structure and function of alphaproteobacterial
Mic60, thereby testing the hypothesis that ICMs were transformed into cristae during the early
evolution of mitochondria (Muñoz-Gómez et al. 2017). To this end, we focused on two distantly
related purple alphaproteobacteria, *R. palustris* and *R. sphaeroides*. We showed that
alphaproteobacterial Mic60 has (1) a conserved tertiary structure and membrane-bending
amphipathic helix, (2) affects photoheterotrophic growth and ICM development and shape, (3) is part
of a higher-order 250 kDa multi-protein complex, and (4) physically interacts with the core BamA
subunit of the outer membrane BAM complex. Furthermore, we also showed that Orf52, which is
encoded by *mic60*'s neighboring gene, also affects photoheterotrophic growth, ICM development and
shape, and, although it does not interact with Mic60, it also physically interacts with the outer
membrane protein BamA.

These findings are consistent with previous observations and experiments that also support a
functional conservation of Mic60 and its involvement in the development of both vesicular and
lamellar ICMs. First, the expression profile of Mic60 follows the development of photosynthetic ICMs
in the absence of oxygen and presence of light (Callister et al. 2006). Second, Mic60 localizes to
photosynthetic ICMs in three phylogenetically disparate alphaproteobacteria, *R. sphaeroides*
(*Rhodobacterales*), *Rhodospirillum rubrum* (*Rhodospirillales*), and *R. palustris* (*Rhizobiales*) (D'Amici
et al. 2010; Jackson et al. 2012; Selao et al. 2011; Fejes et al. 2003), as indicated by the proteomes
of isolated ICMs. Third, *R. sphaeroides*' Mic60 is capable of binding and tubulating membranes *in
vitro*, and its heterologous overexpression induces ICM-like structures in *E. coli* (Tarasenko et al.
2017). Fourth, both *mic60* and *orf52* have paralogs in the MAI of the alphaproteobacterial genus
Magnetospirillum of the order *Rhodospirillales* (Muñoz-Gómez et al. 2015) whose disruption leads to
fewer and larger magnetosomes (Lohße et al. 2011; Lohße et al. 2014). Multiple sources of evidence
thus support the notion that the function of Mic60 has been conserved in alphaproteobacteria.

What is the precise role of Mic60 in ICM development? The capacity of Mic60 to bend membranes
(Tarasenko et al. 2017), presumably through its conserved amphipathic helix (see Fig. 1), suggests
that Mic60 induces curvature at ICMJs. Moreover, alphaproteobacterial Mic60 interacts with BamA,
the central subunit of the BAM complex and homolog of Sam50. Mitochondrial Sam50 interacts with
Mic60 through its intermembrane space-protruding POTRA domains, and it is likely that the same
type of interaction occurs in the periplasm of alphaproteobacteria. This ancient protein-protein
interaction suggests that, in addition to aiding the formation of ICMJs, alphaproteobacterial Mic60 is
also involved in the formation of CJs that anchor ICMs to the alphaproteobacterial envelope. Thus, it
appears that both the formation of CSs and CJs/ICMJs have been conserved in alphaproteobacteria
and mitochondria. The formation of mitochondrial CJs and CS by Mic60 are aided by Mic19
(Hessenberger et al. 2017; Bock-Bierbaum et al. 2022), but there is no evidence for a Mic19-like
protein in alphaproteobacteria; our pull-downs did not reveal proteins with Coiled-Coil-Helix-Coiled-
Coil-Helix (CHCH) motifs. It is plausible that CSs stabilize ICMs by providing an anchor to the OM at
points of high membrane curvature, i.e., ICMJs (Tang et al. 2020). This may contribute to the
biogenesis of ICMs by ensuring their continuity with the CM where protein complex subunits may first
be inserted and assembled, giving rise to the UPB. Moreover, the larger macromolecular complex that
Mic60 is part of in *R. sphaeroides* is reminiscent of the extended interaction network, ERMIONE, that
in *S. cerevisiae* mitochondria involves Mic60 (Wideman and Muñoz-Gómez 2016). It is possible that
BamA, through its interactions with Mic60, Orf52, OM Efflux, and RSP_7515, serves as the hub of a
larger protein-interaction network, that, for example, facilitates lipid transfer and protein export to the
outer membrane. Future studies are required to determine whether alphaproteobacterial Mic60 forms
homodimers and homotetramers as recently reported for fungal Mic60 (Bock-Bierbaum et al. 2022),
and elucidate the molecular mechanisms by which alphaproteobacterial Mic60 interacts with BamA.

The evidence for the structural and functional conservation of alphaproteobacterial Mic60 relative to
its mitochondrial homolog is most compatible with an evolutionary scenario in which ICMs and cristae
are homologous. This view implies that cristae most likely evolved from the ICMs developed by the

last common ancestor of mitochondria and its sister group, the *Alphaproteobacteria*; cristae thus have
a pre-endosymbiotic origin. If this hypothesis turns out to be correct, bioenergetic ICMs might have
pre-adapted the first mitochondrial ancestor to become an efficient bioenergetic or respiratory
organelle (Muñoz-Gómez et al. 2017). The widespread but sporadic phylogenetic distribution of ICMs
across the *Alphaproteobacteria* is then most likely explained by multiple independent losses. This is
conceivable as both cristae and ICMs are known to have been lost repeatedly as a result of
physiological specialization to different environments (e.g., switches to anaerobiosis in mitochondria
or to heterotrophy in photosynthetic bacteria). However, some alphaproteobacteria that have Mic60
homologs are not known to develop ICMs. In these, Mic60 might play a more general function, e.g.,
CS formation by interacting with BamA for lipid transfer or protein export, that is still required in the
absence of ICMs. This more general function might have been ancestral to Mic60 and agrees with the
observation that a distant homologue of Mic60 that lacks the conserved C-terminal mitofilin domain,
namely HemX, is restricted to and widespread in the *Gammaproteobacteria* (Huynen et al. 2016). It is
thus conceivable that the signature mitofilin domain first evolved in a HemX-like protein and that this
coincided with the origin of ICMs in a common ancestor of alphaproteobacteria and mitochondria.
Future endeavors focused on phylogenetically disparate alphaproteobacteria with and without ICMs
will shed light on the functions and mechanisms of Mic60 in prokaryotes. Altogether, the structural
and functional conservation of alphaproteobacterial Mic60 shown here, suggests a role of this protein
in curving membranes at ICMJs and making CSs at envelopes. Therefore, it is probable that the
mitochondrial ancestor was an ICM-bearing alphaproteobacterium.

MATERIALS AND METHODS

<u>Bacterial strains, media and growth conditions</u>

R. palustris TIE-1 and *E. coli* BW29427-λpir-RP4 were kindly provided by Dianne K. Newman
(California Institute of Technology) (Jiao et al. 2005). *R. sphaeroides* 2.4.1, *E. coli* S17-1- λpir, and *E.*
coli DH5α-λpir were kindly provided by Jeanette Johnson-Beatty (University of British Columbia). *R.*
palustris TIE-1 strain was grown both chemo- and photo-heterotrophically at 30 °C in YPS rich
medium (Jiao et al., 2005) unless otherwise noted. *R. palustris* TIE-1 was grown
photoheterotrophically at 30 °C on FEM minimal medium (Ehrenreich and Widdel 1994). *R.*
sphaeroides 2.4.1 strain was grown chemoheterotrophically at 30 °C in LB or RLB rich media (Jun et
al. 2014), unless otherwise noted, and photoheterotrophically at 30 °C on RCVBN minimal medium
(Beatty and Gest 1981). Table S1 lists all strains used this study.

For growth analysis *R. sphaeroides* and *R. palustris* mutants and WT strains were grown in front to
three incandescent 40 W lightbulbs at a temperature of 29 °C, with the distances of the culture tubes
from the light sources adjusted to allow the appropriate light intensities and temperature. The light
intensities were measured to be ~200 µmol photons $m^{-2} s^{-1}$ for high light and ~10 µmol photons $m^{-2} s^{-1}$
$^{-1}$ for low light. The latter was achieved by placing the two layers of approximately ~ 50 % neutral
density filter sheeting between the light source and culture tubes. Prior to growth analysis, cells were
grown in in rich media in triplicate (*R. sphaeroides*, RLB without antibiotics; *R. palustris*, YPS no
antibiotic for WT, 400 µg/ml kanamycin for mutants) until mid-log phase. At mid-log phase equivalent
amounts of cells (turbidity x volume = 20, i.e., if the turbidity value was equal to 5.10, then 3.92 ml of
cells were decanted) were removed from the growth tubes into 15 ml Falcon tubes and pelleted by
centrifugation. The pellet was washed once in minimal media and resuspended in 1 ml of the required
minimal media (*R. sphaeroides* 10 ml RCVBN + 8 ml RLB and *R. palustris* FEM). A 150 µl volume
was of each cell line suspension was inoculated in triplicate; this was time point 0. Turbidity
measurements were performed using a McFarland Densitometer DEN-1B that measures at λ = 565
±15 nm. Turbidity measurements were then taken every 6 h for a total experiment length of 72 h.

<u>DNA methods and plasmid construction</u>

Total genomic DNA (gDNA) was extracted from *R. sphaeroides* and *R. palustris* strains using the ZR
Bacterial DNA Miniprep Kit (Zymo Research) with a BIO101/Savant FastPrep FP120 high-speed bead
beater and a 30 min incubation at 60°C with 20 µl of proteinase K (20 mg/mL), or the Epicentre
MasterPure DNA Purification Kit (Epicentre Biotechnologies). All plasmid constructs were built with
the Gibson Assembly Master Mix or the NEB Builder HiFi DNA Assembly (both New England

Biolabs). Plasmids were isolated using the AxyPrep Plasmid Miniprep Kit (Axygen). All plasmid
constructs were confirmed by Sanger sequencing using both forward and reverse primers. Table S2
lists all plasmids used. Table S3 lists all primers used in this study.

Construction of *R. sphaeroides* and *R. palustris* knock-out strains

To knock out genes in *R. sphaeroides*, a knock-out construct was assembled into the suicide plasmid
vector pZDJ (Brimacombe et al. 2013) with either the Gibson Assembly Cloning Kit or the NEBuilder
HiFi DNA Assembly Master Mix (both New England Biolabs). The antibiotic resistance cassette
flanked by flippase recognition target (FRT) sites used to interrupt the genes to be knocked out
corresponds to that used in the Keio collection (Baba et al. 2006). The resulting suicide vector with
the knock-out construct was cloned into *E. coli* S17-1 λ-pir-RP4, which can replicate the suicide
plasmid. This strain was then conjugated with *R. sphaeroides 2.4.1*. Briefly, donor and recipient cells
grown to stationary phase were mixed in a 1:2 volume ratio and pelleted by centrifugation for 1 min at
4,500-6,000 × g, and then washed twice with antibiotic-free RCVBN minimal medium. After the last
wash, the pellet was resuspended in 50 µl of RCVBN, and 10 µl aliquots were spotted onto antibiotic-
free RCVBN solid medium. The plates were incubated at 30° C overnight to allow conjugation to take
place. Afterwards, an emulsion was made from the several inoculation spots and streaked onto
antibiotic-containing RCVBN+Gm solid medium. *R. sphaeroides* exconjugants were then successively
streaked onto new LB+Gm solid medium until no *E. coli* S17-1 λ-pir-RP4 contamination remained.
The resulting pure *R. sphaeroides* exconjugants contained the pZDJ plasmid, with the knock-out gene
construct, integrated in the chromosome by a first crossing over (recombination) event. To induce a
second crossing over to excise the pZDJ suicide plasmid from the host chromosome, colonies were
picked and grown on liquid LB with no antibiotic selection until late stationary phase (about 2-3 days).
The cultures were then streaked on LB+10% sucrose solid medium, which allows for counter-
selecting of those colonies that have lost the integrated pZDJ plasmid. The counter-selection relies on
the *sacB* gene carried by the suicide plasmid (Jaschke et al. 2011). In order to induce a third crossing
over between the FRT sites of the kanamycin cassette, cells were grown in liquid LB without any
antibiotic selection until late stationary phase and then streaked on LB solid medium. Resultant
colonies were then screened through PCR assays.

To make knock-out strains for *R. palustris*, the same general protocol used for *R. sphaeroides* was
followed (Welander et al. 2012). The suicide plasmid used for R. palustris was pJQ200SK (Quandt
and Hynes 1993) and the host strain was *E. coli* BW29427-λpir-RP4, which requires 300 µM
diaminopimelic acid for growth. This auxotrophy allows to easily remove the plasmid-donor bacterium
from the medium after conjugation.

Construction of *R. sphaeroides* and *R. palustris* overexpression strains

To create Mic60- and Orf52-overexpressing strains, the inducible expression plasmids for *R.*
sphaeroides pIND4 -Km and pIND4-Gm were kindly provided by Judith P. Armitage (University of
Oxford), and Alexander Westbye (University of British Columbia), respectively. The inducible
expression plasmid pSRK was used in *R. palustris*. The coding sequences of the *mic60* and *orf52*
genes of *R. sphaeroides* and *R. palustris* were amplified by PCR and assembled into the expression
vectors pIND4 and pSRK, respectively, with the NEBuilder HiFi DNA Assembly Master Mix (New
England Biolabs). The *mic60* and *orf52* homologs were cloned into kanamycin resistance-conferring
plasmids and gentamycin resistance-conferring plasmids, respectively. To insert the C-terminal 6xHis
tag as part of the coding sequences of *mic60* and *orf52* the BamHI and BglII restriction sites in pIND4
were used, or the tag was incorporated as part of the primer used to amplify the targets in *R.*
palustris. The assembled plasmids were transformed into suitable conjugative *E. coli* hosts, and
conjugation assays with recipient *R. sphaeroides* and *R. palustris* strains were done as described
above. After conjugation, the exconjugants were repeatedly streaked onto antibiotic-containing plates
to remove the plasmid-donor bacterium. The resulting overexpression strains were further verified by
RT qPCR.

RT qPCR

R. sphaeroides and *R. palustris* strains were grown in triplicate until mid-log phase as described for
growth analysis, after which 1 mM IPTG was added, and the cells allowed to grow for 7 h 20 min for
R. sphaeroides and 4 h 30 min for *R. palustris*; cultures were grown in parallel in the absence of
IPTG. Then 1 ml of cell suspension was centrifuged and the pellet resuspended in 1 ml of PGTX
nucleic acid extraction buffer (Pinto et al. 2009). The samples were then flash frozen and stored at -20
°C until the PGTX-mediated nucleic acid extraction procedure, essentially following the classical
phenol/chloroform extraction method (Pinto et al., 2009). Briefly, isolated total nucleic acid
concentration was measured using a NanoDrop DeNovix DS-11 (Thermo Fisher) and diluted to 100
ng/μL in 50 μl DNase/RNase-free H_2O. The digestion of gDNA was performed by addition of DNase I
(Qiagen) and the concentration of total RNA was measured using the Qubit 2.0 Fluorometer (Thermo
Fisher). A total of 100 ng RNA was subsequently reverse transcribed using the Transcriptor First
Strand cDNASynthesis Kit (Roche) at 55 °C for cDNA synthesis. Quantitative PCR (qPCR) was
performed in triplicates in a CFX96 qPCR cycler (Bio-Rad) in 20 μl reactions containing 1x PowerUp
Sybr Green master mix (Applied Biosystems, USA), 8 pmol of each primer, and 10 μg cDNA with the
following program: initial denaturation 10 min at 95 °C; 45 cycles of 15 s at 95 °C and 30 s at 60 °C.
Primers annealing to *mic60*, *orf52*, and the *rpoZ* cDNAs used for qPCR are listed in Table S3. Cycle
threshold values were automatically computed with the CFX Maestro software (Bio-Rad). Additionally,
non-reverse transcribed RNA sample was used as a control to verify complete degradation of
genomic DNA. Relative abundance of transcripts in cells grown in the presence of IPTG compared to
those grown without the inducing agent was calculated using the $2^{-\Delta\Delta CT}$ method (Pfaffl 2001); the
unaffected *rpoZ* housekeeping gene cDNA was used to normalize the *mic60* and *orf52* values.

Transmission electron microscopy

Both classical TEM and electron tomography on 80 nm-thick sections were done as previously
described (Cadena et al., 2021; Kaurov et al., 2018). Scoring of *R. palustris* ICM area/TEM area and
R. sphaeroides tubules/cell per section were performed blinded on images obtained using JEOL 1010
TEM operating at an accelerating voltage of 80 kV and equipped with a MegaView III CCD camera
(SIS). Measurement of *R. palustris* ICM and total cell areas was done using Image J software
(Schneider et al. 2012) by tracing along the outermost electron dense membranes of the ICM network
and the outer membrane, respectively. The occurrence of tubule-like ICMs in *R. sphaeroides* was
counted on 20-21 images that were taken at 40,000x magnification (2.56 nm/pixel) at random parts of
the grid as before (Kaurov et al., 2018). These images were mixed and randomized prior to blind
scoring.

Electron tomograms were collected at a range of ±65° with tilt 1° steps using the JEOL 2100F TEM
working at 200 kV, equipped with Gatan camera K2 Summit and controlled with SerialEM automated
acquisition software (Mastronarde 2005). IMOD software (Mastronarde and Held 2017) was used for
tomogram reconstruction and generating 3D models by segmentation.

SDS PAGE and Western blotting

Bacterial lysates were separated on a Bolt 4-12% Bis-Tris Plus gel (Invitrogen), blotted onto a PVDF
membrane (Amersham), blocked in 5% low-fat, powdered milk (w/v) in phosphate buffered saline with
0.1% Tween 20 (v/v) (PBS-T), and probed with 6x-His Tag Monoclonal Antibody (HIS.H8) (Thermo
Fisher Scientific #MA1-21315) diluted in 5% milk in PBS-T (1:500). This was followed by incubation
with secondary HRP-conjugated anti-mouse antibody (1:2000; Bio-Rad). Proteins were visualized
using the Pierce ECL system (Genetica/Bio-Rad) on a ChemiDoc imager (Bio-Rad).

Isolation of ICMs via high-pressure homogenization and light spectroscopy

R. sphaeroides strains were cultured identically in C-succinate media (Cohen-Bazire et al. 1957) in
flat, glass bottles at 28-30°C under anaerobic conditions. Illumination was provided by one
incandescent 40W bulb providing ~10 μmol photons m^{-2} s^{-1}. The cells were fully adapted to the
incident light-intensity and harvested by centrifugation once they had reached mid-log phase. Cell
pellets were washed in 20 mM MES, 100 mM KCl, pH 6.8, then flash frozen and stored at –80 °C until
required

The cell pellets were re-suspended in 20 mM Tris.Cl, pH 8.0 and homogenised thoroughly with a few
grains of DNAse I (Qiagen) and a few mg of MgCl$_2$. The cells were broken by passage three times
through an Emulsiflex-CS cell disrupter (Avestin). The ruptured cell solution was first subjected to a
low-speed centrifugation step (10 min, 10,000 × g, 4°C) to remove any unbroken cells. The decanted
supernatant was ultra-centrifuged (120 min, 180,000 × g, 4°C), after which the supernatant was
discarded. The resulting chromatophore pellet was gently re-suspended in 20 mM Tris-HCl pH 8.0
before the optical density was adjusted to 10 cm^{-1} at the Qx absorption maximum (~590 nm) using a
Shimadzu UV-Vis-NIR UV2600 spectrophotometer equipped with an integrating sphere.

<u>Blue Native PAGE</u>

BN-PAGE of isolated ICMs was adapted from published protocols (Cadena et al., 2021). Briefly, 1 mg
of total protein was resuspended in 100 µl NativePAGE sample buffer (Invitrogen), lysed with 1.5%
digitonin (v/v) and 0.1% Triton X-100 (v/v) for 1 h on ice then cleared by centrifugation (22,000 × g, 20
min, 4 °C). Subsequently, 5% Coomassie brilliant blue G-250 was added before loading ~100 µg on a
3-12% Bis-Tris BNE gel (Invitrogen). After electrophoresis (2.5 hours, 150 V, 4 °C), the gel was
blotted onto a PVDF membrane (Amersham) and probed as described above.

<u>Affinity purification</u>

Affinity purification (AP) of tagged proteins from 1 mg isolated ICMs were solubilized in IPP50 buffer
(50 mM KCl, 20 mM Tris-HCl pH 7.7, 3 mM MgCl$_2$, 10% glycerol, 1 mM phenylmethanesulfonyl
fluoride, complete EDTA free protease inhibitor cocktail (Roche) supplemented with 1.5% digitonin
(v/v) and 0.1% Triton X-100 (v/v) for 1 h on ice. After centrifugation (22,000 ×g, 20 min, 4 °C) the
supernatant was added to 2.0 mg of cobalt conjugated Dynabeads (Thermo Fisher) to capture the
His-tag. The Dynabeads were pre-washed in 300 µl of IPP50 + 1.5% digitonin for 5 min at RT. The
solubilized ICMs were rotated with beads for 90 min at 4 °C. After removal of the flow-through, the
beads were washed three times in IPP50 + 1.5% digitonin. Prior to eluting, the beads were transferred
into a new tube. Elution was done with 300 mM imidazole in IPP50 for 15 min at RT and shaking at
1000 rpm. The elutes were further processed for LC-MS^2 analysis or resolved by SDS-PAGE. APs
were performed in triplicate.

<u>Protein preparation and mass spectroscopy</u>

Individual bands containing proteins of interest were excised from Coomasie stained SDS PAGE gel
using a razor blade and cut into small pieces (~1 mm^3). Bands were destained by sonication for 30
min in 50% acetonitrile (ACN) and 50 mM ammonium bicarbonate (ABC). After destaining, gels were
dried in ACN. Disulfide bonds were reduced using 10mm DTT in 100mM ABC at 60°C for 30 min.
After that, samples were again dried with ACN and free cysteine residues were blocked using 55 mM
iodoacetamide in 100 mM ABC for 10 min at room temperature in the dark. Samples were dried
thoroughly and then digestion buffer (10% ACN, 40 mM ABC and 13 ng/µl trypsin) was added to
cover gel pieces. Proteins were digested at 37 °C overnight. After digestion, 150 µl of 50% ACN with
0,5% formic acid was added and sonicated for 30 min. Supernatant containing peptides was
transferred to a new microcentrifuge tube and another 150 µl of elution solution was added and
sonicated for 30 min. This solution was removed, combined with the previous solution and dried by
SpeedVac. Dried peptides were reconstitued in 2% ACN with 0,1% TFA and injected into Ultimate
3000 Nano LC coupled to Orbitrap Fusion.

Eluates of co-AP proteins and thin-sliced BN-PAGE gels were processed for MS analysis as
described elsewhere (Cadena et al., 2021). In brief, eluate samples were resuspended in 100 mM
tetraethylammonium bromide containing 2% sodium deoxycholate. Cysteines were reduced with 10
mM tris(2-carboxyethyl)phosphine and subsequently cleaved with 1 µg trypsin overnight at 37 °C.
After digestion, 1% trifluoroacetic acid (TFA) was added to wash twice and eluates were resuspended
in 20 µl TFA per µg of protein. A nano reversed-phased column (EASY-Spray column, 50 cm x 75 µm
inner diameter, PepMap C18, 2 µm particles, 100 Å pore size) was used for LC/MS analysis. Mobile
phase buffer A consisted of water and 0.1% formic acid. Mobile phase D consisted of acetonitrile and
0.1% formic acid. Samples were loaded onto the trap column (Acclaim PepMap300, C18, 5 µm, 300 Å
pore size, 300 µm x 5 mm) at a flow rate of 15 µl/min. The loading buffer consisted of water, 2%

acetonitrile, and 0.1% TFA. Peptides were eluted using a Mobile phase B gradient from 2% to 40%
over 60 min at a flow rate of 300 ml/min. The peptide cations eluted were converted to gas-phase ions
via electrospray ionization and analyzed on a Thermo Orbitrap Fusion (Q-OT- qIT, Thermo Fisher).
Full MS spectra were acquired in the Orbitrap with a mass range of 350-1, 400 *m/z*, at a resolution of
120,000 at 200 *m/z* and with a maximum injection time of 50 ms. Tandem MS was performed by
isolation at 1,5 Th with the quadrupole, high-energy collisional dissociation (HCS) fragmentation with
normalized collision energy of 30, and rapid scan MS analysis in the ion trap. The MS/MS ion count
target was set to 10^4 and the max infection time at 35 ms. Only those precursors with a charge state
of 2-6 were sampled. The dynamic exclusion duration was set to 45 s with a 10 ppm tolerance around
the selected precursor and its isotopes. Monoisotopic precursor selection was on with a top speed
mode of 2 s cycles.

<u>Analysis of mass spectroscopy peptides</u>

Label-free quantification of the data were analyzed using the MaxQuant software (version 1.6.2.1)
(Cox et al. 2014). The false discovery rates for peptides and for proteins was set to 1% with a
specified minimum peptide length of seven amino acids. The Andromeda search engine was used for
the MS/MS spectra against the *R. sphaeroides* 2.4.1 predicted proteome (ASM1290v2) downloaded
from NCBI GenBank on October 2020. Enzyme specificity was set to C-terminal Arg and Lys,
alongside for cleavage at proline bonds with a maximum of two missed cleavages. Dithiomethylation
of cysteine was selected as a fixed modification with N-terminal protein acetylation and methionine
oxidation as variable modifications. The 'match between runs' feature in MaxQuant was used to
transfer identification to other LC-MS/MS runs based on mass and retention time with a maximum
deviation of 0.7 min. Quantifications were performed using a label-free algorithm as previously
described (Cox et al., 2014). Data analysis was performed using Perseus software (version 1.6.1.3).
Eluate co-AP proteins identified with a mean Log2 ratio (protein$_{His}$/WT) >3-fold change and having a
Andromeda confidence score >100 (Cox et al., 2011) in three independent biological replicates were
analyzed, while gel-slice proteins identified with a mean Log2 transformed LFQ score >23 and
present in four biological replicates were analyzed.

<u>Data Availability</u>

The LC-MS/MS data have been deposited to the ProteomeXchange Consortium
(http://www.proteomexchange.org) via the PRIDE partner repository with the data set identifier
PXD032747.

ACKNOWLEDGEMENTS

We would like to dedicate this work to Thomas Cavalier-Smith whose large-scale synthetic work on
evolutionary cell biology stimulated thinking on this topic by SAM-G many years ago. We thank Karel
Harant and Pavel Talacko (Charles University, Prague) for performing LC-MS analysis. SAM-G is
supported by an EMBO Postdoctoral Fellowship (ALTF 21-2020). MML was supported by Nova
Scotia Health Research Foundation (NSHRF) Scotia Scholarship 2012-8781. This work was also
supported by Czech Science Foundation grants 20-23513S to HH, 22-01-26S to JL and 19-28778X to
MK, as well as the Czech Ministry of Education grant OPVVV16_019/0000759 and Czech BioImaging
grant LM2015062 and a Discovery grant RGPIN-2022-05430 to AJR from the Natural Sciences and
Engineering Research Council of Canada.

REFERENCES

Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL,
Mori H. 2006. Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants:
the Keio collection. *Mol Syst Biol* 2:2006.0008.

Barrera NP, Robinson CV. 2011. Advances in the mass spectrometry of membrane proteins: from
individual proteins to intact complexes. *Annu Rev Biochem* 80:247–271.

Beatty JT, Gest H. 1981. Biosynthetic and bioenergetic functions of citric acid cycle reactions in
Rhodopseudomonas capsulata. *J Bacteriol* 148:584–593.

Bock-Bierbaum T, Funck K, Wollweber F, Lisicki E, Laborenz J, Noel JK, Hessenberger M, von der
Malsburg A, von der Malsburg K, Bernert C, et al. 2022. Structural insights into crista junction
formation by the Mic60-Mic19 complex. *bioRxiv*:2022.03.30.486340.

Brimacombe CA, Stevens A, Jun D, Mercer R, Lang AS, Beatty JT. 2013. Quorum-sensing regulation
of a capsular polysaccharide receptor for the *Rhodobacter capsulatus* gene transfer agent
(RcGTA). *Mol Microbiol* 87:802–817.

Cadena LR, Gahura O, Panicucci B, Zíková A, Hashimi H. 2021. Mitochondrial Contact Site and
Cristae Organization System and F_1F_O-ATP Synthase Crosstalk Is a Fundamental Property of
Mitochondrial Cristae. *mSphere*:e0032721.

Callister SJ, Nicora CD, Zeng X, Roh JH, Dominguez MA, Tavano CL, Monroe ME, Kaplan S,
Donohue TJ, Smith RD, et al. 2006. Comparison of aerobic and photosynthetic *Rhodobacter*
sphaeroides 2.4.1 proteomes. *Journal of Microbiological Methods* 67:424–436.

Cohen-Bazire G, Sistrom WR, Stanier RY. 1957. Kinetic studies of pigment synthesis by non-sulfur
purple bacteria. *J Cell Comp Physiol* 49:25–68.

Cox J, Hein MY, Luber CA, Paron I, Nagaraj N, Mann M. 2014. Accurate proteome-wide label-free
quantification by delayed normalization and maximal peptide ratio extraction, termed
MaxLFQ. *Mol Cell Proteomics* 13:2513–2526.

Cox J, Neuhauser N, Michalski A, Scheltema RA, Olsen JV, Mann M. 2011. Andromeda: a peptide
search engine integrated into the MaxQuant environment. *J Proteome Res* 10:1794–1805.

Daems WT, Wisse E. 1966. Shape and attachment of the cristae mitochondriales in mouse hepatic
cell mitochondria. *J Ultrastruct Res* 16:123–140.

D'Amici GM, Rinalducci S, Murgiano L, Italiano F, Zolla L. 2010. Oligomeric Characterization of the
Photosynthetic Apparatus of *Rhodobacter sphaeroides* R26.1 by Nondenaturing
Electrophoresis Methods. *J. Proteome Res.* 9:192–203.

Dietz JV, Willoughby MM, Piel RB, Ross TA, Bohovych I, Addis HG, Fox JL, Lanzilotta WN, Dailey
HA, Wohlschlegel JA, et al. 2021. Mitochondrial contact site and cristae organizing system
(MICOS) machinery supports heme biosynthesis by enabling optimal performance of
ferrochelatase. *Redox Biol* 46:102125.

Doyle MT, Bernstein HD. 2019. Bacterial outer membrane proteins assemble via asymmetric
interactions with the BamA β-barrel. *Nat Commun* 10:3358.

Drews G. 1991. Intracytoplasmic membranes in bacterial cells: organization, function, and
biosynthesis. In: Prokaryotic Structure and Function: A New Perspective. Forty-seventh
Symposium of the Society for General Microbiology Held at the University of Edinburgh.
Cambridge University Press.

Drews G, Golecki JR. 2004. Structure, Molecular Organization, and Biosynthesis of Membranes of
Purple Bacteria. In: Blankenship RE, Madigan MT, Bauer CE, editors. Anoxygenic
Photosynthetic Bacteria. Vol. 2. Advances in Photosynthesis and Respiration. Dordrecht:
Kluwer Academic Publishers. p. 231–257. Available from: http://link.springer.com/10.1007/0-
306-47954-0_12

Drozdetskiy A, Cole C, Procter J, Barton GJ. 2015. JPred4: a protein secondary structure prediction
server. *Nucleic Acids Res* 43:W389–W394.

Dziuba M, Riese CN, Borgert L, Wittchen M, Busche T, Kalinowski J, Uebe R, Schüler D. 2021. The
Complex Transcriptional Landscape of Magnetosome Gene Clusters in *Magnetospirillum*
gryphiswaldense.Ajo-Franklin CM, editor. *mSystems* 6:e00893-21.

Ehrenreich A, Widdel F. 1994. Anaerobic oxidation of ferrous iron by purple bacteria, a new type of
phototrophic metabolism. *Appl. Environ. Microbiol.* 60:4517–4526.

Fan L, Wu D, Goremykin V, Xiao J, Xu Y, Garg S, Zhang C, Martin WF, Zhu R. 2020. Phylogenetic
analyses with systematic taxon sampling show that mitochondria branch within
Alphaproteobacteria. *Nat Ecol Evol* 4:1213–1219.

Fejes AP, Yi EC, Goodlett DR, Beatty JT. 2003. Shotgun proteomic analysis of a chromatophore-
enriched preparation from the purple phototrophic bacterium *Rhodopseudomonas palustris*.
Photosynthesis Research 78:195–203.

Gautier R, Douguet D, Antonny B, Drin G. 2008. HELIQUEST: a web server to screen sequences with
specific alpha-helical properties. *Bioinformatics* 24:2101–2102.

Greening C, Lithgow T. 2020. Formation and function of bacterial organelles. *Nat Rev Microbiol*
18:677–689.

Guan Y, Zhu Q, Huang D, Zhao S, Jan Lo L, Peng J. 2015. An equation to estimate the difference
between theoretically predicted and SDS PAGE-displayed molecular weights for an acidic
peptide. *Sci Rep* 5:13370.

Hammond M, Dorrell RG, Speijer D, Lukeš J. 2022. Eukaryotic cellular intricacies shape mitochondrial
proteomic complexity. *BioEssays* 44:2100258.

Hessenberger M, Zerbes RM, Rampelt H, Kunz S, Xavier AH, Purfürst B, Lilie H, Pfanner N, van der
Laan M, Daumke O. 2017. Regulated membrane remodeling by Mic60 controls formation of
mitochondrial crista junctions. *Nat Commun* 8:15258.

Höhr AIC, Lindau C, Wirth C, Qiu J, Stroud DA, Kutik S, Guiard B, Hunte C, Becker T, Pfanner N, et
al. 2018. Membrane protein insertion through a mitochondrial β-barrel gate. *Science*
359:eaah6834.

Huynen MA, Mühlmeister M, Gotthardt K, Guerrero-Castillo S, Brandt U. 2016. Evolution and
structural organization of the mitochondrial contact site (MICOS) complex and the
mitochondrial intermembrane space bridging (MIB) complex. *Biochim Biophys Acta* 1863:91–
101.

Iba K, Takamiya K, Toh Y, Nishimura M. 1988. Roles of bacteriochlorophyll and carotenoid synthesis
in formation of intracytoplasmic membrane systems and pigment-protein complexes in an
aerobic photosynthetic bacterium, Erythrobacter sp. strain OCh114. *J Bacteriol* 170:1843–
1847.

Imhoff JF, Rahn T, Künzel S, Neulinger SC. 2018. Photosynthesis Is Widely Distributed among
Proteobacteria as Demonstrated by the Phylogeny of PufLM Reaction Center Proteins. *Front.*
Microbiol. 8:2679.

Jackson PJ, Lewis HJ, Tucker JD, Hunter CN, Dickman MJ. 2012. Quantitative proteomic analysis of
intracytoplasmic membrane development in *Rhodobacter sphaeroides*. *Mol Microbiol*
84:1062–1078.

Jaschke PR, Saer RG, Noll S, Beatty JT. 2011. Modification of the Genome of *Rhodobacter*
sphaeroides and Construction of Synthetic Operons. In: Methods in Enzymology. Vol. 497.
Elsevier. p. 519–538. Available from:
https://linkinghub.elsevier.com/retrieve/pii/B9780123850751000238

Jiao Y, Kappler A, Croal LR, Newman DK. 2005. Isolation and characterization of a genetically
tractable photoautotrophic Fe(II)-oxidizing bacterium, *Rhodopseudomonas palustris* strain
TIE-1. *Appl Environ Microbiol* 71:4487–4496.

Jiao Y, Newman DK. 2007. The pio operon is essential for phototrophic Fe(II) oxidation in
Rhodopseudomonas palustris TIE-1. *J Bacteriol* 189:1765–1773.

John GB, Shang Y, Li L, Renken C, Mannella CA, Selker JML, Rangell L, Bennett MJ, Zha J. 2005.
The mitochondrial inner membrane protein mitofilin controls cristae morphology. *Mol Biol Cell*
16:1543–1554.

Jun D, Saer RG, Madden JD, Beatty JT. 2014. Use of new strains of *Rhodobacter sphaeroides* and a
modified simple culture medium to increase yield and facilitate purification of the reaction
centre. *Photosynth Res* 120:197–205.

Kaurov I, Vancová M, Schimanski B, Cadena LR, Heller J, Bílý T, Potěšil D, Eichenberger C, Bruce H,
Oeljeklaus S, et al. 2018. The Diverged Trypanosome MICOS Complex as a Hub for
Mitochondrial Cristae Shaping and Protein Import. *Curr Biol* 28:3393-3407.e5.

Kobayashi K, Masuda T, Tajima N, Wada H, Sato N. 2014. Molecular Phylogeny and Intricate
Evolutionary History of the Three Isofunctional Enzymes Involved in the Oxidation of
Protoporphyrinogen IX. *Genome Biology and Evolution* 6:2141–2155.

Kozjak-Pavlovic V. 2017. The MICOS complex of human mitochondria. *Cell Tissue Res* 367:83–93.

Kühlbrandt W. 2019. Structure and Mechanisms of F-Type ATP Synthases. *Annu Rev Biochem*
88:515–549.

LaSarre B, Kysela DT, Stein BD, Ducret A, Brun YV, McKinlay JB. 2018. Restricted Localization of
Photosynthetic Intracytoplasmic Membranes (ICMs) in Multiple Genera of Purple Nonsulfur
Bacteria.Harwood CS, editor. *mBio* 9:e00780-18.

Lee MD. 2019. GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics* 35:4162–4164.

Lohße A, Borg S, Raschdorf O, Kolinko I, Tompa E, Pósfai M, Faivre D, Baumgartner J, Schüler D.
2014. Genetic dissection of the mamAB and mms6 operons reveals a gene set essential for
magnetosome biogenesis in *Magnetospirillum gryphiswaldense*. *J Bacteriol* 196:2658–2669.

Lohße A, Ullrich S, Katzmann E, Borg S, Wanner G, Richter M, Voigt B, Schweder T, Schüler D.
2011. Functional Analysis of the Magnetosome Island in *Magnetospirillum gryphiswaldense*:
The mamAB Operon Is Sufficient for Magnetite Biomineralization.Battista JR, editor. *PLoS*
ONE 6:e25561.

Mai U, Mirarab S. 2018. TreeShrink: fast and accurate detection of outlier long branches in collections
of phylogenetic trees. *BMC Genomics* 19:272.

Mannella CA, Marko M, Penczek P, Barnard D, Frank J. 1994. The internal compartmentation of rat-
liver mitochondria: tomographic study using the high-voltage transmission electron
microscope. *Microsc Res Tech* 27:278–283.

Martijn J, Vosseberg J, Guy L, Offre P, Ettema TJG. 2018. Deep mitochondrial origin outside the
sampled alphaproteobacteria. *Nature* 557:101–105.

Mastronarde DN. 2005. Automated electron microscope tomography using robust prediction of
specimen movements. *J Struct Biol* 152:36–51.

Mastronarde DN, Held SR. 2017. Automated tilt series alignment and tomographic reconstruction in
IMOD. *J Struct Biol* 197:102–113.

Menardo F, Loiseau C, Brites D, Coscolla M, Gygli SM, Rutaihwa LK, Trauner A, Beisel C, Borrell S,
Gagneux S. 2018. Treemmer: a tool to reduce large phylogenetic datasets with minimal loss
of diversity. *BMC Bioinformatics* 19:164.

Michaud M, Gros V, Tardif M, Brugière S, Ferro M, Prinz WA, Toulmay A, Mathur J, Wozny M,
Falconet D, et al. 2016. AtMic60 Is Involved in Plant Mitochondria Lipid Trafficking and Is Part
of a Large Complex. *Curr Biol* 26:627–639.

Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, Tosatto SCE,
Paladin L, Raj S, Richardson LJ, et al. 2021. Pfam: The protein families database in 2021.
Nucleic Acids Research 49:D412–D419.

Muñoz-Gómez SA, Slamovits CH, Dacks JB, Baier KA, Spencer KD, Wideman JG. 2015. Ancient
homology of the mitochondrial contact site and cristae organizing system points to an
endosymbiotic origin of mitochondrial cristae. *Curr Biol* 25:1489–1495.

Muñoz-Gómez SA, Slamovits CH, Dacks JB, Wideman JG. 2015b. The evolution of MICOS:
Ancestral and derived functions and interactions. *Communicative & Integrative Biology*
8:e1094593.

Muñoz-Gómez SA, Susko E, Williamson K, Eme L, Slamovits CH, Moreira D, López-García P, Roger
AJ. 2022. Site-and-branch-heterogeneous analyses of an expanded dataset favour
mitochondria as sister to known Alphaproteobacteria. *Nat Ecol Evol* 6:253–262.

Muñoz-Gómez SA, Wideman JG, Roger AJ, Slamovits CH. 2017. The Origin of Mitochondrial Cristae
from Alphaproteobacteria. *Mol Biol Evol* 34:943–956.

Myers KS, Vera JM, Lemmer KC, Linz AM, Landick R, Noguera DR, Donohue TJ. 2020. Genome-
Wide Identification of Transcription Start Sites in Two Alphaproteobacteria, *Rhodobacter*
sphaeroides 2.4.1 and *Novosphingobium aromaticivorans* DSM 12444.Cameron Thrash J,
editor. *Microbiol Resour Announc* 9:e00880-20.

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic
algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 32:268–274.

Niederman RA. 2006. Structure, Function and Formation of Bacterial Intracytoplasmic Membranes. In:
Shively JM, editor. Complex Intracellular Structures in Prokaryotes. Vol. 2. Microbiology
Monographs. Berlin, Heidelberg: Springer Berlin Heidelberg. p. 193–227. Available from:
http://link.springer.com/10.1007/7171_025

Noble JM, Lubieniecki J, Savitzky BH, Plitzko J, Engelhardt H, Baumeister W, Kourkoutis LF. 2018.
Connectivity of centermost chromatophores in *Rhodobacter sphaeroides* bacteria. *Mol*
Microbiol 109:812–825.

Pánek T, Eliáš M, Vancová M, Lukeš J, Hashimi H. 2020. Returning to the Fold for Lessons in
Mitochondrial Crista Diversity and Evolution. *Curr Biol* 30:R575–R588.

Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A, Hugenholtz P. 2022. GTDB: an
ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank
normalized and complete genome-based taxonomy. *Nucleic Acids Research* 50:D785–D794.

Perkins G, Renken C, Martone ME, Young SJ, Ellisman M, Frey T. 1997. Electron tomography of
neuronal mitochondria: three-dimensional structure and organization of cristae and
membrane contacts. *J Struct Biol* 119:260–272.

Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic*
Acids Res 29:e45.

Pinevich AV. 1997. Intracytoplasmic membrane structures in bacteria. *Endocytobiosis and Cell*
Research 12:9–40.

Pinto FL, Thapper A, Sontheim W, Lindblad P. 2009. Analysis of current and alternative phenol based
RNA extraction methodologies for cyanobacteria. *BMC Mol Biol* 10:79.

Quandt J, Hynes MF. 1993. Versatile suicide vectors which allow direct selection for gene
replacement in gram-negative bacteria. *Gene* 127:15–21.

Rabl R, Soubannier V, Scholz R, Vogel F, Mendl N, Vasiljev-Neumeyer A, Körner C, Jagasia R, Keil
T, Baumeister W, et al. 2009. Formation of cristae and crista junctions in mitochondria
depends on antagonism between Fcj1 and Su e/g. *J Cell Biol* 185:1047–1063.

Rampelt H, Zerbes RM, van der Laan M, Pfanner N. 2017. Role of the mitochondrial contact site and
cristae organizing system in membrane architecture and dynamics. *Biochim Biophys Acta Mol*
Cell Res 1864:737–746.

Roger AJ, Muñoz-Gómez SA, Kamikawa R. 2017. The Origin and Diversification of Mitochondria. *Curr*
Biol 27:R1177–R1192.

Schavemaker PE, Muñoz-Gómez SA. 2022. The role of mitochondrial energetics in the origin and
diversification of eukaryotes. *bioRxiv*:2021.10.23.465364.

Scheuring S, Nevo R, Liu L-N, Mangenot S, Charuvi D, Boudier T, Prima V, Hubert P, Sturgis JN,
Reich Z. 2014. The architecture of *Rhodobacter sphaeroides* chromatophores. *Biochimica et*
Biophysica Acta (BBA) - Bioenergetics 1837:1263–1270.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis.
Nat Methods 9:671–675.

Selao TT, Branca R, Chae PS, Lehtiö J, Gellman SH, Rasmussen SGF, Nordlund S, Norén A. 2011.
Identification of chromatophore membrane protein complexes formed under different nitrogen
availability conditions in *Rhodospirillum rubrum*. *J Proteome Res* 10:2703–2714.

Sener M, Strumpfer J, Singharoy A, Hunter CN, Schulten K. 2016. Overall energy conversion
efficiency of a photosynthetic vesicle. *eLife* 5:e09541.

Tang J, Zhang K, Dong J, Yan C, Hu C, Ji H, Chen L, Chen S, Zhao H, Song Z. 2020. Sam50-Mic19-
Mic60 axis determines mitochondrial cristae architecture by mediating mitochondrial outer
and inner membrane contact. *Cell Death Differ* 27:146–160.

Tarasenko D, Barbot M, Jans DC, Kroppen B, Sadowski B, Heim G, Möbius W, Jakobs S, Meinecke
M. 2017. The MICOS component Mic60 displays a conserved membrane-bending activity that
is necessary for normal cristae morphology. *Journal of Cell Biology* 216:889–899.

Voulhoux R, Bos MP, Geurtsen J, Mols M, Tommassen J. 2003. Role of a highly conserved bacterial
protein in outer membrane protein assembly. *Science* 299:262–265.

Warnsmann V, Marschall L-M, Meeßen AC, Wolters M, Schürmanns L, Basoglu M, Eimer S,
Osiewacz HD. 2022. Disruption of the MICOS complex leads to an aberrant cristae structure
and an unexpected, pronounced lifespan extension in *Podospora anserina*.
bioRxiv:2022.03.21.485166.

Welander PV, Doughty DM, Wu C-H, Mehay S, Summons RE, Newman DK. 2012. Identification and
characterization of *Rhodopseudomonas palustris* TIE-1 hopanoid biosynthesis mutants.
Geobiology 10:163–177.

Wideman JG, Muñoz-Gómez SA. 2016. The evolution of ERMIONE in mitochondrial biogenesis and
lipid homeostasis: An evolutionary view from comparative cell biology. *Biochim Biophys Acta*
1861:900–912.

Wollweber F, von der Malsburg K, van der Laan M. 2017. Mitochondrial contact site and cristae
organizing system: A central player in membrane shaping and crosstalk. *Biochimica et*
Biophysica Acta (BBA) - Molecular Cell Research 1864:1481–1489.

Zick M, Rabl R, Reichert AS. 2009. Cristae formation-linking ultrastructure and function of
mitochondria. *Biochim Biophys Acta* 1793:5–19.

Supplemental Material

Figure S1. Verification of *mic60* and *orf52* knock-out strains and IPTG-induced overexpression of
Mic60 and Orf52. **(A & B)** PCR assays confirm the disruption of the *mic60* and *orf52* genes in *R.*
palustris **(A)** and *R. sphaeroides* **(B)**. Genomic contexts of the relevant loci for the WT, Δ*mic60* and
Δ*orf52* strains shown on top, with primer pairs and their expected amplicon sizes shown below each
schematic gene arrangement. Lower panel shows each PCR amplicon from each strain (labelled
above gel) after agarose gel electrophoresis. Size markers shown either to the left or right of the gel.
(C) Real time PCR showing relative abundancies of Mic60 and Orf52 mRNAs in *R. palustris* (left) and
R. sphaeroides (right) strains grown in the presence of the expression induction agent IPTG relative
to the same strains grown without IPTG. Error bars show standard deviation from 3 replicates of
assayed induced and non-induced cells.

Figure S2. Proteomic analysis of isolated ICMs plus Mic60 and Orf52 interactomes. **(A)** Blue native
gel resolved detergent-solubilized ICMs from *R. sphaeroides* WT in which the ~250 kDa band (boxed)
was excised in quadruplet for subsequent MS analysis. To right of gel is a histogram of the
Coomassie-stained band intensities (*I*) *along* the vertical axis of the run. The scissors denote the ~250
kDa band intensity signal. **(B & C)** A list of all proteins, including excluded contaminants, found within
the enriched protein area of the volcano plots in in Figure 4A-B for Mic60 **(B)** and Orf52 **(C)**. Columns
as described in legend of Figure 5C-D. Note the presence of likely contaminant found in both APs,
PAS-fold containing histidine kinase (PAS), which is a large protein amenable to LC-MS/MS in
contrast to RSP_7517 and not found in any of our requisite proteomes. **(D)** Kyle and Doolittle
hydropathy plot of RSP_7517, whose amino acid sequence is given below. Predicted transmembrane
domain shaded and a potential oligomerization motif AxxxA underlined. Lysine (K) residues
recognized by trypsin protease are in bold.

Dataset 1. List of proteins found within Blue Native gel slices as indicated in Figures 3C and S4A in
quadruplicates.

Dataset 2. List of all proteins found in the Orf52 and Mic60 interactomes in comparison to WT
negative controls by MS of AP eluates. Related to Figure 4A-B.

Table S1. List of strains used in this study.

Table S2. List of plasmids used in this study.

Table S3. List of primers and their sequences used in this study.

Movie S1. 3D reconstruction model of *R. sphaeroides* WT as rendered from electron tomograms
shown at the start of the video. Red, ICM membranes; yellow, cytoplasmic membrane; green, outer
membrane. Lower right corner, 200 nm scale bar.

Movie S2. 3D reconstruction model of *R. sphaeroides* Δ*mic60* as rendered from electron tomograms
shown at the start of the video. Red, ICM membranes; yellow, cytoplasmic membrane; green, outer
membrane. Lower right corner, 200 nm scale bar.

Movie S3. 3D reconstruction model of *R. sphaeroides* Mic60↑ as rendered from electron tomograms
shown at the start of the video. Red, ICM membranes; yellow, cytoplasmic membrane; green, outer
membrane. Lower right corner, 200 nm scale bar.

Movie S4. 3D reconstruction model of *R. sphaeroides* Δ*orf52* as rendered from electron tomograms
shown at the start of the video. Red, ICM membranes; yellow, cytoplasmic membrane; green, outer
membrane. Lower right corner, 200 nm scale bar.

Movie S5. 3D reconstruction model of *R. sphaeroides* Orf52↑ as rendered from electron tomograms
shown at the start of the video. Red, ICM membranes; yellow, cytoplasmic membrane; green, outer
membrane. Lower right corner, 200 nm scale bar.

1033

A High light

B Low light

C High light

D Low light

R. palustris

R. sphaeroides

E

WT Δ*mic60* Mic60↑ Δ*orf52* Orf52↑

R. palustris

F

G

WT Δ*mic60* Mic60↑ Δ*orf52* Orf52↑

R. sphaeroides

H

	WT	Δ*mic60*	Mic60↑	Δ*orf52*	Orf52↑
Sections (n):	20	21	21	21	20
Total cells:	267	152	80	87	51
Total tubules:	11	66	56	29	30