Modeling foot-and-mouth disease dissemination in Brazil and evaluating the effectiveness of control measures

Nicolas C. Cardenas[1], Francisco P. N. Lopes[2], Alencar Machado[3], Vinicius Maran[3], Celio Trois[3,4], Felipe Amadori Machado[3], Gustavo Machado[1*]

[1]Department of Population Health and Pathobiology, College of Veterinary Medicine, North Carolina State University, Raleigh, NC, USA

[2]Departamento de Defesa Agropecuária, Secretaria da Agricultura, Pecuária e Desenvolvimento Rural, Porto Alegre, Brazil

[3]Laboratory of Ubiquitous, Mobile and Applied Computing (LUMAC), Polytechnic College of Federal University of Santa Maria, Santa Maria, Brazil

[4]Department of Chemical Engineering, Rovira i Virgili University, Tarragona, Spain

* Correspondence:
Corresponding Author: gmachad@ncsu.edu

Keywords: dynamical models, infectious disease control, epidemiology, transmission, targeted control

Abstract

Foot-and-mouth disease (FMD) is known to infect multiple food-animal species and to spread among ungulate populations. We introduce a multiscale compartmental stochastic model that incorporates population dynamics, births, deaths, and species-specific transmission dynamics. The model integrates disease dynamics at both the between-farm and the within-farm levels. We developed four scenarios to evaluate the effectiveness of countermeasures, including movement standstill, vaccination, and depopulation. The base scenario involved vaccinating the animals on 20 farms and depopulating four infected farms. The three alternative control scenarios included increasing vaccination and depopulation capacities and no vaccination. Our findings indicate that bovines were the most frequently infected species, followed by swine and small ruminants. After ten days of initial spread, the number of infected farms ranged from 1 to 123, with most simulations (90.12%) predicting fewer than 50 infected farms. Most of the secondary spread occurred within a range of 25 km. Early response to initiating the control action reduces the number of days spent working on control actions while intensifying daily depopulation and vaccination capacity; this may be worth considering in decision-making processes for future control of FMD. Emergency vaccination proved to be efficient in reducing the magnitude and duration of outbreaks, whereas increasing depopulation without the use of vaccination also proved to be effective in eliminating the outbreaks.

1 Introduction

Foot-and-mouth disease (FMD) is an infectious disease of cloven-hoofed animals that affects multiple species, including bovines, swine, small ruminants, and wildlife (1). During the 2001 FMD epidemic in the U.K. and the Netherlands, more than 6.7 million animals were slaughtered, including healthy ones (preemptive culling) (2). In both outbreaks, multiple species were infected, including goats on mixed dairy goat/veal calf farms (2,3), and there were additional costs to other sectors, such as tourism, with a total expenditure of approximately 2.7 to 3.2 billion euros (4). The distribution of

FMD outbreaks in South America has changed over the years; the most recent outbreaks occurred in 2017 and 2018 in Colombia (5), and no epidemics have been officially reported in Latin America since then, despite Venezuela's absence of official international status for FMD (6). The official World Organization for Animal Health (WOAH) database for 2022 registered more than 57,000 outbreaks in 99 counties from 1996 to 2012; 68% of the FMD cases were associated with cattle, 22% with swine, and 21% with small ruminants and other species, whereas buffalo were not directly mentioned (7). Despite substantial evidence that all susceptible species can contribute to significant FMD epidemics, response plans frequently focus on controlling the spread of FMD among cattle populations. This approach often overlooks the role of other domestic species (8). For example, the large-scale 2001 UK outbreak was linked to swine slaughterhouses (9,10). Similarly, the most recent major outbreak in South America in 2001 affected 2,027 farms in Uruguay, where cattle and small ruminants were the predominant infected species (11).

Notably, FMD pathogenesis and transmission dynamics vary among species, given differences in the viral loads needed to cause infection and variability in the latency and duration of infection (12–15). For example, infected swine shed more viral particles than do cattle and sheep, historically resulting in widespread epidemics expected when swine are infected (14). Thus, it is pivotal to consider heterogeneity in transmission dynamics when modeling within- and/or between-farm FMD dissemination (8). Similarly, field observations and experimental trials have demonstrated that the spread of FMD between farms occurs primarily through direct contact between susceptible and infected animals (12,16) and via indirect contact with fomites and long-distance transport of aerosols, a process known as spatial transmission (16).

Mathematical models have been extensively utilized to investigate the propagation of FMD epidemics (17,18). Despite significant technical and computational advancements, simplifications of complex dynamics are often necessary due to computational costs or the lack of detailed population data, such as herd structure (e.g., number of individuals, number of animals born alive) and animal movements (17). A common simplification in modeling involves restricting the dynamics to a single species (4,19,20). Despite the varied applications and efforts in modeling FMD, outstanding questions persist regarding the measurement of epidemic trajectories and control strategies, particularly given the heterogeneous transmission dynamics among different susceptible species coexisting on the same premises (8,18).

Here, we develop a multihost, single-pathogen, multiscale model designed to capture the dynamics of FMD transmission patterns across different host species. The model i) simulates the spread of FMD within the state of Rio Grande do Sul in Brazil; ii) describes the geodesic distances between the initial outbreak and secondary cases; and iii) compares the effects of control action strategies, including emergency vaccination, depopulation, various restrictions on between-farm movement, and surveillance activities within control zones, taking into account the initial number of infected farms at the onset of control measures.

2   Materials and methods

This section presents the data used to create the dataset that was used to test and evaluate the model in outbreak simulations. The way in which the model was formulated is also described in this section, as are its outputs and the software used in its development.

2.1 Data sources
2.1.1 Population data

A comprehensive dataset was compiled from the official records of 355,676 farms registered in Rio Grande do Sul, Brazil (21), hosted in the Agricultural Defense System (SDA)(22). The dataset included the number of animals per location individually for cattle, buffalo, swine, sheep, and goats. Following stringent criteria, 70,853 premises were excluded due to missing geographical coordinates,

instances without animal stock, and the absence of incoming and/or outgoing movement during the
study period spanning August 24, 2022 to August 24, 2023. Consequently, the final dataset included
284,823 farms with accurate and reliable information. To simplify the analysis, population and
movement data from cattle and buffalo farms were merged into a single category denoted "bovines".
Similarly, sheep and goats were classified as "small ruminants". Supplementary Material Figure S1
presents the farm-level population distribution, and Supplementary Material Figure S2 presents the
geographical farm density distribution.

2.1.2 Birth and death
Producers are required to disclose to the SDA at least once a year the total number of animals born
alive on their farms and the number of deaths, including those due to natural causes. We collected
data on births and deaths from the SDA; these data included 273,787 individual records associated
with births and 268,790 records associated with deaths. The daily numbers of births and deaths,
categorized by species, are depicted in Supplementary Material Figure S3.

2.1.3 Movement data
From August 24, 2022 to August 24, 2023, 763,448 unique between-farm and farm-to-
slaughterhouse movements were recorded; these were collected from the SDA centralized traceability
database. Upon evaluation of the movement data, 106,481 records (13.9%) were removed for various
reasons: a) lack of identification of origin or destination; b) zero animals moved; c) lack of
specification of exact origin and destination premises; and d) movements to or from premises not
registered in the population data or to premises outside the state of Rio Grande do Sul. Ultimately,
413,939 unique between-farm movements and 243,028 farm-to-slaughterhouse movements were
analyzed. The daily farm-to-farm and farm-to-slaughterhouse movements, categorized by species, are
depicted in Supplementary Material Figure S3.

2.2 Outbreak simulation
Over two hundred thousand farms are located in Rio Grande do Sul; from these, a sampling was
made, and the sample farms were used as the initial infected premises. Our sample was multistage
and stratified, with the number of farms and species categorized by municipality (23). The sample
size was determined considering a prevalence of 50%, with a 95% significance level and a 1.1%
margin of error. The sampling yielded a total of 10,294 farms (Supplementary Material Figure S4).
Our model simulation was performed in two steps. First, FMD was seeded randomly into sample
farms via one infected animal between August 24, 2022 and August 24, 2022. For farms with
multiple species, for example, those with bovines, swine, and small ruminants, FMD was seeded into
bovines; for farms with swine and small ruminants, FMD was seeded into swine; and for farms with
cattle and small ruminants, FMD was seeded via one infected bovine. We assumed that all the
animals were susceptible to FMD, as the annual vaccination campaign in Rio Grande do Sul had
been suspended since May 2021(24).

2.3 Model formulation
We developed a multihost, single-pathogen, coupled multiscale model to simulate FMD epidemic
trajectories (25) and subsequently applied countermeasures. The model led to the development of an
R and Python package entitled "MHASpread: A multihost animal spread stochastic multilevel
model" (version 0.3.0); additional details can be found at https://github.com/machado-
lab/MHASPREAD-model. MHASpread allows the explicit specification of species-specific
transmission probabilities and the latent and infectious periods of a disease that infects multiple
species. The within-farm level includes births and deaths for each species. The between-farm level

includes the entry and exit of animals due to between-farm movements and movements to
slaughterhouses (Figure 1).
139
2.3.1 Within-farm dynamics
For within-farm dynamics, we assume that populations are homogeneously distributed. Species were
homogeneously mixed in farms with at least two species, meaning that the probability of contact
among species was homogeneous regardless of whether species were segregated by barns (e.g.,
commercial swine farms are housed in barns with limited changes in direct contact with cattle). The
within-farm dynamics consist of mutually exclusive health states (i.e., an individual can only be in
one state per discrete time step) for animals of each species (bovines, swine, and small ruminants).
These health states (hereafter, "compartments") include susceptible (S), exposed (E), infectious, (I),
and recovered (R) and are defined as follows:
(i) Susceptible: animals that are not infected and are susceptible to infection.
(ii) Exposed: animals that have been exposed but are not yet infected.
(iii) Infectious: infected animals that can successfully transmit the infection.
(iv) Recovered: animals that have recovered and are no longer susceptible.
Our model considers birth and death, and these parameters are used to update the population
of each farm. The total population is calculated as $N = S + E + I + R$. The number of individuals
within each compartment transitions from $S\beta \rightarrow E, 1/\sigma \rightarrow I, 1/\gamma \rightarrow R$ according to the following
equations:

$$\frac{dS_i(t)}{dt} = u_i(t) - v_i(t) - \frac{\beta S_i(t) I_i(t)}{N_i}$$

$$\frac{dE_i(t)}{dt} = \frac{\beta S_i(t) I_i(t)}{N_i} - v_i(t) E_i(t) - \frac{1}{\sigma} E_i(t)$$

$$\frac{dI_i(t)}{dt} = \frac{1}{\sigma} E_i(t) - \frac{1}{\gamma} I_i(t) - v_i(t) I_i(t)$$

$$\frac{dR_i(t)}{dt} = \frac{1}{\gamma} I_i(t) - v_i(t)$$

Transmission depends on the presence of infected and susceptible host species, as reflected by the
species-specific FMD transmission coefficient β (Table 1.) Births are represented by the number of
animals born alive $u_i(t)$ that enter the S compartment on the farm i at time t according to the day-to-
day records; similarly, $v_i(t)$ represents the exit of the animals from any compartment due to death at
time t. The transition from E to I is driven by $1/\sigma$, and the transition from I to R is driven by $1/\gamma$;
these values are drawn from the distribution generated for each specific species according to the
literature (Table 2). The within-farms dynamics are depicted in Figure 1.
164
165
166

167 **Table 1**. Distribution of host-to-host transmission coefficients (β) per animal^{-1} day.

Infected species	Susceptible taxon	Transmission coefficient (β), shape and distribution	Reference
Bovines	Bovines	PERT (0.18, 0.24, 0.56)	Calculated based on the 2000-2001 FMD outbreaks in the state of Rio Grande do Sul [24].
Bovines	Swine	PERT (0.18, 0.24, 0.56)	Assumed.
Bovines	Small ruminants	PERT (0.18, 0.24, 0.56)	Assumed.
Swine	Bovines	PERT (3.7, 6.14, 10.06)	Assumed [26].
Swine	Swine	PERT (3.7, 6.14, 10.06)	[26].
Swine	Small ruminants	PERT (3.7, 6.14, 10.06)	Assumed [26].
Small ruminants	Bovines	PERT (0.044, 0.105, 0.253)	Assumed [27].
Small ruminants	Swine	PERT (0.006, 0.024, 0.09)	[28].
Small ruminants	Small ruminants	PERT (0.044, 0.105, 0.253)	[27].

169 **Table 2**. Within-farm distributions of latent and infectious FMD parameters for each species.

FMD parameter	Species	Mean, median (25th, 75th percentile) in days	Reference
	Bovines	3.6, 3 (2, 5)	[29].

Latent period, σ	Swine	3.1, 2 (2, 4)	(29).
	Small ruminants	4.8, 5 (3, 6)	(29).
Infectious period, γ	Bovines	4.4, 4 (3, 6)	(29).
	Swine	5.7, 5 (5, 6)	(29).
	Small ruminants	3.3, 3 (2, 4)	(29).

Note: Time is given in days.

2.3.2 Kernel transmission dynamics

Spatial transmission encompasses a range of mechanisms that include airborne transmission, animal
contact over fence lines, and sharing of equipment between farms (30,31). Local spread was fitted via
a spatial transmission kernel in which the likelihood of transmission decreased as a function of the
between-farm distance. The probability PE at time t describes the likelihood that the animals on a
farm become exposed and is calculated as follows:

$$PE_j(t) = 1 - \prod_i \left(1 - \frac{I_i(t)}{N_i} \varphi e^{-\alpha d_{ij}}\right) \qquad (5)$$

where j represents the uninfected population and where ad_{ij} represents the distance between farm j
and infected farm i, within a maximum distance of 40 km. Given the extensive literature on distance-
based FMD dissemination and a previous comprehensive mathematical simulation study (32),
distances above 40 km were not considered. Here, $1 - \frac{I_i(t)}{N_i} \varphi e^{-\alpha d_{ij}}$ represents the probability of
transmission between farms i and j scaled by the prevalence of infection on farm i, $\frac{I_i}{N_i}$, given the
distance between the farms in kilometers (Figure 2). The parameters φ and α control the shape of the
transmission kernel, $\varphi = 0.044$, which is the probability of transmission when $d_{ij} = 0$, and $\alpha =$
0.6 controls the steepness with which the probability decreases with distance (30,31). The
relationship of exposure probability to distance is depicted in Figure 2.

2.4 FMD spread and control actions

We first simulate an initial silent spread over ten days. This procedure yielded a wide range of initial
outbreak scenarios, depicted in Figure 2, before the implementation of control actions. Next, we
outline four different control action scenarios that were described by the Brazilian Ministry of
Agriculture and Livestock (33):
The baseline control scenario, which is referred to hereafter as "base", considers the
following measures: i) depopulation of infected farms; ii) emergency vaccination of the animals on
all farms in the infected and buffer zones; iii) restriction of animal movement for 30 days; and iv)
establishment of three distinct control zones around infected farms at the following distances: 3-km
infected zone, 7-km buffer zone, and 15-km surveillance zone (Supplementary Material Figure S5).

Depopulation: Involves the removal of all animals from infected farms located within the
infected zone(s). For this study, the daily depopulation capacity was set to four farms, a value that
aligns with the maximum depopulation capacity observed in Rio Grande do Sul (Dr. Fernando Groff,
personal communication). Farms with relatively high animal populations were prioritized for
depopulation. Once depopulated, these farms are no longer considered in the simulation. If the daily
depopulation capacity was insufficient to cover all identified infected farms within one day, those
farms were scheduled for depopulation on the following day or as soon as possible, respecting the
maximum capacity constraints.
Emergency vaccination: The bovine farms within the infected and buffer zones are
vaccinated. The daily maximum capacity is ten farms in the infected zone(s) and ten farms in the
buffer zone(s) (Dr. Fernando Groff, personal communication). We simulated the delay in starting
vaccination by setting vaccination to begin seven days post-FMD detection. Farms not vaccinated
within a day due to the limited vaccination capacity were vaccinated on the subsequent day(s).
Additionally, vaccine effectiveness was 90% in 15 days (34,35). For details about the implementation
of emergency vaccination, Supplementary Material Control actions.
Traceability: We utilized contact tracing to identify farms that had direct contact with infected
farms within the past 30 days. These farms underwent surveillance, including clinical examination of
the animals and disease detection. Farms with animals that tested positive during traceback were
categorized as infected farms.
Movement standstill: A 30-day restriction on animal movement across all three control zones
was implemented; this restriction prohibited any incoming or outgoing movement of animals. The
control zones were lifted, and the restriction on movement was maintained until depopulation was
complete.
We assumed that we identified 10% of the infected farms when we began controlling them
(10 days after introducing the index case). For example, if 100 farms were infected initially, 10 of
those farms were found. When the number of detected farms was less than one, we rounded up to one
detected farm. Furthermore, the rate of detection was influenced by two primary factors: the total
number of farms within the control zones and the number of infected farms. For example, when
fewer farms are under surveillance but the number of infected farms is higher, the likelihood of
detection increases (Supplementary Material Figure S6). Infected farms located outside the control
zones were also considered to be among the farms subject to detection.
229
2.4.1 Alternative control scenarios
Base x 2: In this scenario, the daily number of vaccinations was increased to 40 farms, and
depopulation was increased to eight farms. *Base x 3:* This scenario differed in that it assumed 60
farms vaccinated daily and 12 farms depopulated. *Depopulation*. This scenario differed from the
baseline control scenario in that the number of infected farms depopulated was increased to 12 and
vaccination was not used.
236
2.5 Model outputs
Our simulations tracked the number of animals in each health compartment and the number of
infected farms at each time step. The epidemic trajectories were used to calculate the geodetic
distances in km between the seeded index infections and the subsequent infections. In addition, we
determined the probability of distance-dependent transmission by calculating the cumulative
empirical spatial distribution. We utilized a generalized additive (GAM) model to plot the
relationship between the number of infected farms and the number of days across different scenarios,
as well as one-way analysis of variance (ANOVA) with a Tukey post hoc test to compare the
scenarios. This enabled us to explore potential nonlinear relationships between the variables,
effectively capturing complex patterns that might exist in the data. In addition, a mixed-effects

regression model was fitted to describe the relationship between days working on control action and
the initial number of infected farms controlled by each scenario.
249
2.6 Sensitivity analysis
We used a combination of Latin hypercube sampling (LHS), developed by (36), and the partial rank
correlation coefficient (PRCC) technique to perform a local sensitivity analysis. LHS is a stratified
Monte Carlo sampling method without replacement that provides unbiased estimates of modeling
output measures subject to combinations of varying parameters. The PRCC approach can be used to
classify how output measures are influenced by changes in a specific parameter value while linearly
accounting for the effects of other parameters (37). As input model parameters, we selected the
following categories and interspecies interactions: β bovine to bovine, β bovine to swine, β bovine to
small ruminants, σ bovine, γ bovine, β swine to swine, β swine to bovine, β swine to small ruminants,
σ swine, γ swine, β small ruminants to small ruminants, β small ruminants to bovine, β small
ruminants to swine, σ small ruminants, and γ small ruminants. In total, 15 parameters were used to
classify the monotonic relationship of infection status to our input variables to classify model
sensitivity. The inputs include one farm at which the initial conditions were varied across 10,000
simulations over the LHS space. A positive PRCC indicates a positive relationship with the number
of infected animals, whereas a negative PRCC indicates an inverse relationship with the number of
infected animals; however, the magnitude of the PRCC does not necessarily indicate the importance
of a parameter (38).
267
268 **2.7 Software**
269 The MHASpread model was developed, and graphics, tables, and maps were created using R v. 4.1.1
270 (39) and Python v. 3.8.12. Sampler (40), tidyverse (41), sf (42), brazilmaps (43), doParallel (44), and
271 lubridate (45) were the packages used with R v. 4.1.1, and Numpy (46), Pandas (47), and SciPy (48)
272 were the packages used with Python v. 3.8.12. The model is available in both R and Python versions.
273
3   Results
3.1 Initial spread and detection
Initially, we explored the variation in initial infection trends within the first ten days (Figure 3). The
median number of infected farms was 52.5 (IQR: 26.75 to 78.25, maximum 123). The majority of the
infected farms were swine farms (median 43.5, IQR: 22.25 to 64.75, maximum 105); the median
number of infected bovine farms was 43 (IQR: 22 to 64, maximum 85), and the median number of
infected small ruminant farms was 20.5 (IQR: 10.75 to 30.25, maximum 42). The full distribution of
the epidemic trajectories is depicted in Supplementary Figure S7.
282
283 **3.2 Distances from the initial outbreak**
284 Among the 284,396 unique simulated FMD trajectories, the median distance from the seeded
285 infection site to the secondarily infected farm within the first ten days was 4.78 km (IQR: 2.64 km to
286 7.98 km, maximum 6.88 km) (Figure 4 A). Furthermore, we observed a linear increase with time in
287 the distance over which the FMD disseminated (Figure 4 B).
288
3.3 Effectiveness of control measures
All the control scenarios were effective in eliminating all outbreaks within 120 days of the start of the
control measures. However, their effectiveness was significantly different (ANOVA, p-value
<0.05), except when we compared *depopulation* with the *base x 3* scenario. In general, the most
effective alternative scenarios were *base x 3* and *depopulation*. The most notable differences in the
average of infected farms by scenario were between the *base* and *depopulation* scenarios, with mean
differences of 2.36 (95% CI: 2.22 to 2.49), followed by *base x 3* and *base* with 2.26 (95% CI: 2.13 to

2.39) and *base x 2* and *base* with 1.35 (95% CI: 1.22 to 1.49) (Supplementary Material Figure S8). In
addition, we used a generalized additive model (GAM) to visualize the course of the simulated
epidemics over time. Notably, scenario *base x 3* consistently exhibited a lower prevalence over time
than the *depopulation*, *base x 2*, and *base* scenarios (Figure 5).
300
3.3.1 Duration of control actions
The median number of days over which the control actions were implemented for the *base* scenario
was 22 (IQR: 17 to 29, maximum 109). While *base x 2* had a median of 16 days (IQR: 14 to 19,
maximum 51), *base x 3* had a median of 15 days (IQR: 13 to 17, maximum 43), and *depopulation*
had a median of 14 days (IQR: 13 to 17, maximum 45) (Figure 6). In addition, we measured the
similarities and disparities between the mean number of days during which control actions were
implemented, meaning the time period during which at least one outbreak response action was still
ongoing. The comparisons between *depopulation* and *base*, *base x 3* and *base*, and *base x 2* and *base*
revealed substantial disparities in the group means: -9.77 (95% CI: -10.04 to -9.49), -9.83 (95% CI: -
10.10 to -9.56), and -8.09 (95% CI: -8.36 to -7.81), respectively. A complete statistical analysis of
these differences is presented in Supplementary Material Figure S9. Our findings indicate that there
is a positive relationship between the time spent in control action and the number of farms infected at
the beginning of the control actions. We found a linear relationship that indicates that, on average, for
each additional farm that is infected at the beginning of the control actions, the number of days spent
working on control actions increases by approximately 1.59 days (GLM, p-value < 0.05).
316
3.3.2 Vaccination
In the *base* scenario, the daily median number of vaccinated animals was 1,928 (IQR: 1,562 to 3,567,
maximum: 20,740). In the *base x 2* scenario, the median increases to 3,959.32 (IQR: 3,067.75 to
5,865.56, maximum: 25,877). Similarly, in the *base x 3* scenario, the median increases to 5,947 (IQR:
4,157 to 8,384, maximum: 25,006). In the initial 30 days, there was a significant increase in the
number of vaccinated animals, after which the amount of vaccine continued to increase with
decreasing step demand (Figure 7).
324
3.3.3 Depopulation
We analyzed the daily average number of depopulated animals over time. The scenarios *base x 2* and
depopulation had the highest cumulative means of 3,071 (IQR: 1,767 to 3,768, maximum: 4,120) and
2,159 (IQR: 1,314 to 3,000, maximum: 3,830), respectively. The *base* and *base x 3* scenarios
followed closely, with means of 2,139 (IQR: 1,798 to 2,541, maximum: 3,039) and 1,151.93 (IQR:
500 to 2,799, maximum: 4,398), respectively. The *depopulation* scenario consistently resulted in the
highest count of affected animals, followed by the *base x 3*, *base x 2*, and *base* scenarios, particularly
for bovines and small ruminants (Figure 8). However, in the case of swine, the *base* and *base x 2*
scenarios presented higher numbers of depopulated animals than did the other species.
334
3.4 Sensitivity analysis
We evaluated the sensitivity of 15 model parameters and found them to have weights ranging from -
0.86 to 0.72; this sensitivity indicates a limited influence of the model parameters on the number of
simulated secondary infections. Specifically, the length of the latent period, σ, had a negative impact
on the number of secondary infections, and the length of the infectious period, γ, had a positive
influence on the number of secondary infections; the results for both σ and γ were significant (p
value <0.05) for the overall simulated species. The complete results of the sensitivity analysis are
presented in Supplementary Material Figure S10.
343
4   Discussion

This study aimed to develop a multiscale, multihost stochastic model of FMD that explicitly
incorporates species-specific transmission interactions. Our model was used to simulate the spread of
FMD among cattle, buffalo, swine, and small ruminants in Rio Grande do Sul, Brazil, and to examine
the effectiveness of countermeasures. Bovine farms were the most infected, followed by farms with
swine and small ruminants, mostly because of the greater number of cattle farms and the connectivity
of the swine contact network. Most secondary infections spread within a distance of 25 km, showing
that proximity plays an important role in the dynamics of disease transmission. Our simulations
demonstrated that tripling the number of daily depopulations and vaccinations eliminated epidemic
trajectories within 15 days: this required the vaccination of 5,947 animals and the depopulation of
2,139 animals.
The number of farms infected within ten days of the introduction of the disease ranged from 1
to 123, with the majority of simulations (90.12%) resulting in fewer than 50 infected farms. Our
study also revealed that when FMD infects swine, the sizes of the resulting epidemics are significant
(Figure 3). This risk is particularly relevant to areas that harbor dense swine populations associated
with commercial swine production; farms engaged in this type of production are typically vertically
integrated, meaning that they move a significant number of swine, facilitating long-distance spread
(49,50). Our study demonstrated that the number of farms initiating control actions has a linear effect
on the duration of these actions, regardless of the implemented scenario. Specifically, each additional
infected farm extended the duration of control actions by an average of 1.6 days. This indicates that
enhancing the sensitivity of animal disease detection is crucial for optimizing the effectiveness of
control strategies (51). Based on this finding, we argue that improving the timing of detection and
optimizing the response to and the management of outbreaks are pivotal in ensuring effective control.
Compared with the other proposed scenarios, scenario *base x 3* demonstrated the best performance,
requiring a median of 15 days of control actions to eliminate the outbreak (Figure 4). In comparison,
the *base* scenario required a median duration of 22 days, and the *base x 2* scenario required a median
duration of 16 days. An average of 5,947 vaccine doses per day were administered in the *base x 3*
scenario, compared with 1,928 per day in the *base* scenario and 3,959 per day in the *base x 2*
scenario. Finally, the depopulation of 12 farms daily was successful in mitigating outbreaks;
however, this scenario poses a significant challenge for official services.
Emergency vaccination presents an alternative to preemptive culling but may also limit the
ability of surveillance systems to detect infected farms, as it may mask clinical signs of disease
(52,53). When the median number of vaccinated animals at the end of the control action scenarios
was examined, the *base* scenario had the highest median number of vaccinated animals at 305,105,
followed by *base x 2* with 210,786 and *base x 3* with 202,471. Interestingly, despite the higher
vaccination rates, the final average number of vaccinated animals was lower in the scenarios with
increased vaccination rates. This occurs because increasing the vaccination rate reduces the duration
of outbreaks, ultimately resulting in fewer animals to be dosed in the latter part of the control actions
period. Our findings are consistent with the results of studies conducted in Australia, Canada, New
Zealand, and the United States, where emergency vaccination correlated with a reduction in the
number of infected farms and a decrease in outbreak duration(54,55,55–57).
By examining the cumulative number of depopulated animals, our results showed that the
depopulation scenario was more effective in controlling epidemics than the *base* and *base x 2*
scenarios. The primary reason for its greater effectiveness was the high intensity of farm
depopulation per day. Our analysis of the daily average number of depopulated animals over time
revealed that the *base x 2* scenario had the highest mean cumulative count, with 3,071 animals culled,
followed closely by the *depopulation* scenario, in which 2,159 animals were culled. The *base*
scenario had a mean of 2,139 culled animals, whereas the *base x 3* scenario had 1,151 culled animals.
When depopulation of individual species was examined, the *depopulation* scenario consistently
recorded the highest number of culled animals, particularly for bovines and small ruminants. In

contrast, in the *base* and *base x 2* scenarios, there was a greater prevalence of culled swine than of
other species. The main reason for this is that prolonged outbreaks tend to affect more pig farms.
Although there are fewer pig farms than cattle farms, the number of animals per pig farm is
significantly greater (Supplementary Figure S1). While depopulation is an effective countermeasure
to address highly contagious diseases such as FMD (58–60) , culling healthy animals raises ethical
concerns. It also increases economic losses due to reduced production and the need for farmer
compensation (61). Other studies have proposed alternatives to ring depopulation; for example, (18)
simulated a target density strategy and demonstrated its advantages in combating FMD; it resulted in
depopulation of a smaller number of healthy animals than traditional total ring depopulation, and the
time required to eliminate the outbreaks was similar. In addition, we emphasize the importance of
timing when initiating depopulation for FMD control to prevent disease outbreaks across all farms in
the area and potential outward spread (62). Moreover, prolonged delays in culling can lead to
recurrent outbreaks in previously controlled areas, and under specific atmospheric conditions, there is
a risk of long-distance airborne spread (60).
408
4.1 Limitations and further remarks
Since there have been no recent FMD outbreaks in the study region, we extracted parameters from
the data on the FMD outbreak in Rio Grande do Sul (2000 and 2021) and utilized the literature to
obtain the remaining parameters. Our sensitivity analysis did not identify a specific number of
infected animals. Thus, our model has an acceptable level of robustness. The virulence, infectivity,
and transmission of FMD can vary among strains (1,13). Although the most recent outbreaks in the
State of Rio Grande do Sul involved serotypes O and A(24), we cannot rule out the possibility that
other strains that exhibit different dissemination patterns might be introduced. Future work could
include transmission scenarios with strains circulating in neighboring countries.
Additionally, other important between-farm transmission routes, such as vehicles and the
movement of farm staff (both of which have been previously associated with FMD dissemination
(63,64), were not included in our model. If such indirect contact networks are considered, the results
would likely change, and the realism of the model would be improved (55). We assumed 100%
compliance with the restriction on between-farm movement from infected farms and farms directly
linked to infected farms and with the restriction on movement into and from control zones; we also
assumed that the method by which the depopulated animals were disposed of eliminated any
possibility of further dissemination of the virus. Nevertheless, real-world compliance with the control
actions was not examined or considered. Our model can provide a distribution of expected FMD
epidemics for any current or future control actions listed in the Brazilian control and elimination plan
(65). Nevertheless, because our results are based on population data and between-farm movement
data from Rio Grande do Sul, the conclusions drawn from our findings should not be extrapolated to
other regions. However, according to the MHASpread model, infrastructure is highly flexible and can
be easily extended to other Brazilian states and other countries.
432
5   Conclusion
In summary, we demonstrated the importance of including species-specific propagation dynamics in
FMD transmission models designed to assist decision-makers in planning control and mitigation
strategies for FMD. We have shown that a quick response in initiating control actions on a lower
number of infected farms is crucial to reduce the necessary duration of control actions. We found that
increasing depopulation capacity was sufficient to eliminate outbreaks without vaccination.
Eliminating infected or likely-infected animals is an optimal strategy for preventing further
epidemics, but culling large numbers of healthy animals raises welfare concerns. Regardless of which
species in which FMD was introduced, the median distance over which the disease spread was within
25 km, a finding that could explain the effectiveness of the simulated countermeasures within the

control areas used for FMD response. Our model projections, along with the necessary software, are available to local animal health officials. Thus, our model can be used as a policy tool for future responses to FMD epidemics through computer-based preparedness drills and capacity building and during emergency responses to FMD epidemics by providing rules of thumb generated from simulated control scenarios.

6   Conflict of interest

The authors declare that the research reported herein was conducted in the absence of any commercial or financial relationships that could be construed as potential conflicts of interest.

7   Author contributions

NCC and GM conceived the study. FPNL coordinated the data collection. NCC designed and developed computer model code, cleaned and processed the population and movement data, and performed computational analysis. FAM and CT translated scripts from R to Python with assistance from VM and AM. GM secured the funding. NCC and GM wrote and edited the manuscript. All the authors discussed the results and critically reviewed the manuscript.

8   Funding

Fundo de desenvolvimento e defesa sanitária animal (FUNDESA-RS) Award number 2021–1318.

9   Acknowledgments

Fundo de desenvolvimento e defesa sanitária animal (FUNDESA-RS).

10   References

1.   Paton DJ, Gubbins S, King DP. Understanding the transmission of foot-and-mouth disease virus at different scales. *Curr Opin Virol* (2018) 28:85–91. doi: 10.1016/j.coviro.2017.11.013

2.   Bouma A, Elbers ARW, Dekker A, de Koeijer A, Bartels C, Vellema P, van der Wal P, van Rooij EMA, Pluimers FH, de Jong MCM. The foot-and-mouth disease epidemic in The Netherlands in 2001. *Prev Vet Med* (2003) 57:155–166. doi: 10.1016/S0167-5877(02)00217-9

3.   Thompson DK, Muriel P, Russell D, Osborne P, Bromley A, Rowland M, Creigh-Tyte S, Brown C. Economic costs of the foot and mouth disease outbreak in the United Kingdom in 2001: -EN- -FR- -ES-. *Rev Sci Tech OIE* (2002) 21:675–687. doi: 10.20506/rst.21.3.1353

4.   Chanchaidechachai T, de Jong MCM, Fischer EAJ. Spatial model of foot-and-mouth disease outbreak in an endemic area of Thailand. *Prev Vet Med* (2021) 195:105468. doi: 10.1016/j.prevetmed.2021.105468

5.   Gomez F, Prieto J, Galvis J, Moreno F, Vargas J. Identification of Super-Spreaders of Foot-and-Mouth Disease in the cattle transportation network: The 2018 outbreak case in Cesar (Colombia). *2019 4th World Conference on Complex Systems (WCCS)*. Ouarzazate, Morocco: IEEE (2019). p. 1–6 doi: 10.1109/ICoCS.2019.8930765

6.   PANAFTOSA-OPS/OMS. Informe de Situación de los Programas de Erradicación de la Fiebre Aftosa en Sudamérica y Panamá, año 2021. (2021) https://www.paho.org/es/documentos/informe-situacion-programas-erradicacion-fiebre-aftosa-sudamerica-panama-ano-2019 [Accessed April 28, 2022]

7.   Santos DV dos, Silva GS e, Weber EJ, Hasenack H, Groff FHS, Todeschini B, Borba MR,
Medeiros AAR, Leotti VB, Canal CW, et al. Identification of foot and mouth disease risk areas
using a multi-criteria analysis approach. *PLOS ONE* (2017) 12:e0178464. doi:
10.1371/journal.pone.0178464

8.   Seymour RG, Kypraios T, O'Neill PD. Bayesian nonparametric inference for heterogeneously
mixing infectious disease models. *Proc Natl Acad Sci* (2022) 119:e2118425119. doi:
10.1073/pnas.2118425119

9.   Bertram MR, Yadav S, Stenfeldt C, Delgado A, Arzt J. Extinction Dynamics of the Foot-and-
Mouth Disease Virus Carrier State Under Natural Conditions. *Front Vet Sci* (2020) 7:276. doi:
10.3389/fvets.2020.00276

10.   Davies G. The foot and mouth disease (FMD) epidemic in the United Kingdom 2001. *Comp
Immunol Microbiol Infect Dis* (2002) 25:331–343. doi: 10.1016/S0147-9571(02)00030-9

11.   Iriarte MV, Gonzáles JL, De Freitas Costa E, Gil AD, De Jong MCM. Main factors associated
with foot-and-mouth disease virus infection during the 2001 FMD epidemic in Uruguay. *Front
Vet Sci* (2023) 10:1070188. doi: 10.3389/fvets.2023.1070188

12.   Alexandersen S, Zhang Z, Donaldson AI, Garland AJM. The Pathogenesis and Diagnosis of
Foot-and-Mouth Disease. *J Comp Pathol* (2003) 129:1–36. doi: 10.1016/S0021-9975(03)00041-
0

13.   Arzt J, Branan MA, Delgado AH, Yadav S, Moreno-Torres KI, Tildesley MJ, Stenfeldt C.
Quantitative impacts of incubation phase transmission of foot-and-mouth disease virus. *Sci Rep*
(2019) 9:2707. doi: 10.1038/s41598-019-39029-0

14.   Moreno-Torres KI, Delgado AH, Branan MA, Yadav S, Stenfeldt C, Arzt J. Parameterization of
the durations of phases of foot-and-mouth disease in pigs. *Prev Vet Med* (2022) 202:105615.
doi: 10.1016/j.prevetmed.2022.105615

15.   Stenfeldt C, Diaz-San Segundo F, de los Santos T, Rodriguez LL, Arzt J. The Pathogenesis of
Foot-and-Mouth Disease in Pigs. *Front Vet Sci* (2016) 3:
https://www.frontiersin.org/article/10.3389/fvets.2016.00041 [Accessed April 28, 2022]

16.   van Andel M, Tildesley MJ, Gates MC. Challenges and opportunities for using national animal
datasets to support foot-and-mouth disease control. *Transbound Emerg Dis* (2021) 68:1800–
1813. doi: 10.1111/tbed.13858

17.   Kirkeby C, Brookes VJ, Ward MP, Dürr S, Halasa T. A Practical Introduction to Mechanistic
Modeling of Disease Transmission in Veterinary Science. *Front Vet Sci* (2021) 7:546651. doi:
10.3389/fvets.2020.546651

18.   Seibel RL, Meadows AJ, Mundt C, Tildesley M. Modeling target-density-based cull strategies
to contain foot-and-mouth disease outbreaks. *PeerJ* (2024) 12:e16998. doi: 10.7717/peerj.16998

19.   Beck-Johnson LM, Gorsich EE, Hallman C, Tildesley MJ, Miller RS, Webb CT. An exploration
of within-herd dynamics of a transboundary livestock disease: A foot and mouth disease case
study. *Epidemics* (2023) 42:100668. doi: 10.1016/j.epidem.2023.100668

20. Sseguya I, Mugisha JYT, Nannyonga B. Outbreak and control of Foot and Mouth Disease
within and across adjacent districts—A mathematical perspective. *Results Control Optim* (2022)
6:100074. doi: 10.1016/j.rico.2021.100074

21. SEAPI. *Secretaria da Agricultura Pecuaria e Irrigação*. (2024).
http://www.agricultura.rs.gov.br/dda-animal [Accessed July 27, 2021]

22. Sistema de Defesa Agropecuária (SDA). *Secr Agric Pecuária Produção Sustentável E Irrig*
(2024) https://www.agricultura.rs.gov.br/sistema-de-defesa-agropecuaria-sda [Accessed May
28, 2024]

23. Ziegel ER, Lohr SL. Sampling: Design and Analysis. *Technometrics* (2000) 42:223. doi:
10.2307/1271491

24. da Costa JMN, Cobellini LG, Cardenas NC, Groff FHS, Machado G. Assessing epidemiological
parameters and dissemination characteristics of the 2000 and 2001 foot-and-mouth disease
outbreaks in Rio Grande do Sul, Brazil. *bioRxiv* (2022)2022–05.

25. Garira W. A primer on multiscale modelling of infectious disease systems. *Infect Dis Model*
(2018) 3:176–191. doi: 10.1016/j.idm.2018.09.005

26. Eblé P, De Koeijer A, Bouma A, Stegeman A, Dekker A. Quantification of within- and
between-pen transmission of Foot-and-Mouth disease virus in pigs. *Vet Res* (2006) 37:647–654.
doi: 10.1051/vetres:2006026

27. Orsel K, Dekker A, Bouma A, Stegeman JA, De Jong MCM. Quantification of foot and mouth
disease virus excretion and transmission within groups of lambs with and without vaccination.
Vaccine (2007) 25:2673–2679. doi: 10.1016/j.vaccine.2006.11.048

28. Goris NE, Eblé PL, Jong MCM de, Clercq KD. Quantifying foot-and-mouth disease virus
transmission rates using published data. *ALTEX - Altern Anim Exp* (2009) 26:52–54. doi:
10.14573/altex.2009.1.52

29. Mardones F, Perez A, Sanchez J, Alkhamis M, Carpenter T. Parameterization of the duration of
infection stages of serotype O foot-and-mouth disease virus: an analytical review and meta-
analysis with application to simulation models. *Vet Res* (2010) 41:45. doi:
10.1051/vetres/2010017

30. Boender GJ, Hagenaars TJ. Common features in spatial livestock disease transmission
parameters. *Sci Rep* (2023) 13:3550. doi: 10.1038/s41598-023-30230-w

31. Boender GJ, van Roermund HJW, de Jong MCM, Hagenaars TJ. Transmission risks and control
of foot-and-mouth disease in The Netherlands: Spatial patterns. *Epidemics* (2010) 2:36–47. doi:
10.1016/j.epidem.2010.03.001

32. Björnham O, Sigg R, Burman J. Multilevel model for airborne transmission of foot-and-mouth
disease applied to Swedish livestock. *PLOS ONE* (2020) 15:e0232489. doi:
10.1371/journal.pone.0232489

33.   MAPA. Plano Estratégico do PNEFA 2017-2026 — Português (Brasil). *Plano Estratég PNEFA*
(2022) https://www.gov.br/agricultura/pt-br/assuntos/sanidade-animal-e-vegetal/saude-
animal/programas-de-saude-animal/febre-aftosa/plano-estrategico-pnefa-2017-2026 [Accessed
May 20, 2022]

34.   Backer JA, Hagenaars TJ, Nodelijk G, van Roermund HJW. Vaccination against foot-and-
mouth disease I: Epidemiological consequences. *Prev Vet Med* (2012) 107:27–40. doi:
10.1016/j.prevetmed.2012.05.012

35.   Ulziibat G, Raizman E, Lkhagvasuren A, Bartels CJM, Oyun-Erdene O, Khishgee B, Browning
C, King DP, Ludi AB, Lyons NA. Comparison of vaccination schedules for foot-and-mouth
disease among cattle and sheep in Mongolia. *Front Vet Sci* (2023) 10:990043. doi:
10.3389/fvets.2023.990043

36.   Mckay MD, Beckman RJ, Conover WJ. A Comparison of Three Methods for Selecting Values
of Input Variables in the Analysis of Output From a Computer Code. *Technometrics* (2000)
42:55–61. doi: 10.1080/00401706.2000.10485979

37.   Marino S, Hogue IB, Ray CJ, Kirschner DE. A methodology for performing global uncertainty
and sensitivity analysis in systems biology. *J Theor Biol* (2008) 254:178–196. doi:
10.1016/j.jtbi.2008.04.011

38.   Teboh-Ewungkem MI, Ngwa GA eds. *Infectious Diseases and Our Planet*. Cham: Springer
International Publishing. (2021). doi: 10.1007/978-3-030-50826-5

39.   R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R
Foundation for Statistical Computing. (2021). https://www.R-project.org/

40.   Baldassaro M. sampler: Sample Design, Drawing & Data Analysis Using Data Frames. (2019)
https://CRAN.R-project.org/package=sampler

41.   Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, Grolemund G, Hayes A,
Henry L, Hester J, et al. Welcome to the Tidyverse. *J Open Source Softw* (2019) 4:1686. doi:
10.21105/joss.01686

42.   Pebesma E. {Simple Features for R: Standardized Support for Spatial Vector Data}. *R J* (2018)
10:439--446. doi: 10.32614/RJ-2018-009

43.   Prado Siqueira R. brazilmaps: Brazilian Maps from Different Geographic Levels. (2021)
http://github.com/rpradosiqueira/brazilmaps

44.   Corporation M, Weston S. doParallel: Foreach Parallel Adaptor for the "parallel" Package.
(2022) https://CRAN.R-project.org/package=doParallel

45.   Grolemund G, Wickham H. Dates and Times Made Easy with {lubridate}. *J Stat Softw* (2011)
40:1--25.

46.   Harris CR, Millman KJ, Walt SJ van der, Gommers R, Virtanen P, Cournapeau D, Wieser E,
Taylor J, Berg S, Smith NJ, et al. Array programming with NumPy. *Nature* (2020) 585:357–
362. doi: 10.1038/s41586-020-2649-2

47. McKinney W. Data Structures for Statistical Computing in Python. In: Walt S van der,
Millman J, editors. *Proceedings of the 9th Python in Science Conference*. (2010). p. 51–56

48. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E,
Peterson P, Weckesser W, Bright J, et al. SciPy 1.0: Fundamental Algorithms for Scientific
Computing in Python. *Nat Methods* (2020) 17:261–272. doi: 10.1038/s41592-019-0686-2

49. Cardenas NC, Sykes AL, Lopes FPN, Machado G. Multiple species animal movements:
network properties, disease dynamics and the impact of targeted control actions. *Vet Res* (2022)
53:14. doi: 10.1186/s13567-022-01031-2

50. Machado G, Galvis JA, Lopes FPN, Voges J, Medeiros AAR, Cárdenas NC. Quantifying the
dynamics of pig movements improves targeted disease surveillance and control plans.
Transbound Emerg Dis (2021) 68:1663–1675.

51. Perez A, AlKhamis M, Carlsson U, Brito B, Carrasco-Medanic R, Whedbee Z, Willeberg P.
Global animal disease surveillance. *Spat Spatio-Temporal Epidemiol* (2011) 2:135–145. doi:
10.1016/j.sste.2011.07.006

52. Barnett PV, Geale DW, Clarke G, Davis J, Kasari TR. A Review of OIE Country Status
Recovery Using Vaccinate-to-Live Versus Vaccinate-to-Die Foot-and-Mouth Disease Response
Policies I: Benefits of Higher Potency Vaccines and Associated NSP DIVA Test Systems in
Post-Outbreak Surveillance. *Transbound Emerg Dis* (2015) 62:367–387. doi:
10.1111/tbed.12166

53. Humphries B, Ward M. Critically appraised topic: the use of vaccination to control the spread of
foot-and-mouth disease in Australian livestock in the event of an incursion. *Aust Vet J*
(2024)avj.13340. doi: 10.1111/avj.13340

54. Porphyre T, Auty HK, Tildesley MJ, Gunn GJ, Woolhouse MEJ. Vaccination against Foot-And-
Mouth Disease: Do Initial Conditions Affect Its Benefit? *PLoS ONE* (2013) 8:e77616. doi:
10.1371/journal.pone.0077616

55. Rawdon TG, Garner MG, Sanson RL, Stevenson MA, Cook C, Birch C, Roche SE, Patyk KA,
Forde-Folle KN, Dubé C, et al. Evaluating vaccination strategies to control foot-and-mouth
disease: a country comparison study. *Epidemiol Infect* (2018) 146:1138–1150. doi:
10.1017/S0950268818001243

56. Sanson R, Rawdon T, Owen K, Hickey K, van Andel M, Yu Z. Evaluating the benefits of
vaccination when used in combination with stamping-out measures against hypothetical
introductions of foot-and-mouth disease into New Zealand: a simulation study. *N Z Vet J* (2017)
65:124–133. doi: 10.1080/00480169.2016.1263165

57. Traulsen I, Rave G, Teuffert J, Krieter J. Consideration of different outbreak conditions in the
evaluation of preventive culling and emergency vaccination to control foot and mouth disease
epidemics. *Res Vet Sci* (2011) 91:219–224. doi: 10.1016/j.rvsc.2010.12.016

58. Roche SE, Garner MG, Wicks RM, East IJ, De Witte K. How do resources influence control
measures during a simulated outbreak of foot and mouth disease in Australia? *Prev Vet Med*
(2014) 113:436–446. doi: 10.1016/j.prevetmed.2013.12.003

59. Sykes AL, Galvis JA, O'Hara KC, Corzo C, Machado G. Estimating the effectiveness of control
actions on African swine fever transmission in commercial swine populations in the United
States. *Prev Vet Med* (2023) 217:105962. doi: 10.1016/j.prevetmed.2023.105962

60. Tao Y, Probert WJM, Shea K, Runge MC, Lafferty K, Tildesley M, Ferrari M. Causes of
delayed outbreak responses and their impacts on epidemic spread. *J R Soc Interface* (2021)
18:rsif.2020.0933, 20200933. doi: 10.1098/rsif.2020.0933

61. De Menezes TC, Ferreira Filho JBDS, Countryman AM. Potential economic impacts of foot-
and-mouth disease in Brazil: A case study for Mato Grosso and Paraná. *J Agric Appl Econ*
Assoc (2023) 2:481–496. doi: 10.1002/jaa2.73

62. Itao K, Omata F, Nishikawa Y, Oda T, Sasaki T, Zhang C, Solomon Maninang J, Yamaguchi T,
1 Department of Basic Science, Graduate School of Arts and Sciences, The University of
Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo, Japan, 2 Kurihara Central Hospital 3-1-1 Miyano-
Chuo Tukidate Kurihara city, Miyagi, Japan, et al. Threshold phenomena with respect to the
initiation of depopulation in a simple model of foot-and-mouth disease. *Math Biosci Eng* (2019)
16:5931–5946. doi: 10.3934/mbe.2019297

63. Muroga N, Kobayashi S, Nishida T, Hayama Y, Kawano T, Yamamoto T, Tsutsui T. Risk
factors for the transmission of foot-and-mouth disease during the 2010 outbreak in Japan: a
case–control study. *BMC Vet Res* (2013) 9:150. doi: 10.1186/1746-6148-9-150

64. Yang Q, Gruenbacher DM, Heier Stamm JL, Amrine DE, Brase GL, DeLoach SA, Scoglio CM.
Impact of truck contamination and information sharing on foot-and-mouth disease spreading in
beef cattle production systems. *PLOS ONE* (2020) 15:e0240819. doi:
10.1371/journal.pone.0240819

65. MAPA M of A Livestock and Food Supply. National Program for the Eradication and
Prevention of Foot and Mouth Disease - PNEFA. *Natl Program Erad Prev Foot Mouth Dis -*
PNEFA (2021) https://www.gov.br/agricultura/pt-br/assuntos/sanidade-animal-e-vegetal/saude-
animal/programas-de-saude-animal/febre-aftosa/brasil-livre-da-
aftosa/PEPNEFA2019versaoingles.pdf

665
666

667
Figure 1. Schematic of state transitions in within-farm and between-farm dynamics. A) Within-
farm dynamics: The green arrows indicate the introduction of animals (births) into the susceptible (S)
compartment at farm i at time t. Each circle indicates a farm containing single or multiple species
with specific host-to-host transition parameters (σ, γ); the dashed lines represent interactions within
and between host species. The red arrows represent the removal of animals (deaths) regardless of
infection status. B) Between-farm dynamics: The layer represents the number of animals moved
(batches; n) from the farm of origin (i) to a destination farm (j) at time t (indicated by the black
dashed arrows). Animals that moved to the slaughterhouse were removed from the simulation
regardless of their infection status and are indicated by red dashed arrows. The kernel distance
dynamics represent the spatial transmission distances.
678

679
680

681 **Figure 2.** A) The y-axis represents the probability of exposure PE , and the x-axis represents the
682 distance in km. B) Representation of farm locations. The color of each dot represents the probability
683 of PE exposure; in this example, the infected farm is located in the center of the radius.
684

686 **Figure 3. Distribution of the initial infected farms.** The y-axis depicts the logarithm (base 10) of
687 the frequency of infected farms across all simulations. The x-axis represents the number of infected
688 farms.
689

691 **Figure 4. Distribution of the cumulative distance of dissemination.** A) Frequency histogram of
692 secondary infections at varying distances from the seeded infection. B) Empirical cumulative
693 distribution function (1-ECDF) of the probability of infection according to the distance from the
694 initial outbreak after disease introduction. Both x-axes are presented on a log10 scale.
695

Figure 5. Estimated number of infected farms on days 11 through 120. The y-axis represents the
number of infected farms, and the x-axis represents the days from start of simulation until the day
with infected farms. The differently colored lines correspond to the individual scenarios.

Figure 6. Comparison of the number of days spent working on control actions. The y-axis shows
the total number of days dedicated to each control action, and the x-axis presents the scenarios.

Figure 7. Vaccination curve by scenario. The y-axis represents the cumulative average number of

vaccinated animals per day on a log10 scale. The x-axis shows the day on which the animals were

vaccination. The red dashed line represents the curve obtained when control actions were initiated

after 15 days of initial control actions.

Figure 8. Depopulation curves by scenario. The y-axis shows the daily cumulative average

number of depopulated animals. The x-axis represents the days since the start of control actions until

the last depopulated farm. The red dashed line represents the day in which control actions started.