AKT1-FOXO4 AXIS RECIPROACLLY REGULATES HEMOCHORIAL

PLACENTATION

Keisuke Kozai[1,*,§], Ayelen Moreno-Irusta[1,*,¥], Khursheed Iqbal[1], Mae-Lan Winchester[2†],

Regan L. Scott[1], Mikaela E. Simon[1], Masanaga Muto[1‡], Marc R. Parrish[2], and Michael

J. Soares[1,2,3,¥]

[1]Institute for Reproduction and Perinatal Research, Department of Pathology &

Laboratory Medicine, University of Kansas Medical Center, Kansas City, KS

[2]Department of Obstetrics and Gynecology, University of Kansas Medical Center,

Kansas City, KS

[3]Center for Perinatal Research, Children's Mercy Research Institute, Children's Mercy,

Kansas City, MO

*Contributed equally

[¥]Correspondence: amoreno2@kumc.edu or msoares@kumc.edu

[§]Present address: Department of Obstetrics and Gynecology, University of Missouri-Kansas
City School of Medicine, Kansas City, MO
22

23 **[†]Present address:** Department of Obstetrics and Gynecology, University Hospital, Case
24 Western Reserve University, Beachwood, OH 44122
25

26 **[‡]Present address:** Department of Stem Cells and Human Disease Models, Research Center
27 for Animal Life Science, Shiga University of Medical Science, Seta, Tsukinowa-cho, Otsu,
28 Shiga 520-2192, Japan
29

30 **Running title:AKT1 and uteroplacental development**

32 **Keywords: AKT1, FOXO4, trophoblast, placenta, pregnancy, rat, genome editing**

SUMMARY STATEMENT

Genome-edited rat models were utilized to investigate roles for AKT1 and FOXO4 in hemochorial placentation. AKT1 and FOXO4 possess reciprocal actions in regulating development of the hemochorial placenta.

ABSTRACT

Hemochorial placentation involves the differentiation of specialized cells called invasive trophoblast cells possessing the capacity to exit the placenta and invade into the uterus where they restructure the vasculature. Invasive trophoblast cells arise from a well-defined compartment within the placenta, referred to as the junctional zone in the rat and the extravillous trophoblast cell column in the human. In this study, we investigated roles for AKT1, a serine/threonine kinase, in placental development using a genome-edited/loss-of-function rat model. Disruption of AKT1 resulted in placental, fetal, and postnatal growth restriction. Forkhead box O4 (*Foxo4*), which encodes a transcription factor and known AKT substrate, was abundantly expressed in the junctional zone and invasive trophoblast cells of the rat placentation site. *Foxo4* gene disruption using genome-editing resulted in placentomegaly, including an enlarged junctional zone. AKT1 and FOXO4 regulate the expression of many of the same transcripts expressed by trophoblast cells; however, in opposite directions. In summary, we have identified AKT1 and FOXO4 as part of a regulatory network that reciprocally controls critical indices of hemochorial placenta development.

INTRODUCTION

The placenta is an extraembryonic structure essential for normal fetal development (**Maltepe & Fisher, 2015; Burton *et al.,* 2016**). Placentas possess two main functions: i) transformation of the maternal environment to support viviparity and ii) regulation of the transfer of nutrients to the fetus (**Gardner & Beddington, 1988; Soares *et al.,* 2018; Knöfler *et al.,* 2019**). These specialized functions are ascribed to trophoblast cells, which differentiate along a multi-lineage pathway and are situated within specific compartments of the placenta (**Gardner & Beddington, 1988; Soares *et al.,* 2018; Knöfler *et al.,* 2019; Aplin & Jones, 2021**). Placentas come in different shapes, sizes, and connectivity to the mother (**Wooding & Burton, 2008; Roberts *et al.,* 2016**). Placentation in some mammalian species is characterized by trophoblast cells migrating into the maternal uterus where they modify the vasculature facilitating maternal nutrient flow to the placenta (**Pijnenborg *et al.,* 1981; Soares *et al.,* 2018**). This type of placenta is referred to as hemochorial (**Wooding & Burton, 2008; Roberts *et al.,* 2016**). The human and rat possess hemochorial placentation where invasive trophoblast cells migrate deep into the uterine parenchyma (**Pijnenborg *et al.,* 1981; Soares *et al.,* 2018**). Regulation of deep hemochorial placentation is poorly understood. The rat represents a useful animal model for investigating the regulation of deep hemochorial placentation (**Pijnenborg & Vercruysse, 2010; Soares *et al.,* 2012; Shukla & Soares, 2022**).

The rat placenta can be divided into two main compartments: i) junctional zone; ii) labyrinth zone (**Ain *et al.,* 2006; Soares *et al.,* 2012**). The junctional zone compartment of the placenta is situated proximal to the uterine endometrium and is responsible for transforming the maternal environment, whereas the labyrinth zone is located between the junctional zone and fetus where it facilitates nutrient delivery to the fetus (**Knipp *et al.,***

1999; Soares *et al.,* 2012). Junctional zone-specific functions include the production of

peptide and steroid hormones that target maternal organs and the generation of invasive

trophoblast cells that migrate into and restructure the uterine parenchyma (**Soares *et al.*, 1996,**

2012). The extravillous trophoblast cell column is a structure within the human placentation

site, which shares some of these same responsibilities (**Soares *et al.,* 2018; Knöfler *et al.,***

2019). The junctional zone and extravillous trophoblast cell column are pivotal to the

regulation of maternal adaptations to pregnancy, yet little is known about how they are

regulated.

In this report, we focus on the phosphatidylinositol 3-kinase (**PI3K**)/AKT pathway and its

involvement in regulating junctional zone biology. AKT1 is one of three AKT

serine/threonine kinases and represents an integral component of signal transduction

pathways regulating cell proliferation, differentiation, migration, survival, and metabolism

(**Manning & Toker, 2017; Cole *et al.,* 2019**). AKT1 has also been implicated in placentation

and trophoblast cell development through rodent mutagenesis experiments and investigations

with human trophoblast cells (**Kamei *et al.,* 2002; Yang *et al.,* 2003; Qiu *et al.,* 2004; Dash**

***et al.,* 2005; Kent *et al.,* 2010, 2011, 2012; Plaks *et al.,* 2011; Haslinger *et al.,* 2013;**

Sharma *et al.,* 2016). Disruptions in AKT signaling have been connected to trophoblast cell

dysfunction leading to recurrent pregnancy loss, preeclampsia, and infertility (**Pollheimer &**

Knöfler, 2005; Ferretti *et al.,* 2007; Fisher, 2015; Burton & Jauniaux, 2018). Herein we

show that AKT1 inactivation leads to placental and fetal growth restriction in the rat. AKT1

acts via phosphorylation of its target proteins leading to functional changes, including

activation or inhibition of the target protein function (**Manning & Toker, 2017; Cole *et al.,***

2019). We identified forkhead box O4 (**FOXO4**), a transcription factor, as an AKT1 substrate

within the rat junctional zone and in rat trophoblast cells and demonstrated FOXO4

involvement in junctional zone development and the regulation of trophoblast cell

differentiation.

RESULTS

Generation of an *Akt1* mutant rat model

We examined the role of AKT1 in regulating deep placentation in the rat using

CRISPR/Cas9 genome editing. A mutant rat strain possessing a 1,332 bp deletion within the

Akt1 gene was generated (**Fig. 1A and B**). The deletion included part of Exon 4, the entire

region spanning Exon 5 through Exon 6, and part of Exon 7 and led to a frameshift and

premature stop codon (**Fig. 1A and B**). The deletion effectively removed the kinase domain

and regulatory regions of AKT1 (**Fig. 1C**). The *Akt1* mutation was successfully transmitted

through the germline. A rat colony possessing the *Akt1* mutation was established and

maintained via heterozygous x heterozygous breeding. Mating of heterozygotes produced the

predicted Mendelian ratio (**Fig. 1D; Table S1**). Placental tissues possessing a homozygous

deletion within the *Akt1* locus (*Akt1*$^{-/-}$) were deficient in AKT1 protein and exhibited

prominent deficits in pan-AKT and phospho-AKT protein expression (**Fig. 1E**). These

findings support the successful disruption of the *Akt1* locus and are consistent with AKT1

being the predominant AKT isoform within the placenta (**Yang *et al.*, 2003; Kent *et al.*,**

2011; Haslinger *et al.*, 2013).

AKT1 deficiency results in placental, fetal, and postnatal growth restriction

Disruption of the *Akt1* locus in the mouse disrupts placental, fetal, and postnatal growth

(**Chen *et al.*, 2001; Cho *et al.*, 2001; Yang *et al.*, 2003; Plaks *et al.*, 2011; Kent et al., 2012**).

We observed a similar phenotype in the rat. Gestation day (**gd**) 18.5 placental and fetal

weights and postnatal pup weights were significantly smaller in the *Akt1*$^{-/-}$ rat model when

compared to $Akt1^{+/+}$ rats (**Fig. 2A-H**). Junctional and labyrinth zone compartments of the

placenta were also significantly smaller in $Akt1^{-/-}$ placentas (**Fig. 2D, E, I**).

AKT1 regulates junctional zone and invasive trophoblast cell phenotypes

Transcript profiles were determined for $Akt1^{+/+}$ and $Akt1^{-/-}$ gd 18.5 junctional zone tissues

using RNA-sequencing (**RNA-seq**). The size and morphological phenotypes associated with

inactivation of AKT1 were associated with distinct transcript profiles (**Fig. 3; Dataset 1**).

Disruption of AKT1 resulted in upregulation of 254 transcripts and downregulation of 333

transcripts (**Dataset 1**). Pathway analysis included signatures for cell cycle, DNA replication,

cellular senescence, and PI3K-AKT signaling pathways (**Fig. 3A**). Transcripts encoding cell

cycle progression were consistently repressed in the $Akt1^{-/-}$ junctional zones (**Fig. 3A and B**).

In addition, we also observed prominent downregulation of a member of the expanded

prolactin (**PRL**) gene family, *Prl8a4*, and the upregulation of cellular communication

network factor 3 (*Ccn3*, also called *Nov*; **Fig. 3B**).

The junctional zone serves as the source of invasive trophoblast cells entering the uterus.

Consequently, we investigated the uterine-placental interface of $Akt1^{+/+}$ and $Akt1^{-/-}$ placentas

and monitored the surface area occupied by intrauterine invasive trophoblast cells and the

expression of invasive trophoblast cell-specific transcripts. $Akt1^{-/-}$ invasive trophoblast cells

exhibited decreased infiltration into the uterus (**Fig. 3C-E**). We also observed approximately

a 50% decrease in the expression of cytokeratin transcripts, which are expressed by invasive

trophoblast cells within the uterine-placental interface (**Fig. 3F**).

Collectively, the data indicate that AKT1 signaling has profound effects on development

of the junctional zone and the invasive trophoblast cell lineage.

FOXO4 is a target of PI3K/AKT signaling

Forkhead box (**FOX**) transcription factors are known targets of PI3K/AKT signaling and have key roles in regulating developmental processes (**Lam *et al.,* 2013; Schmitt-Ney, 2020; Herman *et al.,* 2021**). We interrogated RNA-seq datasets from wild type gd 18.5 junctional zone for FOX family transcription factors. Transcripts for several FOX transcription factors were detected (transcripts per million, TPM value ≥ 1.0) (**Fig. 4A**). *Foxo4* transcripts were striking in their abundance relative to all other FOX family members. AKT1 did not significantly affect expression levels for any of the FOX family transcripts (**Dataset 1**). *Foxo4* transcripts were specifically localized to the junctional zone and a subset of invasive trophoblast cells localized to uterine spiral arteries, termed endovascular invasive trophoblast cells (**Fig. 4B and C**). Total and phosphorylated FOXO4 protein were significantly diminished in *Akt1* null junctional zones (**Fig. 4D**). We next explored FOXO4 in differentiated rat trophoblast stem (**TS**) cells, a model for trophoblast cell lineages found in the junctional zone (**Asanoma *et al.,* 2011**). *Foxo4* transcript and total and phosphorylated FOXO4 protein showed striking increases in abundance following TS cell differentiation (**Fig. 4E-F**). As previously demonstrated, AKT activity increased following trophoblast cell differentiation (**Kamei *et al.,* 2002; Kent *et al.,* 2010, 2011; Fig. 4G**). AKT activation was required for optimal FOXO4 phosphorylation (**Fig. 4G**). Thus, we established a link between PI3K/AKT signaling and FOXO4 in trophoblast cell lineage development.

Generation of a *Foxo4* mutant rat model

We examined the role of FOXO4 in regulating placentation in the rat using CRISPR/Cas9 genome editing. The *Foxo4* gene consists of four exons and resides on the X chromosome (**Liu *et al.,* 2020**). A mutant rat strain possessing a 3,096 bp deletion within the *Foxo4* gene

was generated (**Fig. 5A and B**). The deletion included the 3' part of Exon 2 and the 5' part of

Exon 3 and led to a frameshift and a premature stop codon (**Fig. 5A and B**). The deletion

effectively disrupted the conserved forkhead DNA binding domain and removed nuclear

localization, nuclear export, and transactivation domains of FOXO4 (**Fig. 5C**). The *Foxo4*

mutation was successfully transmitted through the germline (**Fig. 5D; Table S2**). A rat colony

possessing the *Foxo4* mutation was established and maintained via hemizygous male x wild

type female breeding, which produced the predicted Mendelian ratio (**Table S2**). Junctional

zone tissues possessing a maternally inherited *Foxo4* mutation (*Foxo4*$^{Xm-}$) were deficient in

FOXO4 protein (**Fig. 5E**). The results are consistent with paternal silencing of X

chromosome-linked genes expressed in extraembryonic tissues (**Takagi & Sasaki, 1975;**

West et al., 1978; Hemberger, 2002). FOXO4 was successfully disrupted in the rat.

FOXO4 deficiency results in placentomegaly and a modified junctional zone phenotype

Placentation site phenotypes of mice possessing mutations at the *Foxo4* locus have not

been described (**Liu *et al.,* 2020; Hosaka *et al.,* 2004**). Rats possessing a maternally inherited

mutant *Foxo4* allele (*Foxo4*$^{Xm-}$) exhibited placentomegaly and decreased placental efficiency

(fetal/placental weight ratio) when examined on gd 18.5 (**Fig. 6A-C**) and gd 20.5 (**Fig**

S1.A-C). In contrast, a paternally inherited mutant *Foxo4* allele did not significantly affect

placenta or fetal weights (**Fig S1.D and E**). FOXO4 deficiency associated placentomegaly

included significantly larger junctional and labyrinth zones (**Fig. 6D-G, Fig S1.F-H**) but did

not affect the intrauterine invasive trophoblast cell lineage (**Fig S2**). Transcript profiles were

determined for wild type (*Foxo4*$^{Xm+}$) and *Foxo4*$^{Xm-}$ gd 18.5 junctional zone tissues using

RNA-seq (**Fig. 6H**). Disruption of FOXO4 resulted in upregulation of 369 transcripts and

downregulation of 845 transcripts (**Dataset 2**). Pathway analysis included signatures for

PI3K-AKT signaling, cell cycle, DNA replication, extracellular matrix receptor interaction,

and complement and coagulation pathways (**Fig. 6H**). Junctional adhesion molecule-like

(**JAML**), lipoprotein(a) like 2 (**LPAL2**), erb-b2 receptor tyrosine kinase 3 (**ERBB3**), and

growth factor receptor bound protein 7 (**GRB7**) were each conspicuous in their prominent

downregulation in FOXO4 deficient junctional zone tissue (**Fig. 6I**). JAML contributes to

epithelial barrier function, modulates immune cell trafficking, and angiogenesis (**Kummer &**

Ebnet, 2018), whereas LPAL2 is a long noncoding RNA contributing to inflammatory and

oxidative stress responses (**Han *et al.,* 2018**). JAML and LPAL2 have not previously been

linked to trophoblast or placental biology. ERBB3 is a receptor for neuregulin 1 and promotes

trophoblast cell survival (**Fock *et al.,* 2015**) and GRB7 is an adaptor protein participating in

signal transduction activated through ERBB3 (**Fiddes *et al.,* 1998**). Interestingly, numerous

junctional zone transcripts regulated by FOXO4 were reciprocally regulated by AKT1 (88%

of transcripts upregulated in *Akt1* null were downregulated in *Foxo4* mutant tissues; 38% of

transcripts downregulated in *Akt1* null were upregulated in *Foxo4* mutant tissues; **Fig. 7**). The

reciprocal relationship between AKT1 and FOXO4 is evident at structural and molecular

levels.

FOXO4 contributes to the regulation of the trophoblast cell lineage

We next modeled junctional zone cell biology using rat TS cells. The consequences of

FOXO4 disruption in differentiating rat TS cells were examined. FOXO4 expression was

inhibited via ectopic expression of short hairpin RNAs specific to *Foxo4* (**Fig. 8A and B**).

Although, a morphologic phenotype was not evident, prominent differences in the

transcriptomes of TS cells expressing control versus *Foxo4* shRNAs were observed (**Fig. 8C**).

Disruption of FOXO4 resulted in upregulation of 260 transcripts and downregulation of 443

transcripts (**Dataset 3**). Pathway analysis included signatures for PI3K-AKT signaling,

longevity regulating, calcium signaling, and glutathione metabolism pathways (**Fig. 8C**).

Among the dysregulated transcripts was an upregulation of matrix metallopeptidase 12, a

known constituent of endovascular invasive trophoblast cells (**Harris et al., 2010;**

Chakraborty et al., 2016) and downregulation of trophoblast specific protein alpha, a

transcript characteristic of spongiotrophoblast cells within the junctional zone (**Iwatsuki *et***

al., 2000). FOXO4 is a known regulator of responses to oxidative stress (**Liu *et al.,* 2020**).

Several transcripts associated with inflammatory and cellular stress responses, including

thioredoxin interacting protein, glutathoione S-transferase mu 1, arachidonate 5-lipoxygenase

activating protein, interferon kappa, nuclear protein 1, and *Lpal2*, were prominently

downregulated (**Sies & Cadenas, 1985; LaFleur *et al.,* 2001; Mashima & Okuyama, 2015;**

Han *et al.,* 2018; Huang *et al.,* 2021; Nirgude & Choudhary, 2021; Qayyum *et al.,* 2021;

Satapathy & Wilson, 2021).

Key findings

Collectively, the results indicate that AKT1 drives placental growth, including regulation

of deep intrauterine trophoblast cell invasion. These actions are accomplished, at least in part,

through modulation of FOXO4, which acts to restrain placental growth and coordinate

responses to physiological stressors.

DISCUSSION

The rat and human possess a type of hemochorial placentation where specialized

trophoblast cells penetrate deep into the uterus and transform the uterine parenchyma,

including the vasculature (**Pijnenborg *et al.,* 1981; Pijnenborg & Vercruysse, 2010; Soares**

***et al.,* 2012**). Central to deep placentation is the source of invasive trophoblast cells, which in

the rat is a compartment within the placenta referred as the junctional zone and, in the human,

the extravillous trophoblast cell column (**Soares *et al.,* 2012, 2018; Knöfler *et al.,* 2019**). In

addition to an intrauterine role, the cellular constituents of these placental compartments

produce hormones directed to maternal tissues with actions that ensure in utero survival and

promotion of fetal growth (**Soares *et al.,* 1996; John, 2017**). In this report, AKT1 and

FOXO4 were identified as regulators of rat junctional zone development. In vivo disruption

of AKT1 and FOXO4 led to opposite effects on junctional zone development. AKT1

deficiency resulted in growth restriction of the junctional zone, phenotypic alteration of the

invasive trophoblast cell lineage, as well as compromised fetal and postnatal growth, and

AKT1 was capable of phosphorylating FOXO4 in rat trophoblast cells. Deficiency of FOXO4

resulted in an expanded junctional zone. Deficits in AKT1 or FOXO4 also impacted

transcriptomic profiles of the junctional zone. The findings indicate that AKT1 and FOXO4

are part of a gene regulatory network controlling hemochorial placentation.

AKT1 signaling influenced placental development. *Akt1* null mutations in the mouse and

rat yield similar phenotypes characterized by placental, fetal, and postnatal growth restriction

(**Chen *et al.,* 2001; Cho *et al.,* 2001; Yang *et al.,* 2003; Plaks *et al.,* 2011; Kent et al., 2012**).

Smaller junctional zones accompanying AKT1 deficiency were associated with a

downregulation of transcripts encoding proteins driving cell proliferation. These results imply

that size differences of junctional zone compartments in the wild type versus *Akt1* nulls were

related to, at least in part, diminished trophoblast cell proliferation in *Akt1* junctional zone

tissues. The data also fit well with known actions of AKT signaling promoting cell

proliferation in a wide range of cell types (**Manning & Cantley, 2007; Manning & Toker,**

2017; Cole *et al.,* 2019). AKT1 disruption also altered differentiated junctional zone

trophoblast cell phenotypes. As cellular constituents of the junctional zone differentiate, they

acquire the capacity to express several members of the expanded prolactin (**PRL**) family of

hormones/cytokines (**Soares, 2004; Alam *et al.,* 2006; Soares *et al.,* 2007**). The expression

of PRL8A4, a member of the expanded PRL family, was dramatically downregulated in *Akt1* null junctional zones. PRL8A4 is an orphan ligand with little known of its significance to the biology of pregnancy other than as a signature feature of the differentiated junctional zone phenotype (**Iwatsuki *et al.,* 1998; Soares *et al.,* 2007**). AKT signaling has previously been implicated in the regulation of the differentiation of rodent and human trophoblast cells (**Kamei *et al.,* 2003; Kent *et al.,* 2010, 2011; Haslinger *et al.,* 2013**). Disruption of *Akt1* also interfered with invasive trophoblast cell development. Trophoblast cell infiltration into the uterine-placental interface was diminished in *Akt1* nulls. Involvement of AKT signaling has also been implicated in the development of the human extravillous trophoblast cell lineage (**Pollheimer & Knöfler, 2005; Haslinger *et al.,* 2013; Morey *et al.,* 2021**). AKT signaling could affect invasive trophoblast cell development through its actions on their origin in the junctional zone and EVT cell column or alternatively, their maturation as they invade into the uterus. Finally, the impact of AKT signaling in invasive trophoblast cell development may be more profound than observed with AKT1 deficiency due to compensatory activities of AKT2 and AKT3 (**Kent *et al.,* 2011; Haslinger *et al.,* 2013**).

AKT1 regulates cellular function through its actions as a serine/threonine kinase and thus, phosphorylation of its substrates (**Manning & Cantley, 2007; Manning & Toker, 2017; Cole *et al.,* 2019**). The forkhead box (**FOX**) family of transcription factors are well-established targets of AKT action (**Lam *et al.,* 2013; Schmitt-Ney, 2020; Herman *et al.,* 2021**). Among FOX family members, FOXO4 expression was uniquely elevated in the rat junctional zone. FOXO4 phosphorylation state in trophoblast cells was affected by AKT signaling. AKT-mediated phosphorylation of FOXO4 leads to FOXO4 exit from the nucleus and inactivation (**Schmitt-Ney, 2020; Herman *et al*., 2021**), which might suggest that an AKT1 deficiency would result in the stabilization of FOXO4 protein in the junctional zone.

Instead, AKT1 deficiency led to depletion of junctional zone total and phosphorylated

FOXO4 proteins. Consequently, the observed *Akt1* null placental phenotype was associated

with the depletion of both AKT1 and FOXO4 proteins. In addition to regulation by AKT

signaling, FOXO4 activity/stability is stimulated by Jun kinase and monoubiquitylation

(**Essers *et al.,* 2004; van der Horst *et al.,* 2006; Brenkman *et al.,* 2008; Liu *et al.,* 2020**),

while inhibited by acetylation and polyubiquitylation (**Fukuoka *et al.,* 2003; Huang &**

Tindall, 2011; Liu *et al.,* 2020). Whether AKT1 indirectly affects FOXO4 protein via

impacting these other FOXO4 regulators remains to be determined.

FOXO4 is a transcription factor implicated in the regulation of the cell cycle, apoptosis,

responses to oxidative stress, and a range of disease processes (**Liu *et al.,* 2020**). The original

characterization of the *Foxo4* null mouse concluded that FOXO4 did not have a singular role

in the pathophysiology of the mouse (**Hosaka *et al.,* 2004**). The absence of a reported

phenotype for the *Foxo4* null mouse model was attributed to the compensatory actions of

other members of the FOXO family, including FOXO1, FOXO3, and possibly FOXO6

(**Hosaka *et al.,* 2004; Liu *et al.,* 2020**). We describe a prominent placental phenotype for the

Foxo4 null rat model. The placental anomalies associated with FOXO4 deficiency were

compatible with the production of viable offspring. The absence of a fertility defect in the

Foxo4 null mouse model likely precluded a closer examination of placentation (**Hosaka *et al.,***

2004). However, it is also possible that elements of FOXO4 action are species specific.

FOXO4 is prominently expressed in the junctional zone and to a lesser extent in invasive

trophoblast cells. Disruption of FOXO4 led to an expansion of both junctional and labyrinth

zone placental compartments, which probably reflects cell autonomous and non-cell

autonomous actions, respectively. A striking reciprocal pattern of AKT1 versus FOXO4 gene

regulation within the junctional zone was demonstrated and included differentially regulated

transcripts encoding proteins involved in the regulation of cell proliferation and cell death.

We surmise that AKT1 promotes junctional zone growth via stimulating the expression of

transcripts connected to cell cycle progression and inhibited transcripts connected to cell

death, whereas the converse is true for FOXO4. These biological roles are consistent with the

known actions of AKT1 and FOXO4 in other cell systems (**Manning & Toker, 2017; Liu** *et*

al., **2020; Herman** *et al.,* **2021**). A transcriptional regulatory network involving FOXO4 has

also been identified in human extravillous trophoblast cells (**Morey** *et al.,* **2021**). Importantly,

FOXO4 also regulates trophoblast cell responses to oxidative stress and is thus, positioned to

contribute to placental adaptations to a compromised maternal environment and in disease

states affecting placentation.

MATERIALS AND METHODS

Animals

Holtzman Sprague-Dawley rats were maintained in an environmentally controlled facility

with lights on from 0600 to 2000 h with food and water available ad libitum. Time-mated

pregnancies were established by co-housing adult female rats (8-10 weeks of age) with adult

male rats (>10 weeks of age). Detection of sperm or a seminal plug in the vagina was

designated gd 0.5. Pseudopregnant females were generated by co-housing adult female rats

(8-10 weeks of age) with adult male vasectomized males (>10 weeks of age). The detection

of seminal plugs was designated pseudopregnancy day 0.5. Four to five-week-old donor rats

were superovulated by intraperitoneal injection of pregnant mare serum gonadotropin (30

units, G4877, Sigma-Aldrich, St. Louis, MO), followed by an intraperitoneal injection of

human chorionic gonadotropin (30 units, C1063, Sigma-Aldrich) ~46 h later, and

immediately mated with adult males. Zygotes were flushed from oviducts the next morning

(gd 0.5). The University of Kansas Medical Center (**KUMC**) Animal Care and Use

Committee approved all protocols involving the use of rats.

Tissue collection and analysis

Rats were euthanized by CO_2 asphyxiation at designated days of gestation. Uterine

segments containing placentation sites were frozen in dry ice-cooled heptane and stored at

-80ºC until processed for histological analyses. Alternatively, placentation sites were

dissected into placentas, the adjacent uterine-placental interface tissue (also referred to as the

metrial gland), and fetuses as previously described (**Ain *et al.,* 2006**). Placentas were weighed

and dissected into junctional zone and labyrinth zone compartments (**Ain *et al.,* 2006**).

Placental compartments and uterine-placental interfaces were frozen in liquid nitrogen and

stored at -80ºC until used for biochemical analyses. Fetuses were weighed, genotyped, and

sex determined by polymerase chain reaction (**PCR**) (**Dhakal & Soares, 2017**).

Generation of *Akt1* and *Foxo4* mutant rat models

Mutations at *Akt1* and *Foxo4* loci were generated using CRISPR/Cas9 genome editing

(**Kaneko, 2017; Iqbal *et al.,* 2021**). Guide RNAs targeting Exon 4 (target sequence:

GCCGTTTGAGTCCATCAGCC; nucleotides 356-375) and Exon 7 (target sequence:

TTGTCATGGAGTACGCCAAT; nucleotides 712-731) of the *Akt1* gene (NM_033230.3) or

targeting Exon 2 (target sequence: CCAGATATACGAATGGATGGTCC; nucleotides

517-539) and Exon 3 (target sequence: GTTCATCAAGGTACATAACGAGG; nucleotides

631-653) of the *Foxo4* gene (NM_001106943.1) were electroporated into single-cell rat

embryos using the NEPA21 electroporator (Nepa Gene Co Ltd, Ichikawa City, Japan).

Electroporated embryos were transferred to oviducts of day 0.5 pseudopregnant rats. Initially,

offspring were screened for *Akt1* or *Foxo4* mutations from genomic DNA from tail-tip

biopsies using the REDExtract-N-Amp[TM] Tissue PCR kit (XNAT, Millipore Sigma,

Burlington, MA). PCR was performed on the purified DNA samples using primers flanking

the guide RNA sites, and products resolved by agarose gel electrophoresis and ethidium

bromide staining. Genomic DNA containing potential mutations was amplified by PCR, gel

purified, and precise boundaries of deletions determined by DNA sequencing (Genewiz Inc.,

South Plainfield, NJ). Founders with *Akt1* or *Foxo4* mutations were backcrossed to wild type

rats to demonstrate germline transmission. Routine genotyping was performed by PCR on

genomic DNA with specific sets of primers (**Table S3**).

Western blot analysis

Tissue lysates were prepared with radioimmunoprecipitation assay lysis buffer system

(sc-24948A, Santa Cruz Biotechnology, Santa Cruz, CA). Protein concentrations were

determined using the DC^{TM} Protein Assay Kit (5000112, Bio-Rad Laboratories, Hercules,

CA). Proteins (20 μg/lane) were separated by SDS-PAGE. Separated proteins were

electrophoretically transferred to polyvinylidene difluoride membranes (10600023, GE

Healthcare, Milwaukee, WI) for 1 h at 100 V on ice. Membranes were subsequently blocked

with 5% milk or 5% bovine serum albumin for 1 h at room temperature and probed separately

with specific primary antibodies to AKT1 (1:1,000 dilution, 75692, Cell Signaling

Technology, Danvers, MA), pan-AKT (1:1,000 dilution, 4691, Cell Signaling Technology),

phospho-Ser^{473} AKT (1:2,000 dilution, 4060, Cell Signaling Technology), FOXO4 (1:2,000

dilution, 21535-1-AP, Proteintech, Rosemont, IL), phospho-Ser^{262} FOXO4 (1:3,000 dilution,

ab126594, Abcam, Cambridge, MA), and glyceraldehyde 3-phosphate dehydrogenase

(**GAPDH**, 1:5,000 dilution, ab8245, Abcam) in Tris-buffered saline with Tween 20 (**TBST**)

overnight at 4□. After primary antibody incubation, the membranes were washed in TBST

three times for ten min each at room temperature. After washing, the membranes were

incubated with anti-rabbit or anti-mouse immunoglobulin G (**IgG**) conjugated to horseradish

peroxidase [**HRP**, 1:5,000 dilution or 1:20,000 dilution (phospho-Ser^{262} FOXO4), 7074S and

7076S, Cell Signaling Technology] in TBST for 1 h at room temperature, washed in TBST

three times for ten min each at room temperature, immersed in Immobilon Crescendo

Western HRP Substrate (WBLUR0500, Sigma-Aldrich), and luminescence detected using

Radiomat LS film (Agfa Healthcare, Mortsel, Belgium).

Transcript analysis

Total RNA was extracted from tissues using TRI Reagent Solution (AM9738,

Thermo-Fisher, Waltham, MA) according to the manufacturer's instructions. Total RNA (1

µg) was reverse transcribed using a High-Capacity cDNA Reverse Transcription Kit

(4368813, Thermo-Fisher). Complementary DNAs were diluted 1:10 and subjected to reverse

transcription-quantitative PCR (**RT-qPCR**) using PowerUp SYBR Green Master Mix

(A25742, Thermo-Fisher) and primers provided in **Table S4**. QuantStudio 5 Flex Real-Time

PCR System (Applied Biosystems, Foster City, CA) was used for amplification and

fluorescence detection. PCR was performed under the following conditions: 95□ for 10 min,

followed by 40 cycles of 95□ for 15 sec and 60□ for 1 min. Relative mRNA expression was

calculated using the delta-delta Ct method. *Gapdh* was used as a reference transcript.

RNA-seq analysis

Transcript profiles were generated from wild type and *Akt1$^{-/-}$*, and *Foxo4^{Xm-}* junctional

zone tissues, and rat differentiated TS cells expressing control or *Foxo4* shRNAs.

Complementary DNA libraries from total RNA samples were prepared with Illumina TruSeq

RNA preparation kits according to the manufacturer's instructions (Illumina, San Diego, CA).

RNA integrity was assessed using an Agilent 2100 Bioanalyzer (Santa Clara, CA). Barcoded

cDNA libraries were multiplexed onto a TruSeq paired-end flow cell and sequenced (100-bp

paired-end reads) with a TruSeq 200-cycle SBS kit (Illumina). Samples were run on an

Illumina NovaSeq 6000 sequencer at the KUMC Genome Sequencing Facility. Reads from

*.fastq files were mapped to the rat reference genome (Ensembl Rnor_5.0.78) using CLC

Genomics Workbench 12.0 (Qiagen, Germantown, MD). Transcript abundance was expressed

as transcript per million mapped reads (TPM) and a *P* value of 0.05 was used as a cutoff for

significant differential expression. Statistical significance was calculated by empirical

analysis of digital gene expression followed by Bonferroni's correction. Pathway analysis

was performed using Datbase for Annotation, Visualization, and Integrated Discovery

(**DAVID; Huang *et al.,* 2009**).

Immunohistochemistry

Placentation sites were embedded in optimum cutting temperature (**OCT**) compound and

sectioned at 10 μm thickness. Sections were fixed in 4% paraformaldehyde, washed in

phosphate buffered saline (pH 7.4) three times for five min each, blocked with 10% normal

goat serum (50062Z, Thermo-Fisher), and incubated overnight with primary antibodies: pan

cytokeratin (1:300 dilution, F3418, Sigma-Aldrich) to identify trophoblast cells and vimentin

(1:300 dilution, sc-6260, Santa Cruz Biotechnology) to distinguish placental compartments.

After washing with phosphate buffered saline (pH 7.4), sections were incubated with

corresponding secondary antibodies: Alexa 568-conjugated goat anti-rabbit IgG (1:500

dilution, A11011, Thermo-Fisher) or rabbit anti-mouse IgG conjugated to HRP (1:500

dilution, A9044, Sigma-Aldrich) for 3 h at room temperature and color development with a

3-amino-9-ethylcarbazole (**AEC**) substrate kit (SK-4200, Vector Laboratories). Sections were

then mounted with Fluoromount-G mounting media (0100-01, Southern Biotech,

Birmingham, AL) and examined microscopically. Fluorescence images were captured on a

Nikon 80i upright microscope (Nikon) with a Photometrics CoolSNAP-ES monochrome

camera (Roper). The area occupied by cytokeratin-positive cells (invasive trophoblast cells)

within the uterine-placental interface was quantified using ImageJ software, as previously

described (**Nteeba *et al.*, 2020**).

In situ hybridization

Distributions of transcripts for *Foxo4* and *Prl7b1* were determined on cryosections of rat

placentation sites. RNAscope Multiplex Fluorescent Reagent Kit version 2 (Advanced Cell

Diagnostics, Newark, CA) was used for in situ hybridization analysis. Probes were prepared

to detect *Foxo4* (NM_001106943.1, 1038981-C1, target region: 750-1651) and *Prl7b1*

(NM_153738.1, 860181-C2, target region: 28-900). Fluorescence images were captured on a

Nikon 80i upright microscope (Nikon) with a Photometrics CoolSNAP-ES monochrome

camera (Roper).

Rat TS cell culture

Blastocyst-derived rat TS cells (**Asanoma *et al*., 2011**) were cultured in Rat TS Cell

Medium [RPMI 1640 medium (11875093, Thermo-Fisher), 20% (vol/vol) fetal bovine serum

(**FBS**, Thermo-Fisher), 100 µM 2-mercaptoethanol (M7522, Sigma-Aldrich), 1 mM sodium

pyruvate (11360-070, Thermo-Fisher), 100 µM penicillin and 50 U/mL streptomycin

(15140122, Thermo-Fisher)] supplemented with 70% rat embryonic fibroblast

(REF)-conditioned medium prepared as described previously (Asanoma *et al*., 2011), 25

ng/ml fibroblast growth factor 4 (**FGF4**; 100-31, Peprotech), and 1 µg/mL heparin (H3149,

Sigma-Aldrich). For induction of differentiation, rat TS cells were cultured for 15 days in rat

TS medium containing 1% (vol/vol) FBS without FGF4, heparin, and REF-conditioned

medium. In some experiments, rat TS cells were exposed to a phosphatidylinositol 3-kinase

(**PI3K**) inhibitor (LY294002, 10 µM, 9901, Cell Signaling Technology).

Lentivirus construction and production

Lentivirus construction and production were described previously (**Muto *et al.*, 2021; Varberg et al., 2021**). Briefly, the lentivirus encoding the shRNA targeting *Foxo4* was constructed using a pLKO.1 vector. shRNA oligo sequences used in the analyses are provided in **Table S5**. Lentiviral packaging vectors were obtained from Addgene and included pMDLg/pRRE (plasmid 12251), pRSV-Rev (plasmid 12253), pMD2.G (plasmid 12259). Lentiviral particles were produced using Attractene (301005, Qiagen) in human embryonic kidney (HEK) 293FT (Thermo-Fisher) cells.

Lentiviral transduction

Rat TS cells were incubated with lentiviral particles for 24 h followed by selection with puromycin dihydrochloride (5 µg/mL; A11138-03, Thermo-Fisher) for two days. Cells were then cultured for 1-3 days in Rat TS Cell Medium prior to differentiation.

Statistical analysis

Student's *t*-test, Welch's *t*-test, Dunnett's test, or Steel test were performed, where appropriate, to evaluate the significance of the experimental manipulations. Results were deemed statistically significant when $P < 0.05$.

ACKNOWLEDGEMENTS

The research was supported by postdoctoral fellowships from the Kansas Idea Network of Biomedical Research Excellence, P20 GM103418 (A.M-I.), Lalor Foundation (K.K., A.M-I., M.M.), American Heart Association (K.K., M.M.), and an NIH National Research Service Award, HD104495 (R.L.S.) and NIH grants (HD020676, HD079363, HD099638,

HD105734), and the Sosland Foundation. We also thank Stacy Oxley and Brandi Miller for

administrative assistance.

AUTHOR CONTRIBUTIONS

K.K., A.M.-I., and M.J.S. conceived and designed the research; K.K., A.M.-I., K.I.,

M.-L.W., R.L.S., M.E.S., M.M. performed experiments; K.K., A.M.-I., K.I., M.R.P., and

M.J.S. analyzed the data and interpreted results of experiments; K.K., A.M.-I., and M.J.S.

prepared figures and manuscript; All authors read, contributed to editing, and approved the

final version of manuscript.

CONFLICT OF INTEREST

There is no conflict of interest that could be perceived as prejudicing the impartiality of

the research reported.

DATA AVAILABILITY

RNA-seq datasets are available at the Gene Expression Omnibus (**GEO**) database,

https://www.ncbi.nlm.nih.gov/geo/ (accession number GSE205831). All data generated and

analyzed in this study are included in the published article and supporting files. Resources

generated from the research are available from the corresponding author upon reasonable

request. AKT1 and FOXO4 mutant rat models are available through the Rat Resource and

Research Center (Columbia, MO).

REFERENCES

Ain R, Konno T, Canham LN, Soares MJ (2006) Phenotypic analysis of the rat placenta.

Methods Mol Med 121: 295-313

Alam SM, Ain R, Konno T, Ho-Chen JK, Soares MJ (2006) The rat prolactin gene family

locus: species-specific gene family expansion. *Mamm Genome* 17: 858-877

Aplin JD, Jones CJP (2021) Cell dynamics in human villous trophoblast. *Hum Reprod*

Update 27: 904-922

Asanoma K, Rumi MA, Kent LN, Chakraborty D, Renaud SJ, Wake N, Lee DS, Kubota

K, Soares MJ (2011) FGF4-dependent stem cells derived from rat blastocysts differentiate

along the trophoblast lineage. *Dev Biol* 351: 110-119

Brenkman AB, de Keizer PL, van den Broek NJ, Jochemsen AG, Burgering BM (2008)

Mdm2 induces mono-ubiquitination of FOXO4. *PLoS One* 3: e2819

Burton GJ, Fowden AL, Thornburg KL (2016) Placental origins of chronic disease. *Physiol*

Rev 96: 1509-1565

Burton GJ, Jauniaux E (2018) Pathophysiology of placental-derived fetal growth restriction.

Am J Obstet Gynecol 218(Suppl): S745-S761

Chakraborty D, Cui W, Rosario GX, Scott RL, Dhakal P, Renaud SJ, Tachibana M,

Rumi MA, Mason CW, Krieg AJ, Soares MJ (2016) HIF-KDM3A-MMP12 regulatory

circuit ensures trophoblast plasticity and placental adaptations to hypoxia. *Proc Natl Acad Sci*

USA 113: E7212-E7221

Chen WS, Xu PZ, Gottlob K, Chen ML, Sokol K, Shiyanova T, Roninson I, Weng W,

Suzuki R, Tobe K, et al. (2001) Growth retardation and increased apoptosis in mice with

homozygous disruption of the Akt1 gene. *Genes Dev* 15: 2203-2208

Cho H, Thorvaldsen JL, Chu Q, Feng F, Birnbaum MJ (2001) Akt1/PKBalpha is required

for normal growth but dispensable for maintenance of glucose homeostasis in mice. *J Biol*

Chem 276: 38349-38352

Cole PA, Chu N, Salguero AL, Bae H (2019) AKTivation mechanisms. *Curr Opin Struct Biol*

59: 47-53

Dash PR, Whitley GS, Ayling LJ, Johnstone AP, Cartwright JE (2005) Trophoblast

apoptosis is inhibited by hepatocyte growth factor through the Akt and beta-catenin mediated

up-regulation of inducible nitric oxide synthase. *Cell Signal* 17: 571-580

Dhakal P, Soares MJ (2017) Single-step PCR-based genetic sex determination of rat tissues

and cells. *Biotechniques* 62: 232-233

Essers MA, Weijzen S, de Vries-Smits AM, Saarloos I, de Ruiter ND, Bos JL, Burgering

BM (2004) FOXO transcription factor activation by oxidative stress mediated by the small

GTPase Ral and JNK. *EMBO J* 23: 4802–4812

Ferretti C, Bruni L, Dangles-Marie V, Pecking AP, Bellet D (2007) Molecular circuits

shared by placental and cancer cells, and their implications in the proliferative, invasive and

migratory capacities of trophoblasts. *Hum Reprod Update* 13: 121-141

Fiddes RJ, Campbell DH, Janes PW, Sivertsen SP, Sasaki H, Wallasch C, Daly RJ (1998)
Analysis of Grb7 recruitment by heregulin-activated erbB receptors reveals a novel target
selectivity for erbB3. *J Biol Chem* 273: 7717-7724

Fisher SJ (2015) Why is placentation abnormal in preeclampsia? *Am J Obstet Gynecol*
213(Suppl): S115-S122

Fock V, Plessl K, Draxler P, Otti GR, Fiala C, Knöfler M, Pollheimer J (2015)
Neuregulin-1-mediated ErbB2-ErbB3 signalling protects human trophoblasts against
apoptosis to preserve differentiation. *J Cell Sci* 128: 4306-4316

Fukuoka M, Daitoku H, Hatta M, Matsuzaki H, Umemura S, Fukamizu A (2003)
Negative regulation of forkhead transcription factor AFX (Foxo4) by CBP-induced acetylation.
Int J Mol Med 12: 503–508

Gardner RL, Beddington RS (1988) Multi-lineage 'stem' cells in the mammalian embryo. *J
Cell Sci Suppl* 10: 11-27

Han BW, Ye H, Wei PP, He B, Han C, Chen ZH, Chen YQ, Wang WT (2018) Global
identification and characterization of lncRNAs that control inflammation in malignant
cholangiocytes. *BMC Genomics* 19: 735

**Harris LK, Smith SD, Keogh RJ, Jones RL, Baker PN, Knöfler M, Cartwright JE,
Whitley GS, Aplin JD** (2010) Trophoblast- and vascular smooth muscle cell-derived MMP-12

mediates elastolysis during uterine spiral artery remodeling. *Am J Pathol* 177: 2103-2115

Haslinger P, Haider S, Sonderegger S, Otten JV, Pollheimer J, Whitley G, Knöfler M

(2013) AKT isoforms 1 and 3 regulate basal and epidermal growth factor-stimulated SGHPL-5

trophoblast cell migration in humans. *Biol Reprod* 88: 54

Hemberger M (2002) The role of the X chromosome in mammalian extra embryonic

development. *Cytogenet Genome Res* 99: 210-217

Herman L, Todeschini AL, Veitia RA (2021) Forkhead transcription factors in health and

disease. *Trends Genet* 37: 460-475

Hosaka T, Biggs WH 3rd, Tieu D, Boyer AD, Varki NM, Cavenee WK, Arden KC (2004)

Disruption of forkhead transcription factor (FOXO) family members in mice reveals their

functional diversification. *Proc Natl Acad Sci USA* 101: 2975–2980

Huang C, Santofimia-Castaño P, Iovanna J (2021) NUPR1: a critical regulator of the

antioxidant system. *Cancers (Basel)* 13: 3670

Huang D, Sherman B, Lempicki R (2009) Systematic and integrative analysis of large gene

lists using DAVID bioinformatics resources. *Nat Protoc* 4: 44–57

Huang H, Tindall DJ (2011) Regulation of FOXO protein stability via ubiquitination and

proteasome degradation. *Biochim Biophys Acta* 1813: 1961–1964

Iqbal K, Pierce SH, Kozai K, Dhakal P, Scott RL, Roby KF, Vyhlidal CA, Soares MJ (2021) Evaluation of placentation and the role of the aryl hydrocarbon receptor pathway in a rat model of dioxin exposure. *Environ Health Perspect* 129: 117001.

Iwatsuki K, Oda M, Sun W, Tanaka S, Ogawa T, Shiota K (1998) Molecular cloning and characterization of a new member of the rat placental prolactin (PRL) family, PRL-like protein H. *Endocrinology* 139: 4976-4983

Iwatsuki K, Shinozaki M, Sun W, Yagi S, Tanaka S, Shiota K (2000) A novel secretory protein produced by rat spongiotrophoblast. *Biol Reprod* 62: 1352-1359

John RM (2017) Imprinted genes and the regulation of placental endocrine function: pregnancy and beyond. *Placenta* 56: 86-90

Kamei T, Jones SR, Chapman BM, McGonigle KL, Dai G, Soares MJ (2002) The phosphatidylinositol 3-kinase/Akt signaling pathway modulates the endocrine differentiation of trophoblast cells. *Mol Endocrinol* 16: 1469-1481

Kaneko T (2017) Genome editing of rat. *Methods Mol Biol* 1630: 101-108.

Kent LN, Konno T, Soares MJ (2010) Phosphatidylinositol 3 kinase modulation of trophoblast cell differentiation. *BMC Dev Biol* 10: 97

Kent LN, Ohboshi S, Soares MJ (2012) Akt1 and insulin-like growth factor 2 (Igf2) regulate placentation and fetal/postnatal development. *Int J Dev Biol* 56: 255-261

Kent LN, Rumi MA, Kubota K, Lee DS, Soares MJ (2011) FOSL1 is integral to

establishing the maternal-fetal interface. *Mol Cell Biol* 31: 4801-4813

Knipp GT, Audus KL, Soares MJ (1999) Nutrient transport across the placenta. *Adv Drug*

Deliv Rev 38: 41-58

Knöfler M, Haider S, Saleh L, Pollheimer J, Gamage TKJB, James J (2019) Human

placenta and trophoblast development: key molecular mechanisms and model systems. *Cell*

Mol Life Sci 76: 3479-3496

Kummer D, Ebnet K (2018) Junctional adhesion molecules (JAMs): the JAM-integrin

connection. *Cells* 7: 25

LaFleur DW, Nardelli B, Tsareva T, Mather D, Feng P, Semenuk M, Taylor K, Buergin M,

Chinchilla D, Roshke V, et al. (2001) Interferon-kappa, a novel type I interferon expressed in

human keratinocytes. *J Biol Chem* 276: 39765-39771

Lam EW, Brosens JJ, Gomes AR, Koo CY (2013) Forkhead box proteins: tuning forks for

transcriptional harmony. *Nat Rev Cancer* 13: 482-495

Liu W, Li Y, Luo B (2020) Current perspective on the regulation of FOXO4 and its role in

disease progression. *Cell Mol Life Sci* 77: 651-663

Maltepe E, Fisher SJ (2015) Placenta: the forgotten organ. *Annu Rev Cell Dev Biol* 31:

523-552

Manning BD, Cantley LC (2007) AKT/PKB signaling: navigating downstream. *Cell* 129:

1261-1274

Manning BD, Toker A (2017) AKT/PKB signaling: navigating the network. *Cell* 169:

381-405

Mashima R, Okuyama T (2015) The role of lipoxygenases in pathophysiology; new insights

and future perspectives. *Redox Biol* 6: 297-310

Morey R, Farah O, Kallol S, Requena DF, Meads M, Moretto-Zita M, Soncin F, Laurent

LC, Parast MM (2021) Transcriptomic drivers of differentiation, maturation, and polyploidy

in human extravillous trophoblast. *Front Cell Dev Biol* 9: 702046

Muto M, Chakraborty D, Varberg KM, Moreno-Irusta A, Iqbal K, Scott RL, McNally

RP, Choudhury RH, Aplin JD, Okae H, et al. (2021) Intersection of regulatory pathways

controlling hemostasis and hemochorial placentation. *Proc Natl Acad Sci USA* 118:

e2111267118

Nirgude S, Choudhary B (2021) Insights into the role of GPX3, a highly efficient plasma

antioxidant, in cancer. *Biochem Pharmacol* 184: 114365

Nteeba J, Varberg KM, Scott RL, Simon ME, Iqbal K, Soares MJ (2020) Poorly controlled

diabetes mellitus alters placental structure, efficiency, and plasticity. *BMJ Open Diabetes Res*

Care 8: e001243

Pijnenborg R, Robertson WB, Brosens I, Dixon G (1981) Trophoblast invasion and the

establishment of haemochorial placentation in man and laboratory animals. *Placenta* 2:

71-91

Pijnenborg R, Vercruysse L (2010) Animal models of deep trophoblast invasion. In

Placental Bed Disorders, Pijnenborg R, Brosens I, Romero R (eds) pp 127-139. Cambridge:

Cambridge University Press

Plaks V, Berkovitz E, Vandoorne K, Berkutzki T, Damari GM, Haffner R, Dekel N,

Hemmings BA, Neeman M, Harmelin A (2011) Survival and size are differentially regulated

by placental and fetal PKBalpha/AKT1 in mice. *Biol Reprod* 84: 537-545

Pollheimer J, Knöfler M (2005) Signalling pathways regulating the invasive differentiation of

human trophoblasts: a review. *Placenta* 26(Suppl A): S21-S30

Qayyum N, Haseeb M, Kim MS, Choi S (2021) Role of thioredoxin-interacting protein in

diseases and its therapeutic outlook. *Int J Mol Sci* 22: 2754

Qiu Q, Yang M, Tsang BK, Gruslin A (2004) Both mitogen-activated protein kinase and

phosphatidylinositol 3-kinase signalling are required in epidermal growth factor-induced

human trophoblast migration. *Mol Hum Reprod* 10: 677-684

Roberts RM, Green JA, Schulz LC (2016) The evolution of the placenta. *Reproduction* 152:

R179-R189

Satapathy S, Wilson MR (2021) The dual roles of clusterin in extracellular and intracellular

proteostasis. *Trends Biochem Sci* 46: 652-660

Schmitt-Ney M (2020). The FOXO's advantages of being a family: considerations on function

and evolution. *Cells* 9: 787

Sharma N, Kubaczka C, Kaiser S, Nettersheim D, Mughal SS, Riesenberg S, Hölzel M,

Winterhager E, Schorle H (2016) Tpbpa-Cre-mediated deletion of TFAP2C leads to

deregulation of Cdkn1a, Akt1 and the ERK pathway, causing placental growth arrest.

Development 143: 787-798

Shukla V, Soares MJ (2022) Modeling trophoblast cell-guided uterine spiral artery

transformation in the rat. *Int J Mol Sci* 23: 2947

Sies H, Cadenas E (1985) Oxidative stress: damage to intact cells and organs. *Philos Trans R*

Soc Lond B Biol Sci 311: 617-631

Soares MJ (2004) The prolactin and growth hormone families: pregnancy-specific

hormones/cytokines at the maternal-fetal interface. *Reprod Biol Endocrinol* 2: 51

Soares MJ, Chakraborty D, Rumi MAK, Konno T, Renaud SJ (2012) Rat placentation:

an experimental model for investigating the hemochorial maternal-fetal interface. *Placenta*

33: 233-243

Soares MJ, Chapman BM, Rasmussen CA, Dai G, Kamei T, Orwig KE (1996)
Differentiation of trophoblast endocrine cells. *Placenta* 17: 277-289

Soares MJ, Konno T, Alam SM (2007) The prolactin family: effectors of
pregnancy-dependent adaptations. *Trends Endocrinol Metab* 18: 114-121

Soares MJ, Varberg KM, Iqbal K (2018) Hemochorial placentation: development, function,
and adaptations. *Biol Reprod* 99: 196-211

Takagi N, Sasaki M (1975) Preferential inactivation of the paternally derived X chromosome
in the extraembryonic membranes of the mouse. *Nature* 256: 640-642

**van der Horst A, de Vries-Smits AM, Brenkman AB, van Triest MH, van den Broek N,
Colland F, Maurice MM, Burgering BM** (2006) FOXO4 transcriptional activity is regulated
by monoubiquitination and USP7/HAUSP. *Nat Cell Biol* 8: 1064–1073

**Varberg KM, Iqbal K, Muto M, Simon ME, Scott RL, Kozai K, Choudhury RH, Aplin
JD, Biswell R, Gibson M,** *et al.***,** (2021) ASCL2 reciprocally controls key trophoblast lineage
decisions during hemochorial placenta development. *Proc Natl Acad Sci USA* 118:
e2016517118

West JD, Papaioannou VE, Frels WI, Chapman VM (1978) Preferential expression of the
maternally derived X chromosome in extraembryonic tissues of the mouse. *Basic Life Sci* 12:
361-377

Wooding P & Burton G (2008) *Comparative Placentation*. Springer-Verlag, Heidelberg,
Germany

**Yang ZZ, Tschopp O, Hemmings-Mieszczak M, Feng J, Brodbeck D, Perentes E,
Hemmings BA** (2003) Protein kinase B alpha/Akt1 regulates placental development and fetal
growth. *J Biol Chem* 278: 32124-32131

FIGURE LEGENDS

Figure 1. In vivo genome editing of the rat *Akt1* locus. A) Schematic representation of the
rat *Akt1* gene (*Akt1*$^{+/+}$) and guide RNA target sites within Exons 4 and 7 (NM_033230.3). Red
bars beneath Exons 4 and 7 correspond to the 5' and 3' guide RNAs used in the genome
editing. **B)** The mutant *Akt1* allele (*Akt1*$^{-/-}$) possesses a 1,332 bp deletion. Parts of Exons 4
and 7 and all of Exons 5 and 6 are deleted, leading to a frameshift and premature Stop codon
in Exon 7. **C)** Amino acid sequences for AKT1$^{+/+}$ and AKT1$^{-/-}$. The red sequence corresponds
to the frameshift in Exon 7. The blue, red, and green highlighted amino acid sequence regions
correspond to the pleckstrin homology (**PH**), kinase, and regulatory domains, respectively. **D)**
Offspring were backcrossed to wild type rats, and heterozygous mutant rats were intercrossed
to generate homozygous mutants. Wild type (+/+), heterozygous (+/-), and homozygous
mutant (-/-) genotypes were detected by PCR. **E)** Western blot analysis of AKT1, pan-AKT,
phospho (p)-AKT (Ser^{473}) protein in *Akt1*$^{+/+}$ and *Akt1*$^{-/-}$ placentas at gd 18.5. GAPDH was
used as a loading control.

Figure 2. *Akt1*$^{-/-}$ placentas and fetuses are growth restricted, and *Akt1*$^{-/-}$ rats exhibit

postnatal growth restriction. Placentas (**A**) and fetuses (**B**) were dissected from $Akt1^{+/-}$

intercrosses at gd 18.5 and weighed; **C**, fetus/placenta ratio. Placentas were then separated

into junctional zone (**JZ**; **D**) and labyrinth zone (**LZ;E**) compartments, and weighed; **F**, JZ/JZ

ratio. Graphs represent means ± SEM. $Akt1^{+/+}$, n = 22 from 6 dams; $Akt1^{-/-}$, n = 17 from 6

dams. Asterisks denote statistical differences (**$P < 0.01$; ***$P < 0.001$) as determined by

Student's or Welch's t-test. Body weights of $Akt1^{+/+}$ and $Akt1^{-/-}$ pups were measured from two

to eight weeks after birth: males **(G)** and females **(H)**. Graphs represent means ± SEM. n =

13-23/group. Asterisks denote statistical differences (*$P < 0.05$; ***$P < 0.001$) as determined

by Student's or Welch's t-test. **I**) Vimentin immunostaining of gd 18.5 $Akt1^{+/+}$ and $Akt1^{-/-}$

placentation sites. The junctional zone (**JZ**) is negative for vimentin immunostaining,

whereas the uterine-placental interface (**UPI**) and labyrinth zone (**LZ**) stain positive for

vimentin. Scale bars=1000 μm

Figure 3. AKT1 regulates junctional zone and invasive trophoblast cell phenotypes.

A) Heat maps depicting differentially expressed genes between $Akt1^{+/+}$ and $Akt1^{-/-}$ junctional

zones. The heatmap color keys represent z-scores of TPM values. **B**) RT-qPCR validation of

RNA-seq results (n=6/group). **C**) Schematic representation of a late gestation placentation

site. The uterine-placental interface (UPI), site for intrauterine trophoblast invasion, is

highlighted in the boxed area. **D**) AKT1 deficiency affected intrauterine trophoblast cell

invasion. Trophoblast cells were immunostained for pan-cytokeratin. Representative images

are shown. The extent of intrauterine trophoblast invasion is demarcated using a green dashed

line. Scale bars =500 μm. The white dotted line represents the outer border of the uterus, and

the yellow dashed line represents the uterine border with the placenta. **E**) The area of

intrauterine trophoblast invasion is graphically depicted (n = 6/group). Graphs represent

means ± SEM. An asterisk denotes statistical difference (*$P < 0.05$) as determined by

Student's *t*-test. F) RT-qPCR measurements of *Krt7* and *Krt8* transcripts, signature markers

for invasive trophoblast cells, within dissected uterine-placental interface tissue specimens at

gd 18.5 (*Akt1*$^{+/+}$, n = 21; *Akt1*$^{-/-}$, n = 16). Graphs represent means ± SEM. Asterisks denote

statistical difference (*$P < 0.05$; **$P < 0.01$; ***$P < 0.001$) as determined by Student's or

Welch's *t*-test.

Figure 4. FOXO4 is a target of PI3K/AKT signaling. A) Expressions of transcripts for

several FOX transcription factors in the junctional zone. **B**) In situ localization of transcripts

for *Foxo4* with *Prl7b1* (invasive trophoblast-specific transcript) in the placentation site at gd

18.5 of rat pregnancy. Scale bars = 1000 μm. **C**) RT-qPCR measurements of *Foxo4*

transcripts in the uterine-placental interface, junctional and labyrinth zones during the second

half of gestation (n = 6-9/group). Graphs represent means ± SEM. Asterisks denote statistical

difference (**$P < 0.01$) as determined by Steel test. UPI: uterine-placental interface; JZ:

junctional zone; and LZ: labyrinth zone. **D**) Western blot analysis of phospho (p)-FOXO4

(Ser^{262}) and FOXO4 proteins in *Akt1*$^{+/+}$ and *Akt1*$^{-/-}$ placentas at gd 18.5. GAPDH was used as

a loading control. **E)** RT-qPCR measurements of *Foxo4* transcripts in the stem state and

following induction of differentiation (n = 4-6/group). Graphs represent means ± SEM.

Asterisks denote statistical difference (vs Stem, *$P < 0.05$) as determined by Dunnett's test.

Western blot analysis of phospho (p)-FOXO4 (Ser^{262}), FOXO4, p-AKT (Ser^{473}), and

pan-AKT proteins in the stem and differentiating (day 15 of differentiation, D15) states **(F)**,

and in the differentiating state (D15) following treated with vehicle (0.1% DMSO) or a

phosphatidylinositol 3-kinase (**PI3K**) inhibitor (LY294002, 10 μM) for 1 h **(G)**. GAPDH was

used as a loading control.

Figure 5. In vivo genome editing of the rat *Foxo4* locus. A) Schematic representation of

the rat *Foxo4* gene (*Foxo4*$^{Xm+}$) and guide RNA target sites within Exons 2 and 3

(NM_001106943.1). Red bars beneath Exons 2 and 3 correspond to the 5' and 3' guide RNAs

used in the genome editing. **B)** The mutant *Foxo4* allele (*Foxo4*$^{Xm-}$) possesses a 3,096 bp

deletion. Parts of Exons 2 and 3 are deleted, leading to a frameshift and premature Stop

codon in Exon 3. **C)** Amino acid sequences for FOXO4^{Xm+} and FOXO4^{Xm-}. The red sequence

corresponds to the frameshift in Exon 3. The green, red, dark blue, and light blue highlighted

amino acid sequence regions correspond to the forkhead winged-helix DNA-binding domain

(**FHD**), nuclear localization sequence (**NLS**), nuclear export sequence (**NES**), and

transactivation domain (**TAD**), respectively. **D)** Heterozygous mutant female rats were

crossed to wild type male rats to generate hemizygous null male rats. Wild type (+/+),

heterozygous (+/-), and hemizygous null (-/y) genotypes were detected by PCR. **E)** Western

blot analysis of FOXO4 protein in the junctional zone of *Foxo4*$^{Xm+}$ (X^{m+}Y and X^{m+}X^{p+}) and

Foxo4^{Xm-} (X^{m-}Y and X^{m-}X^{p+}) placentas at gd 18.5. GAPDH was used as a loading control.

Figure 6. *Foxo4* hemizygous null and *Foxo4* maternally inherited heterozygous

conceptuses exhibit placental overgrowth, and FOXO4 deficiency alters the

transcriptomes of the junctional zone. Placentas **(A)** and fetuses **(B)** were dissected from

Foxo4 heterozygous females mated with wild type males at gd 18.5 and weighed; **C)**

fetus/placenta ratio. Placentas were then separated into junctional zone (**JZ, D**) and labyrinth

zone (**LZ, E**) compartments, and weighed; **F**, JZ/LZ weight ratio. Graphs represent means ±

SEM. X^{m+}Y, n = 25; X^{m-}Y, n = 31; X^{m+}X^{p+}, n = 14; X^{m-}X^{p+}, n = 22 from 8 dams. Asterisks

denote statistical differences (***$P < 0.001$) as determined by Student's or Welch's *t*-test. **G)**

Vimentin immunostaining of gd 18.5 *Foxo4*$^{Xm+}$ and *Foxo4*^{Xm} placentation sites. The

junctional zone (**JZ**) is negative for vimentin immunostaining, whereas the uterine-placental

interface (**UPI**) and labyrinth zone (**LZ**) stain positive for vimentin. Scale bars=1000 μm. **H)**

Heat maps depicting differentially expressed genes in *Foxo4*$^{Xm+}$ versus *Foxo4*$^{Xm-}$junctional

zones. The heatmap color keys represent z-scores of TPM values. **I**) RT-qPCR validation of

RNA-seq results (n=6/group). Graphs represent means ± SEM. Asterisks denote statistical

difference (**P<0.01; ***P<0.001) as determined by Student's or Welch's t-test.

Figure 7. Reciprocal relationship between AKT1 and FOXO4 in the junctional zone.

Venn diagram and heatmaps representing overlap of differentially expressed genes inversely

regulated by AKT1 and FOXO4. The heatmap color keys represent z-scores of TPM values.

Figure 8. FOXO4 alters the rat TS cell. FOXO4 knockdown efficiency was validated by

RT-qPCR (**A**, n = 4/group) or western blot (**B**) analyses in rat TS cells at day 15 of

differentiation following transduction with lentivirus containing a control shRNA or one of

two independent *Foxo4*-specific shRNAs. Graphs represent means ± SEM. Asterisks denote

statistical difference (vs Control shRNA, *P < 0.05) as determined by Student's or Welch's

t-test. **C**) Heatmap depicting differentially expressed genes between control and *Foxo4*

shRNA-treated rat TS cells. **D**) RT-qPCR validation of RNA-seq results (Control shRNA, n =

4; *Foxo4* shRNA 1, n = 4; *Foxo4* shRNA 2, n = 4). Graphs represent means ± SEM. Asterisks

denote statistical difference (compared to Control shRNA, *P < 0.05; **P < 0.01; ***P <

0.001) as determined by Student's or Welch's *t*-test.

A

B

$Akt1^{-/-}$

Δ1332 bp

C

AKT1$^{+/+}$
MNDVA IVKEGWLHKRGEYIKTWRPRYFLLKNDGTFIGYKERP
QDVEQRESPLNNFSVAQCQLMKTERPRPNTFIIRCLQWTTVIE
RTFHVETPEEREEWTTAIQTVAD GLKRQEEETMDFRSGSPSD
NSGAEEMEVALAKPKHRVTMNE FEYLKLLGKGTFGKVILVKEK
ATGRYYAMKILKKEVIVAKDEVAHTLTENRVLQNSRHPFLTAL
KYSFQTHDRLCFVMEYANGGELFFHLSRERVFSEDRARFYGA
EIVSALDYLHSEKNVVYRDLKLENLMLDKDGHIKITDFGLCKEG
IKDGATMKTFCGTPEYLAPEVLEDNDYGRAVDWWGLGVVMY
EMMCGRLPFYNQDHEKLFELILMEEIRFPRTLGPEAKSLLSGL
LKKDPTQRLGGGSEDAKEIMQHRFF ANIVWQDVYEKKLSPPF
KPQ VTSETDTRYFDEEFTAQMITITPPDQDDSMECVDSERRP
HFPQFSY SASGTA

AKT1$^{-/-}$
MNDVA IVKEGWLHKRGEYIKTWRPRYFLLKNDGTFIGYKERP
QDVEQRESPLNNFSVAQCQLMKTERPRPNTFIIRCLQWTTVIE
RTFHVETPEEREEWTTAIQTVAD GLK QWGRALLPPVS Stop

D

PH domain
Kinase domain
Regulatory region

E

Labyrinth zone at gd 18.5

A

Foxo4^{Xm+}

FHD

NLS NES TAD

1 2 3 4

Fd→ Rv 1← Rv 2←

■ gRNA

gRNA binding site
CCAGATATACGAATGGATGGTCC

gRNA binding site
GTTCATCAAGGTACATAACGAGG

B

Foxo4^{Xm-}

Δ3096 bp

1 2 3 4

C

FOXO4^{Xm+}
MEPENKKSATEAAAIIDLDPDFEPQSRPRSCTWPLPRPELATE
PPEPSEVEPSLGQKVPTEGHSEPTLLPSRLPEPAGGPQPEILG
AVTGPRKGGSR RNAWGNQSYAELISQAIESAPEKRLTLAQIYE
WMVRTVPYFKDKGDSNSSAGWKNSIRHNLSLHSKFIKVHNEA
TGKSSWWMLNPD GG KGGKAPRRRAASMDSSSKLLRGRSKG
PKKK PSVLPAPPEGATPRSPLGHFAKWSSSPCPRNREEADV
WTTFRPRSSSNASTVSTRLSPMRPESEVLAEEEMPASASSYA
GGVPPTLS EDLELLDGLNL ASPHSLLSRSNLSSFSLQHPGLAG
PLHSYGASLFGPIDGSLSAGEGCFSSSQSLEALLTSDTPPPPA
DVLMTQVDPILSQAPTLLLLGGMPSSSKLATGVSLCPTPLEGP
GPSNLVPTLSVMAPPPVMPGAPIPKVLGTPVLASPTEDFSHDR
MPQDLDLDMYMENLECDMDNIISDLMDGEGLDFNFEP DP

FOXO4^{Xm-}
MEPENKKSATEAAAIIDLDPDFEPQSRPRSCTWPLPRPELATE
PPEPSEVEPSLGQKVPTEGHSEPTLLPSRLPEPAGGPQPEILG
AVTGPRKGGSR RNAWGNQSYAELISQAIESAPEKRLTLAQ RP
PARAL GGC

FHD
NLS
NES
TAD

D

bp Ladder X^m+X^p+ X^m-X^p+ X^m-Y

800—

◄ WT

400—

◄ Mutant

E

Foxo4^{Xm+} Foxo4^{Xm-}

kDa X^m+Y X^m+X^p+ X^m-Y X^m-X^p+

70—

60—

50—

FOXO4

40—

◄ GAPDH

35—

Junctional zone at gd 18.5