Neural responses to instructed positive couple interaction: An fMRI study on compliment sharing

Authors:

Monika Eckstein[1#*], Gabriela Stößel[2#], Martin Fungisai Gerchen[2,3,4#], Edda Bilek[5], Peter Kirsch[2,3,4†], Beate Ditzen[1,6†*]

Institutions:

[1]Institute of Medical Psychology, Center for Psychosocial Medicine, Heidelberg University Hospital, Heidelberg, Germany [2]Department of Clinical Psychology, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, Germany, [3]Bernstein Center for Computational Neuroscience Heidelberg/Mannheim, Mannheim, Germany, [4]Department of Psychology, Heidelberg University, Heidelberg, Germany, [5]Department of Psychiatry and Psychotherapy, Central Institute of Mental Health, Mannheim, Germany, [6]Heidelberg University, Heidelberg, Germany

*Corresponding authors: Monika Eckstein, Beate Ditzen

# equally contributing first authors

† equally contributing senior authors

Word Count:5038

Keywords: Social interaction, brain imaging, ventral striatum, couple relationships, compliment reward

Abstract

Love is probably the most fascinating feeling that a person ever experiences. However, little is known about what is happening in the brains of a romantic couple –the central and most salient relationship during adult age– while they are particularly tender and exchanging loving words with one another.

To gain insight into nearly natural couple interaction, we collected data from N=84 individuals (including N=43 heterosexual couples) simultaneously in two functional magnetic resonance imaging scanners, while they sent and received compliments, i.e. short messages about what they liked about each other and their relationship. Activation patterns during compliment sharing in the individuals revealed a broad pattern of activated brain areas known to be involved in empathy and reward processing. Notably, the ventral striatum, including parts of the putamen, was activated particularly when selecting messages for the partner. This provides initial evidence that giving a verbal treat to a romantic partner seems to involve neural reward circuitry in the basal ganglia.

These results can have important implications for the neurobiological mechanisms protecting and stabilizing romantic relationships, which build a highly relevant aspect of human life and health.

Introduction

In almost all human cultures, romantic love is viewed as a central concept to giving meaning and joy to a person's life. Social identity theory states that individuals derive parts of their identity from belonging to a group, a family, or a romantic relationship (Scheepers and Ellemers, 2019) and such social identification is related to less harmful stress as mediated by social support (Haslam *et al.*, 2005). Specifically being in a functional couple relationship is even linked to better health and longer lives (Braithwaite and Holt-Lunstad, 2017).

The health-related impact of couple relationships is very likely centrally mediated with interacting neural networks and structures, such as the limbic reward system and its neurotransmitters. An interaction of the neuromodulator oxytocin and the neurotransmitter serotonin in the nucleus accumbens (Dölen *et al.*, 2013) has been shown to mediate social reward. Oxytocin interacting with dopamine has been suggested to contribute to the formation and maintenance of social bonds in animals (Bosch and Young, 2018) and in humans for instance via a positive evaluation of the own relationship (Scheele *et al.*, 2013; Aguilar-Raab *et al.*, 2019) .

As parts of the dopaminergic reward system, the nucleus accumbens, together with putamen and ventral tegmental area (VTA) among others, are involved in the initiation of joyful behaviors and feelings in general, but especially social reinforcements (Izuma *et al.*, 2008; Dölen *et al.*, 2013). Functional activation of the dopaminergic reward systems might thus be one (of several) underlying mechanism supporting initializing and maintaining human couple relationships (Bartels & Zeki, 2004) and is therefore in the focus of the present study. In previous research, interacting with the partner or observing a partner picture was associated with elevated activation in the VTA, hippocampus, insula (Bartels and Zeki, 2004), anterior

cingulate cortex (ACC) (Aron *et al.*, 2005) posterior superior temporal sulcus (pSTS), and anterior temporal lobe (ATP)(Van der Gaag *et al.*, 2007).

One precondition for functional social interaction and for romantic couple relationships in particular is a theory of mind (ToM), the ability to infer the status of knowledge of another person. ToM is related to activation of the superior temporal brain, temporal, and frontal areas (Dodell-Feder *et al.*, 2015), while actual empathy recruits the anterior insula (Kennedy and Adolphs, 2012; Thornton *et al.*, 2019). During empathy-related processes, the accumbens is also interacting with the ACC (Smith *et al.*, 2021). In addition, the mirror neuron system, which comprises parietal and frontolateral brain areas, is involved in social perception and action (Mier *et al.*, 2010).

Furthermore, social integration and the perception of belonging increase positive affect and self-esteem (Ellemers *et al.*, 1999). Presumably, positive feedback acts as an indicator of social integration. For romantic couples, a constructive way of communication has been shown to be related to better relationship satisfaction and even to buffer a lack of sexual satisfaction (Litzinger and Gordon, 2005) and communicating compliments in everyday life has been linked to better relationship satisfaction with a stronger sensitivity towards compliments specifically in women (Doohan and Manusov, 2004). Imaging studies have shown that receiving compliments from a stranger or from one's own mother involved the dorsolateral prefrontal cortex (DLPFC) (Hooley *et al.*, 2005), ACC, and temporal areas (Miedl *et al.*, 2016). Based on this, receiving compliments from the partner can be considered highly relevant to evaluating the social self, the level of integration and affection and, thereby, act in a particularly rewarding and health-beneficial way. To investigate the neural responses to tender partner compliments, we have adapted the previously established standard instructed partnership appreciation task (Pfeifer *et al.*, 2020; Warth *et al.*, 2020) for a functional imaging (fMRI) paradigm. We compared compliments from the partner to "self compliments" (i.e.,

attributes that the participants defined about themselves), since the mental reflection of positive attributes per se could improve mood (Nicolson *et al.*, 2020) by activating reward-related brain areas (Izuma *et al.*, 2008; Frewen *et al.*, 2020).

For general compliment processing we expected that receiving compliments from the partner would result in elevated activation in a broad network including VTA, hippocampus, insula, ACC, and pSTS, (Van der Gaag *et al.*, 2007). In addition, reward-related task phases as well as phases of reward anticipation (Filimon *et al.*, 2020) should be related to activation in the dopaminergic system: the ventral striatum including the nucleus accumbens. While a participant is actively choosing a compliment, we expected activity known for reading and decision making, and when a participant is sending the compliment, the areas relevant for ToM should be activated, along with mirror neuron areas when they are observing partners' reactions. These activation patterns should become evident using whole brain approaches.

Methods

Participants

Eighty six heterosexual participants (from 43 romantic couples) who were in love and exclusively dating for at least six months were recruited in the Rhine-Neckar metropolitan area, Germany; see Table 1 for sample characteristics. In addition to sociodemographic data, participants provided information on their relationship quality (Partnership Questionnaire, PFB (Hahlweg, 1979), including the scales for tenderness, conflict, and joint interests. Particularly happy couples (reporting at least 5 on a 6-point single-item rating scale on general relationship satisfaction) were included in the study. All participants were eligible for MRI scanning, right-handed, without history of mental disorders and knew sufficient German language to fully understand all instructions. Couples provided written informed consent and were reimbursed 80€ per couple for their participation. The study was conducted in

accordance with the Declaration of Helsinki and approved by the Ethics Committee of

Heidelberg University Medical Faculty (#2011-222N-MA).

Paradigms

In an interview session with the individual participants prior to the MRI session, all

participants were handed a list of 23 areas of individual traits and relationship aspects, based

on factors of the PFB (e.g., trust, humor, intimacy). Based on these areas, participants were

asked to generate up to 18 short positive messages (compliments) about their partner for use

in the upcoming experiment. In addition, participants created up to 18 compliments about

themselves to be viewed as control stimuli. The compliments were kept confidential until the

MRI session. Non-German native speaking couples were allowed to provide compliments in

their native language. The paradigm consisted of 15 trials per condition (receiving, sending,

self compliment). In the send and receive compliment condition, each trial consisted of two

phases, lasting 10s each. In the first phase, the sender chose one of four compliments shown

on his/her screen, and the receiver waited for the partner to select the message. In the second

phase of the trial, the compliment was revealed to both partners. In the self compliment

paradigm, running on both scanners simultaneously, trials consisted of two phases as well: in

the first phase, the text 'Please wait, computer is choosing your compliment' was displayed to

both participants, in the second one, the text appeared: 'Computer has chosen:

compliment_text'. Both phases of each trial were jittered on average by 775ms, one whole

trial lasted 32.5s. All texts were presented on the left-hand side of the screen. On the right-

hand side, a live video of the partner taken with a wide-angle camera (MRC Systems GmbH,

Heidelberg, Germany) in infrared light was shown continually during all paradigms in order

to keep general effects of partner contact constant over conditions. The participants were

randomly assigned to one of the two scanners. The order of which partner sent first, as well as

the assignment of sexes to the scanner and the orders to the scanner were balanced. The temporal order of paradigms (partner vs. self) and the initial sender-sex-scanner matching was randomized and balanced across the sample. However, the first send/receive condition was always followed by the complimentary send/receive condition. The task followed an anatomical measurement and a joint attention paradigm (Bilek *et al.*, 2015).

A follow-up questionnaire directly after the fMRI session assessed the participants' overall evaluation of sending and receiving partner-compliments with: "How much did you enjoy sending and receiving the messages?" and "How much did you enjoy reading your own positive attributes?" on a 9-point scale from "not at all" to "very much".

Data acquisition

Data was acquired with two synchronized 3 Tesla Siemens Tim TRIO scanners, where one scanner was triggered by the other one. Twelve-channel head coils were used. A T2* gradient echo-planar imaging sequence was applied with the following parameters: 28 axial slices, with transversal orientation, oriented first to AC/PC line and then flipped by -25°, 4 mm slice thickness, 1 mm gap, field of view 192 mm, voxel size 3x3x4 mm3. Repetition time (TR) was 1.55s with sampling delay of 10ms, and 1.54s. Echo time was 30ms, flip angle 73°. Slices were acquired in descending order, with A/P phase encoding direction. The GRAPPA method with an acceleration factor of 2 was used. A total of 327 (triggering)/ 324 (triggered scanner) scans were collected per condition. The first 7 (triggering)/4 (triggered scanner) scans were discarded during conversion of DICOM-files into 4d niftis by MRIConvert (version. 2.0 rev. 216) to account for saturation effects, resulting in 320 scans available for analysis per condition.

A high-resolution (voxel size 1x1x1mm) T1 anatomical scan was acquired for individual anatomical registration purposes.

Data analyses and preprocessing

fMRI data were analyzed using SPM12 (v771). The anatomical image was segmented and normalized to the SPM12 TPM MNI template. Preprocessing of the functional data involved slice-time correction, realignment to the mean image, and co-registration of the functional images (mean and others) to the anatomical image. The co-registered functional data was normalized to MNI space, resampled to $3mm^3$ voxels, and smoothed with a Gaussian kernel with full-width-at-half-maximum FWHM=8x8x8 mm. Volumes affected by small movement artefacts were identified with the ART toolbox (http://www.nitrc.org/projects/artifact_detect ; parameters: framewise displacement >0.5mm, image intensity change z>4, and exclusion criterion for a measurement: >25% affected volumes).

Of the original 86 fMRI measurements, we had to exclude nine from the activation analysis of the send-paradigm (resulting in N=77) and seven measurements from the activation analysis of the receive-paradigm (resulting in N=79) due to excessive head motion, technical problems, or aborted measurements due to time constraints. In total, this resulted in 14 participants having to be excluded from the comparison of the receive-paradigm with the self compliment-paradigm (N=72).

First, we analyzed the task-related activation in the individuals' brains by means of general linear modeling. A first-level model with three sessions for the three separate conditions of the experiment was set up to allow for both within-session and across-session contrasts. With the conditions, the individual phases (waiting for and receiving a compliment, as well as selecting and observing shared compliments) were modeled as blocks. Signals from cerebrospinal fluid and white matter, 24 movement parameters (six standard parameters, their backward derivatives, and their squared versions), and ART dummy regressors were included

as nuisance regressors. A high-pass filter with a frequency cutoff of 128s was applied, as well first-degree autoregression.

In the group analyses, age, sex and scanner were included as covariates. Analyses were conducted using one-sample t-tests over the respective contrasts. Contrasts of interests were [Receiving > Waiting] within blocks (partner compliment and self compliment) and [Receiving > Waiting] compared between blocks (partner compliment and self compliment) as well as a contrast between the active block [Choosing compliment > Observing sent compliment] and the passive block [Receiving > Waiting]. All activation results are reported with p<0.05 whole-brain FWE-corrected significance. Beta estimates were additionally extracted, only for visualization of the activity of the ventral striatum (anatomical region-of-interest from the AAL-90 atlas) during conditions (Fig. 4).

Questionnaire data was analyzed using SPSS 27 (IBM).

Results

Activation of individuals when receiving a compliment

When participants were passively receiving compliments (both from the partner and self compliments) as compared to the waiting phases (within blocks), increased activation in a broad network of IFG, DLPFC, VMPFC, midbrain-structures and temporal gyri was observed, see Tables 2 and 3 and Fig. 1a and b.

a [Receiving partner compliment > Waiting for partner compliment]

b [Receiving self compliment > Waiting for self compliment]

L   y = -16 R   x = 5

Fig.1. Higher activation during receiving compliments than during waiting, 1a receiving partner compliments 1b receiving self compliments. All figures p<.05, whole brain FWE corrected (x = 5 y= -16), T-scale applied for both panels.

Activations of individuals when receiving partner vs. self compliments

Contrasting receiving compliments from the partner with self compliments (between the two passive blocks) showed increased VMPFC, ACC, and IFG activity for receiving self compliments (Table 4, Fig 2a) and higher insula, temporal, and amygdala activity when receiving partner compliments. (Table 5, Fig 2b).

a [Receiving self compliment > Waiting for self compliment] >
[Receiving partner compliment > Waiting for partner compliment]

L   y = 24   R   x = -4

b [Receiving partner compliment > Waiting for partner compliment] >
[Receiving self compliment > Waiting for self compliment].

L   y = -10   R   x = 8

Fig.2. Higher brain activation during partner- than during self compliments, 2a receiving self compliments (receive self-compliments (receive > wait) > receive partner-compliments (receive > wait)) (x =-4, y= 24), 2b receive partner compliments (partner-compliments (receive >wait) > self compliments (receive > wait)) (x= 8, y=-10)

Activation of individuals when sending compliments

Comparing brain responses during the blocks of sending and receiving of partner compliments, we found that receiving involves larger insula, and hippocampus activity (Table 6, Fig 3a) but selecting/sending compliments for the partner involved an even broader limbic and reward network, including a large cluster around the ventral striatum, TPJ and the cingulate gyrus (Table 7, Fig 3b and for beta estimates for the activity of the ventral striatum see Fig 4).

Fig. 3. Activation during receiving compared to compliment sending. 3a receiving partner compliment (receive > wait) > sending (choose > observe). 3b sending (choose > observe) > receive partner-compliment (receive > wait), both x 7 = y = 3, t-scale applies to both panels

Fig. 4. Beta estimates on ventral striatum activation during the experimental phases of sending and receiving partner compliments; white dots indicate means, black bars indicate SEM.

We found no sex differences in any comparison.

Taken together, these results suggest similar activation patterns for self compliment and partner compliment in the paradigm: elevated activation in DL/VMPFC, precuneus, and

temporal gyrus when receiving compliments. DLPFC and posterior cingulate are especially sensitive to receiving partner compliments, while temporal lobe and amygdala respond to the anticipation of partner compliments. Interestingly, choosing and sending compliments yielded the strongest activation patterns in the limbic, mentalizing (ToM), and reward systems.

Results of questionnaire data

A Wilcoxon test (with N = 86, who completed the questionnaires) suggest that participants reported significantly (z = -7.223, p <. 001) more subjective joy during the partner compliment phase (Median = 9, Range 5 – 9) than during the self-compliment phase (Median 6, Range 2 – 9).

Discussion

For most adult humans, couple relationships are *the* most relevant social relationship, and interacting with the partner modulates momentary affect and long-term health-related outcomes (Braithwaite and Holt-Lunstad, 2017). Exchanging praise and compliments are one element of positive couple interaction and specific compliments in the relationship are assumed to increase social identity (Ellemers, Kortekaas & Ouwerkerk, 1999). The rationale of the present study was to investigate the neural responses when sending and receiving such compliments, as well as receiving self compliments. In summary, we found that both receiving compliments from the partner and self-generated positive attributes activated the salience and limbic networks, as well as the mirror neuron system, as hypothesized. Differential effects occurred especially during the anticipation of the response to a compliment.

The complex activation pattern to receiving compliments corresponded to the activation seen in previous research investigating the reading of emotionally loaded content (Hsu *et al.*, 2014). Prefrontal and temporal areas, as well as the insula, were involved in both receiving partner compliments and self compliments. This is in line with the notion that the general processing of self-referential information involves the reward circuitry (Frewen *et al.*, 2020) and the dorsal striatum in particular (as part of the nigrostriatal pathway) is involved in comparing predicted and received reward (Oyama *et al.*, 2010).

Amygdaloid responses during the anticipation of partner compliments, relate to the 'emotional' salience network, but also with social reward (Chan *et al.*, 2018). Receiving partner compliments included activation of ACC and temporal gyri. Such activation patterns are part of the "social brain" (Kennedy and Adolphs, 2012) and are involved in successful

communication and mentalizing (Van Overwalle and Baetens, 2009; Laurita *et al.*, 2017). Here, they might serve as an indicator of ToM and the sender's mental engagement with choosing a particular compliment.

The compliment choosing phase was associated with complex activation patterns in the senders' brains which included the dopaminergic reward system. The ventral striatum and neural midline structures showed the strongest activation when choosing a compliment as compared to the other conditions (see Fig 4). While this was not hypothesized, these results are well in line with previous reports indicating that emotion sharing might be rewarding (Wagner *et al.*, 2014), and can be associated with striatal activation during the anticipation of reward (Filimon *et al.*, 2020). Since there was no other experimental condition including non-emotional decision making to compare this data with, we have to interpret this finding with caution, though. Other examples of rewarding anticipation of prosociality include supporting financially family members, which elicits activation in the mesolimbic dopaminergic system (Telzer *et al.*, 2010), as well as deciding to donate to charities, which recruited the ventral and dorsal striatum and VTA (Moll *et al.*, 2006). Similarly, Harbaugh *et al.* (2007) found that both mandatory and voluntary contributions to charities recruited the same areas. Finally, Izuma *et al.* (2010) reported that ventral striatum activity to charitable donations increased in the presence of others, suggesting that this region may be particularly sensitive to social rewards. Our present results add to this line of literature by showing for the first time the differential contributions of dorsal striatum to receiving a treat oneself and of ventral striatum to selecting a treat for someone else during live social interaction.

Our data imply that throughout all conditions, the senders paid close attention to the reaction of their partners during compliment sharing: Activation in oculo-, pre- and motor areas, as well as areas associated with showing emotional, mostly happy, faces such as pSTS and

DMPFC suggest involvement of the emotionally 'extended mirror neuron network' (Van der Gaag *et al.*, 2007). Positive affect and frontal activity during emotional partner interaction were also reported in a recent study using EEG (Packheiser *et al.*, 2021).

In summary, by using a somewhat naturalistic interaction paradigm, the present study design builds on previous research on reward-related brain activation in romantic couples such seeing as pictures from the partner (Acevedo *et al.*, 2012) and extends existing data to a more dynamic couple interaction. To our knowledge, this work is the first to investigate the neural underpinnings of positive emotional interaction between romantic couples using individually meaningful attributes characterizing the relationship and the participants involved, namely self-generated compliments.

The specific areas found to be involved in couple's compliment sharing are known for social cognition processes, social reward processing, ToM, and facial mimicry (Jabbi and Keysers, 2008; Kennedy and Adolphs, 2012). The involvement of the dopaminergic reward system in particular might serve as an important neurobiological mechanism underlying the ever rewarding aspects of lasting couple relationships. Interestingly, these brain areas are also involved in the action of neuropeptides promoting social behavior, such as oxytocin (Riem *et al.*, 2012; Kreuder *et al.*, 2018). Oxytocin has been shown to interact with the reward system, for example when study participants observed the face of their romantic partner (Scheele *et al.*, 2013), and also to influence the appraisal of the relationship (Aguilar-Raab *et al.*, 2019). Furthermore, oxytocin is known to promote health-beneficial effects such as regulation of the stress axes during couple interaction (Ditzen *et al.*, 2009; Zietlow *et al.*, 2018). Therefore, the neural networks reported here and the role of oxytocin might provide a potential neurobiological pathway underlying the association of couple relationships and health.

Our study has strengths but also some limitations. Investigating heterosexual romantic couples only and having them name, choose, and send the compliments helped create an individualized interaction scenario. The paradigm comprised receiving unknown compliments from the partner and known self-compliments while always seeing the partner via video transmission as part of a naturalistic social exchange. Therefore, we can neither rule out that the found differences between partner- and self-compliment are due to novelty nor that some kind of interaction has taken place during the self-compliment phase via facial expressions. Other aspects that differed between task phases were the active or passive role of leading the interaction or making decisions in general.

The selected heterosexual monogamous sample allows no extrapolation to unacquainted individuals, platonic friend dyads or same-sex couples, though. Furthermore, the sample consisted of healthy young couples reporting high relationship satisfaction only. Given inconsistent effects of instructed partnership appreciation in clinical samples (Warth *et al.*, 2020) or couples in therapy (Aguilar-Raab *et al.*, 2018), we cannot extrapolate our findings to marital problems or patient populations (see for instance a study in couple with substance abuse by Flanagan *et al.*, 2018). On the other hand, our findings may still be applicable for some cultures or couple circumstances, since our participants came from Europe and North Africa (15 different nations and 12 mother tongues) therefore generalizability to those parts of the world is given and the individualized compliments have accounted for potential differences. Future studies could systematically investigate cultures and contexts, clinical samples, and couples in the LGBTQIA+ spectrum. We assume similar basic neural effects in all couples though.

In conclusion, our data show substantial involvement of limbic structures during instructed yet individualized couples compliment sharing. The involvement of dopaminergic areas, is evident

not only when receiving compliments but is strongest in the ventral striatum when selecting compliments for the partner. This suggests a role of neural reward processes when giving a treat to the loved one - which might contribute to the maintenance of lasting relationships beyond the mere receipt of affection and support.

FIGURE LEGENDS

Fig. 1. Higher activation during receiving compliments than during waiting, 1a receiving partner compliments 1b receiving self compliments. All figures p<.05, whole brain FWE corrected (x = 5 y= -16), T-scale applied for both panels

Fig. 2.

Fig.2. Higher brain activation during partner- than during self compliments, 2a receiving self compliments (receive self-compliments (receive > wait) > receive partner-compliments (receive > wait)) (x =-4, y= 24), 2b receive partner compliments (partner-compliments (receive >wait) > self compliments (receive > wait)) (x= 8, y=-10)

Fig. 3. Activation during receiving compared to compliment sending. 3a receiving partner compliment (receive > wait) > sending (choose > observe). 3b sending (choose > observe) > receive partner-compliment (receive > wait), both x 7 = y = 3, t-scale applies to both panels.

Fig. 4. Beta estimates on ventral striatum activation during the experimental phases of sending and receiving partner compliments; white dots indicate means, black bars indicate SEM.

Table 1

Sample characteristics. Available data presented for participants included for the corresponding fMRI analyses (number of participants in brackets):

	Sending (77), questionnaires (72)				Receiving partner compliment (79), questionnaires (74)				Receiving self compliments (77) questionnaires (72)			
	mean	std	min	max	mean	std	min	max	mean	std	min	max
Age (years)	24.4	2.8	19	32	24.2	3.0	19	32	24.4	3.1	19	32
Education (years)	12.5	1.6	3	15	12.5	1.6	3	15	12.5	1.6	3	15
Relationship duration (years)	3.1	2.8	0.5	12	3.1	2.7	0.5	12	3.1	2.80	0.5	12

Table 2

Brain responses to (receiving partner compliments > waiting for partner compliments)

peak	peak				
p(FWE-corr)	T	x	y	z	Automated anatomical labeling atlas (Tzourio-Mazoyer *et al.*, 2002)
< 0.001	16.79296303	3	14	65	Superior frontal gyrus/Supplementary motor area
< 0.001	12.37275314	-6	5	77	Superior frontal gyrus
< 0.001	11.77529621	6	17	38	Middle cingulate gyrus
< 0.001	15.70023251	-39	-70	-22	Cerebellum
< 0.001	15.21589851	-42	17	-28	Superior temporal gyrus
< 0.001	15.20484543	-45	17	-7	Inferior frontal gyrus
0.0049	5.602838039	66	-40	26	Inferior partietal lobule/Supramarginal gyrus
0.0065	5.522605896	66	-58	11	Superior temporal gyrus
0.0447	4.955034733	-15	-25	-31	Cerebellum

Whole brain analyses; FWE corr, familywise error correction (applies to all tables)

Table 3

Brain responses to (receiving self compliments > waiting for self compliments)

peak	peak				
p(FWE-corr)	T	x	y	z	
< 0.001	17.5258503	-42	26	-4	Inferior frontal gyrus

p(FWE-corr)	T	x	y	z	
< 0.001	17.06056023	-3	20	59	Superior frontal gyrus/Supplementary motor area
< 0.001	15.37069511	48	26	-1	Inferior frontal gyrus
< 0.001	6.319052696	30	-13	-31	Parahippocampus
0.0042	5.719283581	33	-52	53	Superior parietal gyrus
0.0050	5.673546314	-21	-76	23	Precuneus
0.0054	5.654490948	-3	-52	-16	Vermis/Cerebellum
0.0212	5.255188942	30	-22	35	Postcentral gyrus
0.0353	5.101302147	-9	-16	80	Superior frontal gyrus

Table 4

Brain responses to (receiving self compliment > waiting for self compliment) > (receiving partner compliment > waiting for partner compliment)

peak	peak				
p(FWE-corr)	T	x	y	z	
< 0.001	7.01081753	-9	26	-10	Anterior cingulate
< 0.001	6.95619154	-6	35	-22	Rectal gyrus
< 0.001	6.37565708	-21	41	-13	Middle frontal gyrus
< 0.001	6.97407198	42	50	-4	Middle frontal gyrus
0.0070	5.5989995	24	53	-1	Superior frontal gyrus
0.0002	6.51676655	54	-52	50	Inferior partietal lobule
0.0050	5.69650269	42	-64	53	Angular gyrus

0.0026	5.87921333	3	-28	59	Medial frontal gyrus
0.01062	5.47870684	-9	-31	65	Medial frontal gyrus
0.0064	5.625525	-36	-22	20	Insula
0.0070	5.59888983	36	-19	17	Insula
0.0098	5.50063801	42	23	47	Middle frontal gyrus
0.0122	5.435884	-9	35	-1	Anterior cingulate
0.0189	5.30584526	24	41	-13	Middle frontal gyrus
0.0251	5.21994162	-42	-58	56	Partietal inferior gyrus
0.03185	5.14777374	-54	-55	41	Partietal inferior gyrus

Table 5

Brain responses to (receiving partner compliment > waiting for partner compliment) > (Receiving self compliment > waiting for self compliment)

peak	peak				
p(FWE-corr)	T	x	y	z	
< 0.001	7.21621704	-30	-64	-22	Cerebellum
< 0.001	6.71068478	-42	-61	-25	Cerebellum
< 0.001	6.43157673	-12	-67	-16	Cerebellum
< 0.001	7.18531132	-48	-13	41	Pre/postcentral gyrus
0.0072	5.59245872	-48	-10	59	Precentral gyrus
< 0.001	6.9873867	42	17	-28	Superior temporal gyrus
0.0012	6.0868845	48	8	-7	Insula

0.0140	5.39685678	36	-1	-25	Amygdala
< 0.001	6.43445444	-39	11	-28	Superior temporal gyrus
0.0011	6.10743856	45	-10	41	Precentral gyrus
0.0014	6.04848385	57	-4	44	Precentral gyrus
0.0017	5.99132061	48	-1	59	Frontal middle gyrus
0.0054	5.67191124	27	-7	-13	Amygdala
0.0073	5.58856153	-6	-88	-10	Calcarine/Lingual gyrus
0.0093	5.51773834	3	2	68	Superior frontal gyrus/Supplementary motor area
0.0100	5.4936924	-15	-1	80	Superior frontal gyrus
0.012	5.44215488	9	-13	-13	Hippocampus
0.0125	5.42972374	-6	-82	17	Cuneus
0.0197	5.29384518	3	-85	38	Precuneus
0.0242	5.23110056	-30	-13	-13	Hippocampus
0.0357	5.11234665	9	14	38	Middle cingulate
0.0458	5.03419256	-30	-61	-49	Cerebellum
0.0469	5.02685928	15	-19	-13	Hippocampus

Table 6

Brain responses to (receiving partner's compliment > waiting for partner's compliment) > (choosing compliment for partner > observing sending)

peak	peak				
p(FWE-corr)	T	x	y	z	
< 0.001	12.1697969	21	-91	-34	Cerebellum

< 0.001	6.47661924	3	-88	-19	Cerebellum
< 0.001	11.8543653	-33	-88	-31	Superior/Middle occiptal gyrus
< 0.001	11.2321196	-21	-79	-34	Cerebellum
< 0.001	10.5057125	0	-1	20	Corpus callusum
< 0.001	10.3181181	-12	-4	29	Middle cingulate
< 0.001	8.00352383	-18	-22	29	Nucleus caudatus/Middle cingulate
< 0.001	9.35983181	-54	-67	41	Angular gyrus
< 0.001	8.50362015	-60	-58	29	Supramarginal gyrus
< 0.001	8.44472694	-60	-46	50	Supramarginal gyrus
< 0.001	8.84178734	57	-61	11	Superior temporal gyrus
< 0.001	6.97519064	66	-55	20	Superior temporal gyrus
0.0054	5.63413525	69	-40	29	Supramarginal gyrus
< 0.001	8.6882925	6	56	38	Superior frontal gyrus
< 0.001	8.64810467	6	26	68	Superior frontal gyrus
< 0.001	7.82007265	-6	50	26	Medial Frontal gyrus
< 0.001	7.84036493	-33	-52	2	Lingual gyrus
0.0014	6.01714706	-24	-37	14	Hippocampus
0.0025	5.84849262	-18	-43	14	White matter
< 0.001	7.54454184	27	-7	-7	Globus pallidus/Lentiform nucleus
< 0.001	7.52939987	48	2	-28	Middle temporal gyrus
< 0.001	7.01410103	48	11	-28	Temporal pole
< 0.001	7.29358339	-45	20	53	Middle frontal gyrus
< 0.001	7.24051571	-42	11	-34	Middle frontal gyrus
< 0.001	7.18364382	45	-10	35	Precentral gyrus

< 0.001	6.34821653	33	-46	2	Lingual gyrus
< 0.001	6.16943264	3	-55	32	Middle cingulate/Precuneus
0.0010	6.10089874	-57	-28	-19	Temporal inferior gyrus
0.0023	5.8795042	-63	-19	-19	Temporal middle gyrus
0.0012	6.0625968	0	-28	17	Thalamus
0.0020	5.90837193	57	-1	11	Insula
0.0028	5.81837273	-42	-13	35	Postcentral/Precentral gyrus
0.0042	5.7014122	-36	-13	26	Insula
0.0053	5.63772678	42	5	-4	Insula
0.0069	5.56073284	-27	-1	-22	Amygdala
0.0192	5.25661325	54	-25	-13	Temporal middle
0.0212	5.22633171	57	11	2	Insula
0.0247	5.1791029	51	23	5	Inferior frontal gyrus
0.0339	5.08146048	57	20	-28	Middle temporal pole
0.0399	5.03027153	-33	-7	-19	Parahippocampus
0.04829	4.96986198	-54	-61	-28	Cerebellum

Table 7

Brain responses to (choosing compliment for partner > observing sending) > (receiving partner's compliment > waiting for partner's compliment)

peak	Peak				
p(FWE-corr)	T	x	y	z	
< 0.001	16.525074	-36	-13	59	Precentral gyrus
< 0.001	15.8985653	-39	-31	53	Poscentral gyurs

< 0.001	15.8449984	18	-67	56	Superioral parietal gyrus
< 0.001	9.79416847	45	8	26	Inferior frontal gyrus
< 0.001	7.32125902	57	-46	-16	Inferor Temporal gyrus
< 0.001	7.22525072	6	-31	29	Posterior cingulate
< 0.001	6.43974495	-3	-28	29	Posterior cingulate
0.0026	5.8363061	21	-40	-43	Cerebellum
0.0081	5.51446533	-15	-55	-46	Cerebellum
0.0084	5.50423813	-39	-4	14	Insula
0.0190	5.25947666	-48	11	-13	Superior temporal gyrus
0.0263	5.16038084	-30	-58	-34	Cerebellum
0.0304	5.11576033	-54	8	-10	Superior temporal gyrus

References

Acevedo, B. P., Aron, A., Fisher, H. E. & Brown, L. L. (2012) 'Neural correlates of long-term intense romantic love', *Social Cognitive and Affective Neuroscience*, **7**(2), pp. 145-159.

Aguilar-Raab, C., Eckstein, M., Geracitano, S., Prevost, M., Gold, I., Heinrichs, M., Bilderbeck, A., Ehlert, U. & Ditzen, B. (2019) 'Oxytocin modulates the cognitive appraisal of the own and others close intimate relationships', *Frontiers in neuroscience*, p. 714.

Aguilar-Raab, C., Grevenstein, D., Gotthardt, L., Jarczok, M. N., Hunger, C., Ditzen, B. & Schweitzer, J. (2018) 'Changing me, changing us: relationship quality and collective efficacy as major outcomes in systemic couple therapy', *Family process*, **57**(2), pp. 342-358.

Aron, A., Fisher, H., Mashek, D. J., Strong, G., Li, H. & Brown, L. L. (2005) 'Reward, motivation, and emotion systems associated with early-stage intense romantic love', *Journal of neurophysiology*, **94**(1), pp. 327-337.

Bartels, A. & Zeki, S. (2004) 'The neural correlates of maternal and romantic love', *Neuroimage*, **21**(3), pp. 1155-1166.

Bilek, E., Ruf, M., Schäfer, A., Akdeniz, C., Calhoun, V. D., Schmahl, C., Demanuele, C., Tost, H., Kirsch, P. & Meyer-Lindenberg, A. (2015) 'Information flow between interacting human brains: Identification, validation, and relationship to social expertise', *Proceedings of the National Academy of Sciences*, p. 201421831.

Bosch, O. J. & Young, L. J. (2018) 'Oxytocin and Social Relationships: From Attachment to Bond Disruption', in R. Hurlemann & V. Grinevich (eds), *Behavioral Pharmacology of Neuropeptides: Oxytocin*, Cham, Springer International Publishing.

Braithwaite, S. & Holt-Lunstad, J. (2017) 'Romantic relationships and mental health', *Current Opinion in Psychology*, **13**, pp. 120-125.

Chan, Y.-C., Hsu, W.-C. & Chou, T.-L. (2018) 'Dissociation between the processing of humorous and monetary rewards in the 'motivation'and 'hedonic'brains', *Scientific reports*, **8**(1), p. 15425.

Ditzen, B., Schaer, M., Gabriel, B., Bodenmann, G., Ehlert, U. & Heinrichs, M. (2009) 'Intranasal oxytocin increases positive communication and reduces cortisol levels during couple conflict', *Biological psychiatry*, **65**(9), pp. 728-731.

Dodell-Feder, D., Felix, S., Yung, M. G. & Hooker, C. I. (2015) 'Theory-of-mind-related neural activity for one's romantic partner predicts partner well-being', *Social Cognitive and Affective Neuroscience*, **11**(4), pp. 593-603.

Dölen, G., Darvishzadeh, A., Huang, K. W. & Malenka, R. C. (2013) 'Social reward requires coordinated activity of nucleus accumbens oxytocin and serotonin', *Nature*, **501**(7466), p. 179.

Doohan, E. A. M. & Manusov, V. (2004) 'The communication of compliments in romantic relationships: An investigation of relational satisfaction and sex differences and similarities in compliment behavior', *Western Journal of Communication*, **68**(2), pp. 170-194.

Ellemers, N., Kortekaas, P. & Ouwerkerk, J. W. (1999) 'Self-categorisation, commitment to the group and group self-esteem as related but distinct aspects of social identity', *European journal of social psychology*, **29**(2-3), pp. 371-389.

Filimon, F., Nelson, J. D., Sejnowski, T. J., Sereno, M. I. & Cottrell, G. W. (2020) 'The ventral striatum dissociates information expectation, reward anticipation, and reward receipt', *Proceedings of the National Academy of Sciences*, **117**(26), pp. 15200-15208.

Flanagan, J. C., Fischer, M. S., Nietert, P. J., Back, S. E., Maria, M. M.-S., Snead, A. & Brady, K. T. (2018) 'Effects of oxytocin on cortisol reactivity and conflict resolution behaviors among couples with substance misuse', *Psychiatry Research*, **260**, pp. 346-352.

Frewen, P., Schroeter, M. L., Riva, G., Cipresso, P., Fairfield, B., Padulo, C., Kemp, A. H., Palaniyappan, L., Owolabi, M., Kusi-Mensah, K., Polyakova, M., Fehertoi, N., D'Andrea, W., Lowe, L. & Northoff, G. (2020) 'Neuroimaging the consciousness of self: Review, and conceptual-methodological framework', *Neuroscience & Biobehavioral Reviews*, **112**, pp. 164-212.

Hahlweg, K. (1979) 'Konstruktion und Validierung des Partnerschaftsfragebogens PFB', *Zeitschrift für klinische Psychologie*, **8**(1), pp. 17-40.

Harbaugh, W. T., Mayr, U. & Burghart, D. R. (2007) 'Neural responses to taxation and voluntary giving reveal motives for charitable donations', *Science*, **316**(5831), pp. 1622-1625.

Haslam, S. A., O'Brien, A., Jetten, J., Vormedal, K. & Penna, S. (2005) 'Taking the strain: Social identity, social support, and the experience of stress', *British journal of social psychology*, **44**(3), pp. 355-370.

Hooley, J. M., Gruber, S. A., Scott, L. A., Hiller, J. B. & Yurgelun-Todd, D. A. (2005) 'Activation in dorsolateral prefrontal cortex in response to maternal criticism and praise in recovered depressed and healthy control participants', *Biological psychiatry*, **57**(7), pp. 809-812.

Hsu, C.-T., Conrad, M. & Jacobs, A. M. (2014) 'Fiction feelings in Harry Potter: haemodynamic response in the mid-cingulate cortex correlates with immersive reading experience', *Neuroreport*, **25**(17), pp. 1356-1361.

Izuma, K., Saito, D. N. & Sadato, N. (2008) 'Processing of social and monetary rewards in the human striatum', *Neuron*, **58**(2), pp. 284-294.

Izuma, K., Saito, D. N. & Sadato, N. (2010) 'Processing of the incentive for social approval in the ventral striatum during charitable donation', *Journal of cognitive neuroscience*, **22**(4), pp. 621-631.

Jabbi, M. & Keysers, C. (2008) 'Inferior frontal gyrus activity triggers anterior insula response to emotional facial expressions', *Emotion*, **8**(6), p. 775.

Kennedy, D. P. & Adolphs, R. (2012) 'The social brain in psychiatric and neurological disorders', *Trends in cognitive sciences*, **16**(11), pp. 559-572.

Kreuder, A. K., Wassermann, L., Wollseifer, M., Ditzen, B., Eckstein, M., Stoffel-Wagner, B., Hennig, J., Hurlemann, R. & Scheele, D. (2018) 'Oxytocin enhances the pain-relieving effects of social support in romantic couples', *Human brain mapping*.

Laurita, A. C., Hazan, C. & Spreng, R. N. (2017) 'Dissociable patterns of brain activity for mentalizing about known others: a role for attachment', *Social Cognitive and Affective Neuroscience*, **12**(7), pp. 1072-1082.

Litzinger, S. & Gordon, K. C. (2005) 'Exploring Relationships Among Communication, Sexual Satisfaction, and Marital Satisfaction', *Journal of Sex & Marital Therapy*, **31**(5), pp. 409-424.

Miedl, S. F., Blechert, J., Klackl, J., Wiggert, N., Reichenberger, J., Derntl, B. & Wilhelm, F. H. (2016) 'Criticism hurts everybody, praise only some: Common and specific neural responses to approving and disapproving social-evaluative videos', *Neuroimage*, **132**, pp. 138-147.

Mier, D., Lis, S., Neuthe, K., Sauer, C., Esslinger, C., Gallhofer, B. & Kirsch, P. (2010) 'The involvement of emotion recognition in affective theory of mind', *Psychophysiology*, **47**(6), pp. 1028-1039.

Moll, J., Krueger, F., Zahn, R., Pardini, M., de Oliveira-Souza, R. & Grafman, J. (2006) 'Human fronto–mesolimbic networks guide decisions about charitable donation', *Proceedings of the National Academy of Sciences*, **103**(42), pp. 15623-15628.

Nicolson, N. A., Peters, M. L. & in den Bosch - Meevissen, Y. M. C. (2020) 'Imagining a positive future reduces cortisol response to awakening and reactivity to acute stress', *Psychoneuroendocrinology*, **116**, p. 104677.

Oyama, K., Hernádi, I., Iijima, T. & Tsutsui, K.-I. (2010) 'Reward prediction error coding in dorsal striatal neurons', *Journal of Neuroscience*, **30**(34), pp. 11447-11457.

Packheiser, J., Berretz, G., Rook, N., Bahr, C., Schockenhoff, L., Güntürkün, O. & Ocklenburg, S. (2021) 'Investigating real-life emotions in romantic couples: a mobile EEG study', *Scientific reports*, **11**(1), p. 1142.

Pfeifer, A.-C., Schroeder-Pfeifer, P., Schneider, E., Schick, M., Heinrichs, M., Bodenmann, G., Ehlert, U., Herpertz, S. C., Läuchli, S. & Eckstein, M. (2020) 'Oxytocin and positive couple interaction affect the perception of wound pain in everyday life', *Molecular Pain*, **16**, p. 1744806920918692.

Riem, M. M., Van Ijzendoorn, M. H., Tops, M., Boksem, M. A., Rombouts, S. A. & Bakermans-Kranenburg, M. J. (2012) 'No laughing matter: intranasal oxytocin administration changes functional brain connectivity during exposure to infant laughter', *Neuropsychopharmacology*, **37**(5), p. 1257.

Scheele, D., Wille, A., Kendrick, K. M., Stoffel-Wagner, B., Becker, B., Güntürkün, O., Maier, W. & Hurlemann, R. (2013) 'Oxytocin enhances brain reward system responses in men viewing the face of their female partner', *Proceedings of the National Academy of Sciences*, **110**(50), pp. 20308-20313.

Scheepers, D. & Ellemers, N. (2019) 'Social identity theory', *Social psychology in action*, Springer.

Smith, M. L., Asada, N. & Malenka, R. C. (2021) 'Anterior cingulate inputs to nucleus accumbens control the social transfer of pain and analgesia', *Science*, **371**(6525), pp. 153-159.

Telzer, E. H., Masten, C. L., Berkman, E. T., Lieberman, M. D. & Fuligni, A. J. (2010) 'Gaining while giving: An fMRI study of the rewards of family assistance among White and Latino youth', *Social neuroscience*, **5**(5-6), pp. 508-518.

Thornton, M. A., Weaverdyck, M. E., Mildner, J. N. & Tamir, D. I. (2019) 'People represent their own mental states more distinctly than those of others', *Nature Communications*, **10**(1), p. 2117.

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., Mazoyer, B. & Joliot, M. (2002) 'Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain', *Neuroimage*, **15**(1), pp. 273-289.

Van der Gaag, C., Minderaa, R. B. & Keysers, C. (2007) 'Facial expressions: what the mirror neuron system can and cannot tell us', *Social neuroscience*, **2**(3-4), pp. 179-222.

Van Overwalle, F. & Baetens, K. (2009) 'Understanding others' actions and goals by mirror and mentalizing systems: A meta-analysis', *Neuroimage*, **48**(3), pp. 564-584.

Wagner, U., Galli, L., Schott, B. H., Wold, A., van der Schalk, J., Manstead, A. S., Scherer, K. & Walter, H. (2014) 'Beautiful friendship: social sharing of emotions improves subjective feelings and activates the neural reward circuitry', *Social Cognitive and Affective Neuroscience*, **10**(6), pp. 801-808.

Warth, M., Stoffel, M., Winter, F., Jarczok, M. N., Aguilar-Raab, C. & Ditzen, B. (2020) 'Instructed Partnership Appreciation in Depression: Effects on Mood, Momentary Relationship Satisfaction, and Psychobiological Arousal', *Frontiers in psychiatry*, **11**, p. 701.

Zietlow, A.-L., Eckstein, M., Nonnenmacher, N., Reck, C., Schaer, M., Bodenmann, G., Heinrichs, M. & Ditzen, B. (2018) 'Dyadic coping and its underlying neuroendocrine mechanisms–implications for stress regulation', *Frontiers in psychology*, **9**, p. 2600.

AUTHORS' CONTRIBUTION STATEMENT:

B.D., P.K., G.S. E.B and M.E. designed the study; G.S. and M.E. lead the study; G.S., E.B. and M.E. collected the data; M.F.G. and E.B. established the experimental set-up; G.S. M.F.G and M.E. ran the reported analyses, M.E., and B.D. wrote the manuscript; All authors provided comments on the manuscript.

FUNDING STATEMENT:

This work was partially supported by German Research Foundation DFG through Clinical Research Unit KFO 256, KI 576/15-2, ME 1591/4-2.

The funding agency had no role in the planning of the study design and will not be involved in data collection, analysis, decision to publish or preparation of the manuscript.

COMPETING INTERESTS STATEMENT:

The authors declare that they have no competing interests.

ACKNOWLEDGMENTS: We wish to thank Fabienne Ibert, Jasmin Buchholz, Julia Hein and Laura Kampouridis for their skilled support in data collection and Sarah Fancy for proofreading the manuscript.