An engineered xCas12i with high activity, high specificity and broad PAM range

Hainan Zhang[1, 4], Xiangfeng Kong[1, 4], Mingxing Xue[1, 4], Zikang Wang[1, 4], Yinghui Wei[1, 4],

Haoqiang Wang[1, 4], Jingxing Zhou[1], Weihong Zhang[1], Mengqiu Xu[1], Xiaowen Shen[1],

Jinhui Li[1], Jing Hu[1], Na Zhong[1], Yingsi Zhou[1#], Hui Yang[1, 2, 3#]

[1] HUIEDIT Therapeutics Inc., Shanghai 200031, China.

[2] HUIGENE Therapeutics Inc., Shanghai 200131, China.

[3] Institute of Neuroscience, State Key Laboratory of Neuroscience, Key Laboratory of

Primate Neurobiology, Center for Excellence in Brain Science and Intelligence Technology,

Chinese Academy of Sciences, Shanghai 200031, China.

[4] These authors contributed equally to this work.

Correspondence to: ys@huiedit.com (Y.Z.), huiyang@huidagene.com (H.Y.)

Abstract

The type-V CRISPR effector Cas12i, with its smaller size, short crRNA guiding, and

self-processing features, is a potentially versatile genome editing tool. By screening

Cas12i proteins from a metagenomic database, we identified a natural variant with high

activity in mammalian cells, named as xCas12i. We further engineered the

PAM-interacting, REC, and RuvC domains for enhanced cleavage activity and specificity.

This variant, named as high-fidelity Cas12Max, exhibited robust genome editing activity

and minimal off-target activity with a broad 5'-TN recognition profile. With the fusion of

deaminase TadA8e and further optimization of xCas12i, the base editor dCas12i-Tad8e

also showed the high editing efficiency. This study provides highly efficient and specific

tools for gene therapy.

Keywords: CRISPR, Cas12i, broad PAM recognition, dCas12i base editor

Introduction

The clustered regularly interspaced short palindromic repeats-Cas (CRISPR-Cas) systems, including type II Cas9 and type V Cas12 systems, which serve in the adaptive immunity of prokaryotes against viruses, have been developed into genome editing tools[1-3]. Compared with type II systems, the type V systems including V-A to V-K showed more functional diversity[4, 5]. Amongst them, Cas12i has a relatively smaller size (1033-1093 aa), compared to SpCas9 and Cas12a, and has a 5'-TTN protospacer adjacent motif (PAM) preference[4, 6, 7]. Cas12i is characterized by the capability of autonomously processing precursor crRNA (pre-crRNA) to form short mature crRNA. Cas12i mediates cleavage of dsDNA with a single RuvC domain, by preferentially nicking the non-target strand and then cutting the target strand[8-10]. These intrinsic features of Cas12i enable multiplex high-fidelity genome editing. However, the natural variants of Cas12i (Cas12i1 and Cas12i2), showed low editing efficiency which limits their utility for therapeutic gene editing.

To address these limitations, we screened ten natural Cas12i variants and found one, xCas12i, with robust high activity in HEK293T cells. Engineering of xCas12i by arginine substitutions at the PAM-interacting (PI), REC and RuvC domains led to the production of a variant, high-fidelity Cas12Max (hfCas12Max), with significantly elevated editing activity and minimal off-target cleavage efficiency. In addition, we assessed the base editing efficiency of xCas12i-based base editor, and thus expanded the genome-editing toolbox.

Results

Identification and characterization of type V-I systems xCas12i

In order to identify more Cas12i variants, we developed and employed a bioinformatics pipeline to annotate Cas12i proteins, CRISPR arrays and predicted PAM preferences, and found 10 new CRISPR/Cas12i systems. To evaluate the activity of these Cas12i variants in mammalian cells, we designed a fluorescent reporter system which detected the increased enhanced green fluorescent protein (EGFP) signal intensity activated by Cas-mediated dsDNA cleavage or double strand breaks (Fig.S1a). This system relied on

the co-transfection of a plasmid coding for mCherry, a nuclear localization signal (NLS)

-tagged Cas protein and its guide RNA (gRNA) or crRNA, and one coding BFP and

activatable EGxxFP cassette, which is EGxx-target site-xxFP[11]. EGFP activation was

carried out by Cas mediated DSB and single-strand annealing (SSA)-mediated repair.

Using this system, we observed that a variant, xCas12i, with targeted crRNA induced

significant activation of EGFP expression (Fig. 1a, Fig. S1b), and exhibited a higher

editing frequency than LbCas12a or SpCas9 as determined by Fluorescence Activated

Cell Sorter (FACS) analysis (Fig. 1a). The xCas12i variant was smaller in size compared

to SpCas9 and LbCas12a (Fig. S2a). We explored the effects of spacer length on

cleavage efficiency in xCas12i, and found that 17-22 nt were optimal length for their

activation (Fig. S2b). Considering the 5'-TTN PAM preference of Cas12i, we performed a

NTTN PAM identification assay using the reporter system. We found that xCas12i showed

a consistent high frequency of EGFP activation at target sites with 5'-NTTN PAM

sequences, while LbCas12a had comparable activity at 5'-TTTN PAM, respectively (Fig.

S2c).

To further confirm the dsDNA cleavage activity of xCas12i in mammalian cells, we

transfected an all-in-one plasmid encoding NLS tagged xCas12i with crRNAs targeting 37

sites from *TTR*[12], *PCSK9*[13] in HEK293T or *Ttr* in N2a cells. The editing efficiency, *i.e.* indel

(insertion and deletion) formation at these loci was measured 48 hours after transfection

using FACS and targeted deep sequencing (Fig. S2d). We found that xCas12i mediated a

high frequency, up to 90%, of indel formation at most sites from *Ttr*, *TTR* and *PCSK9*, with

a mean indel formation rate of over 50% (Fig. S2e-f). These data indicate that xCas12i

exhibits a robust genome editing efficiency in mammalian cells, suggesting it has

excellent potential for therapeutic genome editing applications.

Engineered xCas12i mediates high-efficiency editing at 5'-TN PAM sites

To enhance its activity and expand its scope of PAM site recognition, we sought to

engineer xCas12i protein via mutagenesis and screen for variants with higher efficiency

and broader PAM using a reporter system, similar to what is described above. Substitution

of an amino acid in the DNA-binding pocket with positively charged arginine (R) was

shown to enhance the activity of the type V system[14-16]. Combined with predictive

structural analysis of xCas12i, we performed an arginine scanning mutagenesis approach

in the PI, REC-I and RuvC-II domains, generating a library of over 500 mutants (Fig. 1b,

Fig. S3a). We then individually transfected these mutant variants with an activatable

EGFP reporter system in HEK293T cells and analyzed them by FACS (Fig. 1b). Based on

the fluorescence intensity of cells with activated EGFP, over 100 mutants showed an

increased frequency of activated cells relative to wild type (WT) xCas12i, and one mutant,

named as Cas12Max, containing N243R showed a 3.4-fold improvement (Fig. S3a). We

next targeted *DMD* or *Ttr* sites using the fluorescent reporter system, and found that

Cas12Max displayed a markedly increased frequency of EGFP activation, relative to WT

xCas12i (Fig. 1c, Fig. S4a-b). To further test the efficacy of Cas12Max in targeting

genomic loci, we designed a total of eight gRNAs to target sites *TTR* and *PCSK9* in

HEK293T cells and three more targeting *Ttr* in N2a cells. Consistent with our previous

result, Cas12Max exhibited a significantly increased frequency of indels compared to WT

xCas12i (Fig. 1d-e).

Additionally, to investigate Cas12Max's PAM preference, we performed a 5'-NNN

PAM recognition assay by designing reporter plasmids with the same target sequence but

different PAM. Besides showing a consistent or higher cleavage activity at sites with a

5'-TTN PAM, Cas12Max showed a similarly high cleavage activity for targets with TNN,

ATN, GTN and CTN PAM sites, compared with the commonly used Cas12[7, 17] (LbCas12a,

Ultra-AsCas12a) (Fig. 1f). Taken together, these results demonstrate that Cas12Max

exhibits high-efficiency editing activity with highly flexible 5'-TN or 5'-TNN PAM

recognition.

hfCas12Max mediates high-efficient and specific genome editing

To examine the specificity of Cas12Max, we transfected a construct designed to express it

with crRNA targeting *TTR*[12], and performed indel frequency analysis of on- and off-target

(OT) sites predicted by Cas-OFFinder[18]. Using reporter system or targeted deep

sequence analysis, we found that Cas12Max efficiently edited target sites and resulted in

significant indel formation at 2 of 3 predicted off-target sites (Fig. S5). To eliminate the

off-target activity of Cas12Max, we screened these mutants with mutations in the REC

and RuvC domains[19], which have undiminished on-target cleavage activity, for those with

no off-target activity, using two activatable reporter systems, each containing one OT site

(Fig. 1b). We found that four mutants (v4.1-V880R, v4.2-M923R, v4.3-D892R and

v4.4-G883R) maintained a high level of on-target editing activity and showed significantly

reduced off-target EGFP activation (Fig. S6a). We further combined these four amino acid

substitutions with N243R and/or E336R of Cas12Max and found that the variant v6.3

(N243R/D892R/G883R) showed the lowest off-target EGFP activation at OT.1 and OT.2

sites and high on-target at the ON.1 site (Fig. 1g, Fig. S6b-c). Targeted TIDE analysis of

endogenous TTR.2 site and its off-target sites in HEK293T showed that v6.3

(N243R/D892R/G883R) significantly reduced off-target indel frequencies at six OT sites

and retained on-target at ON site, compared to Cas12Max (Fig. 1h). In addition, relative to

Cas12Max (v1.1), v6.3 (N243R/D892R/G883R) retained comparable or even higher

on-target activity at DMD.1, DMD.2 and DMD.3 sties (Fig. S6d). Therefore, we named

v6.3 as high-fidelity Cas12Max (hfCas12Max).

To comprehensively evaluate the performance of hfCas12Max in human cells, we

designed large number of target sites in the exons of *TTR* for various Cas nucleases. In

total, editing activity was monitored over 30 sites for hfCas12Max with TTN PAMs, 40 sites

for SpCas9 with NGG PAMs, 4 sites for LbCas12a with TTTN PAMs, 4 sites for Ultra

AsCas12a, and 7 sites for KKH-saCas9 with NNNRRT PAMs. Indel analysis showed that

hfCas12Max exhibited an average efficiency of 70%, higher activity than other Cas

nucleases, comparable activity with Cas12Max (Fig. 1i, Fig. S7a). Using 5'-NNN PAM

reporter system, hfCas12Max maintained the broad 5'-TN and 5'-TNN PAM recognition

profile similarly to Cas12Max (Fig. 1f). To further evaluate the specificity of hfCas12Max in

human cells, we determined indel frequencies of P2RX5 and NLRC4 on-target and their

corresponding *in silico* predicted off-target sites[20]. TIDE analysis showed that

hfCas12Max had a higher on-target editing efficiency and similarly almost no indel activity

at potential off target sites, compared to Ultra AsCas12a and LbCas12a (Fig. S8a-b).

Overall, these results demonstrate that hfCas12Max has high efficiency and specificity

and is superior to SpCas9 and other Cas12 nucleases.

dxCas12i-based base editors efficiently mediate base editing in mammalian cells

We further explored the base editing of xCas12i by generating a nuclease-deactivation xCas12i (dxCas12i). This was done by first introducing single mutations (D650A, D700A, E875A, or D1049A) in the conserved active site of xCas12i based on alignment to Cas12i1[8] and Cas12i2[10] (Fig. S9a-b). Then, these variants were fused with TadA8e^{V106W} or human APOBEC3A (hA3A^{W104A}) to form the dxCas12i base editors TadA8e.1-dxCas12i and hA3A.1-dxCas12i, respectively[21-23]. The initial versions of TadA8e.1-dxCas12i and hA3A.1-dxCas12i showed low base editing activity with frequencies of 8% A-to-G and 2% C-to-T, respectively (Fig. 2a-d). To address this, we introduced single and combined mutations for high cleavage activity into the PI and Rec domains of dxCas12i, which resulted in significantly increased A-to-G editing activity. Among the improved variants, dxCas12i-TadA8e-v2.2 (N243R/E336R) achieved 50% activity at A9 and A11 sites of the *KLF4* locus, markedly higher than the 30% activity of dLbCas12a-TadA8e (Fig. 2b, Fig. S10a-b). At target sites within *PCSK9*, TadA8e-dxCas12i-v2.2 showed a similarly increased efficiency to mediate A-to-G transitions, higher than dLbCas12a-TadA8e (Fig. S11). We then further engineered the NLS, linker, TadA8e protein and the N-terminus to produce dxCas12i-TadA8e-v4.3 which exhibited a nearly 80% A-to-G editing efficiency, while the editing activities of other dxCas12i-ABE versions were unchanged (Fig. 2c, Fig. S10c-d). In addition, the substitution of dxCas12i-v1.2 (N243R) or dxCas12i-v2.2 (N243R/E336R) showed consistently elevated C-to-T editing at C7 and C10 sites of *DYRK1A*, even higher at C7 by dLbCas12a-hA3A (Fig. 2d). These results together demonstrate that engineered dxCas12i-based editors exhibit the high base editing activity in mammalian cells.

Discussion

In this study, we demonstrate that the Type V-I Cas12i system enables versatile and efficient genome editing in mammalian cells. We found a natural Cas12i variant, xCas12i, that shows high editing efficiency at TTN-PAM sites. By semi-rational design and protein

engineering of its PI, REC, RuvC domains, we obtained a high-efficiency, high-fidelity

variant, hfCas12Max, which contains N243R, E336R, and D892R substitutions. In

agreement with the hypothesis that introducing arginine at key sites could strengthen the

binding between Cas and DNA, the introduction of N243R in the PI domain and E336R at

REC domain significantly increased editing activity and expanded PAM recognition.

Interestingly, D892R or G883R substitutions in the RuvC domain reduced off-target and

retained on-target cleavage activity, whereas alanine substitutions[24, 25], which has been

used to reduce off-target activity, did not (Fig. S6c). The D892R substituted hfCas12Max

was obviously more sensitive to mismatch, which suggests that D892R or G883R

improved sgRNA binding specificity. Our data suggests that a semi-rational engineering

strategy with arginine substitutions based on the EGFP-activated reporter system could

be used as a general approach to improve the activity of CRISPR editing tools.

Through engineering, our Cas12i system has achieved high editing activity, high

specificity and a broad PAM range, comparable to SpCas9, and better than other Cas12

systems. Given its smaller size, short crRNA guide, and self-processing features[4, 8, 10], the

Type V-I Cas12i system is suitable for *in vivo* multiplexed gene editing applications,

including AAV[26] or LNP[12, 13].

In addition, we have confirmed that the Type V-I Cas12i system can be used in base

editing applications. For base editor, the dCas12i system shows high A-to-G editing at

A9-A11 sites even A19 of KLF locus, and C-to-T editing at A7-A10 sites, which is similar to

the dCas12a system but is distinct from the dCas9/nCas9 system. Comparable to

dCas12a, dCas12i-BE exhibited higher base editing activity at KLF4, PCSK9 and

DYRK1A loci (Fig. 4c-d), suggesting it may have more potential as a base editor. This

suggests that the dCas12i system is useful for broad genome engineering applications,

including epigenome editing, genome activation, and chromatin imaging[1, 27-30].

In summary, the Cas12i system described here, which has robust editing activity and

high specificity, is a versatile platform for genome editing or base editing in mammalian

cells and could be useful in the future for *in vivo* or *ex vivo* therapeutic applications.

Note that two groups have recently reported improved Cas12i2 using similar

strategies in this study[31, 32].

Authors' Contributions

H.Z., and Y.Z. jointly conceived and designed the project. H.Z., H.W., J.Z., W.Z., M.X., and X.K., performed experiments with plasmid construction, cell culture and FACS and data analysis. M.X., Z.W., Y.W., X.S., J.L., J.H., and N.Z. assisted with experiments. H.Y. supervised the whole project. H.Z., Y.Z., and H.Y. wrote and revised the manuscript.

Conflict of interest

The authors disclose a patent application relating to aspects of this work. Y.Z. and H.Y. is the founder of HUIEDIT Therapeutics Inc., and H.Y. is also the founder of HUIGENE Therapeutics Inc.

References

1.   Anzalone, A.V., Koblan, L.W. & Liu, D.R. Genome editing with CRISPR-Cas nucleases, base editors, transposases and prime editors. *Nat Biotechnol* **38**, 824-844 (2020).

2.   Doudna, J.A. The promise and challenge of therapeutic genome editing. *Nature* **578**, 229-236 (2020).

3.   Makarova, K.S. et al. Evolutionary classification of CRISPR-Cas systems: a burst of class 2 and derived variants. *Nat Rev Microbiol* **18**, 67-83 (2020).

4.   Yan, W.X. et al. Functionally diverse type V CRISPR-Cas systems. *Science* **363**, 88-91 (2019).

5.   Kleinstiver, B.P. et al. Genome-wide specificities of CRISPR-Cas Cpf1 nucleases in human cells. *Nature Biotechnology* **34**, 869-+ (2016).

6.   Cong, L. et al. Multiplex Genome Engineering Using CRISPR/Cas Systems. *Science* **339**, 819-823 (2013).

7.   Zetsche, B. et al. Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system. *Cell* **163**, 759-771 (2015).

8.   Zhang, B. et al. Mechanistic insights into the R-loop formation and cleavage in CRISPR-Cas12i1. *Nature communications* **12**, 3476 (2021).

9.   Zhang, H., Li, Z., Xiao, R. & Chang, L. Mechanisms for target recognition and cleavage by the Cas12i RNA-guided endonuclease. *Nat Struct Mol Biol* **27**, 1069-1076 (2020).

10.   Huang, X. et al. Structural basis for two metal-ion catalysis of DNA cleavage by Cas12i2. *Nature communications* **11**, 5241 (2020).

11.   Yang, Y. et al. Highly Efficient and Rapid Detection of the Cleavage Activity of Cas9/gRNA via a Fluorescent Reporter. *Appl Biochem Biotechnol* **180**, 655-667

(2016).

12.   Gillmore, J.D. et al. CRISPR-Cas9 In Vivo Gene Editing for Transthyretin Amyloidosis.
The New England journal of medicine **385**, 493-502 (2021).

13.   Musunuru, K. et al. In vivo CRISPR base editing of PCSK9 durably lowers cholesterol
in primates. *Nature* **593**, 429-434 (2021).

14.   Strecker, J. et al. Engineering of CRISPR-Cas12b for human genome editing. *Nature*
communications **10**, 212 (2019).

15.   Kleinstiver, B.P. et al. Engineered CRISPR-Cas12a variants with increased activities
and improved targeting ranges for gene, epigenetic and base editing. *Nat Biotechnol*
37, 276-282 (2019).

16.   Xu, X. et al. Engineered miniature CRISPR-Cas system for mammalian genome
regulation and editing. *Mol Cell* **81**, 4333-4345 e4334 (2021).

17.   Zhang, L. et al. AsCas12a ultra nuclease facilitates the rapid generation of therapeutic
cell medicines. *Nature communications* **12**, 3908 (2021).

18.   Bae, S., Park, J. & Kim, J.S. Cas-OFFinder: a fast and versatile algorithm that
searches for potential off-target sites of Cas9 RNA-guided endonucleases.
Bioinformatics **30**, 1473-1475 (2014).

19.   Yuen, C.T.L. et al. High-fidelity KKH variant of Staphylococcus aureus Cas9 nucleases
with improved base mismatch discrimination. *Nucleic Acids Res* **50**, 1650-1660
(2022).

20.   Kim, D.Y. et al. Efficient CRISPR editing with a hypercompact Cas12f1 and
engineered guide RNAs delivered by adeno-associated virus. *Nat Biotechnol* **40**,
94-102 (2022).

21.   Wang, X. et al. Cas12a Base Editors Induce Efficient and Specific Editing with Low
DNA Damage Response. *Cell Rep* **31**, 107723 (2020).

22.   Richter, M.F. et al. Phage-assisted evolution of an adenine base editor with improved
Cas domain compatibility and activity. *Nat Biotechnol* **38**, 883-891 (2020).

23.   Li, X. et al. Base editing with a Cpf1-cytidine deaminase fusion. *Nat Biotechnol* **36**,
324-327 (2018).

24.   Bravo, J.P.K. et al. Structural basis for mismatch surveillance by CRISPR-Cas9.
Nature **603**, 343-347 (2022).

25.   Kleinstiver, B.P. et al. High-fidelity CRISPR-Cas9 nucleases with no detectable
genome-wide off-target effects. *Nature* **529**, 490-495 (2016).

26.   Wang, D., Zhang, F. & Gao, G. CRISPR-Based Therapeutic Genome Editing:
Strategies and In Vivo Delivery by AAV Vectors. *Cell* **181**, 136-150 (2020).

27.   Wang, H. et al. CRISPR-Mediated Programmable 3D Genome Positioning and
Nuclear Organization. *Cell* **175**, 1405-1417 e1414 (2018).

28.   Konermann, S. et al. Genome-scale transcriptional activation by an engineered
CRISPR-Cas9 complex. *Nature* **517**, 583-588 (2015).

29.   Nakamura, M., Gao, Y., Dominguez, A.A. & Qi, L.S. CRISPR technologies for precise
epigenome editing. *Nat Cell Biol* **23**, 11-22 (2021).

30.   Fellmann, C., Gowen, B.G., Lin, P.C., Doudna, J.A. & Corn, J.E. Cornerstones of
CRISPR-Cas in drug discovery and therapy. *Nat Rev Drug Discov* **16**, 89-100 (2017).

31.   Chen, Y. et al. Synergistic engineering of CRISPR-Cas nucleases enables robust

mammalian genome editing. *Innovation (Camb)* **3**, 100264 (2022).

32.   McGaw, C. et al. Engineered Cas12i2 is a versatile high-efficiency platform for

therapeutic genome editing. *Nature communications* **13**, 2833 (2022).

Material and methods

Plasmid vector construction.

Human codon-optimized *Cas12i*, *TadA8e* and human *APOBEC3A* genes were

synthesized by the GenScript Co., Ltd., and cloned to generate

pCAG_NLS-Cas12i-NLS_pA_pU6_BpiI_pCMV_mCherry_pA by Gibson Assembly. crRNA

oligos were synthesized by HuaGene Co., Ltd., annealed and ligated into *Bpi*I site to

produce the pCAG_NLS-Cas12i-NLS_pA_pU6_crRNA_pCMV_mCherry_pA.

Cell culture, transfection and flow cytometry analysis.

The mammalian cell lines used in this study were HEK293T and N2A. Cells were cultured

in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% FBS,

penicillin/streptomycin and GlutMAX. Transfections were performed using Polyetherimide

(PEI). For variant screening, HEK293T cells were cultured in 24-well plates, and after 12

hours 2 µg of the plasmids (1 µg Cas12i of a mutant plasmid and 1 µg of the reporter

plasmid) were transfected into these cells with 4 µL PEI. 48 hours after transfection,

mCherry and EGFP fluorescence were analyzed using a Beckman CytoFlex

flow-cytometer. For assay of mutations in target sites of endogenous genes, 1 µg plasmid

with Cas12i targeting crRNA was transfected into HEK293T or N2A cells, which were then

sorted using a BD FACS Aria III, BD LSRFortessa X-20 flow cytometer, 48 hours after

transfection.

Detection of gene editing frequency.

Six thousand sorted cells were lysed in 20 µl of lysis buffer (Vazyme). Targeted sequence

primers were synthesized and used in nested PCR amplification by Phanta Max

Super-Fidelity DNA Polymerase (Vazyme). TIDE and targeted deep sequence analysis

was used to determine indel frequencies. Sanger sequencing and EditR were used for

quantification of base substitutions (A-to-G or C-to-T).

Figure Legends

Fig.1 hfCas12Max, an engineered natural variant xCas12i, mediated high-efficient

and -specificity genome editing in mammalian cells. a, Transfection of plasmids

coding Cas12i and sgRNA activates EGFP. **b**, A natural variant xCas12i mediated EGFP

activation efficiency determined by flow cytometry. **c**, Flow diagram for detection of

genome editing efficiency by transfection of an all-in-one plasmid containing xCas12i and

targeted gRNA into HEK293T cells, followed by FACS and NGS analysis. **d**, xCas12i

mediated robust genome editing at the *Ttr* locus in N2a cells and *TTR* and *PCSK9* in

HEK293T cells. **e-g**, Cas12Max exhibited significantly increased cleavage activity than

xCas12i at various genomic or reporter plasmids target sites. **h**, Both Cas12Max and

hfCas12Max exhibited a broader PAM recognition profile than other Cas proteins,

including 5'-TN and 5'-TNN PAM. **i**, v6.3 reduced off-target at OT.1 and OT.2 sites and

retained indel activity at *TTR.2*-ON targets, compared to v1.1-Cas12Max. **j**, TIDE analysis

showed that v6.3-hfCas12Max retained comparable activity at TTR.2-ON targets and

almost no at 6 OT sites, to Cas12Max. **k**, Comparison of indel activity from Cas12Max,

hfCas12Max, LbCas12a, Ultra AsCas12a, SpCas9 and KKH-saCas9 at TTR locus.

HfCas12Max retained the comparable activity of Cas12Max, and higher gene-editing

efficiency than other Cas proteins. Each dot represents one of three repeats of single

target site.

Fig.2 dCas12Max-TadA8e and dCas12Max-hA3A.1 mediated high base editing

activity. a, Schematics of different versions of dCas12Max base editors. **b**, The A-to-G

editing activity of TadA8e.1-dxCas12i-v2.2 was largely increased by combining v1.2 and

v1.3 variant. **c**, Comparison of A-to-G editing frequency at the KLKF4 site from

TadA8e.1-dxCas12i-v1.2, v2.2 and v4.3, ABE-dCas12Max (TadA8e-dxCas12i-v4.3)

showed a high editing activity of 80%. **d**, Comparison of C-to-T editing frequency at the

DYRK1A site from hA3A.1-dxCas12i, -v1.2 and v2.2, CBE-dCas12Max

(hA3A.1-dxCas12i-v2.2) showed a high editing activity of 50%.

Supplementary Figure Legends

Supplementary Figure 1.   Screen for functional Cas12i in HEK293T cells. a, Five of

ten natural Cas12i nuclease mediated EGFP-activated efficiency in HEK293T cells.

Supplementary Figure 2.   Identification and characterization of type V-I systems. a,

Nuclease domain organization of SpCas9, LbCas12a, and xCas12i. **b**, Optimal spacer

length for xCas12i. **c**, PAM scope comparison of LbCas12a, and xCas12i. xCas12i

exhibited a higher editing efficiency at 5'-TTN PAM than Cas12a. **d**, xCas12i mediated

robust genome editing (up to 90%) at the *Ttr* locus in N2a cells and *TTR* and *PCSK9* in

HEK293T cells.

Supplementary Figure 3. Screen for engineered xCas12i mutants with

high-efficiency editing activity. a, Schematics of protein engineering strategy using an

activatable EGFP reporter screening system. **b**, The relative editing frequencies of over

500 rationally engineered xCas12i mutants.

Supplementary Figure 4. Other mutants mediated high-efficiency editing. a-b,

xCas12i mutant with N243R increased 1.2, 5, 20-fold activity at DMD.1, DMD.2 and

DMD.3 locus. **c**, Both Cas12Max (xCas12i-N243R) and Cas12Max-E336R elevated

EGFP-activated fluorescent at different PAM recognition sites.

Supplementary Figure 5. Cas12Max induced off-target editing efficiency at sites

with mismatches using the reporter system (a) and targeted deep sequence (b).

Supplementary Figure 6. hfCas12Max mediates high-efficiency and -specificity

editing. a, Schematics of protein engineering strategy using EGFP activatable reporter

screening system with on-targeted and off-targeted crRNA. **b**, Rational protein

engineering screen of over 200 mutants for highly-fidelity Cas12Max.Four mutants show

significantly decreased activity at both OT (off-target) sites and retains at ON.1 (on-target)

site. **c**, Different versions of xCas12i mutants. **d**, v6.3-hfCas12Max reduced off-target at

OT.1 and OT.2 sites and retained indel activity at *TTR.2*-ON targets, compared to

v1.1-Cas12Max. **e**, v6.3-hfCas12Max exhibited comparable indel activity at DMD.1,

DMD.2, and higher at DMD.3 locus, than v1.1-Cas12Max.

Supplementary Figure 7. hfCas12Max exhibited the high editing activity.   **a**, Indel

frenquencies from Cas12Max, hfCas12Max, LbCas12a, Ultra AsCas12a, SpCas9 and

KKH-saCas9 at *TTR* locus.

Supplementary Figure 8. hfCas12Max mediates high-efficient and -specific editing.

a-b, Off-target efficiency of hfCas12Max, LbCas12a, and UltraAsCas12a at in-silico

predicted off-target sites, determined by targeted deep sequencing. Sequences of

on-target and predicted off-target sites are shown, PAM sequences are in blue and

mismatched bases are in red.

Supplementary Figure 9. Conserved cleavage sites of Cas12i. a, Sequence alignment

of xCas12i, Cas12i1 and Cas12i2 shows that D650, D700, E875 and D1049 are

conserved cleavage sites at RuvC domain. **b**, Introducing point mutations off D650A,

E875A, and D1049A results in abolished activity of xCas12i.

Supplementary Figure 10. Other strategies for high-efficiency ABE-dxCas12i. a,

TadA8e.1-dxCas12i-v1.2 and v1.3 exhibits significantly increased A-to-G editing activity

among various variants at KLKF4 site of genome. **b**, Unchanged or even decreased

editing activity from various dCas12-ABEs carrying different NLS at N-terminal. **c**,

Increased A-to-G editing activity of TadA8e-dxCas12i-v4.3 by combining v2.2,

changed-NLS linker and high-activity Tade8e. **d**, dxCas12i-ABE-N by TadA at the

C-terminus of dCas12 slightly increased editing activity.

Supplementary Figure 11. Comparison of editing frequencies induced by various

dCas12-BEs at different genomic target sites. a, Comparison of A-to-G editing

frequencies induced by indicated TadA8e.1-dxCas12i, v1.2, v2.2, and TadA8e.1-dCas12a

at *PCSK9* genomic locus.

Fig.1 hfCas12Max, an engineered natural variant xCas12i, mediated high-efficient and -specificity genome editing in mammalian cells. a, A natural variant xCas12i mediated EGFP activation efficiency determined by flow cytometry. **b**, Schematics of protein engineering strategy for mutants with high efficiency and high fidelity using an activatable EGFP reporter screening system with on-targeted and off-targeted crRNA. **c-e**, Cas12Max exhibited significantly increased cleavage activity than xCas12i at various genomic or reporter plasmids target sites. **f**, Both Cas12Max and hfCas12Max exhibited a broader PAM recognition profile than other Cas proteins, including 5'-TN and 5'-TNN PAM. **g**, v6.3 reduced off-target at OT.1 and OT.2 sites and retained indel activity at TTR-ON targets, compared to v1.1-Cas12Max. **h**, TIDE analysis showed that v6.3-hfCas12Max retained comparable activity at TTR.2-ON targets and almost no at 6 OT sites, to Cas12Max. **i**, Comparison of indel activity from Cas12Max, hfCas12Max, LbCas12a, Ultra AsCas12a, SpCas9 and KKH-saCas9 at TTR locus. hfCas12Max retained the comparable activity of Cas12Max, and higher gene-editing efficiency than other Cas proteins. Each dot represents one of three repeats of single target site.

Fig.2 dCas12Max-TadA8e and dCas12Max-hA3A.1 mediated high base editing activity. **a**, Schematics of different versions of dCas12Max base editors. **b**, The A-to-G editing activity of TadA8e.1-dxCas12i-v2.2 was largely increased by combining v1.2 and v1.3 variant. **c**, Comparison of A-to-G editing frequency at the KLKF4 site from TadA8e.1-dxCas12i-v1.2, v2.2 and v4.3, ABE-dCas12Max (TadA8e-dxCas12i-v4.3) showed a high editing activity of 80%. **d**, Comparison of C-to-T editing frequency at the DYRK1A site from hA3A.1-dxCas12i, -v1.2 and v2.2, CBE-dCas12Max (hA3A.1-dxCas12i-v2.2) showed a high editing activity of 50%.

Supplementary Figure 1. Screen for functional Cas12i in HEK293T cells. a, Transfection of plasmids coding Cas12i and crRNA mediated EGFP activation. **b,** Five of ten natural Cas12i nuclease mediated EGFP-activated efficiency in HEK293T cells.

Supplementary Figure 2. Identification and characterization of type V-I systems. a, Nuclease domain organization of SpCas9, LbCas12a, and xCas12i. **b,** Optimal spacer length for xCas12i. **c**, PAM scope comparison of LbCas12a, and xCas12i. xCas12i exhibited a higher editing efficiency at 5'-TTN PAM than Cas12a. **d**, Flow diagram for detection of genome editing efficiency by transfection of an all-in-one plasmid containing xCas12i and targeted gRNA into HEK293T cells, followed by FACS and NGS analysis. **e-f**, xCas12i mediated robust genome editing (up to 90%) at the Ttr locus in N2a cells and TTR and PCSK9 in HEK293T cells.

a

Supplementary Figure 3. Screen for engineered xCas12i mutants with high-efficiency editing activity. a, The relative editing frequencies of over 500 rationally engineered xCas12i mutants.

Supplementary Figure 4. Other mutants mediated high-efficiency editing. a-b, xCas12i mutant with N243R increased 1.2, 5, 20-fold activity at DMD.1, DMD.2 and DMD.3 locus. **c,** Both Cas12-Max (xCas12i-N243R) and Cas12Max-E336R elevated EGFP-activated fluorescent at different PAM recognition sites.

Supplementary Figure 5. Cas12Max induced off-target editing efficiency at sites with mismatches using the reporter system (a) and targeted deep sequence (b).

Supplementary Figure 6. hfCas12Max mediates high-efficiency and -specificity editing. a, Rational protein engineering screen of over 200 mutants for highly-fidelity Cas12Max. Four mutants show significantly decreased activity at both OT (off-target) sites and retains at ON.1 (on-target) site. **b**, Different versions of xCas12i mutants. **c**, v6.3-hfCas12Max reduced off-target at OT.1 and OT.2 sites and retained indel activity at TTR-ON targets, compared to v1.1-Cas12-Max. **d**, v6.3-hfCas12Max exhibited comparable indel activity at DMD.1, DMD.2, and higher at DMD.3 locus, than v1.1-Cas12Max.

a

Supplementary Figure 7. hfCas12Max exhibited the high editing activity. a, Indel frenquencies from Cas12Max, hfCas12Max, LbCas12a, Ultra AsCas12a, SpCas9 and KKH-saCas9 at TTR locus.

Supplementary Figure 8. hfCas12Max mediates high-efficient and -specific editing. a-b, Off-target efficiency of hfCas12Max, LbCas12a, and UltraAsCas12a at *in-silico* predicted off-target sites, determined by targeted deep sequencing. Sequences of on-target and predicted off-target sites are shown, PAM sequences are in blue and mismatched bases are in red.

Supplementary Figure 9. Conserved cleavage sites of Cas12i. a, Sequence alignment of xCas12i, Cas12i1 and Cas12i2 shows that D650, D700, E875 and D1049 are conserved cleavage sites at RuvC domain. **b**, Introducing point mutations off D650A, E875A, and D1049A results in abolished activity of xCas12i.

Supplementary Figure 10. Other strategies for high-efficiency ABE-dxCas12i. a, TadA8e.1-dx-Cas12i-v1.2 and v1.3 exhibits significantly increased A-to-G editing activity among various variants at KLKF4 site of genome. **b**, Unchanged or even decreased editing activity from various dCas12-ABEs carrying different NLS at N-terminal. **c**, Increased A-to-G editing activity of TadA8e-dxCas12i-v4.3 by combining v2.2, changed-NLS linker and high-activity Tade8e. **d**, dxCas12i-ABE-N by TadA at the C-terminus of dCas12 slightly increased editing activity.

Supplementary Figure 11. Comparison of editing frequencies induced by various dCas12-BEs at different genomic target sites. a, Comparison of A-to-G editing frequencies induced by indicated TadA8e.1-dxCas12i, v1.2, v2.2, and TadA8e.1-dCas12a at PCSK9 genomic locus.