Not only perception but also grasping actions obey Weber's law

Zoltan Derzsi[1] and Robert Volcic[1,2]

[1] Division of Science, New York University Abu Dhabi, Abu Dhabi, United Arab Emirates

[2] Center for Artificial Intelligence and Robotics, New York University Abu Dhabi, Abu Dhabi, United

Arab Emirates

Contact information:

Dr. Zoltan Derzsi
E-mail: zd8@nyu.edu
 orcid.org/0000-0002-4253-4209

Dr. Robert Volcic
E-mail: robert.volcic@nyu.edu
 orcid.org/0000-0002-0170-4136

New York University Abu Dhabi
P.O. Box 129188, Abu Dhabi, UAE

Abstract

Weber's law, the principle that the uncertainty of perceptual estimates increases proportionally with object size, is regularly violated when considering the uncertainty of the grip aperture during grasping movements. The origins of this perception-action dissociation are debated and are attributed to various reasons, including different coding of visual size information for perception and action, biomechanical factors, the use of positional information to guide grasping, or, sensorimotor calibration. Here, we contrasted these accounts and compared perceptual and grasping uncertainties by asking people to indicate the visually perceived center of differently sized objects (Perception condition) or to grasp and lift the same objects with the requirement to achieve a balanced lift (Action condition). We found that the variability (uncertainty) of contact positions increased as a function of object size in both perception and action. The adherence of the Action condition to Weber's law and the consequent absence of a perception-action dissociation contradict the predictions based on different coding of visual size information and sensorimotor calibration. These findings provide clear evidence that human perceptual and visuomotor systems rely on the same visual information and suggest that the previously reported violations of Weber's law in grasping movements should be attributed to other factors.

Keywords: reaching, grasping, Weber's law, perception, action

Introduction

Weber's law, one of the fundamental principles of perception, states that sensory precision decreases proportionally with the magnitude of a stimulus, or expressed differently, that the uncertainty of stimulus estimates increases with stimulus magnitude [1]. Examples of Weber's law abound in all sensory modalities, with the exception of human grasping movements. When an object is about to be grasped, the variability in the maximum grip aperture between fingers is independent of object size, in contrast to the steady increase in variability with larger objects observed in perceptual estimates [2]. Although the finding that Weber's law is violated in visually guided grasping actions is robust, less so is the agreement about its origin.

According to the Two-Visual Systems (TVS) hypothesis, perception and action rely on different computations of visual size information [3–5]. Vision-for-perception, mediated by the ventral visual stream, is based on relative metrics, taking into consideration not only the object size, but also general aspects of the visual scene. In contrast, vision-for-action, served by the dorsal visual stream, is based on absolute metrics reflecting the actual size of the object, which then leads to the violation of Weber's law [2, 6]. A second account, the biomechanical factors (BF) hypothesis, instead assumes that perception and action rely on the same visual size information and it attributes the dissociation to non-visual sources of noise that contaminate only the variability of the finger aperture in grasping [7–10]. Weber's law in grasping is violated, because, as object size increases, the finger opening is limited more and more by biomechanical factors, which artificially reduce the variability of the maximum grip aperture for larger objects. A third account, the digits-in-space (DIS) hypothesis, suggests that the maximum grip aperture is not influenced by visual size information at all, but it is instead an emergent property of the way the fingers are independently moved toward their contact positions on the object [11–13]. If grasping is influenced by positional information, and not size, it follows that the maximum grip aperture variability should not adhere to Weber's law [14]. A fourth account, the sensorimotor calibration (SMC) hypothesis, states that the dissociation between perception and action arises from differences in the availability of sensory feedback [15–20]. Visual and haptic feedback, which is missing during perceptual estimations, can only calibrate grasping actions and, by this, reduce the variability of the maximum grip aperture [21, 22].

To address this contentious debate, we here introduce an experimental paradigm by which we pit the aforementioned hypotheses against each other by focusing on the precision with which contact positions are selected on an object, a crucial goal of any grasping movement. In the Perception condition, we

asked participants to indicate the visually perceived center of differently sized objects (Fig. 1A). In the Action condition, we asked participants to grasp and lift the same objects with the requirement to achieve a balanced lift (Fig. 1B). Thus, visual size information is essential in both conditions for an effective localization of the object center. The advantages of this paradigm are evident. First, by requiring the same grip aperture to grasp objects of all sizes we can avoid the potential constraints imposed by biomechanical factors. Second, we can evaluate grasping precision when the actual action goal is achieved and not ahead of it as it occurs when precision is measured at the time of the maximum grip aperture.

The fact that the variability of the perceptual estimates of the object center increases proportionally with size has been long known [23]. What is not yet known is whether and how does the variability of grasp contact positions depend on object size. If vision-for-action relies on absolute visual size information, as stated by the TVS hypothesis, we expect the grasping condition to violate Weber's law also in this paradigm (Fig. 2A, solid line in mid panel). On the contrary, if Weber's law is violated because of non-visual sources of noise (BF hypothesis), we expect the variability of the grasp contact positions to increase with object size with the same slope as in the perceptual condition, because biomechanical factors play no role here (Fig. 2A, long-dashed line in mid panel). Similarly, because the object center toward which fingers are steered can only be determined by considering the entire object size, the DIS hypothesis predicts for this paradigm that the variability of grasp contact positions should adhere to Weber's law (Fig. 2A, long-dashed line in mid panel). Lastly, if grasping movements are calibrated based on the visually perceived object rotation around the grasp axis and the haptically perceived tangential torques occurring whenever an object is grasped away from its center of mass (Fig. 1C), the SMC hypothesis predicts that the adherence to Weber's law should be at least reduced, if not completely eliminated (Fig. 2A, dotted line in mid panel). In this case, the sensorimotor calibration might even affect perception, if the perceptual estimates are taken just after grasping (Fig. 2A, dotted line in right panel).

Materials and Methods

Participants

Thirty-one naïve right-handed participants (age 19-35, 20 males) with normal or corrected-to-normal vision and no known history of neurological disorders participated in this study. They received subsistence allowance or course credit for their time. The study was undertaken with the understanding and informed

Figure 1: Methods. *A:* Perception condition: participants indicated the center of differently sized objects with their index finger, before (Pre) and after (Post) the Action condition. *B:* Action condition: participants grasped and lifted differently sized objects. The requirement was to achieve a balanced lift. *C:* Example of an off-center contact position. The position of the finger pad of the index finger was recorded with a virtual marker (gray point) with respect to four markers (orange points) attached on the distal phalanx of the index finger. Two markers (blue points) were attached on the edges of each object. The contact position (dotted line) was calculated with respect to the object center (dashed line). The circular dashed line represents the direction of object rotation and the tangential torque which provided visual and haptic feedback, respectively, whenever the object was lifted.

written consent of each participant and the experimental procedures were approved by the Institutional Review Board of New York University Abu Dhabi.

Apparatus

Participants sat at a table in a normally lit room with their right hand resting at a start position, aligned with their body midline. Stimuli objects were three 12×12 mm square-profile metal rods, 130 mm, 240 mm, and 350 mm in length, and they weighed 134 g, 256 g, and 382 g, respectively. The position of the index finger was recorded with sub-millimeter resolution by using an Optotrak Certus motion capture system (Northern Digital Inc., Waterloo, Ontario, Canada) controlled by the MOTOM toolbox [24]. Four infrared-light-emitting diodes (markers) were attached on the the distal phalanx of the index finger (Fig. 1C). The exact position of the index finger pad (virtual marker) was tracked by determining its location relative to the four markers via an assignment procedure performed at the beginning of the experiment. Two markers were attached on the edges of each object to track the position of the object center (Fig. 1C). A pair of occlusion goggles (Red Scientific, Salt Lake City, UT, USA) was used to prevent vision between trials. The experimental code was written in Matlab (The MathWorks, Inc., Natick, MA, USA).

Procedure

In the Perception condition, participants were asked to use their index finger to indicate the object center (Fig. 1A). In the Action condition, participants were asked to grasp the object with a thumb-index precision grip and lift it while maintaining horizontal balance (Fig. 1B). The Perception condition was performed twice: before (Pre) the Action condition to ensure that no visual or haptic feedback regarding the object true center location did influence perceptual estimates, and, after (Post) the Action condition to capture any effect of visual and haptic feedback on perceptual estimates.

Before each trial, participants placed their hand at the start position. The experimenter then placed one of the objects in front of the participant, 35 cm from the edge of the table and parallel to it. The position of the object was randomized in the range of ±15 cm along this line, to avoid participants reaching always to the same location in space. The trial started when the goggles turned transparent, and the position of all the markers was recorded by the experimenter the moment the task was complete. Participants then placed the object down (in the Action condition) and returned the hand back to the start position. After this, vision was occluded in preparation for the next trial. Each of the three objects was presented 24 times in each condition (216 trials per participant), for a total of 6696 trials.

Data analysis

All analyses were performed in R [25]. The position of each contact position was obtained by performing an orthogonal projection of the fingertip virtual marker onto the line defined by the two markers positioned on the edges of each object. The contact position was then defined relative to the center of the object, with negative and positive values indicating contact positions to the left and to the right of the object center. Trials in which any of the markers were not captured correctly were discarded from further analysis, the exclusion of these trials (72 trials, approximately 1% in total) left us with 6624 trials.

Contact position data were analyzed using a Bayesian mixed-effects distributional regression model, estimated using the *brms* package [26, 27]. Unlike conventional regression models that assume a constant standard deviation across observations, a distributional regression model estimates not only the location parameters, but also the standard deviations of the location parameters. Our model included as predictors the categorical variable Condition (Perception (Pre), Action, Perception (Post)) and the continuous variable Size, which was centered before being entered in the model. Thus, this model estimated the average contact position for each condition (contact position intercept), its change as a function of object

size for each condition (contact position slope), the average standard deviation of the contact position for each condition (standard deviation of contact position intercept), and the standard deviation change of the contact position as a function of object size for each condition (standard deviation of contact position slope), the key measure in this study. The model included also independent random effects for participants to capture the dependency among data due to the repeated measures design. This statistical method allow us to simultaneously model all the raw data of all participants and directly estimate the main variable of interest (changes in standard deviation as a function of object size), in contrast to the multi-step approach normally used in which standard deviations are first calculated separately for each participant, size and condition, then fitted with linear models to extract the participants' slopes, to finally pool them together to draw inferences.

Our model was fitted considering weakly informative prior distributions for each parameter to provide information about their plausible scale. We ran four Markov chains simultaneously, each for 4,000 iterations (1,000 warm-up samples) for a total of 12,000 post-warm-up samples. The obtained posterior distributions quantify the uncertainty about each estimated parameter conditional on the priors, model and data. We have summarized them by computing their means and their 95% credible intervals. To compare parameters between two conditions, we have computed the credible difference distributions by subtracting their respective posterior distributions, which were again summarized by computing their means and their 95% credible intervals.

Data availability

Data and analysis scripts are available at osf.io/e8c92/.

Results

Fig. 2B represents all the contact positions of the index finger relative to the center of each object in the Perception (Pre), Action and in the Perception (Post) conditions. These raw data clearly show that the contact positions were clustered around the center of each object, with no apparent systematic bias in either left (negative values) or right direction (positive values). The estimated average contact positions and their changes as a function of object size confirmed this observation; all posterior distributions of these estimates were centered on zero, equally extended in both positive and negative domains (contact position intercepts, Pre: -0.69 mm, 95% CI $= -1.78$, 0.41; Action: 0.09 mm, 95% CI $= -0.9$, 1.1; Post: 0.05 mm, 95% CI $= -1.27$, 1.36) and did not depend on object size (contact position slopes, Pre:

Figure 2: Predictions and results. *A:* Predicted SD as a function of size for the Two-visual systems (TVS), Biomechanical factors (BF), Digits-in-space (DIS), and Sensorimotor calibration (SMC) accounts in the Perception and Action conditions. *B:* Raw index finger contact positions relative to the center of each object. Points are slightly scattered horizontally for better visualization. *C:* Estimated SD as a function of size. Lines represent the mean and the shaded regions represent the 95% credible intervals of the posterior distributions of the Bayesian regression model. *D:* Estimated slopes of the SD as a function of object size for each condition. Points represent the mean and the error bars the 95% credible intervals of the posterior distributions. *E:* Differences between estimated slopes of the SD for the Action-Pre and Post-Pre comparisons. Points represent the mean and the error bars represent the 95% credible intervals of the differences between posterior distributions.

-0.003, 95% CI $= -0.009$, 0.004; Action: -0.0004, 95% CI $= -0.007$, 0.006; Post: 0.003, 95% CI $= -0.005$, 0.010).

The estimates of the standard deviation, the key measure used to compare the predictions made by the different hypotheses, are shown in Fig. 2C. The average standard deviations were of comparable magnitude in the three conditions (standard deviation of contact position intercepts, Pre: 4.08 mm, 95% CI = 3.70, 4.49; Action: 4.14 mm, 95% CI = 3.67, 4.66; Post: 4.41 mm, 95% CI = 3.93, 4.93). Most importantly, it is evident that the standard deviations increased proportionally for larger object sizes not only in the Perception conditions, but also in the Action condition, contrary to the predictions of the TVS and SMC accounts (compare with Fig. 2A). In adherence to Weber's law, the estimated slopes indicating the change in standard deviation of the contact positions as a function of object size (Pre: 0.016, Action: 0.014, Post: 0.016) and their 95% credible intervals were all positive and substantially above zero (Fig. 2D). No evidence was found of any difference in slopes between the Perception (Pre) and Action conditions, and, between the Perception (Post) and Perception (Pre) conditions. Both 95% credible intervals of the difference distributions overlapped with zero (Fig. 2E). Thus, both Perception and Action conditions conformed equally well to Weber's law, showing no dissociation, but also no sign of sensorimotor calibration and transfer from the Action to the Perception (Post) condition. Computing the average slopes using the more common multi-step approach produces nearly identical slope values relating the standard deviations of contact positions and object size (Pre: 0.017, SD = 0.008; Action: 0.015, SD = 0.008; Post: 0.017, SD = 0.007).

Discussion

The main aim of this study was to investigate Weber's law in relation to object size in both visual perception and visually guided grasping to address the debate about the origin of the perception-action dissociation. Specifically, we compared the uncertainty of perceptual estimates about the center of differently sized objects with the uncertainty with which optimal grasping contact positions are selected on those same objects, a crucial aspect of any visually guided grasping action. Our results show no sign of a perception-action dissociation and provide clear evidence that Weber's law affects not only the variability of perceptual estimates, but also the variability of visually guided grasping.

The finding that the variability of the contact points in the Action condition increased with object size to the same degree as in the perceptual condition is not consistent with the TVS hypothesis [2–6]. If vision-for-action were based on absolute metrics, we would have expected the variability in the Action condition to be constant over object sizes. This was clearly not the case, despite the fact that vision of the hand and the object was constantly available, haptic feedback was provided by touching and lifting

objects, and actions were directed toward real 3D objects. All these factors should have, if anything, favored the emergence of the perception-action dissociation. The range of sizes used for our objects was also not critical as similar ranges have been already used to study Weber's law [28, 29].

The present findings are also in contrast with the SMC hypothesis which predicted a reduction of the variability in the Action condition because of sensorimotor calibration, and, a potential transfer of this calibration on the perceptual estimates obtained just after the Action condition [15–20]. The precision of grasp contact positions definitely did not improve over successive repetitions even though inaccurate grasps provided direct visual and haptic feedback in the form of visually perceived object rotation and haptically perceived tangential torque which requires grip force adjustments to prevent rotational slip [30, 31]. One possible reason for the lack of calibration could be linked to the slow adaptation that the grip force undergoes in the presence of torque load which might, in this particular case, negatively affect the ability to use haptic inputs effectively [32]. Another reason that might have hindered or slowed down calibration was the randomized positioning of objects which required participants to assume slightly different arm postures for each grasping action.

Our study leads to an important implication about the role of Weber's law in grasping actions: the fundamental assumption of the TVS hypothesis that perception and action rely on different computations of the visual input can only hold if we accept that absolute metrics is used exclusively for shaping the grip aperture during grasping, but the selection of optimal grasp contact positions is based on relative metrics. Thus, a recourse to a proposal involving two separate visual streams does not seem to be essential to explain how vision guides grasping movements. Instead, future studies will need to further explore how does the actual grasping precision occasionally conflate with factors intrinsic to motor behavior, how do task constraints determine whether grasping movements are guided by size or positional information, and, and how does sensorimotor calibration affect grip apertures and optimal grasp contact positions over time.

Grants

This work was partially supported by the NYUAD Center for Artificial Intelligence and Robotics, funded by Tamkeen under the NYUAD Research Institute Award CG010.

References

1. Fechner, G. T. *Elemente der Psychophysik* (Breitkopf und Härtel, Leipzig, 1860).

2. Ganel, T., Chajut, E. & Algom, D. Visual Coding for Action Violates Fundamental Psychophysical Principles. *Current Biology* **18,** R599–601. doi:10.1016/j.cub.2008.04.052 (2008).

3. Goodale, M. A. & Milner, A. D. Separate Visual Pathways for Perception and Action. *Trends in Neurosciences* **15,** 20–25. doi:10.1016/0166-2236(92)90344-8 (1992).

4. Goodale, M. A. Transforming Vision into Action. *Vision Research* **51,** 1567–1587. doi:10.1016/j.visres.2010.07.027 (2011).

5. Goodale, M. A. How and Why the Visual Control of Action Differs from Visual Perception. *Proceedings of the Royal Society B: Biological Sciences* **281,** 20140337–20140337. doi:10.1098/rspb.2014.0337 (2014).

6. Ganel, T., Freud, E. & Meiran, N. Action Is Immune to the Effects of Weber's Law throughout the Entire Grasping Trajectory. *Journal of Vision* **14.** doi:10.1167/14.7.11 (2014).

7. Utz, K. S., Hesse, C., Aschenneller, N. & Schenk, T. Biomechanical Factors May Explain Why Grasping Violates Weber's Law. *Vision Research* **111,** 22–30. doi:10.1016/j.visres.2015.03.021 (2015).

8. Bruno, N., Uccelli, S., Viviani, E. & De'Sperati, C. Both Vision-for-Perception and Vision-for-Action Follow Weber's Law at Small Object Sizes, but Violate It at Larger Sizes. *Neuropsychologia* **91,** 327–334. doi:10.1016/j.neuropsychologia.2016.08.022 (2016).

9. Schenk, T., Utz, K. S. & Hesse, C. Violations of Weber's law tell us more about methodological challenges in sensorimotor research than about the neural correlates of visual behaviour. *Vision Research* **140,** 140–143. doi:10.1016/j.visres.2017.05.017 (2017).

10. Uccelli, S., Pisu, V. & Bruno, N. Precision in Grasping: Consistent with Weber's Law, but Constrained by "Safety Margins". *Neuropsychologia* **163,** 108088. doi:10.1016/j.neuropsychologia.2021.108088 (2021).

11. Smeets, J. B. J. & Brenner, E. A New View on Grasping. *Motor Control* **3,** 237–271. doi:10.1123/mcj.3.3.237 (1999).

12. Verheij, R., Brenner, E. & Smeets, J. B. J. Grasping kinematics from the perspective of the individual digits: a modelling study. *PLoS ONE* **7,** e33150–e33150. doi:10.1371/journal.pone.0033150 (2012).

13. Smeets, J. B. J., van der Kooij, K. & Brenner, E. A Review of Grasping as the Movements of Digits in Space. *Journal of Neurophysiology* **122,** 1578–1597. doi:10.1152/jn.00123.2019 (2019).

14. Smeets, J. B. J. & Brenner, E. Grasping Weber's Law. *Current Biology* **18,** R1089–R1090. doi:10.1016/j.cub.2008.10.008 (2008).

15. Bruno, N. & Franz, V. H. When Is Grasping Affected by the Müller-Lyer Illusion? A Quantitative Review. *Neuropsychologia* **47,** 1421–1433. doi:10.1016/j.neuropsychologia.2008.10.031 (2009).

16. Schenk, T. No Dissociation between Perception and Action in Patient DF When Haptic Feedback Is Withdrawn. *Journal of Neuroscience* **32,** 2013–2017. doi:10.1523/JNEUROSCI.3413-11.2012 (2012).

17. Bingham, G. P. & Mon-Williams, M. A. The Dynamics of Sensorimotor Calibration in Reaching-to-Grasp Movements. *Journal of Neurophysiology* **110,** 2857–2862. doi:10.1152/jn.00112.2013 (2013).

18. Volcic, R. & Domini, F. On-line visual control of grasping movements. *Experimental Brain Research* **234,** 2165–2177. doi:10.1007/s00221-016-4620-x (2016).

19. Volcic, R. & Domini, F. The Endless Visuomotor Calibration of Reach-to-Grasp Actions. *Scientific Reports* **8,** 14803. doi:10.1038/s41598-018-33009-6 (2018).

20. Cesanek, E., Campagnoli, C., Taylor, J. A. & Domini, F. Does Visuomotor Adaptation Contribute to Illusion-Resistant Grasping? *Psychonomic Bulletin & Review* **25,** 827–845. doi:10.3758/s13423-017-1368-7 (2018).

21. Davarpanah Jazi, S., Hosang, S. & Heath, M. Memory delay and haptic feedback influence the dissociation of tactile cues for perception and action. *Neuropsychologia* **71,** 91–100. doi:10.1016/j.neuropsychologia.2015.03.018 (2015).

22. Heath, M., Chan, J. & Davarpanah Jazi, S. Tactile-Based Pantomime Grasping: Knowledge of Results is Not Enough to Support an Absolute Calibration. *Journal of Motor Behavior* **51,** 10–18. doi:10.1080/00222895.2017.1408559 (2019).

23. Wolfe, H. K. On the Estimation of the Middle of Lines. *The American Journal of Psychology* **34,** 313–358. doi:10.2307/1413954 (1923).

24. Derzsi, Z. & Volcic, R. MOTOM Toolbox: MOtion Tracking via Optotrak and Matlab. *Journal of Neuroscience Methods* **308,** 129–134. doi:10.1016/j.jneumeth.2018.07.007 (2018).

25. R Core Team. *R: A Language and Environment for Statistical Computing* Vienna, Austria: R Foundation for Statistical Computing, 2022.

26. Bürkner, P.-C. brms: An R Package for Bayesian Generalized Linear Mixed Models Using Stan. *Journal of Statistical Software* **80,** 1–28. doi:10.18637/jss.v080.i01 (2017).

27. Carpenter, B. *et al.* Stan: A Probabilistic Programming Language. *Journal of Statistical Software* **76,** 1–32. doi:10.18637/jss.v076.i01 (2017).

28. Ganel, T., Namdar, G. & Mirsky, A. Bimanual Grasping Does Not Adhere to Weber's Law. *Scientific Reports* **7.** doi:10.1038/s41598-017-06799-4 (2017).

29. Hesse, C., Harrison, R. E., Giesel, M. & Schenk, T. Bimanual Grasping Adheres to Weber's Law. *i-Perception* **12,** 204166952110545. doi:10.1177/20416695211054534 (2021).

30. Kinoshita, H., Bäckström, L., Flanagan, J. R. & Johansson, R. S. Tangential Torque Effects on the Control of Grip Forces When Holding Objects With a Precision Grip. *Journal of Neurophysiology* **78,** 1619–1630. doi:10.1152/jn.1997.78.3.1619 (1997).

31. Goodwin, A. W., Jenmalm, P. & Johansson, R. S. Control of Grip Force When Tilting Objects: Effect of Curvature of Grasped Surfaces and Applied Tangential Torque. *Journal of Neuroscience* **18,** 10724–10734. doi:10.1523/JNEUROSCI.18-24-10724.1998 (1998).

32. Crevecoeur, F., Giard, T., Thonnard, J.-L. & Lefèvre, P. Adaptive Control of Grip Force to Compensate for Static and Dynamic Torques during Object Manipulation. *Journal of Neurophysiology* **106,** 2973–2981. doi:10.1152/jn.00367.2011 (2011).