1 **Conservation of *Nematocida* microsporidia gene expression and host response in**
2 ***Caenorhabditis* nematodes**
3

4 Yin Chen Wan[1], Emily R. Troemel[2], Aaron W. Reinke[1#]
5
6 [1] Department of Molecular Genetics, University of Toronto, Toronto, ON, Canada.
7 [2] School of Biological Sciences, University of California, San Diego, La Jolla, California, United
8 States of America.
9
10 #Author for Correspondence: Aaron W. Reinke, Department of Molecular Genetics, University
11 of Toronto, Toronto, ON, Canada, 416-946-0889, aaron.reinke@utoronto.ca

12
Abstract

Microsporidia are obligate intracellular parasites that are known to infect most types of
animals. Many species of microsporidia can infect multiple related hosts, but it is not known if
microsporidia express different genes depending upon which host species is infected or if the
host response to infection is specific to each microsporidia species. To address these
questions, we took advantage of two species of *Nematocida* microsporidia, *N. parisii* and *N.
ausubeli*, that infect two species of *Caenorhabditis* nematodes, *C. elegans* and *C. briggsae*.
We performed RNA-seq at several time points for each host infected with either microsporidia
species. We observed that *Nematocida* transcription was largely independent of its host. We
also observed that the host transcriptional response was similar when infected with either
microsporidia species. Finally, we analyzed if the host response to microsporidia infection was
conserved across host species. We observed that although many of the genes upregulated in
response to infection are not direct orthologs, the same expanded gene families are
upregulated in both *Caenorhabditis* hosts. Together our results describe the transcriptional
interactions of *Nematocida* infection in *Caenorhabditis* hosts and demonstrate that these
responses are evolutionarily conserved.

29
Key words: *Nematocida* microsporidia*, Caenorhabditis* nematodes*,* immunity, transcriptional
response, host-pathogen interactions

Significance statement

Microsporidia are a powerful model to study pathogen evolution, but much is still unknown
about how these pathogens have evolved to infect multiple host species. We found that
microsporidia express most of their genes similarly even when they are infecting different host
species and that related host species respond similarly to different microsporidia. Our results
suggests that there are conserved transcriptional responses during microsporidia infection.

Introduction
Microsporidia are obligate eukaryotic intracellular pathogens (Vávra and Lukeš, 2013). This
fungal-related phylum contains over 1400 described species that infect a wide range of animal
hosts including invertebrates, vertebrates, and protists (Stentiford *et al.*, 2016; Bojko *et al.*,
2022). Although as a phylum microsporidia infect a wide range of hosts, most species only
infect one or several closely related hosts (Murareanu *et al.*, 2021; Willis and Reinke, 2022).
Throughout evolution, microsporidia have lost many metabolic and biosynthesis genes which
are present in other eukaryotes (Nakjang *et al.*, 2013). These adaptations to survive within
their hosts have resulted in microsporidia having the smallest known eukaryotic genomes
(Katinka *et al.*, 2001). These features of microsporidia being able to specifically infect most
types of animals with a limited coding capacity have made microsporidia a powerful model for
understanding the evolution of intracellular parasites (James *et al.*, 2013; Haag *et al.*, 2014;
Quandt *et al.*, 2017; Wadi and Reinke, 2020).

51
The model nematode *Caenorhabditis elegans* has become an important system for studying
microsporidia infections (Tecle and Troemel, 2022). *Nematocida parisii* and *Nematocida*
ausubeli (referred to as *Nematocia* sp. 1 in some publications) are the two microsporidia
species most commonly observed to infect *Caenorhabditis elegans* (Zhang *et al.*, 2016).
These two species of *Nematocida* are also commonly found to infect *Caenorhabditis briggsae,*
which has been developed as a comparative species to *C. elegans* (Stein *et al.*, 2003). These
two nematode species live in distinct, but overlapping geographical locations (Barrière and
Félix, 2005). *N. parisii* was first found infecting *C. elegans* outside of Paris and *N. ausubeli*
was originally found infecting *C. briggsae* in India (Troemel *et al.*, 2008). Since then, both
microsporidia species have been found infecting both hosts in Europe (Zhang *et al.*, 2016).
There are some similarities of infection shared between these two *Nematocida* species; they
both exclusively infect the intestine, cause intestinal cells to fuse, and have similar life cycles
(Balla *et al.*, 2016). Both species also use similar types of secreted and membrane bound
proteins to interface with host proteins (Reinke *et al.*, 2017). There are also differences in their
infection characteristics, such as *N. ausubeli* displaying faster growth and increased
impairment of host fitness (Balla *et al.*, 2016).
68
In response to microsporidia infection, hosts often display large transcriptional changes. The
gene expression of *C. elegans* in response to *N. parisii* infection is reported to be distinct from
responses to extracellular pathogens, but similar to nodavirus infection (Bakowski *et al.*, 2014).
This transcriptional response has been termed the intracellular pathogen response (IPR)
(Reddy *et al.*, 2017). Among upregulated IPR genes, many contain F-box, FTH, and MATH
domains that are implicated in substrate recognition during ubiquitin-mediated degradation
(Thomas, 2006; Bakowski *et al.*, 2014). The IPR also includes upregulation of several
Caenorhabditis specific families that are still poorly understood, such as the PALS family
(Leyva-Díaz *et al.*, 2017). Mutants that cause the IPR to be upregulated are resistant to
infection and can clear *N. parisii* infections (Reddy *et al.*, 2019; Tecle *et al.*, 2021; Willis *et al.*,
2021).
80
Studies looking at infection of nematodes by different bacterial and fungal pathogens showed
both significant overlap between pathogen infections, as well as species-specific responses
(Wong *et al.*, 2007; Engelmann *et al.*, 2011; Lansdon, Carlson and Ackley, 2022). In addition,
recognition of two species of oomycete displayed a shared transcriptional response (Grover
et al., 2021). Similarities of responses between different host species have also been observed.
For example the response to nodavirus infection is conserved between *C. elegans* and *C.*
briggsae (Chen *et al.*, 2017). An intergenerational transcriptional response to *Pseudomonas*
vranovensis was conserved between some, but not all species of *Caenorhabditis* (Burton *et*
al., 2021). A study looking at different species of bacterial infection in the nematode
Pristionchus pacificus also showed both similarities and differences between the
transcriptional responses in the two hosts (Sinha *et al.*, 2012). Although conservation of
responses between pathogen and hosts in *C. elegans* is observed, different strains of bacteria
can elicit different responses, and different strains of *C. elegans* can have distinct responses
(Zárate-Potes *et al.*, 2020; Lansdon, Carlson and Ackley, 2022).
95
To determine the extent that microsporidia gene expression is influenced by its host and how
conserved the host response is to microsporidia infection, we generated and analysed
transcriptional data of *N. parisii* and *N. ausubeli* infecting *C. elegans* and *C. briggsae*. We
observe similar transcriptional patterns of each *Nematocida* species in the two species of
hosts with only a small set of differentially regulated genes. The host response to either of the
two microsporidia species was also similar. This transcriptional response is conserved across
host species. Altogether, our results suggest that different *Nematocida* species do not have
distinct expression programs depending on the host, and transcriptional responses of
Caenorhabditis hosts to *Nematocida* infection are conserved.
105

Results

Gene expression of *N. parisii* and *N. ausubeli* is similar in *C. elegans* and *C. briggsae*
To understand how gene expression of related microsporidia and hosts species is conserved, we infected *Caenorhabditis* nematodes with *Nematocida* microsporidia (fig. 1*A*). We designed our infection experiment to compare transcriptional differences between related microsporidia and host species (fig. 1*B*). Unlike previous transcriptional profiling experiments of *Nematocida* infection which were done as continuous infections (Bakowski *et al.*, 2014; Chen *et al.*, 2017), we infected the worms for a short period of time to synchronize the infection. We pulse-infected *C. elegans* and *C. briggsae* with either *N. parisii*, *N. ausubeli*, or a mock treatment for 2.5 hours, washed to remove spores from outside the worms, and then replated the animals for a total of 10, 20, or 28 hours of infection. Each condition was done once, except for the 10-hour time point which was performed in duplicate (fig. 1*A*). Samples were stained using a probe specific to the *Nematocida* 18S rRNA and we observed that greater than 75% of each population was infected (figs. 2*A* and *B*). Compared to *N. parisii* infected animals, we observed more parasite in *N. ausubeli* infected hosts at 28 hours, consistent with a previous report that *N. ausubeli* grows faster (Balla *et al.*, 2016). RNA from infected and uninfected animals was extracted and sequenced. To determine the expression of microsporidia genes, we mapped reads of *N. parisii* and *N. ausubeli* to their respective genomes (Supplementary table S1). At 10 hours post infection, the overall percentage of reads from either *N. parisii* or *N. ausubeli* were <1%, which increased to ~3-5% at 28 hours post infection (fig. 2*C*).

To define the transcriptional patterns of *N. parisii* and *N. ausubeli* during infection, we analysed RNA-seq results of each microsporidia species between *C. elegans* and *C. briggsae* at 10, 20, and 28 hours post infection. First, we used principal component analysis to compare microsporidia gene expression in each of these samples. We observed that at the 10-hour time point, there is a larger difference between expression in the two hosts, and at the later time points, expression in the two hosts are similar (figs. 3*A* and *B*). The 20- and 28-hour time points in *N. parisii* cluster closely together, but there is a larger difference between these time points in *N. ausubeli*, likely due to the accelerated growth of this species (Balla *et al.*, 2016). Moreover, we observed the expression of genes in each microsporidia species to be similar between the two *Caenorhabditis* hosts across the timepoints (figs. 3*C* and *D*, S1*A* and S1*B*). Strong correlation of *Nematocida* gene expression between *C. elegans* and *C. briggsae* was also observed, with similar levels of correlation of each microsporidia species in either host than between replicates in the same host (figs. 3*E* and *F*, S1*C* and S1*D*). These similarities between expression in different host species also did not change across time points as the parasite had gone through larger amounts of replication. At 10 hours post infection, over 45% of *Nematocida* genes are within 2-fold of each other, and over 60% are within four-fold of each other. As the time post infection increased to 28 hours, over 75% of *Nematocida* genes are within two-fold of each other, and more than 89% are within four-fold of each other. The similarity is even more pronounced in highly expressed genes (genes with greater than 100 FPKM in either host) at the 20- and 28-hour time point where 95-99% are within four-fold of each other.

To determine genes that are significantly differentially regulated depending upon the host species infected, we compared the 10-hour time points of each *Nematocida* species between the two hosts at 10 hours post infection (fig. 4*A* and *B*). We identified 34 differentially regulated genes in *N. parisii* and 11 in *N. ausubeli* (alpha< 0.05) (Supplementary table S1). Most of these differentially regulated genes had higher expression in *C. elegans* than *C. briggsae* (31/34 in *N. parisii* and 7/11 in *N. ausubeli*). Notably, differentially regulated genes were significantly enriched for ribosomal protein genes in both *N. parisii* (17/34, Fisher's exact test p-value=2.2×10^{-18}) and *N. ausubeli* (6/11, Fisher's exact test p-value=1.2×10^{-8}). Taken together, our results indicate that the transcriptional programs of *N. parisii* and *N. ausubeli* are largely independent of which *Caenorhabditis* hosts they infect.

161
162 **Shared and unique transcriptional responses of each *Caenorhabditis* host to *N. parisii***
163 **and *N. ausubeli* infection**
164
Different microsporidia species may induce similar or distinct transcriptional responses in the
same host. To address this question, we compared how either *C. elegans* or *C. briggsae*
responded to infection with either *N. parisii* or *N. ausubeli* at the 10-hour timepoint for which
we had replicate data. We first mapped the reads from each sample to the corresponding host
genome (Supplementary table S2). In *C. elegans*, we identified 875 genes that are significantly
differentially regulated in *N. parisii* infected samples and 735 genes that are significantly
differentially regulated in *N. ausubeli* infected samples (fig. 5*A-B* and Supplementary table S3).
For *C. briggsae,* 1091 genes are significant in *N. parisii* infected samples while 449 were
significant in *N. ausubeli* infected samples (fig. 5*C-D* and Supplementary table S3). There are
more downregulated than upregulated genes in each sample. Next, we directly compared
shared genes which are upregulated in each of the *Caenorhabditis* hosts when infected by *N.*
parisii or *N. ausubeli*. We observed significant overlap in the transcriptional response to these
infections with 222 upregulated genes shared in *C. elegans* and 117 upregulated genes
shared in *C. briggsae.*
179
We next compared both the overlapping and unique genes between samples to determine if
they are enriched in known biological functions. Using statistical enrichment tests, we
observed that the shared upregulated genes in *N. parisii* and *N. ausubeli* infected *C. elegans*
are enriched for GO Biological Process associated with metabolism; as well as GO Molecular
Process associated with catalytic activity (FDR<0.05, Supplementary table S4). We did not
observe any enrichment in the shared upregulated genes in *C. briggsae*, however the 267
shared downregulated *C. briggsae* genes showed enrichment for small molecule binding and
structural constituent of cuticle (Supplementary table S4). Of the non-shared genes, only the
415 downregulated genes in *C. briggsae* infected with *N. parisii* showed enrichment for GO
Molecular Process associated with structural constituent of cuticle and protein binding
(Supplementary table S4).
191
We then compared our RNA-seq data to previously published studies of the transcriptional
response to *N. parisii* infection (Supplementary table S5). Bakowski *et al.* performed RNA-seq
at multiple time points of a germline-deficient mutant of *C. elegans* continuously infected with
spores at the L3/L4 stage at 25°C. We compared our 10-hour time point to the five time points
in the Bakowski *et al.* study and observed strong overlap especially at the 8-, 16-, and 30-hour
time points (Supplementary fig. S2*A*). We observed significant overlap for all the time points
with both our *N. parisii* and *N. ausubeli* 10-hour post infection samples (fig. S2*B-C*). We also
compared our data to Chen *et al.* where *C. elegans* was infected with *N. parisii* at the L3 stage
at 20°C. We observed significant overlap with both our 10-hour *N. parisii* and *N. ausubeli* data
(fig. S2*D-E*). Chen *et al.* also measured Orsay virus infected *C. elegans* and Santeuil virus
infected *C. briggsae* and we saw significant similarities to these as well (fig. S2*E-D*). Together
our analysis demonstrates that despite the experimental differences between studies, a
largely similar upregulated host response is consistently observed.
205
206 **Evolutionary conserved host response to *Nematocida* infection**
207
To investigate the evolutionary conserved response of *Caenorhabditis* to *Nematocida* infection,
we studied the expression levels of *C. elegans* and *C. briggsae* orthologs. We identified
orthologs using Orthofinder (Emms and Kelly, 2019). Additionally, we determined the subset
of orthologs that only had a single copy present in each nematode species. Of the orthogroups
shared between *C. elegans* and *C. briggsae*, about 76% are single-copy orthologs
(10826/14183). We first clustered the expression of these one-to-one orthologs, which shows
a smaller cluster of upregulated genes and a larger cluster of downregulated genes (fig. 6*A*).
When we compared the host response to infection between the two host species, we observed

a modest but significant corelation of response to infection between *C. elegans* and *C.*
briggsae (fig. 6*B*). We then compared the statistical significance to the fold change of
differentially expressed genes from the different infection conditions (fig. 6*C-D*) and
investigated which of these differentially expressed genes were one-to-one orthologs (fig. 6*E-*
F). We observed that only 8-17% of the differentially upregulated genes were one-to-one
orthologs. Furthermore, of the most highly expressed genes (greater than four-fold), only 0-2%
were one-to-one orthologs. In contrast to less that 20% of significantly upregulated genes not
having one-to-one orthologs, between 29-42% of significantly downregulated genes did.
224
As much of the upregulated response is in the non-single copy orthologs, we analysed these
genes for commonalties between samples. We performed domain enrichment analysis to
determine the types of proteins induced in the non-single copy orthologs by microsporidia
infection. Previous transcriptional analysis of *C. elegans* infected with either *N. parisii* or Orsay
virus has identified serval types of domains enriched in expression, such as F-box,
MATH/BATH, PALS, DUF713, DUF684 and C-type lectins (Bakowski *et al.*, 2014; Chen *et al.*,
2017). In addition to these domains, we also included other types of domains that are enriched
in the significantly upregulated and downregulated non-single copy orthologs, for a total of
thirteen domains that we analysed (Supplementary table S6). Finally, the proportion of each
type of domain or gene family in the upregulated and downregulated non-single copy orthologs
was determined (fig. 7*A-B*). For *C. elegans* infected with *N. parisii*, we observed that ~29% of
the significantly expressed non-single copy orthologs contained at least one of these thirteen
domains, with genes containing PALS domains being the most common. For genes
upregulated after infection with *N. ausubeli*, we observed similar trends with ~35% containing
one of these domains. The downregulated genes showed similar domain patterns for both
microsporidia species, with cuticle collagen and glycosyltransferase being the two most
common domains. Similar patterns regarding types of domains in regulated genes induced by
the two microsporidia species was also observed in *C. briggsae* (fig. 7*B*).
243
As some of the same expanded gene families are upregulated in both *C. elegans* and *C.*
briggsae, we sought to determine the relationships between expression of these families in
response to infection. We first constructed gene trees of PALS and DUF713 families (figs.
S3-S4). Next, we compared the expression levels of these orthologs in our 10-hour samples.
This analysis shows that some phylogenetically related PALS and DUF713 genes are
upregulated in both *C. elegans* and *C. briggsae* in response to *Nematocida* infection (fig. 8*A-*
B).
251
Discussion
To understand the conservation of transcriptional responses during microsporidia infection,
we took the approach of comparing interactions between related microsporidia and host
species. To do this we, used a pair of *Nematocida* species that share ~66% amino acid identity
and have distinct growth and phenotypic characteristics during infection in *C. elegans* (Cuomo
et al., 2012; Balla *et al.*, 2016; Luallen *et al.*, 2016). We used a pair of *Caenorhabditis* species,
for which about 60% of their genes are orthologous and these proteins are 80% identical,
which is approximately the same extent of divergence between humans and mice (Stein *et al.*,
2003). We found that responses between *Caenorhabditis* and *Nematocida* species are largely
conserved. However, there are several limitations to our study. Although we monitored
infection at multiple time points, we only performed duplicates at 10 hours post infection.
Although a small number of replicates limit the ability to detect differentially regulated genes
(Conesa *et al.*, 2016), we saw a large overlap in significantly regulated genes compared to
other *N. parisii* infection datasets (Bakowski *et al.*, 2014; Chen *et al.*, 2017). Our study is also
limited to only two pairs of hosts and microsporidia species. Additionally, differences in
transcriptional responses between species could be dependent upon particular host and
microsporidia strains or environmental conditions.
269

Our results suggest that *Nematocida* microsporidia species do not sense and respond
differently depending upon their host environment. The main difference we observed in
Nematocida gene expression between the two hosts was an upregulation of both small and
large ribosomal subunit genes in *C. elegans* at 10 hours post infection. We also observed
about three-fold more ribosomal genes being differentially regulated between *N. parisii* and *N.*
ausubeli. After invasion both species undergo a lag period before starting replication. This lag
period is slightly shorter in *N. ausubeli* and the larger increase of ribosomal genes in *N. parisii*
is potentially related to these differences in lag time (Balla *et al.*, 2016). The lack of large
differences in the host response also suggests that these *Nematocida* microsporidia are not
differentially regulating host genes for their own benefit.
280
How hosts sense microsporidia infection is not fully understood. In mammals, several toll-like
receptors are necessary for immune activation (Tamim El Jarkass and Reinke, 2020), but the
proteins that *C. elegans* uses to sense and respond to microsporidia infection are mostly
unknown. Several negative regulators of the IPR are known, including PALS-22, LIN-35, and
PNP-1, but none of these are known to be necessary for the response to infection (Reddy *et*
al., 2017, 2019; Tecle *et al.*, 2021; Willis *et al.*, 2021). Recently, a basic-region leucine-zipper
transcription factor, ZIP-1, was found to positively regulate a subset of IPR genes (Lažetić *et*
al., 2022). A viral RNA receptor, DRH-1, is necessary for activation of the IPR by the Orsay
virus, but not for microsporidia infection (Sowa *et al.*, 2019). Whether there is an equivalent
protein that is specifically involved in detecting microsporidia infection in *C. elegans* is
unknown. Although a previous study showed that three IPR genes were induced to a lesser
extent by *N. ausubeli* than *N. parisii* (Zhang *et al.*, 2016) by comparing the full transcriptional
profile we see a largely similar set of genes being induced. As these different microsporidia
species induce a similar transcriptional response, this suggests that some common feature,
such as invasion, is detected by the host.
296
Nematodes, insects, and vertebrates all have strong transcriptional responses to infection by
microsporidia, though the responses appear to be quite diverse (Szumowski and Troemel,
2015; Midttun *et al.*, 2020). For example, antimicrobial peptides are observed to be
upregulated in silkworms and cytokines are induced in infected human cells (Ma *et al.*, 2013;
Flores *et al.*, 2021). The responses seen in these other animals are quite different than what
is observed in *Caenorhabditis*. One reason for this is that many IPR genes are part of large
gene families that are not conserved outside of *Caenorhabditis*, such as PALS (Leyva-Díaz *et*
al., 2017). Little is known about how a host responds to different species of microsporidia. A
study examining two species of microsporidia that infect the same mosquito found that a
horizontal transmitted species elicited a strong transcriptional immune response, but this
response was not enriched during infection with a vertically transmitted species (Desjardins *et*
al., 2015). Further studies will be necessary to know if the similar immune responses we
observe to infection by two horizontally transmitted microsporidia species are common in other
hosts infected by different microsporidia.
311
Microsporidia that infect free-living terrestrial nematodes appear to be common, and there are
other genera and families of nematodes that can be cultured in the laboratory and infected
with microsporidia. Additionally, there are other genera of microsporidia that have been shown
to infect *C. elegans* and other nematodes. Some nematode-infecting microsporidia also infect
multiple genera of nematodes, which could facilitate cross-species host expression (Zhang *et*
al., 2016). These types of broader comparisons would allow for a fuller view of how animals
evolve transcriptional responses to microsporidia infection.
319
320 **Material and Methods**
321
322 **Infection of nematodes with microsporidia**
323

C. elegans strain N2 and *C. briggsae* strain AF16 were maintained at 21°C on 10-cm
nematode growth medium (NGM) plates seeded with *Escherichia coli* OP50-1 for at least three
generations without starvation. Three plates each of mixed stage populations of animals were
washed with M9 and embryos extracted by treating with sodium hypochlorite/1 M NaOH for
2.5 minutes. Embryos were hatched by incubating for 18 hours at 21°C. ~30,000 L1s of each
species were placed on a 10-cm plates along with 1 ml M9 and 10 µl of 10X OP50-1. Three
plates of each species were prepared. Animals were infected with either 20 million *N. parisii*
(ERTm1) or *N. ausubeli* (ERTm2) spores. These spores were prepared as described
previously (Troemel *et al.*, 2008). Uninfected plates were used as the control. Plates were
dried for 1 hour in a clean cabinet and incubated an additional 1.5 hours at 21°C. Animals
were then removed from plates using two 5 ml washes of M9. After washing samples twice
with 10 ml M9, all but 1 ml of M9 was removed from the samples. An additional 3 ml of M9
and 1 ml of 10X OP50-1 was added to each sample, and 1 ml of this solution was plated onto
4 10-cm plates per sample, resulting in ~6,000 animals per plate. At either 10, 20, or 28 hours
post infection, plates were harvested by washing off worms 3 times with 725 µl M9. Samples
were then washed 3 times with 1 ml M9/0.1% Tween-20.
340
341   To determine the extent of infection, ~1,000 animals from each sample were removed, fixed,
342   and stained using an *N. parisii* 18S rRNA fluorescent in situ hybridization probe as previously
343   described (Reinke *et al.*, 2017). Vectashield containing DAPI (Vector Labs) was added and
344   samples imaged using an Zeiss Axioimager 2.
345
346   **RNA extraction and sequencing**
347   After the last wash, 1 ml of TRIzol was added to each sample and RNA was extracted similar
348   to as described previously (Bakowski *et al.*, 2014). mRNA libraries were prepared by the
349   UCSD IGM Genomics Center and sequenced on a single lane of an Illumina HiSeq 4000,
350   using 100 bp paired-end reads.
351
Microsporidia gene expression analysis
Reads from each sample were mapped to either *Nematocida parisii* strain ERTm1 (Genbank:
GCF_000250985.1) or *Nematocida ausubeli* strain ERTm2 (Genbank: GCA_000250695.1)
using TopHat v2.1.2 (Trapnell, Pachter and Salzberg, 2009). The respective *N. parisii* and *N.*
ausubeli genome annotation files were converted into .gtf format using Gffread v0.12.3.
Transcriptome assemblies were then generated using Cufflinks v2.2.1 (Trapnell *et al.*, 2010)
and Cuffmerge in the Cufflinks package. Differentially expressed genes were determined
using Cuffdiff v2.2.1 (Trapnell *et al.*, 2013) and were visualised using the R package
cummeRbund v2.36.0 (L. Goff, 2017).
361
The Fragments Per Kilobase of transcript per Million mapped reads (FPKM) values of *N. parisii*
and *N. ausubeli* transcripts in each *Caenorhabditis* host were calculated by Cuffdiff v2.2.1
(Trapnell *et al.*, 2013). The linear correlation between gene expression in each host was
determined using the R packages ggplot2 v3.3.5 and ggpubr v0.4.0; the ratio of FPKM values
between the two hosts was calculated by dividing the microsporidia's FPKM values in *C.*
elegans by that of *C. briggsae*'s. Hierarchical clustering of log10-transformed FPKM+1 values
was done using the hclust function from the R package stats v4.1.1; heatmaps were generated
by ggplot2 v3.3.5 in R.
370
***Caenorhabditis* RNA-seq analysis**
The paired end reads of each sample 10 hours post infection sample were submitted to Alaska
v1.7.2 (http://alaska.caltech.edu). Briefly, Bowtie2 (Langmead and Salzberg, 2012), Samtools
(Li *et al.*, 2009), RSeQC (Wang, Wang and Li, 2012), FastQC, and MultiQC (Ewels *et al.*, 2016)
were used for quality control of the input files. Then, Kallisto (Bray *et al.*, 2016) was used for
read alignment and quantification, followed by differential analysis using Sleuth (Pimentel *et*
al., 2017). For *C. elegans* samples, the reads were aligned to the N2 reference of the WS268

release (accession: PRJNA13758); the reads from *C. briggsae* samples were aligned to
reference genome of the WS268 release (accession: PRJNA10731).
380
Log2 fold change values of duplicated genes in each of the eight samples were averaged
using the R package dplyr v1.0.8. Genes with FDR-adjusted p-value of <0.05 were regarded
as significant. Differentially upregulated genes were defined as those with an FDR-adjusted
p-value<0.05 and log2 fold change ≥0 (infected vs. control); differentially downregulated
genes were defined as those with an FDR-adjusted p-value <0.05 and log2 fold change ≤0
(infected vs. control).
387
Principal component analysis
The normalised abundance measurements in *C. elegans* and *C. briggsae* generated by Alaska,
were read by the readRDS() function of the R package base v4.1.1. Normalized counts for
each gene in each host species were generated via sleuth_to_matrix() with the "obs_norm"
data and "tpm" units. PCA of gene expression was performed using samples from each
species via the R package pcaexplorer v2.20.2 (Marini and Binder, 2019).
394
Determination of gene expression overlap
Genes with an FDR<0.05 were used to compare expression overlap. Values of duplicated
genes in each sample were averaged using the R package dplyr v1.0.8. Overlap between
samples was determined using the R package gplots v3.1.1. The p-values were calculated
using the Fisher exact test. Statistical enrichment tests of shared and non-shared genes were
performed using PANTHER on pantherdb.org (Mi *et al.*, 2021). Each input list was statistically
tested for enrichment against the annotation sets: "GO biological process complete", "GO
cellular component complete", "GO molecular function complete", "PANTHER GO-Slim
Biological Process", "PANTHER GO-Slim Cellular Component", "PANTHER GO-Slim
Molecular Function", "PANTHER Pathways", "Reactome pathways" and "PANTHER Protein
Class". Results from these tests with p-value<0.05 are in Supplementary table S4.
406
To determine overlap in our samples compared to differentially regulated genes from
Bakowski *et al*., we first used Alaska to process the read files of *C. elegans* infected by *N.*
parisii samples at 8, 16, 30, 40 and 64 hours. To compare our samples' expression with data
from Bakowski *et al*., genes expressed in at least four of those samples were used for
hierarchical clustering by hclust function in the R package stats v4.1.1. The dendrogram in the
was produced using the R package ggdendrogram v0.1.23; the heatmap was generated by
ggplot2 v3.3.5. Overlap between our samples, the Bakowski *et al.,* (2014) samples, and the
Chen *et al.* (2017) samples containing differentially regulated genes for *C. elegans* N2 infected
by Orsay virus or *N. parisii*, and *C. briggsae* infected by Santeuil virus were calculated as
described above.
417
Domain enrichment analysis
The type of gene classes and domains investigated for the enrichment analyses are listed in
Supplementary table S6. Among these, we actively chose to look at F-box, MATH/BATH,
PALS, DUF684, DUF713, C-type lectins, and *skr* domains containing genes, implicated from
differentially regulated genes of *N. parisii* infected *C. elegans* in previous publications
(Bakowski *et al.*, 2014; Chen *et al.*, 2017). We also examined additional types of genes and
domains enriched in our datasets using DAVID Bioinformatics Resources (2021) (Huang, Brad
T. Sherman and Lempicki, 2009; Huang, Brad T Sherman and Lempicki, 2009) for *C. elegans*
and the WormBase Simple Gene Queries tool for *C. briggsae*. From this analysis, we found
additional domains and genes enriched with greater than three domain-containing proteins in
any of our samples. These identified domains are chitinase-like (*chil*) proteins, CUB and CUB-
like domains, cytochrome P450, glucosyltransferase family 92, nematode cuticle collagen N-
terminal domain, and UDP-glucuronosyltransferase. A list of gene names from respective
gene classes were downloaded separately from Wormbase (http://wormbase.org/) and the
corresponding protein-coding sequences were extracted from Wormbase ParaSite

(https://parasite.wormbase.org/). Alignments of other domains or classes were downloaded
directly from Pfam (http://pfam.xfam.org/) separately. After deleting duplicated sequences,
protein sequences of these classes of genes were converted to Stockholm format using
Clustal Omega Multiple Sequence Alignment tool (https://www.ebi.ac.uk/Tools/msa/clustalo/).
Hmmbuild of HMMER v3.3.2 (Eddy, 2011) was used to build respective profile HMMs, which
was then searched against *C. elegans* (accession: PRJNA13758) proteome of WS268 release
using hmmsearch. Output genes with E-value <1e-5 were used as the gene list for the
enrichment analyses. In the output genes, MATH and BATH genes were combined into one
list; *fbxa*, *fbxb* and *fbxc* genes were merged into the gene list for F-box while genes with CUB
and CUB-like domains were categorised together (Supplementary table S6). Using the R
package gplots v3.1.1, the number of genes overlapping between the lists and samples were
computed, then plotted using R package ggplot2 v3.3.5. Genes that do not fall into any of
these gene classes or have any of these domains were categorized as "other".
446

447   **Determination of orthologs between *C. elegans* and *C. briggsae***
448   Orthologous genes were determined between *C. elegans* (accession: PRJNA13758) and *C.*
449   *briggsae* (accession: PRJNA13758) proteomes of the WS268 release using Orthofinder v2.5.2
450   with the default settings (Emms and Kelly, 2019).
451

Single copy orthologs analysis
Genes that are single copy orthologs were extracted, based on the single copy orthologs list
computed by Orthofinder. Single copy orthologs expressed in at least one out of the four
samples were used for hierarchical clustering via the function hclust() from the R package
stats v4.1.1, then plotted using R packages ggdendrogram and ggplot2 v3.3.5. The linear
correlation of each single copy ortholog expression between the two host was calculated using
ggpubr v0.4.0. The scatterplots and volcano plots were generated by ggplot2 v3.3.5.
459

Comparison of gene expression in expanded gene families
The *C. elegans* and *C. briggsae* genes in PALS and DUF713 families were obtained from the
output of HMMER v3.3.2 (Eddy, 2011) using a E-value threshold of <1e-5. The protein
sequences of the genes were obtained from Wormbase ParaSite
(https://parasite.wormbase.org/) and aligned by M-Coffee using default settings (Notredame,
Higgins and Heringa, 2000; Di Tommaso *et al.*, 2011). A phylogenetic tree for each family was
generated using MrBayes (Huelsenbeck and Ronquist, 2001; Ronquist and Huelsenbeck,
2003) with the following settings: lset nst = 1, rates = invgamma, nruns = 2, stoprule = YES,
stopval =0.05 and mcmcdiagn=YES. A dendrogram of the phylogenetic tree of each gene
family was created using the function ReadDendrogram from the R package DECIPHER
v2.22.0. For each gene family, the orthologs' log2 fold change values across the *C. elegans*
and *C. briggsae* samples were plotted as a heatmap using ggplot2 v3.3.5.
472

473   **Data availability**
474   All samples were deposited under NCBI BioProject PRJNA841614 and the sequence reads
475   for all samples were submitted to the NCBI Sequence Read Archive.
476

Acknowledgements
We thank Hala Tamim El Jarkass and Meng A. Xiao for providing helpful comments on the
manuscript. We thank Winnie Zhao for technical help for the comparisons to the Bakowski
data. This work was supported by a Canadian Institutes of Health Research grant no. 400784
(to A.R.), an Alfred P. Sloan Research Fellowship FG2019-12040 (to A.R.), and NIH R01's
AG052622 and GM114139 to E.T. Some strains were provided by the CGC, which is funded
by NIH Office of Research Infrastructure Programs (P40 OD010440) and we thank WormBase.
484

485   **Author contributions:** Y.C.W. and A.R. conceived of the project. Y.C.W. performed all
486   computational analyses. A. R. performed all experiments. E.T. and A. R. provided mentorship
487   and acquisition of funding. Y.C.W. and A.R. co-wrote the paper with edits from E.T.

488
489 **Competing Interests:** The authors declare that they have no competing interests.
490
491 **Figures and legends**
492

Fig. 1. Overview of study design. (*A*) Schematic diagram of the RNA-seq experiment. *C. elegans* and *C. briggsae* were either not infected or infected with *N. parisii* or *N. ausubeli* separately at the L1 stage for two and a half hours. Next, animals were washed to remove any remaining microsporidia spores and replated. Worms were harvested for RNA-sequencing at

498　　10, 20 and 28 hours after infection. (*B*) The three comparisons of transcriptional responses
499　　made in this study are represented. Comparison 1 is microsporidia gene expression between
500　　hosts, comparison 2 is the response of each host species to different microsporidia species,
501　　and comparison 3 is the conserved response between different hosts to each microsporidia
502　　species. Figure was created with biorender (www.biorender.com).
503

504
505
Fig. 2. *Nematocida* infection of *Caenorhabditis* hosts. (A) Pulse-infected *C. elegans* or *C.*
briggsae were fixed and then stained with DAPI to detect nuclei and a FISH probe specific to
Nematocida 18S rRNA to detect either *N. parisii* or *N. ausubeli* at 10, 20 and 28 hours after

infection. Representative images of infection are shown. Scale bars are 23 µm. (B) Percentage
of infected animals at different timepoints. Between 39 to 220 animals were counted for each
sample. (C) Overall read mapping rate of *Nematocida* in each host at different time points. R1
or R2 indicates the replicate samples at the 10-hour timepoint.
513

514
515
Fig. 3. Microsporidia gene expression in different host species. Principal component
analysis of (A) *N. parisii* and (B) *N. ausubeli* in *C. elegans* and *C. briggsae* at 10, 20 and 28
hours. Circles represent confidence ellipses around each strain at 95% confidence interval.

(C-D) Heatmap of transcriptional profiles of genes expressed in *N. parisii* (C) and *N. ausubeli*
(D). Rows represent gene clustered hierarchically. Scale is of differential regulation of infected
compared to uninfected samples. (E-F) Scatterplot of log10 FPKM values in *N. parisii* (E) and
N. ausubeli (F) when infecting *C. briggsae* and *C. elegans*. Each point represents a
microsporidian gene. Pearson correlation value and p-value are indicated on the top left of
each plot. The ratio of FPKM values for each gene between *C. elegans* and *C. briggsae* is
demonstrated by the colour of the points, which is described in the legend at the bottom.
526
527

Fig. 4. Differentially regulated genes of microsporidia. (A-B) Line plots of the FPKM values
with confidence intervals, calculated for the differentially regulated genes (FDR<0.05) in *N.*
parisii (A) and *N. ausubeli* (B) at 10 hours post infection. * indicates ribosomal protein-
encoding genes. The aggregate sample value of the two replicates is noted as a black dot
within the confidence interval, while the value for each replicate is shown as a dot of matching
colour along the confidence interval. Error bars of each gene show the variability of the
distribution of FPKM values.
536
537

538
539
540 **Fig. 5. Transcriptional responses of *Caenorhabditis* hosts to *Nematocida* infection.** (A-
541 D) Venn diagrams of genes in *C. elegans* (A-B) or *C. briggsae* (C-D) that are upregulated (A
542 and C) or downregulated (B and D) when infected by *N. parisii* or *N. ausubeli*. npa (*N. parisii)*
543 and nau (*N. ausubeli*).
544
545

546
Fig. 6. Transcriptional responses of single copy orthologs in *Caenorhabditis* hosts to
***Nematocida* infection.** (A) Heatmap of transcriptional profiles of single-copy orthologs in *C.*
elegans and *C. briggsae* after infection by *N. parisii* or *N. ausubeli*. Rows represent genes
clustered hierarchically. Scale is of differential regulation of infected compared to uninfected

samples. **(B)** Scatterplot of log2 fold change values of single-copy orthologs expressed in *C.*
elegans and *C. briggsae*. Each point represents a single-copy ortholog between the two host
species. Pearson correlation values and p-values are indicated on top left of each plot. The
ratio of log2 fold change values for each single copy ortholog between *C. elegans* and *C.*
briggsae is demonstrated by the colour of the points. (C-D) Volcano plots of genes expressed
in *Nematocida* infected *C. elegans* (C) and *C. briggsae* (D). Upregulated genes are
represented with orange points; downregulated genes are represented with pink points. The
top 5 upregulated and downregulated genes are labelled. (E-F) Volcano plots of genes
expressed in *Nematocida* infected *C. elegans* (E) and *C. briggsae* (F). Single-copy orthologs
are represented with red points and non-single copy orthologs represented with grey points.
npa (*N. parisii)* and nau (*N. ausubeli*).
562
563

Fig. 7. Transcriptional responses of non-single copy orthologs in *Caenorhabditis* hosts to *Nematocida* infection. Domain enrichment analysis of significantly upregulated and downregulated non-single copy ortholog genes in *C. elegans* (A) and *C. briggsae* (B). npa (*N. parisii)* and nau (*N. ausubeli*). Upregulated samples are represented with orange text; downregulated samples are represented with pink text.

571
572
573 **Fig. 8. Conserved response of *Caenorhabditis* hosts to *Nematocida* infection.** (A-B)
574 Heatmap of transcriptional profiles of differentially regulated genes (FDR<0.05) in the PALS
575 (A) and DUF713 (B) gene families. Scale is of differential regulation of infected compared to
576 uninfected samples. npa (*N. parisii)* and nau (*N. ausubeli*).
577

579
Fig. S1. Transcriptional response of *Nematocida* species in *Caenorhabditis* hosts. *(A-*
B) Scatterplot of *N. parisii* (*A*) and *N. ausubeli* (*B*) log10 FPKM values between *C. briggsae*
and *C. elegans* replicates at 10 hours. Pearson correlation values and p-values are indicated

on the top left of each plot. The ratio of FPKM values for each gene between the replicates is
demonstrated by the colour of the points. (*C-D*) Gene expression pattern of *N. parisii (C)* and
N. ausubeli (D) in *C. elegans* and *C. briggsae*.
586
587

A

log2 Fold Change

-5.0 -2.5 0.0 2.5 5.0

Bakowski 8hr · Bakowski 16hr · Bakowski 30hr · Bakowski 40hr · Bakowski 64hr · C. elegans–npa 10hr · C. elegans–nau 10hr

B

3.19e-75 — 266 | 66 | 98 — C. elegans-npa / Bakowski 8hr

7.77e-61 — 289 | 43 | 24 — C. elegans-npa / Bakowski 16hr

1.43e-23 — 275 | 57 | 654 — C. elegans-npa / Bakowski 30hr

4.18e-31 — 288 | 44 | 224 — C. elegans-npa / Bakowski 40hr

6.49e-17 — 301 | 31 | 252 — C. elegans-npa / Bakowski 64hr

C

1.51e-89 — 255 | 74 | 90 — C. elegans-nau / Bakowski 8hr

5.13e-61 — 286 | 43 | 24 — C. elegans-nau / Bakowski 16hr

1.44e-29 — 265 | 64 | 647 — C. elegans-nau / Bakowski 30hr

7.71e-27 — 289 | 40 | 228 — C. elegans-nau / Bakowski 40hr

8.23e-22 — 293 | 36 | 247 — C. elegans-nau / Bakowski 64hr

D

3.55e-89 — 256 | 76 | 109 — C. elegans-npa / Chen npa 12hr

2.72e-65 — 277 | 55 | 74 — C. elegans-npa / Chen orsay 12hr

E

4.81e-93 — 251 | 78 | 107 — C. elegans-nau / Chen npa 12hr

7.63e-71 — 271 | 58 | 71 — C. elegans-nau / Chen orsay 12hr

F

1.44e-42 — 356 | 53 | 168 — C. briggsae-npa / Chen santeuil 12hr

4.39e-37 — 118 | 35 | 186 — C. briggsae-nau / Chen santeuil 12hr

588
Fig. S2. Data from this study has high overlap with previously published intracellular
infection samples. (*A*) Heatmap of transcriptional profiles of differentially regulated genes
(FDR<0.05) across five samples from Bakowski *et al*. and our 10hr *N. parisii* and *N. ausubeli*
infected *C. elegans* samples. Only genes expressed in at least four out of the eight total

samples are included. *(B-C)* Statistical overlap of significant genes between respective *N.*
parisii or *N. ausubeli* infected *C. elegans* samples and Bakowski *et al.* samples across five
timepoints. *(D-E)* Statistical overlap of significant genes between *N. parisii (D)* or *N. ausubeli*
(E) infected *C. elegans* samples and Chen *et al*. Orsay virus infected *C. elegans* sample at 12
hours. *(F)* Statistical overlap of significant genes between respective *N. parisii* or *N. ausubeli*
infected *C. briggsae* sample and Chen *et al*. Santeuil virus infected *C. briggsae* sample at 12
hours. The p-value of each comparison is indicated on top of each Venn diagram. npa *(N.*
parisii) and nau *(N. ausubeli)*.

601
602

603
604   **Fig. S3. Phylogram of *C. elegans* and *C. briggsae* PALS family genes.** Genes with the
605   same name that end with different letters indicate protein isoforms. Node values indicate
606   posterior probabilities for each split as percentage. The scale bar indicates average branch
607   length measured in expected substitutions per site.
608
609

610
611 **Fig. S4. Phylogram of *C. elegans* and *C. briggsae* DUF713 family genes.** Genes with the
612 same name that end with different letters indicate protein isoforms. Node values indicate
613 posterior probabilities for each split as percentage. The scale bar indicates average branch
614 length measured in expected substitutions per site.
620 **Supplementary Table and Materials**
621 **Supplementary Table S1. FPKM values of gene expression and differentially expressed**
622 **microsporidia genes.**
623
624 **Supplementary Table S2. Normalised RNA-seq counts of the *C. elegans* and *C. briggsae***
625 **for the transcriptional analysis generated by Alaska.**
626
627
628 **Supplementary Table S3. Differentially expressed genes of *C. elegans* and *C. briggsae***
629 **exposed to the two species of microsporidia generated by Alaska.**
630
Supplementary Table S4. PANTHER GO enrichment analyses. Sheet S1 contains results
of the statistical enrichment tests of the 222 significantly upregulated genes shared between
N. parisii and *N. ausubeli* infected *C. elegans*. Sheet S2 contains results of the statistical
enrichment tests of the 267 significantly downregulated genes shared between *N. parisii* and
N. ausubeli infected *C. briggsae*. Sheet S3 contains results of the statistical enrichment tests
of the 415 significantly downregulated genes specifically in *N. parisii* infected *C. briggsae*.
637
638 **Supplementary Table S5. List of gene set overlaps with previously published infection**
639 **samples.**

642 **Supplementary Table S6. Gene classes and domains used for enrichment analyses.**

Domain/ Gene Family	Databse	Wormbase gene class OR Pfam ID
F-box	Wormbase	fbxa, fbxb, fbxc
MATH (meprin or Traf homology) or BATH (BTB and MATH domain-containing)	Wormbase	math, bath
PALS (protein containing ALS2CR12 signature)	Wormbase	pals
C-type lectins	Wormbase	clec
DUF713	Pfam	PF015218
DUF684	Pfam	PF05075
chil	Wormbase	chil
Nematode cuticle collagen N-terminal domain	Pfam	PF01484
CUB and CUB-like	Pfam	PF00431, PF02408
Cytochrome P450	Wormbase	cyp
Glucosyltransferase family 92	Pfam	PF00201
UDP-glucuronosyltransferase	Wormbase	ugt
skr (Skp1-related)	Wormbase	Skr
Other (Any genes not in the domain or gene family above)	-	-

References

Bakowski, M.A. *et al.* (2014) 'Ubiquitin-Mediated Response to Microsporidia and Virus

Infection in C. elegans', *PLoS Pathogens*. Edited by D.S. Schneider, 10(6), p. e1004200.

doi:10.1371/journal.ppat.1004200.

Balla, K.M. *et al.* (2016) 'Cell-to-cell spread of microsporidia causes Caenorhabditis elegans

organs to form syncytia', *Nature Microbiology*, 1(11), p. 16144.

doi:10.1038/nmicrobiol.2016.144.

Barrière, A. and Félix, M.-A. (2005) 'High Local Genetic Diversity and Low Outcrossing Rate in

Caenorhabditis elegans Natural Populations', *Current Biology*, 15(13), pp. 1176–1184.

doi:10.1016/j.cub.2005.06.022.

Bojko, J. *et al.* (2022) 'Microsporidia: a new taxonomic, evolutionary, and ecological

synthesis', *Trends in Parasitology*, 0(0). doi:10.1016/j.pt.2022.05.007.

Bray, N.L. *et al.* (2016) 'Near-optimal probabilistic RNA-seq quantification', *Nature*

Biotechnology, 34(5), pp. 525–527. doi:10.1038/nbt.3519.

Burton, N.O. *et al.* (2021) 'Intergenerational adaptations to stress are evolutionarily

conserved, stress-specific, and have deleterious trade-offs.', *eLife*, 10.

doi:10.7554/eLife.73425.

Chen, K. *et al.* (2017) 'An evolutionarily conserved transcriptional response to viral infection

in Caenorhabditis nematodes', *BMC Genomics*, 18(1), p. 303. doi:10.1186/s12864-017-3689-

3.

Conesa, A. *et al.* (2016) 'A survey of best practices for RNA-seq data analysis', *Genome*
Biology, 17(1), p. 13. doi:10.1186/s13059-016-0881-8.

Cuomo, C.A. *et al.* (2012) 'Microsporidian genome analysis reveals evolutionary strategies
for obligate intracellular growth', *Genome Research*, 22(12), pp. 2478–2488.
doi:10.1101/gr.142802.112.

Desjardins, C.A. *et al.* (2015) 'Contrasting host-pathogen interactions and genome evolution
in two generalist and specialist microsporidian pathogens of mosquitoes.', *Nature*
communications, 6, p. 7121. doi:10.1038/ncomms8121.

Di Tommaso, P. *et al.* (2011) 'T-Coffee: a web server for the multiple sequence alignment of
protein and RNA sequences using structural information and homology extension', *Nucleic*
Acids Research, 39(suppl), pp. W13–W17. doi:10.1093/nar/gkr245.

Eddy, S.R. (2011) 'Accelerated Profile HMM Searches', *PLOS Computational Biology*, 7(10), p.
e1002195. doi:10.1371/journal.pcbi.1002195.

Emms, D.M. and Kelly, S. (2019) 'OrthoFinder: phylogenetic orthology inference for
comparative genomics', *Genome Biology*, 20(1), p. 238. doi:10.1186/s13059-019-1832-y.

Engelmann, I. *et al.* (2011) 'A comprehensive analysis of gene expression changes provoked
by bacterial and fungal infection in C. elegans.', *PloS one*, 6(5), p. e19055.
doi:10.1371/journal.pone.0019055.

Ewels, P. *et al.* (2016) 'MultiQC: summarize analysis results for multiple tools and samples in
a single report', *Bioinformatics*, 32(19), pp. 3047–3048. doi:10.1093/bioinformatics/btw354.

Flores, J. *et al.* (2021) 'Human microsporidian pathogen Encephalitozoon intestinalis
impinges on enterocyte membrane trafficking and signaling', *Journal of Cell Science*, 134(5),
p. jcs253757. doi:10.1242/jcs.253757.

Grover, M. *et al.* (2021) 'Infection of C. elegans by Haptoglossa Species Reveals Shared
Features in the Host Response to Oomycete Detection', *Frontiers in Cellular and Infection*
Microbiology, 11, p. 973. doi:10.3389/fcimb.2021.733094.

Haag, K.L. *et al.* (2014) 'Evolution of a morphological novelty occurred before genome
compaction in a lineage of extreme parasites', *Proceedings of the National Academy of*
Sciences, 111(43), pp. 15480–15485. doi:10.1073/pnas.1410442111.

Huang, D.W., Sherman, Brad T. and Lempicki, R.A. (2009) 'Bioinformatics enrichment tools:
paths toward the comprehensive functional analysis of large gene lists', *Nucleic Acids*
Research, 37(1), pp. 1–13. doi:10.1093/nar/gkn923.

Huang, D.W., Sherman, Brad T and Lempicki, R.A. (2009) 'Systematic and integrative analysis
of large gene lists using DAVID bioinformatics resources', *Nature Protocols*, 4(1), pp. 44–57.
doi:10.1038/nprot.2008.211.

Huelsenbeck, J.P. and Ronquist, F. (2001) 'MRBAYES: Bayesian inference of phylogenetic
trees', *Bioinformatics*, 17(8), pp. 754–755. doi:10.1093/bioinformatics/17.8.754.

James, T.Y. *et al.* (2013) 'Shared Signatures of Parasitism and Phylogenomics Unite
Cryptomycota and Microsporidia', *Current Biology*, 23(16), pp. 1548–1553.
doi:10.1016/j.cub.2013.06.057.

Katinka, M.D. *et al.* (2001) 'Genome sequence and gene compaction of the eukaryote
parasite Encephalitozoon cuniculi.', *Nature*, 414(6862), pp. 450–453. doi:10.1038/35106579.

L. Goff, C.T. (2017) *Cummerbund: analysis, exploration, manipulation, and visualization of*
cufflinks high-throughput sequencing data. Bioconductor.
doi:10.18129/B9.BIOC.CUMMERBUND.

Langmead, B. and Salzberg, S.L. (2012) 'Fast gapped-read alignment with Bowtie 2', *Nature*
Methods, 9(4), pp. 357–359. doi:10.1038/nmeth.1923.

Lansdon, P., Carlson, M. and Ackley, B.D. (2022) 'Wild-type Caenorhabditis elegans isolates
exhibit distinct gene expression profiles in response to microbial infection', *BMC Genomics*,
23(1), p. 229. doi:10.1186/s12864-022-08455-2.

Lažetić, V. *et al.* (2022) 'The transcription factor ZIP-1 promotes resistance to intracellular
infection in Caenorhabditis elegans', *Nature Communications*, 13(1), p. 17.
doi:10.1038/s41467-021-27621-w.

Leyva-Díaz, E. *et al.* (2017) 'Silencing of Repetitive DNA Is Controlled by a Member of an
Unusual Caenorhabditis elegans Gene Family', *Genetics*, 207(2), pp. 529–545.
doi:10.1534/genetics.117.300134.

Li, H. *et al.* (2009) 'The Sequence Alignment/Map format and SAMtools', *Bioinformatics*,
25(16), pp. 2078–2079. doi:10.1093/bioinformatics/btp352.

Luallen, R.J. *et al.* (2016) 'Discovery of a Natural Microsporidian Pathogen with a Broad
Tissue Tropism in Caenorhabditis elegans', *PLoS pathogens*, 12(6), p. e1005724.
doi:10.1371/journal.ppat.1005724.

Ma, Z. *et al.* (2013) 'Genome-Wide Transcriptional Response of Silkworm (Bombyx mori) to
Infection by the Microsporidian Nosema bombycis', *PLOS ONE*, 8(12), p. e84137.
doi:10.1371/journal.pone.0084137.

Marini, F. and Binder, H. (2019) 'pcaExplorer: an R/Bioconductor package for interacting
with RNA-seq principal components', *BMC Bioinformatics*, 20(1), p. 331.
doi:10.1186/s12859-019-2879-1.

Mi, H. *et al.* (2021) 'PANTHER version 16: a revised family classification, tree-based
classification tool, enhancer regions and extensive API', *Nucleic Acids Research*, 49(D1), pp.
D394–D403. doi:10.1093/nar/gkaa1106.

Midttun, H.L.E. *et al.* (2020) 'Effects of Pseudoloma neurophilia infection on the brain
transcriptome in zebrafish (Danio rerio)', *Journal of Fish Diseases*, 43(8), pp. 863–875.
doi:10.1111/jfd.13198.

Murareanu, B.M. *et al.* (2021) 'Generation of a Microsporidia Species Attribute Database
and Analysis of the Extensive Ecological and Phenotypic Diversity of Microsporidia', *mBio*.

Nakjang, S. *et al.* (2013) 'Reduction and expansion in microsporidian genome evolution: new
insights from comparative genomics.', *Genome biology and evolution*, 5(12), pp. 2285–2303.
doi:10.1093/gbe/evt184.

Notredame, C., Higgins, D.G. and Heringa, J. (2000) 'T-coffee: a novel method for fast and
accurate multiple sequence alignment 1 1Edited by J. Thornton', *Journal of Molecular
Biology*, 302(1), pp. 205–217. doi:10.1006/jmbi.2000.4042.

Pimentel, H. *et al.* (2017) 'Differential analysis of RNA-seq incorporating quantification
uncertainty', *Nature Methods*, 14(7), pp. 687–690. doi:10.1038/nmeth.4324.

Quandt, C.A. *et al.* (2017) 'The genome of an intranuclear parasite, Paramicrosporidium
saccamoebae, reveals alternative adaptations to obligate intracellular parasitism', *eLife*.
Edited by A. Rokas, 6, p. e29594. doi:10.7554/eLife.29594.

Reddy, K.C. *et al.* (2017) 'An Intracellular Pathogen Response Pathway Promotes
Proteostasis in C. elegans', *Current biology: CB*, 27(22), pp. 3544-3553.e5.
doi:10.1016/j.cub.2017.10.009.

Reddy, K.C. *et al.* (2019) 'Antagonistic paralogs control a switch between growth and
pathogen resistance in C. elegans', *PLoS pathogens*, 15(1), p. e1007528.
doi:10.1371/journal.ppat.1007528.

Reinke, A.W. *et al.* (2017) 'Identification of microsporidia host-exposed proteins reveals a
repertoire of rapidly evolving proteins.', *Nature communications*, 8, p. 14023.
doi:10.1038/ncomms14023.

Ronquist, F. and Huelsenbeck, J.P. (2003) 'MrBayes 3: Bayesian phylogenetic inference
under mixed models', *Bioinformatics*, 19(12), pp. 1572–1574.
doi:10.1093/bioinformatics/btg180.

Sinha, A. *et al.* (2012) 'System Wide Analysis of the Evolution of Innate Immunity in the
Nematode Model Species Caenorhabditis elegans and Pristionchus pacificus', *PLOS ONE*,
7(9), p. e44255. doi:10.1371/journal.pone.0044255.

Sowa, J.N. *et al.* (2019) 'The Caenorhabditis elegans RIG-I Homolog DRH-1 Mediates the
Intracellular Pathogen Response upon Viral Infection', *Journal of Virology* [Preprint].
doi:10.1128/JVI.01173-19.

Stein, L.D. *et al.* (2003) 'The genome sequence of Caenorhabditis briggsae: a platform for
comparative genomics.', *PLoS biology*, 1(2), p. E45. doi:10.1371/journal.pbio.0000045.

Stentiford, G.D. *et al.* (2016) 'Microsporidia – Emergent Pathogens in the Global Food
Chain', *Trends in Parasitology*, 32(4), pp. 336–348. doi:10.1016/j.pt.2015.12.004.

Szumowski, S.C. and Troemel, E.R. (2015) 'Microsporidia–host interactions', *Current Opinion*
in Microbiology, 26, pp. 10–16. doi:10.1016/j.mib.2015.03.006.

Tamim El Jarkass, H. and Reinke, A.W. (2020) 'The ins and outs of host-microsporidia
interactions during invasion, proliferation and exit', *Cellular Microbiology*, 22(11).
doi:10.1111/cmi.13247.

Tecle, E. *et al.* (2021) 'The purine nucleoside phosphorylase pnp-1 regulates epithelial cell
resistance to infection in C. elegans', *PLOS Pathogens*, 17(4), p. e1009350.
doi:10.1371/journal.ppat.1009350.

Tecle, E. and Troemel, E.R. (2022) 'Insights from C. elegans into Microsporidia Biology and
Host-Pathogen Relationships.', *Experientia supplementum (2012)*, 114, pp. 115–136.
doi:10.1007/978-3-030-93306-7_5.

Thomas, J.H. (2006) 'Adaptive evolution in two large families of ubiquitin-ligase adapters in
nematodes and plants', *Genome Research*, 16(8), pp. 1017–1030. doi:10.1101/gr.5089806.

Trapnell, C. *et al.* (2010) 'Transcript assembly and quantification by RNA-Seq reveals
unannotated transcripts and isoform switching during cell differentiation', *Nature*
Biotechnology, 28(5), pp. 511–515. doi:10.1038/nbt.1621.

Trapnell, C. *et al.* (2013) 'Differential analysis of gene regulation at transcript resolution with
RNA-seq', *Nature Biotechnology*, 31(1), pp. 46–53. doi:10.1038/nbt.2450.

Trapnell, C., Pachter, L. and Salzberg, S.L. (2009) 'TopHat: discovering splice junctions with
RNA-Seq', *Bioinformatics*, 25(9), pp. 1105–1111. doi:10.1093/bioinformatics/btp120.

Troemel, E.R. *et al.* (2008) 'Microsporidia are natural intracellular parasites of the nematode
Caenorhabditis elegans.', *PLoS biology*, 6(12), pp. 2736–2752.
doi:10.1371/journal.pbio.0060309.

Vávra, J. and Lukeš, J. (2013) 'Chapter Four - Microsporidia and "The Art of Living Together"',
in Rollinson, D. (ed.) *Advances in Parasitology*. Academic Press, pp. 253–319.
doi:10.1016/B978-0-12-407706-5.00004-6.

Wadi, L. and Reinke, A.W. (2020) 'Evolution of microsporidia: An extremely successful group
of eukaryotic intracellular parasites', *PLOS Pathogens*. Edited by L.J. Knoll, 16(2), p.
e1008276. doi:10.1371/journal.ppat.1008276.

Wang, L., Wang, S. and Li, W. (2012) 'RSeQC: quality control of RNA-seq experiments',
Bioinformatics, 28(16), pp. 2184–2185. doi:10.1093/bioinformatics/bts356.

Willis, A.R. *et al.* (2021) 'A parental transcriptional response to microsporidia infection
induces inherited immunity in offspring', *Science Advances*, 7(19), p. eabf3114.
doi:10.1126/sciadv.abf3114.

Willis, A.R. and Reinke, A.W. (2022) 'Factors That Determine Microsporidia Infection and
Host Specificity.', *Experientia supplementum (2012)*, 114, pp. 91–114. doi:10.1007/978-3-
030-93306-7_4.

Wong, D. *et al.* (2007) 'Genome-wide investigation reveals pathogen-specific and shared
signatures in the response of Caenorhabditis elegansto infection', *Genome Biology*, 8(9), p.
R194. doi:10.1186/gb-2007-8-9-r194.

Zárate-Potes, A. *et al.* (2020) 'The C. elegans GATA transcription factor elt-2 mediates
distinct transcriptional responses and opposite infection outcomes towards different
Bacillus thuringiensis strains', *PLOS Pathogens*, 16(9), p. e1008826.
doi:10.1371/journal.ppat.1008826.

Zhang, G. *et al.* (2016) 'A Large Collection of Novel Nematode-Infecting Microsporidia and
Their Diverse Interactions with Caenorhabditis elegans and Other Related Nematodes', *PLOS
Pathogens*. Edited by J.B. Lok, 12(12), p. e1006093. doi:10.1371/journal.ppat.1006093.

820
821