Inferring CTCF binding patterns and anchored loops across human tissues and cell types

Hang Xu[1,2,#], Xianfu Yi[3,#], Wei Wang[1], Xinlei Chu[1], Shijie Zhang[4], Xiaobao Dong[5], Zhao Wang[4,6], Jianhua Wang[4], Yao Zhou[4], Ke Zhao[4], Hongcheng Yao[7], Zheng Nan[8], Junwen Wang[9], Dariusz Plewczynski[10], Pak Chung Sham[7], Kexin Chen[1], Dandan Huang[11,*] and Mulin Jun Li[1,3,*]

[1]Department of Epidemiology and Biostatistics, Tianjin Key Laboratory of Molecular Cancer Epidemiology, The Province and Ministry Co-sponsored Collaborative Innovation Center for Medical Epigenetics, National Clinical Research Center for Cancer, Tianjin Medical University Cancer Institute and Hospital, Tianjin Medical University, Tianjin, China.

[2]Singapore Immunology Network (SIgN), Agency for Science, Technology and Research (A*STAR), Singapore.

[3]Department of Bioinformatics, Laboratory of Immune Microenvironment and Disease (Ministry of Education), School of Basic Medical Sciences, Tianjin Medical University, Tianjin, China.

[4]Department of Pharmacology, Tianjin Key Laboratory of Inflammation Biology, School of Basic Medical Sciences, Tianjin Medical University, Tianjin, China.

[5]Department of Genetics, School of Basic Medical Sciences, Tianjin Medical University, Tianjin, China.

[6]Institute of Aging, Key Laboratory of Alzheimer's Disease of Zhejiang Province, School of Mental Health and The Affiliated Kangning Hospital, Wenzhou Medical University, Wenzhou, China.

[7]Centre for PanorOmic Sciences-Genomics and Bioinformatics Cores, The University of Hong Kong, Hong Kong.

[8]Department of Network Security and Informatization, Tianjin Medical University, Tianjin, China.

[9]Department of Health Sciences Research and Center for Individualized Medicine, Mayo Clinic, Scottsdale, Arizona, USA.

[10]Faculty of Mathematics and Information Science, Warsaw University of Technology, Warsaw, Poland.

[11]Wuxi School of Medicine, Jiangnan University, Wuxi, China.

[#]The authors contributed equally to this work

*Correspondence: mikey.huang2011@gmail.com (D.H.), mulinli@connect.hku.hk (M.J.L.)

Abstract

CCCTC-binding factor (CTCF) is a transcription regulator which is involved in many cellular processes. How CTCF recognizes DNA sequence to exert chromosome barrier or enhancer blocking effects remains to be fully interrogated. Despite many computational tools were developed to predict CTCF-mediated loops qualitatively or quantitatively, few could specially evaluate the regulatory potential of DNA sequence at CTCF binding sites (CBSs) and how it affects chromatin loop formation. Here, we developed a deep learning model, DeepAnchor, to precisely characterize the binding patterns for different types of CBSs. By incorporating base-wise genomic/epigenomic features, we revealed distinct chromatin and sequence features for CTCF-mediated insulation and looping at a high

resolution, such as two sequence motifs flanking the core CTCF motif at loop-associated
CBSs. Besides, we leveraged the predicted anchor score to optimize the loop extrusion
model and achieved the best performance in predicting CTCF-anchored loops. We
established a compendium of context-specific CTCF-anchored loops across 52 human
tissue/cell types and found that genomic disruption of CTCF-anchored loops may
represent a general causal mechanism of disease pathogenesis. These computational
models, together with the established resource, could facilitate the mechanistic research
on how the CTCF-mediated *cis*-regulatory elements (CREs) shapes context-specific gene
regulation in cell development and disease progression.
52

Introduction

Variable interactions between enhancer and target gene have been systematically profiled
in explaining tissue/cell type-specific transcriptional regulation (Consortium, 2012;

Consortium et al., 2020), but the mechanisms underlying the precise wiring between genes

and the enhancers from varied distances remain unclear. As the main insulator-related
transcription factor (TF) discovered in vertebrates, CTCF assists cohesin to form chromatin
loops, which is believed to be the fundamental biophysical basis for distal gene regulation
(Ali et al., 2016; Braccioli and de Wit, 2019; Gabriele et al., 2022; Ghirlando and Felsenfeld,
2016; Nichols and Corces, 2015; Ong and Corces, 2014; Phillips and Corces, 2009).
According to the loop extrusion model (Davidson et al., 2019; Kim et al., 2019), cohesin
dimers load on DNA at NIPBL binding sites and slide along with DNA until reaching the
anchors, which are usually bound by specific cofactors, such as CTCF, YY1, and ERα
(Beagan et al., 2017; He et al., 2014; Sanborn et al., 2015). Chromatin loops mediated by
YY1 and ERα are found to form enhancer-promoter interactions, while CTCF-anchored
loops generally enclose the formers (Beagan et al., 2017), assembling genome units that
are named insulated neighborhoods (INs). Many lines of evidence also suggest that these
structures constitute the mechanistic underpinnings of higher-order chromosome
organizations (Beagan and Phillips-Cremins, 2020; Hnisz et al., 2016), such as
topologically associating domains (TADs) (Dixon et al., 2012) (Dixon et al., 2012; Nora et
al., 2012). Enhancers tend to regulate genes within the same IN/TAD while the regulation
across different INs/TADs is more likely to be prevented (Dowen et al., 2014; Islam et al.,
2023). Therefore, characterizing CTCF binding patterns and their regulatory
consequences will greatly facilitate the analyses of fine-grained 3D genome regulation.
76
CTCF and its associated *cis*-regulatory elements (CTCF-mediated CREs) play a critical
role in controlling tissue/cell type-specific gene expression by maintaining chromatin
domain boundaries or blocking enhancer activities, however, CTCF has been revealed as
a versatile TF that plays several other roles in different scenarios, including gene activation,

transcriptional repression, pausing, and alternative splicing (Filippova et al., 1996; Guo et

al., 2012; Lobanenkov et al., 1990; Oh et al., 2021; Shukla et al., 2011; Vostrov and
Quitschke, 1997). Therefore, it is necessary to distinguish insulator or looping-related

CTCF binding sites (CBSs) from those with other functions. There are a lot of works that
analyze the functional heterogeneity of CTCF in the nucleus, which revealed that the
binding motif of CTCF could be the key to understand the behavior of CTCF at specific site
(Huang et al., 2021b; Nakahashi et al., 2013; Ribeiro-Dos-Santos et al., 2022). Studies
have uncovered that insulation potency heavily depends on the number of CBSs in tandem
and an upstream flanking sequence at the core CTCF motif (Huang et al., 2021b). Besides,
in the formation of CTCF/cohesin chromatin loops, the CBSs at loop anchors tend to be
arranged in convergent orientation (Guo et al., 2015). However, these studies are either
unable to count the feature patterns of individual CBSs or limited to given genomic loci.
Besides, a large number of computational tools were developed to qualitatively or
quantitatively predict CTCF-mediated loops (Cao et al., 2021; Ibn-Salem and
Andrade-Navarro, 2019; Kai et al., 2018; Kuang and Wang, 2021; Lv et al., 2021;
Matthews and Waxman, 2018; Oti et al., 2016; Wang et al., 2021; Xi and Beer, 2021;
Zhang et al., 2018), but few could specially evaluate the regulatory potential of DNA
sequence at CBS and how it affects the loop formation. Moreover, it remains elusive to
see which factors or combinations of features determine the specificity among different
types of functional CTCF binding events, and whether such information could contribute
the prediction of CTCF-anchored loops across the whole genome.
102
Here, we develop an interpretable deep learning model, DeepAnchor, to ask what types of
genomic/epigenomic features determine a CBS to be insulator/cohesin/loop-associated.
By leveraging large-scale base-wise genomic and epigenomic features, we uncovered
several critical patterns for CTCF-mediated insulation and looping at a high resolution. We
embedded the predicted anchor score to the previous loop extrusion model (Xi and Beer,
2021), and found the optimized model LoopAnchor outperforms the existing methods for
predicting CTCF-anchored loops. Furthermore, for the first time, we established a
landscape of tissue/cell type-specific CTCF-anchored loops across 52 human tissue/cell
types and applied them to interpret disease-causal variants. This method together with the
compiled resources will facilitate the mechanistic research on the dynamic regulation of
3D chromatin during cell development and disease progression.
114
115

Results

DeepAnchor enables high-confidence CTCF binding patterns characterization

The key feature of DeepAnchor is the implementation of a deep learning model that
leverages base-wise features for CTCF-medicated CREs detection. Previous studies
usually analysed the sequence and chromatin status of CBS in particular tissue/cell types
with a one-dimensional feature vector, in which regional feature values are averaged for
each CBS regardless of heterogeneous signals across the locus (Kai et al., 2018; Zhang
et al., 2018). However, a true CTCF-medicated CRE differs from other CBSs in
fine-grained architecture (Huang et al., 2021b; Lee et al., 2022; Ong and Corces, 2014),
which raises new requirements for feature characterization before modelling. One

straightforward solution is to profile a CTCF binding event using high-resolution features
surrounding the CBS. In our implementation, for each CBS, 44 quantitative base-wise
genomic/epigenomic features within the ±500bp region were obtained from CADD
annotation (Rentzsch et al., 2019), and DNA sequence within ±500bp of the CBS was
also extracted and represented by one-hot encoding (Table S1). Thus, by concatenating
large-scale annotations with converted DNA features at the base-wise level, a feature
matrix with a size of 1000×48 was generated for each CBS (Figure 1A top left).

We modelled three major forms of CBSs across the whole human genome, including
insulator-associated CBS (insulator CBS), cohesin-associated CBS (cohesin CBS) and
loop-associated CBS (loop CBS). These CTCF-medicated CREs share common CTCF
binding motif but could display varied regulatory functions. To prepare high-quality ground
truth for different types of CBS learning, we required that each selected positive site
should contain 1) putative CBS by motif scanning; 2) observed CBS by CTCF ChIP-seq at
given tissue/cell type. For insulator CBS, we intersected the selected sites with genomic
segments showing ChromHMM insulator state (15 states model). For cohesin CBS, we
overlapped the selected sites with cohesin ChIP-seq in matched cell type. For loop CBS,
we required the selected sites located in cohesin loop anchor by ChIA-PET at same cell
type (Figure 1A top right). We also randomly selected equal number of negative samples
from the whole CBS pool. To avoid ambiguous results, multiple CBSs shared the same
CTCF/cohesin ChIP-seq peak(s) or the same cohesin loop anchor(s) are excluded (see
Methods). Including base-wise features of the region that spans the selected CBSs will
largely expand the feature volume and scale, which will introduce a flood of information
that is usually beyond the ability of classical machine learning algorithms. For example,
regions with 1000bp can incorporate $1000\times48=48,000$ features if a base-wise value is
applied. To address this challenge, we applied a deep convolutional neural network (CNN)
to extract important patterns from the feature matrixes, which greatly reduces the
calculation burden and could increase prediction performance. After training a classifier on
a positive and negative dataset, the DeepAnchor framework returns the probability of a
CBS region belonging to a true CTCF-mediated CRE, which is called the DeepAnchor
score in the downstream analyses (Figure 1A bottom, see Methods).

We trained DeepAnchor model using high-quality CTCF/RAD21 ChIP-seq and RAD21
ChIA-PET data on the GM12878, K562, and H1-hESC cell lines respectively, and tested
the repeatability of DeepAnchor by cross-validation (see Methods). To avoid the potential
influence of chromosome-specific confounding factors, we divided the train-test-valid set
by splitting CBSs into subsets each on different chromosomes. That is, we selected
chr1-16 as the training set, chr17-18 as the validation set, and others as the test set. The
DeepAnchor models trained on different types of CBS samples all received high area
under the ROC curve (AUC, 0.94-0.99) via five-fold cross-validation across the three cell
lines (Figure 1B). To demonstrate that the DeepAnchor scores derived from different
samples are comparable, we performed cross-cell type-validation across the three cell
lines. Notably, among different types of CBSs, the DeepAnchor models exhibited
consistently high cross-sample prediction performance on all testing pairs (all AUCs > 0.9,

Figure 1C) and the DeepAnchor scores are highly correlated among the three cell
type-specific models (Pearson's r = 0.91-0.96) (Figure 1D), suggesting CTCF-mediated
CREs share conservative features in different contexts and the feasibility to predict
genome-wide functional CBSs at the organismal level.
175
We set a cut-off to define positive and negative CTCF-mediated CREs by finding the
points on the ROC curve with the largest Youden's J statistic. The optimal cut-point was
obtained when the DeepAnchor score is close to 0.5 for different models, where 86%-95%
positive and 85%-88% negative samples in training datasets are correctly assigned. With
the GM12878 DeepAnchor models and the optimal cut-off, we generated positive and
negative CTCF-mediated CREs genome-widely. Using the FANTOM5 enhancer dataset,
we found that positive CBSs colocalize with more enhancers within ±100kb regions than
negative CBSs on average (P-value < 2.22e-16, Mann Whitney U Test) (Figure 1E and
Figure S1A). Besides, by counting the number of enhancers in every 10kb bin within
±100kb of CTCF-mediated CREs, we noted that there is a strand-oriented asymmetric
pattern for positive CBSs, but not negative ones, particularly for loop CBSs and cohesin
CBSs (Figure 1F and Figure S1B). Consistent with previous findings (Clarkson et al.,
2019), the density of CTCF-mediated CREs around the TAD boundaries also shows
strand bias, in which positive CBSs at plus-strand are usually enriched at the left TAD
boundary, while those at minus-strand are aggregated at right TAD boundary. However, no
such patterns were observed for negative CBSs (Figure 1G and Figure S1C). We also
calculated the intersection of the predicted CTCF-mediated CREs (at different thresholds)
among three CBS types. We found that all loop CBSs and most of insulator CBSs are
cohesin-related, which indicates that CBS bound by cohesin could has other functions
than forming chromatin loop or insulation. When increasing the cutoff of DeepAnchor
score, the overlap between the three types of CBSs decreased a lot, and only the overlap
between insulator CBSs and cohesin CBSs have slight changes (Figure 1H). Together,
these results demonstrated that our DeepAnchor model can accurately capture
high-confidence CTCF-mediated CREs across the whole genome.
200

DeepAnchor reveals distinct base-wise chromatin and sequence features
for CTCF-mediated CREs
Although DeepAnchor leverages a deep learning structure to extract high-level
configurations from large-scale features, it also raises challenges in analysing feature
importance and its biological insight. Base-wise representation of various
genomic/epigenomic features in the DeepAnchor model makes it possible to depict a
comprehensive picture of the patterns of CTCF-mediated CREs. To visualize underlying
relationships among these features, we first clustered the feature scores upon the training
dataset and found these features could be generally partitioned into three major subsets,
including transcription-associated, conservation-associated, and chromatin
state-associated feature sets (Figure S2). We used SHapley Additive exPlanations (SHAP)
(Lundberg et al., 2020), a game-theoretic approach to explaining the output of
DeepAnchor. SHAP returns the Shapley value matrix with the same alignment as the input
feature matrix, which shows the contribution of each feature at each position. To inspect

the overall contribution of each feature, we calculated the mean Shapley values at all
positions and then summarized the mean values (Figure 2A). Overall, feature of CTCF
binding evidence at open chromatin (EncOCctcfPval) greatly contribute to the model.
Shapley values of G/T/A/C are also top-ranked, which implies the importance of DNA
sequence. As expected, open chromatin features (e.g., EncOCFairePVal, EncOCC)
display higher importance than other features (Figure 2A and Figure S3). Interestingly,
nucleosome positioning evidence (EncNucleo) displayed more contribution in loop CBS
model, suggesting the CTCF footprinting pattern among nucleosomes could determine
the formulation of CTCF-mediated chromatin loop. Besides, a conservation feature, the
neutral evolution score defined by GERP++ (GerpN) also showed weak preference in loop
CBS model (Figure 2A). Taken together, these results indicate that some informative
features may capture the unique and important patterns that can distinguish
CTCF-mediated CREs from other types of CTCF binding events.
228
To explore the base-wise schema of CTCF-mediated CREs, for several important features,
we investigated the distribution of Shapley values derived from the DeepAnchor model at
every position of CBS and its surrounding region (±500bp) upon the testing dataset with
randomly sampled CBSs. For example, CTCF binding intensity and its occupancy
(EncOCctcfPval) display a periodical pattern at cohesin/loop CBSs instead of insulator
CBSs, in which higher binding intensity of central CBS heavily contributes to positive CBS
discrimination but the Shapley values flip at around 150bp far away from central CBS
(Figure 2B top). Interestingly, we observed that, only at loop CBS, the Shapley values of
nucleosome positioning signature (EncNucleo) are reciprocally distributed with those of
CTCF binding intensity (Figure 2B middle), and the nucleosome positioning level is
anti-correlated with its Shapley value (Figure S4A), indicating that CTCF at loop anchor
could enlarge and protect the linked DNA between its two neighboring nucleosomes to
facilitate the loop formation. Such base-wise visualization of feature contribution also
greatly supports the previous finding that the binding of CTCF provides an anchor point for
positioning nucleosomes thus resulting in an array of well-positioned nucleosomes
flanking sites at particular CBSs (Cuddapah et al., 2009; Fu et al., 2008). We also found
that the Shapley values of the conservation feature (GerpN) showed similar distribution as
those of CTCF binding features, especially in loop CBSs (Figure 2B bottom). In addition,
consistent with previous results (Luan et al., 2021), an investigation of base-wise Shapley
value distribution upon positive or negative training datasets demonstrated that
CTCF-mediated CREs usually demonstrated stronger binding intensity than other types of
CBSs (Figure S4B).
251
Previous CTCF multivalency studies revealed that CTCF reads sequence diversity
through combinatorial clustering of its 11 zinc fingers (ZFs) (Nakahashi et al., 2013; Rhee
and Pugh, 2011), yet the sequence determinants of CTCF-mediated insulation remain
elusive. By applying MEME motif discovery to all predicted positive and negative CBSs,
we revealed two additional weak motifs flanking the core CTCF motif only at positive loop
and insulator CBSs (Figure 2C). In agreement with a sensitive insulator reporter assay
(Huang et al., 2021b), the upstream motif separated from the core sequence by 5 or 6 bp

is recognized by ZFs 9-11 and may contribute to the directionality of CTCF binding to
CBSs (Yin et al., 2017). Interestingly, the downstream motif next to the core sequence is
different from previous findings in terms of distance and sequence context (Nakahashi et
al., 2013). Nonetheless, this downstream sequence associated with CTCF N terminus and
ZFs 1-2 has been documented to stabilize cohesin engagement (Li et al., 2020). Thus, the
distinct sequence features together with multiple lines of existing evidence for CTCF
binding patterns further demonstrate the fidelity of the DeepAnchor model.

Incorporating DeepAnchor score into loop extrusion model improves the performance of CTCF-anchored loop prediction

Although DeepAnchor models the insulative and looping potential of genome-wide CBSs
at the organismal level, accurate prediction of context-specific CTCF-anchored loop will
be essential in understanding 3D chromatin interactions and fine-grained gene regulation.
By assuming that looping interactions found in the CTCF ChIA-PET experiments are due
to the blocking and localization of cohesin at CBSs, a simple but effective loop extrusion
model (LEM) was previously developed (Xi and Beer, 2021). This quantitative model
evaluates CTCF-anchored loop formation with only four features, including CTCF-binding
intensity, CTCF motif orientation, the distance between CTCF-binding events, and loop
competition. Albeit the acquired specificity, we reasoned that incorporating the
loop-associated DeepAnchor score into LEM could further improve the performance of
CTCF-anchored loop prediction. To this end, by considering the looping potential of CTCF
binding (anchor score predicted by loop-associated DeepAnchor model), we weighed the
probability of CTCF binding in the original LEM model and reached a new implementation,
called LoopAnchor (see Methods).

We compared the performance of LoopAnchor with five prevalent algorithms for
CTCF-mediated interaction prediction, including Naïve and Oti methods (Oti et al., 2016),
CTCF-MP (Zhang et al., 2018), Lollipop (Kai et al., 2018), and LEM (Xi and Beer, 2021).
To ensure a fair benchmark, all supervised models were trained on GM12878 RAD21
ChIA-PET data and tested on K562 RAD21 ChIA-PET data. We found that LoopAnchor
outperforms other state-of-art methods in terms of both ROC-AUC (0.936) and PR-AUC
(0.921) (Figure 3A), indicating the effectiveness of our model optimization. Besides, the
correlation analysis between the predicted loop intensities and the observed ChIA-PET
signals on K562 showed that LoopAnchor receives the highest correlation (Pearson's r =
0.653) among the six tools (Figure 3B), which also demonstrates the superiority of
LoopAnchor for quantitative measurement of CTCF-anchored loop.

To investigate whether our LoopAnchor mode could capture the dynamic loop changes
during cell development, we applied our LoopAnchor model to human monocytes
activation data with paired *in situ* Hi-C and CTCF ChIP-seq before and after exposure to
PMA (Phanstiel et al., 2017). We found our LoopAnchor model can accurately predict
25/34 gained loops and all six lost CTCF-anchored loops (Figure 3C). For example, on the
JAG1 locus at chromosome 20:9547732-11575097, there are eight loops detected by
Hi-C, among which one gained loop is identified only at differentiated macrophages

(Figure 3D). Consistently, LoopAnchor not only detected all unchanged loops but also
accurately predicted the gained loop (Figure 3D, see Methods). Notably, in this genomic
region, the difference of loop intensity between monocyte and macrophage cells predicted
by LoopAnchor is highly correlated with the loop score changes measured from Hi-C data
(Pearson's r = 0.796), indicating that LoopAnchor can quantitatively capture the local loop
structure changes at the dynamic cellular process.
309
A landscape of CTCF-anchored loops across 52 human tissue/cell types
Given the simplicity and gained performance of the LoopAnchor model, we sought to
leverage existing large-scale CTCF ChIP-seq data to learn a global picture of
CTCF-mediated chromatin interactions across different human tissue/cell types. We first
uniformly processed 740 CTCF ChIP-seq datasets collected from ENCODE (Consortium
et al., 2020), CistromDB (Zheng et al., 2019), and ChIP-Atlas (Oki et al., 2018) based on
the ENCODE TF ChIP-seq processing pipeline (Table S2, see Methods). According to
data quality control, peak calling filtering, and redundancy removing, we totally retrieved
168 high-quality CTCT ChIP-seq datasets, relating to 113 normal and 55 cancer samples
across 32 tissues and 20 anatomical cell types (Figure 4A and Table S3, see Methods).
Each biosample contains around 100,000 peaks (mean = 104,691, median = 96,296,
Figure S5A). We applied LoopAnchor and LEM to detect genome-wide CTCF-mediated
loops for each biosample respectively, and we found that nearly 20,000 loops were
detected per biosample (mean = 16,255 and median = 16,637 for LoopAnchor, mean =
17,826 and median = 18,252 for LEM, Figure S5B and S5C).
325
Based on the shared pattern of detected loops across 168 biosamples, we first classified
all LoopAnchor loops into four types: (1) 8.2% (n = 6,179) tissue/cell type-shared loops in
more than 75% biosamples; (2) 9.0% (n = 6,802) tissue/cell type-relatively shared loops in
more than 50% but less than 75% of biosamples; (3) 13.3% (n = 10,016) tissue/cell
type-relatively specific loops in more than 25% but less than 50% of biosamples; (4) 69.4%
(n = 52,216) tissue/cell type-specific loops in less than 25% of biosamples (Figure 4B, see
Methods). Based on the same strategy, the LEM loops were also classified into these four
types (Figure S6A). Compared to LEM, LoopAnchor can detect more shared loops but
less specific loops globally (8.2% vs. 3.9% for tissue/cell type-shared loops, and 69.4% vs.
80.3% for tissue/cell type-specific loops, Figure 4B and Figure S6A), suggesting
LoopAnchor could capture more conservative CTCF-anchored loops across different
tissue/cell types. By investigating the distribution of LoopAnchor-predicted loop intensity
scores, we found a significant difference among the four loop categories (P-value <
2.22e-16, two-tailed Mann-Whitney U test, Figure 4C), in which the tissue/cell type-shared
loops received the highest scores and the tissue/cell type-specific loops got the lowest
scores. A similar pattern was also observed for the loop intensity score derived from the
LEM method (Figure S6B). To evaluate the tissue/cell type specificity of classified
CTCF-mediated loops, we leveraged the predicted chromatin loops by Peakachu for 42
human tissue/cell types (Salameh et al., 2020) and compared the number of associated
tissue/cell types among different categories. We found that the tissue/cell type distribution
of the LoopAnchor-predicted loops also showed significant differences among the four

loop categories (P-value < 2.22e-16, two-tailed Mann-Whitney U test, Figure 4D). As
expected, the number of associated tissue/cell types gradually decreased from the
tissue/cell type-shared group (mean = 4, median = 4.743) to the tissue/cell type-specific
group (mean = 2, median = 2.314), indicating the effectiveness of our loop classification. A
similar trend was also identified for the detected loops from the LEM method (Figure S6C).
Together, we established a compendium of tissue/cell type commonality and specificity for
CTCF-anchored loops across large-scale human tissue/cell types. All predicted loops for
168 selected biosamples and all 764 biosamples can be visualized and compared as
separated   tracks   at   UCSC   Track   Data   Hubs
(https://genome.ucsc.edu/cgi-bin/hgHubConnect) by entering customized hub URLs
https://raw.githubusercontent.com/mulinlab/LoopAnchor/master/hubs_landscape.txt   or
https://raw.githubusercontent.com/mulinlab/LoopAnchor/master/hubs_all.txt, respectively.
359

Tissue/cell type-specific loop anchors are highly enriched at
disease-causal loci

Multiple lines of evidence have supported that CTCF/cohesin binding sites are highly
mutated in cancers (Fang et al., 2020; Guo et al., 2018; Katainen et al., 2015) or
constantly shaped via evolutionary selection (Kentepozidou et al., 2020; Sadowski et al.,
2019). However, whether context-specific CTCF-mediated looping and associated loop
anchors are more likely linked to disease-causal genomic loci has not been systematically
tested. By collecting 12,738 causal variants for 54 blood-related autoimmune diseases
from genome-wide association studies (GWASs) (Wang et al., 2020) (Table S4) and
sampling matched control variants, we evaluated the genome-wide enrichment of
autoimmune disease-causal variants on the CTCF-anchored loops predicted by
LoopAnchor in normal tissues with more than three biosamples (see Methods). Notably,
the autoimmune disease-causal variants were significantly enriched in the loop anchors in
blood tissue (mean P-value = 0.00357) compared to other tissues (mean P-value ranges
from 0.14465 (colon) to 0.0055 (prostate), Figure 5A). As the majority of disease-causal
regulatory variants show tissue/cell type specificity in phenotypically relevant contexts,
these results further support that context-dependent loop information could better interpret
GWAS disease-causal variants for complex diseases.
378

To evaluate the relevance of CTCF-mediated loop anchors in the cancer mutation process,
we collected somatic mutations from ICGC whole genome mutation aggregation
(Consortium, 2020) and extracted recurrent mutations in the non-coding regions (see
Methods). By comparing with permuted non-recurrent mutations, we found that mutations
with higher recurrence obtain greater enrichment in CTCF-mediated loop anchors for
cancer biosamples (two-tailed Mann-Whitney U test, Figure 5B). Besides, based on the
whole-genome cancer mutation hotspots collected from three pan-cancer whole-genome
analyses, including ATACseq-AWG (Corces et al., 2018), PCAWG (Rheinbay et al., 2020),
and CNCDriver (Liu et al., 2019), we tested whether the mutation hotspots are more
frequently occurred in CTCF-mediated loop anchors than other CTCF binding loci or
shuffled non-coding genomic regions (see Methods). We uncovered consistent results
among different datasets, in which the loop anchors were highly enriched with cancer

mutation hotspots (two-tailed Mann-Whitney U test, Figure 5C-F). These results suggest
genomic disruption of CTCF-anchored loops may thus represent a general causal
mechanism of disease pathogenesis.
394
395

Discussion

The multiple critical functions of CTCF in the nucleus stimulate constant phenotypic and
mechanistic research in the field, yet how CTCF recognizes insulators to exert
chromosome barrier or enhancer blocking effects remains to be interrogated. In this study,
we developed a novel computational model to accurately predict CTCF-mediated CREs
and CTCF-anchored loops genome-widely. By incorporating large-scale base-wise
genomic and epigenomic features within a deep learning model, we revealed several
distinct chromatin and sequence features for CTCF-mediated insulation at a high
resolution. Importantly, we found two additional sequence motifs flanking the core CTCF
motif at positive loop and insulator-associated CBSs. Besides, we leveraged the predicted
insulator score to optimize the previous loop extrusion model and achieved better
performance in predicting CTCF-anchored loops. Based on the model, for the first time,
we established a compendium of tissue/cell type-specific and -shared CTCF-anchored
loops across 52 human tissue/cell types. Finally, we showed that tissue/cell type-specific
loop anchors are highly enriched at disease-causal loci. Therefore, our results deepen the
understanding of CTCF-mediated insulation and loop formation. The new method together
with the compiled resource also provides useful approaches for studying the dynamic
regulation of 3D chromatin during cell differentiation and disease progression.
414
Although many computational models that can predict CTCF-mediated loops with varied
sensitivity and specificity (Cao et al., 2021; Deng et al., 2022; Ibn-Salem and
Andrade-Navarro, 2019; Kai et al., 2018; Kuang and Wang, 2021; Lv et al., 2021;
Matthews and Waxman, 2018; Oti et al., 2016; Wang et al., 2021; Xi and Beer, 2021;
Zhang et al., 2018), they usually learn from CTCF ChIA-PET data and barely evaluate the
regulatory potential and loop attribute among different types of CBS. Here, we applied
cohesin ChIA-PET/ChIP-seq data, ChromHMM-predicted insulator, and strict CTCF
binding evidence to specifically analyze CTCF binding patterns at three types of
CTCF-mediated CREs. This provides a unique tool for characterizing CTCF binding
consequence and loop formation through the loop extrusion mechanism. Notably, the
limited availability and varied library construction quality of CTCF/cohesin ChIA-PET
restrict the training of context-specific models broadly, however, our cross-cell type
comparisons showed a high agreement among DeepAnchor models trained from different
ChIA-PET data, indicating the discrimination of true CTCF-mediated CREs may rely on
several conservative features revealed by our interpretable deep learning model, such as
higher binding intensity, well-positioned nucleosomes as well as two unique motifs
flanking the central CBS.
432
By introducing the DeepAnchor score into a loop competition and extrusion model (Xi and

Beer, 2021), we observed improved performance in predicting CTCF-anchored loops and
demonstrated that the new LoopAnchor method can achieve better quantitative estimation.
However, in both LEM and LoopAnchor, the quantitative predictions only measure the
contribution of each component independently. For example, the insulation potential of
CBS and loop competition are dependent processes based on dynamic chromatin context
(Davidson and Peters, 2021), which motivates a future optimization direction. Besides,
given existing methods without providing CTCF-associated anchor prediction results, we
cannot evaluate whether incorporating other anchor score would achieve better
performance. Instead, we performed systematic comparisons with five prevalent
algorithms for CTCF-mediated loop prediction. Clearly, our method is limited to inferring
specific CREs and loops bound by CTCF, but recent studies indicated that some TFs
could function as novel architectural proteins in regulating genome organization (Yi et al.,
2021), such as YY1 (Weintraub et al., 2017), ZNF143 (Bailey et al., 2015), MAZ
(Ortabozkoyun et al., 2022), BHLHE40 (Hu et al., 2020), CTCFL (Debruyne et al., 2019),
etc., and some of which are independent of CTCF binding. Our model can be easily
extended to train classifiers to extract base-wise features for binding patterns of these
architectural proteins, given the targeted regions and negative controls. Thus, predicting
full spectrum anchor sites and their associated looping events across the genome deserve
further in-depth investigation.
453

The landscape of CTCF-anchored loops established by applying LoopAnchor to 168
uniformly processed human CTCF ChIP-seq biosamples could be a useful resource in 3D
genome and regulatory genomics studies. Since anchor scores were estimated at the
organism level, LoopAnchor still only requires CTCF binding profiles to accurately predict
CTCF-anchored loops. This will greatly simplifies the application, particularly for studying
dynamic 3D CTCF code during cell development and disease progression. In addition, the
compiled tissue/cell type-shared and -specific loops could facilitate the interpretation of
disease-causal variants identified by GWAS and somatic non-coding driver mutations in
cancers.
463

Methods

Training dataset preparation

We prepared three types of target regions from high quality functional genomics data of three Tier 1 ENCODE cell lines (GM12878, K562, and H1-hESC), including insulators estimated by ChromHMM (Ernst et al., 2011) (https://genome.ucsc.edu/cgi-bin/hgTrackUi?db=hg19&g=wgEncodeBroadHmm), cohesin (RAD21) ChIP-seq narrow peaks (ENCODE: ENCSR000AKB, ENCSR000BPJ, ENCSR000BNH), and cohesin (RAD21) ChIA-PET loops (ENCODE: ENCSR752QCX, ENCSR000CAC, ENCSR543YTV) (Grubert et al., 2020). For ChromHMM insulator segments and RAD21 ChIP-seq narrow peaks, we merged the intervals to avoid overlapping. For RAD21 ChIA-PET data, we preprocessed them with ChIA-PET2 using default parameters (Li et al., 2017). Inter-chromosomal interactions and interactions with less than 2 PETs were filtered out, and the anchor of loops were extracted and merged based on position to get anchor list without overlapped regions. Then, three types of CBSs were prepared by intersecting target intervals with candidate CBSs by motif scanning as well as CTCF ChIP-seq narrow peaks (ENCODE: ENCSR000AKB, ENCSR000BPJ, ENCSR000BNH), including insulator CBS, cohesin CBS, and loop CBS. Thus, for each cell type, the candidate CBSs colocalized with both CTCF ChIP-seq peaks and targeted regions were selected as positive set, while others constituted the pool of negative set. To avoid biases, we removed the CBSs shared the same target region from positive regions, because we cannot distinguish which one is the true positive one.

Feature preparation

DNA binding motif of CTCF (MA0139.1) was downloaded from JASPAR2020 (Fornes et al., 2019). We used FIMO of MEME Suite (v5.1.1) (Grant et al., 2011) to scan the CTCF binding motif across the whole human genome (GRCh37/hg19), resulting in 110,059 CTCF binding sites with P-value < 1e-5. According to the topic relevance, we selected 44 base-wise genomic/epigenomic features from the CADD (v1.4) annotation database (https://cadd.gs.washington.edu/static/ReleaseNotes_CADD_v1.4.pdf) (Kircher et al., 2014; Rentzsch et al., 2019). For each CBS, feature values of ±500kb surrounding region were extracted and stored together to form a feature matrix. All features were scaled by a min-max scaling algorithm. To reduce the influence of extreme high/low values, we extracted 0.1 and 99.9 percentile as the min and max values. Besides, we got the DNA sequence of the ±500kb region of the CBS center using BEDTools (Quinlan and Hall, 2010). A, T, C, G were then transferred into numerical value with one-hot encoding: A (1,0,0,0), T (0,1,0,0), G (0,0,1,0), C (0,0,0,1). Finally, by concatenating large-scale annotations with converted DNA features at the base-wise level, a feature matrix with a size of 1000×48 was generated for each CBS.

DeepAnchor model and fitting

Here we described the statistical model of DeepAnchor to identify insulator/cohesin/loop-associated CBSs. For each CBS $i = 1, \ldots, n$, we extracted

genomic/epigenomic features and DNA sequence features at a base-wise level for the ±500bp region of the CBS center. By concatenating features and one-hot encoding sequence, a two-dimensional 1000×48 signal matrix X_{ij} is generated for each CBS

ij. To extract feature patterns from the signal matrix, DeepAnchor uses 1D convolutional kernels to process the signal matrix. For each layer l, the forward propagation from layer $l-1$ to l is expressed as,

$$a_i^l = b_i^l + \sum_{j=1}^{n_{l-1}} \text{conv1d}(k_{ij}^l, a_j^{l-1}) \qquad (1)$$

Where n_l is the number of neurons at layer l, a_j^{l-1} and a_i^l are the j^{th}

neuron at layer $l-1$ and i^{th} neuron at layer l, k_{ij}^l and the scalar b_i^l are the kernel and the scalar bias from layer $l-1$ to i^{th} neuron at layer l. DeepAnchor implements two 1D-CNN layers, each followed by a max-pooling layer and a drop-out layer. After 1D-CNN layers, all the outputs are fully connected. Three fully connected layers convert the outputs of 1D-CNN layers to lower dimensions and finally train a classifier with sigmoid activation,

$$p_1 = \frac{e^{z_1}}{e^{z_1} + e^{z_0}}$$

$$p_0 = 1 - p_1$$

where z_i is the result of fully connected layers which contain only two elements for positive training and negative target respectively. p_1 is the probability that CBS ij belongs to a positive CBS. The DeepAnchor model was implemented with TensorFlow framework. To avoid overfitting, we extracted a balanced negative set by randomly selecting samples from the pool of negative set. Finally, positive and negative datasets were divided into train, test, and valid set according to chromosomes. Samples on chr1-16 were selected as the training set, chr17-18 as validation set and the others as test set. We considered p_1 and p_2 as the possibility the CBS to be a true target-associated CTCF or not, and we calculated binary cross entropy as the loss of model. Input data were divided into small batches with batch size of 50. The model was trained for 20 epochs, but early stopping was enabled when the validation loss does not decrease any more.

Evaluation of DeepAnchor model
We performed five-fold cross-validation to evaluate the DeepAnchor model and plotted ROC curves and corresponding AUC. We set a cut-off to define positive and negative CBSs by finding the points on the ROC curve with the largest Youden's J statistic. Same

with the training procedure on GM12878, DeepAnchor models were trained for each cell
type and the models were also validated by all test sets belonging to three cell types. After
training the DeepAnchor model, we applied it to all potential CBSs across the whole
human genome. DeepAnchor will calculate a score ranging from [0, 1], with a larger value
meaning more likely to be a positive CTCF-mediated CREs. Human enhancer annotation
was downloaded from the FANTOM5 database (Lizio et al., 2015)
(https://fantom.gsc.riken.jp/5/datafiles/latest/extra/Enhancers/). According to Youden's J
statistic, CBSs with anchor scores > 0.5 are selected as positive CTCF-mediated CREs,
while negative CTCF-mediated CREs are those with anchor scores < 0.5. We also
investigated the performance of DeepAnchor models with different cutoffs. We counted
the number of enhancers for positive/negative insulators if the distance between them is
smaller than 100kb. We tested the differences of involved enhancers between positive
and negative CBSs using the Mann Whitney U Test. The estimated TAD data was derived
from the GM12878 Hi-C assay (Dixon et al., 2012), in which there are 4,386 TADs and the
median length of TADs is 520kb. To test whether CBSs have preference for a position or
not, we counted the distance between each CBS and the 5' end of TAD it belongs. Since
TADs have variable lengths, we normalized the distance by counting the ratio concerning
the total length of the TAD. Therefore, the normalized distance of CBSs will be the
percentage of relative position within TAD.
555

Base-wise feature contribution analyses
We used SHapley Additive exPlanations (SHAP) (Lundberg et al., 2020) to interpret the
output of DeepAnchor. Since the input of the DeepAnchor model is a 48x1000 feature
matrix, SHAP returns the Shapley value matrix with the same alignment as the input
feature matrix. We leveraged the shap package (Lundberg et al., 2018) to load the
well-trained DeepAnchor model and used the training dataset as background examples to
receive an expectation over. Then an explainer was generated based on the model and
background dataset. The explainer was then applied to predict the Shapley values for
every position via testing datasets. Note that the training dataset and testing dataset were
the same as the ones in the training DeepAnchor model. We also interpreted the feature
contribution based on the mean Shapley values for all positions.
567

568 **Optimization of loop extrusion model in LoopAnchor**
569 Enlightened by a recent loop competition and extrusion model (Xi and Beer, 2021), we
570 integrated the loop-associated DeepAnchor score (estimated from loop CBS DeepAnchor
571 model) into the model to predict CTCF-anchored loops. We estimated the probability of
572 insulator-associated CTCF binding at CBS i:

$$p_i = \frac{S_i \cdot t_i}{S_i \cdot t_i + a \cdot \overline{F_{re}}}$$

573 where S_i is the CTCF ChIP-seq signal, and a is a constant which have been estimated
574 in the original paper. In this new implementation, t_i is the loop-associated DeepAnchor
575 score at CBS l, thus, revised mean CTCF signal $\overline{F_{re}}$ is given by:
576

$$\overline{F_{re}} = \frac{\sum_i^n S_i \cdot t_i}{n}$$

Comparison with state-of-art methods for CTCF-mediated loop prediction

We acquired Oti code from Oti et.al (Oti et al., 2016), and the naïve method was derived
from Oti code by setting the recursive round of the Oti method to 1. We used the default
parameters to run Oti and Naïve methods, and loops with a score > 0 were considered
positive loops while others were treated as negative ones. CTCF-MP (Zhang et al., 2018),
Lollipop (Kai et al., 2018), and LEM (Xi and Beer, 2021) were installed and used on
relevant cell types according to their GitHub repositories. To facilitate comparison among
different methods, all methods were trained on the GM12878 cell line and tested on the
K562 cell line. Balanced (Pos/Neg = 1) and unbalanced (Pos/Neg = 1/5) gold datasets
were prepared respectively. ROC curve, PR curve, and corresponding AUC were then
plotted and calculated. We also calculated the Pearson correlation coefficient between
predicted loop intensity with real data to show the performance of all methods.

Application of LoopAnchor for dynamic loop detection during macrophage development

We downloaded CTCF ChIP-seq data for both monocyte and macrophage cells (Phanstiel
et al., 2017) (GEO: GSE96800). CTCF ChIP-seq data with replication were processed
with ENCODE standard pipeline to retrieve bigWig and peak files. LoopAnchor was used
to predict CTCF-anchored loops with peak files for both cell states. We also obtained cell
state-specific based on Hi-C loops detected by HiCCUPS (Rao et al., 2014) (GEO:
GSE63525). Because Hi-C loops are not necessary to be mediated by CTCF and cohesin,
we filtered out loops without CTCF peaks in anchor regions on both sides. We mainly
focused on the 'gained' and 'lost' loops from monocyte to macrophage cells by comparing
the predicted loop intensity L of the same loop between the two cell states. The fold
change of loop intensity for a specific loop between macrophage and monocyte is defined
as L_{macro}/L_{mono} and thresholding at $|L_{macro}/L_{mono}| > 3$ for 'gained' and 'lost' loops.

Landscape construction of CTCF-anchored loops across human tissue/cell types

We systematically collected CTCF ChIP-seq datasets from ENCODE (Consortium et al.,
2020), CistromDB (Zheng et al., 2019), and ChIP-Atlas (Oki et al., 2018), and uniformly
processed them using the ENCODE TF ChIP-seq processing pipeline
(https://github.com/ENCODE-DCC/chip-seq-pipeline2). Briefly, the clean reads were
mapped to the human genome (GRCh37/hg19) with BWA (v0.7.17) (Li and Durbin, 2009)
followed by the post-alignment filtering, and finally, the peaks were called using SPP
(v1.15) (Kharchenko et al., 2008). To select the high-quality ChIP-seq biosamples, we
firstly filtered out the outliers with excessively low or high peaks. Then, all B lymphocyte
biosamples prefixed with "GM" were removed except GM12878. For each mapped tissue
or anatomical cell type, we dropped redundant biosamples according to Jaccard similarity

clustering of called peaks and selected one biosample in each cluster with a maximal
peak number. Finally, we applied LoopAnchor and LEM to detect genome-wide
CTCF-mediated loops for each biosample respectively.
621

Definition of tissue/cell type-sharded and -specific loops

All predicted CTCF-anchored loops were generally classified into four categories
according to their shared patterns across different tissue/cell types. Specifically, loops that
were shared in no less than 75% tissue/cell types were classified as tissue/cell
type-shared loops. Loops that were shared by no less than 50% but less than 75%
tissue/cell types were classified as tissue/cell type-relatively shared loops. Loops that
were shared by no less than 25% but less than 50% tissue/cell types were classified as
tissue/cell type-relatively specific loops. The remaining loops which were shared by less
than 25% tissue/cell types were classified as tissue/cell type-specific loops. The public
Hi-C loops detected by Peakachu (Salameh et al., 2020) for 42 tissue/cell types were
downloaded from the 3D genome browser (Wang et al., 2018)
(http://3dgenome.fsm.northwestern.edu/publications.html). Then, the "pairtopair"
command from BEDTools (Quinlan and Hall, 2010) with parameter "-type both" was used
to compare them with loops predicted by LoopAnchor.
636

GWAS disease-causal variant enrichment

We collected 12,738 causal variants for 54 blood-related autoimmune diseases from
CAUSALdb (Wang et al., 2020). To test the genome-wide enrichment of autoimmune
disease-causal variants on the CTCF-anchored loops predicted by LoopAnchor, we first
sampled the same amount of control variants 10,000 times with matched allele frequency
using vSampler (Huang et al., 2021a). For each normal biosample, the predicted
CTCF-anchored loops were flattened into unique anchors, and the "intersect" command
from BEDTools (Quinlan and Hall, 2010) was used to perform the colocalization between
variants and anchors. The enrichment P-value was calculated by counting the number of
permutations that receive a higher percentage of overlapping variants than the real
dataset in each biosample. This analysis was only performed on normal tissues with more
than three biosamples.
649

Non-coding somatic mutation and hotspot enrichment

The genome-wide somatic mutations were downloaded from ICGC Data Portal (Release
28) (Consortium, 2020). Candidate non-coding somatic mutations were retained by
removing these overlapped with exons and splicing sites and were classified into three
categories according to their mutation recurrence ($\in[2,5)$, $\in[5,10)$, ≥ 10). The somatic
mutations with only one recurrence were used as the background for drawing control
datasets. We generated 10,000 control datasets by randomly sampling the equivalent
mutations for each category. For each cancer biosample, the predicted CTCF-anchored
loops were flattened into unique anchors, and the count of intersected hits between
somatic mutations in each category and anchors in a particular biosample was calculated.
By comparing to the average count value from the control datasets, the fold change of
overlapping mutation percentage was derived. The Mann-Whitney U test was used to test

the difference in fold changes among the three categories. In addition, we also curated
candidate somatic mutation hotspots or driver regions from three previous catalogs,
including ATACseq-AWG (Corces et al., 2018), PCAWG (Rheinbay et al., 2020), and
CNCDriver (Liu et al., 2019). For each cancer biosample, the loops predicted by
LoopAnchor with a score $\geq$ 0.01 were kept and flattened to unique anchors as the
"Anchor" datasets, while the same number of non-anchor regions were sampled from
remaining loops with the smaller predicted scores as "Control" datasets. By excluding real
hotspots and restricting them to non-coding genomic regions, the "shuffle" command from
BEDTools (Quinlan and Hall, 2010) was also used to retrieve shuffled hotspots with 100
permutations as "Shuffled" datasets. The Mann-Whitney U test was used to test the
difference in the overlapped percentage of somatic hotspots among these generated
datasets.
674

Data availability

All predicted loops for 168 selected biosamples and all 764 biosamples can be visualized
and compared as separated tracks at UCSC Track Data Hubs
(https://genome.ucsc.edu/cgi-bin/hgHubConnect) by entering customized hub URLs
https://raw.githubusercontent.com/mulinlab/LoopAnchor/master/hubs_landscape.txt or
https://raw.githubusercontent.com/mulinlab/LoopAnchor/master/hubs_all.txt, respectively.
681

682 **Code availability**

683 The source codes of LoopAnchor are freely available under MIT License at
684 https://github.com/mulinlab/LoopAnchor or https://bitbucket.org/xuhang01/loopanchor.
685

Acknowledgements

Funding

Chinese National Key Research and Development Project [2021YFC2500403]; National
Natural Science Foundation of China [32070675 to M.J.L. and 32270717 to M.J.L.];
Natural Science Foundation of Tianjin [19JCJQJC63600 to M.J.L. and 19JCQNJC09000
to X.Y.].

Competing interests

The authors declare no conflict of interest.
695

Contributions

M.J.L. and H.X. conceived of the project. H.X., X.Y. and D.H. performed the analyses and
implemented the software. W.W., X.C., D.H., S.Z., X.D., Z.W., J.W., Y.Z., K.Z., H.Y. and
N.Z. contributed the data collection and processing. J.W., K.C., D.P. and P.C.S.
contributed to manuscript polishing and provided analysis suggestions. H.X., X.Y. and
M.J.L. wrote the manuscript. The authors read and approved the final submission.
702

References

Ali, T., Renkawitz, R., and Bartkuhn, M. (2016). Insulators and domains of gene expression. Curr Opin Genet Dev *37*, 17-26.

Bailey, S.D., Zhang, X., Desai, K., Aid, M., Corradin, O., Cowper-Sal Lari, R., Akhtar-Zaidi, B., Scacheri, P.C., Haibe-Kains, B., and Lupien, M. (2015). ZNF143 provides sequence specificity to secure chromatin interactions at gene promoters. Nat Commun *2*, 6186.

Beagan, J.A., Duong, M.T., Titus, K.R., Zhou, L., Cao, Z., Ma, J., Lachanski, C.V., Gillis, D.R., and Phillips-Cremins, J.E. (2017). YY1 and CTCF orchestrate a 3D chromatin looping switch during early neural lineage commitment. Genome Res *27*, 1139-1152.

Beagan, J.A., and Phillips-Cremins, J.E. (2020). On the existence and functionality of topologically associating domains. Nat Genet *52*, 8-16.

Braccioli, L., and de Wit, E. (2019). CTCF: a Swiss-army knife for genome organization and transcription regulation. Essays Biochem *63*, 157-165.

Cao, F., Zhang, Y., Cai, Y., Animesh, S., Zhang, Y., Akincilar, S.C., Loh, Y.P., Li, X., Chng, W.J., Tergaonkar, V., *et al.* (2021). Chromatin interaction neural network (ChINN): a machine learning-based method for predicting chromatin interactions from DNA sequences. Genome Biol *22*, 226.

Clarkson, C.T., Deeks, E.A., Samarista, R., Mamayusupova, H., Zhurkin, V.B., and Teif, V.B. (2019). CTCF-dependent chromatin boundaries formed by asymmetric nucleosome arrays with decreased linker length. Nucleic Acids Res *47*, 11181-11196.

Consortium, E.P. (2012). An integrated encyclopedia of DNA elements in the human genome. Nature *489*, 57-74.

Consortium, E.P., Moore, J.E., Purcaro, M.J., Pratt, H.E., Epstein, C.B., Shoresh, N., Adrian, J., Kawli, T., Davis, C.A., Dobin, A., *et al.* (2020). Expanded encyclopaedias of DNA elements in the human and mouse genomes. Nature *583*, 699-710.

Consortium, I.T.P.-C.A.o.W.G. (2020). Pan-cancer analysis of whole genomes. Nature *578*, 82-93.

Corces, M.R., Granja, J.M., Shams, S., Louie, B.H., Seoane, J.A., Zhou, W., Silva, T.C., Groeneveld, C., Wong, C.K., Cho, S.W., *et al.* (2018). The chromatin accessibility landscape of primary human cancers. Science *362*.

Cuddapah, S., Jothi, R., Schones, D.E., Roh, T.Y., Cui, K., and Zhao, K. (2009). Global analysis of the insulator binding protein CTCF in chromatin barrier regions reveals demarcation of active and repressive domains. Genome Res *19*, 24-32.

Davidson, I.F., Bauer, B., Goetz, D., Tang, W., Wutz, G., and Peters, J.M. (2019). DNA loop extrusion by human cohesin. Science *366*, 1338-1345.

Davidson, I.F., and Peters, J.M. (2021). Genome folding through loop extrusion by SMC complexes. Nat Rev Mol Cell Biol *22*, 445-464.

Debruyne, D.N., Dries, R., Sengupta, S., Seruggia, D., Gao, Y., Sharma, B., Huang, H., Moreau, L., McLane, M., Day, D.S., *et al.* (2019). BORIS promotes chromatin regulatory interactions in treatment-resistant cancer cells. Nature *572*, 676-680.

Deng, Y., Tang, L., Zhou, X., Wang, W., and Li, M. (2022). Integrating extrusion complex-associated pattern to predict cell type-specific long-range chromatin loops. iScience *25*, 105687.

Dixon, J.R., Selvaraj, S., Yue, F., Kim, A., Li, Y., Shen, Y., Hu, M., Liu, J.S., and Ren, B. (2012). Topological domains in mammalian genomes identified by analysis of chromatin interactions. Nature *485*, 376-380.

Dowen, J.M., Fan, Z.P., Hnisz, D., Ren, G., Abraham, B.J., Zhang, L.N., Weintraub, A.S., Schujiers, J., Lee,
T.I., Zhao, K., *et al.* (2014). Control of cell identity genes occurs in insulated neighborhoods in
mammalian chromosomes. Cell *159*, 374-387.

Ernst, J., Kheradpour, P., Mikkelsen, T.S., Shoresh, N., Ward, L.D., Epstein, C.B., Zhang, X., Wang, L.,
Issner, R., Coyne, M., *et al.* (2011). Mapping and analysis of chromatin state dynamics in nine human
cell types. Nature *473*, 43-49.

Fang, C., Wang, Z., Han, C., Safgren, S.L., Helmin, K.A., Adelman, E.R., Serafin, V., Basso, G., Eagen, K.P.,
Gaspar-Maia, A., *et al.* (2020). Cancer-specific CTCF binding facilitates oncogenic transcriptional
dysregulation. Genome Biol *21*, 247.

Filippova, G.N., Fagerlie, S., Klenova, E.M., Myers, C., Dehner, Y., Goodwin, G., Neiman, P.E., Collins, S.J.,
and Lobanenkov, V.V. (1996). An exceptionally conserved transcriptional repressor, CTCF, employs
different combinations of zinc fingers to bind diverged promoter sequences of avian and mammalian
c-myc oncogenes. Mol Cell Biol *16*, 2802-2813.

Fornes, O., Castro-Mondragon, J.A., Khan, A., van der Lee, R., Zhang, X., Richmond, P.A., Modi, B.P.,
Correard, S., Gheorghe, M., Baranašić, D., *et al.* (2019). JASPAR 2020: update of the open-access
database of transcription factor binding profiles. Nucleic Acids Research.

Fu, Y., Sinha, M., Peterson, C.L., and Weng, Z. (2008). The insulator binding protein CTCF positions 20
nucleosomes around its binding sites across the human genome. PLoS Genet *4*, e1000138.

Gabriele, M., Brandao, H.B., Grosse-Holz, S., Jha, A., Dailey, G.M., Cattoglio, C., Hsieh, T.S., Mirny, L.,
Zechner, C., and Hansen, A.S. (2022). Dynamics of CTCF- and cohesin-mediated chromatin looping
revealed by live-cell imaging. Science, eabn6583.

Ghirlando, R., and Felsenfeld, G. (2016). CTCF: making the right connections. Genes Dev *30*, 881-891.

Grant, C.E., Bailey, T.L., and Noble, W.S. (2011). FIMO: scanning for occurrences of a given motif.
Bioinformatics *27*, 1017-1018.

Grubert, F., Srivas, R., Spacek, D.V., Kasowski, M., Ruiz-Velasco, M., Sinnott-Armstrong, N., Greenside,
P., Narasimha, A., Liu, Q., Geller, B., *et al.* (2020). Landscape of cohesin-mediated chromatin loops in
the human genome. Nature *583*, 737-743.

Guo, Y., Monahan, K., Wu, H., Gertz, J., Varley, K.E., Li, W., Myers, R.M., Maniatis, T., and Wu, Q. (2012).
CTCF/cohesin-mediated DNA looping is required for protocadherin alpha promoter choice. Proc Natl
Acad Sci U S A *109*, 21081-21086.

Guo, Y., Xu, Q., Canzio, D., Shou, J., Li, J., Gorkin, D.U., Jung, I., Wu, H., Zhai, Y., Tang, Y., *et al.* (2015).
CRISPR inversion of CTCF sites alters genome topology and enhancer/promoter function. Cell *162*,
900-910.

Guo, Y.A., Chang, M.M., Huang, W., Ooi, W.F., Xing, M., Tan, P., and Skanderup, A.J. (2018). Mutation
hotspots at CTCF binding sites coupled to chromosomal instability in gastrointestinal cancers. Nat
Commun *9*, 1520.

He, C., Wang, X., and Zhang, M.Q. (2014). Nucleosome eviction and multiple co-factor binding predict
estrogen-receptor-alpha-associated long-range interactions. Nucleic Acids Res *42*, 6935-6944.

Hnisz, D., Day, D.S., and Young, R.A. (2016). Insulated Neighborhoods: Structural and Functional Units
of Mammalian Gene Control. Cell *167*, 1188-1200.

Hu, G., Dong, X., Gong, S., Song, Y., Hutchins, A.P., and Yao, H. (2020). Systematic screening of CTCF
binding partners identifies that BHLHE40 regulates CTCF genome-wide distribution and long-range
chromatin interactions. Nucleic Acids Res *48*, 9606-9620.

Huang, D., Wang, Z., Zhou, Y., Liang, Q., Sham, P.C., Yao, H., and Li, M.J. (2021a). vSampler: fast and

annotation-based matched variant sampling tool. Bioinformatics *37*, 1915-1917.

Huang, H., Zhu, Q., Jussila, A., Han, Y., Bintu, B., Kern, C., Conte, M., Zhang, Y., Bianco, S., Chiariello,
A.M., *et al.* (2021b). CTCF mediates dosage- and sequence-context-dependent transcriptional
insulation by forming local chromatin domains. Nat Genet *53*, 1064-1074.

Ibn-Salem, J., and Andrade-Navarro, M.A. (2019). 7C: Computational chromosome conformation
capture by correlation of ChIP-seq at CTCF motifs. BMC Genomics *20*, 777.

Islam, Z., Saravanan, B., Walavalkar, K., Farooq, U., Singh, A.K., Sabarinathan, R., Thakur, J., Pandit, A.,
Henikoff, S., and Notani, D. (2023). Active enhancers strengthen insulation by RNA-mediated CTCF
binding at chromatin domain boundaries. Genome Res.

Kai, Y., Andricovich, J., Zeng, Z., Zhu, J., Tzatsos, A., and Peng, W. (2018). Predicting CTCF-mediated
chromatin interactions by integrating genomic and epigenomic features. Nat Commun *9*, 4221.

Katainen, R., Dave, K., Pitkanen, E., Palin, K., Kivioja, T., Valimaki, N., Gylfe, A.E., Ristolainen, H.,
Hanninen, U.A., Cajuso, T., *et al.* (2015). CTCF/cohesin-binding sites are frequently mutated in cancer.
Nat Genet *47*, 818-821.

Kentepozidou, E., Aitken, S.J., Feig, C., Stefflova, K., Ibarra-Soria, X., Odom, D.T., Roller, M., and Flicek, P.
(2020). Clustered CTCF binding is an evolutionary mechanism to maintain topologically associating
domains. Genome Biol *21*, 5.

Kharchenko, P.V., Tolstorukov, M.Y., and Park, P.J. (2008). Design and analysis of ChIP-seq experiments
for DNA-binding proteins. Nat Biotechnol *26*, 1351-1359.

Kim, Y., Shi, Z., Zhang, H., Finkelstein, I.J., and Yu, H. (2019). Human cohesin compacts DNA by loop
extrusion. Science *366*, 1345-1349.

Kircher, M., Witten, D.M., Jain, P., O'Roak, B.J., Cooper, G.M., and Shendure, J. (2014). A general
framework for estimating the relative pathogenicity of human genetic variants. Nat Genet *46*,
310-315.

Kuang, S., and Wang, L. (2021). Deep learning of sequence patterns for CCCTC-binding
factor-mediated chromatin loop formation. J Comput Biol *28*, 133-145.

Lee, R., Kang, M.K., Kim, Y.J., Yang, B., Shim, H., Kim, S., Kim, K., Yang, C.M., Min, B.G., Jung, W.J., *et al.*
(2022). CTCF-mediated chromatin looping provides a topological framework for the formation of
phase-separated transcriptional condensates. Nucleic Acids Res *50*, 207-226.

Li, G., Chen, Y., Snyder, M.P., and Zhang, M.Q. (2017). ChIA-PET2: a versatile and flexible pipeline for
ChIA-PET data analysis. Nucleic Acids Res *45*, e4.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics *25*, 1754-1760.

Li, Y., Haarhuis, J.H.I., Sedeno Cacciatore, A., Oldenkamp, R., van Ruiten, M.S., Willems, L., Teunissen,
H., Muir, K.W., de Wit, E., Rowland, B.D., *et al.* (2020). The structural basis for cohesin-CTCF-anchored
loops. Nature *578*, 472-476.

Liu, E.M., Martinez-Fundichely, A., Diaz, B.J., Aronson, B., Cuykendall, T., MacKay, M., Dhingra, P., Wong,
E.W.P., Chi, P., Apostolou, E., *et al.* (2019). Identification of Cancer Drivers at CTCF Insulators in 1,962
Whole Genomes. Cell Syst *8*, 446-455 e448.

Lizio, M., Harshbarger, J., Shimoji, H., Severin, J., Kasukawa, T., Sahin, S., Abugessaisa, I., Fukuda, S.,
Hori, F., Ishikawa-Kato, S., *et al.* (2015). Gateways to the FANTOM5 promoter level mammalian
expression atlas. Genome Biol *16*, 22.

Lobanenkov, V.V., Nicolas, R.H., Adler, V.V., Paterson, H., Klenova, E.M., Polotskaja, A.V., and Goodwin,
G.H. (1990). A novel sequence-specific DNA binding protein which interacts with three regularly

spaced direct repeats of the CCCTC-motif in the 5'-flanking sequence of the chicken c-myc gene.
Oncogene *5*, 1743-1753.
Luan, J., Xiang, G., Gomez-Garcia, P.A., Tome, J.M., Zhang, Z., Vermunt, M.W., Zhang, H., Huang, A.,
Keller, C.A., Giardine, B.M., *et al.* (2021). Distinct properties and functions of CTCF revealed by a
rapidly inducible degron system. Cell Rep *34*, 108783.
Lundberg, S.M., Erion, G., Chen, H., DeGrave, A., Prutkin, J.M., Nair, B., Katz, R., Himmelfarb, J., Bansal,
N., and Lee, S.-I. (2020). From local explanations to global understanding with explainable AI for trees.
Nature Machine Intelligence *2*, 56-67.
Lundberg, S.M., Nair, B., Vavilala, M.S., Horibe, M., Eisses, M.J., Adams, T., Liston, D.E., Low, D.K.,
Newman, S.F., Kim, J., *et al.* (2018). Explainable machine-learning predictions for the prevention of
hypoxaemia during surgery. Nat Biomed Eng *2*, 749-760.
Lv, H., Dao, F.Y., Zulfiqar, H., Su, W., Ding, H., Liu, L., and Lin, H. (2021). A sequence-based deep learning
approach to predict CTCF-mediated chromatin loop. Brief Bioinform *22*.
Matthews, B.J., and Waxman, D.J. (2018). Computational prediction of CTCF/cohesin-based intra-TAD
loops that insulate chromatin contacts and gene expression in mouse liver. Elife *7*.
Nakahashi, H., Kieffer Kwon, K.R., Resch, W., Vian, L., Dose, M., Stavreva, D., Hakim, O., Pruett, N.,
Nelson, S., Yamane, A., *et al.* (2013). A genome-wide map of CTCF multivalency redefines the CTCF
code. Cell Rep *3*, 1678-1689.
Nichols, M.H., and Corces, V.G. (2015). A CTCF code for 3D genome architecture. Cell *162*, 703-705.
Nora, E.P., Lajoie, B.R., Schulz, E.G., Giorgetti, L., Okamoto, I., Servant, N., Piolot, T., van Berkum, N.L.,
Meisig, J., Sedat, J., *et al.* (2012). Spatial partitioning of the regulatory landscape of the X-inactivation
centre. Nature *485*, 381-385.
Oh, S., Shao, J., Mitra, J., Xiong, F., D'Antonio, M., Wang, R., Garcia-Bassets, I., Ma, Q., Zhu, X., Lee, J.H.,
et al. (2021). Enhancer release and retargeting activates disease-susceptibility genes. Nature *595*,
735-740.
Oki, S., Ohta, T., Shioi, G., Hatanaka, H., Ogasawara, O., Okuda, Y., Kawaji, H., Nakaki, R., Sese, J., and
Meno, C. (2018). ChIP-Atlas: a data-mining suite powered by full integration of public ChIP-seq data.
EMBO Rep *19*.
Ong, C.T., and Corces, V.G. (2014). CTCF: an architectural protein bridging genome topology and
function. Nat Rev Genet *15*, 234-246.
Ortabozkoyun, H., Huang, P.Y., Cho, H., Narendra, V., LeRoy, G., Gonzalez-Buendia, E., Skok, J.A., Tsirigos,
A., Mazzoni, E.O., and Reinberg, D. (2022). CRISPR and biochemical screens identify MAZ as a cofactor
in CTCF-mediated insulation at Hox clusters. Nat Genet *54*, 202-212.
Oti, M., Falck, J., Huynen, M.A., and Zhou, H. (2016). CTCF-mediated chromatin loops enclose
inducible gene regulatory domains. BMC Genomics *17*, 252.
Phanstiel, D.H., Van Bortle, K., Spacek, D., Hess, G.T., Shamim, M.S., Machol, I., Love, M.I., Aiden, E.L.,
Bassik, M.C., and Snyder, M.P. (2017). Static and Dynamic DNA Loops form AP-1-Bound Activation
Hubs during Macrophage Development. Mol Cell *67*, 1037-1048 e1036.
Phillips, J.E., and Corces, V.G. (2009). CTCF: master weaver of the genome. Cell *137*, 1194-1211.
Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics *26*, 841-842.
Rao, S.S., Huntley, M.H., Durand, N.C., Stamenova, E.K., Bochkov, I.D., Robinson, J.T., Sanborn, A.L.,
Machol, I., Omer, A.D., Lander, E.S., *et al.* (2014). A 3D map of the human genome at kilobase
resolution reveals principles of chromatin looping. Cell *159*, 1665-1680.

Rentzsch, P., Witten, D., Cooper, G.M., Shendure, J., and Kircher, M. (2019). CADD: predicting the
deleteriousness of variants throughout the human genome. Nucleic Acids Res *47*, D886-D894.

Rhee, H.S., and Pugh, B.F. (2011). Comprehensive genome-wide protein-DNA interactions detected at
single-nucleotide resolution. Cell *147*, 1408-1419.

Rheinbay, E., Nielsen, M.M., Abascal, F., Wala, J.A., Shapira, O., Tiao, G., Hornshoj, H., Hess, J.M., Juul,
R.I., Lin, Z., *et al.* (2020). Analyses of non-coding somatic drivers in 2,658 cancer whole genomes.
Nature *578*, 102-111.

Ribeiro-Dos-Santos, A.M., Hogan, M.S., Luther, R.D., Brosh, R., and Maurano, M.T. (2022). Genomic
context sensitivity of insulator function. Genome Res *32*, 425-436.

Sadowski, M., Kraft, A., Szalaj, P., Wlasnowolski, M., Tang, Z., Ruan, Y., and Plewczynski, D. (2019).
Spatial chromatin architecture alteration by structural variations in human genomes at the population
scale. Genome Biol *20*, 148.

Salameh, T.J., Wang, X., Song, F., Zhang, B., Wright, S.M., Khunsriraksakul, C., Ruan, Y., and Yue, F.
(2020). A supervised learning framework for chromatin loop detection in genome-wide contact maps.
Nat Commun *11*, 3428.

Sanborn, A.L., Rao, S.S., Huang, S.C., Durand, N.C., Huntley, M.H., Jewett, A.I., Bochkov, I.D.,
Chinnappan, D., Cutkosky, A., Li, J., *et al.* (2015). Chromatin extrusion explains key features of loop and
domain formation in wild-type and engineered genomes. Proc Natl Acad Sci U S A *112*, E6456-6465.

Shukla, S., Kavak, E., Gregory, M., Imashimizu, M., Shutinoski, B., Kashlev, M., Oberdoerffer, P.,
Sandberg, R., and Oberdoerffer, S. (2011). CTCF-promoted RNA polymerase II pausing links DNA
methylation to splicing. Nature *479*, 74-79.

Vostrov, A.A., and Quitschke, W.W. (1997). The zinc finger protein CTCF binds to the APBbeta domain
of the amyloid beta-protein precursor promoter. Evidence for a role in transcriptional activation. J Biol
Chem *272*, 33353-33359.

Wang, J., Huang, D., Zhou, Y., Yao, H., Liu, H., Zhai, S., Wu, C., Zheng, Z., Zhao, K., Wang, Z., *et al.* (2020).
CAUSALdb: a database for disease/trait causal variants identified using summary statistics of
genome-wide association studies. Nucleic Acids Res *48*, D807-D816.

Wang, W., Gao, L., Ye, Y., and Gao, Y. (2021). CCIP: Predicting CTCF-mediated chromatin loops with
transitivity. Bioinformatics.

Wang, Y., Song, F., Zhang, B., Zhang, L., Xu, J., Kuang, D., Li, D., Choudhary, M.N.K., Li, Y., Hu, M., *et al.*
(2018). The 3D Genome Browser: a web-based browser for visualizing 3D genome organization and
long-range chromatin interactions. Genome Biol *19*, 151.

Weintraub, A.S., Li, C.H., Zamudio, A.V., Sigova, A.A., Hannett, N.M., Day, D.S., Abraham, B.J., Cohen,
M.A., Nabet, B., Buckley, D.L., *et al.* (2017). YY1 Is a structural regulator of enhancer-promoter Loops.
Cell *171*, 1573-1588 e1528.

Xi, W., and Beer, M.A. (2021). Loop competition and extrusion model predicts CTCF interaction
specificity. Nat Commun *12*, 1046.

Yi, X., Zheng, Z., Xu, H., Zhou, Y., Huang, D., Wang, J., Feng, X., Zhao, K., Fan, X., Zhang, S., *et al.* (2021).
Interrogating cell type-specific cooperation of transcriptional regulators in 3D chromatin. iScience *24*,
103468.

Yin, M., Wang, J., Wang, M., Li, X., Zhang, M., Wu, Q., and Wang, Y. (2017). Molecular mechanism of
directional CTCF recognition of a diverse range of genomic sites. Cell Res *27*, 1365-1377.

Zhang, R., Wang, Y., Yang, Y., Zhang, Y., and Ma, J. (2018). Predicting CTCF-mediated chromatin loops
using CTCF-MP. Bioinformatics *34*, i133-i141.

Zheng, R., Wan, C., Mei, S., Qin, Q., Wu, Q., Sun, H., Chen, C.H., Brown, M., Zhang, X., Meyer, C.A., *et al.*

(2019). Cistrome Data Browser: expanded datasets and new tools for gene regulatory analysis. Nucleic

Acids Res *47*, D729-D735.

Figure Legends

Figure 1. Structure and performance of DeepAnchor model

A. Schematic view of DeepAnchor model. Base-wise chromatin features and DNA sequences are extracted for all candidate CBSs by motif scanning. Positive (Pos) and negative (Neg) dataset is constructed by considering both CTCF ChIP-seq peaks and targeted chromosome intervals, including ChromHMM insulator-associated CBS (insulator CBS), cohesin ChIP-seq signal-associated CBS (cohesin CBS), and cohesin ChIA-PET loop-associated CBS (loop CBS). 1D-CNN model is then used to train a classifier for distinguishing positive CBSs (CTCF-mediated CREs) versus other ones. Probability of CBSs being insulator/cohesin/loop-associated can be calculated and used as DeepAnchor score for downstream analyses. **Related terminologies**: **Insulator,** a enhancer blocker or a barrier between heterochromatin and euchromatin; **Chromatin loop,** during the interphase of cell, the condensed chromatin form 3D structure within cell nucleus. The basic loop-like structure is called chromatin loop; **Loop anchor,** given a chromatin loop detected by ChIA-PET, we call the endpoints of loop on chromosome as loop anchor; **Insulator/cohesin/loop CBS**; by using different targeted regions, we get different P/N dataset and train different DeepAnchor models. The CBS predicted by different models will be named by particular targeted region.

B. Cross-validation ROC curves based on GM12878, K562 and H1-hESC datasets, respectively, among three types of CBSs.

C. Cross-sample ROC curves for DeepAnchor models on different cell types, among three types of CBSs.

D. Correlation between DeepAnchor scores for three cell type-specific models, among three types of CBSs.

E. Comparison of number of enhancers around CBSs between predicted Pos and Neg CTCF-mediated CREs in loop CBS model.

F. Strand-oriented asymmetric pattern of enhancer enrichment at predicted Pos and Neg CTCF-mediated CREs in loop CBS model.

G. Position and strand preference of TAD boundary enrichment by measuring the distance between each CBS and the 5' end of TAD it belongs at predicted Pos and Neg CTCF-mediated CREs in loop CBS model.

H. The intersection of the predicted CTCF-mediated CREs among three CBS types at different thresholds.

Figure 2. Base-wise analysis of sequence and chromatin patterns across different types of CTCF binding

A. Feature importance analysis of top 20 features at base-wise level among three types of CBSs. Heatmap: Average absolute feature Shapley values of each position at ±500bp of CBS. Barplot: Summation of absolute feature Shapley values across ±500bp of CBS.

B. Base-wise Shapley value distribution for three representative features across different types of CBSs. EncOCctcfPval: P-value (PHRED-scale) of CTCF evidence for open chromatin; EncNucleo: Maximum of ENCODE Nucleosome position track score; GerpN: Neutral evolution score defined by GERP++. Please refer to more feature descriptions

from supplementary table.

C. Comparison of DNA motifs associated with positive CBSs among three types of CBSs

in this study and a commonly used conventional CBS.

Figure 3. Performance evaluation of LoopAnchor for CTCF-anchored loop

prediction

A. ROC curves, Precision-Recall curves and associated AUCs among LoopAnchor and

five state-of-the-art methods. All supervised models, such as LoopAnchor, LEM, Lollipop

and CTCF-MP were trained on GM12878 RAD21 ChIA-PET data and independently

tested on K562 RAD21 ChIA-PET data.

B. Correlation between predicted scores and real RAD21 ChIA-PET loop intensity on

K562.

C. Gain and lost loops from monocyte to macrophage differentiation by comparing

LoopAnchor prediction with Hi-C observation. log10(Fold change) is the transformed fold

change of predicted loop intensity for specific loop between macrophage and monocyte;

log10(Monocyte) is the transformed Hi-C loop score observed in monocyte; blue dot is lost

Hi-C loop; orange dot is gained Hi-C loop.

D. Example of loops predicted by LoopAnchor at *JAG1* locus. For Hi-C loops, line color is

used to distinguish gained and static loops. For LoopAnchor, linewidth represents the

predicted loop intensity and the loop with fold change of intensity >3 was marked with red

color.

Figure 4. The landscape of predicted CTCF-anchored loops across 32 human

tissues and 20 cell types

A. Overview of predicted CTCF-anchored loops across 168 biosamples (columns) and

biological conditions (rows).

B. Classification of CTCF-anchored loops according to their shared patterns. All predicted

loops were classified into four categories, including tissue/cell type-specific, tissue/cell

type-relatively specific, tissue/cell type-relatively shared, and tissue/cell type-shared.

C. Comparison of loop intensity score for loops in four categories. The cumulative

probability was calculated and the two-tailed Mann-Whitney U test was used to test the

significance.

D. Validation of tissue distribution for loops across four categories using predicted

chromatin loops by Peakachu for 42 human tissue/cell types. The cumulative probability

was calculated and the two-tailed Mann-Whitney U test was used to test the significance.

1004

Figure 5. Disease-causal variants and somatic hotspots enrichment

A. Enrichment significance distribution for autoimmune disease-causal variants among

different normal tissues. The tissues were ordered by their average P-values.

B. Comparison of fold change distribution among different somatic mutation recurrence

categories. The two-tailed Mann-Whitney U test was used to test the significance.

C-E. Comparison of overlapped percentage with somatic hotspots using different datasets

from ATACseq-AWG (C), PCAWG (D), and CNCDriver (E). The paired two-tailed

Mann-Whitney U test was used to test the significance.

A

Peak

0   50,000   100,000   150,000   200,000

Loop

0   10,000   20,000   30,000

Source
ChIP-Atlas (41)
CistromDB (15)
ENCODE (112)

State
Cancer (55)
Normal (113)

Tissue
Adipose (3)
Adrenal (1)
Blood (33)
Blood vessel (9)
Bone (2)
Brain (11)
Breast (8)
Bronchus (1)
Cervix (1)
Colon (7)
Esophagus (4)

Eye (3)
Gingival (1)
Heart (4)
Kidney (5)
Liver (4)
Lung (12)
Lymphoid tissue (1)
Muscle (7)
Other (5)
Ovary (2)
Pancreas (4)

Placenta (1)
Prostate (11)
Skin (11)
Spinal cord (1)
Spleen (2)
Stomach (1)
Testis (1)
Thyroid (1)
Uterus (3)
Vagina (1)
Unspecified (7)

Cell type
Adipose cell (1)
Astrocyte (4)
B lymphocyte (12)
Embryonic stem cell (5)
Endothelium (3)
Epithelium (45)
Erythrocyte (3)
Fibroblast (20)
Granulocyte (3)
Hematopoietic stem cell (1)
Hepatocyte (1)

Keratinocyte (3)
Lymphoblast (3)
Mesothelium (1)
Monocyte (2)
Muscle cell (6)
Neural cell (11)
Osteoblast (11)
Plasma cell (4)
T lymphocyte (3)
Unspecified (36)

B Percentage of loops vs Loop commonality

n = 52,216
n = 10,016
n = 6,802
n = 6,179

Loops
8.2%
9.0%
13.3%
69.4%

C Cumulative probability vs Loop intensity score

n	Mean	Median
904,741	0.028	0.024
703,825	0.018	0.016
607,026	0.015	0.014
515,301	0.013	0.012

D Cumulative probability vs Number of tissues

n	Mean	Median
3,694	4.743	4
3,270	3.496	3
3,936	2.986	2
11,122	2.314	2

Tissue/cell type-shared
Tissue/cell type-relatively shared
Tissue/cell type-relatively specific
Tissue/cell type-specific