The effect of *Drosophila attP40* background on the glomerular organization of Or47b olfactory receptor neurons

Qichen Duan[1], Rachel Estrella[1], Allison Carson[1], Yang Chen[1], Pelin C. Volkan[1, #]

[1]Department of Biology, Duke University, Durham, North Carolina 27708

[#]Correspondence: pelin.volkan@duke.edu

Running Title

attP40 background effect on ORN axons

Keywords

Drosophila, *attP40*, Olfactory receptor neuron, Axon terminal organization

Abstract

Bacteriophage integrase-directed insertion of transgenic constructs into specific genomic loci has been widely used by *Drosophila* community. The *attP40* landing site located on the second chromosome gained popularity because of its high inducible transgene expression levels. Here, unexpectedly, we found that homozygous *attP40* chromosome disrupts normal glomerular organization of Or47b olfactory receptor neuron (ORN) class in *Drosophila*. This effect is not likely to be caused by the loss of function of *Msp300*, where the *attP40* docking site is inserted. Moreover, the a*ttP40* background seems to genetically interact with the second chromosome *Or47b-GAL4* driver, which results in a similar glomerular defect. Whether the ORN phenotype is caused by the neighboring genes around *Msp300* locus in the presence of *attP40*-based insertions or a second unknown mutation in the *attP40* background remains elusive. Our findings tell a cautionary tale about using this popular transgenic landing site, highlighting the importance of rigorous controls to rule out the *attP40* landing site-associated background effects.

Introduction

RNA interference (RNAi)-based genetic screens provide scientists with powerful tools to identify genes involved in various biological processes (HOUSDEN *et al.* 2017). Binary expression systems, such as the *GAL4/UAS* system, induce the expression of various effectors in the desired cell populations (BRAND AND PERRIMON 1993). In Drosophila carrying transgenes for both cell-type-specific promoter-driven *GAL4* (driver) and *UAS-RNAi,* GAL4 protein binds *UAS* sites and drives *RNAi* expression, disrupting the expression and function of the target gene (BRAND AND PERRIMON 1993). As RNAi-based knockdown methods were becoming popular, efforts were initiated to make transgenic libraries of flies carrying *UAS-RNAi* targeting all the genes in the genome (DIETZL *et al.* 2007; NI *et al.* 2009; NI *et al.* 2011; PERKINS *et al.* 2015). These genome-wide libraries were then followed by efforts to generate thousands of *GAL4* lines that restrict expression to cellular subpopulations, enabling loss-of-function screens in cells of interest.

Among the RNAi collections, stocks from Transgenic RNAi Project (TRiP) have gained popularity because of their targeted integration of *UAS-RNAi* transgenes into the genome, efficient expression induced by appropriate *GAL4* drivers in different tissues, and high specificity with minimal expected off-target effects (MARKSTEIN *et al.* 2008; NI *et al.* 2008; PERKINS *et al.* 2015). To expedite the generation of transgenic libraries, two predetermined chromosomal docking sites were targeted for recombination events that insert *UAS-RNAi* transgenes: *attP40* on the second chromosome and *attP2* on the third chromosome (MARKSTEIN *et al.* 2008). With the presence of bacteriophage-originated phiC31 integrase (by co-injection of integrase mRNA or germline-expressing transgenic integrase), the *UAS-RNAi* construct can be inserted into the corresponding docking sites (GROTH *et al.* 2004; NI *et al.* 2008). These two sites, *attP40* and *attP2*, are selected because they exhibit optimal inducible expression levels upon binding with diverse tissue-specific *GAL4* drivers (MARKSTEIN *et al.* 2008). Therefore, in addition to the TRiP *UAS-RNAi* library, many other transgenes, including tissue-specific drivers (*GAL4*, *QF*, *LexA*) and *UAS/QUAS/LexAop-effectors/reporters* are also routinely integrated into these two landing sites (ZIRIN *et al.* 2020).

Given the widespread use of transgenic flies with *attP40* and *attP2* backbones, and the lesson learned from another popular *UAS-RNAi* collection with reported non-specific effects due to transgenic docking sites (GREEN *et al.* 2014; VISSERS *et al.* 2016), we must be more cognizant of potential phenotypic influences from these genetic backgrounds. Both *attP40* and *attP2* docking sites are in chromosomal regions populated by many genes. These sites, like any insertion into the genome, can disrupt function of nearby genes. More specifically, the *attP40* site is located within one of the large introns of *Msp300* gene while *attP2* site is inserted in the 5' untranslated region (UTR) of *Mocs1* gene (LARKIN *et al.* 2020). Both Msp300 and Mocs1 have critical biological roles. Specifically, Msp300 is the *Drosophila melanogaster* orthologue of mammalian Nesprins, which organize postsynaptic cytoskeleton scaffold and are required for stabilization of new synapses (ELHANANY-TAMIR *et al.* 2012; MOREL *et al.* 2014; TITLOW *et al.* 2020; ZHENG *et al.* 2020). Mocs1 is involved in Mo-molybdopterin cofactor biosynthetic process and inter-male aggressive behaviors (GAUDET *et al.* 2011; RAMIN *et al.* 2019). It is unclear how the insertion of various transgenic constructs into

attP40 and attP2 docking sites would affect the function of these host genes which may further result in phenotypic defects.

Indeed, recent studies have raised issues related to landing site-associated effects. For example, van der Graaf et al. showed flies bearing two copies of attP40-derived insertions also show decreased Msp300 transcript levels (VAN DER GRAAF et al. 2022). In addition, this study also reported defects in muscle nuclei spacing in larval stages in the attP40 homozygous background, which phenocopies Msp300 mutants (VAN DER GRAAF et al. 2022). These results suggest that the attP40 docking site and attP40-based transgenes are insertional mutations of Msp300 gene (VAN DER GRAAF et al. 2022). Another study reported that attP40 flies show resistance to cisplatin-induced neuronal damage, compared to the attP2 background (GROEN et al. 2022). This study tied the effect to the reduced ND-13A (NADH dehydrogenase 13 kDA subunit, a component of mitochondrial complex I) expression in attP40 homozygous flies (GROEN et al. 2022). It is noteworthy that ND-13A flanks the 5' UTR of Msp300 and is downstream of attP40 docking site. Together, these results imply the integration of attP40 docking site significantly changes the local transcriptional state and interferes with the transcription of surrounding genes.

During a GAL4-driven UAS-RNAi screen for olfactory neuron axon organization, we observed an axon terminal phenotype that is associated with the attP40 background. The phenotype occurs in the flies homozygous for the attP40 docking site alone or with various transgenic insertions, independent of the identity of the transgene. Notably, the phenotype observed in the attP40 background appears to be recessive but is independent of the Msp300 function, possibly implicating other attP40 background mutations nearby or in other locations on the second chromosome. Though the nature of the mutation is unclear, the background effects should be mitigated by designing more rigorous controls to interpret phenotypic data obtained using reagents in concert with the attP40 background.

Results

We used the Drosophila olfactory receptor neurons (ORNs) as a model to understand the molecular mechanisms underlying neuronal circuit assembly. In Drosophila, each class of ORNs expresses a unique olfactory receptor (Or) gene, and ORN axons target to the brain antennal lobe within class-specific and uniquely positioned synaptic units called glomeruli (HONG AND LUO 2014; BARISH AND VOLKAN 2015). To identify the molecular players contributing to the glomerular organization of the ORNs, we genetically screened genes encoding cell adhesion molecules whose expression levels increase over pupal development in the antennae (BARISH et al. 2018). Among these, beat and side gene families drew our attention because they encode the Ig superfamily proteins, form a heterophilic interacting protein network, and have been previously revealed to be involved in neuronal adhesion (FAMBROUGH AND GOODMAN 1996; PIPES et al. 2001; SINK et al. 2001; DE JONG et al. 2005; SIEBERT et al. 2009; ÖZKAN et al. 2013; LI et al. 2017; KINOLD et al. 2021). We obtained a collection of transgenic UAS-RNAi lines from TriP library deposited at the Bloomington Drosophila Stock Center (BDSC) and

crossed these lines with an established recombinant chromosome containing four
different *Or* promoter-driven *GAL4* transgenes (*Or47a-GAL4, Or47b-GAL4, Or23a-*
GAL4, Gr21a-GAL4, 4xOr-GAL4 for short, Figure 1A-D) together with a *UAS-Syt.GFP*
reporter. We examined the knockdown effect of candidate genes on axonal targeting of
these four ORN classes. The parent flies with a single copy of the *GAL4* drivers showed
wild type glomerular organization (Figure 1A). As an additional control, we also crossed
GAL4 driver lines to flies expressing the RNAi against a red fluorescent protein (RFP)
mCherry, which also exhibited no apparent defect in glomerular organization (Figure
1A).

From the screen, we found a strikingly recurrent phenotype, where the axon terminals of
Or47b ORNs invade the neighboring region, leading to an expanded round VA1v
glomerulus in contrast to the crescent shape in control brains (Figure 1A). This
phenotype was observed in two independent RNAi lines targeting the same gene, for
example, *beat-Ia* (Figure 1A). However, screening a list of *beat* and *side* family
members revealed a pattern for the phenotype, which only correlated with the second
chromosome *UAS-RNAi* transgenes, independent of the gene identity. Figure 1B
summarizes the screening results from *beat/side* gene families. All the RNAi lines
inserted at the second chromosome *attP40* site yielded the expanded VA1v glomerulus
phenotype, whereas none of the RNAi lines inserted at the third chromosome *attP2* site
showed this defect. Notably, there is one gene, *beat-IIIc*, with one *attP40*-derived RNAi
line and one *attP2*-derived RNAi line. Only the *attP40 UAS-RNAi* insertion gave rise to
the phenotype (Figure 1B). The same phenotype was also observed with randomly
selected TRiP *UAS-RNAi* lines inserted at the *attP40* site targeting genes without known
roles in ORN development (Figure S1A,B). To test whether this phenotype is caused by
specific effects of RNAi-mediated gene knockdown or simply by the presence of *attP40*-
derived insertions, we first crossed the same *Or47b-GAL4* driver line to a third *UAS-*
RNAi line from Vienna Drosophila Resource Center (VDRC) targeting *beat-Ia*, which
was generated by random P-element-mediated insertions (DIETZL *et al.* 2007). This non-
attP40 UAS-RNAi line could not reproduce the phenotype obtained by the *attP40*-
derived *UAS-RNAi* from the TRiP collection (Figure 1C). In addition, crossing the driver
line to an empty *attP40* site without any transgenes led to the same glomerular
expansion phenotype (Figure 1D). These results suggest that the Or47b ORN-specific
VA1v glomerular defect is independent of the RNAi-based knockdown of the genes
examined but caused by an effect from the *attP40*-derived chromosome.

Since we repeatedly obtained the VA1v glomerular phenotype with the second
chromosome *Or47b-GAL4*-driven *UAS-RNAi*, we also tested if crossing flies carrying
the same *Or47b-GAL4* transgene to various *attP40* derivatives could result in the same
phenotype (Figure 1E). Compared with the no *attP40* control (over a *CyO* balancer
chromosome), the *attP40* landing site with and without *UAS-RNAi* insertion, or a
ubiquitous promoter-driven gRNA targeting the *QUAS* sequence (control gRNA) all
produced the same VA1v glomerular defect when crossed to the second chromosome
Or47b-GAL4-driven *UAS-Syt.GFP* (Figure 1F). We also crossed *Or47b-GAL4 UAS-*
Syt.GFP chromosome to three lines carrying *UAS-rCD2.RFP* transgenic insertion at
three different chromosomal locations, *attP2* (on chr3), *attP5* (on chr2R), and *attP40* (on
chr2L). Only *Or47b-GAL4* over *UAS-rCD2.RFP* insertion at *attP40* resulted in

glomerular expansion phenotype while the insertions at *attP2 and attP5* appeared wild
type. These results again suggest that the VA1v glomerular defect is uniquely linked to
the *attP40*-associated insertions and is independent of the transgene or other attP
landing sites.

The glomerular organization defect could be caused by simply the presence of *attP40*
insertion or the genetic interaction between the *attP40* background and the
chromosome carrying the reporter transgene. To distinguish between these possibilities,
we examined the animals carrying a single copy of *attP40* insertion and a different
Or47b-GAL4 UAS-Syt.GFP reporters on the third chromosome. In these animals, VA1v
glomerulus appeared normal (Figure 1H,I). The observation that a single copy of *attP40*
is not sufficient to produce a glomerular phenotype indicates that the *attP40* effects on
VA1v glomerulus are not dominant. Rather, they point to a combinatorial effect of the
second chromosome with *Or47b-GAL4 UAS-Syt.GFP* over the chromosome with the
empty or transgene-carrying *attP40* docking site on the glomerular phenotype (Figure
1H,I). Thus, we conclude that the *attP40* chromosome genetically interacts with the
second chromosome reporters to disrupt VA1v glomerular organization.

As the *attP40* effect appears to be recessive, we next examined if animals homozygous
for the *attP40* sites display any VA1v glomerular defects. To bypass the glomerular
defects arising from the genetic interactions between *attP40* and the second
chromosome reporters, we used the third chromosome *Or47b-GAL4 UAS-Syt.GFP*
reporter to visualize VA1v glomerulus. Surprisingly, homozygous *attP40* derivatives or
attP40 empty docking site alone produced strong axon terminal defects (Figure 2A-F).
In contrast, flies heterozygous for the *attP40* site with or without transgenes inserted
appeared wild type (Figure 2A-F). Most of the brains homozygous for the *attP40* site
with or without insertions displayed a dorsally positioned VA1v glomerulus (Figure 2B,E,
middle panels; Figure 2C,F), whereas a small proportion also exhibited an expanded
glomerulus (Figure 2B,E, right panels; Figure 2C,F). Given that the *attP40* site is located
within an intron of *Msp300* gene, we posited that it likely disrupts *Msp300* function.
Msp300 encodes a Nesprin-like protein, which is required for proper positioning of
muscle nuclei and neuromuscular junction formation (ELHANANY-TAMIR *et al.* 2012;
MOREL *et al.* 2014). Single-cell RNA-seq datasets from ORNs also show broad
expression of *Msp300* across ORN classes (LI *et al.* 2022). We thus tested if the VA1v
glomerular defect is caused by the loss of *Msp300* function. We analyzed
transheterozygotes of empty *attP40* docking site over other mutant alleles of *Msp300*,
such as *Msp300$^{\Delta KASH}$* (which lacks the KASH domain (XIE AND FISCHER 2008;
ELHANANY-TAMIR *et al.* 2012)), *Msp300^{MI00111}, Msp300^{MI01145}* (two MIMIC-based alleles
predicted to disrupt most splice isoforms of *Msp300* transcripts (VENKEN *et al.* 2011)),
and *Msp300^{KG03631}* (a P-element-based insertion which is close to *attP40* landing site
(BELLEN *et al.* 2004)) (Figure 2G). However, none of these genetic combinations
recapitulated VA1v glomerular phenotype (Figure 2H). This indicates that the VA1v
glomerular defect is independent of the *Msp300* function and is likely caused by other
genes nearby affected by the *attP40* insertion or a second recessive mutation linked to
the *attP40* docking site.

We next sought to figure out the molecular basis of the genetic interaction between that
specific *Or47b-GAL4*-bearing chromosome and *attP40* chromosome. Flies

transheterozygous for *attP40* (empty or with insertions) over *4xOr-GAL4 UAS-Syt.GFP*
or *Or47b-GAL4 UAS-Syt.GFP* robustly exhibit VA1v glomerular phenotype. We infer
that the putative genetic lesion is directly caused by or genetically linked to the *Or47b-*
GAL4 transgene for three reasons: 1) a farther second site mutation would likely be lost
during meiotic recombination events to generate these stocks; 2) this *Or47b-GAL4*
recombined with other *UAS-reporters*, *UAS-mCD8.GFP* or *UAS-RFP*, over the *attP40*
derivatives exhibited the same phenotype (Figure S2); and 3) crossing the second
chromosome *Or47b-GAL4* transgene alone to the *UAS-rCD2.RFP* reporter inserted at
attP40 also reproduced the expanding glomerulus (Figure 3A), which rules out the
confounding effect from *UAS-Syt.GFP* transgene. Indeed, two copies of this *Or47b-*
GAL4 chromosome also results in similar glomerular phenotypes (Figure 3A). *Or47b-*
GAL4 transgene was generated by P-element-mediated genomic integration (Figure
3B) and the exact site of the insertion was not mapped (VOSSHALL *et al.* 2000). We used
inverse PCR (HUANG *et al.* 2009) to identify the insertion site and determine the gene
whose function is potentially disrupted (Figure 3C). We successfully recovered a piece
of genomic sequence immediately flanking the 3' end of the inserted P-element, which
is within the first intron of *Bacc* gene (chr2L:2753160, Figure 3D,E) and also a P-
element insertion-enriched region. We performed genomic PCR to further validate the
Bacc intronic insertion from both 3' and 5' ends of the P-element using different primer
pairs targeting the P-element ends and flanking *Bacc* sequences (Figure 3E). We
amplified the desired DNA fragments from *Or47b-GAL4*-bearing flies but not from w^{1118}
flies (Figure 3E,F). In addition, primer pairs targeting only *Bacc* sequences flanking the
insertion amplified the expected fragment only from w^{1118} files but not from *Or47b-GAL4*
flies (Figure 3E,F). We next tested if this intronic insertion affects *Bacc* transcriptional
levels using Quantitative Reverse Transcription-PCR (qRT-PCR). We extracted mRNA
from whole heads of the adult *Or47b-GAL4* homozygotes, as well as homozygous
attP2, *attP40*, and w^{1118} adults. qRT-PCR results showed that *Bacc* transcripts
normalized to the housekeeping gene *RpL13A* decrease by ~two fold in flies
homozygous for *Or47b-GAL4* but are not significantly altered in *attP2* or *attP40* animals
compared to w^{1118} controls (Figure 3G). In contrast, other housekeeping genes *Act5C*
and *Tbp* remain unchanged across these genotypes (Figure 3G). These results indicate
that: 1) second chromosome *Or47b-GAL4* transgene insertion disrupts *Bacc* gene
function; 2) homozygous and transheterozygous combinations of *Or47b-GAL4* and
attP40 backgrounds likely utilize distinct mechanisms to disrupt VA1v glomerular
organization.

Figure 1. Genetic interactions between *attP40* and *Or47b-GAL4* backgrounds on the second chromosome disrupt the glomerular organization of Or47b ORNs in the antennal lobes. (A, C & D) Confocal images of representative brains from a genetic screen to identify adhesion molecules involved in the glomerular organization of the Drosophila olfactory receptor neuron axon terminals. We crossed a second chromosome containing four different Drosophila olfactory receptor promoter-driven *GAL4s (Or47a-GAL4, Or47b-GAL4, Or23a-GAL4, Gr21a-GAL4)* together with a *UAS-Syt.GFP* reporter (*4xOr-GAL4>Syt.GFP*) to the indicated *UAS-RNAi* lines or *attP40* background flies. The parental driver chromosome over the *CyO* balancer was used as a no-RNAi control. The invading Or47b ORN axons are denoted with white arrowheads. **(B)** Summary of the phenotypical results from the genetic screen focusing on *beat/side* gene families. The Bloomington stock number and the transgenic docking site of each line are also listed. **(E, F)** Schematic in **(E)** shows the genotype of animals used in **(F)**, where each fly has one copy of the second chromosome carrying an *Or47b-GAL4* driver

and a *UAS-Syt.GFP* reporter, and one copy of the indicated second chromosome, either
a *CyO* balancer or *attP40* docking site derivatives. *attL* and *attR* sites are generated as
a result of transgene integration into *attP40* docking site. Confocal images of
representative brains are shown in **(F)**. **(G)** Confocal brain images of the indicated
genotypes. **(H, I)** Schematic in **(H)** shows the genotype of animals used in **(I)**, where
each fly has one copy of the second chromosome *UAS-beat-Ia RNAi* transgene inserted
into the *attP40* docking site, with one copy of *Or47b-GAL4 UAS-Syt.GFP*, either on the
second or third chromosome. Confocal images of representative brains are shown in **(I)**.
10-25 brains were examined in each genotype and the phenotypical penetrance is close
to 100% in each *attP40*-derived group.

A

attL *transgene* *attR* chr2

+

Or47b-GAL4 *UAS-Syt.GFP* chr3

attP40 *transgene* chr2

B *UAS-beat-Ia RNAi attP40 /CyO*

UAS-beat-Ia RNAi attP40 /UAS-beat-Ia RNAi attP40

VA1v

Syt.GFP
Ncad

VA1v expansion

VA1v dorsal shift

C

N= 12 28

****

Penetrance (%)

□ Wild-type
■ VA1v expansion
■ VA1v dorsal shift

UAS-RNAi attP40/CyO
UAS-RNAi attP40/UAS-RNAi attP40

D

chr2

+

Or47b-GAL4 *UAS-Syt.GFP* chr3

attP40 chr2

E *attP40/CyO*

attP40/attP40

Syt.GFP
Ncad

VA1v expansion

VA1v dorsal shift

F

N= 38 32

****

Penetrance (%)

□ Wild-type
■ VA1v expansion
■ VA1v dorsal shift

attP40/CyO
attP40/attP40

G

ΔKASH

Msp300-RB
Msp300-RD
Msp300-RE
Msp300-RF
Msp300-RG
Msp300-RH
Msp300-RI
Msp300-RJ
Msp300-RK
Msp300-RL
Msp300-RM

Msp300

KG03631
attP40
MI01145
MI00111

H *attP40/Msp300^{ΔKASH}*   *attP40/Msp300^{MI00111}*   *attP40/Msp300^{MI01145}*   *attP40/Msp300^{KG03631}*

Syt.GFP
Ncad

Figure 2. Homozygous *attP40* chromosome affects glomerular organization of Or47b ORNs independent of the *Msp300* function. (A-C) Schematic in **(A)** shows the genotype of animals used in **(B)**, where each fly has one or two copies of the second chromosome *UAS-beat-Ia RNAi* transgene inserted at the *attP40* docking site, with the third chromosome *Or47b-GAL4 UAS-Syt.GFP* transgenes. Confocal images of representative brains are shown in **(B)**. The percentage of the phenotypes is shown in **(C)**. ****, p<0.0001 after Fisher's exact test. **(D-F)** Schematic in **(D)** shows the genotype of animals used in **(E)**, where each fly has one or two copies of the second chromosome empty *attP40* docking site, with the third chromosome *Or47b-GAL4 UAS-Syt.GFP* transgenes. Confocal images *of* representative brains are shown in **(E)**. The percentage of the phenotypes is shown in **(F)**. ****, p<0.0001 after Fisher's exact test. N in **(C)** and **(F)** denotes the antennal lobes examined. **(G)** Schematic showing the *Msp300* genomic locus, the *attP40* docking site, three insertional *Msp300* mutations (*Msp300^{MI00111}, Msp300^{MI01145}, Msp300^{KG03631}*), and one deletion allele (*Msp300^{ΔKASH}*), each denoted with red lines. **(H)** Confocal images of representative brains of the indicated transheterozygous animals, with the *attP40* docking site over the indicated *Msp300* alleles. N = 11, 8, 4, 12 brains in each genotype group, from left to right.

Figure 3. Identification of the *Or47b-GAL4* transgene insertion site. (A) Confocal images of the indicated genotypes suggesting the second chromosome *Or47b-GAL4* transgene is accounting for the glomerular expansion phenotype. N = 16 brains in each genotype and the phenotypical penetrance is 100%. **(B)** The structure of the P-element vector containing *Or47b-GAL4* transgene (*Or47b* promoter fused with GAL4 coding sequence, denoted as *Or47bp::GAL4* in this and other panels of this figure) and a *mini-white* selectable marker. **(C)** Schematic illustrating the recovery of genomic sequences flanking P-element insertion by inverse PCR. **(D)** Schematic showing the sequencing results from one ligated genomic DNA template digested by Sau3AI restriction enzyme. A region within *Bacc* gene was recovered as the immediate sequencing flanking the 3'

end of the inserted P-element containing *Or47b-GAL4* transgene. **(E)** Schematic
showing the P-element insertion site within the first intron of *Bacc* gene. The arrows
indicate the primers used in a PCR assay to validate this insertion. **(F)** DNA gel showing
the amplicons by each indicated primer pairs from *Or47b-GAL4* transgenic flies or w^{1118}
control flies. Primer pairs FWD-1/REV-1 and FWD-2/REV-2 can amplify the specific
fragments from both *Or47b-GAL4* transgenic flies and w^{1118} control flies as expected
because the corresponding sequences exist in both. Primer pairs FWD-1/Plac1 and
Pry4/REV-2 only amplify specific products from *Or47b-GAL4* transgenic flies while fail to
work from w^{1118} control flies as one primer in each pair targets the sequence of P-
element which is not present in w^{1118} flies. Primer pair FWD-1/REV-2 works in w^{1118}
control flies but fails in *Or47b-GAL4* flies as the expected amplicon is too long to be
amplified in transgenic animals because of the integration of large P-element sequence.
(G) qPCR results comparing the indicated gene expression levels normalized by
RpL13A transcripts across different genotypes. ns, not significant. *, p-value < 0.05 after
two-way ANOVA followed by multiple comparison test. Error bars indicate mean ± SEM.
N = 3 biological replicates per genotype. Each biological replicate contain 14~20 fly
heads from equal number of males and females.

A *4xOr-GAL4, UAS-Syt.GFP/+;*
UAS-SMC3 RNAi attP2/+

4xOr-GAL4, UAS-Syt.GFP
/UAS-SMC3 RNAi attP40; +/+

Syt.GFP
Ncad

B *4xOr-GAL4, UAS-Syt.GFP/+;*
UAS-vtd RNAi attP2/+

4xOr-GAL4, UAS-Syt.GFP
/UAS-vtd RNAi attP40; +/+

**Figure S1. Additional *TRiP UAS-RNAi* results showing the *attP40* but not *attP2-*
specific glomerular organization phenotype. Similar to *Beat/Side* screening results in
Figure 1A,B, crossing *4xOr-GAL4, UAS-Syt.GFP* chromosome to two *UAS-SMC3 RNAi*
lines **(A)** and *UAS-vtd RNAi* lines **(B)** respectively gave rise to glomerular expansion
with *attP40* insertion but not *attP2* insertion. 5-9 brains were examined in each genotype
and the phenotypical penetrance is 100% in *attP40* groups.

**Figure S2. *attP40* over *Or47b-GAL4* recombined with *UAS-reporters* other than
UAS-Syt.GFP exhibits the same glomerular organization phenotype.** *Or47b-GAL4,
UAS-mCD8.GFP* **(A)** and *Or47b-GAL4, UAS-RFP* **(B)** over the *attP40* derivative
showed VA1v glomerular expansion compared with the respective no-*attP40* control. 5-
15 brains were examined in each genotype and the phenotypical penetrance is 100% in
attP40 groups.

Discussion

Here, we found that homozygous *attP40* chromosome leads to defective glomerular organization of ORNs. This defect is likely not caused by the loss of *Msp300* function, where the *attP40* site is inserted. Moreover, the a*ttP40* chromosome genetically interacts with a second chromosome carrying the *Or47b-GAL4* transgene, resulting in a similar ORN axon terminal defect. Though the exact genetic reasons and molecular mechanisms are unknown, our finding raises the critical issue with using this popular transgene landing site. Rigorous controls are needed to rule out the *attP40*-associated background effects, as discussed below.

The genetics underlying the Or47b ORN phenotypes

A recent study reported that flies homozygous for the *attP40*-derived insertions had 50% reduction in *Msp300* transcript levels and phenocopied the defects in larval muscle nuclei clustering in *Msp300* mutants (VAN DER GRAAF *et al.* 2022). As homozygotes of the *attP40* chromosome are defective in Or47b ORN axon terminal organization, we hypothesized that the *attP40*-affected *Msp300* gene is responsible for the defect. However, this is not the case as *attP40* over various *Msp300* mutations appeared phenotypically wild type, suggesting the *attP40* chromosome may carry an unannotated mutation responsible for *Msp300*-independent ORN glomerular disorganization.

The *attP40* docking site with or without transgene insertions may also disrupt other genes in the vicinity of *Msp300*. For example, in addition to *Msp300, attP40* docking site is flanked on the opposing side by *ND-13A,* which encodes a component of the mitochondria electron transport chain complex I. Thus, the *attP40* docking site alone or with transgene insertions may lead to a variety of phenotypes as a result of disrupted *ND-13A*. Indeed, Groen et al. reported that *attP40* flies exhibit resistance to cisplatin-induced neuronal damage mediated by the reduced expression of *ND-13A* (GROEN *et al.* 2022). Whether the glomerular defect is dependent on the *ND-13A* function is beyond the scope of this paper but needs to be tested in the future studies.

Surprisingly, we found transheterozygous animals with an *attP40* chromosome over the second chromosome *Or47b-GAL4* transgene produced similar but not identical glomerular abnormalities to *attP40* homozygotes. Additionally, *Or47b-GAL4* homozygotes exhibit comparable phenotypes with *Or47b-GAL4/attP40* transheterozygotes. These suggest several possible underlying genetic mechanisms. 1) *Or47b-GAL4* and *attP40* backgrounds harbor common mutations; 2) *Or47b-GAL4* and *attP40* backgrounds possess completely separate genetic lesions that genetically interact. The genetic interaction model is favored due to qualitatively distinguishable phenotypes between *Or47b-GAL4/attP40* animals and *attP40/attP40* ones. Furthermore, *Or47b-GAL4* transgene is inserted into an intron of *Bacc* gene, which encodes a tyramine-dependent nuclear regulator (CHEN *et al.* 2013), reducing its expression levels by about 50% in homozygotes. No change in transcript levels were observed in *attP40* homozygotes. *Bacc* mRNAs are abundant in brain tissues, comparable to housekeeping genes *Act5C* and *RpL13A* (Figure 3G). Our results imply its potential novel role in ORN axon pathfinding or glomerular patterning. Future

functional studies will determine whether the disruption of *Bacc* expression is causative
to the VA1v glomerular phenotype and the mechanisms by which *Bacc* mutations and
their genetic interactors in the *attP40* background result in glomerular defects.

Unique genetic sensitivity of VA1v glomerulus architecture.

One of the most peculiar observations from our study is that Or47b ORNs seem to be
particularly sensitive to changes in genetic background. In fact, VA1v glomerular
disruptions are not only restricted to the *attP40* background, but can be seen in many
other mutants with effects on ORN axon and synapse organization in the antennal lobes
(ANG *et al.* 2003; YAO *et al.* 2007; HONG *et al.* 2012; LI *et al.* 2013; HUESTON *et al.* 2016;
WU *et al.* 2017; XIE *et al.* 2019; HING *et al.* 2020). In addition to the variability of VA1v
glomerular architecture to genetic background effects, pheromone sensing Or47b ORNs
and the trichoid at4 sensillum that houses Or47b, Or88a, and Or65a/b/c ORNs are
developmentally special. At4 sensillum appears to be a developmentally default state
for all trichoid sensilla. For example, loss of transcription factor Rn function, normally
expressed in at1 and at3 ORNs, leads to a loss of at1 and at3 sensilla identity, and their
conversion to at4 sensillum identity (LI *et al.* 2013; LI *et al.* 2015; LI *et al.* 2016).
Similarly, Or47b ORNs in at4 sensillum appear to have a default identity, as mutants
in *Alh,* a chromatin factor, result in the conversion of Or88a and Or65a ORNs to Or47b
ORN fate (HUESTON *et al.* 2016). In addition to these findings that point to a
developmentally special state for pheromone sensing Or47b ORNs or "at" sensilla, the
glomeruli targeted by the trichoid ORNs are morphologically plastic. In Drosophila, they
are sexually dimorphic, appearing larger in males (STOCKINGER *et al.* 2005). In insects
such as moths trichoid glomeruli can form separate macro-glomerular complex outside
the antennal lobe of male brains (BERG *et al.* 1998). Given the developmental plasticity
of trichoid pheromone sensing ORNs and the developmental ground state of at4
sensilla, Or47b developmental trajectory might be particularly sensitive to genetic
background effects to accommodate adaptive developmental, behavioral and
evolutionary processes. On the other hand, we only examined Or47b VA1v glomerulus
in the *attP40* background, and *attP40* homozygotes possibly display structural defects in
other ORN classes and their glomeruli. Future studies will help identify these
phenotypes and the genetic lesions leading to *attP40*-associated phenotypes.

Addressing genetic background issues when using genetic reagents

To summarize, we found unexpected background effects of the *Drosophila attP40*
landing site on the ORN glomerular organization. In parallel with other recent studies
reporting other phenotypes arising from the *attP40* background, ranging from muscle
development to neuronal stress responses, such background effects should be
seriously considered in using *attP40*-derived flies. It is recommended to avoid using
homozygotes/double-copies of the *attP40*-based insertions. Researchers should also be
aware of the potential genetic interactions between the *attP40*-bearing chromosome
and the other homologous second chromosomes even if it doesn't contain any *attP40*
derivatives. Appropriate controls should be applied to override these caveats. For

example, when working with *GAL4/UAS-effector* binary system, it is better to use a GAL4-driven *UAS-neutral effector* (such as *UAS-RNAi* against neutral or non-fly genes inserted at the same docking site) as a negative control, rather than the widespread use of *GAL4* alone or *UAS-effector* alone controls. Transgenic rescue of RNAi-based gene knockdowns is not feasible due to targeting of rescue transgenes by the RNAi. Thus, use of full animal mutants or MARCM based clonal mutant analysis should be coupled with RNAi-based phenotypic analyses. Though the underlying genetic reasons remain elusive, studies demonstrated that the *attP40* landing site on the second chromosome affects the expression of multiple genes (GROEN *et al.* 2022; VAN DER GRAAF *et al.* 2022). Additional omics-based experiments in the future will be needed to determine all the genetic lesions in *attP40* strains that underly many phenotypic defects observed in this background. These studies will also reveal potential genetic alterations associated with glomerular defects, providing new insights into ORN axon pathfinding and glomerular organization.

Materials and Methods

Drosophila stocks and genetics

Drosophila were raised in classic molasses media provided by Archon Scientific. For the RNAi screen experiments, flies were raised at 28 °C to maximize the knockdown efficiency. Most of the other crosses were also kept at 28 °C, except for the experiments shown in Figure 1F,G and Figure 3A, which were conducted at room temperature (23 °C). After eclosion, the flies are aged for 5-7 days before dissection. In addition to the *UAS-RNAi* stocks from Bloomington Drosophila Stock Center (listed in Figure 1B), the following stocks are used: *UAS-RFP RNAi attP2* (BDSC# 35785), *UAS-beat-Ia RNAi #3* GD1386 (VDRC# 4544), *UAS-SMC3 RNAi attP2* (BDSC# 60017), *UAS-SMC3 RNAi attP40* (BDSC# 50899), *UAS-vtd RNAi attP2* (BDSC# 36786), *UAS-vtd RNAi attP40* (BDSC# 65229), *attP40* (BDSC# 36304), *attP2* (BDSC# 36303), *ctrl-gRNA attP40* (BDSC# 67539), *UAS-RFP attP40* (BDSC# 32222), *UAS-rCD2.RFP attP2* (BDSC# 56179), *UAS-rCD2.RFP attP5* (BDSC# 56180), *UAS-rCD2.RFP attP40* (BDSC# 56181), *Msp300^{ΔKASH}* (BDSC# 26781), *Msp300^{MI01145}* (BDSC# 53050), *Msp300^{MI00111}* (BDSC# 30623), *Msp300^{KG03631}* (BDSC# 13024); *Or47b-GAL4* (chr2, BDSC#9983), *Or47b-GAL4* (chr3, BDSC#9984), *Or43a-GAL4* (chr2), *Or47a-GAL4* (chr2) (VOSSHALL *et al.* 2000; FISHILEVICH AND VOSSHALL 2005), and *Gr21a-GAL4* (chr2) (SCOTT *et al.* 2001)) are gifts from Dr. Leslie Vosshall; *UAS-Syt.GFP* (chr2 or chr3), *UAS-mCD8.GFP*, *UAS-RFP* are Volkan lab stocks (BARISH *et al.* 2018). The line *Or47b-GAL4, Or47a-GAL4, Or43a-GAL4, Gr21a-GAL4, UAS-Syt.GFP/CyO* (short for *4xOr-GAL4>Syt.GFP*) was recombined and balanced from the above components.

Immunocytochemistry

Flies were sacrificed in 70% ethanol. Fly brains were then dissected in PBST buffer (0.2% Triton X-100 in 1X PBS), fixed in 4% paraformaldehyde for 30 mins, followed by washing with PBST for three 10-min cycles. Brains were incubated in the primary

antibody mix at 4 °C overnight, followed by three 20-min washes with PBST at room
temperature, then incubated in the secondary antibody mix at 4 °C overnight. The
brains were washed again by three 20-min wash with PBST before being mounted on
the slide for imaging. The blocking was done together with each antibody incubation,
with 1% natural goat serum mixed with primary and secondary antibodies, respectively.
The following primary antibodies were used: 1:1000 rabbit anti-GFP (Invitrogen), 1:20
rat anti-Ncad (DSHB); the following secondary antibodies were used: 1:1000 Alexa
Fluor 488 goat anti-rabbit IgG (Invitrogen), 1:200 Alexa Fluor 647 goat anti-rat IgG
(Invitrogen); all antibodies are diluted in PBST.

Confocal imaging and phenotypic quantification

Confocal imaging was performed by either Olympus Fluoview FV1000 microscope or
Zeiss 880 microscope. Brains were imaged across Z-axis from the posterior side to the
most anterior side of the antennal lobes, and all confocal sections were overlayed for
phenotypical analysis. The same set of imaging parameters was used between
experimental and control groups. The phenotype was qualitatively determined by
glomerular morphology, i.e., whether Or47b ORN axons appear in the dorsal antennal
lobe region, in contrast to the typical V-shaped glomerulus in wild-type controls. The
phenotype shown in Figure 1 (glomerular expansion) is largely consistent from brain to
brain, while the phenotypes shown in Figure 2B,E exhibit variability, which were
categorized into expansion or dorsal shift. The phenotype was quantified by the
percentage of antennal lobes exhibiting each defect among all the brains examined in
respective groups. P-value was calculated by two-tailed Fisher's exact test through the
built-in functions of GraphPad Prism 9 software.

Inverse PCR to recover the genomic DNA sequence flanking the *Or47b-GAL4* transgenic insertion

Or47b-GAL4 transgene was inserted into an unknown region on the second
chromosome by P-element-mediated method (VOSSHALL *et al.* 2000). The P-element
structure of *Or47b-GAL4* transgene was shown in the Figure 3B. We used previously
described inverse PCR method (HUANG *et al.* 2009) to identify the genomic sequence
flanking the insertion site. Briefly, genomic DNA was first extracted from 30 *Or47b-*
GAL4 flies (BDSC# 9983), followed by overnight digestion with any of three restriction
enzymes, MspI, HinP1I, or Sau3AI (New England BioLabs). Each digest was then
ligated by T4 DNA ligase (New England BioLabs) in larger volume (400 μL, see (HUANG
et al. 2009) for mix details) to promote intramolecular ligation while minimizing
intermolecular ligation. Ligation was performed at 16 °C overnight. The unknown
flanking sequence was then amplified from the ligated genomic DNA by inverse PCR
using forward and reverse primers targeting the 3'- or 5'-end sequences of the P-
element (Figure 3C). PCR reaction: 10 μL ligated DNA, 2 μL 5 μM forward and reverse
primer mix, 10 μL 5X *myTaq* reaction buffer, 0.5 μL *Taq* DNA polymerase. Thermal
cycling program: 3 min initial denaturation at 95 °C + 35 cycles (30 s denaturation at
95 °C, 1 min annealing at 55 °C, 2 min extension at 72 °C) + 10 min final extension at

72 °C. PCR products were then cleaned with Qiagen QIAquick PCR purification kit
followed by commercial DNA sequencing service.

A genomic region on the second chromosome (starting from 2L: 2753160) within the
first intron of *Bacc* gene was identified (Figure 3D,E) as the immediate sequence
flanking the inserted P-element 3' end. To verify this identified genomic insertion site, a
PCR assay was designed to amplify the genomic regions from *Or47b-GAL4* transgenic
flies and *w^{1118}* control flies respectively with different primer pairs (Figure 3E,F). PCR
mix recipe and thermal cycling programs were the same as abovementioned. Amplicons
were also purified and sequenced for validation.

Quantitative Reverse Transcription-PCR

To measure whether the insertion of *Or47b-GAL4* transgene into *Bacc* gene locus
affects its expression, we used the quantitative Reverse Transcription-PCR (qRT-PCR)
to extract and quantify the mRNA levels from whole heads of fruit flies. RNA extraction,
cDNA preparation, and qPCR protocols were described previously (ZHAO *et al.* 2020;
DEANHARDT *et al.* 2022).

Briefly, for each biological replicate, equal number of male and female heads (7 to 10,
5-7 days old post eclosion) were dissected in RNase-free environment with Trizol,
followed by tissue homogenization, cell lysis, and filtration with Qiagen QIAshredder
spin column. Three biological replicates were analyzed for each genotype. RNA was
then extracted and purified by Qiagen RNeasy Kit per manufacturer's instructions and
eluted in 60 µL RNase-free water. Genomic DNA was then removed using Invitrogen
TURBO DNA-free Kit per manufacturer's instructions. RNA concentration was
measured using NanoDrop after DNase treatment. Reverse transcription was performed
by Invitrogen SuperScript IV First-Strand cDNA synthesis Reaction Kit per
manufacturer's instructions. Notably, approximately equal amount of template RNA
across different samples were added based on RNA concentration.

Lastly, qPCR reactions were run on Roche LightCycler 96 Instrument with FastStart
Essential DNA Green Master (2X) in 20 µL volume with technical triplicates per
manufacturer's instructions. Thermal cycling program: 600 s pre-incubation at 95 °C +
40 three-step amplification cycles (10 s denaturation at 95 °C, 10 s annealing at 55 °C,
15 s extension at 72 °C with Single acquisition) + Melting Curve (10 s 95 °C, 60 s 65 °C,
1 s 97 °C with 5 Readings/°C). qPCR primers were designed to span two adjacent
exons when possible, with target amplicon length of 120 bp. Primer pairs were tested to
generate standard curves to evaluate amplification efficiency before being used for
expression comparison experiments. Expression levels of the gene of interest were
normalized to the house keeping gene *RpL13A* in each sample by $2^{-\Delta Ct}$ method. Two-
way ANOVA followed by multiple comparison test was performed in Prism 9 software to
determine statistical significance.

Table 1. Primers used in this study.

Experiment	Primer	Sequence (5' to 3')	Source
Inverse PCR	*For amplify P-element 3'-end flanking region*		(HUANG *et al.* 2009)
	Pry1	CCTTAGCATGTCCGTGGGGTTTGAAT	
	Pry4	CAATCATATCGCTGTCTCACTCA	
	For amplify P-element 5'-end flanking region		
	Plac1	CACCCAAGGCTCTGCTCCCACAAT	
	Plac4	ACTGTGCGTTAGGTCCTGTTCATTGTT	
Sequencing	*For sequencing P-element 3'-end product*		(HUANG *et al.* 2009)
	Spep1	GACACTCAGAATACTATTC	
	For sequencing P-element 5'-end product		
	Sp1	ACACAACCTTTCCTCTCAACAA	
PCR Validation assay	**FWD-1**	GAGCACATATCGGTGGTTAG	This study
	REV-1	GGCCCATACAATACACTCAA	
	FWD-2	CGAGAGGCAGTGCTTAAATA	
	REV-2	CTTGAGATCGTCCTTGACAG	
qPCR	**Bacc-F**	AGGCTCTGGAGGAAATCA	This study
	Bacc-R	CCGGAACCGTCATCATTATC	
	Act5C-F	GGCGCAGAGCAAGCGTGGTA	(ZHAO *et al.* 2020)
	Act5C-R	GGGTGCCACACGCAGCTCAT	
	RpL13A-F	GCGAGGAGCTGAACCTCTC	
	RpL13A-R	GGAAGTGGAATGGACCACGG	
	Tbp-F	TAAGCCCCAACTTCTCGATTCC	
	Tbp-R	GCCAAAGAGACCTGATCCCC	

Data availability

The authors affirm that all the data necessary for drawing the conclusions are present in the text, figures, and figure legends. Most of the *Drosophila* stocks are obtained from Bloomington or Vienna stock center, with identifiers listed in the materials and methods section. All the other lines are available upon request.

Acknowledgments

We would like to thank Bloomington Drosophila Stock Center and Vienna Drosophila Resource Center for providing all the fly stocks. We thank Duke Light Microscopy Core Facility for help with imaging. We thank Chengcheng Du for help with molecular biology.

Funding

This study is funded by grant NSF 2006471 and NIH 5R01NS109401 (both to P.C.V.). Q.D. is supported by Duke Biology Department Ph.D. program.

Conflicts of interest

The authors declare no conflict of interest with the contents of this paper.

Author contribution

Q.D. and P.C.V. conceived the study and designed the experiments; Q.D. did most of the experiments with help from R.E., A.C., and Y.C.; Q.D. analyzed the data and prepared the figures; Q.D. and P.C.V. wrote and edited the manuscript.

References

Ang, L. H., J. Kim, V. Stepensky and H. Hing, 2003 Dock and Pak regulate olfactory axon pathfinding in Drosophila. Development 130**:** 1307-1316.

Barish, S., S. Nuss, I. Strunilin, S. Bao, S. Mukherjee *et al.*, 2018 Combinations of DIPs and Dprs control organization of olfactory receptor neuron terminals in Drosophila. PLoS Genet 14**:** e1007560.

Barish, S., and P. C. Volkan, 2015 Mechanisms of olfactory receptor neuron specification in Drosophila. Wiley Interdiscip Rev Dev Biol 4**:** 609-621.

Bellen, H. J., R. W. Levis, G. Liao, Y. He, J. W. Carlson *et al.*, 2004 The BDGP Gene Disruption Project: Single Transposon Insertions Associated With 40% of Drosophila Genes. Genetics 167**:** 761-781.

Berg, B., T. Almaas, J. Bjaalie and H. Mustaparta, 1998 The macroglomerular complex of the antennal lobe in the tobacco budworm moth Heliothis virescens: specified subdivision in four compartments according to information about biologically significant compounds. Journal of comparative Physiology A 183**:** 669-682.

Brand, A. H., and N. Perrimon, 1993 Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. Development 118**:** 401-415.

Chen, J., Y. Wang, Y. Zhang and P. Shen, 2013 Mutations in Bacchus reveal a tyramine-dependent nuclear regulator for acute ethanol sensitivity in Drosophila. Neuropharmacology 67**:** 25-31.

de Jong, S., J. A. Cavallo, C. D. Rios, H. A. Dworak and H. Sink, 2005 Target recognition and synaptogenesis by motor axons: responses to the sidestep protein. Int J Dev Neurosci 23: 397-410.

Deanhardt, B., Q. Duan, C. Du, C. Soeder, A. Morlote *et al.*, 2022 Social experience and pheromone receptor activity reprogram behavioral switch and neuromodulatory gene expression in sensory neurons. bioRxiv**:** 2021.2006. 2018.449021.

Dietzl, G., D. Chen, F. Schnorrer, K.-C. Su, Y. Barinova *et al.*, 2007 A genome-wide transgenic RNAi library for conditional gene inactivation in Drosophila. Nature 448**:** 151-156.

Elhanany-Tamir, H., Y. X. V. Yu, M. Shnayder, A. Jain, M. Welte *et al.*, 2012 Organelle positioning in muscles requires cooperation between two KASH proteins and microtubules. Journal of Cell Biology 198**:** 833-846.

Fambrough, D., and C. S. Goodman, 1996 The Drosophila beaten path gene encodes a novel secreted protein that regulates defasciculation at motor axon choice points. Cell 87**:** 1049-1058.

Fishilevich, E., and L. B. Vosshall, 2005 Genetic and functional subdivision of the Drosophila antennal lobe. Curr Biol 15**:** 1548-1553.

Gaudet, P., M. S. Livstone, S. E. Lewis and P. D. Thomas, 2011 Phylogenetic-based propagation of functional annotations within the Gene Ontology consortium. Brief Bioinform 12**:** 449-462.

Green, E. W., G. Fedele, F. Giorgini and C. P. Kyriacou, 2014 A Drosophila RNAi collection is subject to dominant phenotypic effects. Nat Methods 11**:** 222-223.

Groen, C. M., J. L. Podratz, J. Pathoulas, N. Staff and A. J. Windebank, 2022 Genetic Reduction of Mitochondria Complex I Subunits is Protective against Cisplatin-Induced Neurotoxicity in Drosophila. Journal of Neuroscience 42**:** 922-937.

Groth, A. C., M. Fish, R. Nusse and M. P. Calos, 2004 Construction of transgenic Drosophila by using the site-specific integrase from phage φC31. Genetics 166**:** 1775-1782.

Hing, H., N. Reger, J. Snyder and L. G. Fradkin, 2020 Interplay between axonal Wnt5-Vang and dendritic Wnt5-Drl/Ryk signaling controls glomerular patterning in the Drosophila antennal lobe. Plos Genetics 16.

Hong, W., and L. Luo, 2014 Genetic control of wiring specificity in the fly olfactory system. Genetics 196**:** 17-29.

Hong, W., T. J. Mosca and L. Luo, 2012 Teneurins instruct synaptic partner matching in an olfactory map. Nature 484: 201-207.

Housden, B. E., M. Muhar, M. Gemberling, C. A. Gersbach, D. Y. Stainier *et al.*, 2017 Loss-of-function genetic tools for animal models: cross-species and cross-platform differences. Nature Reviews Genetics 18: 24-40.

Huang, A. M., E. J. Rehm and G. M. Rubin, 2009 Recovery of DNA sequences flanking P-element insertions in Drosophila: inverse PCR and plasmid rescue. Cold Spring Harb Protoc 2009: pdb prot5199.

Hueston, C. E., D. Olsen, Q. Li, S. Okuwa, B. Peng *et al.*, 2016 Chromatin Modulatory Proteins and Olfactory Receptor Signaling in the Refinement and Maintenance of Fruitless Expression in Olfactory Receptor Neurons. PLoS Biol 14: e1002443.

Kinold, J. C., M. Brenner and H. Aberle, 2021 Misregulation of Drosophila Sidestep Leads to Uncontrolled Wiring of the Adult Neuromuscular System and Severe Locomotion Defects. Front Neural Circuits 15: 658791.

Larkin, A., S. J. Marygold, G. Antonazzo, H. Attrill, G. dos Santos *et al.*, 2020 FlyBase: updates to the Drosophila melanogaster knowledge base. Nucleic Acids Research 49: D899-D907.

Li, H., J. Janssens, M. De Waegeneer, S. S. Kolluru, K. Davie *et al.*, 2022 Fly Cell Atlas: A single-nucleus transcriptomic atlas of the adult fruit fly. Science 375: eabk2432.

Li, H., A. Watson, A. Olechwier, M. Anaya, S. K. Sorooshyari *et al.*, 2017 Deconstruction of the beaten Path-Sidestep interaction network provides insights into neuromuscular system development. Elife 6.

Li, Q., S. Barish, S. Okuwa, A. Maciejewski, A. T. Brandt *et al.*, 2016 A Functionally Conserved Gene Regulatory Network Module Governing Olfactory Neuron Diversity. PLoS Genet 12: e1005780.

Li, Q., S. Barish, S. Okuwa and P. C. Volkan, 2015 Examination of Endogenous Rotund Expression and Function in Developing Drosophila Olfactory System Using CRISPR-Cas9-Mediated Protein Tagging. G3 (Bethesda) 5: 2809-2816.

Li, Q., T. S. Ha, S. Okuwa, Y. Wang, Q. Wang *et al.*, 2013 Combinatorial rules of precursor specification underlying olfactory neuron diversity. Curr Biol 23: 2481-2490.

Markstein, M., C. Pitsouli, C. Villalta, S. E. Celniker and N. Perrimon, 2008 Exploiting position effects and the gypsy retrovirus insulator to engineer precisely expressed transgenes. Nature genetics 40: 476-483.

Morel, V., S. Lepicard, A. N Rey, M.-L. Parmentier and L. Schaeffer, 2014 Drosophila Nesprin-1 controls glutamate receptor density at neuromuscular junctions. Cellular and molecular life sciences 71: 3363-3379.

Ni, J.-Q., L.-P. Liu, R. Binari, R. Hardy, H.-S. Shim *et al.*, 2009 A Drosophila resource of transgenic RNAi lines for neurogenetics. Genetics 182: 1089-1100.

Ni, J.-Q., M. Markstein, R. Binari, B. Pfeiffer, L.-P. Liu *et al.*, 2008 Vector and parameters for targeted transgenic RNA interference in Drosophila melanogaster. Nature methods 5: 49-51.

Ni, J.-Q., R. Zhou, B. Czech, L.-P. Liu, L. Holderbaum *et al.*, 2011 A genome-scale shRNA resource for transgenic RNAi in Drosophila. Nature methods 8: 405-407.

Özkan, E., R. A. Carrillo, C. L. Eastman, R. Weiszmann, D. Waghray *et al.*, 2013 An extracellular interactome of immunoglobulin and LRR proteins reveals receptor-ligand networks. Cell 154: 228-239.

Perkins, L. A., L. Holderbaum, R. Tao, Y. Hu, R. Sopko *et al.*, 2015 The transgenic RNAi project at Harvard Medical School: resources and validation. Genetics 201: 843-852.

Pipes, G. C., Q. Lin, S. E. Riley and C. S. Goodman, 2001 The Beat generation: a multigene family encoding IgSF proteins related to the Beat axon guidance molecule in Drosophila. Development 128: 4545-4552.

664   Ramin, M., Y. Li, W. T. Chang, H. Shaw and Y. Rao, 2019 The peacefulness gene promotes aggression in
665   Drosophila. Mol Brain 12**:** 1.
666   Scott, K., R. Brady, Jr., A. Cravchik, P. Morozov, A. Rzhetsky *et al.*, 2001 A chemosensory gene family
667   encoding candidate gustatory and olfactory receptors in Drosophila. Cell 104**:** 661-673.
668   Siebert, M., D. Banovic, B. Goellner and H. Aberle, 2009 Drosophila motor axons recognize and follow a
669   Sidestep-labeled substrate pathway to reach their target fields. Genes Dev 23**:** 1052-1062.
670   Sink, H., E. J. Rehm, L. Richstone, Y. M. Bulls and C. S. Goodman, 2001 sidestep encodes a target-derived
671   attractant essential for motor axon guidance in Drosophila. Cell 105**:** 57-67.
672   Stockinger, P., D. Kvitsiani, S. Rotkopf, L. Tirian and B. J. Dickson, 2005 Neural circuitry that governs
673   Drosophila male courtship behavior. Cell 121**:** 795-807.
674   Titlow, J., F. Robertson, A. Järvelin, D. Ish-Horowicz, C. Smith *et al.*, 2020 Syncrip/hnRNP Q is required for
675   activity-induced Msp300/Nesprin-1 expression and new synapse formation. Journal of Cell
676   Biology 219.
677   van der Graaf, K., S. Srivastav, P. Singh, J. A. McNew and M. Stern, 2022 The Drosophila attP40 docking
678   site and derivatives are insertion mutations of MSP300. bioRxiv.
679   Venken, K. J., K. L. Schulze, N. A. Haelterman, H. Pan, Y. He *et al.*, 2011 MiMIC: a highly versatile
680   transposon insertion resource for engineering Drosophila melanogaster genes. Nat Methods 8**:**
681   737-743.
682   Vissers, J. H., S. A. Manning, A. Kulkarni and K. F. Harvey, 2016 A Drosophila RNAi library modulates
683   Hippo pathway-dependent tissue growth. Nat Commun 7**:** 10368.
684   Vosshall, L. B., A. M. Wong and R. Axel, 2000 An olfactory sensory map in the fly brain. Cell 102**:** 147-159.
685   Wu, B., J. Li, Y. H. Chou, D. Luginbuhl and L. Luo, 2017 Fibroblast growth factor signaling instructs
686   ensheathing glia wrapping of Drosophila olfactory glomeruli. Proc Natl Acad Sci U S A 114**:** 7505-
687   7512.
688   Xie, Q., B. Wu, J. Li, C. Xu, H. Li *et al.*, 2019 Transsynaptic Fish-lips signaling prevents misconnections
689   between nonsynaptic partner olfactory neurons. Proc Natl Acad Sci U S A 116**:** 16068-16073.
690   Xie, X., and J. A. Fischer, 2008 On the roles of the Drosophila KASH domain proteins Msp-300 and
691   Klarsicht. Fly 2**:** 74-81.
692   Yao, Y., Y. Wu, C. Yin, R. Ozawa, T. Aigaki *et al.*, 2007 Antagonistic roles of Wnt5 and the Drl receptor in
693   patterning the Drosophila antennal lobe. Nat Neurosci 10**:** 1423-1432.
694   Zhao, S., B. Deanhardt, G. T. Barlow, P. G. Schleske, A. M. Rossi *et al.*, 2020 Chromatin-based
695   reprograming of a courtship regulator by concurrent pheromone perception and hormone
696   signaling. Sci Adv 6**:** eaba6913.
697   Zheng, Y., R. A. Buchwalter, C. Zheng, E. M. Wight, J. V. Chen *et al.*, 2020 A perinuclear microtubule-
698   organizing centre controls nuclear positioning and basement membrane secretion. Nature cell
699   biology 22**:** 297-309.
700   Zirin, J., Y. Hu, L. Liu, D. Yang-Zhou, R. Colbeth *et al.*, 2020 Large-Scale Transgenic Drosophila Resource
701   Collections for Loss- and Gain-of-Function Studies. Genetics 214**:** 755-767.