Cardiovirus leader proteins retarget RSK kinases toward alternative substrates to perturb nucleocytoplasmic traffic

Belén Lizcano-Perret[1], Cécile Lardinois[1], Fanny Wavreil[1], Philippe Hauchamps[2], Gaëtan Herinckx[3], Frédéric Sorgeloos[1], Didier Vertommen[3], Laurent Gatto[2] and Thomas Michiels[1*]

[1] VIRO unit, de Duve Institute, Université Catholique de Louvain, Brussels, 1200, Belgium

[2] CBIO unit, de Duve Institute, Université Catholique de Louvain, Brussels, 1200, Belgium

[3] PHOS unit and MASSPROT platform, de Duve Institute, Université Catholique de Louvain, Brussels, 1200, Belgium

[*]Correspondence: thomas.michiels@uclouvain.be

Summary

Proteins from unrelated pathogens, including some RNA or DNA viruses and bacteria can recruit and activate cellular p90-ribosomal protein S6 kinases (RSKs) through a common linear motif. Data suggested a model where pathogens' proteins act to dock the recruited RSKs toward specific substrates, which then act as effectors to the benefit of the pathogens. Using cardiovirus leader protein (L) as a paradigm, we show that pathogens' proteins can modify the spectrum of RSK substrates in infected cells. L triggers nucleocytoplasmic trafficking perturbation and phenylalanine-glycine (FG)-nucleoporin hyperphosphorylation in an RSK-dependent fashion. Biotin ligase experiments identified FG-nucleoporins as common partners of L and RSK in infected cells. Using cells expressing an analog-sensitive RSK2 mutant, we show that L triggers direct phosphorylation of NUP98 and NUP214 by RSK2 in infected cells. Our data therefore demonstrate a novel virulence mechanism where pathogens' proteins hijack and retarget cellular protein kinases of the RSK family.

Keywords: Linear motifs, p90 ribosomal S6 kinases, Picornavirus, leader protein, nucleoporins, nuclear pore complex, phosphorylation, virulence, innate immunity escape, nucleocytoplasmic traffic

Introduction

Proteins encoded by several unrelated pathogens, including RNA viruses, DNA viruses and bacteria, were recently shown to use a common short linear motif (D/E-D/E-V-F, referred to as DDVF hereafter)

to recruit members of the cellular p90-ribosomal S6 protein kinases (RSKs) family: RSK1, RSK2, RSK3 and RSK4 (Sorgeloos et al., 2022, Alexa et al., 2022). Interestingly, competition and cross-linking experiments, as well as crystallography data show that these pathogens' proteins, the leader (L) protein (cardioviruses), ORF45 (Kaposi sarcoma-associated herpes virus - KSHV) and YopM (*Yersinia*) use a common interface to recruit RSKs. Binding of the pathogens' proteins prevents RSK dephosphorylation by cellular phosphatases, thereby maintaining RSK in an active state (Sorgeloos et al., 2022, Alexa et al., 2022). Although infection with all three pathogens leads to RSK activation, the outcome of this activation differs according to the protein bound to RSK. YopM association with RSK was proposed to inhibit the inflammasome and to lead to IL-10 production (Chung et al., 2016, Ratner et al., 2016); RSK recruitment by ORF45 enhances lytic replication of KSHV (Kuang et al., 2011, Li et al., 2015), whereas RSK recruitment by cardiovirus L protein leads to the inhibition of the antiviral eukaryotic initiation factor 2 alpha kinase 2 (EIF2AK2), better known as PKR (Sorgeloos et al., 2022). PKR inhibition by L was shown to depend on L interaction with RSK but also on L protein's C-terminal domain. Indeed, the M60V mutation in the C-terminal domain of L does not affect RSK recruitment but abolishes PKR inhibition. These observations led us to propose the "model of the clamp" whereby pathogens' proteins would act as adaptor proteins that force a given enzyme (here RSKs) to act on a specific substrate. Through their DDVF motif, pathogens' proteins recruit and maintain RSKs in an activated state. Through another domain (i.e. the C-terminal domain for L), they recruit proteins that could serve as substrate for RSKs. After phosphorylation, such proteins would act as effectors to the benefit of the pathogen (Figure 1A) (Sorgeloos et al., 2022). In an alternative model, proteins interacting with RSK through the conserved DDVF motif could directly act as preferential substrates for RSK-mediated phosphorylation (Figure 1B), as was suggested for protein RHBDF1 (Rhomboid 5 homolog 1) (Alexa et al., 2022).

Cardioviruses belong to the *Picornaviridae* family and include encephalomyocarditis virus (EMCV), Theiler's murine encephalomyelitis virus (TMEV) and the human Saffold virus (SAFV), closely related to TMEV. Despite its very small size (67-76 amino acids) and lack of enzymatic activity, the L protein encoded by these viruses was shown to be multifunctional as, beside inhibiting PKR, it blocks interferon gene transcription, and triggers an extensive diffusion of nuclear and cytoplasmic proteins across the nuclear membrane (Freundt et al., 2018, Brahic et al., 2005, van Pesch et al., 2001, Zoll et al., 2002, Hato et al., 2007, Borghese and Michiels, 2011, Borghese et al., 2019, Delhaye et al., 2004, Lidsky et al., 2006, Lizcano-Perret and Michiels, 2021, Sorgeloos et al., 2022). This nucleocytoplasmic trafficking perturbation was associated with the hyperphosphorylation of phenylalanine-glycine-nucleoporins (FG-NUPs) such as NUP98, NUP153 or NUP62 (Porter and Palmenberg, 2009, Porter et al., 2010, Ricour et al., 2009b, Bardina et al., 2009). FG-NUPs are proteins of the nuclear pore complex (NPC) that possess intrinsically disordered domains, rich in phenylalanine and glycine residues. These domains form a mesh in the center of the nuclear pore, which enables interaction with karyopherins, thus allowing selective transport of proteins and RNA through the NPC (Aramburu and Lemke, 2017,

Beck and Hurt, 2017, Hampoelz et al., 2019, Hülsmann et al., 2012). Molecular dynamic simulations

predict that FG-NUP phosphorylation drastically decreases the density of FG-NUPs inside the pore

(Mishra et al., 2019). Accordingly, electron microscopy analysis of EMCV-infected cells displayed

such a density loss in the center of the NPC (Bardina et al., 2009). Through perturbation of the NPC,

cardioviruses trigger a diffusion of nuclear proteins to the cytoplasm which can be used by the virus for

viral replication and translation (Lizcano-Perret and Michiels, 2021).

Preliminary data suggested that nucleocytoplasmic trafficking perturbation by cardiovirus L proteins

depended on RSK and on the recruitment of a cellular target by the L-RSK complex (model of the

clamp). The aim of this work was to assess whether RSK could indeed be retargeted by L to a new

substrate in infected cells and to identify such a substrate that could trigger nucleocytoplasmic

trafficking perturbation. Our data show that L can recruit both RSK and FG-NUPs and that FG-NUPs

can act as direct RSK substrates in infected cells. This work provides strong support to the model of the

clamp and elucidates a novel virulence mechanism.

Results

Cardiovirus L-mediated nucleocytoplasmic trafficking perturbation depends on RSK

recruitment by L

We tested whether L-mediated redistribution of nuclear and cytoplasmic proteins across the nuclear

envelope and the hyperphosphorylation of FG-NUPs depends on both RSK and L. In agreement with

previous reports (Delhaye et al., 2004), cytosolic diffusion of the nuclear protein PTB occurred in HeLa

cells infected for 10 hours with TMEV expressing a wild type L protein (L^{WT}) but not in non-infected

cells or in cells infected with viruses carrying the M60V mutation in the C-terminal part of L

(L^{M60V})(Figures 2A and S1A). In the latter cells, PTB however partly appeared in cytosolic punctae,

likely corresponding to stress granules that were shown to form in the absence of PKR inhibition by L

(Borghese and Michiels, 2011). Importantly, L^{WT}-mediated PTB diffusion was almost abrogated in

RSK-deficient HeLa cells (HeLa-RSK-TKO). Viral replication was also decreased in the absence of

RSK, in agreement with increased PKR activation in these cells. Interestingly, PTB diffusion was

restored after transduction of HeLa-RSK-TKO cells with lentiviruses expressing any of the four human

RSK isoforms (Figures 2A and S1B). Accordingly, NUP98 hyperphosphorylation was dramatically

decreased in HeLa-RSK-TKO cells and restored in cells transduced to express any of the four RSK

isoforms (Figure 2B). Similar observations were made in the case of EMCV as soon as 5 hours post-

infection, although dependence on RSK was less pronounced (Figures 2C-D and S1C-D). Thus, PTB

diffusion out of the nucleus and NUP98 hyperphosphorylation depend on both L and RSK.

In addition, TMEV and EMCV expressing L^{WT} but not mutant L proteins triggered an RSK-dependent

disruption of nucleocytoplasmic trafficking of other proteins harboring canonical nuclear export (NES)

and import (NLS) sequences, as shown by diffusion of GFP-NES and RFP-NLS proteins in live cells that stably express these proteins (Figures 2E-G and S1E). This observation demonstrates that nucleocytoplasmic redistribution of proteins was not specific to PTB. Again, RSK dependence of protein redistribution was slightly less prominent for EMCV than for TMEV.

Ectopic expression of TMEV L^{WT} but not the L^{F48A} mutant (affected in the RSK-binding motif) was sufficient to trigger nucleocytoplasmic redistribution of proteins (Figures 2H-J). Also, expression of L^{M60V} or YopM did not affect nucleocytoplasmic trafficking although these proteins can activate RSK (Sorgeloos et al., 2022). Taken together, these data show that L-mediated nucleocytoplasmic trafficking alteration does not dependent on virus replication per se or on the expression of other viral proteins and requires both interaction with RSK and, at least for TMEV L, integrity of the C-terminal part of the protein.

Identifying RSK partners in TMEV-infected cells using BioID2

According to the model of the clamp, TMEV L would recruit RSK through its central DDVF motif and would recruit potential RSK substrates through its C-terminal domain. As suggested above, the M60V mutation of L would prevent the recruitment of such a target, without affecting RSK binding.

In order to identify target proteins that are recruited by L as potential RSK targets, we used the modified bacterial biotin ligase BioID2 (BioID) which, when fused to a protein of interest, allows promiscuous biotinylation of proximal proteins in the cell (Kim et al., 2016).

A BioID-RSK2 (BioID-RSK) fusion was constructed and expressed in RSK1/2-deficient HeLa cells (HeLa-RSK-DKO) by lentiviral transduction to identify RSK's potential partners during TMEV infection (Figure 3A). After testing that the BioID-RSK construct was activated (i.e. phosphorylated) in the presence of L in infected cells and able to rescue L activities such as L-mediated PKR inhibition (Figure S2A), we performed biotinylation experiments in order to identify RSK partners in the context of infected cells. HeLa BioID-RSK cells were infected with either L^{WT} or L^{M60V} viruses for 16 hours in presence of biotin. Biotinylated proteins were pulled down using streptavidin beads and processed for mass spectrometry analysis. Western blot detection of biotinylated proteins (Figure 3B) confirmed the concentration of biotinylated proteins in the pulled down fraction and revealed the presence of a 100kDa band specific to the L^{WT}-infected sample, suggesting that the L protein status indeed influenced target biotinylation by the BioID-RSK fusion.

Identifying L protein partners using BioID

Several attempts to identify proteins that are recruited by TMEV L were made in our lab by coimmunoprecipitation. However, beside RSK, no clear L partner was consistently identified. It is not unlikely that proteins recruited by L would interact with L only transiently and with low affinity. We thus used the BioID system to identify such proteins that may transiently interact with L in infected cells and then serve as RSK substrates. However, due to packaging limitations in picornaviruses, the

BioID coding sequence (696 nt) would be too long to be accommodated into the viral genome. We thus

generated *trans*-encapsidated replicons where a sequence coding for BioID-L and eGFP was substituted

for the capsid-coding region (Figure S2B). These replicons were encapsidated in 293T cells stably

expressing a capsid coding sequence carrying synonymous mutations in the VP2 region that affect *CRE*

function (Lobert et al., 1999) to prevents the selection of replication-competent wild type viruses that

might emerge through recombination. *Trans*-encapsidated replicons coding for BioID fused to L^{WT} ,

L^{M60V} and L^{F48A} were produced (Figure 3A). In HeLa cells infected with these replicons, L activities

(i.e. RSK phosphorylation, PKR activation and NUP98 hyperphosphorylation) were as expected (Figure

S2C). HeLa cells were then infected with BioID-L replicons for 14 hours in presence of biotin.

Biotinylated proteins were pulled down in stringent conditions. Western blot confirmed protein

biotinylation in infected cell extracts and showed some bands occurring in L^{WT} and L^{F48A} but not in

L^{M60V} samples (Figure 3C). These results confirm specific biotinylation of proteins by BioID-L.

FG-NUPs are enriched in BioID-L and BioID-RSK proxeomes

We expected that RSK targets recruited by L would be present in both BioID-RSK and BioID-L screens.

Pulled down biotinylated proteins from both screens (n=3 for each screen) were identified by mass

spectrometry (MS) and sorted according to their peptide spectrum match (PSM) number (for more detail

on the calculations see material and methods). For BioID-RSK, proteins were sorted according to their

abundance in L^{WT}-infected cells relative to that in non-infected cells and in cells infected with the L^{M60V}

mutant virus (Figure 3D, vertical axis). For BioID-L (replicons), proteins were sorted according to their

abundance in L^{WT} and L^{F48A} samples relative to that in L^{M60V}-infected and in non-infected samples

(Figure 3D, horizontal axis). Hence, the proteins that would be recruited by the C-terminal domain of

L are those that have high ratios in both screens. Interestingly, many of these proteins were FG-NUPs

(Figures 3D and E). ProDA (Probabilistic Dropout Analysis) statistical analysis (Ahlmann-Eltze and

Anders, 2020) was used to identify proteins whose abundance differs in pairwise comparisons: L^{WT}

versus L^{M60V} for the BioID-RSK experiments, and L^{WT} versus L^{M60V} or L^{F48A} versus L^{M60V} for the BioID-

L experiment. The table in Figure 3E shows adjusted P-values obtained from this analysis for the 20

best-ranked proteins. In the BioID-RSK experiments, FG-NUPs had the lowest adjusted P-values

(though not significant) and highest fold change (Figures 3E and F). In the BioID-L experiments many

FG-NUPs reached significant scores when pairwise comparisons were made between L^{WT} and L^{M60V} or

between L^{M60V} and L^{F48A} (Figures 3E and F). Most importantly, FG-NUPs that had previously been

shown to be hyperphosphorylated during cardiovirus infection: NUP62, NUP98, NUP153 and NUP214

(Porter and Palmenberg, 2009, Porter et al., 2010, Ricour et al., 2009b, Bardina et al., 2009) exhibited

statistical significance (Figure 3E). From these results we conclude that L recruits RSK through its

DDVF motif and then docks RSK toward FG-NUPs when the C-terminal domain of L is intact.

Proteins biotinylated by BioID-RSK and BioID-L are localized at the NPC

We analyzed the subcellular localization of proteins that were biotinylated by BioID-RSK and by

BioID-L in infected cells. To this end, biotinylation experiments were performed as above but infected

cells were fixed after 10h of infection, permeabilized and stained them with Alexa Fluor conjugated

streptavidin. For BioID-RSK, mock-infected cells show staining of biotinylated proteins in the nucleus,

in agreement with the mostly nuclear localization of RSKs (Chen et al., 1992) (Figures 4A-B and S3A).

In cells infected with the L^{WT} virus, biotinylated proteins form a faint rim around the nucleus, in addition

to the diffuse nuclear staining, showing that part of the (BioID-)RSK molecules are recruited to the

nuclear envelope. In cells infected with the L^{M60V} mutant virus, this nuclear rim is lost, in agreement

with a lack of L^{M60V} interaction with nucleoporins. For BioID-L replicons, staining occurs in the nucleus

and at the nuclear envelope for L^{WT}, in agreement with an interaction of L with both RSK and NUPs

(Figures 4C, D and S3B). Proteins biotinylated by BioID-L^{M60V} show diffuse staining in the nucleus, in

agreement with L^{M60V} interaction with RSK but not with nucleoporins. Proteins biotinylated by BioID-

L^{F48A} show extensive colocalization with nucleoporin POM121 at the nuclear envelope (Figure S3B),

in agreement with the lack of interaction with RSK, which would likely sequester part of the L protein

in the nucleoplasm. In conclusion, labeling of biotinylated proteins support our model where L interacts

with RSK in the nucleus and recruits a portion of RSK molecules to the NPC, via an interaction between

the C-terminal domain of L and NUPs.

FG-NUPs are direct substrates for RSK in cardiovirus infected cells

Since FG-NUPs are known to be hyperphosphorylated during cardiovirus infection and since our results

showed that RSK is in their close proximity, we aimed to test if RSK can directly phosphorylate these

FG-NUPs in infected cells. For this, we used the « analog-sensitive kinase system » developed by the

group of Shokat (Allen et al., 2007). This system is based on the use of an ATP analog: N6-alkylated

ATP-γ-S (A*TP-S), bulkier than ATP, that only fits in ATP binding pockets of kinases that have been

mutated to accommodate such bulkier analogs. The advantage of this technique is that the mutated

kinase (RSK in our case) is the only kinase in the cell that can use A*TP-S and will therefore

thiophosphorylate its substrates. After alkylation, such target proteins can be recognized using a specific

anti-thiophosphate ester antibody (Figure 5A). We identified Leu147 as the RSK2 "Gatekeeper" residue

(i.e. residue in the ATP-binding pocket that, when mutated to a smaller residue, allows the access of the

A*TP-S) by homology to Thr338 in the consensus sequence defined for c-SRC (Lopez et al., 2014)

(Figure S4A). Leu147 was mutated either to Gly (As1-RSK) or to Ala (As2-RSK). To test for ATP

analog usage by the modified RSK kinases, WT, As1- and As2-RSKs were stably expressed in HeLa-

RSK-TKO cells by lentiviral transduction, yielding WT-, As1- and As2-RSK cells. Expression of both

As1- and As2-RSK allowed L functions (i.e. PKR inhibition, RSK activation and NUP98

hyperphosphorylation)(Figure S4B). The ATP analog N6-Bn-ATP-γ-S was better incorporated than

N6-PhEt-ATP-γ-S by the As-RSKs (Figure S4C). The As2-RSK kinase was better expressed than As1-

RSK and readily accommodated this analog (Figures S4B and C). Therefore, As2-RSK cells and N6-Bn-ATP-γ-S were chosen for further experiments. HeLa WT-RSK or As2-RSK cells were then infected with L^{WT} and L^{M60V} viruses for 8h30min. Cells were then permeabilized with digitonin and treated with A*TP-S (N6-Bn-ATP-γ-S) for 1 hour. Then, NUP98 was immunoprecipitated and its thiophosphorylation status was analyzed, after alkylation, by western blot with the anti-thiophosphate ester antibody. The blot shown in Figure 5B shows a band highly enriched in the immunoprecipitated sample, migrating at the expected molecular mass for NUP98 (98kDa). This band was most prominent in the samples from as2-RSK cells, confirming the specificity of thiophosphate ester detection. Importantly, this band was detected much more in samples of cells infected with the L^{WT} virus than in samples of cells that were not infected or infected with the L^{M60V} virus. These data indicate that NUP98 can be directly phosphorylated by RSK and, as expected, that this phosphorylation only occurs after infection with the L^{WT} virus. For EMCV, as this virus replicates much faster than TMEV, NUP98 was immunoprecipitated after 3h30 of infection. Thiophosphorylation of immunoprecipitated proteins was tested by western blot using the anti-thiophosphate ester antibody (Figure 5C). As for TMEV, a ~100kDa protein likely corresponding to NUP98 was thiophosphorylated by As2-RSK only in L^{WT} infected conditions. It is noteworthy that additional bands were detected, which likely correspond to other FG-NUPs such as NUP62, NUP153 and NUP214 that coimmunoprecipitated with NUP98.

In order to confirm FG-NUPs thiophosphorylation by As2-RSK in infected cells, experiments were performed the other way around. Therefore, thiophosphorylated proteins were immunoprecipitated from cells infected with TMEV variants, using the thiophosphate ester antibody. Then the IP fraction (i.e. thiophosphorylated proteins) was analyzed by western blot for the presence of NUP98 (Figure 5D) and NUP214 (Figure S4D). Again, these NUPs were most prominently detected in the conditions where the L protein was WT and RSK was mutated (As2-RSK). These data confirm that RSK directly phosphorylates FG-NUPs when the L^{WT} protein is present.

Taken together, these results confirm that part of the cellular RSKs can be redirected to the nuclear pore complex by the L protein, where RSKs phosphorylate FG-NUPs. By interacting with RSK through its DDVF motif and with FG-NUPs through its C-terminal domain, L serves as a viral adapter protein to modulate RSK's activity and redirects these kinases toward new substrates. thus supporting the model of the clamp. Hyperphosphorylation of FG-NUPs by RSK likely triggers a global perturbation of nucleocytoplasmic trafficking and may therefore facilitate cytoplasmic replication of cardioviruses.

Discussion

Recent work showed that some proteins expressed by unrelated viruses and bacteria hijack host RSK kinases through a conserved DDVF linear motif that likely emerged in those proteins by convergent

evolution (Sorgeloos et al., 2022, Alexa et al., 2022). Occurrence of the DDVF motif in RHBDF1, a

known target of RSKs suggests that some proteins, either from pathogens or from the host cell, can

associate with RSK through the DDVF motif to promote their own phosphorylation by RSKs (Alexa et

al., 2022). In the case of TMEV L protein, a point mutation in the C-terminal domain of the protein

(L^{M60V}) abrogated L activities although this mutation affected neither L-RSK interaction nor RSK

activation by L (Ricour et al., 2009a, Sorgeloos et al., 2022). The phenotype of this mutant supported

the model of the clamp where the pathogens' protein would act to bridge RSK to specific substrates.

The two models are not mutually exclusive.

Our data show that L proteins from TMEV and EMCV indeed act by redirecting part of the cellular

RSK kinases toward the nuclear pore complex where RSKs directly phosphorylate FG-NUPs.

Bacteria of the genus *Yersinia* use a type III secretion system to inject Yop proteins into contacted

eukaryotic cells (Cornelis and Wolf-Watz, 1997). YopM was shown to interact with RSK through a C-

terminal DDVF motif (Sorgeloos et al., 2022) and with protein kinase PKNs through central leucin-rich

motifs. Formation of the tripartite (YopM-RSK-PKN) complex results in the phosphorylation and

activation of PKN, which requires active RSK (McDonald et al., 2003, Hentschke et al., 2010),

suggesting that RSK might also be redirected toward PKN as a specific substrate. This scenario is very

likely although direct phosphorylation of PKN by RSK was not formally proven.

Another question regarding the clamp model is whether the substrates to which RSKs are redirected by

the pathogens' proteins are physiological RSK substrates, the phosphorylation of which is increased by

their forced association with RSKs or whether they represent unconventional RSK substrates. From the

blots of analog-sensitive RSK experiments (Figures 5B-D and S4D), it appears that basal

thiophosphorylation of NUP98 or NUP214 might occur, even in non-infected cells that do not contain

the L protein. Also, given that the main binding site of pathogens' proteins is located in a surface-

exposed loop of RSK, relatively remote from the catalytic site, it is unlikely that this interaction can

modify the catalytic site enough to accommodate structurally divergent substrates. Thus, the role of the

bridging proteins is likely to increase the frequency of encounters between RSKs and specific

substrates, and possibly also to drive these kinases to specific subcellular locations.

Viruses of the *Picornaviridae* family disturb nucleocytoplasmic traffic, likely to recruit host nuclear

RNA-binding proteins to the cytoplasm of the infected cells, where their genome translation and

replication takes place. Perturbation of the traffic was also reported to inhibit the activation of innate

immunity genes such as genes coding for interferon or chemokines, because these genes rely on the

translocation of transcription factors like IRF3/7 or NFκB (Lizcano-Perret and Michiels, 2021, De

Jesús-González et al., 2021). Enteroviruses encode proteases $2A^{pro}$ and $3C^{pro}$, which directly cleave FG-

NUPs (Saeed et al., 2020), thereby opening the central channel of the nuclear pore complex (Belov et

al., 2004, Bardina et al., 2009). Cardioviruses, however, induce the hyperphosphorylation of FG-NUPs,

thus mimicking a process occurring during mitosis, which ends up in the dismantling of the NPCs and

the free diffusion of nuclear and cytoplasmic proteins. Here, we show that TMEV L promotes direct

FG-NUPs phosphorylation by RSKs. During mitosis, FG-NUP phosphorylation was proposed to involve mostly cyclin-dependent kinase 1 (CDK1), polo-like kinase 1 (PLK1) and NIMA-related kinases (Linder et al., 2017, Hampoelz et al., 2019). A phosphoproteomic analysis by Kosako et al. (Kosako et al., 2009) also suggested the involvement of the ERK pathway, ERK1/2 being the upstream kinase of RSK. We hypothesize that RSKs might indeed contribute, to some extent, to NUP phosphorylation during mitosis and that L triggers a stronger RSK-mediated FG-NUP phosphorylation during infection.

Our work illustrates a new virulence mechanism whereby pathogens' proteins not only activate but also redirect host kinases toward specific substrates and decipher how cardioviruses trigger RSK-mediated FG-NUP hyperphosphorylation to perturb nucleocytoplasmic trafficking in the host cell.

Acknowledgments

We are grateful to Melissa Drappier for suggesting the analog-sensitive kinase experiments. We are much indebted to Kevan Shokat for quick support and detailed protocols concerning this great technique.

We thank Frank J.M. van Kuppeveld for the gift of anti-EMCV capsid antibodies. We thank Stéphane Messe for excellent technical assistance, Nicolas Dauguet for expert help in cell sorting, and Eric Freundt and Frank J.M. van Kuppeveld for critical reading of this manuscript.

BLP and CL were the recipients of FRIA fellowship from the belgian FNRS. Work was supported by the EOS joint programme of Fonds de la recherche scientifique-FNRS and Fonds wetenschappelijk onderzoek-Vlaanderen-FWO (EOS ID: 30981113 and 40007527), Belgian fund for Scientific Research (PDR T.0185.14), Loterie Nationale through support to the de Duve Institute and Actions de Recherches Concertées (ARC).

Author Contributions https://www.cell.com/pb/assets/raw/shared/guidelines/CRediT-taxonomy.pdf

Conceptualization: B.L.P. and T.M.

Methodology: B.L.P, C.L., D.V., F.S. and T.M.

Software:

Validation:

Formal analysis: P.H., F.S., L.G.

Investigation: B.L.P, F.W., G.H., D.V. and T.M.

Resources:

Data curation:

Writing – original draft: B.L.P.

Writing – review & editing: B.L.P., D.V., F.S. and T.M.

Visualization:

Supervision: T.M.

Project administration: T.M.

Funding acquisition: T.M.

Declaration of Interests

The authors declare no conflicts of interests

Figure legends

Figure 1. Pathogen's proteins hijack RSK kinases.

(A) Indirect substrate recruitment - model of the clamp: a pathogen's protein acts as an adaptor protein,

which binds and activates RSK through its DDVF motif and recruits a target protein to be

phosphorylated by RSK through another domain. (B) Direct substrate recruitment. A pathogen's protein

interacts with RSK through its DDVF motif and is directly phosphorylated by RSK.

Figure 2. Nucleocytoplasmic traffic perturbation induced by cardioviruses depends on both the

L protein and RSK kinases.

(A-D) PTB diffusion out of the nucleus and hyperphosphorylation of NUP98 depend on both L and

RSK. Graphs showing the percentage of cells with PTB diffusion (mean ± SD), among cells infected

by TMEV (3D-positive) for 10h (A) or by EMCV (capsid-positive) for 5h (C). Counts: 25-50 infected

cells per experiment (n=4). **** denotes significant differences (One-way ANOVA, p < 0.0001)

between HeLa-RSK-TKO cells transduced with an empty vector (HeLa TKO) and cells re-expressing

indicated RSK isoforms, for L^{WT}-infected samples. (B and D) Western blots showing the RSK-

dependent hyperphosphorylation of NUP98 (shift upwards) or the inhibition of PKR activation induced

by TMEV (B) or EMCV (D). 3D or 2A were detected as control of infection, and ß-actin as loading

control. (E-G) Graphs and confocal microscopy pictures showing the quantification of RFP-NLS and

GFP-NES diffusion (mean ± SD) in live HeLa-LVX cells infected with TMEV for 10h (E-F) or EMCV

for 4h30. HeLa-LVX cells were either WT, or RSK-TKO cells transduced with an empty vector (HeLa-

TKO) or re-expressing RSK1. (G). Counts: 50 cells per experiment (n=4). ****: denotes significant

differences as above. Scale bar: 20μm

(H-J) Impact of L on nucleocytoplasmic traffic. HeLa-LVX cells transfected with an empty plasmid or

with plasmids expressing L^{WT}, L mutants or YopM. (H) Graph showing the percentage of cells with

GFP-NES and RFP-NLS diffusion (mean ± SD). Counts: 70±5 cells per experiment (n=6). One-way

ANOVA test was used to compare all samples to L^{WT}. (I) Confocal microscopy of HeLa-LVX cells

24hours post-transfection. Scale bar: 20μm. (J) Western blot of HeLa-LVX cells 24h post-transfection. FLAG staining shows the expression of L proteins and YopM.

Figure 3. BioID-RSK and BioID-L proxeomes identify FG-NUPs as targets recruited by the L-RSK complex.

(A) Cartoon showing the expected biotinylation by BioID-RSK or BioID-L fusion proteins of a target recruited by the C-terminal domain of L. (B) Western blot of proteins biotinylated by BioID-RSK in lysates and pulled down samples of TMEV infected cells. HeLa BioID-RSK cells were incubated for 2 days without biotin. Cells were then infected for 16h (MOI 2.5) with L^{WT} or L^{M60V} viruses in medium containing 5μM biotin. Biotinylated proteins were pulled-down using streptavidin-magnetic beads. (C) Western blot of proteins biotinylated by BioID-L in lysates and pulled down samples of TMEV infected cells. HeLa cells were infected for 14h (MOI 2.5) with BioID-L^{WT}, BioID-L^{M60V} or BioID-L^{F48A} replicons in medium containing 5μM biotin. Biotinylated proteins were then pulled-down using streptavidin-magnetic beads. (D) Proteins identified in the proximity of BioID-RSK (y axis) and BioID-L (x axis). Y-axis shows proteins detected after infection with L^{WT} but not L^{M60V} viruses (=PSMs in $L^{WT}/(L^{M60V}+NI+1)$). X-axis shows proteins detected after infection with BioID-L^{WT} or BioID-L^{F48A} but not BioID-L^{M60V} replicons (= PSMs in $(L^{WT}+ L^{M60V})/(L^{F48A}+NI+1)$). FG-NUPs are identified by green dots. (E) Table showing the adjusted P-values for the 20 best ranked proteins. Ranking was attributed by multiplying the BioID-RSK ratio by the BioID-L ratio. Statistical analysis of pairwise comparisons made between L^{WT} and L^{M60V} for BioID-RSK and between L^{WT} and L^{M60V} or L^{F48A} and L^{M60V} for BioID-L. Adjusted P-values < 0.05 are colored in red, FG-NUPs are colored in green. (F) Volcano plots showing the same pairwise comparisons as in E. Proteins having an adjusted P-value < 0.05 and a Log_2 fold change (LFC) > 1 are colored in red (= proteins in Q2). FG-NUPs are colored in green.

Figure 4. Subcellular localization of proteins biotinylated by BioID-RSK and BioID-L.

(A) Confocal microscopy images of biotinylated proteins (green) and 3D viral polymerase (red) detected in BioID-RSK expressing HeLa cells infected with L^{WT} or L^{M60V} viruses for 10h (MOI 5). (B) Quantification of cells showing a visible nuclear rim of biotinylated proteins in HeLa BioID-RSK cells infected with L^{WT} or L^{M60V} viruses (mean ± SD). Counts: 42 ± 10 infected cells per experiment (n=4). (C) Confocal microscopy images of biotinylated proteins stained with streptavidin-Alexa Fluor 594 (orange) and 3D viral polymerase (green) in HeLa cells infected for 10h with BioID-L^{WT}, BioID-L^{M60V} or BioID-L^{F48A} replicons (MOI 5). (D) Quantification of nuclear rim staining in HeLa cells infected with BioID-L^{WT}, BioID-L^{M60V} or BioID- L^{F48A} viruses (mean ± SD). Counts: 31 ± 11 infected cells per experiment (n=4). 3D viral polymerase was stained as a control of infection. (A and C) White arrows point to examples of cells exhibiting a nuclear rim staining. Scale bar: 10μm. (B and D) One-way ANOVA tests were used to compare all samples with each other.

Figure 5. RSK directly phosphorylates NUP98 and NUP214 in TMEV and EMCV infected cells.
(A) Cartoon showing the principle of the analog-sensitive kinase system. The kinase mutated in the ATP binding pocket (As-RSK) can use regular ATP as well as a bulkier ATP analog (A*TP-S). Use of the ATP analog induces the thiophosphorylation of the substrate. After an alkylation reaction with paranitrobenzylmesylate (PNBM), thiophosphates are converted to thiophosphate esters, which are specifically recognized by an anti-thiophosphate ester antibody. All other wild-type kinases in the cell, have normal ATP binding pockets that cannot accommodate the ATP analog. (B-D) HeLa cells expressing As2-RSK or WT-RSK were infected with TMEV for 8h (MOI 5) (B and D) or with EMCV for 3h30min (MOI 5) (C). Cells were then permeabilized with digitonin, and N6-Bn-ATP-γ-S was added for 1 hour. Cells were then lysed and either NUP98 (B and D) or thio-phosphate-ester containing proteins were immunoprecipitated (D). (B-C) Immunoblots showing a concentrated amount of NUP98 in the immunoprecipitation samples and thiophosphate-ester proteins. (D) Immunoblots showing NUP98 in the thiophosphate ester IP fraction when L^{WT} is present. Detection of viral capsid in the lysates was made as control for infection.

STAR Methods

RESOURCE AVAILABILITY

Lead contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Thomas Michiels (thomas.michiels@uclouvain.be)

Materials availability
All reagents generated in this study are available upon request after completion of a Materials Transfer Agreement

Data and code availability
The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD034604"..
MaxQuant outputs are available in the study repository https://github.com/UCLouvain-CBIO/2022-RSK-Nups-VIRO
Codes: Processed data and reproducible proDA analyses scripts are available in the study repository at https://github.com/UCLouvain-CBIO/2022-RSK-Nups-VIRO

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Cells

The subclone of HeLa cells used in this work were HeLa M cells that reportedly have low endogenous RNase L activity (kindly offered by R.H. Silvermann) (Dong et al., 2001).

HeLa-RSK-DKO, (RSK1- and RSK2-deficient) and HeLa-RSK-TKO (RSK-1, RSK2- and RSK3-deficient) cells were obtained from HeLa M cells using the CRISPR-Cas9 technology (Sorgeloos et al., 2022). As HeLa cells express very little RSK3 and virtually no RSK4 transcripts, both cell lines can be considered as RSK-KO cells. One transcribed allele is however detected carrying a 81nt in-frame deletion in RSK1 (Sorgeloos et al., 2022).

HeLa-LVX and HeLa-RSK-TKO-LVX were obtained by transduction of HeLa M cells and of HeLa-RSK-TKO cells with pLVX-EF1alpha-2xGFP:NES-IRES-2xRFP:NLS. Clones showing regular 2xGFP:NES and 2xRFP:NLS expression levels under the fluorescent microscope were selected for further use. HeLa BioID-RSK cells were obtained by transduction of HeLa-RSK-DKO cells with BLP10 and transduced cell populations were then selected with 1mg/ml of G418 (Roche). BioID2-RSK2 expression was then verified by western blot. HeLa WT-RSK, HeLa As1-RSK and HeLa As2-RSK cells were obtained by transduction of HeLa-RSK-TKO cells with TM1117, BLP20 and BLP21 respectively and transduced cell populations were selected with 1mg/ml of G418 (Roche). WT, As1 or As2-RSK expression was then tested by western blot.

HeLa-RSK-TKO cells were transduced with the empty lentiviral vector TM952 or with TM1116-19 derivatives expressing the four isoforms of human RSK. Transduced cell populations were selected with 2mg/ml of G418.

BHK-CL13 and 293T-CL13 cells were obtained by lentiviral transduction of CL13 (TMEV capsid precursor - IRES - mCherry) in the corresponding cells. mCherry-positive cells were then sorted by FACS.

293T cells were kindly provided by F. Tangy (Pasteur Institute, Paris). Both HeLa M and 293T cells and their derivatives were maintained in Dulbecco's modified Eagle medium (DMEM) (Lonza) supplemented with 10% of fetal calf serum (FBS, Sigma), 100 U/ml penicillin and 100μg/ml streptomycin (Thermo Fisher). BHK-21 cells (ATCC), used for viral production and titration, were maintained in Glasgow's modified Eagle medium (GMEM) (Gibco) supplemented with 10% newborn calf serum (Gibco), 100 U/ml penicillin, 100μg/ml streptomycin and 2.6g/l tryptose phosphate broth (Difco). All cells were cultured at 37°C, in 5% CO2.

Viruses

TMEV viruses used in this study are derivatives of KJ6, a variant of the persistent DA strain (DA1 molecular clone) adapted to grow on L929 cells (Jnaoui and Michiels, 1998). FB09 carries the M60V mutation in L (L^{M60V}), which was shown to abrogate L toxicity (Ricour et al., 2009a, Borghese and Michiels, 2011). KJ6 and FB09 are indicated as TMEV L^{WT} and L^{M60V} respectively.

EMCV viruses used is this study are derived from pMC24, a Mengo virus molecular clone carrying a

shortened (24C) polyC tract (Duke and Palmenberg, 1989). The virus denoted EMCV^{WT} (used in Figure

5) is the virus produced from the pMC24 plasmid. Viruses denoted EMCV L^{WT} and L^{Zn} (used in Figure

1 and S1) are produced from plasmids pFS269 and pTM1098 and express N-terminally Flag-tagged L

proteins. L^{Zn} carries two point mutations (C19A and C22A) in the zinc finger of L, which likely affect

the overall structure of the protein (Cornilescu et al., 2008).

These viruses were produced by reverse genetics from plasmid constructions containing the full-length

viral cDNA sequences. To this end, BHK-21 cells were electroporated (1500V, 25μFd, no shunt

resistance) using a Gene pulser apparatus (Bio-Rad) with viral RNA produced by *in vitro* transcription

(RiboMax transcription kit P1300, Promega). Supernatants were collected 48-72h after electroporation,

when cytopathic effect was complete. After 2 to 3 freeze-thaw cycles, the supernatants containing the

virus were clarified by centrifugation at 1258g for 20 min and viruses were stored at -80°C. Viruses

were titrated by plaque assay in BHK-21 cells.

Trans-encapsidated viral replicons

FW12 is the replicon encoding for BioID2-HA-linker(3xGGGGS)-L^{WT}. This replicon derives from

TMEV DA1. It was constructed by replacing the capsid-coding region of the virus by a sequence coding

sequentially (5'-to-3'), in a single frame, for: BioID2-HA-linker(3xGGGGS)-L, the L/VP4 boundary

cleaved by protease 3C, eGFP, the VP2 coding segment encompassing the CRE sequence, the 3' end of

the 2A coding region and the following part of the viral genome (see Figure S2B). The BioID-L^{M60V}

(FW16) and BioID-L^{F48A} (FW17) viral replicons are derivates of FW12 carrying the indicated mutation.

Trans-encapsidation of the replicon occurred in 293T cells transduced with the lentiviral vector CL13

to stably express the viral capsid precursor protein. pCL13, the plasmid carrying the lentiviral vector,

is a pTM945 derivative expressing the 10 C-terminal aminoacids of L followed by the entire capsid

precursor of virus DA1. C-terminal residues of L were inserted in the construct to restore a genuine

L/VP4 clivage site for protease 3C, which allows the capsid protein to start with an N-terminal glycine,

potentially undergoing myristoylation (Chow et al., 1987). Importantly, 4 silent mutations were

introduced in the VP2 coding sequence corresponding to the *CRE* region to prevent the replication of

recombinant wild type viruses that would emerge through recombination between replicon RNA and

capsid-coding mRNA in packaging cells.

293T-CL13 cells were seeded (600,000/well) in 6-well plates. 24h after seeding, cells were transfected

(TransIT-mRNA, Mirus) with 5μg of replicon RNA previously obtained by *in vitro* transcription. Six

hours post-transfection, medium was changed to DMEM without serum. The infected cell population

was passaged 8 times and reseeded with part of the collected supernatant to amplify viral production.

Then, 150ml of replicon-containing supernatant was collected and filtered (0.45μm). SDS was then

added to a final concentration of 0.5% and, after 2h incubation at RT, supernatants were centrifuged for

5min, at 1258g at 20°C. Next, 13ml of clarified supernatant was added to a 2.6ml sucrose 30% cushion

in polyallomer tubes (Beckman) and centrifuged for 16h at 20°C, 25.000rpm in a SW28 swinging bucket rotor. After ultracentrifugation, pellets containing the replicons were resuspended in 150µl of 10mM Tris-HCl pH 7.5. and further dialyzed in 10mM Tris-HCl pH 7.5. Replicons were then titrated by plaque assay using BHK-21 cells stably expressing viral capsid proteins (BHK-21 CL13), and kept at -80°C.

Lentiviral vectors and cell transduction

- pLVX-EF1alpha-2xGFP:NES-IRES-2xRFP:NLS was a gift from Fred Gage (Addgene plasmid # 71396) (Mertens et al., 2015)

Other lentiviral vectors used for protein expression were derived from pCCLsin.PPT.hPGK.GFP.pre. (Follenzi et al., 2000)

- TM945 is a Prom_{CMV}-MCS-IRES-mCherry construct (Hermant et al., 2013)

- TM952 is a Prom_{CMV}-MCS-IRES-neo construct (Cesaro et al., 2021)

- BLP10 is a TM952 derivative coding for BioID2-HA-linker(3xGGGS)-MuRSK2. Note that this construct expresses murine RSK2, which differs from human RSK2 by a single amino acid.

- BLP20 is a TM952 derivative coding for As1-RSK2 (hu-RSK2 with Analog-sensitive kinase 1 (As1) mutation (Leu 147 → Gly).

- BLP21 is a TM952 derivative coding for As2-RSK2 (hu-RSK2 with Analog-sensitive kinase 2 (As2) mutation (Leu 147 → Ala).

- CL13 is a TM945 derivative expressing the 10 C-terminal aminoacids of L followed by the entire capsid precursor of virus DA1.

- TM1116, TM1117, TM1118 and TM1119 are TM952 derivatives carrying the sequences coding for 3xHA-Human RSK1, RSK2, RSK3 and RSK4 respectively (Sorgeloos et al., 2022). They were constructed using the Gateway technology (Invitrogen) from donor plasmids Hs.RPS6KA1, Hs.RPS6KA2, Hs.RPS6KA3 and Hs.RPS6KA6 kindly provided by Dominic Esposito through the Addgene collection (Addgene refs: 70573, 70575, 70577, and 70579, respectively)

Lentiviruses were produced in 293T cells by co-transfection of the following plasmid, using TransIT-LT1 reagent (Mirus Bio): 2.5 µg of lentiviral vector, 0.75 µg of pMD2-VSV-G (VSV-glycoprotein), 1.125 µg of pMDLg/pRRE (Gag-Pol), and 0.625 µg of pRSV-Rev (Rev). DNA quantities are for transfection of 1 well of a 6-well plate. Supernatants were typically collected 72h post transfection and filtered (porosity: 0.45µM). For transduction, cells were typically seeded in a 24-well plate as of 5,000 - 10,000 cells/well and infected 1 or 2 times with 100µL of filtered lentivirus.

METHOD DETAILS

Biotinylation experiments

<u>BioID-RSK</u> : HeLa BioID-RSK cells were seeded in 6-well plates at a density of 90,000 cells per well.
Two 6-well plates were used per condition. 24h after seeding the cells, the medium was changed to
OptiMEM (Gibco) depleted from biotin (previously incubated with streptavidin beads for 2h at 4°C,
then filtered). Cells were kept in OptiMEM without biotin for 48h, then they were infected with 600µl
of virus per well at an MOI of 2.5. One hour post-infection 2ml/well of DMEM containing 5µM biotin
was added. Infection proceeded for 16h. For immunostaining: 1,000 HeLa BioID-RSK cells were
seeded in wells of a 96-well plate. 24h after seeding the cells, the medium was changed to OptiMEM
without biotin. Cells were then incubated without biotin for 48h, then they were infected with 50µL of
virus per well at an MOI of 5. One hour post-infection 150µL of DMEM containing 5µM biotin was
added. Infection proceeded for 10h.
<u>BioID-L</u> : HeLa cells were seeded in 6-well plates as of 160,000 cells per well. Two 6-well plates were
used per condition. 24h after seeding, the cells were infected with 600µl of BioID-L replicons at an
MOI of 2.5. One hour post-infection 2ml/well of DMEM containing 5µM biotin was added. Infection
proceeded for 14h. For immunostaining: 3,500 HeLa cells were seeded in wells of a 96-well plate. 24h
after seeding, the cells were infected with 50µL of virus per well at an MOI of 5. One hour post-infection
150µL of DMEM containing 5µM biotin was added. Infection proceeded for 10h.
565

Streptavidin Pulldown
Cells were washed with PBS 3 times and lysed with 200µL/well of stringent lysis buffer (50mM Tris-
HCl pH 7.6, 500mM NaCl, 0.4% SDS, 1mM DTT, 1 tablet of Pierce phosphatase/protease inhibitor
(Thermo Scientific) per 10ml of lysis buffer) for 15 min at room temperature (RT). Lysates were then
diluted twice by addition of 200µl/well of 50mM Tris-HCl pH7.6 and homogenized by 10 passages
through 21G needles. Lysates were then cleared by centrifugation at 12,000g for 10 min at RT.
Supernatants were then transferred to new tubes, 200µL (per condition) of protein A/G magnetic beads
(Pierce) were added to remove non-specific binding and incubated for 30 min at RT. Supernatants were
then transferred to a new tube and a sample of 160µL per condition was mixed with 80µL of 3x Laemmli
buffer (cell lysate control). The rest of the supernatant was incubated for 2h at RT with 260µL (per
condition) of Streptavidin magnetic beads (Pierce). Streptavidin beads were then washed once with 2%
SDS, and twice with "normal lysis buffer" (50mM Tris-HCl pH 7.5, 100mM NaCl, 2mM EDTA, 0.5%
NP40, 1 tablet of phosphatase/protease inhibitor (Thermo Scientific) per 10ml of lysis buffer) for 5 min
at RT. Beads were resuspended in 40µl of 1x Laemmli buffer and heated for 5min at 100°C to allow
protein's separation from the beads. Supernatants were then conserved at -20°C.
581

Immunostaining
Cells seeded in 96-well screenstar plates (Greiner, 655866) were fixed with PBS-4% PFA for 5 min at
RT. Cells were then permeabilized with PBS-0.2% Triton X-100 (ICN Biomedicals Inc.) for 5 min at
RT. Cells were then blocked with TNB blocking reagent (Perkin Elmer) for 1h at RT. Antibodies were

diluted in TNB blocking reagent: rabbit anti-TMEV 3D polymerase 1/1000 (rabbit, kindly provided by

M.Brahic), anti-2A 1/1200 (rabbit, home-made), anti-NUP98 (rat, N1038, Sigma) 1/400, anti-POM121

(rabbit, 15645-1-AP, Proteintech) 1/400 or anti-PTB (mouse, 324800, Invitrogen) 1/400; and added to

the cells and incubated for 1h at RT. Cells were then washed three times with PBS-0.1% Tween 20 for

5 min. Cells were then incubated with secondary antibodies (anti-rat Alexa Fluor 594, anti-rabbit Alexa

Fluor 488 or anti-rabbit Alexa Fluor 594, 1/800, anti-mouse Alexa Fluor 488 1/400) or streptavidin

(Alexa fluor 488 or Alexa fluor 594, 1/500) for 1h at RT. Finally, cells were washed 3 times with PBS-

0.1% Tween 20 and kept in PBS-Azide 0.02% until analysis. Fluorescence analysis was made with a

spinning disk confocal microscope (Zeiss) and adjustment of the images was performed using the Zen

image analysis software (Zeiss). Exposure time, image brightness and contrast were adjusted equally

within an experiment. For LVX cells (qualitative analysis), brightness and contrast were adjusted using

the automatic "best fit" setting of the the Zen software.

Western blotting

Proteins in Laemmli buffer were heated for 5 minutes at 100°C. They were then run in 8 or 10% tris-

glycine SDS polyacrylamide gels and then transferred to either polyvinylidenefluoride or nitrocellulose

membranes (only for Mab414 detection). Membranes were blocked with TBS-5% non-fat dry milk

(Regilait) or TBS-5% BSA (for streptavidin-HRP blots) for 1 hour at RT. Primary antibodies: anti-2A

(TMEV, rabbit home-made 1/4000), anti-Phospho-S380-RSK(rabbit, CST11989 – Cell signaling

technology, 1/5000), anti-PKR (rabbit, 18244-1-AP – Proteintech, 1/4000), anti-Phospho-T446-PKR

(rabbit, AB32036 – Abcam, 1/4000), anti-ß-actin (mouse, A5441 – Sigma, 1/10,000), anti-TMEV 3D

polymerase (rabbit, kindly provided by M.Brahic, 1/2000), anti-NUP98 (rat, N1038 – Sigma, 1/2000),

, anti-HA (mouse, MMS101P – Covance, 1/4000), anti-phospho-Akt-substrates: RxxS*/T* (phospho-

RSK substrates, rabbit, CST9614 – Cell signaling technology, 1/1000), anti-thio-P-ester (rabbit, NBP2-

67738 – Novusbio, 1/1000), anti-TMEV capsid (home-made, 1/2000), anti-FLAG M2 (mouse, Sigma,

F1804 1/5000) were diluted in blocking solution and were incubated with membranes over-night.

Membranes were then washed 3 times for 15min with TBS-0.1% Tween 20. Secondary antibodies (anti-

rabbit, anti-mouse or anti-rat coupled to HRP, Dako, 1/5000) or Streptavidin-HRP (Dako, 1/1000) were

then added for 1h at RT. Then, membranes were washed three times with TBS-0.1% Tween 20, once

with TBS and revelation was made with SuperSignal West chemiluminescence substrate (Pico or Dura,

Thermoscientific).

Mass spectrometry

Streptavidin pulldown samples were resolved using a 10% Tris-Glycine SDS gel run until 6mm

migration in the separating gel. Proteins were colored using PageBlue (Thermo Scientific, 24620). The

6mm bands containing whole proteins were cut into 3 different slices and trypsin digested. Peptides

were dissolved in solvent A (0.1% TFA in 2% ACN), directly loaded onto reversed-phase pre-column

(Acclaim PepMap 100, Thermo Scientific) and eluted in backflush mode. Peptide separation was performed using a reversed-phase analytical column (Acclaim PepMap RSLC, 0.075 x 250 mm,Thermo Scientific) with a linear gradient of 4%-27.5% solvent B (0.1% FA in 98% ACN) for 35 min, 27.5%-40% solvent B for 5 min, 40%-95% solvent B for 5 min and holding at 95% for the last 5 min at a constant flow rate of 300 nl/min on an EASY-nLC 1000 RSLC system. The peptides were analyzed by an Orbitrap Fusion Lumos tribrid mass spectrometer (ThermoFisher Scientific). The peptides were subjected to NSI source followed by tandem mass spectrometry (MS/MS) in Fusion Lumos coupled online to the nano-LC. Intact peptides were detected in the Orbitrap at a resolution of 120,000. Peptides were selected with the quadrupole for MS/MS using HCD setting at 35; ion fragments were detected in the ion trap. A data-dependent procedure that alternated between one MS scan followed by MS/MS scans was applied for 3 seconds for ions above a threshold ion count of 5.0E3 in the MS survey scan with 30.0s dynamic exclusion. The electrospray voltage applied was 2.1 kV. MS1 spectra were obtained with an AGC target of 4E5 ions and a maximum injection time of 50ms, and MS2 spectra were acquired with an AGC target of 1E4 ions and a maximum injection set to 35ms. For MS scans, the m/z scan range was 375 to 1800. The resulting MS/MS data was processed using Sequest HT search engine within Proteome Discoverer 2.5 SP1 against a *Homo Sapiens* protein database obtained from Uniprot (78 787 entries) and containing the sequences of viral proteins and BioID2-RSK. Trypsin was specified as cleavage enzyme allowing up to 2 missed cleavages, 4 modifications per peptide and up to 5 charges. Mass error was set to 10 ppm for precursor ions and 0.1 Da for fragment ions. Oxidation on Met (+15.995 Da), phosphorylation on Ser, Thr and Tyr (+79.966 Da), conversion of Gln (-17.027 Da) or Glu (- 18.011 Da) to pyro-Glu at the peptide N-term, biotinylation of Lys (+ 226.077) were considered as variable modifications. False discovery rate (FDR) was assessed using Percolator and thresholds for protein, peptide and modification site were specified at 1%.

PSM Ratio calculations were as followed:

$$BioID-RSK: \frac{PSM(L^{WT})}{PSM(L^{M60V})+PSM(mock)+1} \qquad BioID-L: \frac{PSM(L^{WT})+PSM(L^{F48A})}{PSM(L^{M60V})+PSM(mock)+1}$$

GST-S6 recombinant production and purification

ER2566 electrocompetent bacteria were electroporated with pFS208 (plasmid coding for GST-S6) and cultured in 10mL of TSB at 37°C O/N. Bacteria were then amplified at 37°C in 300mL of TSB until DO600 reaches 1. At this point, we added IPTG at a final concentration of 0.2mM and let it at 37°C for 2 hours. Then bacteria were centrifuged at 6,000rpm 15min 4°C and resuspended in PBS, 1% Triton, 0.2mM PMSF, 2mM EDTA, 0.1% ß-mercaptoethanol. Bacteria were lysed with French press and centrifuged at 21,000g for 20 min at 4°C. Supernatant containing soluble protein was then kept for GST-pulldown. GST-pulldown was performed with agarose-glutathion beads. GST-S6 was eluted from

658 the beads with 50mM Tris-HCl pH8 – 10mM glutathion and then purified by dialysis with 25mM Hepes
659 buffer at 4°C.
660

***In vitro* kinase assay**
293T cells were seeded (500,000 cells/well) in 6-well plates. 24h after seeding, cells were transfected
with TransIT-LT1 reagent (Mirus Bio) with 2.5 µg of plasmids pTM1117 (coding for WT-RSK2),
pBLP20 (coding for As1-RSK2) and pBLP21 (coding for As2-RSK2). 6 hours post-transfection, cells
were treated with 32nM PMA over-night. HA-RSKs were then immunoprecipated with anti-HA
magnetic beads. Therefore, cells were washed once with PBS and lysed with 150µL of lysis buffer
(50mM Tris-HCl pH8, 100mM NaCl, 0.5% NP40, 2mM EDTA, 1 tablet of phosphatase/protease
inhibitor (Thermo Scientific) per 10ml of lysis buffer) for 15min at 4°C. Lysates were then
homogenized by 10 passages through 21G needles and cleared by centrifugation at 12,000g for 10min
at 4°C. Supernatants were then transferred to new tubes, 12,5µl (per condition) of protein A/G magnetic
beads (Pierce) were added to remove non-specific binding and incubated for 30 min at 4°C.
Supernatants were then transferred to a new tube and a sample of 20µl/ condition was mixed with 10µL
of 3x Laemmli buffer (cell lysate control). The rest of the supernatant was incubated with 12.5µl of
HA-magnetic beads (Pierce) for 2h at 4°C. Beads were then washed 3 times with lysis buffer for 5 min
at 4°C and resuspended in 30µl of TBS, 1mM PMSF, 1mM Na_3VO_4, 1 tablet of phosphatase/protease
inhibitor (Thermo Scientific) per 10ml.
For one *in vitro* kinase reaction we put: 2.5µg of GST-S6 (recombinant substrate), 10µL of kinase buffer
5x (125mM Hepes pH 7.5, 250mM NaCl, 100mM ß-glycerophosphate, 5mM DTT, 100mM $MgCl_2$,
500µM Na_3VO_4), 0.5 µL of 10mM ATP (Roth, HN35.1) or A*TP analogs (BioLog), and water to a
final volume of 30µL. To this we added 10µL of the immunoprecipitated RSK and incubated everything
at 30°C for 30 min with shacking. Following this, the samples were alkylated with PNBM (Abcam) at
a final concentration of 2.5mM for 2 hours at room temperature with shacking. Reaction was stopped
by the addition of Laemmli buffer 3x. Samples were then heated at 100°C for 5 min, separated from the
magnetic beads and kept at -20°C.
685

Thiophosphorylation in permeabilized cells
1.5 x10^6 HeLa-RSK-TKO TM1117(WT-RSK2) or BLP21 (As2-RSK2) cells were seeded in 10-cm
dishes, 1 10-cm dish per condition. The next day, cells were infected at an MOI of 5 for 8h30min (for
TMEV) and 3h30min (for EMCV). After that, cells were permeabilized with 500µL of the analog-
kinase buffer: 20mM Hepes pH 7.5, 100mM KOAc, 5mM NaOAc, 2mM $MgOAc_2$, 1mM EGTA,
20µg/ml digitonin, 10mM $MgCl_2$, 0.5mM DTT, 1x phosphatase inhibitor cocktail 2 (P5726, Merck),
1x cOmplete protease inhibitor (REF), 57µg/ml creatin kinase (Calbiochem 23895), 5mM Creatin
phosphate (Calbiochem, 2380), 0.1mM ATP, 0.1 mM N6-Bn-A*TP analog (BioLog), 3mM GTP (Roth,

K056.4) in order to get the ATP analog inside the cells and the thiophosphorylation reaction to happen.

Reaction proceeded for 1h at 37°C, 5% CO2 with the dishes on a rocking plate.

Immunoprecipitation after thiophosphorylation reactions.

NUP98 IP: cells were lysed by adding 500µL of 2x salty lysis buffer (100mM Tris-HCl pH8, 800mM

NaCl, 2% Triton X-100, 4mM EDTA, 2mM DTT, 1 tablet of phosphatase/protease inhibitor (Thermo

Scientific) per 10ml of lysis buffer) to the analog-kinase buffer for 15 min at 4°C. Lysates were diluted

4x by addition of 3ml of regular lysis buffer (50mM Tris-HCl pH8, 100mM NaCl, 0.5% NP40, 2mM

EDTA, 1 tablet of phosphatase/protease inhibitor (Thermo Scientific) per 10ml of lysis buffer). Lysates

were then homogenized by 10 passages through 21G needles and cleared by centrifugation at 12,000g

for 10min at 4°C. Supernatants were then transferred to new tubes, 30µL (per condition) of protein A/G

magnetic beads (Pierce) were added to remove non-specific binding and incubated for 30 min at 4°C.

Supernatants were then transferred to a new tube and a sample of 200µL/ condition was alkylated with

PNBM at a final concentration of 2.5mM for 2h at RT. Then alkylation reaction was stopped by addition

of 3x Laemmli buffer (cell lysate control). The rest of the supernatant was incubated O/N at 4°C with

8µg (per condition) of anti- NUP98 (rat, N1038 – Sigma). 62.5µl of A/G beads per condition were

added and incubated for 2h at 4°C. A/G beads were then washed three times with regular lysis buffer

for 5 min at 4°C. Beads were resuspended in 40µl of kinase buffer 1x. The IP samples were then

alkylated with PNBM at a final concentration of 2.5mM for 2h at RT. Alkylation reaction was stopped

by the addition of 3x Laemmli, and immunoprecipitated proteins were separated from the beads after

heating 5min at 100°C.

Thiophosphate ester IP: cells were lysed by adding 500µL of 2x lysis buffer (100mM Tris-HCl pH8,

200mM NaCl, 1% NP40, 40mM EDTA, 1 tablet of phosphatase/protease inhibitor (Thermo Scientific)

per 10ml of lysis buffer) for 15min at 4°C. Lysates were then homogenized by 10 passages through

21G needles and cleared by centrifugation at 12,000g for 10min at 4°C. Supernatants were then

alkylated with PNBM at a final concentration of 2.5mM for 1h30min at RT. As PNBM inhibits

immunoprecipitation, lysates were run through PD10 columns to remove PNBM prior to the

immunoprecipitation procedure. Fractions containing protein were kept and incubated with 30µl per

condition of A/G magnetic beads (to avoid non-specific binding) for 30 min at 4°C. Supernatants were

then transferred to a new tube and 1/10th of the lysate (200µl) was mixed with 100µl of 3x Laemmli

(cell lysate control). The rest of the supernatant was incubated with 8µg per condition of anti-

thiophosphate ester antibody (rabbit, NBP2-67738 – Novusbio) for 2h at 4°C. Then 62.5µl per condition

of A/G magnetic beads were added and incubated for 2h at 4°C. A/G beads were then washed three

times with regular lysis buffer for 5 min at 4°C and resuspended in 60µl of Laemmli 1x.

Immunoprecipitated proteins were separated from the beads after heating 5min at 100°C.

QUANTIFICATION AND STATISTICAL ANALYSIS

Data processing and statistical analysis (for MS)

Protein identification and label-free quantitation were performed with MaxQuant version 1.6.7.0.(Cox and Mann, 2008). Database searching was performed against the UniProt FASTA database, using a false-discovery rate at the peptide and protein level was set to 0.01 and allowing a maximum of two missed cleavages. All search parameters are available in the parameter file available in the study repository (https://github.com/UCLouvain-CBIO/2022-RSK-Nups-VIRO). After filtering out of identified contaminants, identified reverse sequences and proteins having missing values in all samples, Max Quant protein intensities were log-2 transformed and normalized using sample median alignment. In order to cope with the numerous drop-outs in protein intensities, the proDA method, as described in (Ahlmann-Eltze and Anders, 2020) was applied.

For the analysis of the BioID-L experiment, the backbone linear model included one indicator variable representing a potential batch effect between 2 groups of replicates, and 2 condition indicator variables, representing the effect of resp. WT and F48A conditions with respect to the reference condition, i.e. M60V condition. Potential batch effect was tested for each protein using two-sided Wald test on the batch indicator variable. Since adjusted p-values were > 0.2 for all proteins, this indicator variable was subsequently removed from the model. Then, the effects of WT and F48 conditions (w.r.t. M60V) were tested, for each protein, using one-sided Wald tests, since the anticipated sign of the effect was known. All p-values were adjusted using Benjamini-Hochberg corrections (Benjamini and Hochberg, 1995).

For the analysis of the BioID-RSK experiment, the backbone linear model included one indicator variable representing a potential batch effect between 2 groups of replicates, and 1 condition indicator variable, representing the effect of WT condition with respect to the reference condition, i.e. M60V condition. Potential batch effect was tested for each protein using two-sided Wald test on the batch indicator variable. Since adjusted p-values were > 0.6 for all proteins, this indicator variable was subsequently removed from the model. Then, the effect of WT condition (w.r.t. M60V) was tested, for each protein, using a one-sided Wald test, since the anticipated sign of the effect was known. All p-values were adjusted using Benjaminin-Hochberg corrections.

Statistical analysis on IF

763 Statistical analysis on immunofluorescence experiments was done using GraphPad Prism v9 A one-
764 way ANOVA test was used. The number of independent experiments (n) and statistical comparison
765 groups are indicated in the figures and figure legends.
766

References

AHLMANN-ELTZE, C. & ANDERS, S. 2020. proDA: Probabilistic Dropout Analysis for
Identifying Differentially Abundant Proteins in Label-Free Mass Spectrometry.
bioRxiv, 661496.
ALEXA, A., SOK, P., GROSS, F., ALBERT, K., KOBORI, E., PÓTI Á, L., GÓGL, G.,
BENTO, I., KUANG, E., TAYLOR, S. S., ZHU, F., CILIBERTO, A. & REMÉNYI,
A. 2022. A non-catalytic herpesviral protein reconfigures ERK-RSK signaling by
targeting kinase docking systems in the host. *Nat Commun,* 13, 472.
ALLEN, J. J., LI, M., BRINKWORTH, C. S., PAULSON, J. L., WANG, D., HÜBNER, A.,
CHOU, W. H., DAVIS, R. J., BURLINGAME, A. L., MESSING, R. O.,
KATAYAMA, C. D., HEDRICK, S. M. & SHOKAT, K. M. 2007. A semisynthetic
epitope for kinase substrates. *Nat Methods,* 4, 511-6.
ARAMBURU, I. V. & LEMKE, E. A. 2017. Floppy but not sloppy: Interaction mechanism of
FG-nucleoporins and nuclear transport receptors. *Semin Cell Dev Biol,* 68, 34-41.
BARDINA, M. V., LIDSKY, P. V., SHEVAL, E. V., FOMINYKH, K. V., VAN
KUPPEVELD, F. J., POLYAKOV, V. Y. & AGOL, V. I. 2009. Mengovirus-induced
rearrangement of the nuclear pore complex: hijacking cellular phosphorylation
machinery. *J Virol,* 83, 3150-61.
BECK, M. & HURT, E. 2017. The nuclear pore complex: understanding its function through
structural insight. *Nat Rev Mol Cell Biol,* 18, 73-89.
BELOV, G. A., LIDSKY, P. V., MIKITAS, O. V., EGGER, D., LUKYANOV, K. A., BIENZ,
K. & AGOL, V. I. 2004. Bidirectional increase in permeability of nuclear envelope
upon poliovirus infection and accompanying alterations of nuclear pores. *J Virol,* 78,
10166-77.
BENJAMINI, Y. & HOCHBERG, Y. 1995. Controlling the False Discovery Rate: A Practical
and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society.*
Series B (Methodological), 57, 289-300.
BORGHESE, F. & MICHIELS, T. 2011. The leader protein of cardioviruses inhibits stress
granule assembly. *J Virol,* 85, 9614-22.
BORGHESE, F., SORGELOOS, F., CESARO, T. & MICHIELS, T. 2019. The Leader Protein
of Theiler's Virus Prevents the Activation of PKR. *J Virol,* 93.
BRAHIC, M., BUREAU, J. F. & MICHIELS, T. 2005. The genetics of the persistent infection
and demyelinating disease caused by Theiler's virus. *Annu Rev Microbiol,* 59, 279-98.
CESARO, T., HAYASHI, Y., BORGHESE, F., VERTOMMEN, D., WAVREIL, F. &
MICHIELS, T. 2021. PKR activity modulation by phosphomimetic mutations of serine
residues located three aminoacids upstream of double-stranded RNA binding motifs.
Sci Rep, 11, 9188.
CHEN, R. H., SARNECKI, C. & BLENIS, J. 1992. Nuclear localization and regulation of erk-
and rsk-encoded protein kinases. *Mol Cell Biol,* 12, 915-27.
CHOW, M., NEWMAN, J. F., FILMAN, D., HOGLE, J. M., ROWLANDS, D. J. & BROWN,
F. 1987. Myristylation of picornavirus capsid protein VP4 and its structural
significance. *Nature,* 327, 482-6.
CHUNG, L. K., PARK, Y. H., ZHENG, Y., BRODSKY, I. E., HEARING, P., KASTNER, D.
L., CHAE, J. J. & BLISKA, J. B. 2016. The Yersinia Virulence Factor YopM Hijacks

Host Kinases to Inhibit Type III Effector-Triggered Activation of the Pyrin
Inflammasome. *Cell Host Microbe,* 20**,** 296-306.

CORNELIS, G. R. & WOLF-WATZ, H. 1997. The Yersinia Yop virulon: a bacterial system
for subverting eukaryotic cells. *Mol Microbiol,* 23**,** 861-7.

CORNILESCU, C. C., PORTER, F. W., ZHAO, K. Q., PALMENBERG, A. C. & MARKLEY,
J. L. 2008. NMR structure of the mengovirus Leader protein zinc-finger domain. *FEBS*
Lett, 582**,** 896-900.

COX, J. & MANN, M. 2008. MaxQuant enables high peptide identification rates,
individualized p.p.b.-range mass accuracies and proteome-wide protein quantification.
Nat Biotechnol, 26**,** 1367-72.

DE JESÚS-GONZÁLEZ, L. A., PALACIOS-RÁPALO, S., REYES-RUIZ, J. M., OSUNA-
RAMOS, J. F., CORDERO-RIVERA, C. D., FARFAN-MORALES, C. N.,
GUTIÉRREZ-ESCOLANO, A. L. & DEL ÁNGEL, R. M. 2021. The Nuclear Pore
Complex Is a Key Target of Viral Proteases to Promote Viral Replication. *Viruses,* 13**,**
706.

DELHAYE, S., VAN PESCH, V. & MICHIELS, T. 2004. The leader protein of Theiler's virus
interferes with nucleocytoplasmic trafficking of cellular proteins. *J Virol,* 78**,** 4357-62.

DONG, B., NIWA, M., WALTER, P. & SILVERMAN, R. H. 2001. Basis for regulated RNA
cleavage by functional analysis of RNase L and Ire1p. *Rna,* 7**,** 361-73.

DUKE, G. M. & PALMENBERG, A. C. 1989. Cloning and synthesis of infectious cardiovirus
RNAs containing short, discrete poly(C) tracts. *J Virol,* 63**,** 1822-6.

FOLLENZI, A., AILLES, L. E., BAKOVIC, S., GEUNA, M. & NALDINI, L. 2000. Gene
transfer by lentiviral vectors is limited by nuclear translocation and rescued by HIV-1
pol sequences. *Nat Genet,* 25**,** 217-22.

FREUNDT, E. C., DRAPPIER, M. & MICHIELS, T. 2018. Innate Immune Detection of
Cardioviruses and Viral Disruption of Interferon Signaling. *Front Microbiol,* 9**,** 2448.

HAMPOELZ, B., ANDRES-PONS, A., KASTRITIS, P. & BECK, M. 2019. Structure and
Assembly of the Nuclear Pore Complex. *Annu Rev Biophys,* 48**,** 515-536.

HATO, S. V., RICOUR, C., SCHULTE, B. M., LANKE, K. H., DE BRUIJNI, M., ZOLL, J.,
MELCHERS, W. J., MICHIELS, T. & VAN KUPPEVELD, F. J. 2007. The
mengovirus leader protein blocks interferon-alpha/beta gene transcription and inhibits
activation of interferon regulatory factor 3. *Cell Microbiol,* 9**,** 2921-30.

HENTSCHKE, M., BERNEKING, L., BELMAR CAMPOS, C., BUCK, F.,
RUCKDESCHEL, K. & AEPFELBACHER, M. 2010. Yersinia virulence factor YopM
induces sustained RSK activation by interfering with dephosphorylation. *PLoS One,* 5.

HERMANT, P., FRANCIUS, C., CLOTMAN, F. & MICHIELS, T. 2013. IFN-ε is
constitutively expressed by cells of the reproductive tract and is inefficiently secreted
by fibroblasts and cell lines. *PLoS One,* 8**,** e71320.

HÜLSMANN, B. B., LABOKHA, A. A. & GÖRLICH, D. 2012. The permeability of
reconstituted nuclear pores provides direct evidence for the selective phase model. *Cell,*
150**,** 738-51.

JNAOUI, K. & MICHIELS, T. 1998. Adaptation of Theiler's virus to L929 cells: mutations in
the putative receptor binding site on the capsid map to neutralization sites and modulate
viral persistence. *Virology,* 244**,** 397-404.

KIM, D. I., JENSEN, S. C., NOBLE, K. A., KC, B., ROUX, K. H., MOTAMEDCHABOKI,
K. & ROUX, K. J. 2016. An improved smaller biotin ligase for BioID proximity
labeling. *Mol Biol Cell,* 27**,** 1188-96.

KOSAKO, H., YAMAGUCHI, N., ARANAMI, C., USHIYAMA, M., KOSE, S., IMAMOTO,
N., TANIGUCHI, H., NISHIDA, E. & HATTORI, S. 2009. Phosphoproteomics reveals

new ERK MAP kinase targets and links ERK to nucleoporin-mediated nuclear
transport. *Nat Struct Mol Biol,* 16**,** 1026-35.
KUANG, E., FU, B., LIANG, Q., MYOUNG, J. & ZHU, F. 2011. Phosphorylation of
eukaryotic translation initiation factor 4B (EIF4B) by open reading frame 45/p90
ribosomal S6 kinase (ORF45/RSK) signaling axis facilitates protein translation during
Kaposi sarcoma-associated herpesvirus (KSHV) lytic replication. *J Biol Chem,* 286**,**
41171-41182.
LI, X., DU, S., AVEY, D., LI, Y., ZHU, F. & KUANG, E. 2015. ORF45-Mediated Prolonged
c-Fos Accumulation Accelerates Viral Transcription during the Late Stage of Lytic
Replication of Kaposi's Sarcoma-Associated Herpesvirus. *J Virol,* 89**,** 6895-906.
LIDSKY, P. V., HATO, S., BARDINA, M. V., AMINEV, A. G., PALMENBERG, A. C.,
SHEVAL, E. V., POLYAKOV, V. Y., VAN KUPPEVELD, F. J. & AGOL, V. I. 2006.
Nucleocytoplasmic traffic disorder induced by cardioviruses. *J Virol,* 80**,** 2705-17.
LINDER, M. I., KÖHLER, M., BOERSEMA, P., WEBERRUSS, M., WANDKE, C.,
MARINO, J., ASHIONO, C., PICOTTI, P., ANTONIN, W. & KUTAY, U. 2017.
Mitotic Disassembly of Nuclear Pore Complexes Involves CDK1- and PLK1-Mediated
Phosphorylation of Key Interconnecting Nucleoporins. *Dev Cell,* 43**,** 141-156.e7.
LIZCANO-PERRET, B. & MICHIELS, T. 2021. Nucleocytoplasmic Trafficking Perturbation
Induced by Picornaviruses. *Viruses,* 13**,** 1210.
LOBERT, P. E., ESCRIOU, N., RUELLE, J. & MICHIELS, T. 1999. A coding RNA sequence
acts as a replication signal in cardioviruses. *Proc Natl Acad Sci U S A,* 96**,** 11560-5.
LOPEZ, M. S., KLIEGMAN, J. I. & SHOKAT, K. M. 2014. The logic and design of analog-
sensitive kinases and their small molecule inhibitors. *Methods Enzymol,* 548**,** 189-213.
MCDONALD, C., VACRATSIS, P. O., BLISKA, J. B. & DIXON, J. E. 2003. The yersinia
virulence factor YopM forms a novel protein complex with two cellular kinases. *J Biol*
Chem, 278**,** 18514-23.
MERTENS, J., PAQUOLA, A. C. M., KU, M., HATCH, E., BÖHNKE, L., LADJEVARDI,
S., MCGRATH, S., CAMPBELL, B., LEE, H., HERDY, J. R., GONÇALVES, J. T.,
TODA, T., KIM, Y., WINKLER, J., YAO, J., HETZER, M. W. & GAGE, F. H. 2015.
Directly Reprogrammed Human Neurons Retain Aging-Associated Transcriptomic
Signatures and Reveal Age-Related Nucleocytoplasmic Defects. *Cell Stem Cell,* 17**,**
705-718.
MISHRA, A., SIPMA, W., VEENHOFF, L. M., VAN DER GIESSEN, E. & ONCK, P. R.
2019. The Effect of FG-Nup Phosphorylation on NPC Selectivity: A One-Bead-Per-
Amino-Acid Molecular Dynamics Study. *Int J Mol Sci,* 20.
PORTER, F. W., BROWN, B. & PALMENBERG, A. C. 2010. Nucleoporin phosphorylation
triggered by the encephalomyocarditis virus leader protein is mediated by mitogen-
activated protein kinases. *J Virol,* 84**,** 12538-48.
PORTER, F. W. & PALMENBERG, A. C. 2009. Leader-induced phosphorylation of
nucleoporins correlates with nuclear trafficking inhibition by cardioviruses. *J Virol,* 83**,**
1941-51.
RATNER, D., ORNING, M. P., PROULX, M. K., WANG, D., GAVRILIN, M. A., WEWERS,
M. D., ALNEMRI, E. S., JOHNSON, P. F., LEE, B., MECSAS, J., KAYAGAKI, N.,
GOGUEN, J. D. & LIEN, E. 2016. The Yersinia pestis Effector YopM Inhibits Pyrin
Inflammasome Activation. *PLoS Pathog,* 12**,** e1006035.
RICOUR, C., BORGHESE, F., SORGELOOS, F., HATO, S. V., VAN KUPPEVELD, F. J. &
MICHIELS, T. 2009a. Random mutagenesis defines a domain of Theiler's virus leader
protein that is essential for antagonism of nucleocytoplasmic trafficking and cytokine
gene expression. *J Virol,* 83**,** 11223-32.

RICOUR, C., DELHAYE, S., HATO, S. V., OLENYIK, T. D., MICHEL, B., VAN
KUPPEVELD, F. J., GUSTIN, K. E. & MICHIELS, T. 2009b. Inhibition of mRNA
export and dimerization of interferon regulatory factor 3 by Theiler's virus leader
protein. *J Gen Virol,* 90**,** 177-86.
SAEED, M., KAPELL, S., HERTZ, N. T., WU, X., BELL, K., ASHBROOK, A. W., MARK,
M. T., ZEBROSKI, H. A., NEAL, M. L., FLODSTRÖM-TULLBERG, M.,
MACDONALD, M. R., AITCHISON, J. D., MOLINA, H. & RICE, C. M. 2020.
Defining the proteolytic landscape during enterovirus infection. *PLoS Pathog,* 16**,**
e1008927.
SORGELOOS, F., PEETERS, M., HAYASHI, Y., BORGHESE, F., CAPELLI, N.,
DRAPPIER, M., CESARO, T., COLAU, D., STROOBANT, V., VERTOMMEN, D.,
DE BODT, G., MESSE, S., FORNÉ, I., MUELLER-PLANITZ, F., COLLET, J. F. &
MICHIELS, T. 2022. A case of convergent evolution: Several viral and bacterial
pathogens hijack RSK kinases through a common linear motif. *Proc Natl Acad Sci U S*
A, 119.
VAN PESCH, V., VAN EYLL, O. & MICHIELS, T. 2001. The leader protein of Theiler's
virus inhibits immediate-early alpha/beta interferon production. *J Virol,* 75**,** 7811-7.
ZOLL, J., MELCHERS, W. J., GALAMA, J. M. & VAN KUPPEVELD, F. J. 2002. The
mengovirus leader protein suppresses alpha/beta interferon production by inhibition of
the iron/ferritin-mediated activation of NF-kappa B. *J Virol,* 76**,** 9664-72.
929

Figure 1.

Figure 2.

Figure 3.

Figure 4.

A

Mock   L^WT   L^M60V

Biotinylated proteins 3D

B

Cells with perinuclear staining (%)

****   ****

TMEV   Mock   L^WT   L^M60V

C

Mock   BioID-L^WT   BioID-L^M60V   BioID-L^F48A

Biotinylated proteins 3D

D

Cells with perinuclear staining (%)

****

****   ****   ****

TMEV   Mock   BioID-L^WT   BioID-L^M60V   BioID-L^F48A

Figure 5.

A

B

C

D