Different approaches to processing environmental DNA samples in turbid

waters have distinct effects for fish, bacterial and archaea communities.

Authors: Rachel Turba[1*], Glory H. Thai[1], David K. Jacobs[1]

[1]Department of Ecology and Evolutionary Biology, University of California, Los Angeles,

California 90095 USA

*Corresponding author: rturba@outlook.com

Keywords: metabarcoding; turbidity; method comparison; fish

Abstract

Coastal lagoons are an important habitat for endemic and threatened species in California that

have suffered impacts from urbanization and increased drought. Environmental DNA has been

promoted as a way to aid in the monitoring of biological communities, but much remains to be

understood on the biases introduced by different protocols meant to overcome challenges

presented by unique systems under study. Turbid water is one methodologic challenge to eDNA

recovery in these systems as it quickly clogs filters, preventing timely processing of samples. We

investigated biases in community composition produced by two solutions to overcome slow

filtration due to turbidity: freezing of water prior to filtration (for storage purposes and long-term

processing), and use of sediment (as opposed to water samples). Bias assessments of community

composition in downstream eDNA analysis was conducted for two sets of primers, 12S (fish)

and 16S (bacteria and archaea). Our results show that freezing water prior to filtration had no

effects on community composition for either primer, even when using a filter of larger pore size

(3 μm), and therefore it is a viable approach in this system for comparison of water borne fish,

bacteria and archaea. However, the 16S primer showed significantly different community

composition in sediments compared to water samples, although still recovering eDNA of

organisms from the water column. Sediment sample replicates were heterogeneous, and therefore

increasing the number of replicates would be recommended for similar habitats.

Introduction

Coastal lagoons in California are the numerically dominant form of coastal wetland (Jacobs et

al., 2011; Stein et al., 2014) and are important in many other Mediterranean climates and

subtropical environments. These lagoons are characterized by seasonal and episodic breaching

(opening of the lagoon to the sea, usually by stream flow) and closure (isolation of the lagoon by

a high sandbar), which provide a suite of ecological services: from groundwater infiltration to

support of unique biodiversity (Ballard et al., n.d.). This system serves as important habitat and

nursery for endemic and endangered fishes and amphibians, such as the steelhead

(*Oncorhynchus mykiss*), red-legged frog (*Rana aurora draytonii*), and the tidewater goby

(*Eucyclogobius newberryi*) (Earl et al., 2010; Shaffer et al., 2004; Swift et al., 1993, 2016). Thus,

California lagoons are spatially and temporally variable systems with unique biodiversity and

biodiversity assessment challenges.

Coastal lagoons have been drastically reduced in numbers along the California coastline,

driven mostly by the impact of coastal land use for transport structures, agriculture, and

development. These are further exacerbated by ongoing changes in the hydrological cycles due

to climate change (SCWRP, 2018). While these sites are critical for endangered species

conservation, they are also subject to frequent invasion and their response to environmental

variation is poorly documented. However, monitoring of this habitat can be limited by a variety

of issues, ranging from limited human power and access to challenges driven by the natural

complexity and dynamism of these lagoons.

The use of environmental DNA (eDNA) has been advocated as an alternative for

monitoring communities and target species (Thomsen & Willerslev, 2015), and can overcome

and complement certain field limitations from traditional methods (e.g. seining, trapping). On-

site collection can be relatively fast, and therefore allow field workers to cover more ground. It

can also recover the DNA signal of species that are rare, cryptic and/or hard to capture by

traditional methods, and being non-intrusive, it offers an alternative when working with

endangered species for which permits are necessary (Deiner et al., 2017; Dejean et al., 2012;

Sard et al., 2019). In addition, metabarcoding approaches allow the investigation of multiple

species from a single collection (Taberlet, Coissac, et al., 2012).

Nevertheless, it is important to recognize that this approach also brings its own

limitations and biases (van der Loos & Nijland, 2021). In some circumstances, eDNA sampling

can be more expensive than traditional, more established methods (Smart et al., 2016). Since

there are no voucher specimens from collections, contamination is a major issue that needs to be

addressed early on, following best practices in the field (Goldberg et al., 2016). The lack of

voucher specimens also leads to an overdependence on the use of barcodes and genetic databases

for taxonomic identification, which introduces another set of biases, from misidentification to

lack of species representation (Taberlet, Coissac, et al., 2012). Other challenges arise from the

non-universality of sampling methods and downstream processing, with the probability of

detection varying depending on the species and their density, as well as the type of environment,

which affects rates of DNA degradation (Deiner et al., 2015; Nagarajan et al., 2022; Rees et al.,

2014; Williams et al., 2017).

Coastal lagoons can vary drastically in their environmental properties. One major

challenge is the high and variable turbidity of the water. High turbidity often occurs during high

stream flow, or when lagoons are closed to the ocean by a sandbar leading to an accumulation of

organic and inorganic matter. In this scenario, filtering turbid water on-site is problematic.

However, filtration is widely used for handling water samples (Laramie et al., 2015; Tsuji et al.,

2019). Typically, set volumes of water are run through a small pore size filter to concentrate

DNA before extractions. However, high concentration of fine sediment or organic matter in

water quickly obstructs these filters, making the filtration process time-consuming (although it

could actually aid recovery by binding DNA to suspended particles: Kumar et al., 2022; Liang &

Keeley, 2013; Torti et al., 2015).

To overcome this issue, some stakeholders have relied on a tiered filtration step

(prefiltration) to reduce particles and avoid clogging filters (Tsuji et al., 2019), but this approach

increases costs, labor and opportunities for potential contamination (Li et al., 2018; Majaneva et

al., 2018; Robson et al., 2016). The use of filters with bigger pore sizes (up to 20 µm) has been

previously tested and in cases of turbid waters is generally preferred, but requires filtering larger

volumes of water to capture the same amount of DNA recovered in smaller pore size filters

(Robson et al., 2016; Turner et al., 2014).

Freezing water for storage purposes prior to filtration can mitigate the issue of slow

filtration in the field and allow it to be done in batches in the laboratory at a later time, but this

type of sample storage could also introduce bias on eDNA capture and community composition

(Kwambana et al., 2011; Sekar et al., 2009). Freezing can disrupt cells and extrude their DNA in

the environment, which would then make it easier for it to pass through the filter pores and be

lost, an issue that has been demonstrated in certain cases (e.g. Suomalainen et al., 2006).

Increasing the pore size of filters to speed the filtration process could worsen this problem by

letting DNA in solution flow through the pores more easily.

When dealing with turbid waters, some stakeholders have opted to use the centrifugation

approach (e.g. Williams et al., 2017). Extracellular DNA (i.e. DNA not contained within a cell

wall) can be bound to particles (Torti et al., 2015) and consequently be captured and detected

following centrifugation of particles into pellets. However, the amount of water used is limited

by centrifuge size, usually around 15-30 mL per replicate (Doi et al., 2017; Ficetola et al., 2008),

which might limit recovery of diluted DNA (Deiner et al., 2015).

Processing sediment samples may be preferable to processing highly turbid water

samples. However, it is important to understand how DNA recovery from these different media

compare to one another. Turner et al. (2015) and Perkins et al. (2014) have shown that sediment

can have a higher concentration of fish eDNA and some bacteria, respectively. This may relate to

the organic-particle binding and sinking properties, and longer persistence of DNA in sediment

compared to water samples. However, as is the case with water samples, there is no consensus on

the rate of degradation of eDNA in soil and sediment (Dell'Anno & Corinaldesi, 2004; Levy-

Booth et al., 2007; Torti et al., 2015), and this will depend on multiple local biotic and abiotic

factors. In addition, biological communities can naturally differ between water column and

sediments, even when we expect some level of overlap due to both DNA sinking and suspension.

Previous works have compared different approaches to processing eDNA, including

assessment of filtration and storage methods (Hinlo et al., 2017; Takahara et al., 2015),

comparisons between water and sediment eDNA recovery (Sales et al., 2019; Turner et al.,

2015), as well as some work on turbid waters (Kumar et al., 2022; Robson et al., 2016; Williams

et al., 2017). But results have been contradictory, or limited to looking at just DNA

concentration, or at a single targeted species.

The goal of the present study is to compare how freezing water prior to filtration and

using water versus sediment samples induce and/or exacerbate biases in taxa detection for a set

of universal primers targeting different biological communities–12S (fish) and 16S (bacteria and

archaea)—in coastal lagoons. By understanding the biases introduced when processing

environmental samples, we will be able to inform decisions regarding experimental design for

monitoring such a dynamic and challenging habitat, which has invaluable importance for the

maintenance of ecosystem services for both wild and urban populations. We expect these results

will be of interest relative to eDNA sampling in other aquatic systems as well, such as rivers,

streams, and ponds, especially those with turbid waters.

Material and Methods

Site - Topanga Lagoon

To determine the variability of species detection for each protocol, water and sediment samples

were collected from a south-facing coastal lagoon in southern California, located in Malibu, a

stretch of coast that runs from Santa Monica to Point Mugu. This lagoon is part of the Topanga

State Park and is currently undergoing plannings for restoration. It is the only lagoon on this

stretch of coast that still harbors a stable population of tidewater goby (*E. newberryi*), a federally

endangered species, and is relatively less impacted than other lagoons in the same region. The

endangered southern steelhead trout (*O. mykiss*) is also found in this system during anadromy

when the lagoon is breached. Due to the presence of these species, Topanga lagoon has been

periodically surveyed by the Jacobs' lab members and collaborators such as researchers at the

Resource Conservation District of The Santa Monica Mountains (RCDSMM), and therefore its

macrobiota is regularly studied, especially the fish fauna. The lagoon was sampled on September

6th, 2018, at the end of the Summer season, and as is typical of this time of the year, the weather

was dry with no record of precipitation since June (WeatherSpark.com, n.d.). The lagoon was

closed to the ocean by a sandbar and the water was murky (Fig. 1), which in the author's

experience, such turbidity slowed filtration and easily clogged 0.45 µm cellulose nitrate filters.

Protocols and samples

A sterilized water jug was used to collect a single water sample in the lagoon, at a mid-point

between the mouth margin and the road bridge (Fig. 1). The sample was then placed on ice and

brought to the laboratory (~1 hr car ride). This method of "grab-and-hold" has proven to be

similarly effective as on-site filtration in a previous study (Pilliod et al., 2013). Once in the

laboratory, the total volume was divided for two separate protocols: centrifugation; and filtration.

The centrifugation protocol followed Doi et al. (2017) using five replicates of 50 mL

tubes (with 27 mL water samples each). Besides extracting the pellet, we also included a

filtration step of the supernatant using a 0.45 µm filter. For the filtration protocol, water was

separated into ten 500 mL bottles (Fig. 2) and these were used in two separate protocols. Half

(i.e. five bottles) were frozen in the -20°C for three days before thawing for filtration (hereafter

referred to 'pre-freezing (PF) protocol'), and half was not frozen and immediately filtered

(hereafter referred to 'no freezing (NF) protocol'). Filtration was done in two sequential steps for

both treatments (pre- and no freezing) using an adapted vacuum pump in the pre-PCR room of

the laboratory (Fig. S1). First, the water sample was filtered using a 3 µm filter, then the filtrate

was passed through a 0.45 µm filter. All filters used in this work were cellulose nitrate. Here,

however, we will focus only on the results from the first filtration step of the water filtration

protocol (3 µm filters). More details on that are further explained in the supplemental material.

Surficial sediment (within the first 5 cm) was also collected at the same location where

water was sampled (hereafter referred to 'sediment (Sed) protocol'). We used five collection kits,

and each kit consisted of three 2 mL tubes (15 tubes total). Subsamples of these were pooled in

triplicates prior to DNA extraction following instructions as defined by the CALeDNA program

(https://ucedna.com/methods-for-researchers). These were also kept on ice during field work and

stored in a -80°C freezer upon arrival at the laboratory until DNA extractions.

In order to test if freezing water is a viable process to store and manage water samples,

we compared the results of pre-freezing water prior to filtration with the no freezing protocol.

Results from the sediment protocol were compared against both filtration protocols: (pre-

freezing versus no freezing) to test if sediments can be used as an alternative to water samples.

DNA Extraction

DNA from sediments and filters were extracted following the PowerSoil extraction protocol.

Filters were chopped into thin strips before being added to the bead tubes, and sediment

triplicates were pooled in small batches to reach 0.25-0.3 g before processing. We used the soil

extraction kit on the filters as well to reduce potential PCR inhibition caused by the water

turbidity (Kumar et al., 2022), but also to limit the number of variables in the research design by

adding another extraction protocol.

Contamination best practices

Care was taken to avoid contamination both in the field and the lab. Before collection, bottles

and the water jug were cleaned and bleached and then handled with clean gloves on site.

Extractions and PCR were done in a separate pre-PCR room. Utensils and bench top were

cleaned with 10% bleach, followed by 70% ethanol. Forceps and scissors for handling filters

were seared and cleaned with bleach and ethanol after dealing with each sample. PCR reagents

were prepared in a clean, PCR-free, positive pressure hood. Sediment samples were collected

with new 2 mL cryotubes and following field protocol as recommended by the CALeDNA

program. Blanks were made for the field collection, laboratory filtration and PCR (5 blanks in

total) and included in the library for sequencing.

Sequencing

Library preparation followed CALeDNA protocols (https://ucedna.com/methods-for-

researchers). Metabarcode libraries were generated for bacteria and archaea (16S rRNA), fish

(12S rRNA) and metazoans (CO1). Sequences for each primer can be found at Table 1. All

libraries consisted of triplicate PCR reactions. PCR products were visualized using gel

electrophoresis, and for each barcode, PCR triplicates were pooled by sample. After bead

cleaning, all markers were pooled by sample and tagged for sequencing (single indexing).

Libraries were pooled and run on a MiSeq SBS Sequencing v3 in a pair-end 2x300 bp format

[Technology Center for Genomics & Bioinformatics (TCGB), UCLA] with a target sequencing

depth of 25,000 reads/sample/metabarcode. Two sequencing runs were conducted, but the CO1

primer was still below the sequencing depth threshold and therefore its results will not be

discussed here (see Figs. S2-3). For each run, our library was pooled with different samples from

different collaborators to maximize efficiency of the sequencing run.

Bioinformatics and data pre-processing

Sequence data was bioinformatically processed in Hoffman2, the High Performance Computing

cluster at UC Los Angeles, using the Anacapa Toolkit (Curd, Gomer, et al., 2018) with default

settings. Briefly, reads are demultiplexed and trimmed for adapters (cutadapt, Martin, 2013) and

low-quality reads (FastX Toolkit, *FASTX-Toolkit*, n.d.). Dada2 (Callahan et al., 2016) is used to

denoise, dereplicate, merge and remove chimeras, and the resulting clean Amplicon Sequence

Variants (ASVs) have their taxonomy assigned using Bowtie2 (Langmead & Salzberg, 2012),

matched to a custom reference library (CRUX, Curd, Kandlikar, et al., 2018). Confidence levels

are determined by the BLCA algorithm (Gao et al., 2017) to generate a table of best taxonomic

hits, from super-kingdom to species level. The pipeline was designed to process not only paired,

but also unmerged and unpaired reads.

Taxonomic tables with a bootstrap confidence cutoff score of 0.6 were used for

downstream analyses. Except when noted, all bioinformatic analyses mentioned beyond this

point were performed using R v.3.6.2 (R Core Team, 2018) in RStudio v.1.2.1335 (RStudio

Team, 2020). Decontamination was done separately for each primer set and each run (since the

dataset was pooled with different combinations of samples for sequencing). We used the package

metabaR (Zinger et al., 2020) to lower tag-jumping and remove contaminants through detection

of ASVs whose relative abundance is highest in negative controls. We also ran a modification of

the gruinard pipeline (https://github.com/zjgold/gruinard_decon), including only steps 4 (site

occupancy modeling) and 5 (dissimilarity between replicates), since previous steps were

redundant with the metabaR decontamination steps. Lastly, taxa classified as "Not_found",

"Unclassified", "*Canis lupus*", "*Bos taurus*", and "*Homo sapiens*" were removed from the final

tables before being merged and used in downstream analyses.

Diversity analysis

We used the laboratory's own sampling record and the Global Biodiversity Information Facility

database (Gbif.Org, 2022) to manually check the 12S primer final taxonomic table. The number

of species captured by each protocol was visualized using Venn Diagrams (package

VennDiagram, Chen, 2018). Species rarefaction curves were made for each metabarcode to

inspect the level of species saturation in each protocol replicate. The slope of each curve was

calculated using the rareslope function in the vegan package (Oksanen et al., 2019), and the

confidence interval was calculated using pairwiseCI (Schaarschmidt & Gerhard, 2019) with

confidence level at 95%. Rarefaction curves were plotted using the ggrare function from the

ranacapa package (using step = 5).

Differential abundance

The raw dataset was analyzed using DESeq2 and ALDEx2 to look at differential abundance

between protocols (Fernandes et al., 2013; Love et al., 2014), and the output of both analyses

were compared to look for consistency in results. While DESeq2 uses a series of negative-

binomial generalized linear models to count data and estimate the log2-fold change in

abundance, ALDEx2 (ANOVA-like Differential Expression) accounts for community

composition when calculating differential expression by performing a scale invariant

transformation on read counts, which are modeled as distributions of posterior probabilities

sampled from a Dirichlet distribution. This method has been found to produce more consistent

and reproducible results (Nearing et al., 2022).

For DESeq2, the default testing framework was used (test = "Wald", fitType =

"parametric"), which includes the Benjamini-Hochberg multiple inference correction. The

sfType option was defined as poscounts since this estimator is able to handle zeros. The log2

fold change of each pairwise comparison for which there were significant differences in

abundances was plotted. For ALDEx2, we performed a two sample T-test for each pairwise

comparison: between the different filtration protocols (pre- versus no freezing), and between

each filtration protocol and sediment protocol (i.e. PF vs. Sed; and NF vs. Sed). We used 500

Monte Carlo samples (mc.samples = 500) for estimation of the posterior distribution. We plotted

both the between- (M) versus within-condition fold change (W) (MW) and Bland-Altman (BA)

plots using the aldex.plot function.

Beta diversity

For the beta diversity analysis, samples were standardized by using either the eDNA index

(Kelly et al., 2019) or by rarefying them as a way to equalize sequencing effort and minimize

stochasticity and bias. For the eDNA index, we followed the Wisconsin double standardization

method in the vegan package. The custom_rarefaction function in the R package ranacapa

(Kandlikar, 2020) was used to rarefy the dataset with 10 replicates.

For the 12S primer, samples were rarefied to 20 000 reads. Three sediment replicates

were excluded due to very low read numbers (<100). For the 16S, samples were rarefied to

15 000 and one sediment replicate that had ~5000 reads was excluded. The number of reads per

taxa for each protocol replicate was plotted using the phyloseq package (McMurdie & Holmes,

2013), for both the raw and rarefied dataset.

The rarefied dataset followed a Constrained Analysis of Principal Coordinates (CAP)

using the capscale function in vegan and Bray-Curtis distance. This ordination method, which

can be used with non-Euclidean dissimilarity indices, explains the ordination of assemblage

composition based on species abundances. The difference in community composition for each

treatment was then analyzed using a PERMANOVA and Bray-Curtis dissimilarity, followed by a

pairwise PERMANOVA comparison (all with the vegan package). P-values were adjusted using

the FDR (False Discovery Rate) approach.

Results

The first run generated a total of 6 407 371 reads: 3 817 216 reads for the 12S primer, 2 393 627

for 16S, and 196 528 for CO1. In the second run there were a total of 9 088 496 reads: 6 685 673

reads for the 12S metabarcode, 1 904 283 reads for 16S and 498 540 for the CO1. For the 12S

and 16S primers, we were able to reach our threshold of 25 000 reads/sample in most cases,

while that was not the case for all except one sample of the CO1 primer. Because of this

limitation on the number of reads/sample, the CO1 metabarcode will not be discussed further in

the main paper (but check the supplemental material for more details).

The number of reads per sample after decontamination and combining both runs is illustrated in

Figure S3. We manually checked the final taxonomic tables of each separate run for the 12S

primer to look for signs of contamination and evaluate how well the bioinformatic

decontamination steps worked (metabaR and gruinard). The taxonomic tables for the 12S primer

have substantially less species than the 16S, and the local fish fauna is relatively well known,

making the process more tractable.

For the run that was pooled with samples from Palmyra Atoll, the output still retained

some tropical reef and pelagic fish and elasmobranch species that are not found in coastal

lagoons in California. We can expect that tag-jumping contamination is also present in the other

sequencing runs and primers as well. Interestingly, eight out of 28 of those tropical species (ca.

28%) were found exclusively on the sediment protocol and not the filtration protocols (e.g.

Acanthurus achilles, Scarus altipinnis, Lutjanus russellii).

Barplots for both the raw and rarefied dataset (Figs. S3-4, respectively) show that

sediment replicates had greater variability amongst themselves, both in number of reads and

community composition, compared to the replicates of either filtration protocols. Water

replicates were more consistent within and between protocols, and had an overall higher number

of reads than the sediment replicates.

Diversity

After the decontamination steps (metabaR and gruinard) and removing specific, uninformative

ASVs (as listed above), the total number of species assigned to 12S was 39, distributed in 20

orders and 22 families. Of these 39 species, only four had been previously recorded for the site

(Table S1). For 16S, the total number of taxa assigned to species was 2 625, distributed in 45

phyla and 335 families.

We also noticed some dubious taxonomic assignments. For example, for the 12S primer,

we had one hit for *Fundulus diaphanus*, which is a species of killifish native to the northeast of

North America. However, the californian species *F. parvipinnis* has been previously documented

in Topanga by lab members sampling at the site. Similarly, there were two hits for *Phoxinus*

phoxinus, which has a European distribution with a closely related North American counterpart,

P. eos, although this species has not been identified in collections from Topanga lagoon. Another

dubious identification occurred for two species of *Odontesthes*, *O. incisa* and *O. smitti*, which

were among the most abundant hits in our dataset but are native to the southwest Atlantic. These

two species, however, are South American relatives of topsmelt (*Atherinops affinis*), commonly

found in coastal lagoons and estuaries in California (Table S1).

The Venn Diagram (Fig. 3) shows that even though the sediment protocol had lower

numbers of reads overall (Figs. S2-3), they had the highest number of species recovered (12S

primer: N=27, 19 unique; 16S primer: N=1 929, 1 178 unique). The species overlap between

protocols for the 12S was only 1.2% (n=1), and for the 16S primer it was 3.5% (n=402).

318   Species rarefaction curves also show that the sediment protocol is further from reaching

319   saturation compared to both filtration protocols, for both the 12S and 16S primers (Fig. 4),

320   although there was more variation between the replicates for the 12S sediment protocol. For the

321   12S primer, there is a significant difference in the slope of the species curves between the

322   sediment and no freezing protocols (Fig. 5, Table 2), while for the 16S, all pairwise comparisons

323   showed significant differences.

324   Differential Abundance

In the DESeq2 analysis, there was no significant difference between species abundance for the

12S primer in any of the protocols' pairwise comparisons. In ALDEx2, the tidewater goby (*E.*

newberryi) was the only species with an effect size > 1 in the comparisons of filtration protocols

(NF and PF) against the sediment protocol, being underrepresented in the latter (Fig. 6). The

proportion of overlap of the 95% CI of the effect size was not zero, but they were small

nonetheless (NF: 0.010; PF: 0.014) (Table S4).

331   In the case of the 16S primer, the DESeq2 analysis showed no significant difference in

comparison between the filtration protocols (NF versus PF). However, there were significant

differences in the pairwise comparisons between filtration protocols and sediment protocol (NF

vs. Sed; PF vs. Sed; Fig. 6, Tables S2-3). The five most differentially abundant species with

highest abundance in the filtration protocols, relative to the sediment, were representatives of the

families Aphanizomenonaceae, Comamonadaceae and Flavobacteriaceae (in both NF and PF

comparisons); and Hemiselmidaceae and Geminigeraceae (PF protocol only). These comprise

groups of cyanobacteria (Aphanizomenonaceae) and algae (Hemiselmidaceae and

Geminigeraceae), as well as environmental bacteria (Comamonadaceae and Flavobacteriaceae).

The most differentially abundant species with highest abundance in the sediments were

representatives of the families Catenulaceae, Fragilariaceae and an archaea assigned to the

Thaumarchaeota phylum (in both NF and PF comparisons); plus Woeseiaceae and Elphidiidae

(NF protocol only); and Anaerolineaceae and Desulfobacteraceae (PF protocol only). These

comprise groups of diatoms (Catenulaceae and Fragilariaceae), environmental bacteria

(Woeseiaceae, Anaerolineaceae and Desulfobacteraceae) and archaea (Thaumarchaeota), and

foraminiferans (Elphidiidae).

In the ALDEx2 analysis, there were many species that showed an effect size > 1 (Fig. 8),

but only a few showed proportions of overlap < 0.001 (none in the NF versus PF comparison)

(Table S5). For the species that fall within these criteria (effect > 1, overlap < 0.001), in the

comparisons between filtration protocols and sediment protocol (NF vs. Sed; PF vs. Sed), there

were 61 species overrepresented in the NF protocol compared to 39 in the sediment; and 73

species overrepresented in the PF protocol compared to the sediment (there was no species

overrepresented in the sediment that met the criteria above) (Table S5). Taxa overrepresented in

both filtration protocols (NF and PF) relative to sediment are in the families Comamonadaceae,

Flavobacteriaceae and Rhodobacteraceae. Both Comamonadaceae and Flavobacteriaceae were

also represented in the DESeq2 results and comprise groups of environmental bacteria.

Rhodobacteraceae are heterotrophic bacteria usually found in association with algae in marine

environments (Bischoff et al., 2021). In the NF protocol, Cryomorphaceae is the most

represented family, with seven species, and comprises aquatic bacteria generally found in

locations rich in organic carbon (Bowman, 2014).

In the sediments, when compared to the NF protocol, the most overrepresented family is

the Anaerolineaceae with 12 species, followed by Desulfobacteraceae (5), Nonionidae (2) and

Elphidiidae (2). Anaerolineaceae, Desulfobacteraceae and Elphidiidae were also found in the

DESeq2 results. The first two comprise environmental bacteria, while the latter is a family of

foraminiferans (such as the Noniodidae family).

Beta diversity

For the rarefied dataset, there were no discernible differences in assemblage composition in the

CAP analysis for the 12S primer (Fig.9A). One sediment replicate is driving most of the

difference (CAP1=86%) with the overrepresentation of many tropical species, likely tag-jump

contaminants. The PERMANOVA result showed a relatively weak effect overall ($R^2 = 0.38$; p =

0.05), with the pairwise test showing a relatively weak effect for the NF versus PF comparison

($R^2 = 0.20$), followed by stronger effects in the sediment comparisons (NF vs. Sed: $R^2 = 0.41$; PF

vs. Sed: $R^2 = 0.31$) (Table 3). The lack of significant differences between the filtration and

sediment protocols could have been driven by the loss of three sediment replicates when

rarefying the dataset.

When using the eDNA index, the CAP analysis for the 12S primer showed that many of

the species driving the differences in assemblage composition were the tropical species that are

coming from the tag-jumping contamination (Fig. S5A). For example, we see overrepresentation

in the sediment protocol of Stegastes nigricans and Caranx melampygus; and in the NF protocol,

Sphyraena barracuda. Nevertheless, we also see some other species that are known to be found in

the lagoon, such as the Eucyclogobius newberryi, being mostly overrepresented in the filtration

protocols (NF and PF) compared to the sediments; and Gila orcutii, overrepresented in the NF

protocol. Two species of dubious taxonomic assignment are also overrepresented in the

sediment: Phoxinus phoxinus (as discussed in the previous 'Diversity' section); and

Acanthogobius flavimanus, which is a species of goby native to Asia, but that has been recorded

previously in California estuaries (Nico et al., 2022). The PERMANOVA result showed a

weaker effect than in the rarefied dataset (R2 = 0.21; p = 0.055), including for the pairwise

comparisons (Table 4).

For the rarefied 16S primer dataset, the different protocols showed discernible differences

in assemblage composition in the CAP analysis. The first axis explains most of the total variation

(CAP1=96%), with the tidewater goby being the most underrepresented in the sediment

compared to the filtration protocols (NF and PF), especially in the NF protocol (Fig. 9B). The

sediment protocol was also slightly overrepresented by a few other species. One of them was

identified as *Nitrosopelagicus brevis*, a species of uncultured "Candidatus", ammonia-oxidizing

archaea (Thaumarchaeota) found mainly in the epi- and upper mesopelagic environments of the

open oceans (Santoro et al., 2015). There are also two species of *Monomorphina*, (*M. pyrum* and

M. pseudonordstedti) that belong to the Euglenaceae family, a group of eukaryotic flagellates

found in freshwater environments. Lastly, there is *Elphidium williamsoni*, a foraminifera

belonging to the family Elphidiidae found in tidal flats of the North Sea. CAP2 identifies the

variation (14%) differentiating the filtration protocols, with the most distinguishing species being

the *Guillardia theta*, a species of flagellate algae belonging to the family Geminigeraceae,

overrepresented in the PF protocol. Most of these species were also recovered in the DESeq2 and

ALDEx2 analyses, with the exception of both Monomorphina species. The PERMANOVA

result showed a relatively large effect ($R^2 = 0.69$; $p = 0.001$), as well as for all the pairwise

comparisons ($R^2 > 0.5$; Table 3).

The species represented in the rarefied dataset differ from the ones found when using the

eDNA index for the 16S primer. Most of the community assemblage difference in the eDNA

index (CAP1=85%) is driven by differences between filtration and sediment protocols, with six

species being underrepresented in the latter: *Burkholderiales bacterium* TP637, *Curvibacter* sp.

UKPF8, *Diaphorobacter ruginosibacter* and *Verminephrobacter aporrectodeae*

(Comamonadaceae); *beta proteobacterium Mzo1* (Oxalobacteraceae); and *Stella humosa*

(*Peptostreptococcaceae*). Most of these species were also recovered in the DESeq2 and ALDEx2

analyses, with the exception of *V. aporrectodeae*. The PERMANOVA result also showed

relatively high effect ($R^2 = 0.44$; $p = 0.001$), but for the pairwise comparisons the effect between

the filtration protocols (NF vs. PF) was relatively small (R2 = 0.15;Table 4).

Discussion

Standardized protocols to process eDNA are under development (e.g. Bohmann et al., 2021), but

to implement these efficiently it is necessary to compare biases in taxa detection associated with

different protocols. Here, we have explored the detection biases in community composition

introduced by freezing water samples prior to filtration (for storage purposes), and the use of

sediment samples as an alternative to sampling turbid waters. We find that pre-freezing water

does incur some effect on the recovery of community composition, but most strongly for the 16S

primer, while having a relatively small effect for the 12S when using larger, cellulose nitrate

pore size filters (3 μm).The sediment protocol was able to recover eDNA from organisms that

inhabit the water column, however, due to high variability in read abundance among replicates,

we suggest increasing the number of biological replicates and/or the volume sampled in the field.

Tag-jumping contamination

Contamination concerns are usually centered around pre-sequencing, during the field and wet laboratory work. These are of fundamental importance and care should be taken by sterilizing equipment and using negative controls. However, previous literature shows that the sequencing phase can be another source of contamination, generating up to 10% of contaminated reads by tag-jumping (Larsson et al., 2018; Schnell et al., 2015), which can skew analyses of taxa abundance and composition towards the rare taxa. There are ways to help minimize this issue by making use of dual indexing (Kircher et al., 2012)—although see Caroe and Bohmann (2020) for a library approach without dual indexing—, and amplification of positive controls. The latter can be used to track the rate and level of contamination after sequencing to guide read cutoffs on samples (Deiner et al., 2017; Port et al., 2016).

Bioinformatics and data pre-processing

We relied on a bioinformatic approach developed by the metabaR package, adapted from Esling et al. (2015), to reduce the issue of contamination from tag-jumping, since it does not rely solely on the use of positive controls (which we lacked in this analysis) to make the estimated cutoff thresholds. However, after manually checking the fish dataset (12S primer), the final taxonomic tables still contained reads assigned to taxa that are not found in coastal lagoons in California (Table S1). Some of it might be contamination from tag-jumping, although we cannot rule out the possibility that for a few of these species the eDNA could have come from local aquaria, as some are known in the pet trade (e.g. *Acanthurus achilles*). We also cannot disregard the limitations of the reference database, especially related to the absence of estuarine and lagoonal taxa that may lead to dubious assignments to non-local related species (Nagarajan et al., 2022). Due to inability to completely remove potential tag-jump contaminants from the dataset, we can

expect a bias towards the rare taxa that will inflate diversity metrics in our samples for all primer

sets.

The sediment protocol generally showed higher variability among replicates compared to

filtration protocols for both primer sets, both in number of reads and community composition

(Fig. S3-4). The greater consistency of filtration replicates is a result of the single source for the

water samples (the large jug). This means that potentially, multiple biological replicates could be

sampled at the site and pooled for fewer downstream analysis, optimizing cost and labor (Sato et

al., 2017; Dickie et al., 2018), however, since this was not the focus of this work this design

would require further investigation. Sediment replicates were done by individually sampling the

bottom of the lagoon. Although replicates were done a few centimeters apart, the bottom of the

lagoon appears to have small-scale heterogeneity. The spatial variation of soil and sediment

samples is recognized in the literature (Perkins et al., 2014; Taberlet, Prud'Homme, et al., 2012),

and can be caused by sediment composition but also by the flow dynamic and distribution of

eDNA in the water column. While this variability has been shown to occur for water samples as

well in lentic environments (Harper et al., 2019 and references therein), the heterogeneity of

water replicates in this system still requires further investigation.

The sediment protocol also had an overall lower number of reads compared to the

filtration protocols for both primer sets (Fig. S3). The lower number of reads seems to go against

the expectations that eDNA can be more concentrated in sediments (Dell'Anno & Corinaldesi,

2004; Harper et al., 2019; Turner et al., 2015). This could be due to a few issues, some of which

may interact. First, it could be related to a faster degradation and/or turn-over rates of eDNA in

the sediment, which are determined by the soil and eDNA characteristics, as well as enzymatic

and microbial activities (Levy-Booth et al., 2007; Pietramellara et al., 2009; Torti et al., 2015).

The overall lower abundance of eDNA in the sediments could also be driven by increased

inhibition (Buxton et al., 2017; Pawlowski et al., 2022). Even though we used a specific soil

extraction kit for both sediment and filtration protocols, the purification steps in the protocol

could still not have been enough to reduce inhibition in the sediment as well as for the water

samples. Lastly, this could have been driven by the much lower volume of sediment used: 0.25-

0.3 g versus 500 mL for water samples.

There is also the fact that this type of environment is affected by scouring (purging of

sediment to the ocean) during high precipitation events and increased flow of freshwater.

However, since the sediment collection was done out of the rainy season and the lagoon was

closed by a sandbar with no signs of scouring, we are confident that this was not a factor that

could have caused the decreased ability to recover eDNA from the sediments. Therefore, we

expect that this difference in read abundance between sediment and water samples would be

more related to the other factors mentioned above, such as eDNA degradation and turn-over

rates, inhibition, and different processed volumes. Considering both the high variability and the

lower sequencing throughput of the sediment replicates, we advise using a modified sampling

protocol, e.g. the one developed by Taberlet, Prud'Homme, et al. (2012) that includes increasing

the number of replicates and mixing larger volumes before downstream processing.

Diversity
Considering that contamination through tag-jumping could be inflating the numbers of rare

species in the dataset, the steepness and lack of a plateau for many of the species rarefaction

curves could be artificial (Fig. 4). This is especially evident for the 12S primer since we were

able to manually investigate the taxonomy tables (Table S1). However, this lack of a plateau is

an expected outcome from environmental samples (Alberdi et al., 2018), and has been shown to

occur more acutely in a coastal lagoon in California when compared to other environments in

California (Shirazi et al., 2021)—albeit the authors were looking specifically at plants and fungi.

The high number of species recovered from the sediment for the 16S primer (Fig. 3) is likely

driven by the recovery of a rich and complex sediment biota that is not paralleled in the water

column.

The low taxonomic assignment to the species level for some of the dubious fish species

found in our dataset, e.g *Phoxinus phoxinus, Odontesthes* spp. And *Sebastes pachycephalus*, also

highlight the need to expand barcoding efforts to the local estuarine taxa to improve reference

databases. On the other hand, *Fundulus diaphanus*, the northeastern killifish, did receive a few

high taxonomic scores at the species level, which merit further consideration for biomonitoring

of coastal lagoons in the region.

Pre-freezing water prior to filtration had an effect on the species curves of the 16S primer

dataset, but not so much on the 12S. This could be explained by how differently eDNA

molecules are found in the environment for these two different groups of organisms, and how

freezing and thawing water would impact them. In the case of the fish fauna, the DNA that is

shed from the organisms would be either found within cells, or adsorbed to colloids (Liang &

Keeley, 2013; Torti et al., 2015; Turner et al., 2014). Even if cell walls were to disintegrate from

the freezing and thawing process, they could still release intact mitochondria (which range from

1-8 μm in length) that could still be captured by our 3 μm pore size filters. On the other hand,

bacteria and archaea, which are prokaryotic and often single celled organisms, would have their

DNA released directly to the medium and pass through the larger pore size filters (>0.2 μm).

Nevertheless, this freezing effect on cell walls has been shown to not always occur and likely be

species-dependent (Sekar et al., 2009; Suomalainen et al., 2006).

    Differential abundance

Pre-freezing water did introduce bias in species abundance compared to the no freezing protocol,

although the effect was distinctively larger for the 16S than for the 12S primer dataset when

using larger, cellulose nitrate pore size filters (3 μm). Our results seem to align to other reports,

where it was shown that freezing had differential effects on detection and relative abundance of

different prokaryotic taxa (Kwambana et al., 2011; Sekar et al., 2009; Suomalainen et al., 2006).

The smaller effect seen for the 12S dataset could be related to the different qualities and

properties of eDNA in the environment, as mentioned in the previous section ('Diversity').

Larger, mitochondrial molecules from Eukaryotes may be more easily captured in the filters

compared to the smaller celled, prokaryotic organisms targeted by the 16S primer.

Naturally, due to eDNA precipitation and resuspension, we expect to capture some

community overlap between water and surficial sediment samples, however abundances should

be different as consequence of the origin and fate of the eDNA in the environment and the

processes acting on it throughout (Torti et al., 2015). As expected, in the differential abundance

analyses, we see overrepresentation of families of algae (Hemiselmidaceae and Geminigeraceae),

environmental and aquatic bacteria (Comamonadaceae, Flavobacteriaceae, Cryomorphaceae and

Rhodobacteraceae), and cyanobacteria (Aphanizomenonaceae) in the water samples (NF and PF

protocols).There is also significantly higher representation of presumptively benthic diatoms

(Catenulaceae and Fragilariaceae), environmental bacteria (Anaerolineaceae and

Desulfobacteraceae) and foraminiferans (Elphidiidae and Nonionidae) in the sediment. In

addition, the types of environmental bacteria most abundant in the sediments were typical of soil

and sediments elsewhere. Of particular note are those from anoxic environments (e.g.

Anaerolineaceae and Desulfobacteraceae) as lagoon sediments in this region are often dark and

sulfide-rich, consistent with anoxia and sufler-cycling.

In the family Flavobacteriaceae there are important pathogens of fish and humans that

belong to the genus *Flavobacterium*. Suomalainen et al.(2006) found that *F. columnare* was

more susceptible to having its cell walls disrupted to freezing due to high amounts of DNAases,

lyases and proteases, likely connected to its pathogenicity, which then led to lower rates of DNA

recovery. The species found in our dataset was *F. johnsoniae*, a species not known to be

pathogenic–albeit with low species taxonomic score. Given that there was no difference in

abundance for this species in our NF and PF protocols, which is in contrast with the results for

the pathogenic species, *F. columnare*, this might relate to a true non-pathogenic organism.

However, considering that the endangered northern tidewater goby often achieves high

abundance in this lagoon, more detailed assessment of the *Flavobacterium* species inhabiting this

site would be of interest.

The other species assignment that draws our attention is the archea Candidatus

Nitrosopelagicus brevis (Thaumarchaeota), which was shown as more abundant in sediment than

water samples. As mentioned earlier, this is a pelagic species, normally found in the open ocean

worldwide. Although coastal lagoons are subject to marine input, the relatively high

concentration in sediment is unexpected and merits inquiry, especially considering that the

confidence in its taxonomic assignment was low across reads. More likely, this represents a new

environmental archaea that is abundant in coastal lagoon sediments.

Beta diversity
McMurdie and Holmes (2014) recommends against rarefying datasets due to the risk of

removing true, rare ASVs. However, in our case, where we were unable to completely remove

tag-jumping contaminants, this pre-process could help alleviate some of the noise caused by

contaminants. Nevertheless, the CAP and PERMANOVA results on both the rarefied and

standardized (eDNA index) dataset are mostly in agreement, and show greater differences in

assemblage composition for the 16S primer than for the 12S primer, especially between the

filtration and sediment protocols (NF vs. Sed; PF vs. Sed). Although results from the 12S primer

were highly affected by the tag-jump contaminants in the dataset standardized using the eDNA

index (Fig. S5A), the CAP analysis was able to capture the overrepresentation of tidewater goby

in the filtration protocols compared to the sediment, a signal that was lost in the rarefied data

(Fig. 9A).

For the rarefied 16S primer dataset, most species that were over- and underrepresented in

the CAP analysis were also captured by the differential abundance analyses, such as *Guillardia*

theta (Geminigeraceae), which was overrepresented in the PF protocol compared to the NF

protocol. In addition, the species of foraminifera, *Elphidium williamsoni* (Elphidiidae) and the

archea Candidatus *Nitrosopelagicus brevis* (Thaumarchaeota) were found to be overrepresented

in the sediment compared to the filtration protocols. The CAP results on the dataset standardized

using the eDNA index (Fig. S5B) showed different species as underrepresented in the sediment

compared to the filtration protocols, but those also showed up as significantly different in the

differential abundance analyses, with the exception of one, *Verminephrobacter aporrectodeae*.

Interestingly, the CAP analysis was also able to capture the underrepresentation of

tidewater gobies (*E. newberryi*) in the sediment protocol on the 16S primer when compared to

the NF protocol (Fig. 9B). The 16S ribosomal RNA region have been used for fish species

detection, and is not uncommon that different primer sets are taxonomically congruent (Shu,

Lidwig & Peng, 2020; Alberdi et al., 2018). This reinforces the idea discussed earlier

('Bioinformatics and data pre-processing' section) that fish eDNA, at least in this environment,

is less concentrated in the sediment than in the water column, which contradicts other findings

from the literature (Perkins et al., 2014; Turner et al., 2015). But it is worth noting that this

underrepresentation of fish eDNA in the sediment was found to be of small effect for the 12S

primer, though, and there could be some bias related to how these two genes behave and degrade

differently in the environment for the fish fauna.

Lessons Learned

Here is a list of recommendations and best practices for eDNA sampling and analysis in coastal

environments that we have learned throughout this work and believe will be useful for others

working in similar environments with turbid water and highly heterogeneous sediment/soil:

1. Filtered water samples had an overall higher number of reads compared to sediment for

both primer sets. Therefore, we recommend the use of this protocol as it will increase

chances of species detection;

2. If using sediment samples, we recommend increasing the number of replicates and

mixing larger volumes before processing for DNA extractions (as in Taberlet,

Prud'Homme, et al., 2012);

3. Pre-freezing water samples prior to filtration can be an effective storage solution, at least

for Eukaryotes, and when used in combination with cellulose nitrate filters of up to 3 μm

pore size filters;

4. The use of dual-indexing and positive controls during library preparation will help

minimize and address cross-contamination from tag-jumping, as is now widely

recognized in many best-practice protocols (e.g. Deiner et al., 2017; Goldberg et al.,

2016);

5. Although rarefying the dataset is not recommended (McMurdie & Holmes, 2014), we

recognize that it can aid in reducing the noise of contaminants from your dataset, as long

as they are rare. Otherwise, the use of eDNA index (Kelly et al., 2019) can be an

alternative to standardize your dataset.

Conclusions

In this work, we assessed environmental DNA protocols for use in coastal lagoons, a highly

dynamic habitat at the intersection of terrestrial, freshwater and marine environments. Pre-

freezing water combined with the use of larger pore size filters (at least up to 3 μm) can be a

viable alternative for storage and processing of turbid water samples, at least in the case of fish

communities (12S, MiFish). However, the use of sediment samples as an alternative to

processing water samples should be done with caution, and at minimum the number of biological

replicates and/or volume sampled should be increased. Also, while sediment samples were able

to recover eDNA from organisms commonly found in the water column, such as the tidewater

goby, this was achieved during a period of relatively long lagoon closure, when there was no

recent scouring of sediments to the ocean.

While we expect these guidelines to be helpful in the development of strategies to use

eDNA as a monitoring resource in similar environments, protocol testing is still strongly advised

whenever possible, especially when working in a new system. Much work is necessary to

understand the full potential eDNA brings for the conservation and restoration of endangered

species and habitats.

Acknowledgements

Funding was provided by the National Council for Scientific and Technological Development of

Brazil (Rachel Turba) under Grant No. 209261/2014-5 and by NOAA Sea Grant 120651698:1.

Funding for the CALeDNA sample processing, infrastructure, and personnel was provided by the

University of California Research Initiatives (UCRI) Catalyst grant CA-16-376437 and Howard

Hughes Medical Institute (HHMI) Professors Grant GT10483. We are very grateful for all the

help provided by the CALeDNA team, but would like to give special thanks to Teia Schweizer,

who personally trained us in the bench work. Huge thanks to Ryan Kelly, for helping streamline

the design and analysis of this paper and for always being so responsive via email, and to

reviewer David Murray-Stoker for a much detailed and comprehensive feedback that helped

improve this manuscript.

References

Alberdi, A., Aizpurua, O., Gilbert, M. T. P., & Bohmann, K. (2018). Scrutinizing key steps for

reliable metabarcoding of environmental samples. *Methods in Ecology and Evolution*,

9(1), 134–147. https://doi.org/10.1111/2041-210X.12849

Ballard, J., Pezda, J., Spencer, D., & Plantinga, A. (n.d.). *An Economic Valuation of Southern*

California Coastal Wetlands. http://scwrp.org/wp-

content/uploads/2017/06/SoCalWetlands_FinalReport.pdf

Bischoff, V., Zucker, F., & Moraru, C. (2021). Marine Bacteriophages. In D. H. Bamford & M.

Zuckerman (Eds.), *Encyclopedia of Virology (Fourth Edition)* (pp. 322–341). Academic

Press. https://doi.org/10.1016/B978-0-12-809633-8.20988-6

Bohmann, K., Chua, P., Holman, L. E., & Lynggaard, C. (2021). DNAqua☐Net conference

unites participants from around the world with the quest to standardize and implement

DNA☐based aquatic biomonitoring. *Environmental DNA*, *3*(5), 884–888.

https://doi.org/10.1002/edn3.207

Bowman, J. P. (2014). The Family Cryomorphaceae. In E. Rosenberg, E. F. DeLong, S. Lory, E.

Stackebrandt, & F. Thompson (Eds.), *The Prokaryotes* (pp. 539–550). Springer Berlin

Heidelberg. https://doi.org/10.1007/978-3-642-38954-2_135

Buxton, A. S., Groombridge, J. J., & Griffiths, R. A. (2017). Is the detection of aquatic

environmental DNA influenced by substrate type? *PLOS ONE*, *12*(8), e0183371.

https://doi.org/10.1371/journal.pone.0183371

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.

(2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature*

Methods, *13*(7), 581–583. https://doi.org/10.1038/nmeth.3869

Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens,

S. M., Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J. A., Smith, G., & Knight,

R. (2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq

and MiSeq platforms. *The ISME Journal*, *6*(8), 1621–1624.

https://doi.org/10.1038/ismej.2012.8

Caroe, C., & Bohmann, K. (2020). Tagsteady: A metabarcoding library preparation protocol to

avoid false assignment of sequences to samples. *BioRxiv*.

Chen, H. (2018). *VennDiagram: Generate High-Resolution Venn and Euler Plots* (1.6.20).

https://CRAN.R-project.org/package=VennDiagram

Curd, E., Gomer, J., Kandlikar, G., Gold, Z., Ogden, M., & Shi, B. (2018). *The Anacapa Toolkit*.

https://github.com/limey-bean/Anacapa

Curd, E., Kandlikar, G., & Gomer, J. (2018). *CRUX: Creating Reference libraries Using*

eXisting tools. https://github.com/limey-bean/CRUX_Creating-Reference-libraries-

Using-eXisting-tools

Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., Creer,

S., Bista, I., Lodge, D. M., & De Vere, N. (2017). Environmental DNA metabarcoding:

Transforming how we survey animal and plant communities. *Molecular Ecology*, *26*(21),

5872–5895.

Deiner, K., Walser, J.-C., Mächler, E., & Altermatt, F. (2015). Choice of capture and extraction

methods affect detection of freshwater biodiversity from environmental DNA. *Biological*

Conservation, *183*, 53–63.

Dejean, T., Valentini, A., Miquel, C., Taberlet, P., Bellemain, E., & Miaud, C. (2012). Improved

detection of an alien invasive species through environmental DNA barcoding: The

example of the American bullfrog *Lithobates catesbeianus*: *Alien invasive species*

detection using eDNA. Journal of Applied Ecology, *49*(4), 953–959.

https://doi.org/10.1111/j.1365-2664.2012.02171.x

Dell'Anno, A., & Corinaldesi, C. (2004). Degradation and Turnover of Extracellular DNA in

Marine Sediments: Ecological and Methodological Considerations Degradation and

Turnover of Extracellular DNA in Marine Sediments: Ecological and Methodological

Considerations. *Applied and Environmental Microbiology*, *70*(7), 4384–4386.

https://doi.org/10.1128/AEM.70.7.4384

Doi, H., Uchii, K., Matsuhashi, S., Takahara, T., Yamanaka, H., & Minamoto, T. (2017).

Isopropanol precipitation method for collecting fish environmental DNA. *Limnology and*

Oceanography: Methods, *15*(2), 212–218. https://doi.org/10.1002/lom3.10161

Earl, D. A., Louie, K. D., Bardeleben, C., Swift, C. C., & Jacobs, D. K. (2010). Rangewide

microsatellite phylogeography of the endangered tidewater goby, a genetically

subdivided coastal fish with limited marine dispersal. *Conservation Genetics*, *11*, 103–

104. https://doi.org/10.1007/s10592-009-0008-9

Esling, P., Lejzerowicz, F., & Pawlowski, J. (2015). Accurate multiplexing and filtering for high-

throughput amplicon-sequencing. *Nucleic Acids Research*, *43*(5), 2513–2524.

https://doi.org/10.1093/nar/gkv107

FASTX-Toolkit. (n.d.). Retrieved January 11, 2018, from http://hannonlab.cshl.edu/fastx_toolkit/

Fernandes, A. D., Macklaim, J. M., Linn, T. G., Reid, G., & Gloor, G. B. (2013). ANOVA-Like

Differential Expression (ALDEx) Analysis for Mixed Population RNA-Seq. *PLoS ONE*,

8(7), e67019. https://doi.org/10.1371/journal.pone.0067019

Ficetola, G. F., Miaud, C., Pompanon, F., & Taberlet, P. (2008). Species detection using

environmental DNA from water samples. *Biology Letters*, *4*(4), 423–425.

https://doi.org/10.1098/rsbl.2008.0118

Gao, X., Lin, H., Revanna, K., & Dong, Q. (2017). A Bayesian taxonomic classification method

for 16S rRNA gene sequences with improved species-level accuracy. *BMC*

Bioinformatics, *18*(1), 247. https://doi.org/10.1186/s12859-017-1670-4

Gbif.Org. (2022). *Occurrence Download* (p. 170487) [Darwin Core Archive]. The Global

Biodiversity Information Facility. https://doi.org/10.15468/DL.HTJ3HT

Goldberg, C. S., Turner, C. R., Deiner, K., Klymus, K. E., Thomsen, P. F., Murphy, M. A.,

Spear, S. F., McKee, A., Oyler-McCance, S. J., & Cornman, R. S. (2016). Critical

considerations for the application of environmental DNA methods to detect aquatic

species. *Methods in Ecology and Evolution*, *7*(11), 1299–1307.

Harper, L. R., Buxton, A. S., Rees, H. C., Bruce, K., Brys, R., Halfmaerten, D., Read, D. S.,

Watson, H. V., Sayer, C. D., & Jones, E. P. (2019). Prospects and challenges of

environmental DNA (eDNA) monitoring in freshwater ponds. *Hydrobiologia*, *826*(1),

25–41.

Hinlo, R., Gleeson, D., Lintermans, M., & Furlan, E. (2017). Methods to maximise recovery of

environmental DNA from water samples. *PloS One*, *12*(6), e0179251.

Jacobs, D. K., Stein, E. D., & Longcore, T. (2011). Classification of California Estuaries Based

on Natural Closure Patterns: Templates for Restoration and Management Management.

Technical Report, *August*, 1–72.

Kandlikar, G. (2020). *ranacapa: Utility Functions and "shiny" App for Simple Environmental*

DNA Visualizations and Analyses (0.1.0). https://github.com/gauravsk/ranacapa

Kelly, R. P., Shelton, A. O., & Gallego, R. (2019). Understanding PCR Processes to Draw

Meaningful Conclusions from Environmental DNA Studies. *Scientific Reports*, *9*(1),

12133. https://doi.org/10.1038/s41598-019-48546-x

Kircher, M., Sawyer, S., & Meyer, M. (2012). Double indexing overcomes inaccuracies in

multiplex sequencing on the Illumina platform. *Nucleic Acids Research*, *40*(1), e3–e3.

https://doi.org/10.1093/nar/gkr771

Kumar, G., Farrell, E., Reaume, A. M., Eble, J. A., & Gaither, M. R. (2022). One size does not

fit all: Tuning eDNA protocols for high□ and low□turbidity water sampling.

Environmental DNA, *4*(1), 167–180. https://doi.org/10.1002/edn3.235

Kwambana, B. A., Mohammed, N. I., Jeffries, D., Barer, M., Adegbola, R. A., & Antonio, M.

(2011). Differential effects of frozen storage on the molecular detection of bacterial taxa

that inhabit the nasopharynx. *BMC Clinical Pathology*, *11*(1), 2.

https://doi.org/10.1186/1472-6890-11-2

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature*

Methods, *9*(4), 357–359. https://doi.org/10.1038/nmeth.1923

Laramie, M. B., Pilliod, D. S., Goldberg, C. S., & Strickler, K. M. (2015). Environmental DNA

sampling protocol—Filtering water to capture DNA from aquatic organisms. *U.S*

Geological Survey Techniques and Methods, *Book 2*(Chapter A13), 15 p.

https://doi.org/10.3133/TM2A13

Larsson, A. J. M., Stanley, G., Sinha, R., Weissman, I. L., & Sandberg, R. (2018).

Computational correction of index switching in multiplexed sequencing libraries. *Nature*

Methods, *15*(5), 305–307. https://doi.org/10.1038/nmeth.4666

Leray, M., Yang, J. Y., Meyer, C. P., Mills, S. C., Agudelo, N., Ranwez, V., Boehm, J. T., &

Machida, R. J. (2013). A new versatile primer set targeting a short fragment of the

mitochondrial COI region for metabarcoding metazoan diversity: Application for

characterizing coral reef fish gut contents. *Frontiers in Zoology*, *10*(1), 34.

https://doi.org/10.1186/1742-9994-10-34

Levy-Booth, D. J., Campbell, R. G., Gulden, R. H., Hart, M. M., Powell, J. R., Klironomos, J.

N., Pauls, K. P., Swanton, C. J., Trevors, J. T., & Dunfield, K. E. (2007). Cycling of

extracellular DNA in the soil environment. *Soil Biology and Biochemistry*, *39*(12), 2977–

2991. https://doi.org/10.1016/j.soilbio.2007.06.020

Li, J., Lawson Handley, L.-J., Read, D. S., & Hänfling, B. (2018). The effect of filtration method

on the efficiency of environmental DNA capture and quantification via metabarcoding.

Molecular Ecology Resources, *18*(5), 1102–1114.

Liang, Z., & Keeley, A. (2013). Filtration Recovery of Extracellular DNA from Environmental

Water Samples. *Environmental Science & Technology*, *47*(16), 9324–9331.

https://doi.org/10.1021/es401342b

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.

https://doi.org/10.1186/s13059-014-0550-8

Majaneva, M., Diserud, O. H., Eagle, S. H., Boström, E., Hajibabaei, M., & Ekrem, T. (2018).

Environmental DNA filtration techniques affect recovered biodiversity. *Scientific*

Reports, *8*(1), 1–11.

Martin, M. (2013). Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.Journal, *17*(1), 10–12.

McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. *PLoS ONE*, *8*(4), e61217.

https://doi.org/10.1371/journal.pone.0061217

McMurdie, P. J., & Holmes, S. (2014). Waste Not, Want Not: Why Rarefying Microbiome Data

Is Inadmissible. *PLoS Computational Biology*, *10*(4), e1003531.

https://doi.org/10.1371/journal.pcbi.1003531

Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J. Y., Sato, K., Minamoto, T., Yamamoto,

S., Yamanaka, H., Araki, H., Kondoh, M., & Iwasaki, W. (2015). MiFish, a set of

universal PCR primers for metabarcoding environmental DNA from fishes: Detection of

more than 230 subtropical marine species. *Royal Society Open Science*, *2*(7).

http://rsos.royalsocietypublishing.org/content/2/7/150088.abstract

Nagarajan, R. P., Bedwell, M., Holmes, A. E., Sanches, T., Acuña, S., Baerwald, M., Barnes, M.

A., Blankenship, S., Connon, R. E., Deiner, K., Gille, D., Goldberg, C. S., Hunter, M. E.,

Jerde, C. L., Luikart, G., Meyer, R. S., Watts, A., & Schreier, A. (2022). Environmental

DNA Methods for Ecological Monitoring and Biodiversity Assessment in Estuaries.

Estuaries and Coasts, *45*(7), 2254–2273. https://doi.org/10.1007/s12237-022-01080-y

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,

O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H.

(2019). *vegan: Community Ecology Package* (2.5-6). https://CRAN.R-

project.org/package=vegan

Pawlowski, J., Bruce, K., Panksep, K., Aguirre, F. I., Amalfitano, S., Apothéloz-Perret-Gentil,

L., Baussant, T., Bouchez, A., Carugati, L., Cermakova, K., Cordier, T., Corinaldesi, C.,

Costa, F. O., Danovaro, R., Dell'Anno, A., Duarte, S., Eisendle, U., Ferrari, B. J. D.,

Frontalini, F., … Fazi, S. (2022). Environmental DNA metabarcoding for benthic

monitoring: A review of sediment sampling and DNA extraction methods. *Science of The*

Total Environment, *818*, 151783. https://doi.org/10.1016/j.scitotenv.2021.151783

Perkins, T. L., Clements, K., Baas, J. H., Jago, C. F., Jones, D. L., Malham, S. K., & McDonald,

J. E. (2014). Sediment Composition Influences Spatial Variation in the Abundance of

Human Pathogen Indicator Bacteria within an Estuarine Environment. *PLoS ONE*, *9*(11),

e112951. https://doi.org/10.1371/journal.pone.0112951

Pietramellara, G., Ascher, J., Borgogni, F., Ceccherini, M. T., Guerri, G., & Nannipieri, P.

(2009). Extracellular DNA in soil and sediment: Fate and ecological relevance. *Biology*

and Fertility of Soils, *45*(3), 219–235. https://doi.org/10.1007/s00374-008-0345-8

Pilliod, D. S., Goldberg, C. S., Arkle, R. S., Waits, L. P., & Richardson, J. (2013). Estimating

occupancy and abundance of stream amphibians using environmental DNA from filtered

water samples. *Canadian Journal of Fisheries and Aquatic Sciences*, *70*(8), 1123–1130.

https://doi.org/10.1139/cjfas-2013-0047

Port, J. A., O'Donnell, J. L., Romero☐Maraccini, O. C., Leary, P. R., Litvin, S. Y., Nickols, K.

J., Yamahara, K. M., & Kelly, R. P. (2016). Assessing vertebrate biodiversity in a kelp

forest ecosystem using environmental DNA. *Molecular Ecology*, *25*(2), 527–541.

https://doi.org/10.1111/mec.13481

R Core Team. (2018). *R: A language and environment for statistical computing*. R Foundation

for Statistical Computing. https://www.R-project.org/

Rees, H. C., Maddison, B. C., Middleditch, D. J., Patmore, J. R. M., & Gough, K. C. (2014).

REVIEW: The detection of aquatic animal species using environmental DNA – a review

of eDNA as a survey tool in ecology. *Journal of Applied Ecology*, *51*(5), 1450–1459.

https://doi.org/10.1111/1365-2664.12306

Robson, H. L. A., Noble, T. H., Saunders, R. J., Robson, S. K. A., Burrows, D. W., & Jerry, D.

R. (2016). Fine-tuning for the tropics: Application of eDNA technology for invasive fish

detection in tropical freshwater ecosystems. *Molecular Ecology Resources*, *16*(4), 922–

932. https://doi.org/10.1111/1755-0998.12505

RStudio Team. (2020). *RStudio: Integrated Development for R*. RStudio, PBC.

http://www.rstudio.com/

Sales, N. G., Wangensteen, O. S., Carvalho, D. C., & Mariani, S. (2019). Influence of

preservation methods, sample medium and sampling time on eDNA recovery in a

neotropical river. *Environmental DNA*, *1*(2), edn3.14. https://doi.org/10.1002/edn3.14

Santoro, A. E., Dupont, C. L., Richter, R. A., Craig, M. T., Carini, P., McIlvin, M. R., Yang, Y.,

Orsi, W. D., Moran, D. M., & Saito, M. A. (2015). Genomic and proteomic

characterization of "Candidatus Nitrosopelagicus brevis": An ammonia-oxidizing

archaeon from the open ocean. *Proceedings of the National Academy of Sciences*, *112*(4),

1173–1178. https://doi.org/10.1073/pnas.1416223112

Sard, N. M., Herbst, S. J., Nathan, L., Uhrig, G., Kanefsky, J., Robinson, J. D., & Scribner, K. T.

(2019). Comparison of fish detections, community diversity, and relative abundance

using environmental DNA metabarcoding and traditional gears. *Environmental DNA*,

1(4), 368–384. https://doi.org/10.1002/edn3.38

Schaarschmidt, F., & Gerhard, D. (2019). *PairwiseCI: Confidence Intervals for Two Sample*

Comparisons (0.1-27). https://CRAN.R-project.org/package=pairwiseCI

Schnell, I. B., Bohmann, K., & Gilbert, M. T. P. (2015). Tag jumps illuminated—Reducing

sequence-to-sample misidentifications in metabarcoding studies. *Molecular Ecology*

Resources, *15*(6), 1289–1303. https://doi.org/10.1111/1755-0998.12402

SCWRP. (2018). *Wetlands on the Edge: The Future of Southern California's Wetlands:*

Regional Strategy 2018 (p. 142). California State Coastal Conservancy.

scwrp.databasin.org

Sekar, R., Kaczmarsky, L. T., & Richardson, L. L. (2009). Effect of Freezing on PCR

Amplification of 16S rRNA Genes from Microbes Associated with Black Band Disease

of Corals. *Applied and Environmental Microbiology*, *75*(8), 2581–2584.

https://doi.org/10.1128/AEM.01500-08

Shaffer, H. B., Fellers, G. M., Randal Voss, S., Oliver, J. C., & Pauly, G. B. (2004). Species

boundaries, phylogeography and conservation genetics of the red-legged frog (Rana

aurora/draytonii) complex. *Molecular Ecology*, *13*(9), 2667–2677.

https://doi.org/10.1111/j.1365-294X.2004.02285.x

Shirazi, S., Meyer, R. S., & Shapiro, B. (2021). Revisiting the effect of PCR replication and

sequencing depth on biodiversity metrics in environmental DNA metabarcoding. *Ecology*

and Evolution, *11*(22), 15766–15779. https://doi.org/10.1002/ece3.8239

Smart, A. S., Weeks, A. R., Rooyen, A. R., Moore, A., McCarthy, M. A., & Tingley, R. (2016).

Assessing the cost‐efficiency of environmental DNA sampling. *Methods in Ecology and*

Evolution, *7*(11), 1291–1298. https://doi.org/10.1111/2041-210X.12598

Stein, E. D., Cayce, K., Salomon, M., Bram, D. L., De Mello, D., Grossinger, R., & Dark, S.

(2014). *Wetlands of the Southern California Coast: Historical Extent and Change Over*

Time (SFEI Report 720; SCCWRP Technical Report 826; p. 58). Southern California

869   Coastal Water Research Project and San Francisco Estuary Institute.

870   https://www.caltsheets.org/socal/download.html

871 Suomalainen, L.-R., Reunanen, H., Ijäs, R., Valtonen, E. T., & Tiirola, M. (2006). Freezing

872   Induces Biased Results in the Molecular Detection of *Flavobacterium columnare*.

873   *Applied and Environmental Microbiology*, *72*(2), 1702–1704.

874   https://doi.org/10.1128/AEM.72.2.1702-1704.2006

875 Swift, C. C., Haglund, T. R., Ruiz, M., & Fisher, R. N. (1993). The Status and Distribution of the

876   Freshwater Fishes of Southern California. *Bulletin of the Southern California Academy of*

877   *Sciences*, *92*(3), 101–167.

878 Swift, C. C., Spies, B., Ellingson, R. A., & Jacobs, D. K. (2016). A New Species of the Bay

879   Goby Genus Eucyclogobius, Endemic to Southern California: Evolution, Conservation,

880   and Decline. *PloS One*, *11*(7), e0158543. https://doi.org/10.1371/journal.pone.0158543

881 Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C., & Willerslev, E. (2012). Towards next-

882   generation biodiversity assessment using DNA metabarcoding: NEXT-GENERATION

883   DNA METABARCODING. *Molecular Ecology*, *21*(8), 2045–2050.

884   https://doi.org/10.1111/j.1365-294X.2012.05470.x

885 Taberlet, P., Prud'Homme, S. M., Campione, E., Roy, J., Miquel, C., Shehzad, W., Gielly, L.,

886   Rioux, D., Choler, P., Clément, J.-C., Melodelima, C., Pompanon, F., & Coissac, E.

887   (2012). Soil sampling and isolation of extracellular DNA from large amount of starting

888   material suitable for metabarcoding studies: EXTRACTION OF EXTRACELLULAR

889   DNA FROM SOIL. *Molecular Ecology*, *21*(8), 1816–1820.

890   https://doi.org/10.1111/j.1365-294X.2011.05317.x

Takahara, T., Minamoto, T., & Doi, H. (2015). Effects of sample processing on the detection rate

of environmental DNA from the Common Carp (Cyprinus carpio). *Biological*

Conservation, *183*, 64–69. https://doi.org/10.1016/j.biocon.2014.11.014

Thomsen, P. F., & Willerslev, E. (2015). Environmental DNA – An emerging tool in

conservation for monitoring past and present biodiversity. *Biological Conservation*, *183*,

4–18. https://doi.org/10.1016/j.biocon.2014.11.019

Torti, A., Lever, M. A., & Jørgensen, B. B. (2015). Origin, dynamics, and implications of

extracellular DNA pools in marine sediments. *Marine Genomics*, *24*, 185–196.

https://doi.org/10.1016/j.margen.2015.08.007

Tsuji, S., Takahara, T., Doi, H., Shibata, N., & Yamanaka, H. (2019). The detection of aquatic

macroorganisms using environmental DNA analysis—A review of methods for

collection, extraction, and detection. *Environmental DNA*, *1*(2), 99–108.

https://doi.org/10.1002/edn3.21

Turner, C. R., Barnes, M. A., Xu, C. C. Y., Jones, S. E., Jerde, C. L., & Lodge, D. M. (2014).

Particle size distribution and optimal capture of aqueous macrobial eDNA. *Methods in*

Ecology and Evolution, *5*(7), 676–684. https://doi.org/10.1111/2041-210X.12206

Turner, C. R., Uy, K. L., & Everhart, R. C. (2015). Fish environmental DNA is more

concentrated in aquatic sediments than surface water. *Biological Conservation*, *183*, 93–

102. https://doi.org/10.1016/j.biocon.2014.11.017

van der Loos, L. M., & Nijland, R. (2021). Biases in bulk: DNA metabarcoding of marine

communities and the methodology involved. *Molecular Ecology*, *30*(13), 3270–3288.

https://doi.org/10.1111/mec.15592

WeatherSpark.com. (n.d.). *Historical Weather Summer 2018 at Point Mugu Naval Air Warfare*

Center. WeatherSpark.Com. Retrieved June 8, 2022, from

https://weatherspark.com/h/s/145310/2018/1/Historical-Weather-Summer-2018-at-Point-

Mugu-Naval-Air-Warfare-Center;-California;-United-States#Figures-Rainfall

Williams, K. E., Huyvaert, K. P., & Piaggio, A. J. (2017). Clearing muddied waters: Capture of

environmental DNA from turbid waters. *PLOS ONE*, *12*(7), e0179282.

https://doi.org/10.1371/journal.pone.0179282

Zinger, L., Lionnet, C., Benoiston, A.-S., Donald, J., Mercier, C., & Boyer, F. (2020). *metabaR:*

An R package for the evaluation and improvement of DNA metabarcoding data quality

[Preprint]. Bioinformatics. https://doi.org/10.1101/2020.08.28.271817

923
924

925   Tables

926   Table 1: Detailed information of the primers used.

Primer	Targets	Forward Primer	Reverse Primer	Reference
12S	Fish	GTCGGTAAAACTCGTG CCAGC	CATAGTGGGGTATCT AATCCCAGTTTG	Miya et al., 2015
16S	Bacteria and archaea	GTGYCAGCMGCCGCGG TAA	GGACTACNVGGGTWT CTAAT	Caporaso et al., 2012 (F: 515F and R: 806R)
CO1	Animals	GGWACWGGWTGAACW GTWTAYCCYCC	TANACYTCnGGRTGN CCRAARAAYCA	Leray et al., 2013

928   Table 2: Confidence interval estimates and upper/lower limits of species rarefaction curves (Fig.
929   4). NF: no freezing; PF: pre-freezing; Sed: sediment.

Primer	Comparison	Estimate	Lower	Upper
12S	PF-NF	-1.43693	-3.77705	0.903185
	Sed-NF	-6.24818	-11.198	-1.29838
	Sed-PF	-4.81125	-9.76679	0.144296
16S	PF-NF	-0.2004	-0.29947	-0.10132
	Sed-NF	-0.54775	-0.73983	-0.35568
	Sed-PF	-0.34736	-0.54064	-0.15408

930
931

932 Table 3: PERMANOVA and pairwise PERMANOVA (rarefied dataset) between all three
933 protocols: no-freezing, pre-freezing and sediment. P.adjusted is the adjusted p-value after FDR
934 correction.

Primer	PERMANOVA		Pairwise PERMANOVA				
	R2	Pr(>F)	Comparison	F.Model	R2	p.value	p.adjusted
12S	0.3803	0.05	No-freezing vs Pre-freezing	2.07252	0.20576	0.169	0.295
			No-freezing vs Sediment	3.56051	0.41592	0.295	0.295
			Pre-freezing vs Sediment	2.25713	0.31102	0.229	0.295
16S	0.6922	0.001	No-freezing vs Pre-freezing	10.3356	0.56369	0.006	0.009
			No-freezing vs Sediment	12.1022	0.63355	0.007	0.009
			Pre-freezing vs Sediment	12.5474	0.6419	0.009	0.009

936 Table 4: Pairwise PERMANOVA and pairwise PERMANOVA (eDNA index dataset) between
937 all three protocols: no-freezing, pre-freezing and sediment pre- and no freezing water prior to
938 filtration and sediment samples. P.adjusted is the adjusted p-value after FDR correction.

Primer	PERMANOVA		Pairwise PERMANOVA				
	R2	Pr(>F)	Comparison	F.Model	R2	p.value	p.adjusted
12S	0.2113	0.055	No-freezing vs Pre-freezing	1.25387	0.1355	0.297	0.297
			No-freezing vs Sediment	1.76131	0.18044	0.112	0.168
			Pre-freezing vs Sediment	1.69357	0.17471	0.077	0.168
16S	0.4417	0.001	No-freezing vs Pre-freezing	1.47905	0.15603	0.005	0.0135
			No-freezing vs Sediment	5.96537	0.42715	0.017	0.017
			Pre-freezing vs Sediment	6.51459	0.44883	0.009	0.0135

939
940

Figure legends

Figure 1: Photo of Topanga lagoon taken on August 22nd, 2018, a few weeks after collection. There was no record of precipitation for the previous three months and the lagoon was closed to the ocean by a sandbar. There was also no sign of recent waves topping over the sandbar and reaching the lagoon.

Figure 2: Flowchart of the study design. Water was taken as a single sample in a 18 L water jug from the middle of the lagoon (Fig. 1). In the laboratory, the water was split in batches to follow either the centrifugation or the filtration protocol. Sediment samples were taken in 2 μL cryotubes at the same location where water was taken. Subsamples of these were then pooled in triplicates prior to extraction (total 0.25-0.3 g). The dotted box indicates which part of the study design is addressed in the main paper, and which is addressed in the supplemental material. The black 'X' indicates the stage at which DNA extraction was performed and sequenced. Figures were obtained from Biorender.

Figure 3: Venn diagrams of A) 12S and B) 16S primers showing the number of species found at and between each protocol. Sediment samples showed the highest number of unique species for both primers, although for the 12S dataset, about 28% are the result of contamination from tag-jumping.

Figure 4: Species rarefaction curves based on sequencing effort for each protocol. A) 12S primer; B) 16S primer. With the exception of the water samples for the 12S primer, none of the curves have reached a plateau, although we expect the high diversity seen for the 12S sediment samples be due to contamination from tag-jumping.

Figure 5: Confidence interval (CI) for slopes of rarefaction curves (Fig. 3) for each pairwise comparison of the different protocols. Only the comparison between pre- versus no freezing water samples, and pre-freezing versus sediment samples for the 12S primer (A) have come out non significant. The remaining comparisons showed significant differences between rarefaction slopes.

Figure 6: Plots of log2fold change of families of bacteria and archaea (16S primer) for the pairwise comparison between A) no freezing versus sediment; and B) pre-freezing versus sediment. Circles are colored by phylum. Species present above zero are overrepresented in the pre- or no freezing protocol, and species below the zero threshold are overrepresented in the sediments.

Figure 7: MW (left) and BA (right) plots for 12S primer dataset. In both plots, gray represents features that are abundant, but not differentially abundant; black are rare, but not differentially abundant. Diagonal dashed lines are shown for zero-intercept with slope of $\pm$ 1.

Figure 8: MW (left) and BA (right) plots for 16S primer dataset. In both plots, gray represents
features that are abundant, but not differentially abundant; black are rare, but not differentially
abundant. Red represents features called as differentially abundant with q < 0.1. Diagonal dashed
lines are shown for zero-intercept with slope of ± 1.

Figure 9: Constrained Analysis of Principal Coordinates (CAP) of rarefied A) 12S and B) 16S
primer datasets. Circles are colored by protocol. While there were no discernible differences in
assemblage composition for the 12S primer, in the 16S primer *Guillardia theta* is
overrepresented in the PF protocol in comparison to the NF protocol. *Eucyclogobius newberyii* is
underrepresented in the sediments compared to the filtration protocols, especially the NF
protocol.

A) 12S

B) 16S

A) 12S

B) 16S