Single-cell atlas of bronchoalveolar lavage from preschool cystic fibrosis reveals new cell phenotypes

Jovana Maksimovic[1,2,9]*, Shivanthan Shanthikumar[2,3,4]*, George Howitt[1,9], Peter F Hickey[5,6], William Ho[5], Casey Anttila[5], Daniel V. Brown[5], Anne Senabouth[7,8], Dominik Kaczorowski[7,8], Daniela Amann-Zalcenstein[5], Joseph E. Powell[7,8], Sarath C. Ranganathan[2,3,4], Alicia Oshlack[1,9,10], Melanie R. Neeland[2,3,11#].

[1] Computational Biology Program, Peter MacCallum Cancer Centre, Parkville, VIC, Australia
[2] Respiratory Diseases, Murdoch Children's Research Institute, Parkville, VIC, Australia
[3] Department of Paediatrics, University of Melbourne, Parkville, VIC, Australia
[4] Respiratory and Sleep Medicine, Royal Children's Hospital, Parkville, VIC, Australia
[5] Cellular Genomics Projects Team, Advanced Technology and Biology Division, WEHI, Parkville, VIC, Australia
[6] Department of Medical Biology, University of Melbourne, Parkville, VIC, Australia
[7] Garvan-Weizmann Centre for Cellular Genomics, Garvan Institute of Medical Research, Sydney, NSW, Australia
[8] UNSW Cellular Genomics Futures Institute, University of New South Wales, Sydney, NSW, Australia
[9] Sir Peter MacCallum Department of Oncology, University of Melbourne, Parkville, VIC, Australia
[10] School of Mathematics and Statistics, University of Melbourne, Parkville, VIC, Australia
[11] Molecular Immunity, Murdoch Children's Research Institute, Parkville, VIC, Australia
*Contributed equally
#Corresponding author

Corresponding author details:
Dr Melanie Neeland, PhD
Royal Children's Hospital, 50 Flemington Road, Parkville, VIC, Australia
Email: melanie.neeland@mcri.edu.au

Author contributions: JM, SS, SCR, AO and MRN designed the study. SS and SCR recruited the patients and collected samples. SS, CA, DVB, DK and MRN performed experiments and acquired the data. JM, SS, GH, PFH, WH, AS, AO and MRN analysed the data. JM, SS, DA-K, JEP, SCR, AO and MRN interpreted the data. All authors edited, revised, and approved the final manuscript.

Funding: The authors would like to acknowledge funding support from the Chan Zuckerberg Initiative (Inflammation Network), National Health and Medical Research Council of Australia Synergy Grant (1183640), and a Vertex Mentored Research Innovation Award. S.S is supported by a clinician-scientist fellowship from the Melbourne Children's Campus.

Running title: Single-cell atlas of preschool cystic fibrosis BAL

ABSTRACT

Inflammation is a key driver of cystic fibrosis (CF) lung disease, not addressed by current standard care. Improved understanding of the mechanisms leading to aberrant inflammation may assist the development of effective anti-inflammatory therapy. Single-cell RNA sequencing (scRNA-seq) allows profiling of cell composition and function at previously unprecedented resolution. Herein, we seek to use multimodal single-cell analysis to comprehensively define immune cell phenotypes, proportions and functional characteristics in preschool children with CF. We analyzed 42,658 cells from bronchoalveolar lavage of 11 preschool children with CF and a healthy control using scRNA-seq and parallel assessment of 154 cell surface proteins. Validation of cell types identified by scRNA-seq was achieved by assessment of samples by spectral flow cytometry. Analysis of transcriptome expression and cell surface protein expression, combined with functional pathway analysis, revealed 41 immune and epithelial cell populations in BAL. Spectral flow cytometry analysis of over 256,000 cells from a subset of the same patients revealed high correlation in major cell type proportions across the two technologies. Macrophages consisted of 13 functionally distinct sub populations, including previously undescribed populations enriched for markers of vesicle production and regulatory/repair functions. Other novel cell populations included CD4 T cells expressing inflammatory IFNα/β and NFκB signalling genes. Our work provides a comprehensive cellular analysis of the pediatric lower airway in preschool children with CF, reveals novel cell types and provides a reference for investigation of inflammation in early life CF.

INTRODUCTION

Inflammation is a key hallmark of lung disease in cystic fibrosis (CF). A proinflammatory state is present in early life and associated with the development of irreversible structural lung disease (1). Current treatments do not directly target pulmonary inflammation in CF, and the search for effective anti-inflammatory treatment is ongoing. A better understanding of the mechanisms underlying the aberrant inflammation seen in CF would help to develop novel approaches for anti-inflammatory therapy. This could, in turn, improve outcomes for people with CF even in the era of highly effective modulator therapy (2).

Single-cell technologies offer the opportunity to better understand cell composition and function using small volume samples. Such technologies include single-cell RNA sequencing (scRNA-seq), multiomic single-cell sequencing (including cellular indexing of transcriptomes and epitopes by sequencing (CITE-seq)), and spectral flow cytometry (an advance on conventional flow cytometry that allows analysis across the whole fluorescence spectrum). The use of scRNA-seq in CF was recently reviewed by Januska and Walsh (3) who found that the majority of studies had recruited adult participants. The authors highlighted that it was "*crucial to include infant and young children in future investigations*" (3). Further, they reported that most studies had focused on the respiratory epithelium, with only two studies using sample types primarily comprised of immune cells. Schupp *et al*, compared sputa collected from adults with CF and healthy controls (4). They found sputa from those with CF had an increased proportion of recruited monocytes and neutrophils, with a corresponding reduction in alveolar macrophages. The authors also reported functional differences in cell types in people with CF, namely immature and proinflammatory neutrophils as well as monocytes demonstrating increased heat shock responses. Li *et al* compared bronchoalveolar lavage (BAL) from adults with a mild CF lung disease to healthy controls (5). While they described a novel alveolar macrophage "supercluster" based on expression of the genes *IFI27* and *APOC2*, no differences in gene expression were observed between those with CF and controls.

To improve our understanding of pulmonary inflammation in early life CF lung disease, and to address the knowledge gaps identified by Januska and Walsh (3), we aimed to apply novel multimodal single-cell analyses to BAL samples collected from children with CF as part of clinical care in the first 6 years of life. We also included one healthy control for comparison. We applied scRNA-seq, CITE-seq and spectral flow cytometry. The results offer several

94 advancements in our understanding of lung immune cell phenotype and composition in
95 children with CF.

MATERIALS AND METHODS

Study participants

All subjects (n=11 with CF and n=1 healthy control) are enrolled in the AREST CF cohort at the Royal Children's Hospital, Melbourne, Australia. All families gave written and informed consent for their involvement in the AREST CF research program (HREC #25054), which includes collection of samples and clinical data. The healthy control participant had no history of lower airway disease and underwent bronchoscopy to investigate upper airway pathology. Supplementary Table E1 describes the demographics of study participants, and Figure 1A outlines the experimental design.

BAL sample collection and cryopreservation

All subjects underwent clinically indicated bronchoscopy. BAL was performed under general anaesthesia. Each BAL aliquot consisted of 1mL/kg (maximum 20mL) of normal saline being inserted via the working channel of the bronchoscope and then suctioned for return. The BAL samples were sent for routine bacterial culture is a clinical laboratory used to processing samples from children with CF. Viral polymerase chain reaction testing was also performed by the clinical laboratory.

The portion of the BAL samples used for research were placed on ice and processed immediately after the procedure. Samples were centrifuged at 300 x g for 10 min at 4°C. Cell-free BAL supernatant was then collected and stored at -80°C for cytokine analysis. The cell pellet was resuspended in 10mL of media (RPMI supplemented with 2% fetal calf serum (FCS)), filtered through a 70uM filter and centrifuged at 300 x g for 7mins at 4°C. Supernatant was discarded and the cell pellet resuspended in media of for cell counting. The sample was then centrifuged at 300 x g for 10 min and resuspended in equal parts media and chilled freezing solution (FCS with15% dimethyl sulfoxide (DMSO)). The freezing solution was added to resuspended cells drop by drop on ice. The samples were transferred to cryovials, and then cooled at -1°C per minute to -80 °C overnight before being transferred to liquid nitrogen for storage.

Single-cell sequencing and flow cytometry processing

  1. BAL cell thawing and live single cell sorting

Cryopreserved BAL cells were thawed in 10mL media (RPMI supplemented with 10% heat-
inactivated FCS) with 25U/mL benzonase at 37°C and centrifuged at 300xg for 10 min. The
pellet was resuspended in 1mL PBS for cell counting. Following cell count, 9mL PBS was
added to the tube and cells were centrifuged at 300 x g for 10 min. Supernatant was discarded
and the cell pellet resuspended in PBS for viability staining using live/dead fixable near infra-
red viability dye (Invitrogen) according to manufacturers' instructions. The viability dye
reaction was stopped by the addition of FACS buffer (2% heat-inactivated FCS in PBS 2mM
EDTA) and cells were centrifuged at 400 x g for 5 min. Cells were resuspended in FACS
buffer for live, single cell sorting using a BD FACSAria Fusion according to the gating
strategy outlined in Supplementary Figure E1.

2.   <u>Sample and raw data processing for scRNA-seq (participants A1-D1)</u>

Viable cells were sorted on a BD Influx cell sorter (Becton-Dickinson) into PBS + 0.1%
bovine serum albumin and retained on ice. Sorted cells were counted and assessed for
viability with Trypan Blue using a Countess automated counter (Invitrogen), and then
resuspended at a concentration of 800-1000 cells/µL ($8x10^5$ to $1x10^6$ cells/mL). Final cell
viability estimates ranged between 92-96%. Single cell suspensions were loaded onto 10X
Genomics Single Cell 3' Chips along with the reverse transcription (RT) mastermix as per the
manufacturer's protocol for the Chromium Single Cell 3' Library (10X Genomics; PN-
120233), to generate single cell gel beads in emulsion (GEMs). Reverse transcription was
performed using a C1000 Touch Thermal Cycler with a Deep Well Reaction Module (Bio-
Rad) as follows: 55°C for 2h; 85°C for 5min; hold 4°C. cDNA was recovered and purified
with DynaBeads MyOne Silane Beads (Thermo Fisher Scientific; Cat# 37002D) and
SPRIselect beads (Beckman Coulter; Cat# B23318). Purified cDNA was amplified as
follows: 98°C for 3min; 12x (98°C for 15s, 67°C for 20s, 72°C for 60s); 72°C for 60s; hold
4°C. Amplified cDNA was purified using SPRIselect beads and sheared to approximately
200bp with a Covaris S2 instrument (Covaris) using the manufacturer's recommended
parameters. Sequencing libraries were generated with unique sample indices (SI) for each
chromium reaction. Libraries for all samples were multiplexed and sequenced across on
2×150 cycle flow cells on an Illumina NovaSeq 6000 (26bp (Read 1), 8bp (Index), and 98 bp
(Read 2)).
156

The Cell Ranger Single Cell Software Suite (version 6.0.2) by 10x Genomics was used to
process raw sequence data into FASTQ files. First, raw base calls from multiple flow cells
were demultiplexed into separate pools of samples. Reads from each pool were then mapped
to the GRCh38/hg38 genome (version 12) using STAR. The count data was processed and
analyzed in *R* as described below. This dataset will be henceforth referred to as the "scRNA-
seq data".

3. Sample and raw data processing for CITE-seq (participants E2-L2)

Sorted live single cells were centrifuged at 400 x g for 5min at 4°C and resuspended in 25µL
of cell staining buffer (BioLegend). Human TruStain FcX FC blocking reagent (BioLegend)
was added according to manufacturers' instructions for 10min on ice. Each tube was made up
to 100µL with cell staining buffer and TotalSeq Hashtag (HTO 1-8) reagents (BioLegend)
were added to each sample for 20min on ice. Cells were washed with 3mL cell staining
buffer and centrifuged at 400xg for 5min at 4°C. Supernatant was discarded and each sample
resuspended at 62,500 cells/100µL following which 100µL of each sample were pooled into
one tube. Pooled cells were centrifuged at 400xg for 5min at 4°C, supernatant discarded, and
resuspended in 25µL cell staining buffer and 25ul of TotalSeqA Human Universal Cocktail
v1.0 (BioLegend) for 30min on ice. This cocktail contains 154 immune related surface
proteins. Cells were washed in 3mL cell staining buffer and centrifuged at 400xg for 5min at
4°C. Following two more washes, cells were resuspended in PBS + 0.04% BSA for
Chromium captures.

Single-cell captures and library preparations were processed with the 10x Genomics
Chromium single-cell Platform using the 10x Chromium Next GEM single-cell 3' Reagent
V3.1 Dual Index kits (10x Genomics, USA) following the manufacturer's manual. In brief,
pooled cells were counted and an estimated 40,000 cells were loaded per lane onto the 10x
Chromium controller to form single-cell Gel Beads-in-Emulsion (GEMs) in duplicate.
Captured cells were then lysed and barcoded and mRNA molecules were reverse transcribed
to generate cDNA within the single GEMs. The barcoded cDNA was PCR-amplified and
scRNA-seq libraries were constructed using the 10x 3'v3.1 library kits. The TotalSeq-A
scADT-seq and scHTO-seq libraries were constructed as per manufacturer's instructions
(BioLegend). The duplicate scRNA-seq, scADT-seq and scHTO-seq libraries were quantified
by Tapestation 4200 D1000 chip (Agilent). Upon input normalisation, the 6 libraries were

pooled and sequenced on the Illumina NextSeq500 sequencing platform to generate 400

million 2x75-bp paired-end reads.

Reads from each sample were processed using 10x Genomics Cell Ranger software (version

5.0.0). 'cellranger mkfastq' was used to demultiplex the Illumina sequencer's BCL files into

FASTQ files. Next, 'cellranger count' was used to generate single-cell gene-count matrices

against the 10x Genomics pre-built GRCh38 reference genome and transcriptome (2020-A

(July 7, 2020) version). All subsequent analysis was performed in R (version 4.0.3) (R Core

Team. R. A language and environment for statistical computing, 2020) with Bioconductor

(version 3.12) (6). The DropletUtils package (version 1.10.3) (7) was used to create a

SingleCellExperiment object from the Cell Ranger output directories and to identify non-

empty droplets. Samples were demultiplexed based on the hashtag oligo (HTO) counts using

the 'hashedDrops' function from DropletUtils with the default parameters. This was

performed separately for each capture. In parallel, samples were demultiplexed using their

genetic data by running cellsnp-lite (v1.2.0) (8) and Vireo (v0.5.6) (9). Specifically, reads

from non-empty droplets were genotyped at 36.6M SNPs with minor allele frequency (MAF)

>   0.0005   in   the   1000   Genomes   Project

(http://ufpr.dl.sourceforge.net/project/cellsnp/SNPlist/genome1K.phase3.SNP_AF5e4.chr1to

X.hg38.vcf.gz) using cellsnp-lite and then vireo assigned each cell barcode to 1 of 8 donors,

doublets, or unassigned based on these genotypes. This was performed separately for each

capture and then pairs of donors were matched across captures by identifying the best match

between captures based on the genotype profile of each donor. As more cells were

confidently assigned to donors using genetic demultiplexing, the genetic assignments were

used for downstream analysis. Almost all genetic assignments corresponded to a single HTO,

which was then used to link each cell's genetic assignment to a study participant

(Supplementary Table E2). This dataset will be henceforth referred to as the "CITE-seq

data".

4. Flow cytometry protocol and analysis (participants F2-L2)

BAL samples from seven participants had remaining cells for flow cytometry analysis.

Following viability stain as described above, cells were then resuspended in human FC-block

according to manufacturers' instructions for 5 minutes at room temperature. The antibody

cocktail (Supplementary Table E3) made up at 2X concentration was added 1:1 with the cells

and incubated for 30 minutes on ice. Following staining, cells were washed with 2 mL FACS

buffer and centrifuged at 400 x g for 5min. Cells were then resuspended in 2% PFA for a

20min fixation on ice, washed, and resuspended in 150µl FACS buffer for acquisition using a

5L Cytek Aurora. Our protocols for the collection and processing of pediatric BAL for

single-cell analysis are publicly available at https://www.protocols.io/workspaces/earlyAIR.

Flow cytometry results were analysed (manual gating, UMAP) using FlowJo Version 10.8.1

software. Manual gating was performed according to the gating strategy depicted in

Supplementary Figure E1. UMAP analysis was conducted using a concatenated file

containing all live single cells (total 256, 278) from each individual using default parameters

within FlowJo. Mean expression levels of each protein were exported from FlowJo for each

manually gated cell type, following which heatmap plotting and unsupervised hierarchical

clustering was performed using the Morpheus heatmap tool

(https://software.broadinstitute.org/Morpheus).

Single-cell sequencing data analysis

The "scRNA-seq data" was generated from 4 individuals' samples (participants A1-D1),

across 4 captures (1 control sample, 3 CF samples). The "CITE-seq data" was generated from

a further 8 individuals' samples (participants E2-L2) multiplexed across 2 captures (8 CF

samples). Both datasets were similarly processed using the R statistical programming

language (version 4.1.0). As previously described for the "CITE-seq data" in Methods

Section 2 of *Single-cell sequencing and flow cytometry processing*, the DropletUtils package

(version 1.14.1) (10) was used to identify non-empty droplets in the "scRNA-seq data". This

was performed separately for each capture. There were 33,538 non-empty droplets across the

4 samples for the "scRNA-seq data" and 36,601 across the 8 samples for the "CITE-seq

data". To detect within-sample doublets that cannot be identified from the genetic and HTO

demultiplexing data, additional doublets were called for each capture using both scds (version

1.10.0) (10) and scDblFinder (version 1.8.0) (11). scds was run in cxds, bcds and hybrid

mode with the *estDbl* parameter set to TRUE, scDblFinder was run without cluster mode

using the doublet rate calculation as implemented in Demuxafy (12). Quality control was

performed on each sample independently by visually examining the total cell number, the

total unique molecular identifier (UMI) count distributions, the number of unique genes

detected, and the proportions of ribosomal and mitochondrial gene counts per cell. Outlier

droplets in terms of mitochondrial read % were detected based on being more than 3 median

absolute deviations (MADs) from the median value of the metric across all droplets, in each

sample. Droplets that were identified as mitochondrial read % outliers were then removed. DropletQC (13) was used to detect and tag additional empty droplets; there was convincing evidence of additional empty droplets present in the "scRNA-seq data". The filtered droplets from the "scRNA-seq data" and "CITE-seq data" were then combined across 32,732 unique genes, resulting in a total of 54,106 unique droplets. 1,558 "uninformative" genes, such as mitochondrial genes, ribosomal genes, sex chromosome genes and pseudogenes, were then removed. Droplets that were unassigned or heterogenic doublets, along with HTO cross-sample doublets and the additional DropletQC empty droplets originating from the "scRNA-seq data", leaving 45,590 droplets. 19,120 genes were retained for downstream analysis after genes expressed in less than 20 cells were discarded.

The droplets were then automatically annotated with cell type labels from the Human Lung Cell Atlas (HLCA) versions 1 (https://app.azimuth.hubmapconsortium.org/app/human-lung) and 2 (https://app.azimuth.hubmapconsortium.org/app/human-lung-v2) using versions 0.4.1 and 0.4.4 of the Azimuth online application, respectively. The annotated cells were then split into 3 separate data subsets based on their HLCA level 3 labels: "macrophage" (33,161 cells), "T/NK" (6,462 cells) and all "other" cells (5,922 cells), to facilitate confirmation of cell identity and identification of cellular subpopulations. Genes without associated Entrez identifiers were removed at this stage, leaving 16,001 genes for subsequent analyses.

Each data subset ("macrophage", "T/NK" and "other" cells) was independently normalised with SCTransform (14), integrated, scaled, and clustered using Seurat (version 4.0.6) (15-18). Data integration of the 12 samples for each subset of cells was performed using reciprocal principal components analysis (RPCA) with the 4 "scRNA-seq data" samples set as references, 30 dimensions, 3000 features and 20 neighbours for anchor selection. For the "other" cells subset, the number of neighbours to consider when weighting anchors (k.weight) was set to the smallest number of cells in a single sample minus one. The cells in each subset were clustered using the smart local moving (SLM) algorithm with 30 principal components. Ten resolutions between 0.1 and 1 were explored using the clustree (19) package (version 0.4.4); a resolution of 1 was selected for downstream analysis of each data subset. This strategy resulted in 23, 18, and 23 subclusters in the "macrophage", "T/NK" cell, and "other" cell data subsets, respectively. The clusters were visualised using Uniform Manifold Approximation and Projection (UMAP). The quality of each cluster was assessed

by examining their Azimuth prediction score distributions, total UMI count distributions and

the distributions of the total number of unique genes detected.

Marker gene analysis was performed for each cluster, in each of the data subsets, as

previously described in Sim et al. 2021 (20), except that a 1.5-fold change cut-off was used

for the TREAT test. Gene set enrichment analysis of the REACTOME gene sets

(https://www.gsea-msigdb.org/gsea/msigdb/collections.jsp) was performed for each cluster in

each data subset using 'camera' (21). The antibody derived tag (ADT) data associated with

the "CITE-seq data" was normalised using the dsb (22) method (version 1.0.1) following

their   suggested   workflow   (https://cran.r-

project.org/web/packages/dsb/vignettes/end_to_end_workflow.html). The combination of the

top marker genes, REACTOME pathways and ADT expression for each cluster was used to

assign them cellular and/or functional identities. The clusters were then consolidated using

these manually assigned labels. Cells in clusters with no obvious cellular and/or functional

identity and poor-quality control metrics were removed from subsequent analyses. A

"macrophage" subset cluster displaying marker genes for both macrophage and T-cell

lineages was found to be overrepresented for within-sample doublets called by *both* scds and

scDblFinder (Fisher's Exact Test p-value < 0.05). All the cells in this cluster were removed,

along with all other "macrophage" subset doublets called by *both* scds and scDblFinder. The

remaining "macrophage" subset cells were then re-clustered and manually annotated as

previously described. This strategy resulted in 12, 12 and 15 manually annotated clusters in

the "macrophage", "T/NK" and "other" subsets, respectively. Broad cell type labels were also

manually assigned to each cluster, resulting in a total of 13 broad cell type labels across the

three data subsets. Marker genes for the consolidated subpopulations in each data subset were

identified using the 'Cepo' (23) method.

The "macrophage", "T/NK" and "other" cell data subsets were subsequently combined

resulting in a total of 42,658 cells. The data were then normalised, integrated, scaled and

clustered as previously described herein. The subpopulation and broad cell type labels were

visualised using UMAP. Marker genes for the broad cell types were identified using the

'Cepo' method. Cell type proportions estimated by flow cytometry and scRNA-seq were

compared using the 'propeller' approach; the proportions were arcsin transformed and linear

models fitted using limma, taking the paired individual samples into account. The Benjamini-

Hochberg procedure (24) was used to adjust for multiple testing.

All of the code, figures and outputs for the analyses described herein can be viewed at the

following   workflowr   (25)   (version   1.7.0)   analysis   website   on   GitHub:

https://oshlacklab.com/paed-cf-cite-seq/. The code, as well as all necessary inputs and outputs

can be cloned from the GitHub repository associated with the analysis website:

https://github.com/Oshlack/paed-cf-cite-seq.

RESULTS AND DISCUSSION

1. Broad immune and epithelial cell profile of BAL

Unsupervised clustering of single-cell sequencing data and cell annotation using marker gene analysis revealed the following transcriptionally distinct broad cell populations in pediatric BAL: macrophages, proliferating macrophages, monocytes, dendritic cells, neutrophils, B cells, CD4 T cells, CD8 T cells, NK cells, NK-T cells, proliferating NK/T cells, innate lymphocytes, γδ T cells, mast cells, epithelial cells and endothelial cells (Figure 1B). Macrophages were the most abundant cell type identified, followed by CD4 T cells, proliferating macrophages, CD8 T cells, dendritic cells, innate lymphocytes, B cells, monocytes, NK cells and neutrophils. Epithelial cells, NK-T cells, γδ T cells, proliferating NK/T cells, mast cells, and endothelial cells were rarer, and not detected in all individuals (11/12, 11/12, 9/12, 10/12, 6/12, and 2/12 respectively) (Figure 1C).

The five most significant marker genes for each broad cell population are shown in Figure 1D. These marker genes included those described previously for respective cell populations; macrophages (CYP27A1, MS4A4A and PPARG), proliferating macrophages and proliferating NK/T cells expressing cyclic genes (PCLAF and MKI67), B cells (CD19, MS4A1), CD4 T cells (CD3D, CD3E), CD8 T cells (CD8A, CD8B), innate lymphocytes (LEF1, XCL1), γδ T cells (TRGC1, GZMK), NK cells (KLRC1, XCL1), NK-T cells (KLRC1, CD3D), dendritic cells (CLEC10A, CD1E), monocytes (CSF1R, FCGR2B), mast cells (TPSAB1, CPA3), epithelial cells (AGR2, KRT17), and endothelial cells (SPARCL1, ACKR1) (26-30).

Annotation of each cell population was further confirmed by analysis of expression of TotalSeq-A ADTs (Figure 1E). Macrophages expressed myeloid lineage proteins CD172α, CD11c, CD71, HLA-DR, and CD169 as previously reported for alveolar macrophages in adults (31, 32). Monocytes were distinguished from macrophages based on the pan monocyte marker CD14, as well as the lack of CD71 and CD169. Dendritic cells expressed CD11c, CD172α and CD1c; and mast cells expressed known protein markers FcεR1α, CD63, and IgE (33). Cells annotated as lymphoid lineage by marker genes expressed expected lineage proteins, including B cells (CD19$^+$CD20$^+$), CD4 T cells (CD3$^+$CD5$^+$CD4$^+$), CD8 T cells (CD3$^+$CD5$^+$CD8$^+$), γδ T cells (CD3$^+$TCRVδ2$^+$), NK cells (CD3$^-$CD56$^+$) and NK-T cells (CD3$^+$CD56$^+$). Epithelial cells were positive for tissue resident marker CD49b. Endothelial

cells were not identified by ADTs as they were only detected in transcriptomic data from two
participants, neither of whom were included in the CITE-seq experiment (Figure 1C).

2. Comparison of cell proportions by single-cell sequencing and flow cytometry

To investigate how results obtained by single-cell sequencing compared to those obtained by
flow cytometry, we directly compared the 7 BAL samples analysed with both techniques.
Our flow cytometry panel permitted identification of macrophages, monocytes, dendritic
cells, neutrophils, B cells, CD4 T cells, CD8 T cells and airway epithelial cells across
256,000 live, single cells (Figure 2A, Supplementary Figure E1). Proportions of
macrophages, monocytes, B cells, CD4 T cells, CD8 T cells and airway epithelial cells were
generally consistent between the two techniques for each individual (Figure 2B-C). However,
we observed statistically significant differences in the proportion of neutrophils (fold change
flow vs scRNA-seq: 1.2, FDR=0.006) and dendritic cells (fold change flow vs scRNA-seq: -
1.17, FDR=0.019) (Figure 2C). Due to the low RNA content of neutrophils and previously
reported challenges in characterising these cells using single-cell technologies such as 10x
(34), the higher proportion of neutrophils detected by flow cytometry was unsurprising. The
reduced proportion of dendritic cells detected by flow cytometry compared to scRNA-seq
was likely due to the absence of key dendritic cell subset markers CD1c and CD123 in our
flow cytometry panel. Comparable protein expression patterns were observed for common
surface markers CD3, CD8, CD4, CD19, CD16, CD45, HLADR, CD11c, CD14, CD47 and
CD279 across the two techniques (Figure 2D). Of note, our flow cytometry panel also
included several surface proteins not available in the TotalSeq-A cocktail, including CD206
for macrophages, CD15 and CD66b for neutrophil subtyping, and EPCAM for airway
epithelial cells (Figure 2C). Airway macrophages showed auto-fluorescent signatures in flow
cytometry data as we and others have described previously (35, 36).

3. Identification of distinct lung macrophage subpopulations
As macrophages are the most abundant immune cell in BAL and are known to undergo
significant functional development in early life (37), we performed a subclustering analysis
of our broad macrophage population (Figure 3A-B). Subclusters were annotated based on
marker gene analysis (Figure 3C), ADT protein expression (Figure 3D), and with reference to
what has been described in previous scRNA-seq literature. To further confirm annotations
and understand subpopulation functionality based on cell population marker genes, we also
performed REACTOME pathway enrichment analyses for each subtype (38). A list of

enriched REACTOME pathways in each population is provided in Supplementary Table E4,

and a full list of cluster marker genes can be found in extended data file 1.

Our analysis revealed 13 distinct subpopulations of macrophages, of which several have

recently been described in adults. All macrophage subclusters were identified in all

participants including the healthy control (Figure 3B). Subtypes characterised in the adult

lung and also identified here include intermediate macrophages expressing both monocyte

and macrophage genes (macro-int) (27), chemokine-expressing macrophages (macro-CCL)

(27), interstitial macrophages (26), proliferating (or cycling) macrophages (5), macrophages

associated with lipid digestion and transport (NUPR1, RBP4) denoted as macro-lipid (39),

macrophages expressing cholesterol biosynthesis related genes (MSM01, FDFT1) denoted as

macro-cholesterol (5), macrophages expressing IFN-α/β signalling genes (macro-IFN-α/β)

(5), macrophages expressing genes associated with viral response (macro-viral)(5) and

metallothionein-expressing macrophages (macro-MT) (27) (Figure 3B). Novel subtypes

identified were macrophages expressing genes associated with; vesicle production (AZU1,

PLAC8) denoted as macro-vesicle; regulatory function (ATF4, GDF15) denoted as macro-

reg; and response to DNA damage (CDKN1A, MDM2) denoted macro-repair (Figure 3C).

Interestingly, macro-interstitial, macro-CCL, and macro-int cells expressed high protein

levels of CD54 (ICAM1), a transmembrane glycoprotein receptor known to play an important

role in migration of leukocytes to sites of inflammation (40) (Figure 3D).

The most abundant macrophage subcluster in our data, at a median of 16.1% of macrophages,

was a population expressing IFN-α/β signalling genes that we denoted as macro-IFN-α/β.

This cluster was characterised by expression of IFI27 and IGF1 and was enriched for

REACTOME pathways including interferon-alpha-beta signalling. It has previously been

identified in analysis of BAL from adults with CF (5).

The macro-lipid (9.4% of macrophage) subcluster was defined based on enrichment of genes

(NUPR1, RBP4) and REACTOME pathways associated with lipid digestion and transport.

Based on the results of gene set enrichment analysis, this cluster appears to be similar to one

identified previously. Further, RBP4 was identified as a marker gene in both the current and

previous study (39). Similarly, the macro-cholesterol subpopulation appears similar to a

previously reported subcluster in terms of gene set enrichment analysis and a shared marker

gene (MSM01)(5). The macro-viral subcluster was a rare population, which was also

identified in a prior study of BAL from adults from CF where it was referred to as IFN-
reacting (5).

The macro-int population was previously reported in a study of BAL from adults with
COVID-19, where those with more severe disease had an increased proportion of this subtype
(39). In our study this population accounted for 11.2% of the macrophages. In addition to
expressing genes associated with both monocytes and macrophages, the macro-int population
also expressed proteins associated with both populations, including CD11b, CD48 and
intermediate levels of CD14 (Figure 3D). Similarly, the macro-interstitial population also
expressed CD14 and CD11b protein (Figure 3D). CD14 expressing macrophages were also
identified in our flow cytometry data (Supplementary Figure E2A). In general, proportions of
$CD14^+$ macrophages identified by flow cytometry and sc-RNAseq were similar
(Supplementary Figure E2B) although this requires further validation using markers such as
CD11b, CD48, and CD54 to differentiate the macro-int from macro-interstitial using flow
cytometry.

The macro-CCL population in our dataset was enriched for the chemokine-receptor-bind-
chemokines REACTOME pathway, and expressed genes for chemokines CCL4, CCL20,
CCL23, CXCL5, CXCL8, CXCL9 and CXCL10, further confirming its annotation. This
subcluster has been reported in all prior BAL based scRNA-seq studies, however we did not
identify the 4 macro-CCL subclusters that were reported by Li *et al* (5).

The macro-MT population has been previously identified by scRNA-seq evaluation of BAL
samples (5, 41). It was proposed that the gene expression profile of this subpopulation
indicates these cells have been exposed to heavy metals (such as zinc, cadmium, or iron) (41).
If this is correct, the presence of this population in preschool children indicates that this
exposure, and subsequent development of the subpopulation of macrophages, occurs in early
life.

Overall, our data suggest macrophages in the preschool CF lung are highly activated. This is
evidenced by the identified subclusters, but also enrichment for a range of inflammatory
pathways including IL-1 signalling, IFNα/β activation, chemokine signalling, and antiviral
responses. This is consistent with prior studies showing that the upper airway of children
demonstrates a baseline pre-activated state, characterised by upregulation of antiviral and

inflammatory signatures compared to adults (42). Our findings demonstrate that children with

CF may have these pre-activated signatures in the lower airway as well.

4. Neutrophil-like population identified by scRNA-seq

Given the importance of neutrophilic inflammation in CF we attempted to identify

neutrophils in the single-cell data (1). Despite the aforementioned challenges of studying

neutrophils by scRNA-seq, we identified a population of 252 neutrophil-like cells, to which 7

donors contributed more than 10 cells and the remaining 5 fewer than 10 cells. This contrasts

with the only study which has used scRNA-seq to profile BAL from people with CF, which

did not identify a neutrophil population (5). The population of cells we identified mapped to

neutrophils identified by a publicly available and annotated scRNA-seq dataset of the adult

lung (43) and expressed inflammatory genes including those encoding for S100 proteins and

the IL-1 signalling pathway. Protein expression on these cells included CD35, CD55,

CLEC12A, CD48, and a lack of HLA-DR (Figure 1E). CD55 and CD35 are known to be

involved in neutrophil phagocytosis (44). Cells in this neutrophil-like population also shared

gene and protein features with monocytes (genes: FCN1, CD300E; proteins: CD11b and

CD14), which has been described for neutrophils previously (45-48). A more extensive panel

of neutrophil-specific ADTs that are not currently included in the TotalSeq A cocktail could

be used to further characterise this cell type; for example, CD15 and CD66b are protein

markers that we used to distinguish BAL neutrophils from monocytes by flow cytometry

(36).

We show by flow cytometry that BAL neutrophils can be further subtyped based on CD16

and CD66b expression, with CD16$^-$CD66b^{high}, CD16$^+$CD66b^{high} and CD16$^+$CD66b^{low} subsets

identified (Supplementary Figure E1, Figure E2D). This classification has previously been

used to identify Granule-Releasing Immunomodulatory Metabolically active neutrophils

(CD16$^-$CD66B^{high}) which has been implicated in CF related lung inflammation (49). These

cells are characterised by primary granule release and show impaired capacity for bacterial

clearance.

An inherent limitation of our data is that our samples were cryopreserved and thawed prior to

analysis. This process is known to deplete granulocytes; however, we have shown in previous

work that cryopreservation of BAL should not affect the yield of other immune cell

populations (35). As is the case with most clinical studies, our sampling times are

unpredictable in nature and cryopreservation is unavoidable. However, the use of fresh
samples would not have obviated issues with identifying neutrophil populations in scRNA-
seq data, as evidenced by the absence of neutrophils in the prior study of CF BAL which used
fresh samples (5).

5. Inflammatory CD4 T cell subsets in the pediatric lower airway

Our initial gene and protein expression analysis in Figure 1 showed broad populations of
CD4 T cells, CD8 T cells, innate lymphocytes, NK cells, NK-T cells and γδ T cells in BAL
of preschool children with CF. As with macrophages, we next explored heterogeneity within
these T/NK subsets using a subclustering approach (Figure 4). Our analysis revealed both
known and novel subtypes, using marker genes (Figure 4C, extended data file 2), ADT
protein expression (Figure 4D) and REACTOME pathway analysis (Supplementary Table
E5). Previously reported subtypes observed by single-cell RNA sequencing of adult lung
samples that we also observed in our analysis were CD8 Trm, CD8-GZMK, CD4 Treg, γδ T
cells, NK cells, NK-T cells, innate lymphocytes and proliferating NK/T cells (26, 27, 43).

Two subtypes of CD8 T cells (CD8 Trm and CD8-GZMK) and NK-T cells showed gene and
protein markers of lung tissue residency, including surface expression of CD103 and CD49a
(50) (Figure 4C-D). Conversely, CD4 T cell subsets, NK cells, innate lymphocytes, and γδ T
cells were negative or only weakly positive for CD103 and CD49a, suggesting they may have
been recently recruited from the circulation. Both CD8 Trm and CD8-GZMK subtypes
expressed genes encoding for cytotoxic proteases GZMA, GZMB, GZMH, GZMM however
CD8-GZMK was the only CD8 subtype to express GZMK. Recent work has shown that
patterns of cytotoxic molecule expression relate to CD8 T cell differentiation stage, with a
lack of GZM expression observed in naïve CD8 T cells, the majority of memory CD8 T cells
co-expressing GZMA/B/M, and a low number of intermediately-differentiated CD8 T cells
expressing GZMK (51). CD4 T follicular helper (CD4-TFH) cells expressing TFH genes
SCGB3A1 and PD1 signalling (52) were also identified, however only in one individual
(Figure 4B).

To our knowledge, we are the first to describe subtypes of CD4 T cells enriched for IFN and
NFκB signalling genes in the lung. Features of the CD4 T-IFN subtype include marker genes
IFIT1, IFIT3, RSAD2 (Figure 4C) as well as REACTOME enrichment for interferon and
antiviral pathways. Features of the CD4 T-NFκB subtype include marker genes TNFRSF4,

TNFRSF18, NFKB2 (Figure 4C) as well as REACTOME enrichment for several NFκB
activation and signalling pathways (Supplementary Table E5). These inflammatory CD4 T
cells were observed in both the healthy control and children with CF (Figure 4B).

This highlights that these cells may represent a unique characteristic of the early life CF lung
and adds further evidence to the concept that the pediatric CF lower airway is primed for
inflammatory responses.

6. Defining rare cells

A final subclustering analysis of all other cells (not macrophages, not T/NK) revealed further
heterogeneity within the rarer DC, B cell and epithelial cell clusters (Figure 5, Supplementary
Table E6, extended data file 3). DCs were separated into conventional DC (cDC1 and cDC2),
plasmacytoid DC, and migratory DC, based on previously described marker genes (26) and
protein expression. cDC1 expressed CLEC9A gene and CD141 protein, whilst cDC2
expressed CD1c (53-55). Plasmacytoid DC expressed plasmacytoid surface marker CD123,
were negative for conventional markers CD1c and CD141, and marker genes included
CLEC4C and GZMB. Plasma B cells were also identified, characterised by key marker genes
(JCHAIN, MZB1, TNFRSF17) as well as protein expression of plasma markers CD27 and
CD38 (56) (Figure 5C-D).

Whilst BAL predominantly samples immune cells of the lower airway, a small fraction of
airway epithelial cells was also captured. We identified 4 subtypes of airway epithelial cells:
ciliated epithelial cells, basal epithelial cells and 2 clusters containing cells of a goblet/club
phenotype. These were annotated using the Human Lung Cell Atlas (HCLA) (26). The
ciliated epithelial cell cluster expressed marker genes which are exclusively expressed by
ciliated columnar cells of the tracheobronchial tree, and multiciliated epithelial cells in the
HLCA. The basal epithelial cells expressed well-validated marker genes KRT17, KRT5, and
KRT6a. Two distinct populations expressing a mixture of goblet and club cell marker genes
were identified. One cluster, denoted as goblet/club cells, expressed marker genes including
MUC5AC, TFF1, AZGP1, FST, and SYT8 which are expressed by club, goblet and mucus
secreting cells. Another cluster, denoted as tracheobronchial goblet/club cells, expressed the
marker gene TSPAN8 which is a HCLA marker gene for both club and goblet cells from the
lower airway.

CONCLUSIONS

This is the first single-cell analysis of bronchoalveolar lavage from preschool children with CF and includes both RNA and protein. These novel data fill a substantial gap in existing pulmonary scRNA-seq data and in CF lung disease.

Early life is a crucial period of development for both epithelial and immune cells. For example, unlike other organs such as the skin, liver and brain where the transcriptome of resident macrophages from fetal and adult life are very similar, there are substantial differences in the gene expression profiles of fetal and adult lung resident macrophages (57). This highlights childhood as a period where fetal lung resident macrophages must develop before eventually exhibiting an adult phenotype.

Previous scRNA-seq studies involving people with CF have characterised adult samples and primarily focused on respiratory epithelium rather than immune cells. This is also true for the broader pulmonary scRNA-seq field (3). For example, the recent release of the HLCA contained a core dataset derived from samples of 107 individuals, however the youngest of these individuals was 10 years old (26). The HLCA also included an extended dataset, however only 6 of the 338 (1.7%) samples were from individuals 6 years or younger and these samples were not analysed separately to identify specific developmental changes.

Integration of transcriptome-wide data, assessment of highly multiplexed surface proteins, and functional pathway analysis, enabled us to extensively characterise 41 cell populations in the bronchoalveolar lavage of preschool children, predominantly with CF. We revealed several novel cell subtypes, most notably functional macrophage subpopulations and inflammatory CD4 T cells. These novel cell types were found in both children with CF and the one control participant, implying they are unlikely to be disease specific, although this needs to be confirmed in larger studies. Our analysis of paired cell surface protein expression using TotalSeq ADTs and spectral flow cytometry further extends current knowledge of lung cell types and provides within-study validation of our transcriptomic findings.

There are several future directions resulting from this work. First, replication in a larger cohort of children with CF and control participants will allow both validation of our findings as well as investigation of differences in health and disease. In addition, given the variability of the CF lung disease phenotype, it may be possible to identify transcriptomic signatures

568 associated with disease severity. It will also be important to further validate the novel cell
569 populations identified in our study in other cohorts and by different technologies.

570 This study provides the first transcriptional profile of early life lower pulmonary samples and
571 provides a reference dataset for researchers investigating the pulmonary immune system in
572 early life cystic fibrosis lung disease.

 FIGURE LEGENDS

 Figure 1. Broad immune and epithelial cell profile of the pediatric lung. **(A)** Multimodal
 experimental design: 12 cryopreserved bronchoalveolar (BAL) samples from children aged 3-
 6 years were thawed and sorted by FACS for live, single cells. These cells underwent
 scRNA-seq, CITE-seq, or flow cytometry as specified and were analysed to create a cell atlas
 of the pediatric lower airway. **(B)** UMAP visualisation of broad immune and epithelial cell
 populations, coloured by cell type. **(C)** Proportions of each broad population in each
 participant **(D)** Most significant marker genes for each broad population. **(E)** Expression of
 TotalSeq antibody derived tags (ADT) for immune cell lineage markers in each broad cell
 population for 8 samples that were analysed by CITE-seq. Endothelial cells were only
 identified in 2 participants whose samples were analysed by scRNA-seq, so there is no ADT
 data for this cell type.

 Figure 2. Flow cytometry analysis of BAL samples and comparison with scRNA-seq. **(A)**
 UMAP depicting major cell populations identified by spectral flow cytometry of 7 samples:
 B cells, CD4 T cells, CD8 T cells, dendritic cells, macrophages, monocytes, neutrophils and
 airway epithelial cells. **(B)** Cell proportions identified by scRNA-seq and flow cytometry in
 samples from individuals that received both techniques (participants F2-L2). Common cell
 types identified in both datasets are shown. All cell types identified by scRNA-seq are shown
 in Figure 1. **(C)** Statistical comparison of the difference in proportions identified by flow
 cytometry and scRNA-seq. *indicates cell types where a significant difference was observed.
 (D) Cell type specific protein expression by flow cytometry for markers included in the
 panel: CD15, CD66b, CD16, CD11c, CD14, CD45, CD19, CD47, HLA-DR, CD63, CD172α,
 CD206, CD4, CD4, CD8, CD279 (PD1), and EPCAM.

 Figure 3. Identification of functionally distinct macrophage subpopulations. **(A)** UMAP
 visualisation of macrophage subpopulations identified by a subclustering analysis of cells
 within the total macrophage pool. **(B)** Proportions (as a percent of all macrophages) of each
 macrophage subpopulation in each participant. **(C)** Most significant marker genes for each
 macrophage subpopulation. **(D)** Expression of relevant TotalSeq antibody derived tags
 (ADT) in each macrophage subpopulation for 8 samples that were analysed by CITE-seq.

Figure 4. Characterisation of distinct T and NK cell subsets. **(A)** UMAP visualisation of T
and NK cell subsets identified by a subclustering analysis of cells within the 'T/NK' group.
(B) Proportions (as a percent of all T/NK cells) of each T/NK subpopulation in each
participant. **(C)** Most significant marker genes for each identified T and NK cell subset. **(D)**
Expression of TotalSeq antibody derived tags (ADT) for T and NK cell markers in each cell
subset for 8 samples that were analysed by CITE-seq. CD4 TFH cells were only identified in
1 participant whose sample was analysed by scRNA-seq, so there is no ADT data for this cell
type.

Figure 5. Characterisation of rare myeloid, B cell and epithelial cell population. **(A)** UMAP
visualisation of cell subsets identified by a subclustering analysis of cells within the 'not
macrophage, not T/NK' population. **(B)** Proportions (as a percent of all "other" cells) of each
"other" subpopulation in each participant. **(C)** Most significant marker genes for each cell
subset. **(D)** Expression of relevant TotalSeq antibody derived tags (ADT) in each cell subset
for 8 samples that were analysed by CITE-seq. Endothelial cells were only identified in 2
participants whose samples were analysed by scRNA-seq, so there is no ADT data for this
cell type.

ACKNOWLEDGEMENTS

The authors would like to acknowledge the generous support and guidance provided by

Professor Daniel Chambers and Dr Brendan O'Sullivan regarding experimental methods

when designing the first study. We thank the children and parents who participated in the

AREST-CF study, without whom this work would not have been possible. Finally, we thank

the Chan Zuckerberg Initiative Single-cell Biology community for helpful discussions and

support.

REFERENCES

1. Sly PD, Gangell CL, Chen L, Ware RS, Ranganathan S, Mott LS, Murray CP, Stick SM. Risk factors for bronchiectasis in children with cystic fibrosis. *The New England journal of medicine* 2013; 368: 1963-1970.

2. Shanthikumar S, Ranganathan S, Neeland MR. Ivacaftor, not ivacaftor/lumacaftor, associated with lower pulmonary inflammation in preschool cystic fibrosis. *Pediatric pulmonology* 2022; 57: 2549-2552.

3. Januska MN, Walsh MJ. Single-Cell RNA-Sequencing Reveals New Basic and Translational Insights in the Cystic Fibrosis Lung. *American journal of respiratory cell and molecular biology* 2022.

4. Schupp JC, Khanal S, Gomez JL, Sauler M, Adams TS, Chupp GL, Yan X, Poli S, Zhao Y, Montgomery RR, Rosas IO, Dela Cruz CS, Bruscia EM, Egan ME, Kaminski N, Britto CJ. Single-Cell Transcriptional Archetypes of Airway Inflammation in Cystic Fibrosis. *American journal of respiratory and critical care medicine* 2020; 202: 1419-1429.

5. Li X, Kolling FW, Aridgides D, Mellinger D, Ashare A, Jakubzick CV. ScRNA-seq expression of IFI27 and APOC2 identifies four alveolar macrophage superclusters in healthy BALF. *Life science alliance* 2022; 5.

6. Huber W, Carey VJ, Gentleman R, Anders S, Carlson M, Carvalho BS, Bravo HC, Davis S, Gatto L, Girke T, Gottardo R, Hahne F, Hansen KD, Irizarry RA, Lawrence M, Love MI, MacDonald J, Obenchain V, Oleś AK, Pagès H, Reyes A, Shannon P, Smyth GK, Tenenbaum D, Waldron L, Morgan M. Orchestrating high-throughput genomic analysis with Bioconductor. *Nature methods* 2015; 12: 115-121.

7. Lun ATL, Riesenfeld S, Andrews T, Dao TP, Gomes T, participants in the 1st Human Cell Atlas J, Marioni JC. EmptyDrops: distinguishing cells from empty droplets in droplet-based single-cell RNA sequencing data. *Genome biology* 2019; 20: 63-63.

8. Huang X, Huang Y. Cellsnp-lite: an efficient tool for genotyping single cells. *Bioinformatics* 2021.

9. Huang Y, McCarthy DJ, Stegle O. Vireo: Bayesian demultiplexing of pooled single-cell RNA-seq data without genotype reference. *Genome Biol* 2019; 20: 273.

10. Bais AS, Kostka D. scds: computational annotation of doublets in single-cell RNA sequencing data. *Bioinformatics* 2020; 36: 1150-1158.

11. Germain PL, Lun A, Garcia Meixide C, Macnair W, Robinson MD. Doublet identification in single-cell sequencing data using scDblFinder. *F1000Res* 2021; 10: 979.

12. Neavin D, Senabouth A, Hang Lee JT, Ripoll A, Franke L, Prabhakar S, Ye CJ, McCarthy DJ, Melé M, Hemberg M, Powell JE. Demuxafy: Improvement in droplet assignment by integrating multiple single-cell demultiplexing and doublet detection methods. *bioRxiv* 2022: 2022.2003.2007.483367.

13. Muskovic W, Powell JE. DropletQC: improved identification of empty droplets and damaged cells in single-cell RNA-seq data. *Genome Biology* 2021; 22: 329.

14. Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq data using regularized negative binomial regression. *Genome Biology* 2019; 20: 296.

15. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, Lee MJ, Wilk AJ, Darby C, Zager M, Hoffman P, Stoeckius M, Papalexi E, Mimitou EP, Jain J, Srivastava A, Stuart T, Fleming LM, Yeung B, Rogers AJ, McElrath JM, Blish CA, Gottardo R, Smibert P, Satija R. Integrated analysis of multimodal single-cell data. *Cell* 2021; 184: 3573-3587.e3529.

16. Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nat*
Biotechnol 2018; 36: 411-420.
17. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, Hao Y,
Stoeckius M, Smibert P, Satija R. Comprehensive Integration of Single-Cell Data.
Cell 2019; 177: 1888-1902.e1821.
18. Stuart T, Satija R. Integrative single-cell analysis. *Nat Rev Genet* 2019; 20: 257-272.
19. Zappia L, Oshlack A. Clustering trees: a visualization for evaluating clusterings at
multiple resolutions. *GigaScience* 2018; 7.
20. Sim CB, Phipson B, Ziemann M, Rafehi H, Mills RJ, Watt KI, Abu-Bonsrah KD,
Kalathur RKR, Voges HK, Dinh DT, Ter Huurne M, Vivien CJ, Kaspi A,
Kaipananickal H, Hidalgo A, Delbridge LMD, Robker RL, Gregorevic P, Dos
Remedios CG, Lal S, Piers AT, Konstantinov IE, Elliott DA, El-Osta A, Oshlack A,
Hudson JE, Porrello ER. Sex-Specific Control of Human Heart Maturation by the
Progesterone Receptor. *Circulation* 2021.
21. Wu D, Smyth GK. Camera: a competitive gene set test accounting for inter-gene
correlation. *Nucleic acids research* 2012; 40: e133-e133.
22. Mulè MP, Martins AJ, Tsang JS. Normalizing and denoising protein expression data from
droplet-based single cell profiling. *Nat Commun* 2022; 13: 2099.
23. Kim HJ, Wang K, Chen C, Lin Y, Tam PPL, Lin DM, Yang JYH, Yang P. Uncovering
cell identity through differential stability with Cepo. *Nature Computational Science*
2021; 1: 784-790.
24. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate - a Practical and
Powerful Approach to Multiple Testing. *J R Stat Soc B* 1995; 57: 289-300.
25. Blischak JD, Carbonetto P, Stephens M. Creating and sharing reproducible research code
the workflowr way. *F1000Research* 2019; 8: 1749.
26. Sikkema L, Strobl D, Zappia L, Madissoon E, Markov N, Zaragosi L, Ansari M, Arguel
M, Apperloo L, Bécavin C, Berg M, Chichelnitskiy E, Chung M, Collin A, Gay A,
Hooshiar Kashani B, Jain M, Kapellos T, Kole T, Mayr C, von Papen M, Peter L,
Ramírez-Suástegui C, Schniering J, Taylor C, Walzthoeni T, Xu C, Bui L, de Donno
C, Dony L, Guo M, Gutierrez A, Heumos L, Huang N, Ibarra I, Jackson N, Kadur
Lakshminarasimha Murthy P, Lotfollahi M, Tabib T, Talavera-Lopez C, Travaglini
K, Wilbrey-Clark A, Worlock K, Yoshida M, Consortium LBN, Desai T, Eickelberg
O, Falk C, Kaminski N, Krasnow M, Lafyatis R, Nikolíc M, Powell J, Rajagopal J,
Rozenblatt-Rosen O, Seibold M, Sheppard D, Shepherd D, Teichmann S, Tsankov A,
Whitsett J, Xu Y, Banovich N, Barbry P, Duong T, Meyer K, Kropski J, Pe'er D,
Schiller H, Tata P, Schultze J, Misharin A, Nawijn M, Luecken M, Theis F. An
integrated cell atlas of the human lung in health and disease. *bioRxiv* 2022:
2022.2003.2010.483747.
27. Madissoon E, Oliver AJ, Kleshchevnikov V, Wilbrey-Clark A, Polanski K, Orsi AR,
Mamanova L, Bolt L, Richoz N, Elmentaite R, Pett JP, Huang N, He P, Dabrowska
M, Pritchard S, Tuck L, Prigmore E, Knights A, Oszlanczi A, Hunter A, Vieira SF,
Patel M, Georgakopoulos N, Mahbubani K, Saeb-Parsy K, Clatworthy M, Bayraktar
OA, Stegle O, Kumasaka N, Teichmann SA, Meyer KB. A spatial multi-omics atlas
of the human lung reveals a novel immune cell survival niche. *bioRxiv* 2021:
2021.2011.2026.470108.
28. Liao Z, Jin Y, Chu Y, Wu H, Li X, Deng Z, Feng S, Chen N, Luo Z, Zheng X, Bao L, Xu
Y, Tan H, Zhao L. Single-cell transcriptome analysis reveals aberrant stromal cells
and heterogeneous endothelial cells in alcohol-induced osteonecrosis of the femoral
head. *Commun Biol* 2022; 5: 324.

29. Crinier A, Dumas P-Y, Escalière B, Piperoglou C, Gil L, Villacreces A, Vély F, Ivanovic
Z, Milpied P, Narni-Mancinelli É, Vivier É. Single-cell profiling reveals the
trajectories of natural killer cell differentiation in bone marrow and a stress signature
induced by acute myeloid leukemia. *Cellular & molecular immunology* 2021; 18:
1290-1304.
30. Mattiola I, Tomay F, De Pizzol M, Silva-Gomes R, Savino B, Gulic T, Doni A, Lonardi
S, Astrid Boutet M, Nerviani A, Carriero R, Molgora M, Stravalaci M, Morone D,
Shalova IN, Lee Y, Biswas SK, Mantovani G, Sironi M, Pitzalis C, Vermi W,
Bottazzi B, Mantovani A, Locati M. The macrophage tetraspan MS4A4A enhances
dectin-1-dependent NK cell-mediated resistance to metastasis. *Nat Immunol* 2019; 20:
1012-1022.
31. Bissonnette EY, Lauzon-Joset J-F, Debley JS, Ziegler SF. Cross-Talk Between Alveolar
Macrophages and Lung Epithelial Cells is Essential to Maintain Lung Homeostasis.
Frontiers in immunology 2020; 11.
32. Morrell ED, Wiedeman A, Long SA, Gharib SA, West TE, Skerrett SJ, Wurfel MM,
Mikacenic C. Cytometry TOF identifies alveolar macrophage subtypes in acute
respiratory distress syndrome. *JCI Insight* 2018; 3.
33. Kraft S, Jouvin M-H, Kulkarni N, Kissing S, Morgan ES, Dvorak AM, Schröder B, Saftig
P, Kinet J-P. The tetraspanin CD63 is required for efficient IgE-mediated mast cell
degranulation and anaphylaxis. *Journal of immunology (Baltimore, Md : 1950)* 2013;
191: 2871-2878.
34. Grieshaber-Bouyer R, Radtke FA, Cunin P, Stifano G, Levescot A, Vijaykumar B,
Nelson-Maney N, Blaustein RB, Monach PA, Nigrovic PA, Aguilar O, Allan R,
Astarita J, Austen KF, Barrett N, Baysoy A, Benoist C, Brown BD, Buechler M,
Buenrostro J, Casanova MA, Chowdhary K, Colonna M, Crowl T, Deng T, Desland
F, Dhainaut M, Ding J, Dominguez C, Dwyer D, Frascoli M, Gal-Oz S, Goldrath A,
Johanson T, Jordan S, Kang J, Kapoor V, Kenigsberg E, Kim J, Kim Kw, Kiner E,
Kronenberg M, Lanier L, Laplace C, Lareau C, Leader A, Lee J, Magen A, Maier B,
Maslova A, Mathis D, McFarland A, Merad M, Meunier E, Monach PA, Mostafavi S,
Muller S, Muus C, Ner-Gaon H, Nguyen Q, Novakovsky G, Nutt S, Omilusik K,
Ortiz-Lopez A, Paynich M, Peng V, Potempa M, Pradhan R, Quon S, Ramirez R,
Ramanan D, Randolph G, Regev A, Rose SA, Seddu K, Shay T, Shemesh A, Shyer J,
Smilie C, Spidale N, Subramanian A, Sylvia K, Tellier J, Turley S, Vijaykumar B,
Wagers A, Wang C, Wang PL, Wroblewska A, Yang L, Yim A, Yoshida H, ImmGen
C. The neutrotime transcriptional signature defines a single continuum of neutrophils
across biological compartments. *Nat Commun* 2021; 12: 2856.
35. Shanthikumar S, Burton M, Saffery R, Ranganathan SC, Neeland MR. Single Cell Flow
Cytometry Profiling of Bronchoalveolar Lavage in Children. *American journal of*
respiratory cell and molecular biology 2020.
36. Shanthikumar S, Ranganathan SC, Saffery R, Neeland MR. Mapping Pulmonary and
Systemic Inflammation in Preschool Aged Children With Cystic Fibrosis. *Frontiers in*
immunology 2021; 12: 733217.
37. Tan SYS, Krasnow MA. Developmental origin of lung macrophage diversity.
Development 2016; 143: 1318-1327.
38. Gillespie M, Jassal B, Stephan R, Milacic M, Rothfels K, Senff-Ribeiro A, Griss J,
Sevilla C, Matthews L, Gong C, Deng C, Varusai T, Ragueneau E, Haider Y, May B,
Shamovsky V, Weiser J, Brunson T, Sanati N, Beckman L, Shao X, Fabregat A,
Sidiropoulos K, Murillo J, Viteri G, Cook J, Shorser S, Bader G, Demir E, Sander C,
Haw R, Wu G, Stein L, Hermjakob H, D'Eustachio P. The reactome pathway
knowledgebase 2022. *Nucleic acids research* 2021; 50: D687-D692.

39. Liao M, Liu Y, Yuan J, Wen Y, Xu G, Zhao J, Cheng L, Li J, Wang X, Wang F, Liu L,
Amit I, Zhang S, Zhang Z. Single-cell landscape of bronchoalveolar immune cells in
patients with COVID-19. *Nature medicine* 2020; 26: 842-844.
40. Figenschau SL, Knutsen E, Urbarova I, Fenton C, Elston B, Perander M, Mortensen ES,
Fenton KA. ICAM1 expression is induced by proinflammatory cytokines and
associated with TLS formation in aggressive breast cancer subtypes. *Scientific reports*
2018; 8: 11720.
41. Mould KJ, Moore CM, McManus SA, McCubbrey AL, McClendon JD, Griesmer CL,
Henson PM, Janssen WJ. Airspace Macrophages and Monocytes Exist in
Transcriptionally Distinct Subsets in Healthy Adults. *American journal of respiratory*
and critical care medicine 2021; 203: 946-956.
42. Loske J, Röhmel J, Lukassen S, Stricker S, Magalhães VG, Liebig J, Chua RL, Thürmann
L, Messingschlager M, Seegebarth A, Timmermann B, Klages S, Ralser M, Sawitzki
B, Sander LE, Corman VM, Conrad C, Laudi S, Binder M, Trump S, Eils R, Mall
MA, Lehmann I. Pre-activated antiviral innate immunity in the upper airways controls
early SARS-CoV-2 infection in children. *Nat Biotechnol* 2022; 40: 319-324.
43. Zilionis R, Engblom C, Pfirschke C, Savova V, Zemmour D, Saatcioglu HD, Krishnan I,
Maroni G, Meyerovitz CV, Kerwin CM, Choi S, Richards WG, De Rienzo A, Tenen
DG, Bueno R, Levantini E, Pittet MJ, Klein AM. Single-Cell Transcriptomics of
Human and Mouse Lung Cancers Reveals Conserved Myeloid Populations across
Individuals and Species. *Immunity* 2019; 50: 1317-1334.e1310.
44. Vandendriessche S, Cambier S, Proost P, Marques PE. Complement Receptors and Their
Role in Leukocyte Recruitment and Phagocytosis. *Frontiers in Cell and*
Developmental Biology 2021; 9.
45. Abdel-Salam BK, Ebaid H. Expression of CD11b and CD18 on polymorphonuclear
neutrophils stimulated with interleukin-2. *Cent Eur J Immunol* 2014; 39: 209-215.
46. Haziot A, Tsuberi BZ, Goyert SM. Neutrophil CD14: biochemical properties and role in
the secretion of tumor necrosis factor-alpha in response to lipopolysaccharide. *J*
Immunol 1993; 150: 5556-5565.
47. Silva-Gomes R, Mapelli SN, Boutet MA, Mattiola I, Sironi M, Grizzi F, Colombo F,
Supino D, Carnevale S, Pasqualini F, Stravalaci M, Porte R, Gianatti A, Pitzalis C,
Locati M, Oliveira MJ, Bottazzi B, Mantovani A. Differential expression and
regulation of MS4A family members in myeloid cells in physiological and
pathological conditions. *J Leukoc Biol* 2022; 111: 817-836.
48. Munthe-Fog L, Hummelshoj T, Honoré C, Moller ME, Skjoedt MO, Palsgaard I,
Borregaard N, Madsen HO, Garred P. Variation in FCN1 affects biosynthesis of
ficolin-1 and is associated with outcome of systemic inflammation. *Genes &*
Immunity 2012; 13: 515-522.
49. Margaroli C, Moncada-Giraldo D, Gulick DA, Dobosh B, Giacalone VD, Forrest OA,
Sun F, Gu C, Gaggar A, Kissick H, Wu R, Gibson G, Tirouvanziam R.
Transcriptional firing represses bactericidal activity in cystic fibrosis airway
neutrophils. *Cell reports Medicine* 2021; 2: 100239.
50. Zheng MZM, Wakim LM. Tissue resident memory T cells in the respiratory tract.
Mucosal Immunol 2021.
51. Bengsch B, Ohtani T, Herati RS, Bovenschen N, Chang K-M, Wherry EJ. Deep immune
profiling by mass cytometry links human T and NK cell differentiation and cytotoxic
molecule expression patterns. *J Immunol Methods* 2018; 453: 3-10.
52. Weinstein JS, Lezon-Geyda K, Maksimova Y, Craft S, Zhang Y, Su M, Schulz VP, Craft
J, Gallagher PG. Global transcriptome analysis and enhancer landscape of human
primary T follicular helper and T effector lymphocytes. *Blood* 2014; 124: 3719-3729.

53. Minoda Y, Virshup I, Leal Rojas I, Haigh O, Wong Y, Miles JJ, Wells CA, Radford KJ.
Human CD141+ Dendritic Cell and CD1c+ Dendritic Cell Undergo Concordant Early
Genetic Programming after Activation in Humanized Mice In Vivo. *Frontiers in*
Immunology 2017; 8.
54. Tullett KM, Lahoud MH, Radford KJ. Harnessing Human Cross-Presenting
CLEC9A+XCR1+ Dendritic Cells for Immunotherapy. *Frontiers in immunology*
2014; 5.
55. Collin M, McGovern N, Haniffa M. Human dendritic cell subsets. *Immunology* 2013;
140: 22-30.
56. da Silva FAR, Pascoal LB, Dotti I, Setsuko Ayrizono MdL, Aguilar D, Rodrigues BL,
Arroyes M, Ferrer-Picon E, Milanski M, Velloso LA, Fagundes JJ, Salas A, Leal RF.
Whole transcriptional analysis identifies markers of B, T and plasma cell signaling
pathways in the mesenteric adipose tissue associated with Crohn's disease. *J Transl*
Med 2020; 18: 44.
57. Bian Z, Gong Y, Huang T, Lee CZW, Bian L, Bai Z, Shi H, Zeng Y, Liu C, He J, Zhou J,
Li X, Li Z, Ni Y, Ma C, Cui L, Zhang R, Chan JKY, Ng LG, Lan Y, Ginhoux F, Liu
B. Deciphering human macrophage development at single-cell resolution. *Nature*
2020; 582: 571-576.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5