Landscape drives zoonotic malaria prevalence in non-human primates

Emilia Johnson[1,2,3] *, Reuben Sunil Kumar Sharma[4], Pablo Ruiz Cuenca[2,5,6], Isabel Byrne[2], Milena Salgado-Lynn[7,8,9], Zarith Suraya Shahar[4], Lee Col Lin[4], Norhadila Zulkifli[4], Nor Dilaila Mohd Saidi[4], Chris Drakeley[10], Jason Matthiopoulos[1], Luca Nelli[1], Kimberly Fornace[1,2,3,11]

[1] Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, F12 8QQ, Scotland

[2] Department of Disease Control, London School of Hygiene & Tropical Medicine, London WC1E 7HT, UK

[3] Centre on Climate Change and Planetary Health, London School of Hygiene & Tropical Medicine, London WC1E 7HT, UK

[4] Faculty of Veterinary Medicine, Universiti Putra Malaysia, 43400, UPM, Serdang, Selangor, Malaysia

[5] Lancaster University, Bailrigg, Lancaster LA1 4YW

[6] Liverpool School of Tropical Medicine, Pembroke Place Liverpool, L3 5QA, UK

[7] School of Biosciences, Cardiff University, Cardiff CF10 3AX, UK

[8] Wildlife Health, Genetic and Forensic Laboratory, Sabah Wildlife Department, Wisma Muis, 88100, Kota Kinabalu, Malaysia

[9] Danau Girang Field Centre, Sabah Wildlife Department, Wisma MUIS 88100, Kota Kinabalu Sabah, Malaysia

[10] Department of Infection Biology, London School of Hygiene & Tropical Medicine, London WC1E 7HT, UK

[11] Saw Swee Hock School of Public Health, National University of Singapore, Singapore 117549

*corresponding author: emilia.johnson@glasgow.ac.uk

ABSTRACT

In wildlife and humans, parasite infection has been linked to land conversion and fragmented

habitat. However, zoonotic disease dynamics in wildlife hosts are rarely quantified at

macroecological scales due to the lack of systematic surveys. Non-human primates (NHPs)

host *Plasmodium knowlesi,* a zoonotic malaria with increasing public health impact and the

main barrier to malaria elimination in Southeast Asia. Understanding of regional *P. knowlesi*

infection dynamics in wildlife is limited. Here, we systematically assemble reports of NHP *P.*

knowlesi and investigate how landscape impacts prevalence. Meta-analysis of 6322 NHPs

from 148 sites highlights that prevalence is heterogeneous across Southeast Asia, with low

overall prevalence and high estimates for Malaysian Borneo. Broadly, we find that regions

exhibiting higher prevalence in NHPs overlap with human infection hotspots. By assembling

remote sensing data and fitting generalised linear mixed models to prevalence at multiple

spatial scales, we identify novel relationships between *P. knowlesi* in NHPs and forested

landscapes. Preliminary evidence suggests that higher prevalence may be contingent on

habitat complexity, which might explain geographic variation in parasite burden. Findings

address critical gaps in understanding *P. knowlesi* epidemiology and indicate that

prevalence in simian reservoirs may be a key spatial driver of human spillover risk.

Keywords

INTRODUCTION

Zoonotic infectious diseases arise from the spillover of pathogens into human populations, typically from a reservoir in wildlife hosts. Anthropogenic land use and land cover changes (LULCC) have now been widely linked to infectious disease outbreaks(Brock et al., 2019; Davidson et al., 2019; Loh et al., 2016). Such practices, including deforestation, logging, clearing for cash-crop plantations or conversion of intact forest into arable land, are accelerating across tropical forests of Southeast Asia(Fornace et al., 2021; Imai et al., 2018). Mechanisms that underly the association between habitat disturbance and spillover risk from wildlife hosts are complex and occur over multiple spatial scales(Brock et al., 2019). In Brazil, re-emergence of Yellow Fever Virus in both simians and humans has been linked to areas with highly fragmented forest(Ilacqua et al., 2021). In part, an increase in forest fringes or 'edge' habitat in fragmented landscapes can facilitate spatial overlap and altered contact patterns between wildlife, vectors and humans(Lehman et al., 2006). Such ecological interfaces are also thought to contribute to parasite spillover in other vector-borne diseases including Zika(J. Li et al., 2021), Babesiosis and Lyme disease(Simon et al., 2014), *Trypanosoma cruzi*(Vaz et al., 2007) and zoonotic malaria(Brock et al., 2019; Grigg et al., 2017). At the same time, habitat complexity can have detrimental impact on wildlife population viability, with reduced host species occupancy and reduced disease burden in highly disturbed habitats(Hanski and Ovaskainen, 2000). Disentangling this interplay is essential to inform ecological strategies for surveillance and mitigation of diseases in regions undergoing landscape change(Fornace et al., 2021).

Zoonotic *P. knowlesi* is a public health threat of increasing importance across Southeast Asia, following the identification of a prominent infection foci in Borneo in 2004(Singh et al., 2004). *P. knowlesi* is a zoonosis, with a sylvatic cycle circulating in non-human primates (NHPs). Human cases are thought to occur only from spillover events(Cuenca et al., 2022; Fornace, 2022; Lee et al., 2011). Human transmission requires bites from infective mosquitos, primarily anopheline mosquitos of the Leucosphyrus Complex (*Anopheles balabacensis, An. latens, An. introlactus)* and Dirus Complex (*An. dirus, An. cracens)*(Moyes et al., 2016; Vythilingam et al., 2006; Wong et al., 2015)*.* Natural hosts for *P. knowlesi* are typically Long-tailed macaques (*Macaca fascicularis*) and Southern Pig-tailed macaques (*M. nemestrina)*(Moyes et al., 2016), both occurring widely across Southeast Asia. Currently, distribution of *P. knowlesi* cases is thought to be restricted to the predicted ranges of known vector and host species(Speldewinde et al., 2019), though recent studies have also identified other NHPs found to be harbouring *P. knowlesi.* This includes Stump-tailed macaques *(M. arctoides),* which are now considered to be another natural reservoir(Fungfuang et al., 2020).

Progress towards malaria elimination in Malaysia has been stymied by a recent rise in
human incidence of *P. knowlesi* malaria. Accounting for increases in surveillance and
diagnostic improvements it is now recognised as the most common cause of clinical malaria
in Malaysia(Cooper et al., 2020). Indeed, Malaysia was the first country not to qualify for
malaria elimination due to ongoing presence of zoonotic malaria and the World Health
Organisation (WHO) updated the guidelines to reflect zoonotic malaria as a public health
threat ("Global Malaria Programme," n.d.). Emergence of *Plasmodium knowlesi* infections
has been linked to LULCC(Fornace et al., 2021). While sporadic cases have been reported
across Southeast Asia, including in Indonesia(Setiadi et al., 2016), the Philippines(Fornace
et al., 2018), Vietnam(Maeno et al., 2015), Brunei(Koh et al., 2019) and Myanmar(Ghinai et
al., 2017), the majority of *P. knowlesi* cases are found in East Malaysia (Borneo) with
hotspots in the states of Sabah and Sarawak(Jeyaprakasam et al., 2020), areas that have
seen extensive deforestation and landscape modification. In Sabah, human prevalence of *P.*
knowlesi infection has recently been shown to be specifically associated with recent loss of
intact forest, agricultural activities and fragmentation across multiple localised spatial
scales(Brock et al., 2019; Fornace et al., 2019b, 2016)*.*

Prevalence of the pathogen in reservoir hosts is considered to be one of three crucial factors
determining the force of infection in zoonotic spillover events(Murray and Daszak, 2013).
Despite this, very little is known of the impact of rapid LULCC on the distribution of *P.*
knowlesi in NHPs. Literature on the impacts of fragmentation on primates tends to focus on
primate density and abundance(Link et al., 2010; ZUNINO et al., 2007). What is known is
that effects of LULCC on primate-pathogen dynamics are highly variable and context-
specific. Although the vector species responsible for sylvatic transmission remain unknown,
the *Anopheles leucospryphus* group, the only vector group implicated in *P. knowlesi*
transmission, is widely associated with secondary, disturbed forest(Brant, 2011; Hawkes et
al., 2019; Wong et al., 2015). Macaques have been known to preferentially rely on fringe
habitat, behaviour which may be exaggerated in response to forest fragmentation and
facilitate exposure to vectors(Lehman et al., 2006; Stark et al., 2019). Changes to land
composition can also create the biosocial conditions for higher rates of parasitism in
primates. Under conditions of limited resources and reduction in viable habitat, conspecific
primate density may increase as troops compete for available space. In turn, this can favour
transmission via intra-species contact or allow the exchange of pathogens between troops
dwelling in intact forest versus edge habitat(Faust et al., 2018; Stark et al., 2019). Habitat
use may also become more intensive, preventing parasite avoidance behaviours(Nunn and
Dokey, 2006). Land use change is also known to favour more adaptable, synanthropic
species such as *M. fascicularis*(McFarlane et al., 2012). Considering the spillover risk posed

by wildlife reservoirs of *P. knowlesi,* clarifying any relationships between environmental

factors and parasitaemia in key host species may contribute to a more comprehensive

understanding of *P. knowlesi* transmission patterns.

Earth Observation (EO) data provides novel opportunities to investigate epidemiological

patterns of diseases which are linked to environmental drivers(Kalluri et al., 2007). In relation

to *P. knowlesi,* utility of fine-scale remote-sensing data has been demonstrated: examples

include satellite-derived data used to examine household–level exposure risk in relation to

proximate land configuration(Fornace et al., 2019b), UAV-imagery used to link real–time

deforestation to macaque host behavioural change(Stark et al., 2019), and remote-sensing

data used to interrogate risk factors for vector breeding sites(Byrne et al., 2021). Though

macroecological studies that utilise geospatial data are often confounded by issues of

matching temporal and spatial scales, as well as by the quality and accuracy of available

georeferencing, measures can be taken to account for this when examining the role of

environmental factors in modulating disease outcomes. Furthermore, ecological processes

occur and interact over a range of distances, or 'spatial scales'(Brock et al., 2019; Fornace

et al., 2016; Loh et al., 2016). This applies to determinants of vector-borne disease ecology,

from larval breeding microclimate to wildlife host foraging behaviour. As multiple influential

variables are rarely captured by a single scale(Cohen et al., 2016), data-driven methods can

be applied to examine risk factors over multiple scales and identify covariates at their most

influential extent(Byrne et al., 2021).

We hypothesise that higher prevalence of *P. knowlesi* in primate host species is driven by

LULCC, contributing to the strong associations described between LULCC and human *P.*

knowlesi risk. In conceptual frameworks and transmission models, it is often assumed that *P.*

knowlesi infections in NHPs are chronic and ubiquitous(Brock et al., 2016; Jeyaprakasam et

al., 2020). No studies have systematically assessed the extent and quality of available data

on *P. knowlesi* in NHPs. Independent studies investigating *P. knowlesi* in primates are

typically constrained by small sample sizes and confined geographic areas, limiting

inference that can be made about relationships between infection dynamics and landscape

characteristics. Systematic tools developed for epidemiological studies of disease

prevalence in human populations are rarely applied to the study of wildlife disease

prevalence; however, such tools can be used to capture the scale and contrast required in

macroecological studies to quantify disease burdens regionally. Furthermore, while recent

research has shown the impact of deforestation on the distribution of macaques in the

context of *P. knowlesi* (Moyes et al., 2016; Stark et al., 2019), associations between LULCC

and variation in the prevalence of simian *Plasmodium* spp. in primates have not been

explored. This study is the first to systematically assess *P. knowlesi* prevalence in NHPs at a

regional scale, and across a wide range of habitats. We aimed to 1) systematically assemble

a georeferenced dataset of *P. knowlesi* in NHPs; 2) evaluate variation in NHP *P. knowlesi*

prevalence by geographic region; and 3) assess environmental and spatial risk factors for *P.*

knowlesi prevalence in NHPs across Southeast Asia.

RESULTS

A systematic literature review was conducted in Medline, Embase and Web of Science to identify articles reporting prevalence of naturally acquired *Plasmodium knowlesi* in NHPs. 23 research articles were identified(Akter et al., 2015; Amir et al., 2020; Chang et al., 2011; Fungfuang et al., 2020; Gamalo et al., 2019; Ho et al., 2010; Jeslyn et al., 2011; Lee et al., 2011; M. I. Li et al., 2021; Muehlenbein et al., 2015; Putaporntip et al., 2010; Saleh Huddin et al., 2019; Seethamchai et al., 2008; Unpublished, 2015, 2013; Vythilingam et al., 2008; Zarith et al, 2021; Zhang et al., 2016), containing 148 unique primate survey records to form the dataset for analyses (see SI for details of JBI Critical Assessment, Table S5)(Munn et al., 2015). Year of sampling ranges from 2004–2019. No primatological studies were identified from Vietnam, Brunei or Timor-Leste. Full characteristics of the articles and individual study methodologies are reported in Supplementary Information (Table S2). Spatial resolution of the survey sites varied from GPS point coordinates to country-level administrative boundaries (Supplementary Table S7). Geographic distribution of sampling is illustrated in Figure 1.

Overall, records report on a cumulative total of 6322 primates, with the largest proportion sampled from Peninsular Malaysia (48.5%, n=3069/6322). Primate surveys were primarily conducted on Long-tailed macaques (*Macaca fascicularis*) (90.5%, n=5720/6322) followed by Pig-tailed macaques (*M. nemestrina)* (n=532/6322)(Amir et al., 2020; Lee et al., 2011; Muehlenbein et al., 2015; Putaporntip et al., 2010) (Table S3). Reported prevalence of *Plasmodium knowlesi* in NHPs ranged from 0%–100%. Only 87 of the surveys (58.8%, n=87/148) reported a positive diagnosis*,* with the remaining 61 sites finding no molecular evidence of *P. knowlesi* infection (41.2%) in any primates tested. A full breakdown of *P. knowlesi* infection rates according to reported primate characteristics can be found in SI, Table S4.

188 **Figure 1.** *Sampling sites and primate species sampled across Southeast Asia. 'Other' includes*
189 Trachypithecus obscurus *and undefined species from the genus* Presbytis. *Total surveys = 148.*

Meta-analysis of *P. knowlesi* prevalence

To quantify regional heterogeneity in simian cases of *P. knowlesi,* a one-stage meta-analysis of prevalence (number positive out of the number sampled) was conducted on primate malaria survey data. Overall pooled estimate for *P. knowlesi* prevalence was 11.99% (CI95% 9.35–15.26). Overall heterogeneity was assessed using the I^2 statistic. Substantial between-study heterogeneity ($I^2 \geq 75\%$) was found across all prevalence records ($I^2=80.5\%$; CI95% 77.3–83.1). In the sub-group analysis by region, pooled prevalence estimates are consistently low for Thailand (2.0%, CI95% 1.1–3.5%), moderate in Peninsular Malaysia (14.3%, CI95% 11.1–18.2) and elevated in Singapore (23.3%, CI95% 11.0–42.8) and Malaysian Borneo (41.1%, CI95% 20.8–64.9) (Figure 2). Sub-group heterogeneity was assessed using prediction intervals, derived from τ^2 statistic used to describe between-study variability. Prediction intervals indicate high heterogeneity of estimates within regions, consistent with expectations of high variability of prevalence across individual study sites. Detailed forest plots for individual prevalence estimates can be found in Supplementary Figures S6.

	Studies (*k*)	Primates (n)		Estimate (%)	CI95%	Pred. Interval	Weight
Borneo	10	251		41.1%	(20.8–64.9)	[0.02–0.96]	0.08
Singapore	13	1092		23.3%	(11.0–42.8)	[0.01–0.90]	0.10
Pen. Malaysia	91	3069		14.3%	(11.1–18.2)	[0.02–0.60]	0.64
Sumatra	5	108		9.8%	(3.4–25.1)	[0.01–0.66]	0.03
Philippines	5	163		4.6%	(0.7–25.3)	[0.00–0.97]	0.03
Laos	1	44		2.3%	(0.3–14.5)	·	0.01
Thailand	21	1496		2.0%	(1.1–3.5)	[0.01–0.06]	0.11
Myanmar	1	45		1.1%	(0.1–15.1)	·	0.01
Cambodia	1	54		0.9%	(0.1–12.9)	·	0.01
						*I*² (CI 95%)	
Overall	148	6322		12.0%	(9.4–15.3)	80.5% (77.3; 83.1)	

210 **Figure 2 (A)** *Forest plot of pooled estimates for* P. knowlesi *prevalence (%) in all non-human primates*
211 *tested (n=6322) across Southeast Asia, disaggregated by species and sampling site (k=148).*
212 *Random-effects meta-analysis, sub-grouped by region.* **(B)** *Map of regional prevalence estimates for*
213 *P. knowlesi prevalence in NHP in Southeast Asia from meta-analysis. Point colour denotes pooled*
214 *estimate (%). Size denotes total primates tested per region (n). Shading indicates data availability.*

Risk factor analysis

Covariate data and *P. knowlesi* prevalence data were used to fit additional models to explore the relationships between localised landscape configuration and NHP malaria prevalence. Environmental covariates were extracted from satellite-derived remote sensing datasets (Table 1) at either true sampling sites (GPS coordinates) or 10 random pseudo-sampling sites to account for geographic uncertainty in prevalence data. Host species was grouped as '*Macaca fascicularis'* or 'Other' due to sample counts of <10 for certain primate species. Only 57.4% (n=85/148 records) of data included year of sampling, deemed to be insufficient to assess temporal patterns in prevalence. Forest cover ranged from negligible to near total cover (100%) within buffer radii (Supplementary Table S12). Details of covariate data processing is illustrated in Supplementary Information (Figure S7–8).

Table 1. *Spatial and temporal resolution and sources for environmental covariates. PARA=Perimeter: Area ratio of forest class.*

Covariate	Spatial res.	Temporal res.	Source
Human density (p/km^2)	1km	2012	WorldPop(WorldPop, 2018)
Elevation (m)	1km	2003	SRTM 90m Digital Elevation v4.1(Jarvis, A., H.I. Reuter, A. Nelson, 2008)
Forest cover (%)	30m	Annual	Hansen's Global Forest Watch(Hansen et al., 2013)
Fragmentation (PARA>0)	30m	Annual	Hansen's Global Forest Watch(Hansen et al., 2013)

Following a two-stage approach for selection of explanatory variables, forest cover and PARA were retained at 5km as linear terms, human population density was retained at both 5km and 20km and primate species was retained as a categorical variable. Spearman's rank tests for residual correlation between final variables at selected scales indicates a strong negative correlation between forest cover and fragmentation index (PARA) (ρ= –0.75) (SI, Figure S10).

Adjusting for all other covariates in the model, we identified strong evidence of an effect between increasing forest cover and higher prevalence of *P. knowlesi* in NHPs within a 5km radius (aOR=1.38, CI95% 1.19–1.60; p<0.0001). Evidence was also found for an association between likelihood of *P. knowlesi* and higher degrees of forest fragmentation, measured by perimeter: area ratio (PARA) within 5km (aOR=1.17, CI95% 1.02–1.34, p<0.0281). Evidence suggests that human population density within a 5km radius is associated with risk of *P. knowlesi* in NHP (aOR=1.36, CI95% 1.16–1.58, p=0.0001) whilst human density within 20km has an inverse effect on likelihood of *P. knowlesi* (aOR=0.56, CI95% 0.46–0.67, p<0.0001).

M. fascicularis is also associated with higher prevalence relative to all other non-human

primate species (aOR=2.50, CI95% 1.31–4.85; *p*=0.0051). Additional complexity did not

provide a more optimal model fit and effect modification was not pursued. Final adjusted OR

for the multivariable model can be visualised in Figure 4 (Table S14).

Figure 4. *Multivariable regression results. Spatial scale denoted in square bracket. Forest cover = %.*
Adjusted OR (dots) and CI95% (whiskers) for factors associated with P. knowlesi *in NHPs at*
significant spatial scales. N=1354, accounting for replicate pseudo-sampling.

DISCUSSION

LULCC are widely linked to spillover of zoonotic pathogens from sylvatic reservoirs into human populations, and pathogen prevalence in wildlife host species is key in driving the force of infection in spillover events. Our initial analyses found that for *Plasmodium knowlesi,* there is substantial spatial heterogeneity and prevalence in non-human primates varies markedly between regions of Southeast Asia(Zhang et al., 2016). Consistent with our hypothesis that parasite density in primate hosts would be higher in areas experiencing habitat disturbance, we identified strong links between *P. knowlesi* in NHPs and measures of contemporaneous forest cover and forest fragmentation. To our knowledge, this is the first systematic study to find evidence of landscape changes influencing the distribution of *P. knowlesi* prevalence in NHPs. Results offer evidence that *P. knowlesi* infection rates in NHPs are linked to changes in land use across broad spatial scales, and that prevalence of *P. knowlesi* in reservoir species may be driving spillover risk across Southeast Asia. These findings could provide insight to improving surveillance of *P. knowlesi* and to the development of ecologically targeted interventions.

While previous studies have estimated that *P. knowlesi* infection would be chronic in all macaques, or as high as 50–90% for modelling *P. knowlesi* transmission in Malaysia(Brock et al., 2016), this data strongly suggests that this is not the case. Overall prevalence of *P. knowlesi* infection in all NHPs is markedly lower than usual estimates, emphasising the importance of accounting for 'negative' data in estimations of prevalence. Considerable heterogeneity was identified between and within regional estimates for *P. knowlesi* across Southeast Asia, which likely reflects genuine differences according to distinct climates and distinct habitats(Shearer et al., 2016). Malaysian Borneo was found to have an estimated prevalence over five-fold higher than West Malaysia. Crucially, such extreme prevalence estimates for NHPs in Borneo align with the known hotspot for human incidence of *P. knowlesi*(Cooper et al., 2020). By comparison, for Peninsular Malaysia, estimated prevalence is far lower than anticipated. Cases of human *P. knowlesi* do occur in West Malaysia, though transmission has been found to exhibit spatial clustering(Phang et al., 2020) which may correspond to pockets of high risk within the wider context of low prevalence of *P. knowlesi* in macaque populations. Regional trends in *P. knowlesi* also mask differences in infection rates between sample locations, driven by more localised factors. Multiple studies reported finding *P. knowlesi* infections in wild macaques to be low or absent in peri-domestic or urbanised areas, attributed to the absence of vector species typically found in forest fringes(Brant et al., 2016; Chua et al., 2019; Manin et al., 2016). This pattern is seen in reports from Peninsular Malaysia(Saleh Huddin et al., 2019; Vythilingam et al., 2008) , Singapore(Jeslyn et al., 2011; M. I. Li et al., 2021) and Thailand(Fungfuang et al.,

2020; Putaporntip et al., 2010). The high heterogeneity of reports here suggests that the
picture is even more complex. *P. knowlesi* infections may even vary between troops within a
single study site, as was seen in the Philippines(Gamalo et al., 2019). Fine-scale interactions
are unlikely to be captured by the scale of this study.

Ecological processes determining *P. knowlesi* infection are influenced by dynamic factors
over multiple spatial scales(Cohen et al., 2016). We utilised a data-driven methodology to
select variables at distances that capture maximum impact on *P. knowlesi* prevalence(Byrne
et al., 2021; Fornace et al., 2019b), with forest cover and forest fragmentation influential at
localised scales and human population density also exerting influence within wider radii.
Contrary to previous studies on risk factors for human incidence of *P. knowlesi*(Fornace et
al., 2019b, 2016), elevation was not found to be associated with *P. knowlesi* in NHPs at any
scale. Vector and host species composition vary substantially across tropical ecotones, and
it is likely that the study extent encompasses a range of putative vectors across different
landscapes, such as those of the Minimus Complex in northern regions(Parker et al., 2015)
or the recently incriminated *An.-collessi* and *An.-roperi* from the Umbrosus Group(de Ang et
al., 2021). Given that the vector species driving sylvatic transmission remain elusive, it is
conceivable that the elevation range covers multiple vector and host species niches and
explains the lack of observed relationship between elevation and *P. knowlesi* in NHPs.
Human population density was found to be significant at multiple distances, with contrasting
effects on parasite prevalence in NHP. Previous studies have found a negative association
between human density and vector density and biting rates in forested landscapes(Fornace
et al., 2019a). Across wide spatial scales, increased vector density in less populated, more
forested areas could generate higher parasite prevalence in NHPs. At the same time Long-
tailed macaques, a species shown here to subject to higher prevalence rates, are notorious
as nuisance animals and many of the available samples were collected opportunistically in
urban areas, which might underly the observed positive association between localised high
human density and higher prevalence in NHP. Whilst more data would be needed to
understand this interaction, this further demonstrates the importance of using approaches to
identify disease dynamics across multiple spatial scales(Brock et al., 2019).

A key finding is the link between high prevalence of *P. knowlesi* in primate host species with
high degrees of habitat fragmentation. Forest fragmentation is a key aspect of landscape
modification, where large contiguous areas of forest are broken into a mosaic of smaller
patches. This disturbs the ecological structure by increasing the density of forest fringes or
'edges', dynamic habitat at the boundaries between natural ecosystems and human-
modified landscapes(Borremans et al., 2019). Other studies have linked habitat

fragmentation to increased generalist parasite density in primates. In Uganda, a higher
prevalence and infection risk of protozoal parasites was observed in wild populations of red
colobus primates (*Procolobus rufomitratus*) inhabiting fragmented forests compared to those
in undisturbed habitat(Gillespie and Chapman, 2008). For *P. knowlesi,* creation of edge
habitat is thought to favour vectors of the Leucosphyrus Complex(Davidson et al., 2019;
Hawkes et al., 2019). *Anopheles* spp. presence can be predicted by indices of fragmentation
in Sabah, Borneo, with changing land use creating more suitable micro-climate for larval
habitats(Byrne et al., 2021), and an increased abundance of *An. balabacensis* found in
forest fringes(Hawkes et al., 2019; Wong et al., 2015). Increasing complexity of forest
boundaries results in increased density of edge habitat, with conceivably higher density of
vectors in forest fringes. Therefore, preferential use of fringe habitat and high exposure to
vectors in forest fringes may contribute to higher conspecific transmission of *P. knowlesi*
between primates in increasingly fragmented habitats. This finding also lends clarity to
landscape fragmentation as a risk factor for human exposure to *P. knowlesi* in Malaysian
Borneo(Brock et al., 2019; Fornace et al., 2019b), with changes in relative host density,
vector density and wildlife parasite prevalence in nascent forest fringes potentially enhancing
the spillover of this disease system into human populations in fragmented habitats.

Conversely, we saw a strong association between high parasite prevalence and high forest
coverage. Given an inverse relationship with fragmentation, with high forest density
correlating to low fragmentation indices and vice versa, this speaks to a trade-off between
dense forest cover and high habitat complexity and suggests an 'ideal' amount of forest
fragmentation that facilitates prevalence in primate hosts. For animals with larger home
ranges, individual-based disease models combined with movement ecology approaches
have shown that the most highly fragmented areas are less favourable for maintaining
parasite transmission(White et al., 2018). In Sabah, individual macaques were shown to
increase ranging behaviour in response to deforestation(Stark et al., 2019). Forest edge
density also peaks at intermediate levels of land conversion(Borremans et al., 2019). With
smaller forest patches in maximally fragmented landscapes potentially insufficient to support
macaque troops, this interplay between disease ecology and metapopulation theory may
explain why both forest density and habitat complexity appear to pose a greater risk for
simian *P. knowlesi.* Likewise, this may relate to the finding that in Borneo, larger forest
patches (lower fragmentation indices) were associated with *P. knowlesi* spillover in
Borneo(Fornace et al., 2019b). Overall, this finding offers an insight to mechanisms that
underpin the increased force of infection of *P. knowlesi* that is associated with LULCC.

There are limitations to consider in the available data and interpretation of these findings.

'Small-study effects' were observed in the dataset, suggestive of a bias toward positive

effect estimates(Stewart et al., 2012). This may be a result of data disaggregation and small

studies creating artefactually higher estimates or may reflect true bias in data collection

toward areas known to be endemic for *P. knowlesi* and convenience sampling of macaques.

Assumptions have also been made that sample site equates to habitat, which may not reflect

actual habitat use, and even accurate georeferenced datapoints are unlikely to entirely

reflect surrounding habitat within the macaque home range. Variability in study designs and

data reporting also impacted geospatial accuracy. Steps were taken to account for spatial

bias by extracting covariates at randomly generated pseudo-sampling points. Whilst

uncertainty cannot be eliminated, we demonstrate a robust methodology to accommodate

for geographical uncertainty in ecological studies. Future investigations should prioritise

systematic, georeferenced sampling across a range of landscape scenarios.

Results are based on broad regional ecological trends and may not be generalisable at

individual levels, or to all putative host species in all geographic contexts(Zhang et al., 2016).

Effects of fragmentation are likely to be dependent on land conversion type, species

composition and surrounding matrix habitat. Since specific land configurations have been

linked to *P. knowlesi* exposure in Borneo(Fornace et al., 2019b), broad forest classifications

used here may mask important differences in *P. knowlesi* between land classes. Use of

perimeter: area ratio (PARA) as a fragmentation index was justified given high forest

coverage in study sites(Wang et al., 2014), though Edge Density (ED) or normalised

Landscape Shape Index (nLSI) might be more appropriate in future analyses to account for

variation in forest abundance. As it was not possible to include contemporary land cover

classifications in this analysis, future studies would benefit from looking at forest type (e.g.,

primary forest, agroforest, plantation).

CONCLUDING REMARKS

Strong links have been identified between land use changes and ecosystem perturbation that favours the transmission of vector-borne diseases(Loh et al., 2016). Prevalence of *P. knowlesi* in macaques is likely to be a crucial determinant of human infection risk, and more representative estimates of *P. knowlesi* prevalence derived here can better inform regionally-specific transmission risk models. This study characterises landscape risk factors for heightened prevalence of *P. knowlesi* in NHPs. Findings provide evidence in support of the premise that *P. knowlesi* in primate hosts is partly driven by LULCC across Southeast Asia. While the full complexity is not captured by the covariates used, it is clear that *P. knowlesi* infection in NHPs is not restricted to densely forested areas. This study also demonstrates the utility of systematic meta-analysis tools and remote-sensing datasets in the investigation of macroecological disease trends, in conjunction with methods to standardise a spatially heterogeneous dataset and data-driven selection of spatial scales. Gaps identified in data reporting should inform more systematic and localised primatological surveys to disentangle precise mechanisms. Notwithstanding limitations, this study highlights the marked spatial heterogeneity and role of landscape complexity in driving *P. knowlesi* infection rates in NHPs. Given the clear intersection between human epidemiology and wildlife ecology, it is essential that infection dynamics within wildlife reservoirs are considered in future public health interventions.

Study site

This study focused on the simian malaria *Plasmodium knowlesi* across Southeast Asia, within 28°30'00.0"N, 92°12'00.0"E and 11°00'00.0"S, 141°00'00.0"E. Climate mainly corresponds to the equatorial tropical zone, with high temperatures and high humidity.

Data assembly

A systematic literature review was conducted under the CoCoPop framework (Condition, Context, Population)(Cuenca et al., 2022; Munn et al., 2015). All studies identified in the literature review were screened for data on NHPs with a confirmed *P. knowlesi* diagnosis or absence data (zero counts of *P. knowlesi* with appropriate diagnostic methods). Exclusion criteria included (a) studies exclusively relying on microscopy(Antinori et al., 2013) (b) laboratory, animal model or experimental infection studies (c) data from outside of Southeast Asia. No limit was set on the temporal range. Duplicate records reporting results from the same surveys were removed, with one record per survey retained. Critical appraisal of the studies was conducted using the Joanna Briggs Institute (JBI) checklist for prevalence studies(Munn et al., 2015)(Munn et al., 2015) (see Supplementary Information (SI) for details and criteria). A flowchart of the selection process is illustrated in Figure S3, with a full list of articles included provided in Table S2.

Primary outcome was defined as *P. knowlesi* prevalence (*p,* proportion positive for *P. knowlesi* infection from *n* sampled NHPs). For each independent primate study, the following variables were extracted: year of data collection, primate species sampled, primate status (wild/captive), diagnostic test (PCR/sequencing) and target gene(s), sampling method (routine/purposive), number of *P. knowlesi* positive samples, number of *Plasmodium* spp. positive samples, total number of primates tested and geographical information.

In most studies identified, study site was only geolocated to a geographic area or descriptive location. Geolocation was assigned at the lowest available level of administrative polygon (i.e., district/state/country) by cross-referencing reported sampling location with GADM (v3.6) administrative boundaries. If specific location was given, GPS coordinates were assigned via Google Maps. For data visualisation, centroids were extracted per polygon and point coordinates were plotted in QGIS (3.10.14) and R (4.1.0) software.

Meta-analysis of *P. knowlesi* prevalence

Meta-analysis was conducted using methods that are standard in the analysis of human disease prevalence in individual participant datasets (IDP)(Liberati et al., 2009; Stewart et

al., 2012). Data were disaggregated by geographic location (site) and primate species, to

illustrate variance in prevalence by survey unit(Stewart et al., 2012). One-stage meta-

analysis is considered appropriate for studies where the outcome may be infrequent, so data

was included in a single model under the 'DerSimonian and Laird' variance estimator (Munn

et al., 2015). Sensitivity analyses were conducted to compare methods for the back-

transformation of prevalence estimates. For studies where prevalence estimates tend

towards 0% or 100%, variance tends towards 0. To stabilise the variance and enable back-

transformation of zero prevalence records, logit method was selected for the transformation

of prevalence, with the inverse variance method used for individual study weights (see SI for

details).

Overall heterogeneity of prevalence records was assessed using the I^2 statistic(Hippel,

2015), a relative estimate of true between-study variance. Sub-group analysis was

conducted according to geographic region, with the heterogeneity of reported prevalence

within regional sub-groups assessed using prediction intervals derived from the τ^2 statistic.

Small-study effects, including selection and publication biases, were assessed by examining

funnel plots and imputing 'missing' estimates using the trim-and-fill method(Lin and Chu,

2018). Full rationale and details of small-study effect assessments can be found in

Supplementary Information.

Remote sensing data

Satellite-derived remote sensing datasets were used to assemble local environmental and

anthropogenic covariates. Gridded UN-adjusted human population estimates were

assembled at 1km resolution from WorldPop(WorldPop, 2018). Elevation data was obtained

from NASA SRTM 90m Digital Elevation Database v4.1 (CGIAR-CSI)(Jarvis, A., H.I. Reuter,

A. Nelson, 2008) with a spatial resolution of 1km. Contemporaneous tree cover was derived

from Hanson's Global Forest Watch (30m)(Hansen et al., 2013). Tree cover was classified

as ≥50% crown density, and then matched to primate data by sample site geolocation and

by year of sample collection to account for rapid forest loss (SI, Figure S7). Where a broad

timeframe of sampling was provided (≥3 years), median year was used. Full details can be

found in Supplementary Information (Table S11–12, Figure S8).

Perimeter: area ratio (PARA, ratio of patch perimeter length to patch surface area) of given

land class is a key metric for habitat conversion, where a higher PARA provides a measure

of boundary complexity and indicates a more fragmented landscape(McGarigal and

Cushman, 2012). Mean PARA was extracted from forest cover within circular buffers.

Habitat fragmentation has been shown to correlate with disease transmission

parameters(Borremans et al., 2019; Faust et al., 2018), but definitions often lack precision

and can be considered with respect to 'separation effects' (division and isolation of patches)
and 'geometric effects' (changes to ratios of perimeter and core habitat)(Wilkinson et al.,
2018). PARA provides a measure of forest edge density within the buffer area (PARA>0)
and has been shown to provide a good index of fragmentation and discrimination of spatial
aggregation across areas where habitat abundance is high(Wang et al., 2014) (SI, Figure
S9).

For every georeferenced sampling point, mean values for all selected covariates were
extracted within buffer radii at 5km, 10km and 20km (SI, Figure S9). Buffer area sizes were
selected to investigate multiple spatial scales over which associations between risk factors
and *P. knowlesi* prevalence might occur. A minimum radius of 5km was chosen to
approximate the maximum ranging distance for *M. fascicularis*(Waxman et al., 2014), with
wider radii (10–20km) included to account for the geographic uncertainties in areal data.

Analysis of environmental risk factors

Generalised linear mixed-effect regression models (GLMM) were fitted using a binomial
distribution with a logit link. To account for within-study correlation, a unique identifier
combining author and study was included as a random intercept in all models.

For studies with exact GPS coordinates, precise environmental data at a single site could be
obtained. For surveys published without GPS coordinates, there is considerable geographic
uncertainty in the exact sampling location (SI, Table S7). Uncertainty in the spatial and
environmental determinants of prevalence generates a sampling bias, with the precision of
covariates correlated to certain studies. Strategies were employed to account for and
mitigate the effect of spatial uncertainty. Each prevalence observation was replicated and
assigned a random sample of environmental realisations. 10 random sampling points were
generated within the sampling area provided by the study, and covariates were extracted at
each proxy sampling site (SI, Table S8). Selection of random points was validated by visual
inspection of the stability of model coefficients with the inclusion of an increasing number of
points. Number of points was selected conservatively at the point where coefficients
stabilised (n=10), minimising the risk of retaining covariates in model selection due to
artificially inflated sample sizes. Artificial inflation of sample size in the replicated data was
then accommodated for by reducing individual observation weights to $1/10^{th}$ within the
model.

Each covariate at each spatial scale was assessed for inclusion in the multivariable model
based on bivariable analysis and a criterion of $p > 0.2$ under likelihood ratio tests (LRT)
(Table S13). A quadratic term for the fragmentation index 'PARA' was included to account
for possible nonlinearity. Multicollinearity among independent predictors at multiple scales

was examined according to calculation of variance inflation factor (VIF). The VIF of each

predictor variable was examined following a stepwise procedure, starting with a saturated

model and sequentially excluding the variable with the highest VIF score from the model.

Stepwise selection continued in this manner until the entire subset of explanatory variables

in the global model satisfied a moderately conservative threshold of VIF ≤6(Rogerson,

2001). Qualifying variables obtained were then assessed for model inclusion using a

backward stepwise strategy, removing variables with the highest p value (Wald test) until a

pre-defined threshold of α <0.05. Model parsimony was examined according to reduction in

Akaike's Information Criterion (AIC) value. Spearman's rank tests were conducted on the

selected variables to observe residual correlation, plotted as a correlation matrix. Covariate

pairs with correlation coefficient ρ>0.6 were considered highly correlated but were retained.

Fully adjusted OR for associations between environmental covariates and *P. knowlesi*

prevalence were derived from the final multivariable GLMM with p values derived from LRT.

Ethics

Ethics approval was not required for this research, following assessment by the LSHTM

Research Governance & Integrity Office (LSHTM MSc Ethics Ref: 25429).

ACKNOWLEDGEMENTS

Research was supported by the Sir Henry Dale Fellowship, jointly funded by the Wellcome Trust and the Royal Society (Grant Number 221963/Z/20/Z). Data collected in Peninsular Malaysia by the Faculty of Veterinary Medicine, Universiti Putra Malaysia, was funded by the Ministry of Higher Education Malaysia (Grant LRGS/1/2018/UM/01/1/). Additional funding was supported by the World Health Organization.

REFERENCES

Akter R, Vythilingam I, Khaw LT, Qvist R, Lim YA-LL, Sitam FT, Venugopalan B, Sekaran SD, Vythilingam I, Khaw LT, Lim YA-LL, Sitam FT, Venugopalan B, Sekaran SD, Qvist R, Lim YA-LL, Sitam FT, Venugopalan B, Sekaran SD, Vythilingam I, Khaw LT, Lim YA-LL, Sitam FT, Venugopalan B, Sekaran SD. 2015. Simian malaria in wild macaques: first report from Hulu Selangor district, Selangor, Malaysia. *Malar J* **14**:386. doi:http://dx.doi.org/10.1186/s12936-015-0856-3

Amir A, Shahari S, Liew JWK, de Silva JR, Khan MB, Lai MY, Snounou G, Abdullah ML, Gani M, Rovie-Ryan JJ, Lau YL. 2020. Natural Plasmodium infection in wild macaques of three states in peninsular Malaysia. *Acta Trop* **211**:1–6. doi:10.1016/j.actatropica.2020.105596

Antinori S, Galimberti L, Milazzo L, Corbellino M. 2013. Plasmodium knowlesi: The emerging zoonotic malaria parasite. *Acta Trop* **125**:191–201. doi:http://dx.doi.org/10.1016/j.actatropica.2012.10.008

Borremans B, Faust C, Manlove KR, Sokolow SH, Lloyd-Smith JO. 2019. Cross-species pathogen spillover across ecosystem boundaries: Mechanisms and theory. *Philosophical Transactions of the Royal Society B: Biological Sciences* **374**:20180344. doi:10.1098/rstb.2018.0344

Brant HL. 2011. Changes in abundance, diversity and community composition of mosquitoes based on different land use in Sabah, Malaysia. *Imperial College London*.

Brant HL, Ewers RM, Vythilingam I, Drakeley C, Benedick S, Mumford JD. 2016. Vertical stratification of adult mosquitoes (Diptera: Culicidae) within a tropical rainforest in Sabah, Malaysia. *Malar J* **15**:1–9. doi:10.1186/s12936-016-1416-1

Brock PM, Fornace KM, Grigg MJ, Anstey NM, William T, Cox J, Drakeley CJ, Ferguson HM, Kao RR, Fornace KM, Cox J, Drakeley CJ, Grigg MJ, Anstey NM, William T, Kao RR. 2019. Predictive analysis across spatial scales links zoonotic malaria to deforestation. *Proc Biol Sci* **286**:20182351. doi:http://dx.doi.org/10.1098/rspb.2018.2351

Brock PM, Fornace KM, Parmiter M, Cox J, Drakeley CJ, Ferguson HM, Kao RR. 2016. Plasmodium knowlesi transmission: integrating quantitative approaches from epidemiology and ecology to understand malaria as a zoonosis. *Parasitology* **143**:389–400. doi:10.1017/S0031182015001821

Byrne I, Drakeley CJ, Fornace KM, Manin BO, Chua TH, Aure W, Vythilingam I, Ferguson HM. 2021. Environmental and spatial risk factors for the larval habitats of Plasmodium knowlesi vectors in Sabah, Malaysian Borneo. *Sci Rep* **11**:11810. doi:http://dx.doi.org/10.1038/s41598-021-90893-1

Chang Q, Sun X, Wang J, Yin J, Song J, Peng S, Lu H, Zhou H, Jiang N, Chen Q. 2011. Identification of Hepatocystis species in a macaque monkey in northern Myanmar. *Res Rep Trop Med* **2**:141–146.

Chua TH, Manin BO, Vythilingam I, Fornace K, Drakeley CJ. 2019. Effect of different habitat types on abundance and biting times of Anopheles balabacensis Baisas (Diptera: Culicidae) in Kudat district of Sabah, Malaysia. *Parasit Vectors* **12**:364. doi:http://dx.doi.org/10.1186/s13071-019-3627-0

Cohen JM, Civitello DJ, Brace AJ, Feichtinger EM, Ortega CN, Richardson JC, Sauer EL, Liu X, Rohr JR. 2016. Spatial scale modulates the strength of ecological processes driving disease distributions. *Proceedings of the National Academy of Sciences* **113**:E3359–E3364. doi:10.1073/pnas.1521657113

Cooper DJ, Rajahram GS, William T, Jelip J, Mohammad R, Benedict J, Alaza DA, Malacova E, Yeo TW, Grigg MJ, Anstey NM, Barber BE. 2020. Plasmodium knowlesi Malaria in Sabah, Malaysia, 2015-2017: Ongoing increase in incidence despite nearelimination of the human-only plasmodium species. *Clinical Infectious Diseases* **70**:361–367. doi:10.1093/cid/ciz237

Cuenca R, Journal M, Cuenca PR, Key S, Lindblade KA, Vythilingam I, Drakeley C, Fornace K. 2022. Is there evidence of sustained human - mosquito - human transmission of the zoonotic malaria Plasmodium knowlesi ? A systematic literature review. *Malar J* 1–19. doi:10.1186/s12936-022-04110-z

Davidson G, Chua TH, Cook A, Speldewinde P, Weinstein Tock H.; ORCID: http://orcid.org/0000-0002-8984-8723 PAO-C, Weinstein P, Weinstein Tock H.; ORCID: http://orcid.org/0000-0002-8984-8723 PAO-C, Weinstein P. 2019. The Role of Ecological Linkage Mechanisms in Plasmodium knowlesi Transmission and Spread. *Ecohealth* **16**:594–610. doi:http://dx.doi.org/10.1007/s10393-019-01395-6

de Ang JX, Yaman K, Kadir KA, Matusop A, Singh B. 2021. New vectors that are early feeders for Plasmodium knowlesi and other simian malaria parasites in Sarawak, Malaysian Borneo. *Sci Rep* **11**:1–12. doi:10.1038/s41598-021-86107-3

Faust CL, McCallum HI, Bloomfield LSP, Gottdenker NL, Gillespie TR, Torney CJ, Dobson AP, Plowright RK. 2018. Pathogen spillover during land conversion. *Ecol Lett* **21**:471–483. doi:10.1111/ele.12904

Fornace K. 2022. No evidence of sustained nonzoonotic Plasmodium knowlesi transmission: estimating species-specic spatiotemporally varying reproduction numbers for malaria in Malaysia. *Nature Portfolio*. doi:10.21203/rs.3.rs-2278329/v1

Fornace KM, Abidin TR, Alexander N, Brock P, Grigg MJ, Murphy A, William T, Menon J, Drakeley CJ, Cox J, Abidin TR, Brock P, Grigg MJ, Murphy A, William T, Menon J. 2016. Association between landscape factors and spatial patterns of Plasmodium knowlesi infections in Sabah, Malaysia. *Emerg Infect Dis* **22**:201–208. doi:http://dx.doi.org/10.3201/eid2202.150656

Fornace KM, Alexander N, Abidin TR, Brock PM, Chua TH, Vythilingam I, Ferguson HM, Manin BO, Wong ML, Ng SH, Cox J, Drakeley C. 2019a. Local human movement patterns and land use impact exposure to zoonotic malaria in Malaysian borneo. *Elife* **8**. doi:10.7554/ELIFE.47602

Fornace KM, Brock PM, Abidin TR, Grignard L, Herman LS, Chua TH, Daim S, William T, Patterson CLEBEB, Hall T, Grigg MJ, Anstey NM, Tetteh KKAA, Cox J, Drakeley CJ. 2019b. Environmental risk factors and exposure to the zoonotic malaria parasite Plasmodium knowlesi across northern Sabah, Malaysia: a population-based cross-sectional survey. *Lancet Planet Health* **3**:e179--e186. doi:https://doi.org/10.1016/S2542-5196(19)30045-2

Fornace KM, Herman LS, Grignard L, Cox J, Tetteh KKA, Drakeley CJ, Abidin TR, Chua TH, Daim S, Lorenzo PJ, Espino F, Nuin NA, Ying LT, Grigg MJ, William Kimberly M.; ORCID: http://orcid.org/0000-0002-5484-241X AO - Grignard, Lynn; ORCID:

http://orcid.org/0000-0001-8305-3981 AO - Lorenzo, Pauline J.; ORCID: http://orcid.org/0000-0003-0563-4582 TAO-F. 2018. Exposure and infection to Plasmodium knowlesi in case study communities in Northern Sabah, Malaysia and Palawan, The Philippines. *PLoS Negl Trop Dis* **12**:e0006432. doi:http://dx.doi.org/10.1371/journal.pntd.0006432

Fornace KM, Lines J, Drakeley CJ, Diaz Kimberly M.; ORCID: http://orcid.org/0000-0002-5484-241X AVAO-F, Diaz A v., Lines J, Drakeley CJ. 2021. Achieving global malaria eradication in changing landscapes. *Malar J* **20**:69. doi:http://dx.doi.org/10.1186/s12936-021-03599-0

Fungfuang W, Udom C, Tongthainan D, Kadir KA, Singh B. 2020. Malaria parasites in macaques in Thailand: Stump-tailed macaques (Macaca arctoides) are new natural hosts for Plasmodium knowlesi, Plasmodium inui, Plasmodium coatneyi and Plasmodium fieldi. *Malar J* **19**:1–7. doi:10.1186/s12936-020-03424-0

Gamalo LE, Dimalibot J, Kadir KA, Singh B, Paller VG. 2019. Plasmodium knowlesi and other malaria parasites in long-tailed macaques from the Philippines. *Malar J* **18**:1–7. doi:10.1186/s12936-019-2780-4

Ghinai I, Cook J, Hla TTW, Htet HMT, Hall T, Lubis IND, Ghinai R, Hesketh T, Naung Y, Lwin MM, Latt TS, Heymann DL, Sutherland CJ, Drakeley C, Field N. 2017. Malaria epidemiology in central Myanmar: identification of a multi-species asymptomatic reservoir of infection. *Malar J* **16**. doi:10.1186/s12936-016-1651-5

Gillespie TR, Chapman CA. 2008. Forest fragmentation, the decline of an endangered primate, and changes in host-parasite interactions relative to an unfragmented forest. *Am J Primatol* **70**:222–230. doi:10.1002/ajp.20475

Global Malaria Programme. n.d. https://www.who.int/teams/global-malaria-programme/elimination

Grigg MJ, Barber BE, Anstey NM, Cox J, Fornace KM, Drakeley CJ, William T, Jelip J, Brock PM, von Seidlein L, Yeo TW, Fornace KM, Brock PM, von Seidlein L, Barber BE, Anstey NM, Yeo TW, Drakeley CJ, Cox J, Fornace KM, Drakeley CJ, William T, Jelip J, Brock PM, von Seidlein L, Yeo TW, Fornace KM, Brock PM, von Seidlein L, Barber BE, Anstey NM, Yeo TW, Drakeley CJ. 2017. Individual-level factors associated with the risk of acquiring human Plasmodium knowlesi malaria in Malaysia: a case-control study. *Lancet Planet Health* **1**:e97–e104. doi:http://dx.doi.org/10.1016/S2542-5196%2817%2930031-1

Hansen MC, Potapov P v., Moore R, Hancher M, Turubanova SA, Tyukavina A, Thau D, Stehman S v., Goetz SJ, Loveland TR, Kommareddy A, Egorov A, Chini L, Justice CO, Townshend. JRG. 2013. High-Resolution Global Maps of 21st-Century Forest Cover Change. *Science 342 (15 November)* 850–53.

Hanski I, Ovaskainen O. 2000. The metapopulation capacity of a fragmented landscape. *Nature* **404**:755–758. doi:10.1038/35008063

Hawkes FM, Manin BO, Cooper A, Daim S, R H, Jelip J, Husin T, Chua TH, Cooper A, Jelip J, Husin Benny O.; ORCID: http://orcid.org/0000-0003-0726-6146 AO - Chua, Tock H.; ORCID: http://orcid.org/0000-0002-8984-8723 TAO-M, Homathevi R, Jelip J, Husin T, Chua TH. 2019. Vector compositions change across forested to deforested ecotones in emerging areas of zoonotic malaria transmission in Malaysia. *Sci Rep*, Erratum for: Sci Rep. 2019 Sep 16;9(1):13312; PMID: 31527622

[https://www.ncbi.nlm.nih.gov/pubmed/31527622] **9**:13312.
doi:http://dx.doi.org/10.1038/s41598-019-49842-2

Hippel PT von. 2015. The heterogeneity statistic I2 can be biased in small meta-analyses. *BMC Med Res Methodol* **15**. doi:10.1186/S12874-015-0024-Z

Ho GC, Lee CL, Sharma RSK. 2010. Association of Institutions for Tropical Veterinary Medicine (AITVM) Conference 23-26 August 2010 Bangkok,Thailand.

Ilacqua RC, Medeiros-Sousa AR, Ramos DG, Obara MT, Ceretti-Junior W, Mucci LF, Marrelli MT, Laporta GZ. 2021. Reemergence of Yellow Fever in Brazil: The Role of Distinct Landscape Fragmentation Thresholds. *J Environ Public Health* **2021**:8230789. doi:10.1155/2021/8230789

Imai N, Furukawa T, Tsujino R, Kitamura S, Yumoto T. 2018. Factors affecting forest area change in southeast Asia during 1980-2010. *PLoS One* **13**:1–14. doi:10.1371/journal.pone.0197391

Jarvis, A., H.I. Reuter, A. Nelson EG. 2008. Hole-filled SRTM for the globe Version 4, available from the CGIAR-CSI SRTM 90m Database (http://srtm.csi.cgiar.org).

Jeslyn WPS, Huat TC, Vernon L, Irene LMZ, Sung LK, Jarrod LP, Singh B, Ching NL, Huat TC, Singh B, Vernon L, Jarrod LP, Irene LMZ, Sung LK, Jarrod LP, Singh B, Ching NL, Huat TC, Singh B, Vernon L, Jarrod LP. 2011. Molecular epidemiological investigation of Plasmodium knowlesi in humans and macaques in Singapore. *Vector-Borne and Zoonotic Diseases* **11**:131–135. doi:http://dx.doi.org/10.1089/vbz.2010.0024

Jeyaprakasam NK, Liew JWK, Low VL, Wan-Sulaiman WY, Vythilingam I. 2020. Plasmodium knowlesi infecting humans in southeast asia: What's next? *PLoS Negl Trop Dis* **14**:1–16. doi:10.1371/journal.pntd.0008900

Kalluri S, Gilruth P, Rogers D, Szczur M. 2007. Surveillance of arthropod vector-borne infectious diseases using remote sensing techniques: A review. *PLoS Pathog* **3**:1361–1371. doi:10.1371/journal.ppat.0030116

Koh GJN, Ismail PK, Koh D. 2019. Occupationally Acquired Plasmodium knowlesi Malaria in Brunei Darussalam. *Saf Health Work* **10**:122–124. doi:http://dx.doi.org/10.1016/j.shaw.2018.09.002

Lee K-SS, Divis PCSS, Zakaria SK, Julin RA, Singh B, Matusop A, Conway DJ, Cox-Singh J, Julin RA, Conway DJ, Cox-Singh J, Singh B, Matusop A, Conway DJ, Cox-Singh J. 2011. Plasmodium knowlesi: Reservoir hosts and tracking the emergence in humans and macaques. *PLoS Pathog* **7**:e1002015. doi:10.1371/journal.ppat.1002015

Lehman SM, Rajaonson A, Day S. 2006. Edge effects and their influence on lemur density and distribution in southeast Madagascar. *Am J Phys Anthropol* **129**:232–241. doi:10.1002/ajpa.20241

Li J, Jia K, Liu Y, Yuan B, Xia M, Zhao W. 2021. Spatiotemporal distribution of zika virus and its spatially heterogeneous relationship with the environment. *Int J Environ Res Public Health* **18**:1–14. doi:10.3390/ijerph18010290

Li MI, Mailepessov D, Vythilingam I, Lee V, Lam P, Ng LC, Tan CH. 2021. Prevalence of simian malaria parasites in macaques of Singapore. *PLoS Negl Trop Dis* **15**:1–12. doi:10.1371/journal.pntd.0009110

Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, Clarke M,
Devereaux PJ, Kleijnen J, Moher D. 2009. The PRISMA statement for reporting
systematic reviews and meta-analyses of studies that evaluate health care
interventions: Explanation and elaboration. *PLoS Med* **6**.
doi:10.1371/JOURNAL.PMED.1000100

Lin L, Chu H. 2018. Quantifying Publication Bias in Meta-Analysis. *Biometrics* **74**:785.
doi:10.1111/BIOM.12817

Link A, de Luna A, Alfonso F, Giraldo-Beltran P, Ramirez F. 2010. Initial effects of
fragmentation on the density of three neotropical primate species in two lowland forests
of Colombia. *Endanger Species Res* **13**:41–50. doi:10.3354/esr00312

Loh EH, Murray KA, Nava A, Daszak P. 2016. Evaluating the Links between Biodiversity,
Land-Use Change, and Infectious Disease Emergence. *Tropical Conservation* 79–88.

Maeno Y, Quang NT, Marchand RP, Culleton R, Kawai S, Masuda G, Nakazawa S. 2015.
Humans frequently exposed to a range of non-human primate malaria parasite species
through the bites of Anopheles dirus mosquitoes in South-central Vietnam. *Parasit*
Vectors **8**:376. doi:http://dx.doi.org/10.1186/s13071-015-0995-y

Manin BO, Ferguson HM, Vythilingam I, Fornace K, William T, Torr SJ, Drakeley C, Chua
TH, Ferguson HM, Vythilingam I, Fornace K, Drakeley C, William T, Torr SJ, Drakeley
C, Chua TH, Ferguson HM, Vythilingam I, Fornace K, Drakeley C, William T, Torr SJ,
Drakeley C, Chua TH. 2016. Investigating the Contribution of Peri-domestic
Transmission to Risk of Zoonotic Malaria Infection in Humans. *PLoS Negl Trop Dis*
10:1–14. doi:10.1371/journal.pntd.0005064

McFarlane R, Sleigh A, McMichael T. 2012. Synanthropy of wild mammals as a determinant
of emerging infectious diseases in the Asian-Australasian region. *Ecohealth* **9**:24–35.
doi:10.1007/s10393-012-0763-9

McGarigal K, Cushman SA. 2012. FRAGSTATS v4: Spatial Pattern Analysis Program for
Categorical and Continuous Maps. Computer software program produced by the
authors at the University of Massachusetts, Amherst.
http://www.umass.edu/landeco/research/fragstats/fragstats.html

Moyes CL, Shearer FM, Huang Z, Wiebe A, Gibson HS, Nijman V, Mohd-Azlan J, Brodie JF,
Malaivijitnond S, Linkie M, Samejima H, O'Brien TG, Trainor CR, Hamada Y, Giordano
AJ, Kinnaird MF, Elyazar IRFF, Sinka ME, Vythilingam I, Bangs MJ, Pigott DM, Weiss
DJ, Golding N, Hay SI, Huang Z, Gibson HS, Sinka ME, Weiss DJ, Nijman V, Mohd-
Azlan J, Brodie JF, Malaivijitnond S, Linkie M, Samejima H, O'Brien TG, Kinnaird MF,
Trainor CR, Hamada Y, Giordano AJ, Elyazar IRFF, Vythilingam I, Bangs MJ, Pigott
DM, Hay SI. 2016. Predicting the geographical distributions of the macaque hosts and
mosquito vectors of Plasmodium knowlesi malaria in forested and non-forested areas.
Parasit Vectors **9**:242. doi:http://dx.doi.org/10.1186/s13071-016-1527-0

Muehlenbein MP, Pacheco MA, Taylor JE, Prall SP, Ambu L, Nathan S, Alsisto S, Ramirez
D, Escalante AA, Pacheco MA, Taylor JE, Ambu L, Nathan S, Alsisto S, Ramirez D,
Escalante AA. 2015. Accelerated diversification of nonhuman primate malarias in
Southeast Asia: Adaptive radiation or geographic speciation? *Mol Biol Evol* **32**:422–
439. doi:10.1093/molbev/msu310

Munn Z, MClinSc SM, Lisy K, Riitano D, Tufanaru C. 2015. Methodological guidance for
systematic reviews of observational epidemiological studies reporting prevalence and

cumulative incidence data. *Int J Evid Based Healthc* **13**:147–153.
doi:10.1097/XEB.0000000000000054

Murray KA, Daszak P. 2013. Human ecology in pathogenic landscapes: two hypotheses on
how land use change drives viral emergence. *Curr Opin Virol* **3**:79–83.
doi:10.1016/j.coviro.2013.01.006

Nunn CL, Dokey ATW. 2006. Ranging patterns and parasitism in primates. *Biol Lett* **2**:351–
354. doi:10.1098/rsbl.2006.0485

Parker DM, Carrara VI, Pukrittayakamee S, McGready R, Nosten FH. 2015. Malaria ecology
along the Thailand-Myanmar border. *Malar J* **14**:388.
doi:http://dx.doi.org/10.1186/s12936-015-0921-y

Phang WK, Hamid MHA, Jelip J, Mudin RN, Chuang T-W, Lau YL, Fong MY. 2020. Spatial
and Temporal Analysis of Plasmodium knowlesi Infection in Peninsular Malaysia, 2011
to 2018. *Int J Environ Res Public Health* **17**. doi:10.3390/ijerph17249271

Putaporntip C, Jongwutiwes S, Thongaree S, Seethamchai S, Grynberg P, Hughes AL,
Chaturong Putaporntip1,*, Somchai Jongwutiwes1, Siriporn Thongaree2, Sunee
Seethamchai3, Priscila Grynberg4 and ALH. 2010. Ecology of malaria parasites
infecting Southeast Asian macaques: evidence from cytochrome b sequences
Chaturong. *Mol Ecol* **19**:3466–3476. doi:10.1111/j.1365-294X.2010.04756.x.Ecology

Rogerson PA. 2001. Statistical Methods for Geography. SAGE, London.

Saleh Huddin A, Md Yusuf NA, Razak MRMA, Ogu Salim N, Hisam S. 2019. Genetic
diversity of Plasmodium knowlesi among human and long-tailed macaque populations
in Peninsular Malaysia: The utility of microsatellite markers. *Infection, Genetics and*
Evolution **75**:103952. doi:10.1016/j.meegid.2019.103952

Seethamchai S, Putaporntip C, Malaivijitnond S, Cui L, Jongwutiwes S, Malaivijitnond S, Cui
L. 2008. Malaria and Hepatocystis species in wild macaques, southern Thailand.
American Journal of Tropical Medicine & Hygiene **78**:646–653.
doi:http://dx.doi.org/10.4269/ajtmh.2008.78.646

Setiadi W, Sudoyo H, Trimarsanto H, Wangsamuda S, Asih PBS, Syafruddin D, Sihite BA,
Saragih RJ, Juliawaty R. 2016. A zoonotic human infection with simian malaria,
Plasmodium knowlesi, in Central Kalimantan, Indonesia. *Malar J* **15**:218.
doi:http://dx.doi.org/10.1186/s12936-016-1272-z

Shearer FM, Huang Z, Weiss DJ, Wiebe A, Gibson HS, Battle KE, Pigott DM, Brady OJ,
Putaporntip C, Jongwutiwes S, Lau YL, Manske M, Amato R, Elyazar IRF, Vythilingam
I, Bhatt S, Gething PW, Singh B, Golding N, Hay SI, Moyes CL. 2016. Estimating
Geographical Variation in the Risk of Zoonotic Plasmodium knowlesi Infection in
Countries Eliminating Malaria. *PLoS Negl Trop Dis* **10**:1–17.
doi:10.1371/journal.pntd.0004915

Simon JA, Marrotte RR, Desrosiers N, Fiset J, Gaitan J, Gonzalez A, Koffi JK, Lapointe FJ,
Leighton PA, Lindsay LR, Logan T, Milord F, Ogden NH, Rogic A, Roy-Dufresne E,
Suter D, Tessier N, Millien V. 2014. Climate change and habitat fragmentation drive the
occurrence of Borrelia burgdorferi, the agent of Lyme disease, at the northeastern limit
of its distribution. *Evol Appl* **7**:750–764. doi:10.1111/eva.12165

Singh B, Sung LK, Matusop A, Radhakrishnan A, Shamsul SSG, Cox-Singh J, Thomas A,
Conway DJ. 2004. A large focus of naturally acquired Plasmodium knowlesi infections
in human beings. *The Lancet* **363**:1017–1024. doi:10.1016/S0140-6736(04)15836-4

Speldewinde P, Davidson G, Cook A, Chua TH, Weinstein Gael; ORCID:
http://orcid.org/0000-0002-9447-9125 PAO-D, Cook A, Speldewinde P, Weinstein P.
2019. Defining the ecological and evolutionary drivers of Plasmodium knowlesi
transmission within a multi-scale framework. *Malar J* **18**:66.
doi:https://dx.doi.org/10.1186/s12936-019-2693-2

Stark DJ, Fornace KM, Brock PM, Abidin TR, Gilhooly L, Jalius C, Goossens B, Drakeley
CJ, Salgado-Lynn M. 2019. Long-Tailed Macaque Response to Deforestation in a
Plasmodium knowlesi-Endemic Area. *Ecohealth* **16**:638–646. doi:10.1007/s10393-019-
01403-9

Stewart GB, Altman DG, Askie LM, Duley L, Simmonds MC, Stewart LA. 2012. Statistical
Analysis of Individual Participant Data Meta-Analyses: A Comparison of Methods and
Recommendations for Practice. *PLoS One* **7**. doi:10.1371/JOURNAL.PONE.0046042

Unpublished. 2015. Personal Correspondance.

Unpublished. 2013. Personal Correspondance.

Vaz VC, D'Andrea. PS, Jansen AM. 2007. Effects of habitat fragmentation on wild mammal
infection by Trypanosoma cruzi. *Parasitology* **134**:1785–93.

Vythilingam I, Noorazian YM, Huat TC, Jiram AI, Yusri YM, Azahari AH, Norparina I,
Noorrain A, Lokmanhakim S, Huat TC, Azahari AH, Noorrain A, Lokmanhakim S, Jiram
AI, Yusri YM, Azahari AH, Norparina I, Noorrain A, Lokmanhakim S, Huat TC, Azahari
AH, Noorrain A, Lokmanhakim S. 2008. Plasmodium knowlesi in humans, macaques
and mosquitoes in peninsular Malaysia. *Parasit Vectors* **1**:26.
doi:http://dx.doi.org/10.1186/1756-3305-1-26

Vythilingam I, Tan CH, Asmad M, Chan ST, Lee KS, Singh B. 2006. Natural transmission of
Plasmodium knowlesi to humans by Anopheles latens in Sarawak, Malaysia. *Trans R
Soc Trop Med Hyg* **100**:1087–1088. doi:10.1016/j.trstmh.2006.02.006

Wang X, Blanchet FG, Koper N. 2014. Measuring habitat fragmentation : An evaluation of
landscape pattern metrics 634–646. doi:10.1111/2041-210X.12198

Waxman D, Weinert LA, Welch JJ. 2014. Inferring host range dynamics from comparative
data: the protozoan parasites of new world monkeys. *Am Nat* **184**:65–74.
doi:http://dx.doi.org/10.1086/676589

White LA, Forester JD, Craft ME. 2018. Disease outbreak thresholds emerge from
interactions between movement behavior, landscape structure, and epidemiology. *Proc
Natl Acad Sci U S A* **115**:7374–7379. doi:10.1073/pnas.1801383115

Wilkinson DA, Marshall JC, French NP, Hayman DTS. 2018. Habitat fragmentation,
biodiversity loss and the risk of novel infectious disease emergence. *J R Soc Interface*
15. doi:10.1098/rsif.2018.0403

Wong ML, Chua TH, Leong CS, Khaw LT, Fornace K, Wan-Sulaiman W-Y, William T,
Drakeley C, Ferguson HM, Vythilingam I, Chua TH, Fornace K, Drakeley C, William T,
Ferguson HM. 2015. Seasonal and Spatial Dynamics of the Primary Vector of

Plasmodium knowlesi within a Major Transmission Focus in Sabah, Malaysia. *PLoS*
Negl Trop Dis **9**:e0004135. doi:http://dx.doi.org/10.1371/journal.pntd.0004135

WorldPop. 2018. Global High Resolution Population Denominators Project - Funded by The
Bill and Melinda Gates Foundation (OPP1134076).
https://dx.doi.org/10.5258/SOTON/WP00671

Zarith et al. 2021. Molecular Prevalence, Spatial Distribution and Epidemiological Risk
Factors of Plasmodium knowlesi infection among Macaca fascicularis in Peninsular
Malaysia (pre-publication).

Zhang X, Kadir KA, Quintanilla-Zariñan LF, Villano J, Houghton P, Du H, Singh B, Smith DG.
2016. Distribution and prevalence of malaria parasites among long-tailed macaques
(Macaca fascicularis) in regional populations across Southeast Asia. *Malar J* **15**:1–8.
doi:10.1186/s12936-016-1494-0

ZUNINO GABRIEL, KOWALEWSKI1 M, OKLANDER L. 2007. Habitat Fragmentation and
Population Size of the Black and Gold Howler Monkey (Alouatta caraya) in a
Semideciduous Forest in Northern Argentina. *Am J Primatol* **69**:966–975.
doi:10.1002/ajp

846