Group A *Streptococcus* Induces Lysosomal Dysfunction in THP-1 Macrophages

Scott T. Nishioka[1][†], Joshua Snipper[1][†], Jimin Lee[1], Joshua Schapiro[1], Robert Z. Zhang[1], Hyewon Abe[1], Andreas Till[2,3], Cheryl Y.M. Okumura[1]*

[1]Biology Department, Occidental College, Los Angeles, CA 90041, USA

[2]Division of Biological Sciences and The San Diego Center for Systems Biology, University of California San Diego, La Jolla, CA 92093-0688, USA

[3]University Hospital of Bonn, Bonn, Germany

[†]These authors have contributed equally to this work and share first authorship

* Correspondence:
Cheryl Okumura
okumura@oxy.edu

Keywords: Group A *Streptococcus*, streptolysin O (SLO), macrophage, lysosome, vacuolar ATPase, cathepsin B

Abstract

The human-specific bacterial pathogen Group A *Streptococcus* (GAS) is a significant cause of morbidity and mortality. Macrophages are important to control GAS upon infection, but previous data indicate that GAS can persist in macrophages. In this study, we detail the molecular mechanisms by which GAS survives in THP-1 macrophages. Our fluorescence microscopy studies demonstrate that GAS are readily phagocytosed by macrophages, but persist within phagolysosomes. These phagolysosomes are not acidified, which is in agreement with the inability of the GAS to survive in low pH environments. We find that the secreted pore-forming toxin Streptolysin O (SLO) perforates the phagolysosomal membrane, allowing leakage of not only protons, but large proteins including the lysosomal protease cathepsin B. Additionally, GAS blocks the activity of vacuolar ATPase (v-ATPase) to prevent acidification of the phagolysosome. Thus, while GAS does not inhibit fusion of the lysosome with the phagosome, it has multiple mechanisms to prevent proper phagolysosome function, allowing for persistence of the bacteria within the macrophage. This has important implications for not only the initial response, but the overall function of the macrophages and resulting pathology in GAS infection and suggests that therapies aimed at improving macrophage function may improve outcomes in GAS infection.

1 Introduction

GAS is a significant cause of morbidity and mortality worldwide, but especially in medically underserved areas where antibiotic therapy may not be promptly available (1). Infection produces wide-ranging clinical manifestations including pharyngitis, acute rheumatic fever (ARF) often leading to rheumatic heart disease, and invasive disease such as necrotizing fasciitis (1). Acting as the first line of defense against such infections, macrophages are critical for the early control and resolution of GAS infection (2,3). However, GAS has been shown to survive intracellularly in macrophages during both acute invasive soft tissue infection and asymptomatic carriage (3–5). Although GAS remains highly sensitive to beta-lactam antibiotics, the gold standard of treatment (6),

the ability of bacteria to persist in macrophages and re-emerge after antibiotic treatment poses a
therapeutic challenge (4).

As both phagocytes and antigen-presenting cells, macrophages function at the intersection of
innate and adaptive immunity and are therefore critical for identifying and defending against
infection. Macrophages engulf bacterial pathogens into phagosomes, which fuse with lysosomes that
deliver proteolytic enzymes to facilitate bacterial destruction. These enzymes have optimal
proteolytic activity under acidic conditions, and therefore lysosomal acidification is important for
macrophage bactericidal function and subsequent antigen presentation (7). Inhibition of either
lysosomal fusion with bacteria-containing phagosomes or phagolysosomal acidification is an evasion
tactic employed by several bacterial pathogens including *Mycobacterium tuberculosis*, *Legionella*
pneumophila, and *Staphylococcus aureus* (7–9). GAS may employ similar tactics to survive
intracellularly within macrophages and to evade the adaptive immune response upon host re-
infection, which may contribute to the development of pathologies such as toxic shock syndrome
(10).

GAS is a human-specific pathogen and therefore has evolved mechanisms to survive the
immune response (11). Previous data have indicated that phagocytosed GAS can prevent fusion with
destructive organelles such as azurophilic granules and lysosomes (12,13). Other reports have shown
that while GAS-containing phagosomes do fuse with lysosomes, GAS survives within macrophages
via the secretion of virulence factors such as Streptolysin O (SLO) and NADase (14,15). SLO is a
cholesterol-dependent cytolysin that oligomerizes to form large pores (~25-30 nm) within the host
cell membrane (16). As a result, SLO is hypothesized to permeabilize the phagolysosomal
membrane, permitting free flow of protons out of the phagolysosome and limiting the activity of
lysosomal proteases that facilitate intracellular killing (15,17). Interestingly, it has also been reported
that GAS is capable of preventing lysosomal acidification by preventing the recruitment of the
vacuolar ATPase (v-ATPase) (18). In the context of these data, it is unclear what mechanisms are
contributing to GAS intracellular survival in macrophages. In this paper, we used the human
monocytic cell line THP-1 differentiated into macrophages to understand the molecular mechanism
by which GAS survives in macrophages. We find that not only does GAS induce leakage of large
proteins from the phagolysosome into the cytosol, but that acidification of the phagolysosome is also
limited. This has important consequences for the survival of the bacteria within the macrophage and
underscores the need to mount a proper macrophage response in GAS infection to improve patient
outcomes.

2   Materials and Methods

2.1   Antibodies and chemicals

Antibodies to the following proteins were used in this study: EEA-1 (Abcam, ab2900), LAMP-2
(Abcam, ab25631), V_0D_1 (Abcam, ab56441), Cathepsin B (Cell Signaling Technology, D1C7Y),
LAMP-1 (Cell Signaling Technology, #9091), V_1A (Abnova, H00000523-A01), and beta-actin
(Thermo Fisher Scientific, MA5-15739). Fluorescent secondary antibodies were purchased from
Thermo Fisher Scientific. The following fluorescent probes were purchased from Thermo Fisher
Scientific: Alexa Fluor 488 Hydrazide (A10436), Oregon Green 488 Anionic, Lysine Fixable
Dextran,10,000 MW (D7171) and 70,000 MW (D7173), and LysoTracker DeepRed (L12492).
40,000 and 70,000 MW Fluorescein isothiocyanate (FITC) dextrans were purchased from Millipore
Sigma. To generate antibody-coupled beads, 0.1um Carboxylate-Modified Microspheres (Thermo
Fisher Scientific) were coupled to approximately 400mg normal pooled human serum (Thermo

Fisher Scientific) using 2% 1-Ethyl-3-(3-dimethylaminopropyl) carbodiimide hydrochloride
(QBiosciences). The reaction was quenched with 40mM ethanolamine and resuspended in PBS.
Successful coupling of the beads was assessed by fluorescence microscopy and increased
phagocytosis by THP-1 macrophages compared with uncoupled beads (data not shown). 1mM L-
leucyl L-leucine O-methyl ester (LLOMe, Cayman Chemical, #16008) was incubated with cells to
induce lysosomal damage.

2.2   Bacterial strains

Wild-type (WT) GAS strain M1T1 5448 (M1 GAS) was originally isolated from a patient with
necrotizing fasciitis and toxic shock syndrome (19). Isogenic mutant strains lacking Streptolysin O
(ΔSLO), Streptolysin S (via lack of SagA, ΔSagA) and Emm1 (ΔM1) were previously described (20–
22). Bacterial strains were cultivated in Todd-Hewitt broth (THB) at 37˚C. For all experiments,
bacteria were grown to log phase in the presence of 1:200 pooled normal human serum (Thermo
Fisher Scientific) to opsonize bacteria. The gene for monomeric WASABI (mWASABI) (23,24) was
codon-optimized for expression in bacteria and synthesized in a shuttle vector (GenScript; see
supplemental data for amino acid sequence). The gene was subcloned into pDCerm (25) and the
resulting plasmid (pWASABI) was transformed into WT M1 GAS and the ΔSLO mutant. Plasmid-
bearing strains were maintained in THB supplemented with 5ug/mL erythromycin. Heat-killed (HK)
bacteria were prepared by incubating a known concentration of bacteria at 95°C for 10 min. followed
by a 15 min. opsonization in 1:200 pooled normal human serum at room temperature.

2.3   Cell Culture

104   THP-1 cells were purchased from Sigma (cat. 88081201) and cultured in RPMI supplemented with
105   10% heat inactivated fetal bovine serum (FBS) (Corning), 2mM L-glutamine and 100 U/mL
106   penicillin/100 µg/mL streptomycin at 37°C/5% CO_2. Cells were differentiated to macrophages using
107   20nM phorbol 12-myristate 13-acetate (PMA) (MilliporeSigma) 24-48 hours prior to experiments.

2.4   Immunofluorescence

2.5×10^5 THP-1 cells were seeded on coverslips in the presence of 20nM PMA 24-48 hours prior to
experiments. For dextran leakage assays, cells were incubated in media containing 20ug/mL of the
indicated dextran or 500ug/mL 70kD dextran overnight, washed and chased in cell culture media for
2 hours prior to bacterial infection. Bacteria were combined with cells at an MOI = 10 in RPMI
supplemented with 2% FBS only (no antibiotics). At the indicated time points, cells were fixed with
4% paraformaldehyde or 3.7% formaldehyde for 10 min. at room temperature. Cells were incubated
with blocking solution (10% goat serum, 3% BSA and 0.1% Triton X-100 in PBS), then incubated
with the indicated primary antibodies for 1 hour at room temperature in block solution. Cells were
washed and incubated with the indicated secondary antibodies for 1 hour at room temperature. Cells
were washed and mounted onto slides with ProLong antifade reagent with DAPI (Thermo Fisher
Scientific). Slides were imaged on an inverted Leica TCS SP5 confocal microscope using a 63x/1.40
oil objective with 2-3x digital zoom at calibrated magnifications and recorded with LAS AF software
(Leica). Quantitation of dextran and LAMP-1 colocalization with bacteria was analyzed with
CellProfiler 3.1.9 (26) using the Otsu thresholding method. For dextran, colocalization is expressed
as the percentage of dextran-labeled lysosomes that colocalized with bacteria compared with the total
number of lysosomes. For LAMP-1, colocalization is expressed as the percentage of LAMP-1 that
colocalized with bacteria compared with the total number of bacteria. For quantitation of GFP-
expressing bacteria within phagosomes, images were coded and counted manually in a blind fashion
by at least 4 independent reviewers. For all imaging data, at least 3 independent experiments were

performed, at least 16 images containing >10 cells per image were analyzed and at least 100 cells
were counted per time point.

2.5 Verification of pH sensitivity of mWASABI in GAS

To assess the fluorescence intensity of mWASABI in live GAS, log phase cultures of bacteria were
incubated on coverslips in THB with 10mM HEPES adjusted to the indicated pH for 1 hour at 37°C.
Bacteria were fixed with 2% paraformaldehyde and analyzed by confocal microscopy. Gain settings
were adjusted to bacteria incubated in pH 7 media, and the same settings were applied to all other
conditions. Images were thresholded and the corrected total cell fluorescence (integrated density –
average background integrated density) of each bacterial cell was measured using ImageJ.

2.6 Acidification assay

10^5 THP-1 cells were seeded with 20nM PMA in 96-well black plates with clear bottoms overnight.
Cells were fed 500ug/mL 70kD FITC-dextran overnight. The following day, cells were chased with
normal cell culture media for 2 hours prior to infection. Cells were infected at an MOI of 10 with
either the indicated bacterial strain or Ab-conjugated latex beads. Fluorescence intensity at
480nm/535nm (Em/Ex) was monitored at 37°C in a fluorescent plate reader every 15 min. for 3
hours.

2.7 Bacterial survival in different pH media

145 Bacteria were incubated in THB with 10mM HEPES adjusted to the indicated pH. For bacterial
146 growth, cultures were adjusted to an OD_{600nm} of 0.1 and incubated at 37°C. OD_{600nm} was measured
147 every 15 min. For bacterial survival, 2 x 10^5 cfu log phase bacteria were added to 250ul pH-adjusted
148 culture media in 96 well plates. At the indicated time point, a 25ul aliquot was removed from the
149 culture and quantitated by plating on agar plates.

2.8 Cell fractionation and Western blot

Following a 30 min. infection period, a total of 10^7 PMA-differentiated cells were scraped and
collected in 1 mL of fractionation buffer (50 mM KCl, 90 mM K-Gluconate, 1 mM EGTA, 5 mM
MgCl2, 50 mM sucrose, 20 mM HEPES, pH 7.4, 5 mM glucose, 1X HALT phosphatase/protease
inhibitor cocktail (Thermo Fisher Scientific), 1ug/mL pepstatin (Millipore Sigma), and 1mM PMSF
(Millipore Sigma)). Cells were lysed by nitrogen cavitation equilibrated on ice for 30 min. at 400 psi.
The resulting lysates were fractionated into membrane and cytosolic fractions by centrifugation at
16,000 xg at 4˚C for 15 min. (pellet = membrane fraction, supernatant = cytosolic fraction). Protein
concentrations were measured by BCA assay (Thermo Fisher Scientific). 10ug of each sample was
run on SDS-PAGE gels and transferred to a 0.2um PVDF membrane. Blots were blocked with 5%
(w/v) non-fat dry milk in 1X Tris-buffered saline with 0.1% Tween 20 (TBST). Blots were probed
with the indicated antibodies overnight at 4˚C. Relative protein concentrations were quantified by
densitometry analysis in ImageLab v. 6.0.1 (BioRad). v-ATPase assembly was measured by
normalizing V_1A densitometry values in the membrane fraction to the V_0D_1 loading controls (in
membrane fractions) and calculating the ratio of V_1A in the sample compared with uninfected
macrophages (27).

2.9 Cathepsin B activity assay

4 x 10^6 PMA-differentiated THP-1 cells were infected at an MOI = 10 for 60 min. followed by lysis
with 30ug/mL digitonin in acetate buffer (50 mM Na-acetate pH 5.6, 150 mM NaCl, 0.5 mM EDTA)

and protease inhibitors (1X HALT phosphatase/protease inhibitor cocktail (Thermo Fisher
Scientific), 100 μM PMSF, 1 μg/mL pepstatin)) (28). The cell lysate was separated into cytosolic and
membrane fractions as described above. Cathepsin B activity in cytosolic fractions was measured
using the SensoLyte 520 Cathepsin B Assay Kit (AnaSpec) according to the manufacturer's
instructions. Relative fluorescence units were normalized to the corresponding protein concentration
for each sample and compared with the untreated cells. Data from 3 independent experiments were
combined.

2.10 v-ATPase activity assay

Cell membrane fractions containing lysosomes were prepared using 30ug/mL digitonin in acetate
buffer as described above. 10ug of each sample was equilibrated in 200ul assay buffer (10mM
HEPES, 5mM MgCl2, 125mM KCl, pH = 7.0) (29) at room temperature for 30 min. in the presence
or absence of 200nM bafilomycin (MilliporeSigma). 1mM fresh ATP in 200mM Tris pH = 7 was
added to each sample and incubated at room temperature for 2.5 hours. 25ul of the reaction was
collected and diluted in ultrapure water, and liberation of free phosphate was measured using the
Malachite Green Phosphate Assay Kit according to the manufacturer's instructions (MilliporeSigma).
Corrected absorbance values (= A_{sample} - $A_{substrate\ blank}$ - $A_{buffer\ blank}$) were compared with a phosphate
standard curve to determine phosphate concentration. v-ATPase activity was calculated as the
difference in phosphate concentration between samples incubated with and without bafilomycin.

2.11 Statistical analysis

Growth curve data was analyzed using the Growthcurver package in R (30) and comparison of the
intrinsic growth rate (r) and carrying capacity (K) was analyzed by one-way ANOVA and Tukey's
post-hoc test. All other data were analyzed using Prism v. 9.3.1 (GraphPad Software) by one-way
ANOVA with Tukey's (normally distributed data) or Kruskal-Wallis (nonparametric data) multiple
comparison tests. Outliers were assessed by the ROUT method (Q = 1%). For all data presented,
****$P<0.0001$, ***$P<0.001$, *$P<0.05$, n.s., not significant.

3   Results

3.1   Bacterial trafficking in THP-1 macrophages

We first wanted to determine the intracellular fate of GAS in THP-1 macrophages. To study
bacterial trafficking, we performed a short time course infection experiment with WT M1 GAS and
tracked phagolysosomal maturation. In agreement with other reports (13,15,31), bacteria were rapidly
engulfed by THP-1 cells within 7.5 minutes (Fig. 1A, B) and colocalized with the phagolysosomal
marker EEA-1 (Fig. 2). Interestingly, bacteria in phagosomes began colocalizing with the lysosomal
marker LAMP-1 as early as 7.5 minutes, with maximal colocalization at 30 minutes (Fig. 1A, B).
GAS remained in phagolysosomes throughout the later time points (Fig. 1B), and the number of
bacteria per cell was relatively constant throughout the experiment (Fig. 1C). These data indicated
that bacteria were trafficked normally to phagolysosomes, but bacteria were held in these
compartments without observable bacterial killing.

3.2   GAS reside in non-acidified phagolysosomes in THP-1 macrophages

Because bacteria appeared to be held in phagolysosomes, we wondered whether
phagolysosomes containing GAS were acidified. Acidification is critical for optimal activity of the
proteolytic enzymes in the lysosome (32). Lysotracker is an acidotropic dye commonly used to track

acidic compartments such as lysosomes in the cell. However, because GAS is an acid-producing
bacterial species, we tested whether Lysotracker would be an accurate indicator of lysosomal pH in
GAS-infected cells. Although Lysotracker colocalized with lysosomal markers in THP-1 cells, the
staining and uptake was inconsistent (Supp. Fig. 1A). Furthermore, bacteria were brightly stained
with Lysotracker in both infected cells and in the absence of cells (Supp. Fig. 1A-C). This made it
difficult to determine whether and to what extent phagolysosomes containing GAS were acidified.
We therefore expressed a pH-sensitive green fluorescent protein variant mWASABI (24) in GAS to
monitor phagolysosomal pH of GAS-infected cells (Fig. 2A). Green fluorescence produced by live
bacteria was appropriately quenched in low pH environments (Fig. 2A, B). Notably, pH was
distinguishable by fluorescence intensity between pH 5.0 and 4.5 (Fig. 2B).

We infected THP-1 macrophages with mWASABI-expressing GAS and performed a detailed
trafficking assay of GAS through the phagolysosomal pathway (Fig. 2C, D). Because bacteria were
tracked by mWASABI expression instead of immunofluorescent staining, we manually counted
bacteria that colocalized with phagolysosomal markers (Fig. 2C, D). We again found that GAS was
initially located in phagosomes, as indicated by EEA-1 staining, which was followed by rapid
bacterial colocalization with the mature lysosomal marker LAMP-2 within 15 minutes of infection
(Fig. 2C, D), similar to the colocalization dynamics observed with LAMP-1 staining (Fig. 1). A
significant proportion of the total intracellular bacteria were colocalized with LAMP-2 within 60
minutes post-infection (Fig. 2C, D). Although there were some fluctuations in the number of bacteria
in phagolysosomes (Fig. 2E), the number of intracellular bacteria remained relatively constant at all
time points, consistent with our initial data (Fig. 1C). This corroborated our finding that little to no
bacterial eradication or replication was occurring within this time frame (Fig. 2E). Surprisingly, we
also found that the green fluorescence signal did not diminish over time (Fig. 2C, F). Fluorescence
signal intensity of intracellular bacteria in phagolysosomes was not significantly different than that of
extracellular bacteria (Fig. 2F). As the lysosomal lumen is typically between pH 4.5–5 when
acidified (33), a pH level that our mWASABI probe can differentiate (Fig. 2A, B), these data
indicated that GAS were maintained in phagolysosomes that are not appropriately acidified.

We confirmed that GAS-infected phagosomes were not acidified with a second method using
pH-sensitive FITC-conjugated 70kD dextran particles. Acidification of the lysosomes resulted in
quenching of the fluorescence signal in uninfected and control cells infected with immunoglobulin-
conjugated beads (Fig. 2G). However, the fluorescence signal was increased in cells infected with all
tested strains of GAS, indicating no pH change (Fig. 2G). These results confirmed that GAS-infected
phagolysosomes are not properly acidified, which may account for the persistence of the bacteria in
THP-1 cells (Fig. 1C, 2E).

3.3   GAS does not tolerate lysosomal acidic conditions

The pH of the lysosomal lumen has been reported to range between pH 4.5-5 (33). Because
GAS is a lactic acid-producing bacterial species capable of acidifying its environment and possesses
systems such as the arginine deiminase (ADI) pathway that confer acid tolerance (34,35), we
wondered whether a pH similar to that found in the lysosome adversely affected bacterial growth and
survival. In growth experiments, bacteria grown in media adjusted to pH 5 or lower did not
demonstrate measurable growth (Fig. 3A). Bacteria grown in media adjusted to pH 6 grew at a
significantly slower rate and did not reach a similar maximum population size in the time frame of
our experiment compared with the unbuffered media control (Fig. 3A). Although growth rates were
slower, the ability of bacteria to grow in pH 6 media is consistent with the ability of bacteria to grow
in increasingly acidified media, as indicated by the final pH of the stationary phase cultures in

unbuffered culture media (Table 1). For bacteria grown in buffered media adjusted to pH 7, there was
no significant difference in maximum population size, but a statistically significant increase in the
intrinsic growth rate of the population compared with unbuffered media (Fig. 3A). Thus, bacterial
growth is inhibited by low pH. As expected, we found that the final pH of both buffered and
unbuffered culture medium after bacterial growth was acidic (Table 1).
260
Quantitative plating of cultures at various time points throughout the growth experiment
indicated that bacteria in pH 5 media were present, but not measurably growing (data not shown). We
therefore performed a detailed time course to monitor bacterial survival in low pH media (Fig. 3B).
Bacteria incubated in unbuffered media did not significantly grow during the time frame of the
experiment (data not shown). Similar to other studies (36,37), we found that GAS does not survive in
low pH media (Fig. 3B). Although bacteria incubated in up to pH 5.3 media could survive for a short
time, bacterial numbers declined at later time points (Fig. 3B), in agreement with pH 5 being the limit
at which bacteria can survive (35,38) and the final pH of bacteria grown in media (Table 1).
Combined, our data affirm that GAS cannot survive in the acidified lysosome. Thus, in order to
persist in the phagolysosome, GAS likely prevents its acidification.
271

3.4   Streptolysin O creates large perforations in the phagolysosome

Although GAS has several acid stress response strategies, a better tactic may be to avoid the
acidification of the phagolysosome altogether (39). The inability of bacteria to survive in low pH
environments (Fig. 3) led us to examine whether GAS infection has a mechanism to prevent
phagolysosomal acidification. Pore-forming toxins such as SLO create pores as large as 25-30nm in
size (16). Streptolysin S (SLS) can similarly perforate membranes (40). Others have shown that pore-
forming toxins, including SLO, allow escape of hydrogen ions that prevent acidification of the
lysosome (15,41). To confirm these data, we fed THP-1 macrophages fluorescent-conjugated
molecules and dextrans of various sizes to assess pore size and leakage from phagolysosomes in
GAS-infected cells. The ability to measure phagosomal leakage was confirmed using a fluorescent
dye approximately 570Da in size, a 10kD dextran and the lysosomal permeabilization agent LLOMe
(Supp. Fig. 2A). LLOMe permeabilizes lysosomes to molecules up to at least 4.4KDa, but not to
molecules greater than 10kD (28). Retention of the 10kD, but not the 570Da probe in LLOMe-treated
cells confirmed uptake of the probe in phagolysosomes and appropriate release into the cytosol upon
permeabilization (Supp. Fig. 2A). Cells loaded with probes were infected with various strains of
GAS (Fig. 4A, B). As expected, WT GAS caused probes up to 40kD in size, approximately 9nm
diameter (42), to leak from phagolysosomes (Fig. 4A, B). ΔSLO GAS-infected cells retained the
probes, while cells infected with ΔSagA GAS (lacking SLS) allowed leakage of the probes, similar to
the WT strain (Fig. 4A, B). Mutant bacteria lacking the surface protein M1 (ΔM1), but with unaltered
expression of SLO and SLS behaved similarly to WT strains (Fig. 4A, B). Heat-killed bacteria (HK)
that retain surface structure but are unable to secrete proteins such as SLO and SLS were included as
a comparison and corroborated the data from cells infected with the live GAS strains (Fig. 4A, B).
Compared with heat-killed bacteria, WT bacteria could retain probes up to 70kD in size
(approximately 13nm diameter) in agreement with the literature (42), though it is possible that
leakage of probes of this size still occurs (Fig. 4B, compare with ΔSLO). In our assays, SLS did not
significantly damage the phagolysosomal membrane, as indicated by the lack of leakage of even the
small 570Da probe (Supp. Fig. 2B). These data indicate that SLO is the primary pore-forming toxin
that perforates the phagolysosome and allows leakage of not only protons, but relatively large
molecules.
301

Because such large molecules can leak from GAS-infected phagosomes, we wondered whether
lysosomal enzymes could escape into the cytosol through SLO-mediated pores. We infected cells
with WT or ΔSLO GAS and collected membrane and cytosolic cell fractions. Lack of lysosomal
markers in the cytosolic fraction indicated little to no contamination of the cytosolic fraction with
lysosomes (Fig. 2C). The membrane fraction contained the majority of the lysosomal enzyme
cathepsin B as expected (Fig. 2C). However, we noted a small, but visible increase in the amount of
cathepsin B present in the cytosolic fraction of WT-infected cells compared with uninfected and
ΔSLO-infected cells (Fig. 2C). In order to verify that cathepsin B was present in the cytosol, we
measured cathepsin B activity in the cytosolic fraction. Cytosolic fractions from WT-infected cells
contained the highest amount of cathepsin B activity compared with uninfected cells (Fig. 2D).
Cytosolic fractions from ΔSLO bacteria incapable of leaking large proteins from the phagolysosome
(Fig. 2B) did not exhibit cathepsin B activity higher than background levels in uninfected cells (Fig.
2D). These data confirmed that SLO secreted by WT bacteria perforates the phagolysososmal
membrane and allows the leakage of large proteins into the cytosol. This loss of proteolytic enzymes
from the phagolysosome likely also contributes to the ability of GAS to persist in the phagolysosome
during infection.

3.5   GAS prevents proper function of vacuolar ATPase

Although SLO-mediated pores could cause leakage of protons from the phagolysosome,
reports in the literature suggest that the v-ATPase that is required for acidification of the lysosome is
non-functional in GAS-infected cells (18). Supporting this, we found that the phagolysosomes of
macrophages infected with ΔSLO and heat-killed bacteria were also not acidified (Fig. 2G).
Furthermore, ΔSLO bacteria are trafficked to phagolysosomes and are maintained at consistent
numbers in THP-1 cells similar to WT bacteria (Supp. Fig. 3), consistent with the idea that
phagolysosomes infected with ΔSLO bacteria were not acidified. We therefore investigated whether
v-ATPase is properly assembled and functional in GAS-infected cells. v-ATPase consists of the
membrane-bound F_0 subunit (containing the V_0D_1 protein) that must assemble with the cytosolic F_1
subunit (containing the V_1A) for proper function (27). We monitored v-ATPase assembly by tracking
movement of the V_1A subunit from the cytosolic to membrane fractions. Cells fed Ig-conjugated
beads (Ab beads) were included as a positive control (Fig. 5A, B). We observed no significant
difference in v-ATPase subunit assembly in cells infected with GAS compared to Ab-bead fed or
uninfected cells (Fig. 5A, B). This observation was consistent for cells infected with either WT or
ΔSLO GAS, as well as heat-killed bacteria (Fig. 5A, B). However, when we measured the v-ATPase
activity in membrane fractions of GAS-infected cells, we found significantly decreased v-ATPase
activity in fractions from cells infected with all strains of GAS compared with uninfected cells (Fig.
5C). These data are in agreement with the lack of acidification in GAS-infected cells (Fig. 2G).
Therefore, though the v-ATPase is properly assembled in GAS-infected cells, the activity of this
enzyme is decreased, preventing proper acidification of the phagolysosome. Thus, our data suggests
that GAS persists in macrophages by employing a multi-pronged approach to avoid the destructive
mechanisms of the phagolysosome.

4   Discussion

The human-specific pathogen GAS has co-evolved with the human immune system, and
therefore has multiple mechanisms for survival within the host (1,11). Although GAS is an
extracellular bacterial pathogen, phagocytic cells such as macrophages can readily engulf bacteria
(13,15,31). The ability of GAS to not only survive in macrophages, but to potentially alter their

function, escape downstream immune responses, and be sheltered from antibiotic treatment provides
a basis for successful persistence in humans.

In our experiments, we found that GAS was readily phagocytosed and phagosomes quickly
fused with lysosomes, but that GAS remained intact in these compartments (Fig. 1, 2). There is
conflicting evidence in the literature about the fusion of lysosomes with GAS-containing phagosomes
in macrophages, which may be due to a difference in cell origin or antibodies (13,15). In addition, we
corroborated previous data that SLO induces perforation and damage of the phagolysosomal
membrane (Fig. 4) (15,43). Although ΔSLO GAS also prevented phagolysosomal acidification (Fig.
2, 5), others have shown that ΔSLO GAS are less fit for survival in macrophages and other cells
(15,44). ΔSLO GAS is likely still susceptible to other macrophage killing mechanisms such as
reactive oxygen species or xenophagy (44). However, all data are in agreement that in macrophages,
GAS does not escape into the cytosol and remains in a membrane-bound structure (Fig. 2, Supp. Fig.
3) (13,15). In THP-1 macrophages, bacteria do not replicate in this structure, but they are also not
eradicated (Fig. 1, 2, Supp. Fig. 3) (31). This is different from what has previously been observed in
epithelial cells, where GAS escapes into the cytosol and replicates (14,43). The downstream fate of
GAS in non-functional lysosomes in THP-1 macrophages remains to be elucidated.

The pH of the lysosomal lumen has been reported to range between pH 4.5-5 (33). However,
we found that acidified medium was sufficient to kill GAS (Fig. 3). These results are surprising since
not only does GAS produce lactic acid, but previous reports suggest GAS has multiple mechanisms
for acid tolerance, including a F_0/F_1 proton pump and the arginine deiminase pathway which
produces alkali (34,35,39,45). However, there may be a limit to which these systems work, especially
within an enclosed environment such as in vitro conditions or in the phagolysosome, as demonstrated
by the threshold for bacterial survival in low pH medium (Fig. 3) (35). Additionally, GAS lacks
enzymes such as glutamate decarboxylase and urease (39), making neutralization of highly acidified
environments a less viable strategy. Therefore, two mechanisms to prevent acidification, perforating
the phagolysosomal membrane (Fig. 4) and blocking v-ATPase activity (Fig. 5) provide a means to
survive within the phagolysosomal compartment. Other bacteria such as *Legionella pneumophila*
have proteins such as SidK, which binds v-ATPase subunits to prevent activity (29). Our data
indicate that neither SLO nor any other secreted factor is responsible for interfering with v-ATPase
activity (Fig. 5c), and the identity of the responsible GAS protein(s) remains to be elucidated.

Previous data and our results indicate SLO causes leakage of lysosomal contents (Fig. 4).
These proteins include active lysosomal enzymes such as cathepsin B (Fig. 4) but could also include
bacteria proteases such as SpeB, whose presence in the cytosol could lead to NLRP3 inflammasome
activation and release of the proinflammatory cytokine IL-1β (46–50). Persistent IL-1β activation as
a result of GAS infection has been linked with ARF and rheumatic heart disease (51). Thus, besides
enabling bacterial survival and blunting the normal macrophage response, the resulting macrophage
response may lead to pathologies such as ARF that are observed during GAS infection. Furthermore,
SLO promotes macrophage apoptosis by releasing SLO from the lysosome and causing
mitochondrial damage (20,52). Loss of macrophages may also have detrimental effects on the long-
term response to GAS infection, including the inability to generate protective antibodies. Therapies
aimed at improving macrophage function may therefore improve GAS infection outcomes. Although
SLO is a promising target, our work shows that additional drugs to restore v-ATPase activity would
be necessary to restore macrophage function.

5   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

6   Author Contributions

SN, JS, AT and CO contributed to conception and design of the study. All authors performed the
experiments and analyzed data. CO wrote the first draft of the manuscript. All authors contributed to
manuscript revision, read, and approved the submitted version.

7   Funding

8   Acknowledgments

We thank Cheldon Alcantara and Ryan Hino for contributions to experiments, and Kelvin Zheng,
Jessica Kim, Stephanie Peacock, Isabel Perez, Kevin Kim, and the Fall 2021 Bio 395 GRR course for
contributions to data analysis in this paper.

9   References

1.   Sims Sanyahumbi A, Colquhoun S, Wyber R, Carapetis JR. Global Disease Burden of Group A
Streptococcus. In: Ferretti JJ, Stevens DL, Fischetti VA, editors. Streptococcus pyogenes : Basic
Biology to Clinical Manifestations [Internet]. Oklahoma City (OK): University of Oklahoma
Health Sciences Center; 2016 [cited 2022 Apr 11]. Available from:
http://www.ncbi.nlm.nih.gov/books/NBK333415/

2.   Goldmann O, Rohde M, Chhatwal GS, Medina E. Role of Macrophages in Host Resistance to
Group A Streptococci. Infect Immun. 2004 May;72(5):2956–63.

3.   Mishalian I, Ordan M, Peled A, Maly A, Eichenbaum MB, Ravins M, et al. Recruited
macrophages control dissemination of group A Streptococcus from infected soft tissues. J
Immunol Baltim Md 1950. 2011 Dec 1;187(11):6022–31.

4.   Thulin P, Johansson L, Low DE, Gan BS, Kotb M, McGeer A, et al. Viable group A
streptococci in macrophages during acute soft tissue infection. PLoS Med. 2006 Mar;3(3):e53.

5.   Österlund A, Popa R, Nikkilä T, Scheynius A, Engstrand L. Intracellular Reservoir of
Streptococcus pyogenes In Vivo: A Possible Explanation for Recurrent Pharyngotonsillitis. The
Laryngoscope. 1997;107(5):640–7.

6.   Johnson AF, LaRock CN. Antibiotic Treatment, Mechanisms for Failure, and Adjunctive
Therapies for Infections by Group A Streptococcus. Front Microbiol. 2021;12:760255.

7.   Flannagan RS, Heit B, Heinrichs DE. Antimicrobial Mechanisms of Macrophages and the
Immune Evasion Strategies of Staphylococcus aureus. Pathogens. 2015 Dec;4(4):826–68.

8.   Carranza C, Chavez-Galan L. Several Routes to the Same Destination: Inhibition of
Phagosome-Lysosome Fusion by Mycobacterium tuberculosis. Am J Med Sci. 2019
Mar;357(3):184–94.

9.   Misch EA. Legionella: virulence factors and host response. Curr Opin Infect Dis. 2016
Jun;29(3):280–6.

10. Ganem MB, De Marzi MC, Fernández-Lynch MJ, Jancic C, Vermeulen M, Geffner J, et al.
Uptake and intracellular trafficking of superantigens in dendritic cells. PloS One.
2013;8(6):e66244.

11. Siemens N, Snäll J, Svensson M, Norrby-Teglund A. Pathogenic Mechanisms of Streptococcal
Necrotizing Soft Tissue Infections. Adv Exp Med Biol. 2020;1294:127–50.

12. Staali L, Bauer S, Mörgelin M, Björck L, Tapper H. Streptococcus pyogenes bacteria modulate
membrane traffic in human neutrophils and selectively inhibit azurophilic granule fusion with
phagosomes. Cell Microbiol. 2006 Apr;8(4):690–703.

13. Hertzén E, Johansson L, Wallin R, Schmidt H, Kroll M, Rehn AP, et al. M1 protein-dependent
intracellular trafficking promotes persistence and replication of Streptococcus pyogenes in
macrophages. J Innate Immun. 2010;2(6):534–45.

14. Walker MJ, Barnett TC, McArthur JD, Cole JN, Gillen CM, Henningham A, et al. Disease
Manifestations and Pathogenic Mechanisms of Group A Streptococcus. Clin Microbiol Rev.
2014 Apr 1;27(2):264–301.

15. Bastiat-Sempe B, Love JF, Lomayesva N, Wessels MR. Streptolysin O and NAD-
glycohydrolase prevent phagolysosome acidification and promote group A Streptococcus
survival in macrophages. mBio. 2014 Sep 16;5(5):e01690-01614.

16. Bhakdi S, Tranum-Jensen J, Sziegoleit A. Mechanism of membrane damage by streptolysin-O.
Infect Immun. 1985 Jan;47(1):52–60.

17. Håkansson A, Bentley CC, Shakhnovic EA, Wessels MR. Cytolysin-dependent evasion of
lysosomal killing. Proc Natl Acad Sci U S A. 2005 Apr 5;102(14):5192–7.

18. Nordenfelt P, Grinstein S, Björck L, Tapper H. V-ATPase-mediated phagosomal acidification is
impaired by Streptococcus pyogenes through Mga-regulated surface proteins. Microbes Infect.
2012 Nov;14(14):1319–29.

19. Chatellier S, Ihendyane N, Kansal RG, Khambaty F, Basma H, Norrby-Teglund A, et al.
Genetic relatedness and superantigen expression in group A streptococcus serotype M1 isolates
from patients with severe and nonsevere invasive diseases. Infect Immun. 2000 Jun;68(6):3523–
34.

20. Timmer AM, Timmer JC, Pence MA, Hsu LC, Ghochani M, Frey TG, et al. Streptolysin O
promotes group A Streptococcus immune evasion by accelerated macrophage apoptosis. J Biol
Chem. 2009 Jan 9;284(2):862–71.

21. Lauth X, von Köckritz-Blickwede M, McNamara CW, Myskowski S, Zinkernagel AS, Beall B,
et al. M1 protein allows Group A streptococcal survival in phagocyte extracellular traps through
cathelicidin inhibition. J Innate Immun. 2009;1(3):202–14.

22. Datta V, Myskowski SM, Kwinn LA, Chiem DN, Varki N, Kansal RG, et al. Mutational
analysis of the group A streptococcal operon encoding streptolysin S and its virulence role in
invasive infection. Mol Microbiol. 2005 May;56(3):681–95.

23. Ai H wang, Olenych SG, Wong P, Davidson MW, Campbell RE. Hue-shifted monomeric
variants of Clavularia cyan fluorescent protein: identification of the molecular determinants of
color and applications in fluorescence imaging. BMC Biol. 2008 Mar 6;6:13.

24. Zhou C, Zhong W, Zhou J, Sheng F, Fang Z, Wei Y, et al. Monitoring autophagic flux by an
improved tandem fluorescent-tagged LC3 (mTagRFP-mWasabi-LC3) reveals that high-dose
rapamycin impairs autophagic flux in cancer cells. Autophagy. 2012 Aug;8(8):1215–26.

25. Jeng A, Sakota V, Li Z, Datta V, Beall B, Nizet V. Molecular genetic analysis of a group A
Streptococcus operon encoding serum opacity factor and a novel fibronectin-binding protein,
SfbX. J Bacteriol. 2003 Feb;185(4):1208–17.

26. McQuin C, Goodman A, Chernyshev V, Kamentsky L, Cimini BA, Karhohs KW, et al.
CellProfiler 3.0: Next-generation image processing for biology. PLoS Biol. 2018
Jul;16(7):e2005970.

27. McGuire CM, Forgac M. Glucose starvation increases V-ATPase assembly and activity in
mammalian cells through AMP kinase and phosphatidylinositide 3-kinase/Akt signaling. J Biol
Chem. 2018 Jun 8;293(23):9113–23.

28. Repnik U, Borg Distefano M, Speth MT, Ng MYW, Progida C, Hoflack B, et al. L-leucyl-L-
leucine methyl ester does not release cysteine cathepsins to the cytosol but inactivates them in
transiently permeabilized lysosomes. J Cell Sci. 2017 Sep 15;130(18):3124–40.

29. Xu L, Shen X, Bryan A, Banga S, Swanson MS, Luo ZQ. Inhibition of host vacuolar H+-
ATPase activity by a Legionella pneumophila effector. PLoS Pathog. 2010 Mar
19;6(3):e1000822.

30. Sprouffske K, Wagner A. Growthcurver: an R package for obtaining interpretable metrics from
microbial growth curves. BMC Bioinformatics. 2016 Apr 19;17:172.

31. O'Neill AM, Thurston TLM, Holden DW. Cytosolic Replication of Group A Streptococcus in
Human Macrophages. mBio. 2016 Apr 12;7(2):e00020-00016.

32. Bright NA, Davis LJ, Luzio JP. Endolysosomes Are the Principal Intracellular Sites of Acid
Hydrolase Activity. Curr Biol CB. 2016 Sep 12;26(17):2233–45.

33. Canton J, Grinstein S. Chapter 5 - Measuring lysosomal pH by fluorescence microscopy. In:
Platt F, Platt N, editors. Methods in Cell Biology [Internet]. Academic Press; 2015 [cited 2021
Jul 1]. p. 85–99. (Lysosomes and Lysosomal Diseases; vol. 126). Available from:
https://www.sciencedirect.com/science/article/pii/S0091679X14000223

34.   Degnan BA, Fontaine MC, Doebereiner AH, Lee JJ, Mastroeni P, Dougan G, et al.
499   Characterization of an isogenic mutant of Streptococcus pyogenes Manfredo lacking the ability
500   to make streptococcal acid glycoprotein. Infect Immun. 2000 May;68(5):2441–8.

501   35.   Cusumano ZT, Caparon MG. Citrulline protects Streptococcus pyogenes from acid stress using
502   the arginine deiminase pathway and the F1Fo-ATPase. J Bacteriol. 2015 Apr;197(7):1288–96.

503   36.   Maudsdotter L, Jonsson H, Roos S, Jonsson AB. Lactobacilli reduce cell cytotoxicity caused by
504   Streptococcus pyogenes by producing lactic acid that degrades the toxic component lipoteichoic
505   acid. Antimicrob Agents Chemother. 2011 Apr;55(4):1622–8.

506   37.   Daglia M, Papetti A, Grisoli P, Aceti C, Dacarro C, Gazzani G. Antibacterial activity of red and
507   white wine against oral streptococci. J Agric Food Chem. 2007 Jun 27;55(13):5038–42.

508   38.   Jin H, Agarwal S, Agarwal S, Pancholi V. Surface export of GAPDH/SDH, a glycolytic
509   enzyme, is essential for Streptococcus pyogenes virulence. mBio. 2011;2(3):e00068-00011.

510   39.   Cotter PD, Hill C. Surviving the acid test: responses of gram-positive bacteria to low pH.
511   Microbiol Mol Biol Rev MMBR. 2003 Sep;67(3):429–53, table of contents.

512   40.   Molloy EM, Cotter PD, Hill C, Mitchell DA, Ross RP. Streptolysin S-like virulence factors: the
513   continuing sagA. Nat Rev Microbiol. 2011 Sep;9(9):670–81.

514   41.   Shaughnessy LM, Hoppe AD, Christensen KA, Swanson JA. Membrane perforations inhibit
515   lysosome fusion by altering pH and calcium in Listeria monocytogenes vacuoles. Cell
516   Microbiol. 2006 May;8(5):781–92.

517   42.   Ambati J, Canakis CS, Miller JW, Gragoudas ES, Edwards A, Weissgold DJ, et al. Diffusion of
518   high molecular weight compounds through sclera. Invest Ophthalmol Vis Sci. 2000
519   Apr;41(5):1181–5.

520   43.   Cheng YL, Wu YW, Kuo CF, Lu SL, Liu FT, Anderson R, et al. Galectin-3 Inhibits Galectin-
521   8/Parkin-Mediated Ubiquitination of Group A Streptococcus. mBio. 2017 Jul 25;8(4):e00899-
522   17.

523   44.   O'Seaghdha M, Wessels MR. Streptolysin O and its co-toxin NAD-glycohydrolase protect
524   group A Streptococcus from Xenophagic killing. PLoS Pathog. 2013;9(6):e1003394.

525   45.   Cusumano ZT, Watson ME, Caparon MG. Streptococcus pyogenes arginine and citrulline
526   catabolism promotes infection and modulates innate immunity. Infect Immun. 2014
527   Jan;82(1):233–42.

528   46.   Keyel PA, Roth R, Yokoyama WM, Heuser JE, Salter RD. Reduction of streptolysin O (SLO)
529   pore-forming activity enhances inflammasome activation. Toxins. 2013 Jun 6;5(6):1105–18.

530   47.   Chu J, Thomas LM, Watkins SC, Franchi L, Núñez G, Salter RD. Cholesterol-dependent
531   cytolysins induce rapid release of mature IL-1beta from murine macrophages in a NLRP3
532   inflammasome and cathepsin B-dependent manner. J Leukoc Biol. 2009 Nov;86(5):1227–38.

48. Richter J, Monteleone MM, Cork AJ, Barnett TC, Nizet V, Brouwer S, et al. Streptolysins are
the primary inflammasome activators in macrophages during Streptococcus pyogenes infection.
Immunol Cell Biol. 2021 Nov;99(10):1040–52.

49. LaRock DL, Russell R, Johnson AF, Wilde S, LaRock CN. Group A Streptococcus Infection of
the Nasopharynx Requires Proinflammatory Signaling through the Interleukin-1 Receptor.
Infect Immun. 2020 Sep 18;88(10):e00356-20.

50. Campden RI, Zhang Y. The role of lysosomal cysteine cathepsins in NLRP3 inflammasome
activation. Arch Biochem Biophys. 2019 Jul 30;670:32–42.

51. Kim ML, Martin WJ, Minigo G, Keeble JL, Garnham AL, Pacini G, et al. Dysregulated IL-1β-
GM-CSF Axis in Acute Rheumatic Fever That Is Limited by Hydroxychloroquine. Circulation.
2018 Dec 4;138(23):2648–61.

52. Cortés G, Wessels MR. Inhibition of dendritic cell maturation by group A Streptococcus. J
Infect Dis. 2009 Oct 1;200(7):1152–61.

10   Tables

Table 1: Final pH of media after bacterial growth, mean ± SD

starting pH	4.0	5.0	6.0	7.0	media (6.6)
ending pH	4.10 ± 0.02	5.07 ± 0.02	4.99 ± 0.01	5.23 ± 0.08	5.63 ± 0.06

11   Figure legends

Figure 1: GAS persists in phagolysosomes of THP-1 cells. THP-1 cells were infected with GAS
for the indicated times, fixed and probed with anti-LAMP-1 (green) and anti-human IgG (bacteria,
red) antibodies. **(A)** Representative image of LAMP-1 colocalized with GAS in THP-1 cells at 60
min. post-infection. Arrowheads indicate examples of colocalization. **(B)** Quantitation of LAMP-1
positive bacteria at the indicated time points. **(C)** Average number of bacteria per cell at the indicated
time points. For all graphs, data from three independent experiments were combined and results are
given as mean ± 95% CI. Data were analyzed by one-way ANOVA with Tukey's multiple
comparisons test.

Figure 2: GAS persists in non-acidified phagolysosomes of THP-1 cells. (A) GAS expressing
mWASABI were incubated at the indicated pH. Representative green fluorescence (mWASABI) and
brightfield images are shown. **(B)** Corrected total cell fluorescence of green fluorescence signal from
bacteria expressing mWASABI incubated at the indicated pH for 1 hour. **(C)** Representative
fluorescence microscopy images of GAS-infected THP-1 cells at 7.5 and 60 min. post-infection.
Arrowheads denote examples of bacteria encapsulated in early phagosomes (EEA-1, magenta),
arrows denote examples of bacteria encapsulated in phagolysosomes (LAMP-2, red). Fluorescence

images merged with brightfield images are shown in the bottom row. All images were taken with a
63x objective with 2x digital zoom. **(D)** Quantitation of bacteria colocalized with phagosomes (EEA-
1) or phagolysosomes (LAMP-2) at the indicated time points. **(E)** Average number of bacteria per
cell at the indicated time points. **(F)** Violin plot of fluorescence signal intensity from intracellular
(clear bars) or extracellular (shaded bars) bacteria expressing mWASABI. **(G)** Fluorescence signal
from cells fed 70kD FITC dextran and infected with immunoglobulin-conjugated beads (Ab beads),
the indicated GAS strain or heat-killed (HK) bacteria. Significance indicates difference from Ab
beads (positive control). Data from at least three independent experiments were combined. For (D)
and (E), data from 5 individual counters of three independent experiments were combined. Results
are given as mean ± 95% CI and analyzed by one-way ANOVA with Kruskal-Wallis multiple
comparison test.

Figure 3: GAS does not survive in low pH environments. (A) Growth curves of bacteria grown in
buffered media or unbuffered bacterial media (THB) at the indicated pH. Results are given as mean ±
95% CI. Statistics in (A) indicate differences in intrinsic growth rate compared with bacteria grown
in unbuffered media. **(B)** Bacterial survival in buffered media at the indicated pH compared with
survival of bacteria in unbuffered bacterial media (THB). Results are given as mean ± SEM.
Statistics in (B) are shown only for pH = 4.9 and 5.3 media and indicate differences in bacterial
survival at the corresponding time point compared with bacteria in unbuffered media. Differences for
bacteria grown in pH = 4.7 or lower media were statistically significant compared with unbuffered
media at all time points after 30 min. For each experiment, samples were prepared in triplicate and
data from at least three independent experiments were combined.

Figure 4: SLO induces phagolysosomal perforation and protein leakage. (A) Representative
images of the 40kD dextran probe (green) with indicated bacterial strains (magenta). Arrowheads
indicate examples of colocalization. All images were taken with a 63x objective with 2x digital
zoom. **(B)** Quantitation of colocalization of indicated bacterial strains with 10, 40 and 70kD dextran
probes. Results are given as mean ± 95% CI and analyzed by one-way ANOVA with Tukey's
multiple comparison test. **(C)** Cells were infected with the indicated bacterial strains, fractionated
into membrane (left) and cytosolic (right) fractions, and probed with the indicated antibodies:
LAMP-2 (lysosomal marker; loading control for membrane fraction), GAPDH (cytosolic marker;
loading control for cytosolic fraction) and cathepsin B (lysosomal enzyme). **(D)** Cathepsin B activity
was measured in cytosolic cell fractions collected from cells infected with the indicated strains. Data
in (B) and (D) are combined from at least three independent experiments.

Figure 5: GAS infection limits v-ATPase activity. (A) Cells were infected with immunoglobulin-
conjugated beads (Ab beads, control), the indicated GAS strain or heat-killed (HK) bacteria and
fractionated into membrane (M) and cytosolic (C) fractions. Fractions were probed with antibodies to
V_1A, V_0D or GAPDH proteins. Representative data are shown. **(B)** Quantitation of data shown in
(A). V_1A band intensities were quantified and normalized to V_0D (membrane) or GAPDH (cytosol).
Relative assembly was calculated as the amount of V_1A in the membrane fraction of the indicated
infected sample compared with the uninfected sample for each experiment. **(C)** v-ATPase activity
assay in membrane fractions from cells infected with the indicated GAS strain. Data in (B) and (C)
are combined from at least three independent experiments and results are given as mean ± 95% CI
and analyzed by one-way ANOVA with Tukey's multiple comparison test.

Supplementary Material

1   Supplementary Data

mWASABI amino acid sequence:

MVSKGEETTMGVIKPDMKIKLKMEGNVNGHAFVIEGEGEGKPYDGTNTINLEVKEGAPLPF
SYDILTTAFSYGNRAFTKYPDDIPNYFKQSFPEGYSWERTMTFEDKGIVKVKSDISMEEDSFI
YEIHLKGENFPPNGPVMQKETTGWDASTERMYVRDGVLKGDVKMKLLLEGGGHHRVDFK
TIYRAKKAVKLPDYHFVDHRIEILNHDKDYNKVTVYEIAVARNSTDGMDELYK

2 Supplementary Figures

Supplemental Figure 1: Lysotracker detects GAS. Cells were loaded with 100nM LysoTracker Deep Red (Thermo Fisher Scientific) prior to infection and infected with GAS for 30 min. Cells were fixed and stained with LAMP-2 (lysosome) and anti-human IgG (bacteria) antibodies as indicated: **(A)** LysoTracker (blue), LAMP-2 (red), and GAS (green). **(B)** Lysotracker (red) and GAS (green). **(C)** Bacteria alone incubated with 100nM Lysotracker (red). All images were taken with a 63x objective with 2x digital zoom.

Supplemental Figure 2: Fluorescent probes appropriately monitor phagolysosomal leakage.
Cells were loaded with Alexa-fluor 488 dye (570 Da) or 10kD Oregon Green dextran (10kDa) for 24 hrs., then either untreated, treated with 1mM LLOMe or infected with the indicated GAS strain. **(A)** Representative images of fluorescent probe-loaded cells in the presence or absence of LLOMe. All images were taken with a 63x objective with 2x digital zoom. **(B)** Quantitation of colocalization of Alexa-fluor 488 (570Da) with indicated bacterial strains. Data from three independent experiments were combined and are given as mean ± 95% CI.

Supplemental Figure 3: ΔSLO GAS persist in THP-1 phagolysosomes. (A) Representative fluorescence microscopy images of ΔSLO-infected THP-1 cells at 7.5 and 60 min. post-infection. Arrowheads denote examples of ΔSLO bacteria (green) encapsulated in early phagosomes (EEA-1, magenta) or in phagolysosomes (LAMP-2, red). All images were taken with a 63x objective with 2x digital zoom. **(B)** Quantitation of bacteria in intracellular compartments. **(C)** Average number of bacteria per cell at the indicated time points. For B and C, data from at least 5 individual counters of three independent experiments were combined. Results are given as mean ± 95% CI and analyzed by one-way ANOVA with Kruskal-Wallis multiple comparison test.