Single-cell characterization of human GBM reveals regional differences in tumor-infiltrating leukocyte activation

Philip Schmassmann[1,*], Julien Roux[2,3], Steffen Dettling[4], Sabrina Hogan[1], Tala Shekarian[1], Tomás A. Martins[1], Marie-Françoise Ritz[1], Sylvia Herter[5], Marina Bacac[5], and Gregor Hutter[1,6,*]

[1]Brain Tumor Immunotherapy Lab, Department of Biomedicine, University of Basel, Basel, Switzerland

[2]Bioinformatics Core Facility, Department of Biomedicine, University of Basel, Basel, Switzerland

[3]Swiss Institute of Bioinformatics, Basel, Switzerland

[4]Roche Pharmaceutical Research and Early Development, Roche Innovation Center Munich, Penzberg, Germany

[5]Roche Pharmaceutical Research and Early Development, Roche Innovation Center Zürich, Schlieren, Switzerland

[6]Department of Neurosurgery, University Hospital Basel, Basel, Switzerland

[*]Correspondence: p.schmassmann@unibas.ch, gregor.hutter@usb.ch

Abstract

20
Glioblastoma (GBM) harbors a highly immunosuppressive tumor microenvironment
(TME) which influences glioma growth. Major efforts have been undertaken to describe the
TME on a single-cell level. However, human data on regional differences within the TME re-
main scarce. Here, we performed high-depth single-cell RNA sequencing (scRNAseq) on
paired biopsies from the tumor center, peripheral infiltration zone and blood of five primary
GBM patients. Through analysis of > 45'000 cells, we revealed a regionally distinct transcrip-
tion profile of microglia (MG) and monocyte-derived macrophages (MdMs) and an impaired
activation signature in the tumor-peripheral cytotoxic-cell compartment. Comparing tumor-in-
filtrating CD8$^+$ T cells with circulating cells identified CX3CR1^{high} and CX3CR1^{int} CD8$^+$ T cells
with effector and memory phenotype, respectively, enriched in blood but absent in the TME.
Tumor CD8$^+$ T cells displayed a tissue-resident memory phenotype with dysfunctional features.
Our analysis provides a regionally resolved mapping of transcriptional states in GBM-associ-
ated leukocytes, serving as an additional asset in the effort towards novel therapeutic strategies
to combat this fatal disease.

Introduction

Glioblastoma (GBM) is a fatal disease without effective long-term treatment options. The current standard of care consists of tumor resection followed by adjuvant chemoradiotherapy resulting in a median overall survival of only 14 months [1]. One of the hallmarks in GBM progression is the high rate of neovascularization. The GBM-induced aberrant vessels not only nourish glioma cells, but also provide a specialized niche for tumor-associated stromal and immune cells such as monocyte-derived macrophages (MdMs), yolk sac-derived microglia (MG) (together termed glioma-associated macrophages/microglia, GAMs), and peripheral adaptive immune cells. This immune tumor microenvironment (iTME) paradoxically acts in an immunosuppressive manner and promotes tumor progression [2]. For example, clinical trials of systemic T cell checkpoint blockade showed only disappointing results [3, 4], which was attributed in part to the immunosuppressive components of the GBM iTME. The origin of GAMs, infiltration of peripherally derived macrophages across the blood-brain-barrier (BBB) or recruitment of tissue-resident MG to the tumor site, as well as their contribution to gliomagenesis are studied intensively [2, 5-7]. Hence, major efforts have been undertaken to describe the GBM iTME on a single cell level [5, 6, 8], or dissect the composition and changes upon disease stages, recurrence and immunotherapy specifically within the GAM compartment [9-11]. However, human data on the composition of the iTME in different tumor regions (contrast enhancing tumor center versus peripheral infiltration zone) remain scarce [12, 13].

To study the region-dependent cellular diversity within individual GBMs, we performed single-cell RNA sequencing (scRNA-seq) on patient-matched biopsies from the tumor center and the peripheral infiltration zone of five primary GBM patients. Additionally, peripheral blood mononuclear cells (PBMC) of the same patients were included to explore the transcriptional changes occurring during tumor infiltration of circulating immune cells.

Our analysis revealed a regionally distinct transcription profile of MG and MdMs and an impaired activation signature in the tumor-peripheral cytotoxic-cell compartment. Comparing tumor-infiltrating CD8$^+$ T cells with PBMC-derived, identified CX3CR1^{high} and CX3CR1^{int} CD8$^+$ T cells with effector and memory phenotype, respectively, enriched in blood but absent in the iTME. Tumor CD8$^+$ T cells displayed features of tissue-resident memory T cells and were characterized by an exhaustion phenotype. This work provides a regionally-resolved map of transcriptional states in glioma-associated cell types complemented by patient-matched PBMCs, dissecting the composition and molecular diversity of the iTME in GBM.

     Results

scRNA-seq analysis of paired tumor center, periphery and PBMC samples.

Fresh, neurosurgically resected tissue from five primary, treatment naïve GBM patients

were harvested (Figure 1A, Supplementary File 1). According to the 2021 WHO Classification

of Tumors of the Central Nervous System [14], in which the term glioblastoma designates only

IDH-wildtype grade 4 tumors, we will hence use the term grade 4 glioma, as we included as

well IDH-mutant grade 4 tumors (Supplementary File 1). The tumor center was defined as con-

trast enhancing, whereas the tumor periphery was defined as T2 hyperintense by magnetic res-

onance imaging (MRI)-guided, navigated surgical resection (Figure 1B). Increased cellular

density of the center vs. periphery samples was confirmed by nuclear DAPI staining on matched

histological micrographs of the resected tissue specimens used for scRNA-seq (Figure 1C). As

outlined in Figure 1A, we separately processed patient tumor and blood samples and enriched

them for immune cells by fluorescence-activated cell sorting (FACS) (Figure 1 – figure sup-

plement 1A and 1B). The three samples per patient (center, periphery and PBMC) were loaded

on different wells of a 10x Genomics Chromium system for a targeted recovery of 10,000 cells.

Due to technical issues cells from the center sample of patient BTB 609 could not be collected.

In total we analyzed 45,466 cells that passed initial quality control and filtering, com-

prising 8,254 cells from tumor center, 5,954 cells from tumor periphery and 31,258 PBMCs,

with 6,354 to 10,957 cells per patient (Supplementary File 2; Figure 1 – figure supplement 1C-

1E, Supplementary File 3). All cells were projected onto a two dimensions *t*-distributed sto-

chastic neighbor embedding (tSNE) [15]. As we observed a good overlap of cells across patients

for most of the dataset (see Methods) (Figure 1 – figure supplement 2B – 2F), we chose not to

perform any correction for patient-specific effects. Using hierarchical clustering, the cells were

partitioned into clusters (Figure 2 – figure supplement 1A, 1B) which were then annotated into

nine distinct cell types for the immune subset, including two transcriptionally distinct MG sub-

sets (MG_1 and MG_2) and four cell types for the CD45 negative subset (Figure 2A; Figure 2

– figure supplement 1C, 1D; Supplementary File 3).

In more detail, our annotation strategy made use of the relative expression patterns of

known marker genes (Figure 2B), and of cluster-specific genes (Figure 2 – figure supplement

2). Additionally, unbiased cell-type annotation using whole-transcriptome comparisons to ref-

erence bulk and single-cell datasets was performed with the *SingleR* package. We used as ref-

erence (i) a public bulk RNA-seq dataset of sorted immune cell types from human PBMC sam-

ples [16], which helped in identifying the major immune cell lineages: B cells, CD4$^+$ and CD8$^+$

T cells, dendritic cells (DCs), monocytes and NK cells (Figure 2 – figure supplement 1E). (ii)

A bulk RNA-seq dataset of sorted immune cell types from the TME of human gliomas [5] was helpful to separate GBM-associated CD4$^+$ and CD8$^+$ T cells, as well as microglia and MdMs (Figure 2 – figure supplement 1F). (iii) A 10X genomics scRNA-seq dataset of the innate immune TME of seven newly diagnosed GBM patients [9] confirmed the annotation of the innate immune subset (Figure 2 – figure supplement 1I), which was also supported by a microglia and a macrophage signature scores defined using gene lists obtained from [7] (Figure 2 – figure supplement 1J, 1K). (v) Finally a Smartseq2 scRNA-seq dataset of *IDH1^{wt}* tumors [17] was used to characterize the CD45^{neg} population (Figure 2 – figure supplement 1N).

To perform a differential expression analysis between tumor sites, we stratified the analysis by cell type to decrease the influence of differential abundance patterns (analyzed separately, see below). A principal component analysis (PCA) on the pseudo-bulk aggregated samples (see Methods) confirmed that the major source of variation was due to differences between lymphoid, MG/myeloid cells, and CD45$^-$ cells (PCs 1 and 2; Figure 2C). There was no clear association to the *IDH1* status of patients on these or deeper components. However, our study was neither designed nor powered to find regional signatures within the iTME depending on *IDH1* status, but rather to identify common transcriptional differences within the iTME between tumor center, periphery and PBMC of *IDH1^{wt}* and *IDH1^{mut}* grade 4 glioma.

Differential abundance analysis between tumor center and periphery did not reveal significant changes, but some populations displayed suggestive differences: the abundance of both CD4$^+$ T cell and monocytic clusters increased in periphery compared to center, while the abundance of MdM clusters decreased (Figure 2D). This has been observed by others as well [12, 18]. Testing the abundance at a finer scale with *miloR* provided us interesting insights. Indeed, within the innate immune population (Figure 2E) we confirmed the decreased abundance of MdMs in the tumor periphery and noticed an increased abundance of the MG_1 subset of MG in the tumor periphery. (Figure 2F, 2G).

MG display regionally resolved transcriptional profiles that differ from those of DCs and MdMs.

Differential expression analysis between MG from tumor center and periphery revealed downregulation of inflammatory-related genes in the peripheral MG (*FCGBP* and *CCL20*), downregulation of genes associated with canonical interferon (IFN) responses (*IFI6, IFI27, STAT1, ISG15)*, and cell proliferation (*STMN1*) as well as downregulation of scavenger receptor gene *CD163* (Figure 3A and Supplementary File 4)*. The latter was shown to have a potential

role in the phagocytic response of MG to beta-amyloid depositions identified in single-nucleus RNA-seq of postmortem human brain[19].

Among many upregulated genes whose function is not yet known, we found Inhibitor of DNA-Binding 1, also known as Inhibitor of Differentiation 1 (*ID1*) to be increased in the peripheral MG (Figure 3A). ID1 is well described in GBM progression, treatment resistance and glioma stem cell biology [20]. However, new evidence has emerged, linking *ID1* to suppression of the anti-tumor immune response in the myeloid compartment and promoting tumor progression [21].

To further explore the underlying biological processes differing between MG in the two compartments, we conducted a gene set enrichment analysis (GSEA) on the results of the differential expression analysis using Gene Ontology (GO) database (Biological Processes). This revealed overall a significant downregulation of GO categories involved in antigen processing and presentation via MHC-I and MHC-II in the peripheral MG relative to the center MG, as well as downregulation of amino acid metabolism and TNF-α signaling pathway (Figure 3B), which supported the conclusion of an impaired activation state of peripheral MG. When comparing the transcriptional profiles of the other innate immune phagocytic populations, DCs and MdMs, we observed for both cell types upregulation of pro-inflammatory related gene sets in the tumor periphery compared to tumor center (Figure 3C, 3D, Figure 3 - figure supplement 1A, 1B and Supplementary File 4). Within DCs, the induced pro-inflammatory phenotype was also observed when comparing DCs from tumor periphery to PBMC-derived ones (Figure 3 -figure supplement 1C, 1D and Supplementary File 6). Taken together, DCs and MdMs seem to have a proinflammatory phenotype in the glioma periphery as opposed to MG. However, they are less abundant there, at least as far as MdMs are concerned (Figure 2E-2G).

The iTME of grade 4 glioma harbors two transcriptionally distinct MG subpopulations.

Unsupervised hierarchical sub-clustering of the MG population revealed two transcriptionally distinct iTME MG subsets, which we termed MG_1 and MG_2, respectively (Figure 3E, Figure 3 – figure supplement 1E and Supplementary File 5). The MG_1 cluster was highly enriched for the activator protein-1 (AP-1) family of transcription factors including *FOS, FOSB, JUN, JUNB, MAF* and *MAFB* (Figure 3F and Supplementary File 5), which convey a surveilling phenotype to adult MG, but are also involved in numerous processes including cell growth, differentiation and immune activation [22]. Specifically, *FOSB* gene products were implicated in the excitotoxic MG activation by regulating complement C5a receptor expression

[23]. Yet, concomitant upregulation of anti-inflammatory Krüppel-like factor 2 (*KLF2*) [24]

and Dual Specificity Protein Phosphatase 1 (*DUSP1*), an inhibitor of innate inflammation by

negatively regulating the mitogen-activated protein kinase (MAPK) pathway [25], together

with increased expression of anti-proliferative genes like *RHOB, BTG1* and *BTG2* paint a more

complex picture of these cells. Particularly, *BTG1* was identified as an activation-induced apop-

totic sensitizer in MG after exposure to inflammatory stimuli [26], serving as an autoregulatory

mechanism and possibly hinting towards an exhausted state in these MG_1 cells. GSEA for

differences between MG_1 and MG_2 clusters using the MSigDB Hallmark collection of major

biological pathways [27] revealed downregulation of many MG effector functions in the MG_1

population including (1) inflammation ("Complement", "Allograft Rejection", "Reactive Oxy-

gen Species Pathway"), (2) immune cell activation ("IFN-α Response", "IFN-γ Response",

"IL6 JAK STAT3 Signaling") and (3) immunometabolism ("Fatty Acid Metabolism", Oxida-

tive Phosphorylation", "Glycolysis") (Figure 3G). As we examined the expression of previ-

ously described reactivity markers of MG including *C1QA, FCGR1A, CD14, HLA-DRA*,

TREM2 and Ferritin (*FTH1*) [28-31] (Figure 3H and Figure 3 – figure supplement 1F), and

established MG homeostatic genes like *CX3CR1, HEXB* and *SPI1* (PU.1) (Supplementary File

5), we noted a reduced expression of these genes in the MG_1 cluster compared to MG_2 cells,

while the anti-inflammatory transcription factors *NR4A1* [32] *NR4A2* [33] were highly upreg-

ulated (Figure 3 – figure supplement 1G). Additionally, when comparing the distribution of

MG subsets between sites among total MG, we observed an increased abundance of MG_1

cells in the tumor periphery, which might at least partially explain the previously observed

impaired activation state of total peripheral MG (Figure 3 – figure supplement 1H). To rule out

this explanation we stratified the differential expression analysis between tumor periphery and

center by MG subtype and observed similar genes downregulated in the tumor periphery in

each subtype (e.g. *CCL20, FCGBP*) (Supplementary File 5), arguing that the non-reactive phe-

notype is a common feature for all peripheral MG and is not only driven by a differential abun-

dance pattern of one MG subpopulation relative to the other.

The tumor peripheral cytotoxic cell compartment exhibits an impaired activation signature.

Next, we investigated the regional differences in the lymphoid compartment composed

of CD4$^+$ and CD8$^+$ T cells and natural killer (NK) cells. We observed only very few significant

changes in the transcriptomic profiles of CD4$^+$ T cells between tumor center and periphery

(Figure 4 – figure supplement 1A and Supplementary File 4). Yet, comparing peripheral CD8$^+$

T cells with CD8$^+$ T cells from tumor center revealed 110 differentially expressed genes (43

genes upregulated and 67 genes downregulated) (Figure 4A and Supplementary File 4). Many downregulated genes in the peripheral CD8$^+$ T cells associated with canonical IFN responses (*IFI6, IFI27, MX1, STAT1, EPSTI1 PARP9, ISG15*) [34] cell proliferation (*STMN1, CENPF, HELLS, NUSAP1* and *DNPH1*) and T cell co-stimulation (*CD28, TMIGD2* (CD28H), *TNFRSF4* (OX40), *CD27* and *TNFRSF18* (GITR)) (Figure 4A). Contrary to our expectations, we saw upregulation of *CTLA4* in the center CD8$^+$ T cells which acts as a negative costimulatory molecule. However, unlike other costimulatory receptors, such as CD27 and CD28, CTLA-4 is not constitutively expressed on T lymphocytes [35]. but only induced following T cell activation, along with positive costimulatory molecules such as OX40 and GITR. In addition, upregulation of CTLA-4 requires entry into the cell cycle [35]. In line with that, we detected an upregulation of proliferative genes in center CD8$^+$ T cells. In summary, CTLA-4 induction in center CD8$^+$ T cells rather suggested T cell activation than exhaustion, especially since other inhibitory receptors like *PDCD1* (PD-1)*, LAG3* and *HAVCR2* (TIM-3) were not differentially expressed between sites. Moreover, we did not observe differential expression of genes involved in CD8$^+$ T cell effector functions like cytotoxicity (e.g., *GZMK, GZMB, KLRG1,* PRF1) or cytokines (e.g., *CCL5, XCL1, XCL2, IL10*). Yet, we noted upregulation of inhibitory genes (*TGFB1* and *FCRL6* [36]) in the peripheral CD8$^+$ T cells, suggesting that a pool of activated, proliferating and IFN-responsive CD8$^+$ T cells is present in the tumor center, however, absent in the tumor periphery.

Similar trends were observed for the peripheral NK cell population with peripherally reduced IFN response (*MX1* and *IFI44L*), and proliferative genes (*STMN1, HELLS, CENPF, PTTG1* and *DNPH1*), downregulated stimulatory receptors (*TMIGD2* (CD28H) and *TNFRSF18* (GITR)), and reduced NF-κB signaling (*NFKB1* and *RELB*) (Figure 4B and Supplementary File 4). Although, we observed upregulation of key genes associated with NK cell effector function in the periphery (e.g., *FCGR3A* (CD16), *FGFBP2, ITGB2, GZMH* and *KIR2DS4*), increased expression of inhibitory receptors like *LILRB1* and *KLRG1*, the latter especially in co-expression with chemokine receptor *CX3CR1*, identified the peripheral NK cells rather to be terminally differentiated with impaired cytotoxic capabilities [37]. This was in line with the observed abrogated cytokine activity profile in the peripheral NK cells with reduced expression of key factors like *XCL1, XCL2, LTB* and *CKLF*. In summary, our data revealed an impaired activation signature in the peripheral cytotoxic cell compartment.

CX3CR1 labels a specific CD8$^+$ T cell population in the circulation of grade 4 glioma patients.

Next, we investigated the relationships between circulating CD8$^+$ T cells and those from

the tumor milieu and, more specifically, the peripheral, infiltration zone characterized by an

abrogated CD8$^+$ T-cell IFN response and activation signature. Strikingly, there were large tran-

scriptomic differences between PBMC and periphery CD8$^+$ T cells (Figure 5A), with 1,417

differentially expressed genes (864 genes upregulated in the tumor periphery and 553 genes

upregulated in PBMC) (Figure 5B, Supplementary File 6).

Interestingly, one of the key genes upregulated in PBMC CD8$^+$ T cells was the chemo-

kine receptor *CX3CR1* (Figure 5B). Flow cytometry of an additional matched glioma grade 4

patient cohort confirmed an increased abundance of CX3CR1$^+$ CD8$^+$ T cells in PBMC com-

pared to almost absent CX3CR1$^+$ CD8$^+$ T cells in tumor periphery (Figure 5C, Figure 5 – figure

supplement 1A, Supplementary File 1). Unsupervised hierarchical sub-clustering of CD8$^+$ T

cells revealed that *CX3CR1* expression labelled a specific subpopulation among PBMC CD8$^+$

T cells (Figure 5D, 5E).

Recently, expression of CX3CR1 was demonstrated to distinguish memory CD8$^+$ T cells

with cytotoxic effector function in healthy donors and different inflammation related conditions

[38-41]. Further characterization of classical central memory (T$_{cm}$) and effector memory (T$_{em}$)

populations by varying surface expression levels of CX3CR1 identified a novel CX3CR1^{int}

subpopulation, termed peripheral memory (T$_{pm}$). T$_{pm}$ cells underwent frequent homeostatic di-

visions, re-acquired CD62L, homed to lymph nodes, and predominantly surveyed peripheral

tissues compared to T$_{cm}$ and T$_{em}$ [39]. In our dataset, the circulating CX3CR1$^+$ CD8$^+$ T cells

indeed displayed a core signature of memory CD8$^+$ T cells with effector function, comprising

expression of LFA-1 (*IGAL- ITGB2*)*, TBX2* (Tbet), *SELL* (CD62L)*, GZMB* (Granzyme B) and

PRF1 (Perforin 1) (Figure 5F), separating them from circulating *CX3CR1$^-$ CD28^{high}*, *IL7R^{high}*

and *CD27^{high}* naïve CD8$^+$ T cells (Figure 5G and Figure 5 – figure supplement 1B). Flow cy-

tometric analysis confirmed T$_{eff}$ to be CX3CR1^{high}, with negligible expression levels in the na-

ïve CD8$^+$ T cells, whereas the identified memory CD8$^+$ T cells (T$_{em}$ and T$_{pm}$) were CX3CR1^{int}

(Figure 5H, 5I). Collectively, surface expression analysis of CX3CR1 identified a subset of

CX3CR1^{high} T$_{eff}$ and CX3CR1^{int} memory (T$_{em}$, T$_{pm}$) CD8$^+$ T cells in the circulation of grade 4

glioma patients with potentially elevated tissue surveilling properties in the case of T$_{pm}$, which

are, however, largely absent in the tumor microenvironment.

CD8$^+$ T cells in the tumor periphery share features with tissue-resident memory T cells (T$_{rm}$)

We next examined the differing transcriptional and surface-specific features between

tumor infiltrating and circulating CD8$^+$ T cells. Surface staining for CD45RA and CD45RO,

discriminating naïve/effector from memory T cells, attributed a predominant CD45RO$^+$ memory phenotype to the tumor infiltrating CD8$^+$ T cells (Figure 6A, 6B). Interrogation of the transcriptomic profile of these cells revealed a key marker expression signature consistent with tissue-resident memory T cells (T$_{rm}$): Expression of cellular adhesion molecules (integrins) *ITGA1* (CD49a) and *ITGAE* (CD103), tissue retention marker *CD69*, chemokine receptors implicated in tissue-homing *CXCR3, CXCR6* and *CCR5* [42] and transcription factors, *ZNF683* (Hobit) and *PRDM1* (Blimp1) as well as reduced expression of *TBX21* (Tbet) and *EOMES* [43], strongly suggested a T$_{rm}$ phenotype for these cells (Figure 6C, 6D and Figure 6 – figure supplement 1A). Co-expression analysis of paired PBMC and tumor periphery samples using flow cytometry showed that CD69$^+$ CD103$^-$ and CD69$^+$ CD103$^+$ cells are the dominant CD8$^+$ T cell populations in the tumor periphery (Figure 6E, 6F). Combined, these data strongly suggest a T$_{rm}$ phenotype for the CD8$^+$ T cells in the tumor periphery.

Previous reports of T$_{rm}$ populating the brain in the aftermath of central or peripheral infections concluded that brain T$_{rm}$ cells surveil the brain tissue and mediate protection by rapid activation and enhanced cytokine production [42]. Indeed, CD8$^+$ T cells in the tumor periphery showed increased expression of genes belonging to costimulatory pathways, including *ICOS, TNFRSF4* (OX40) and *TNFRSF9* (4-1BB) (Figure 6 – figure supplement 1B, Supplementary File 6), albeit accompanied by high levels of inhibitory receptors *PDCD1* (PD-1), *LAG3, HAVCR2* (TIM-3) and *CTLA4* (Figure 6G). Moreover, expression of genes coding for cytotoxic molecules, including Granzyme B and Perforin 1 were decreased in the peripheral CD8$^+$ T cells, suggesting a compromised killing capacity of these cells. And lastly, CD8$^+$ T cells in the tumor periphery exhibited a transcription factor profile of exhausted T cells with high expression of *NR4A1, MAF* and *IRF4* (Figure 6G and Figure 6 – figure supplement 1C), which were implicated in T cell dysfunction and exhaustion [44, 45]. Collectively, these data indicate that CD8$^+$ T cells in the glioma periphery share features with T$_{rm}$ cells. However, inhibitory receptor expression, functional molecules and transcriptional signature ascribe an exhausted phenotype to these cells.

Noteworthy, we observed high upregulation of similar genes in the comparison tumor periphery vs. PBMC for CD4$^+$ T cells as for CD8$^+$ T cells (Figure 6 – figure supplement 1D). These included transcription factor family *NRA41-3,* identified as key mediator of T cell dysfunction [45], Dual Specificity Protein Phosphatase 2/4 (*DUSP2, DUSP4*) described as negative regulators of mitogen-activated protein (MAP) kinase superfamily and associated with impaired T cell effector activity [46] and T cell senescence [47], and transcription factor *CREM*

which was implicated in IL-2 suppression [48]. These genes could potentially identify pan T cell dysfunction markers within the GBM iTME [49].

Interrogation of cell-cell interactions revealed critical role of SPP1-mediated crosstalk between MG and lymphocytes in the tumor periphery

We finally investigated cell-cell interactions based on ligand-receptor expression levels using the CellChat platform [50]. Considering that MG and lymphocytes displayed an impaired activation signature in the tumor periphery, we focused our analysis on the tumor-peripheral crosstalk between these cells (Figure 7A). This revealed SPP1 (Osteopontin) as a leading potential cell-cell interaction mediator between MG and lymphocytes (Figure 7A and 7B). MG SPP1-mediated signaling was as well among the most significant interactions, when investigating cell-cell communication across all cell types and both sites (Figure 7 – figure supplement 1A, 1B). Further, we found that *SPP1* is mainly expressed by MG rather than glioma cells, contrary to previous reports [51] (Figure 7C, Figure 7 – figure supplement 1C, 1D). MG SPP1 conveys different interactions, depending on the recipient cell binding receptor expression profile. The predicted interactions of MG SPP1 with NK cells could be mediated via the integrin complex *ITGA4-ITGB1* (CD49d-CD29) (Figure 7A, 7C), whereas CD4[+] and CD8[+] T cells exhibited strong interactions with MG via SPP1/CD44 interaction (Figure 7A, 7B). The SPP1/CD44 axis was recently described to suppress T cell activation and proliferation [52]. The identified MG SPP1-mediated interactions might represent potential targets to modulate MG-lymphocyte crosstalk in the tumor periphery.

Discussion

In this study, we combined single-cell RNA sequencing and flow cytometry to interro-

gate the regional leukocyte activation signature in patient-matched biopsies from contrast-en-

hancing tumor center, infiltrative peripheral rim, and blood PBMCs of grade 4 glioma patients.

Our analyses revealed a distinct, regionally dependent transcriptional profile for most of the

investigated cell populations. While peripheral MG and cytotoxic cells predominantly dis-

played an impaired activation signature, MdMs showed pro-inflammatory traits in the tumor

periphery, however, were less abundant there compared to the tumor center, which was reported

by others as well [12, 13]. Supplemented with transcriptional analysis of paired PBMC samples,

we provide an in-depth characterization of the three main immunological compartments of

grade 4 glioma.

Previous studies focused on the description of the TME of grade 4 glioma, which also

considered regional differences, yet they focused primarily at neoplastic cells rather than the

immune compartment [12]. Others investigated the differences in the iTME composition be-

tween primary and metastatic brain tumors [5, 6]. Interestingly, the two latter ones reported

differences in the iTME composition between *IDH1^wt* and *IDH1^mut* glioma. Of note, both au-

thors included low-grade and even pre-treated recurrent glioma patients into the *IDH1^mut* group,

representing a quite heterogenous patient cohort. In this study, we aimed at providing a repre-

sentative selection of primary, treatment-naïve grade 4 glioma patients including *IDH1^wt* and

IDH1^mut to identify common transcriptional differences within the iTME between tumor center,

periphery and PBMC of grade 4 glioma. How far these regional differences vary between

IDH1^wt and *IDH1^mut* grade 4 glioma merits further investigation.

We identified a transcriptionally distinct MG subcluster, MG_1, which was enriched in

the infiltrative tumor periphery and displayed an anti-inflammatory/non-reactive phenotype. A

similar MG subpopulation expressing a comparable gene signature was recently described to

be enriched in Alzheimer's disease patients [53].

Additionally, the peripheral cytotoxic cell compartment exhibited an impaired activa-

tion state, including a downregulated IFN response signature in CD8$^+$ T cells. Induction of an

IFN response state was described as a consequence of T cell receptor-mediated IFN-γ produc-

tion, likely serving as an autocrine response and inducing the proliferative program [34]. Hence,

the reduced autocrine IFN-responsive state in the tumor peripheral CD8$^+$ T cells, together with

downregulated proliferative and co-stimulatory genes emphasized their impaired activation in

the peripheral infiltration zone. Recently, we showed that the response to immunotherapy in

GBM is indeed region dependent. For this, we cultured GBM explants in perfusion bioreactors

and treated with anti-CD47, anti-PD1, or their combination which induced an IFN-γ response

only in the tumor center, but not periphery [54]. Adding experimental support to the here de-

scribed impaired activation signature in the tumor periphery.

By exploring the transcriptional trajectory of $CD8^+$ T cells from the blood circulation

into the immunosuppressive TME of the tumor periphery, we uncovered $CX3CR1^{high}$ and

$CX3CR1^{int}$ effector and memory $CD8^+$ T cells, respectively, to be highly enriched in the PBMC,

but absent in the iTME. Recently, adoptive transfer studies of $CX3CR1^+ CD8^+$ T cells in a mel-

anoma mouse model significantly suppressed tumor growth [41]. Others identified increased

frequencies of $CX3CR1^+ CD8^+$ T cells in non-small cell lung and melanoma patients who re-

sponded to anti-PD-1 therapy, where these cells exhibited migratory capabilities into the tumor

site followed by potent tumor rejection [40, 41]. Thus, the authors proposed T cell CX3CR1

expression as a predictor of response to ICI therapy. Even though $CX3CR1^+ CD8^+$ T cells might

not be specific to GBM, as they are found in healthy donor and different inflammation related

conditions [38-41], the absence of these effector and potentially ICI therapy responsive $CD8^+$

T cells in the glioma TME could serve as an additional explanation for the disappointing out-

come of clinical trials using ICI in glioma patients.

The observed T_{rm} exhaustion phenotype of the glioma residing $CD8^+$ T cells was re-

cently reported as well for tumor-infiltrating $PD-1^{high} CD8^+$ T cells in hepatocellular carcinoma

[44]. Whether these glioma-associated $CD8^+$ T cells really possess tumor-specificity requires

further study. Particularly in the light of a recent study by Smolders and colleagues who re-

ported a consistent brain-resident $CD8^+$ T cell population in a miscellaneous autopsy cohort of

patients with neurological disorders excluding brain malignancies (Alzheimer's disease, Par-

kinson's disease, dementia, depression, multiple sclerosis), as well as patients with no known

brain disease. These brain-resident $CD8^+$ T cells displayed a remarkably consistent T_{rm} pheno-

type [55]. The authors further showed high expression of inhibitory receptors CTLA-4 and PD-

1 on the brain-resident $CD8^+ T_{rm}$ cells, which is in line with the core phenotypic signature of

T_{rm} cells from other tissues [56, 57]. Yet, the brain $CD8^+ T_{rm}$ cells showed a preserved inflam-

matory potential with substantial production of IFN-γ and TNF-α upon *ex vivo* stimulation.

They concluded that extensive immune activation with release of highly neurotoxic lytic en-

zymes, such as perforin and granzyme B, harmfully impacts the brain parenchyma and should

be tightly controlled, whilst maintaining the capability to elicit a fast inflammatory response

when a neurotropic virus threatens the CNS [55]. Therefore, inhibitory receptors like PD-1 and

CTLA-4 on brain $CD8^+ T_{rm}$ cells may support CNS homeostasis by preventing uncontrolled T

cell reactivity, and the availability of the receptor ligands may determine their inhibitory effect.

While this may represent a well-balanced equilibrium under healthy conditions, the tumor set-
ting leads to its disruption with upregulation of inhibitory ligands like PD-L1 on glioma cells
and CD86 on GAMs, leading to the dysfunctional state seen in the glioma-residing CD8$^+$ T$_{rm}$
cells.

Another study comprehensively showed, that peripheral infections generate antigen-
specific CD8$^+$ T$_{rm}$ cells in the brain, mediating protection against CNS infections [42]. These
data could implicate that the glioma-associated CD8$^+$ T cells are devoid of tumor-specific reac-
tivity, but rather represent a pre-existing T cell population generated after peripheral infections,
which acquired a dysfunctional state upon glioma formation. To test this hypothesis, further
characterization of these cells is required, including analysis of T cell receptor clonality and
tumor-specificity by patient-matched T cell/glioma-sphere co-culture assays.

Lastly, our cell-cell interaction analysis revealed signaling pathways between peripheral
MG and lymphocytes potentially inducing the observed impaired activation signature. The pre-
dicted interaction between MG SPP1 and NK cells integrin complex *ITGA4-ITGB1* (CD49d-
CD29), might mediate NK cell adhesion and migration [58]. This may facilitate interaction of
inhibitory NK receptors KLRB1 and CD94/NKG2A with MG C-type lectin-related ligands and
HLA-E, respectively, which could explain the observed impaired activation state of peripheral
NK cells. Moreover, SPP1/CD44 interaction in T lymphocytes was described to suppress cell
activation and proliferation [52]. In a comprehensive approach where transcriptional states of
human MG were mapped during aging and disease, *SPP1* was found to be differentially ex-
pressed in aging-microglia. This was associated to a doubling of the abundance of SPP1$^+$ GAMs
in GBM samples compared to MG from age-matched controls [59]. These observations support
a possible role of MG SPP1 in glioma progression. More recently, myeloid-derived osteopontin
(encoded by *SPP1*) was shown to trigger a chronic activation of NFAT2 in tumor-reactive CD8$^+$
T cells, leading to T cell dysfunction and exhaustion in an experimental mouse model of GBM
[60]. This adds an additional layer of mechanistic evidence for the *SPP1*/Osteopontin-mediated
signaling axis leading to CD8$^+$ T cell dysfunction in GBM.

Limitations of our study include the limited patient number, thereby our study was nei-
ther designed nor powered to explore differences in neoplastic cells, given the high inter- and
intra-patient variability in glioma cells [12]. Importantly, our dataset establishes a starting point
for further interrogation and provides a first analysis of the transcriptional landscape of the
major immune populations in grade 4 glioma within three important regional compartments.
Further, we confirmed the observed phenotype of CD8$^+$ T cells in the blood and tumor periphery
by flow cytometry in a cohort of ten additional patients, addressing possible generalization

concerns. Together, we provide a regionally resolved map of leukocyte activation in the TME

and blood circulation from grade 4 glioma patients, helping the research community to uncover

novel therapeutic strategies to combat this fatal disease.

Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
biological sample (Human adult GBM tissue samples)	Tumor Center; Tumor Periphery	Neurosurgical Clinic of the University Hospital of Basel, Switzerland		
biological sample (Human Peripheral Blood Buffy Coat)	PBMC	Neurosurgical Clinic of the University Hospital of Basel, Switzerland		
antibody	Rat monoclonal anti-human/mouse CD11b (clone M1/70), FITC	BioLegend	Cat# 101206	FACS: 1:50
antibody	Mouse monoclonal anti-human CD45 (clone 2D1), FITC	BioLegend	Cat# 368508	FACS: 1:50
antibody	Fc-Block anti-human CD16/CD32, TruStain FcX	BioLegend	Cat# 422302	FACS: 1:50
antibody	Mouse monoclonal anti-human CD45RO (clone UCHL1), APC	BioLegend	Cat# 304210	FACS: 1:25
antibody	Mouse monoclonal anti-human CD45RA (clone HI100), PE	BioLegend	Cat# 304108	FACS: 1:25
antibody	Mouse monoclonal anti-human CD3e (clone UCHT1), BV650	BioLegend	Cat# 300468	FACS: 1:25
antibody	Mouse monoclonal anti-human CD8a (clone RPA-T8), BV421	BioLegend	Cat# 301036	FACS: 1:25

antibody	Mouse monoclonal anti-human CCR7 (clone G043H7), FITC	BioLegend	Cat# 353216	FACS: 1:25
antibody	Mouse monoclonal anti-human CD62L (clone DREG-56), AF700	BioLegend	Cat# 304820	FACS: 1:50
antibody	Mouse monoclonal anti-human CD69 (clone FN50), APC/Cy7	BioLegend	Cat# 310914	FACS: 1:25
antibody	Mouse monoclonal anti-human CD103 (clone Ber-ACT8), PerCP/Cy5.5	BioLegend	Cat# 350226	FACS: 1:25
antibody	Rat monoclonal anti-human CX3CR1 (clone 2A9-1), PE-Cy7	BioLegend	Cat# 341612	FACS: 1:25
commercial assay or kit	Chromium Single Cell 3' Reagent Kits v3	10x Genomics	Cat# CG000183	
commercial assay or kit	BioAnalyzer High Sensitivity DNA Analysis kit	Agilent	Cat# 5067-4626	
commercial assay or kit	Qubit dsDNA High Sensitiivity assay kit	ThermoFisher	Cat# Q33230	
chemical compound, drug	Collagenase-4	Worthington Bio-chemical Cooperation	Cat# LS004188	
chemical compound, drug	DNAse1	Roche	Cat# 10104159001	
chemical compound, drug	ACK lysis buffer	Gibco	Cat# A1049201	

chemical compound, drug	Bambanker	Nippon Genetics	Cat# BB01	
chemical compound, drug	Sucorse	Sigma Aldrich	Cat# 84100	
chemical compound, drug	Ficoll-Paque PLUS	Cytiva	Cat# 17144002	
chemical compound, drug	Live/Dead Fixable Near IR Dead Stain Kit, APC-Cy7	ThermoFisher	Cat# L34976	
chemical compound, drug	Zombie Aqua Fixable Viability Kit	BioLegend	Cat# 423102	
software, algorithm	R environment, version 4.1	R Core Team	https://www.r-project.org/	
software, algorithm	GraphPad Prism 9	GraphPad Software Inc.	N/A	
software, algorithm	Flow Jo, version 10.8.1	Tree Star	N/A	

Ethics statement

Human adult GBM tissue samples were obtained at the Neurosurgical Clinic of the Uni-
versity Hospital of Basel, Switzerland in accordance with the Swiss Human Research Act and
institutional ethics commission (EKNZ 02019-02358). All patients gave written informed con-
sent for tumor biopsy collection and signed a declaration permitting the use of their biopsy
specimens in scientific research, including storage in our brain tumor biobank (Req-2019-
00553). All patient identifying information was removed and tissue was coded for identifica-
tion.

Glioma tissue dissociation

Resected glioma tissue samples were immediately placed on ice and transferred to the

laboratory for single cell dissociation within 2-3 h after resection. Human brain tissue was man-

ually minced using razor blades and enzymatically dissociated at 37°C for 30 minutes with 1

mg/ml collagenase-4 (#LS004188, Worthington Biochemical Corporation, USA) and 250 U/ml

DNAse1 (#10104159001, Roche, Switzerland) in a buffer containing Hank's Balanced Salt

Solution (HBSS) with Ca^{2+}/Mg^{2+}, 1% MEM non-essential amino acids (Gibco, USA), 1 mM

sodium pyruvate (Gibco), 44 mM sodium bi-carbonate (Gibco), 25 mM HEPES (Gibco), 1%

GlutaMAX (Gibco) and 1% antibiotic-antimycotic (Sigma-Aldrich, USA). Cells were filtered

and separated from dead cells, debris and myelin by a 0.9 M sucrose (#84100, Sigma Aldrich)

density gradient centrifugation. Upon ACK-lysis for removal of erythrocytes (#A1049201,

Gibco) the now generated single-cell suspension (SCS) was washed, counted and frozen in

Bambanker (#BB01, Nippon Genetics, Germany) in liquid nitrogen until use.

PBMCs (Peripheral blood mononuclear cells) preparation

Patient blood samples were directly placed on ice and transferred to the laboratory for

PBMC isolation. Blood samples were centrifuged to separate buffy coat from plasma and eryth-

rocytes, followed by standard density gradient centrifugation protocol (#17144002, Ficoll-

Paque PLUS, Cytiva, USA) to isolate PBMCs. PBMCs were frozen in Bambanker (#BB01,

Nippon Genetics, Germany) in liquid nitrogen until use.

FACS sorting for single cell RNA sequencing (scRNA-seq)

Cryopreserved tumor digests from glioma samples (center and periphery), as well as

autologous PBMCs were thawed and washed with excess ice-cold 1xPBS and spun down at

350xg for 5 min. Subsequently, the cells were stained with Live/Dead (APC-Cy7 (Near IR), #

L34976, Thermo Fischer) and a cocktail of fluorescently-conjugated antibodies CD11b (FITC,

clone M1/70, #101206, BioLegend) and CD45 (FITC, clone 2D1, #368508, BioLegend), and

large debris were removed with a 40-μm strainer. All samples were acquired on the BD FACS

ARIA Fusion III (Becton Dickinson GmbH, Germany). For single-cell RNA-seq experiments,

live and single gated cells were sorted into non-immune cell ($CD45^-CD11b^-$) and immune cell

($CD45^+CD11b^+$) populations. Both populations were directly sorted into Eppendorf tubes with

1xPBS supplemented with 1% BSA for single cell RNA sequencing.

Single cell RNA sequencing (scRNA-seq) – Library preparation and sequencing

473 Single-cell RNA-seq was performed using Chromium Single Cell 3' GEM, Library &

474 Gel Bead Kit v3 (#CG000183, 10x Genomics, Pleasanton, CA, USA) following the manufac-

475 turer's protocol. Briefly, non-immune cells and immune cells were mixed at a defined ratio of

476 1:4. Roughly 8000-10000 cells per sample, diluted at a density of 100–800 cells/μL in PBS plus

477 1% BSA determined by Cellometer Auto 2000 Cell Viability Counter (Nexelom Bioscience,

478 Lawrence, MA USA), and were loaded onto the chip. The quality and concentration of both

479 cDNA and libraries were assessed using an Agilent BioAnalyzer with High Sensitivity kit

480 (#5067-4626, Agilent, Santa Clara, CA USA) and Qubit Fluorometer with dsDNA HS assay

481 kit (#Q33230, Thermo Fischer Scientific, Waltham, MA USA) according to the manufacturer's

482 recommendation. For sequencing, samples were mixed in equimolar fashion and sequenced on

483 an Illumina HiSeq 4000 with a targeted read depth of 50,000 reads/cell and sequencing param-

484 eters were set for Read1 (28 cycles), Index1 (8 cycles), and Read2 (91 cycles).

486 *Single cell RNA sequencing (scRNA-seq) - Computational analysis*

487 The dataset was analyzed by the Bioinformatics Core Facility, Department of Biomed-

488 icine, University of Basel. Read quality was controlled with the FastQC tool (version 0.11.5).

489 Sequencing files were processed using the Salmon Alevin tool (v 1.3.0) [61] to perform quality

490 control, sample demultiplexing, cell barcode processing, pseudo-alignment of cDNA reads to

491 the human Gencode v35 reference and counting of UMIs. Parameters *--keepCBFraction 1* and

492 *--maxNumBarcodes 100000* were used.

493 Processing of the UMI counts matrix was performed using the Bioconductor packages

494 DropletUtils (version 1.8.0) [62, 63], scran (version 1.16.0) [64, 65] and scater (version 1.16.2)

495 [66], following mostly the steps illustrated in the OSCA book (http://bioconduc-

496 tor.org/books/release/OSCA/) [65, 67]. Filtering for high-quality cells was done based on li-

497 brary size (at least 2,000 UMI counts per cell), the number of detected genes (at least 700 genes

498 detected) and the percentage of reads mapping to mitochondrial genes (larger than 0% and

499 lower than 15%), based on the distribution observed across cells. Low-abundance genes with

500 average counts per cell lower than 0.006 were filtered out. The presence of doublet cells was

501 investigated with the scDblFinder package (version 1.2.0), and suspicious cells were filtered

502 out (score > 0.6). After quality filtering, the resulting dataset consisted of UMI counts for

503 15,523 genes and 45,466 cells, ranging from 803 to 9,121 per sample.

504 UMI counts were normalized with size factors estimated from pools of cells created

505 with the scran package *quickCluster()* function [64, 68]. To distinguish between genuine bio-

506 logical variability and technical noise we modeled the variance of the log-expression across

genes using a Poisson-based mean-variance trend. The scran package *denoisePCA()* function
was used to denoise log-expression data by removing principal components corresponding to
technical noise. Consistent to the *findElbowPoint()* function from the PCAtools Bioconductor
package, this led us to retain the top 5 principal components (PCs), explaining 48.3% of the
total variance in the dataset, for the clustering and dimensionality reduction steps.

Since the excluded deeper PCs (PC8 and 10) were associated to patient-specific effects
(which in our experimental setup are also confounded with batch effects), this choice dispensed
us from performing an additional correction for patient-specific effects. A quantitative test of
overlap of cells across patients was made using the CellMixS package [69] (version 1.12.0)
developed to quantify the effectiveness of batch correction methods. We used the cell-specific
mixing (CMS) score, which highlighted a very good overlap across cells from different patients
in the lymphoid compartment and for the monocytes. The myeloid compartment displayed a
slightly elevated patient-specific structure, while it was most pronounced for the CD45 negative
subset.

A t-stochastic neighbor embedding (t-SNE) was built with a perplexity of 50 using the
top most variable genes (141 genes with estimated biological variance > 0.3, excluding genes
with highest proportion of reads in the ambient RNA pool estimated from empty droplets), and
the denoised principal components as input. Clustering of cells was performed with hierarchical
clustering on the Euclidean distances between cells (with Ward's criterion to minimize the total
variance within each cluster [70]; package cluster version 2.1.0). The number of clusters used
for following analyses was identified by applying a dynamic tree cut (package dynamicTreeCut,
version 1.63-1) [71], resulting in 10 with argument *deepSplit* set to 1, or 22 clusters with argu-
ment *deepSplit* set to 2. This clustering was validated with an alternative clustering approach
using a graph-based approach and Louvain algorithm for community detection (using the *Find-*
Neighbors() and *FindClusters()* functions from the *Seurat* package [72] (version 4.3.0), with a
resolution of 0.7) which showed a good correspondence to our hierarchical clustering.

The *findMarkers* function of the scran package was used to find the best markers across
annotated cell types (parameters *direction="up"* and *pval.type="any"*). The top 10 markers
for each cell type were extracted and pooled to from a list of 68 markers.

The Bioconductor package *SingleR* (version 1.2.4) was used for cell-type annotation of
the cells [73] using as references (i) a public bulk RNA-seq dataset of sorted immune cell types
from human PBMC samples [16], available through the *celldex* Bioconductor package; (ii) a
bulk RNA-seq dataset of sorted immune cell types from the tumor microenvironment of human
gliomas [5] (UMI count matrix and annotation downloaded from

); (iii) a 10X genomics scRNA-seq dataset of TAMs from the tumor micro-environment of glioblastoma tumors from 7 newly diagnosed human patients [9]; (iv) a microglia and a macrophage signature scores were defined by averaging the center and scaled expression levels of gene lists obtained in [7] and (v) a Smartseq2 scRNA-seq dataset of IDH-wild-type glioblastoma tumors [17] (downloaded from GEO accession GSE131928). An endothelial score was defined by averaging the center and scaled expression levels of the genes *CDH5*, *VWF*, *CD34* and *PECAM1*.

The SingleR high-quality assignments (pruned scores) from the comparisons to the multiple references (some being better at enlightening some immune subsets than others) were used to manually derive a consensus cell type annotation for each cluster. This manual annotation was done jointly with the input from the signature scores and relative expression patterns of known marker genes and of cluster-specific genes.

When we judged that the resolution was not sufficient to discriminate well the subtypes of cells within a subset (in particular for the CD45neg and myeloid cells subsets), we isolated these cells and re-performed selection of hyper-variable genes, dimensionality reduction and clustering on the isolated subset.

Differential abundance analysis of the identified cell types between tumor sites was performed using *limma-voom* [74]. This method, implemented in the *diffcyt* package, was shown to be able to handle well the overdispersion in the cell type proportions estimates typically observed with single cell technologies [75]. Differential abundance of cell types was considered to be significant at a false discovery rate (FDR) lower than 5 %. To provide increased resolution into local trends within the tumor innate immune subset we used the *miloR* bioconductor package [76] (version 1.4.0) to test for differential abundance on neighborhoods of a *k*-nearest neighbor graph. Testing was done with *limma-voom* similarly to above. Cell type labels were assigned to each neighborhood by finding the most abundant cell type across cells and requiring that at least 70% of the cells come from one cell type.

Differential expression between tumor sites, or between PBMC cells and tumor periphery cells, stratified by annotated cell type, was performed using a pseudo-bulk approach, which was shown to be the best-performing approach for differential expression testing in recent benchmarks [77]. UMI counts were summed across cells from each cell type in each sample when at least 20 cells could be aggregated. The aggregated samples were then treated as bulk RNA-seq samples [78] and for each pairwise comparison genes were filtered to keep genes detected in at least 5% of the cells aggregated. The package edgeR (version 3.30.3) [79] was used to perform TMM normalization [80] and to test for differential expression with the

Generalized Linear Model (GLM) framework, using a model accounting for patient-specific

effects. Genes with a FDR lower than 5 % were considered differentially expressed. To validate

our differential expression analysis, we rerun a pseudo-bulk expression analysis using DESeq2

(version 1.38.3) [81] for the main immune populations in the periphery versus center and pe-

riphery versus PBMC comparison. Although, the results were not identical, they overall agreed

with edgeR (Supplementary File 7). Gene set enrichment analysis was performed with the func-

tion camera [82] on gene sets from the Molecular Signature Database (MSigDB, version 7.4)

[27, 83]. We retained only sets containing more than 5 genes, and gene sets with a FDR lower

than 5% were considered as significant.

Cell chat analysis

The R package CellChat (1.1.3) [50] was used to analyze cell-cell interactions in our

dataset (with previously annotated 9 cell types). We followed the recommended workflow to

infer the cell state-specific communications (using *identifyOverExpressedGenes*, *identifyOver-*

ExpressedInteractions and *projectData* with the default parameters). We performed 3 separate

analyses, on the center and the periphery subsets and a comparison analysis as described in the

official workflow. We visualized the significant interactions for the microglia cluster using

netVisual_chord_gene and used *plotGeneExpression* to display of the expression of all genes

involved SPP1 signaling pathway in the cell populations. Finally, *netAnalysis_signalin-*

gRole_scatter was used to calculate and visualize incoming and outgoing signaling strength.

Flow cytometry analysis of paired PBMC and periphery samples

Cryopreserved samples were thawed and washed with excess ice-cold 1xPBS and spun

down at 350xg for 5 min. Cells were resuspended in FACS buffer (PBS plus 2% FBS) and

blocked with monoclonal antibody to CD16/32 (Human TruStain FcX, #422302, Biolegend)

for 10 min at 4°C before staining with surface antibodies: CD45RA (PE, clone HI100,

#304108), CD45RO (APC, clone UCHL1, #304210), CD3e (BV650, clone UCHT1, #300468),

CD8a (BV421, clone RPA-T8, #301036), CCR7 (FITC, clone G043H7, #353216), CD62L

(AF700, clone DREG-56, #304820), CD69 (APC-Cy7, clone FN50, #310914), CD103

(PerCP/Cy5.5, clone Ber-ACT8, #350226) and CX3CR1 (PE/Cy7, clone 2A9-1, #341612). All

antibodies were purchased from BioLegend, USA. Cells were stained for 30 min at 4°C, and

subsequently washed with FACS buffer. To exclude dead cells Zombie Aqua Fixable Viability

Kit (#423102, 1:100, BioLegend) was added. Acquisition was performed on a CytoFLEX

(Beckman). Data was analyzed using FlowJo software, version 10.8.1 (TreeStar). Gates were

drawn by using Fluorescent Minus One (FMO) controls.

Statistical analysis of flow cytometry data

Data analysis and graph generation was performed using GraphPad Prism 9 (GraphPad

Prism Software Inc.). Paired comparisons between two groups were performed using Wilcoxon

matched-pairs signed rank test. Differences of more than two paired groups were assessed using

repeated measures one-way ANOVA test, followed by post-hoc Šidák's multiple comparisons

correction. A *p* value < 0.05 was considered statistically significant. *$p \le 0.05$, **$p \le 0.01$, ***p

≤ 0.001, ****$p \le 0.0001$.

Graphical illustrations

All graphical illustrations were created with BioRender.com.

Data availability

622
623   The UMI count matrix and cell metadata from the scRNA-seq dataset are available on
624   GEO under accession number GSE197543. The remaining data are available within the Article
625   and Supplementary Information.

References

1. Stupp, R., W.P. Mason, M.J. van den Bent, M. Weller, B. Fisher, M.J. Taphoorn, K. Belanger, A.A. Brandes, C. Marosi, U. Bogdahn, J. Curschmann, R.C. Janzer, S.K. Ludwin, T. Gorlia, A. Allgeier, D. Lacombe, J.G. Cairncross, E. Eisenhauer, and R.O. Mirimanoff, *Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma.* N Engl J Med, 2005. **352**(10): p. 987-96.

2. Bowman, R.L., F. Klemm, L. Akkari, S.M. Pyonteck, L. Sevenich, D.F. Quail, S. Dhara, K. Simpson, E.E. Gardner, C.A. Iacobuzio-Donahue, C.W. Brennan, V. Tabar, P.H. Gutin, and J.A. Joyce, *Macrophage Ontogeny Underlies Differences in Tumor-Specific Education in Brain Malignancies.* Cell Rep, 2016. **17**(9): p. 2445-2459.

3. Reardon, D.A., T.J. Kaley, J. Dietrich, J.L. Clarke, G.P. Dunn, M. Lim, T.F. Cloughesy, H.K. Gan, A.J. Park, P. Schwarzenberger, T. Ricciardi, M.J. Macri, A. Ryan, and R.R. Venhaus, *Phase 2 study to evaluate safety and efficacy of MEDI4736 (durvalumab [DUR]) in glioblastoma (GBM) patients: An update.* Journal of Clinical Oncology, 2017. **35**(15_suppl): p. 2042-2042.

4. Reardon, D.A., A. Omuro, A.A. Brandes, J. Rieger, A. Wick, J. Sepulveda, S. Phuphanich, P. de Souza, M.S. Ahluwalia, M. Lim, G. Vlahovic, and J. Sampson, *OS10.3 Randomized Phase 3 Study Evaluating the Efficacy and Safety of Nivolumab vs Bevacizumab in Patients With Recurrent Glioblastoma: CheckMate 143.* Neuro-Oncology, 2017. **19**(Suppl 3): p. iii21-iii21.

5. Klemm, F., R.R. Maas, R.L. Bowman, M. Kornete, K. Soukup, S. Nassiri, J.P. Brouland, C.A. Iacobuzio-Donahue, C. Brennan, V. Tabar, P.H. Gutin, R.T. Daniel, M.E. Hegi, and J.A. Joyce, *Interrogation of the Microenvironmental Landscape in Brain Tumors Reveals Disease-Specific Alterations of Immune Cells.* Cell, 2020. **181**(7): p. 1643-1660.e17.

6. Friebel, E., K. Kapolou, S. Unger, N.G. Núñez, S. Utz, E.J. Rushing, L. Regli, M. Weller, M. Greter, S. Tugues, M.C. Neidert, and B. Becher, *Single-Cell Mapping of Human Brain Cancer Reveals Tumor-Specific Instruction of Tissue-Invading Leukocytes.* Cell, 2020. **181**(7): p. 1626-1642.e20.

7. Müller, S., G. Kohanbash, S.J. Liu, B. Alvarado, D. Carrera, A. Bhaduri, P.B. Watchmaker, G. Yagnik, E. Di Lullo, M. Malatesta, N.M. Amankulor, A.R. Kriegstein, D.A. Lim, M. Aghi, H. Okada, and A. Diaz, *Single-cell profiling of human gliomas reveals macrophage ontogeny as a basis for regional differences in macrophage activation in the tumor microenvironment.* Genome Biology, 2017. **18**(1): p. 234.

8. Abdelfattah, N., P. Kumar, C. Wang, J.-S. Leu, W.F. Flynn, R. Gao, D.S. Baskin, K. Pichumani, O.B. Ijare, S.L. Wood, S.Z. Powell, D.L. Haviland, B.C. Parker Kerrigan, F.F. Lang, S.S. Prabhu, K.M. Huntoon, W. Jiang, B.Y.S. Kim, J. George, and K. Yun, *Single-cell analysis of human glioma and immune cells identifies S100A4 as an immunotherapy target.* Nature Communications, 2022. **13**(1): p. 767.

9. Pombo Antunes, A.R., I. Scheyltjens, F. Lodi, J. Messiaen, A. Antoranz, J. Duerinck, D. Kancheva, L. Martens, K. De Vlaminck, H. Van Hove, S.S. Kjølner Hansen, F.M. Bosisio, K. Van der Borght, S. De Vleeschouwer, R. Sciot, L. Bouwens, M. Verfaillie, N. Vandamme, R.E. Vandenbroucke, O. De Wever, Y. Saeys, M. Guilliams, C. Gysemans, B. Neyns, F. De Smet, D. Lambrechts, J.A. Van Ginderachter, and K.

Movahedi, *Single-cell profiling of myeloid cells in glioblastoma across species and*
disease stage reveals macrophage competition and specialization. Nature
Neuroscience, 2021. **24**(4): p. 595-610.

10. Goswami, S., T. Walle, A.E. Cornish, S. Basu, S. Anandhan, I. Fernandez, L. Vence, J.
Blando, H. Zhao, S.S. Yadav, M. Ott, L.Y. Kong, A.B. Heimberger, J. de Groot, B.
Sepesi, M. Overman, S. Kopetz, J.P. Allison, D. Pe'er, and P. Sharma, *Immune profiling*
of human tumors identifies CD73 as a combinatorial target in glioblastoma. Nature
Medicine, 2020. **26**(1): p. 39-46.

11. Chen, A.X., R.D. Gartrell, J. Zhao, P.S. Upadhyayula, W. Zhao, J. Yuan, H.E. Minns,
A. Dovas, J.N. Bruce, A. Lasorella, A. Iavarone, P. Canoll, P.A. Sims, and R. Rabadan,
Single-cell characterization of macrophages in glioblastoma reveals MARCO as a
mesenchymal pro-tumor marker. Genome Medicine, 2021. **13**(1): p. 88.

12. Darmanis, S., S.A. Sloan, D. Croote, M. Mignardi, S. Chernikova, P. Samghababi, Y.
Zhang, N. Neff, M. Kowarsky, C. Caneda, G. Li, S.D. Chang, I.D. Connolly, Y. Li,
B.A. Barres, M.H. Gephart, and S.R. Quake, *Single-Cell RNA-Seq Analysis of*
Infiltrating Neoplastic Cells at the Migrating Front of Human Glioblastoma. Cell Rep,
2017. **21**(5): p. 1399-1410.

13. Landry, A.P., M. Balas, S. Alli, J. Spears, and Z. Zador, *Distinct regional ontogeny and*
activation of tumor associated macrophages in human glioblastoma. Scientific Reports,
2020. **10**(1): p. 19542.

14. Louis, D.N., A. Perry, P. Wesseling, D.J. Brat, I.A. Cree, D. Figarella-Branger, C.
Hawkins, H.K. Ng, S.M. Pfister, G. Reifenberger, R. Soffietti, A. von Deimling, and
D.W. Ellison, *The 2021 WHO Classification of Tumors of the Central Nervous System:*
a summary. Neuro-Oncology, 2021. **23**(8): p. 1231-1251.

15. Linderman, G.C., M. Rachh, J.G. Hoskins, S. Steinerberger, and Y. Kluger, *Fast*
interpolation-based t-SNE for improved visualization of single-cell RNA-seq data. Nat
Methods, 2019. **16**(3): p. 243-245.

16. Monaco, G., B. Lee, W. Xu, S. Mustafah, Y.Y. Hwang, C. Carré, N. Burdin, L. Visan,
M. Ceccarelli, M. Poidinger, A. Zippelius, J. Pedro de Magalhães, and A. Larbi, *RNA-*
Seq Signatures Normalized by mRNA Abundance Allow Absolute Deconvolution of
Human Immune Cell Types. Cell Rep, 2019. **26**(6): p. 1627-1640.e7.

17. Neftel, C., J. Laffy, M.G. Filbin, T. Hara, M.E. Shore, G.J. Rahme, A.R. Richman, D.
Silverbush, M.L. Shaw, C.M. Hebert, J. Dewitt, S. Gritsch, E.M. Perez, L.N. Gonzalez
Castro, X. Lan, N. Druck, C. Rodman, D. Dionne, A. Kaplan, M.S. Bertalan, J. Small,
K. Pelton, S. Becker, D. Bonal, Q.-D. Nguyen, R.L. Servis, J.M. Fung, R. Mylvaganam,
L. Mayr, J. Gojo, C. Haberler, R. Geyeregger, T. Czech, I. Slavc, B.V. Nahed, W.T.
Curry, B.S. Carter, H. Wakimoto, P.K. Brastianos, T.T. Batchelor, A. Stemmer-
Rachamimov, M. Martinez-Lage, M.P. Frosch, I. Stamenkovic, N. Riggi, E. Rheinbay,
M. Monje, O. Rozenblatt-Rosen, D.P. Cahill, A.P. Patel, T. Hunter, I.M. Verma, K.L.
Ligon, D.N. Louis, A. Regev, B.E. Bernstein, I. Tirosh, and M.L. Suvà, *An Integrative*
Model of Cellular States, Plasticity, and Genetics for Glioblastoma. Cell, 2019. **178**(4):
p. 835-849.e21.

18. Pinton, L., E. Masetto, M. Vettore, S. Solito, S. Magri, M. D'Andolfi, P. Del Bianco, G.
Lollo, J.P. Benoit, H. Okada, A. Diaz, A. Della Puppa, and S. Mandruzzato, *The immune*
suppressive microenvironment of human gliomas depends on the accumulation of bone
marrow-derived macrophages in the center of the lesion. J Immunother Cancer, 2019.
7(1): p. 58.

19. Nguyen, A.T., K. Wang, G. Hu, X. Wang, Z. Miao, J.A. Azevedo, E. Suh, V.M. Van
Deerlin, D. Choi, K. Roeder, M. Li, and E.B. Lee, *APOE and TREM2 regulate amyloid-*
responsive microglia in Alzheimer's disease. Acta Neuropathol, 2020. **140**(4): p. 477-
493.

20. Soroceanu, L., R. Murase, C. Limbad, E. Singer, J. Allison, I. Adrados, R. Kawamura,
A. Pakdel, Y. Fukuyo, D. Nguyen, S. Khan, R. Arauz, G.L. Yount, D.H. Moore, P.Y.
Desprez, and S.D. McAllister, *Id-1 is a key transcriptional regulator of glioblastoma*
aggressiveness and a novel therapeutic target. Cancer Res, 2013. **73**(5): p. 1559-69.

21. Papaspyridonos, M., I. Matei, Y. Huang, M. do Rosario Andre, H. Brazier-Mitouart,
J.C. Waite, A.S. Chan, J. Kalter, I. Ramos, Q. Wu, C. Williams, J.D. Wolchok, P.B.
Chapman, H. Peinado, N. Anandasabapathy, A.J. Ocean, R.N. Kaplan, J.P. Greenfield,
J. Bromberg, D. Skokos, and D. Lyden, *Id1 suppresses anti-tumour immune responses*
and promotes tumour progression by impairing myeloid cell maturation. Nature
Communications, 2015. **6**(1): p. 6840.

22. Holtman, I.R., D. Skola, and C.K. Glass, *Transcriptional control of microglia*
phenotypes in health and disease. The Journal of Clinical Investigation, 2017. **127**(9):
p. 3220-3229.

23. Nomaru, H., K. Sakumi, A. Katogi, Y.N. Ohnishi, K. Kajitani, D. Tsuchimoto, E.J.
Nestler, and Y. Nakabeppu, *Fosb gene products contribute to excitotoxic microglial*
activation by regulating the expression of complement C5a receptors in microglia. Glia,
2014. **62**(8): p. 1284-98.

24. Sweet, D.R., N.T. Vasudevan, L. Fan, C.E. Booth, K.S. Keerthy, X. Liao, V.
Vinayachandran, Y. Takami, D. Tugal, N. Sharma, E.R. Chan, L. Zhang, Y. Qing, S.L.
Gerson, C. Fu, A. Wynshaw-Boris, P. Sangwung, L. Nayak, P. Holvoet, K. Matoba, Y.
Lu, G. Zhou, and M.K. Jain, *Myeloid Krüppel-like factor 2 is a critical regulator of*
metabolic inflammation. Nature Communications, 2020. **11**(1): p. 5872.

25. Salojin, K.V., I.B. Owusu, K.A. Millerchip, M. Potter, K.A. Platt, and T. Oravecz,
Essential role of MAPK phosphatase-1 in the negative control of innate immune
responses. J Immunol, 2006. **176**(3): p. 1899-907.

26. Lee, H., S. Cha, M.S. Lee, G.J. Cho, W.S. Choi, and K. Suk, *Role of antiproliferative B*
cell translocation gene-1 as an apoptotic sensitizer in activation-induced cell death of
brain microglia. J Immunol, 2003. **171**(11): p. 5802-11.

27. Liberzon, A., C. Birger, H. Thorvaldsdóttir, M. Ghandi, J.P. Mesirov, and P. Tamayo,
The Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell
systems, 2015. **1**(6): p. 417-425.

28. Walker, D.G. and L.F. Lue, *Immune phenotypes of microglia in human*
neurodegenerative disease: challenges to detecting microglial polarization in human
brains. Alzheimers Res Ther, 2015. **7**(1): p. 56.

29.   Hopperton, K.E., D. Mohammad, M.O. Trépanier, V. Giuliano, and R.P. Bazinet,
Markers of microglia in post-mortem brain samples from patients with Alzheimer's
disease: a systematic review. Molecular Psychiatry, 2018. **23**(2): p. 177-198.

30.   Hammond, T.R., C. Dufort, L. Dissing-Olesen, S. Giera, A. Young, A. Wysoker, A.J.
Walker, F. Gergits, M. Segel, J. Nemesh, S.E. Marsh, A. Saunders, E. Macosko, F.
Ginhoux, J. Chen, R.J.M. Franklin, X. Piao, S.A. McCarroll, and B. Stevens, *Single-*
Cell RNA Sequencing of Microglia throughout the Mouse Lifespan and in the Injured
Brain Reveals Complex Cell-State Changes. Immunity, 2019. **50**(1): p. 253-271.e6.

31.   McQuade, A., Y.J. Kang, J. Hasselmann, A. Jairaman, A. Sotelo, M. Coburn, S.K.
Shabestari, J.P. Chadarevian, G. Fote, C.H. Tu, E. Danhash, J. Silva, E. Martinez, C.
Cotman, G.A. Prieto, L.M. Thompson, J.S. Steffan, I. Smith, H. Davtyan, M. Cahalan,
H. Cho, and M. Blurton-Jones, *Gene expression and functional deficits underlie*
TREM2-knockout microglia responses in human models of Alzheimer's disease. Nature
Communications, 2020. **11**(1): p. 5370.

32.   Rothe, T., N. Ipseiz, M. Faas, S. Lang, F. Perez-Branguli, D. Metzger, H. Ichinose, B.
Winner, G. Schett, and G. Krönke, *The Nuclear Receptor Nr4a1 Acts as a Microglia*
Rheostat and Serves as a Therapeutic Target in Autoimmune-Driven Central Nervous
System Inflammation. Journal of immunology (Baltimore, Md. : 1950), 2017. **198**(10):
p. 3878-3885.

33.   Saijo, K., B. Winner, C.T. Carson, J.G. Collier, L. Boyer, M.G. Rosenfeld, F.H. Gage,
and C.K. Glass, *A Nurr1/CoREST pathway in microglia and astrocytes protects*
dopaminergic neurons from inflammation-induced death. Cell, 2009. **137**(1): p. 47-59.

34.   Szabo, P.A., H.M. Levitin, M. Miron, M.E. Snyder, T. Senda, J. Yuan, Y.L. Cheng,
E.C. Bush, P. Dogra, P. Thapa, D.L. Farber, and P.A. Sims, *Single-cell transcriptomics*
of human T cells reveals tissue and activation signatures in health and disease. Nature
Communications, 2019. **10**(1): p. 4706.

35.   Alegre, M.L., P.J. Noel, B.J. Eisfelder, E. Chuang, M.R. Clark, S.L. Reiner, and C.B.
Thompson, *Regulation of surface and intracellular expression of CTLA4 on mouse T*
cells. J Immunol, 1996. **157**(11): p. 4762-70.

36.   Johnson, D.B., M.J. Nixon, Y. Wang, D.Y. Wang, E. Castellanos, M.V. Estrada, P.I.
Ericsson-Gonzalez, C.H. Cote, R. Salgado, V. Sanchez, P.T. Dean, S.R. Opalenik, D.M.
Schreeder, D.L. Rimm, J.Y. Kim, J. Bordeaux, S. Loi, L. Horn, M.E. Sanders, P.B.
Ferrell, Jr., Y. Xu, J.A. Sosman, R.S. Davis, and J.M. Balko, *Tumor-specific MHC-II*
expression drives a unique pattern of resistance to immunotherapy via LAG-3/FCRL6
engagement. JCI insight, 2018. **3**(24): p. e120360.

37.   Sciumè, G., G. De Angelis, G. Benigni, A. Ponzetta, S. Morrone, A. Santoni, and G.
Bernardini, *CX3CR1 expression defines 2 KLRG1+ mouse NK-cell subsets with distinct*
functional properties and positioning in the bone marrow. Blood, 2011. **117**(17): p.
4467-4475.

38.   Böttcher, J.P., M. Beyer, F. Meissner, Z. Abdullah, J. Sander, B. Höchst, S. Eickhoff,
J.C. Rieckmann, C. Russo, T. Bauer, T. Flecken, D. Giesen, D. Engel, S. Jung, D.H.
Busch, U. Protzer, R. Thimme, M. Mann, C. Kurts, J.L. Schultze, W. Kastenmüller, and

P.A. Knolle, *Functional classification of memory CD8+ T cells by CX3CR1 expression.*
Nature Communications, 2015. **6**(1): p. 8306.

39. Gerlach, C., E.A. Moseman, S.M. Loughhead, D. Alvarez, A.J. Zwijnenburg, L.
Waanders, R. Garg, J.C. de la Torre, and U.H. von Andrian, *The Chemokine Receptor*
CX3CR1 Defines Three Antigen-Experienced CD8 T Cell Subsets with Distinct Roles
in Immune Surveillance and Homeostasis. Immunity, 2016. **45**(6): p. 1270-1284.

40. Yamauchi, T., T. Hoki, T. Oba, V. Jain, H. Chen, K. Attwood, S. Battaglia, S. George,
G. Chatta, I. Puzanov, C. Morrison, K. Odunsi, B.H. Segal, G.K. Dy, M.S. Ernstoff, and
F. Ito, *T-cell CX3CR1 expression as a dynamic blood-based biomarker of response to*
immune checkpoint inhibitors. Nature Communications, 2021. **12**(1): p. 1402.

41. Yan, Y., S. Cao, X. Liu, S.M. Harrington, W.E. Bindeman, A.A. Adjei, J.S. Jang, J. Jen,
Y. Li, P. Chanana, A.S. Mansfield, S.S. Park, S.N. Markovic, R.S. Dronca, and H.
Dong, *CX3CR1 identifies PD-1 therapy-responsive CD8+ T cells that withstand*
chemotherapy during cancer chemoimmunotherapy. JCI Insight, 2018. **3**(8).

42. Urban, S.L., I.J. Jensen, Q. Shan, L.L. Pewe, H.-H. Xue, V.P. Badovinac, and J.T. Harty,
Peripherally induced brain tissue–resident memory CD8+ T cells mediate protection
against CNS infection. Nature Immunology, 2020. **21**(8): p. 938-949.

43. Mackay, L.K., M. Minnich, N.A. Kragten, Y. Liao, B. Nota, C. Seillet, A. Zaid, K. Man,
S. Preston, D. Freestone, A. Braun, E. Wynne-Jones, F.M. Behr, R. Stark, D.G. Pellicci,
D.I. Godfrey, G.T. Belz, M. Pellegrini, T. Gebhardt, M. Busslinger, W. Shi, F.R.
Carbone, R.A. van Lier, A. Kallies, and K.P. van Gisbergen, *Hobit and Blimp1 instruct*
a universal transcriptional program of tissue residency in lymphocytes. Science, 2016.
352(6284): p. 459-63.

44. Ma, J., B. Zheng, S. Goswami, L. Meng, D. Zhang, C. Cao, T. Li, F. Zhu, L. Ma, Z.
Zhang, S. Zhang, M. Duan, Q. Chen, Q. Gao, and X. Zhang, *PD1(Hi) CD8(+) T cells*
correlate with exhausted signature and poor clinical outcome in hepatocellular
carcinoma. Journal for immunotherapy of cancer, 2019. **7**(1): p. 331-331.

45. Liu, X., Y. Wang, H. Lu, J. Li, X. Yan, M. Xiao, J. Hao, A. Alekseev, H. Khong, T.
Chen, R. Huang, J. Wu, Q. Zhao, Q. Wu, S. Xu, X. Wang, W. Jin, S. Yu, Y. Wang, L.
Wei, A. Wang, B. Zhong, L. Ni, X. Liu, R. Nurieva, L. Ye, Q. Tian, X.-W. Bian, and
C. Dong, *Genome-wide analysis identifies NR4A1 as a key mediator of T cell*
dysfunction. Nature, 2019. **567**(7749): p. 525-529.

46. Dan, L., L. Liu, Y. Sun, J. Song, Q. Yin, G. Zhang, F. Qi, Z. Hu, Z. Yang, Z. Zhou, Y.
Hu, L. Zhang, J. Ji, X. Zhao, Y. Jin, M.A. McNutt, and Y. Yin, *The phosphatase PAC1*
acts as a T cell suppressor and attenuates host antitumor immunity. Nature
Immunology, 2020. **21**(3): p. 287-297.

47. Bignon, A., A. Régent, L. Klipfel, A. Desnoyer, P. de la Grange, V. Martinez, O.
Lortholary, A. Dalloul, L. Mouthon, and K. Balabanian, *DUSP4-mediated accelerated*
T-cell senescence in idiopathic CD4 lymphopenia. Blood, 2015. **125**(16): p. 2507-18.

48. Maine, C.J., J.R. Teijaro, K. Marquardt, and L.A. Sherman, *PTPN22 contributes to*
exhaustion of T lymphocytes during chronic viral infection. Proceedings of the National
Academy of Sciences, 2016. **113**(46): p. E7231-E7239.

49.   Li, H., A.M. van der Leun, I. Yofe, Y. Lubling, D. Gelbard-Solodkin, A.C.J. van
840   Akkooi, M. van den Braber, E.A. Rozeman, J.B.A.G. Haanen, C.U. Blank, H.M.
841   Horlings, E. David, Y. Baran, A. Bercovich, A. Lifshitz, T.N. Schumacher, A. Tanay,
842   and I. Amit, *Dysfunctional CD8 T Cells Form a Proliferative, Dynamically Regulated*
843   *Compartment within Human Melanoma.* Cell, 2019. **176**(4): p. 775-789.e18.

844   50.   Jin, S., C.F. Guerrero-Juarez, L. Zhang, I. Chang, R. Ramos, C.-H. Kuan, P. Myung,
845   M.V. Plikus, and Q. Nie, *Inference and analysis of cell-cell communication using*
846   *CellChat.* Nature Communications, 2021. **12**(1): p. 1088.

847   51.   Wei, J., A. Marisetty, B. Schrand, K. Gabrusiewicz, Y. Hashimoto, M. Ott, Z. Grami,
848   L.Y. Kong, X. Ling, H. Caruso, S. Zhou, Y.A. Wang, G.N. Fuller, J. Huse, E. Gilboa,
849   N. Kang, X. Huang, R. Verhaak, S. Li, and A.B. Heimberger, *Osteopontin mediates*
850   *glioblastoma-associated macrophage infiltration and is a potential therapeutic target.*
851   J Clin Invest, 2019. **129**(1): p. 137-149.

852   52.   Klement, J.D., A.V. Paschall, P.S. Redd, M.L. Ibrahim, C. Lu, D. Yang, E. Celis, S.I.
853   Abrams, K. Ozato, and K. Liu, *An osteopontin/CD44 immune checkpoint controls*
854   *CD8+ T cell activation and tumor immune evasion.* J Clin Invest, 2018. **128**(12): p.
855   5549-5560.

856   53.   Olah, M., V. Menon, N. Habib, M.F. Taga, Y. Ma, C.J. Yung, M. Cimpean, A.
857   Khairallah, G. Coronas-Samano, R. Sankowski, D. Grün, A.A. Kroshilina, D. Dionne,
858   R.A. Sarkis, G.R. Cosgrove, J. Helgager, J.A. Golden, P.B. Pennell, M. Prinz, J.P.G.
859   Vonsattel, A.F. Teich, J.A. Schneider, D.A. Bennett, A. Regev, W. Elyaman, E.M.
860   Bradshaw, and P.L. De Jager, *Single cell RNA sequencing of human microglia uncovers*
861   *a subset associated with Alzheimer's disease.* Nature Communications, 2020. **11**(1): p.
862   6129.

863   54.   Shekarian, T., C.P. Zinner, E.M. Bartoszek, W. Duchemin, A.T. Wachnowicz, S.
864   Hogan, M.M. Etter, J. Flammer, C. Paganetti, T.A. Martins, P. Schmassmann, S.
865   Zanganeh, F. Le Goff, M.G. Muraro, M.-F. Ritz, D. Phillips, S.S. Bhate, G.L. Barlow,
866   G.P. Nolan, C.M. Schürch, and G. Hutter, *Immunotherapy of glioblastoma explants*
867   *induces interferon-γ responses and spatial immune cell rearrangements in*
868   *tumor center, but not periphery.* Science Advances, 2022. **8**(26): p. eabn9440.

869   55.   Smolders, J., K.M. Heutinck, N.L. Fransen, E.B.M. Remmerswaal, P. Hombrink, I.J.M.
870   ten Berge, R.A.W. van Lier, I. Huitinga, and J. Hamann, *Tissue-resident memory T cells*
871   *populate the human brain.* Nature Communications, 2018. **9**(1): p. 4593.

872   56.   Kumar, B.V., W. Ma, M. Miron, T. Granot, R.S. Guyer, D.J. Carpenter, T. Senda, X.
873   Sun, S.H. Ho, H. Lerner, A.L. Friedman, Y. Shen, and D.L. Farber, *Human Tissue-*
874   *Resident Memory T Cells Are Defined by Core Transcriptional and Functional*
875   *Signatures in Lymphoid and Mucosal Sites.* Cell Rep, 2017. **20**(12): p. 2921-2934.

876   57.   Mackay, L.K., A. Rahimpour, J.Z. Ma, N. Collins, A.T. Stock, M.L. Hafon, J. Vega-
877   Ramos, P. Lauzurica, S.N. Mueller, T. Stefanovic, D.C. Tscharke, W.R. Heath, M.
878   Inouye, F.R. Carbone, and T. Gebhardt, *The developmental pathway for*
879   *CD103(+)CD8+ tissue-resident memory T cells of skin.* Nat Immunol, 2013. **14**(12): p.
880   1294-301.

58. Gandoglia, I., F. Ivaldi, P. Carrega, E. Armentani, G. Ferlazzo, G. Mancardi, N. Kerlero
de Rosbo, A. Uccelli, and A. Laroni, *In vitro VLA-4 blockade results in an impaired NK*
cell-mediated immune surveillance against melanoma. Immunology Letters, 2017. **181**:
p. 109-115.

59. Sankowski, R., C. Böttcher, T. Masuda, L. Geirsdottir, Sagar, E. Sindram, T.
Seredenina, A. Muhs, C. Scheiwe, M.J. Shah, D.H. Heiland, O. Schnell, D. Grün, J.
Priller, and M. Prinz, *Mapping microglia states in the human brain through the*
integration of high-dimensional techniques. Nature Neuroscience, 2019. **22**(12): p.
2098-2110.

60. Kilian, M., R. Sheinin, C.L. Tan, M. Friedrich, C. Krämer, A. Kaminitz, K. Sanghvi, K.
Lindner, Y.-C. Chih, F. Cichon, B. Richter, S. Jung, K. Jähne, M. Ratliff, R.M. Prins,
N. Etminan, A. von Deimling, W. Wick, A. Madi, L. Bunse, and M. Platten, *MHC class*
II-restricted antigen presentation is required to prevent dysfunction of cytotoxic T cells
by blood-borne myeloids in brain tumors. Cancer Cell, 2023. **41**(2): p. 235-251.e9.

61. Srivastava, A., L. Malik, T. Smith, I. Sudbery, and R. Patro, *Alevin efficiently estimates*
accurate gene abundances from dscRNA-seq data. Genome Biol, 2019. **20**(1): p. 65.

62. Griffiths, J.A., A.C. Richard, K. Bach, A.T.L. Lun, and J.C. Marioni, *Detection and*
removal of barcode swapping in single-cell RNA-seq data. Nat Commun, 2018. **9**(1): p.
2667.

63. Lun, A.T.L., S. Riesenfeld, T. Andrews, T.P. Dao, T. Gomes, J. participants in the 1st
Human Cell Atlas, and J.C. Marioni, *EmptyDrops: distinguishing cells from empty*
droplets in droplet-based single-cell RNA sequencing data. Genome Biol, 2019. **20**(1):
p. 63.

64. Vallejos, C.A., D. Risso, A. Scialdone, S. Dudoit, and J.C. Marioni, *Normalizing single-*
cell RNA sequencing data: challenges and opportunities. Nat Meth, 2017. **14**(6): p. 565-
571.

65. Lun, A., D. McCarthy, and J. Marioni, *A step-by-step workflow for low-level analysis*
of single-cell RNA-seq data with Bioconductor [version 2; referees: 3 approved, 2
approved with reservations]. F1000Research, 2016. **5**(2122).

66. McCarthy, D.J., K.R. Campbell, Q.F. Wills, and A.T.L. Lun, *Scater: pre-processing,*
quality control, normalization and visualization of single-cell RNA-seq data in R.
Bioinformatics, 2017. **33**(8): p. 1179-1186.

67. Amezquita, R.A., A.T.L. Lun, E. Becht, V.J. Carey, L.N. Carpp, L. Geistlinger, F.
Martini, K. Rue-Albrecht, D. Risso, C. Soneson, L. Waldron, H. Pages, M.L. Smith, W.
Huber, M. Morgan, R. Gottardo, and S.C. Hicks, *Orchestrating single-cell analysis with*
Bioconductor. Nat Methods, 2019.

68. Lun, A.T., D.J. McCarthy, and J.C. Marioni, *A step-by-step workflow for low-level*
analysis of single-cell RNA-seq data with Bioconductor. F1000Res, 2016. **5**: p. 2122.

69. Lütge, A., J. Zyprych-Walczak, U. Brykczynska Kunzmann, H.L. Crowell, D. Calini,
D. Malhotra, C. Soneson, and M.D. Robinson, *CellMixS: quantifying and visualizing*
batch effects in single-cell RNA-seq data. Life Science Alliance, 2021. **4**(6): p.
e202001004.

70. Murtagh, F. and P. Legendre, *Ward's Hierarchical Agglomerative Clustering Method:*
Which Algorithms Implement Ward's Criterion? J. Classif., 2014. **31**(3): p. 274–295.

71. Langfelder, P., B. Zhang, and S. Horvath, *Defining clusters from a hierarchical cluster*
tree: the Dynamic Tree Cut package for R. Bioinformatics, 2008. **24**(5): p. 719-20.

72. Hao, Y., S. Hao, E. Andersen-Nissen, W.M. Mauck, S. Zheng, A. Butler, M.J. Lee, A.J.
Wilk, C. Darby, M. Zager, P. Hoffman, M. Stoeckius, E. Papalexi, E.P. Mimitou, J.
Jain, A. Srivastava, T. Stuart, L.M. Fleming, B. Yeung, A.J. Rogers, J.M. McElrath,
C.A. Blish, R. Gottardo, P. Smibert, and R. Satija, *Integrated analysis of multimodal*
single-cell data. Cell, 2021. **184**(13): p. 3573-3587.e29.

73. Aran, D., A.P. Looney, L. Liu, E. Wu, V. Fong, A. Hsu, S. Chak, R.P. Naikawadi, P.J.
Wolters, A.R. Abate, A.J. Butte, and M. Bhattacharya, *Reference-based analysis of lung*
single-cell sequencing reveals a transitional profibrotic macrophage. Nat Immunol,
2019. **20**(2): p. 163-172.

74. Law, C.W., Y. Chen, W. Shi, and G.K. Smyth, *voom: precision weights unlock linear*
model analysis tools for RNA-seq read counts. Genome Biology, 2014. **15**(2): p. R29.

75. Weber, L.M., M. Nowicka, C. Soneson, and M.D. Robinson, *diffcyt: Differential*
discovery in high-dimensional cytometry via high-resolution clustering.
Communications Biology, 2019. **2**(1): p. 183.

76. Dann, E., N.C. Henderson, S.A. Teichmann, M.D. Morgan, and J.C. Marioni,
Differential abundance testing on single-cell data using k-nearest neighbor graphs.
Nature Biotechnology, 2022. **40**(2): p. 245-253.

77. Junttila, S., J. Smolander, and L.L. Elo, *Benchmarking methods for detecting*
differential states between conditions from multi-subject single-cell RNA-seq data.
Briefings in Bioinformatics, 2022. **23**(5).

78. Lun, A.T.L. and J.C. Marioni, *Overcoming confounding plate effects in differential*
expression analyses of single-cell RNA-seq data. Biostatistics, 2017. **18**(3): p. 451-464.

79. Robinson, M.D., D.J. McCarthy, and G.K. Smyth, *edgeR: a Bioconductor package for*
differential expression analysis of digital gene expression data. Bioinformatics, 2010.
26(1): p. 139-140.

80. Robinson, M.D. and A. Oshlack, *A scaling normalization method for differential*
expression analysis of RNA-seq data. Genome Biol, 2010. **11**(3): p. R25.

81. Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and*
dispersion for RNA-seq data with DESeq2. Genome Biology, 2014. **15**(12): p. 550.

82. Wu, D. and G.K. Smyth, *Camera: a competitive gene set test accounting for inter-gene*
correlation. Nucleic Acids Res, 2012. **40**(17): p. e133.

83. Subramanian, A., P. Tamayo, V.K. Mootha, S. Mukherjee, B.L. Ebert, M.A. Gillette,
A. Paulovich, S.L. Pomeroy, T.R. Golub, E.S. Lander, and J.P. Mesirov, *Gene set*
enrichment analysis: A knowledge-based approach for interpreting genome-wide
expression profiles. Proc Natl Acad Sci U S A, 2005. **102**(43): p. 15545-15550.

Acknowledgements

We are grateful to the patients and their families for their consent to donate tissue to our brain tumor biobank. We thank Tamara Hüssen, Alison Riberio and Florian Limani for experimental support and Cecile Buenter for critical revision of the manuscript. Calculations were performed at sciCORE (http://scicore.unibas.ch/) scientific computing center at the University of Basel.

Author contribution

G.H., S. Herter, M.B., S.D. and P.S., conceived and planned the project. J.R. and S. Hogan analyzed the scRNA-sequencing data and performed statistical analysis. P.S. and S.D. performed experiments and interpreted the results. T.S., T.A.M. and M.-F.R. helped coordinating experiments. P.S. wrote the manuscript. G.H. supervised and coordinated the study and critically revised the manuscript. All authors reviewed the paper and approved its final version.

Competing interests

G.H. has equity in, and is a cofounder of Incephalo Inc. S.D., S.H. and M.B. are affiliated with Roche Pharmaceutical Research and Early Development. The authors have no financial interests to declare.

Main Figures

 Figure 1 Single-cell RNA-seq of cells from tumor center, periphery and blood.

(**A**) Experimental workflow for single-cell analysis of cells isolated from tumor center, periphery and peripheral blood mononuclear cells (PBMC), including fluorescent-activated cell sorting and 3'-scRNA-seq. (**B**) Axial T1 with contrast (left) and T2 (right) MRI brain in a patient with a left temporal GBM. Fresh tumor biopsies were taken according to neuronavigation (green cross). The tumor center was defined as contrast enhancing, whereas the tumor periphery was defined as T2 hyperintense. (**C**) Nuclear DAPI staining of resected tissue specimens. 40x magnification (scale bar = 20 μm). n = 3 patients, 4 field of view (FOV) per patient. Statistics: ***$p < 0.001$, two-tailed Mann Whitney U test (Figure 1 – source data 1).

991 **Figure 2 Single-cell RNA-seq analysis identifies main immune cell populations.**

992 **(A)** Dimensionally reduced tSNE projection of the scRNAseq data showing the annotated cell

993 types. **(B)** Heatmap displaying centered and scaled normalized average expression values of

994 characteristic cell-type specific genes used to annotate clusters. Columns are ordered by site

and cell type, and rows show centered and scaled expression values, hierarchically clustered. Heatmap displaying genes whose expression is most specific to each cell type is shown in Figure 2 – figure supplement 2. (**C**) Principal component analysis of pseudo-bulk scRNAseq samples aggregated by patient and cell type. Symbols represent individual patients and cell lineage is displayed by different colors. (**D)** Relative frequencies of immune populations among leukocytes between tumor center and periphery. Symbols represent individual patients and paired samples are indicated by connecting lines. *P*-values were calculated using *diffcyt-DA-voom* method (Figure 2 – source data 1). (**E-G**) Differential abundance testing of the tumor innate immune compartment (**E**) using the *miloR* package which tests the abundance of each neighborhood of cells separately between tumor center und periphery (**F, G**).

Figure 3 MG display regionally resolved transcriptional profiles that differ from those of DCs and MdMs.

(A) Microglia cluster highlighted on tSNE map and scatterplots showing differentially expressed genes (FDR<5%, indicated by blue and yellow) in Microglia (MG) cells from tumor periphery versus center. Volcano plot showing p value versus fold-change (left) and MA plot showing fold-change versus mean expression (right). For a complete list of differentially expressed genes per cell cluster between tumor periphery and center, please refer to Supplementary File 4. **(B)** Heatmap representation of Gene set enrichment analysis (GSEA) results between peripheral and center microglia using Gene Ontology (GO) collection (Biological Processes). The fraction of overlap between gene sets is calculated as Jaccard coefficient of overlap between the gene sets. **(C, D)** Heatmap representation of GSEA of DCs **(C)** and monocyte-derived macrophages (MdMs) **(D)** from tumor periphery versus tumor center using Hallmark collection of major biological categories. **(E)** Unsupervised hierarchical sub-clustering of the MG population revealed two transcriptionally distinct subsets of MG, termed MG_1 and MG_2, displayed on the tSNE map. **(F)** Heatmap displaying the cluster-specific genes identifying MG_1 subcluster. Columns are ordered by site and cell type, and rows show centered and scaled normalized average expression values, hierarchically clustered. A complete list of cluster specific genes for MG_1 and MG_2 subcluster is provided in Supplementary File 5. **(G)** Heatmap representation of GSEA between MG_1 and MG_2 subclusters using Hallmark collection of major biological categories. **(H)** Heatmap displaying previously described reactivity markers of MG. Columns are ordered by site and cell type, and rows show centered and scaled normalized average expression values, hierarchically clustered.

1028

Figure 4 The peripheral cytotoxic cell compartment exhibits an impaired activation sig-

nature.

(A, B) Volcano plots showing differentially expressed genes (FDR corrected *p* value < 0.05,

indicated by blue and yellow) in CD8$^+$ T cells **(A)** and NK cells **(B)** from tumor periphery versus

tumor center. Colored rings mark genes belonging to selected GSEA Hallmark or Gene Ontol-

ogy (GO) pathways as indicated. For a complete list of differentially expressed genes per cell

cluster between tumor periphery and center, please refer to Supplementary File 4.

Figure 5 CX3CR1 labels a specific CD8+ T cell population in the circulation of grade 4 glioma patients.

(**A**) CD8$^+$ T cell cluster highlighted on tSNE map (left). CD8$^+$ T cell cluster colored by site of origin (right). (**B**) Volcano plot showing differentially expressed genes (FDR corrected *p* value < 0.05, indicated by blue and green) in CD8$^+$ T cells from tumor-periphery versus PBMC. For a complete list of differentially expressed genes per cell cluster between tumor periphery and PBMC, please refer to Supplementary File 6. (**C**) Frequency of CX3CR1$^+$ CD8$^+$ T cells among all CD8$^+$ T cells in flow cytometry data (Figure 5 – source data 1). (**D**) Unsupervised hierarchical sub-clustering of CD8$^+$ T cells from PBMC and Periphery revealed two transcriptionally

distinct subsets of PBMC CD8$^+$ T cells, displayed on the tSNE map. (**E**) Expression of *CX3CR1*

overlaid on tSNE CD8$^+$ T cell cluster. (**F**) Expression of genes associated with effector memory

phenotype overlaid on tSNE CD8$^+$ T cell cluster. Displayed genes are significantly, differen-

tially expressed genes (DEGs) between tumor periphery and PBMC, as identified by differen-

tial gene expression analysis shown in panel (B). (**G**) Expression of selected genes associated

with naive phenotype overlaid on tSNE CD8$^+$ T cell cluster (**H**) Gating procedure applied to

identify CD3$^+$ CD8$^+$ naïve, T effector cells (T$_{eff}$), effector memory (T$_{em}$), peripheral memory

(T$_{pm}$) and central memory (T$_{cm}$), eluted from PBMCs. (**I**) Expression of CX3CR1 in PBMC

CD8$^+$ T cell subpopulations identified in (H) (Figure 5 – source data 2). $n = 6$ donors (**C**), $n =$

11 donors (**I**). Statistics: Wilcoxon matched-pairs signed rank test (**C**); repeated measures one-

way ANOVA with post-hoc Šidák's correction for multiple comparisons (**I**). *$p \leq 0.05$, **$p \leq$

0.01, ***$p \leq 0.001$, ****$p \leq 0.0001$, no brackets indicate no significant difference.

1058

Figure 6 CD8$^+$ T cells in the tumor periphery share features with tissue-resident memory

T cells (Trm)

(**A**) Representative dot plot of tumor-periphery CD8$^+$ T cells stained for CD45RA and CD45RO. (**B**) Quantification of tumor-periphery CD8$^+$ T cells expressing CD45RA or CD45RO (Figure 6 – source data 1). (**C**) Expression of genes associated with tissue-resident memory (T$_{rm}$) phenotype overlaid on tSNE CD8$^+$ T cell cluster. (**D**) Average expression levels of selected T$_{rm}$ markers between CD8$^+$ T cells from PBMC versus tumor-periphery.

Significance testing based on differential gene expression analysis shown in panel (Fig. 5B)

(**E**) Representative dot plots of CD69 and CD103 co-expression in CD8$^+$ T cells from PBMC

and tumor-periphery. (**F**) Quantification of CD69 and CD103 co-expression revealed CD69$^-$

CD103$^-$ in PBMC and CD69$^+$ CD103$^-$ and CD69$^+$ CD103$^+$ in tumor-periphery as the dominant

phenotypes (Figure 6 – source data 2). (**G**) Expression of selected markers associated with T

cell exhaustion/dysfunction, shown as boxplots between CD8$^+$ T cell from PBMC and tumor-

periphery and overlaid on tSNE CD8$^+$ T cell cluster. n = 6 donors (**B, F**). Statistics: Wilcoxon

matched-pairs signed rank test (**B**); repeated measures one-way ANOVA with post-hoc Šidák's

correction for multiple comparisons (**F**). *$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$, ****$p \leq 0.0001$,

no brackets indicate no significant difference.

1076

Figure 7 Cell-cell communication analysis using CellChat reveals critical role for SPP1-mediated crosstalk in tumor periphery.

(A) Chord diagram showing significant interactions from microglia to lymphocyte cell clusters. The inner bar colors represent the targets that receive signal from the corresponding outer bar. The inner bar size is proportional to the signal strength received by the targets. Chords indicate ligand-receptor pairs mediating interaction between two cell clusters, size of chords is proportional to signal strength of the given ligand-receptor pair. **(B)** Comparison of incoming and outgoing interaction strength allows identification of main senders and receivers. **(C)** Violin plots showing the expression distribution of signaling genes involved in the inferred SPP1 signaling network.