The mechanosensitive ion channel Piezo1 modulates the migration and immune response of microglia

Ting Zhu[1], Shashwati Kala[1], Jinghui Guo[1], Yong Wu[1], Hui Chen[2], Jiejun Zhu[1], Kin Fung Wong[1], Chi Pong Cheung[1], Xiaohui Huang[1], Xinyi Zhao[1], Ting Lei[1], Minyi Yang[1] and Lei Sun[1]*

[1]Department of Biomedical Engineering, The Hong Kong Polytechnic University, Hung Hom, Hong Kong SAR, P. R. China, 999077

[2]Biotherapy Centre, The Third Affiliated Hospital, Sun Yat-sen University, Guangzhou, China; Cell-gene Therapy Translational Medicine Research Centre, The Third Affiliated Hospital, Sun Yat-sen University, Guangzhou, China.

*Correspondence to:

Lei Sun, Ph.D

Room ST409, Department of Biomedical Engineering,

The Hong Kong Polytechnic University,

Hong Kong SAR, China, People's Republic of.

Phone: (852) 2766 7663

Email: lei.sun@polyu.edu.hk

Abstract

Microglia are the brain's resident immune cells, performing surveillance to promote homeostasis and healthy functioning. While chemical signaling in microglia is well-studied, the mechanical cues regulating their function are less well-understood. Here, we investigate the role of the mechanosensitive ion channel Piezo1 in microglia migration, pro-inflammatory cytokine production and stiffness sensing. We demonstrated the functional expression of Piezo1 in microglia and identified genes whose expression was affected by conditional Piezo1 knockout in transgenic mice. Functional assays revealed that Piezo1-deficiency in microglia enhanced migration towards amyloid β-protein, and decreased levels of pro-inflammatory cytokines produced upon stimulation by lipopolysaccharide, both *in vitro* and *in vivo*. The phenomenon could be mimicked or reversed using a Piezo1-specific agonist or antagonist. Finally, we also showed that Piezo1 mediated the effect of substrate stiffness-induced migration and cytokine expression. Altogether, we show that Piezo1 is an important molecular mediator for microglia, its activation modulating their migration and immune responses.

Keywords

Microglia, migration, immune, Piezo1, mechanosensing

Introduction

As the resident innate immune cells of the central nervous system (CNS), microglia play pivotal roles in maintaining brain homeostasis, pathogen defense and repairing injuries (Matcovitch-Natan, Winter et al. 2016). Microglia can recognize pathogens via pathogen-associated molecular patterns (PAMPs) on their surface (Kigerl, de Rivero Vaccari et al. 2014) and migrate to the sites of damage along a chemo-attractant gradient (Fan, Xie et al. 2017). They are able to secrete chemicals, e.g., inflammatory cytokines (Aloisi 2001) and neurotrophic factors (Nakajima, Tohyama et al. 2007) in order to protect the brain. In addition to responding to biochemical stimuli, physical cues including tissue stiffness and mechanical stimulation are also thought to contribute to microglial activity (Moshayedi, Ng et al. 2014, Bollmann, Koser et al. 2015, Ayata and Schaefer 2020). Previous studies have shown that stiff surgical implants in the CNS led to a localized foreign body reaction (FBR) by enhancing inflammatory activation of glial cells (Moshayedi, Ng et al. 2014). Putting aside the artificial manipulation of CNS microenvironmental stiffness, brain tissue itself is mechanically heterogeneous (Elkin, Azeloglu et al. 2007), and these mechanical properties may alter with age (Segel, Neumann et al. 2019) or pathological conditions (Murphy, Huston et al. 2011, Riek, Millward et al. 2012, Schregel, Wuerfel et al. 2012, Streitberger, Sack et al. 2012). Thus, microglia are exposed to varying mechanical conditions through their life cycle, but little is known about molecular mechanism how these stimuli may modulate microglial activity.

A growing body of evidence implicates the roles of the cytoskeleton, cell

adhesion and Rho signaling in the process of mechanosensation and

mechanotransduction (Ohashi, Fujiwara et al. 2017). However, located in lipid bilayers

of cell membranes, mechanically gated ion channels serve as primary mediators of

cellular mechanosensation, and merit deeper examination as well (Lim, Jang et al.

2018). Upon the increase of membrane tension, mechanically-activated ion channels

undergo conformational changes and allow the flow of ions across the membrane,

sensing and transducing external physical stimuli into electrochemical activity that

further influences cell signaling and behavior (Mobasheri, Carter et al. 2002, Wang and

Thampatty 2006). Of particular note is Piezo1, a mechanosensitive cation ion channel

that detects mechanical forces with high sensitivity and broad dynamic range (Syeda,

Florendo et al. 2016). Piezo1 has been reported to be expressed at high levels in

endothelial cells and microglia of the brain according to previous RNA sequencing

results (Zhang, Chen et al. 2014). Substantial recent research reveals that Piezo1

transduces mechanical cues to cells (Coste, Mathur et al. 2010) and is involved in many

physiological and pathological process (Pathak, Nourse et al. 2014, Cahalan, Lukacs et

al. 2015, Chen, Wanggou et al. 2018, Nonomura, Lukacs et al. 2018, Romac, Shahid et

al. 2018, Solis, Bielecki et al. 2019). Therefore, Piezo1 is a potential mechanosensor

and may be involved in coordinating the functions of microglia. In light of this evidence,

we hypothesized that Piezo1 may play important roles in the functioning of microglia.

In the present study, microglia from mouse brains, *in vivo* and *ex vivo*, and a

mouse microglial cell line were used as models for examining Piezo1 expression and

its role in microglial functions. RT-PCR analysis showed that Piezo1 was expressed at

high levels in primary microglia compared to other mechanosensitive ion channels.

Gene clusters affected by the conditional knockout (KO) of Piezo1 in microglia were

identified by RNA sequencing (RNAseq) and were found to include genes associated

with cell motility and extracellular matrix composition. Functional analyses confirmed

that Piezo1 deficiency in microglia resulted in enhanced migration ability towards $A\beta_{1-42}$, a major component of Alzheimer's Disease plaques. Moreover, activation of Piezo1

could suppress the migration of microglia, while blockade of Piezo1 enhanced it. We

also found that microglia lacking Piezo1 showed decreased pro-inflammatory cytokines

(IL-1β, IL-6 and TNF-α) production when challenged by lipopolysaccharide (LPS).

Activation of Piezo1 by Yoda1, a Piezo1-specific agonist, increased production of these

pro-inflammatory cytokines production by microglia when stimulated by LPS.

Variation in microglial substrate stiffness was mimicked *in vitro* through modifiable

gels, and we found that the expected stiffness-dependent reduction in migration and

increase in pro-inflammatory cytokines production by microglia were both Piezo1-

dependent. Interestingly, we also found the expression of Piezo1 was itself enhanced in

microglia cultured on stiffer substrates. In summary, our study reveals that Piezo1 is a

mechanosensor of stiffness in microglia, and that its activity modulates the migratory

tendencies and immune response of microglia.

Results

Piezo1 is functionally expressed in microglial cells

We first attempted to confirm the expression of Piezo1 in primary microglia. Microglia from C57BL/6J mouse pups at postnatal day 3 (P3) were isolated by magnetic-activated cell sorting (MACS) and the purity was checked using flow cytometry (Fig. S1). The expression of various well-established mechanosensitive ion channels was then compared through RT-PCR (Ohana, Newell et al. 2009, Konno, Shirakawa et al. 2012, Shi, Du et al. 2013, Echeverry, Rodriguez et al. 2016, Cojocaru, Burada et al. 2021). Piezo1 was the channel most abundantly expressed, and its expression was significantly higher than its homolog Piezo2 (Coste, Mathur et al. 2010) (Fig. 1A). Functional analysis of Piezo1 in primary microglia cultured for 5 days was performed by patch clamping cells treated with a Piezo1-specific agonist Yoda1 (Syeda, Xu et al. 2015) and the Piezo1 blocker GsMTx4 (Bae, Sachs et al. 2011). 30 µM Yoda1 could induce strong inward current in microglia (462.7 ± 182.1 pA), but pre-treatment with 1 µM GsMTx-4 significantly reduced the currents induced (40.4 ± 15.36 pA) (Fig. 1B-C). Piezo1 being a cation ion channel with high permeability to Ca^{2+}, calcium imaging was performed on cells loaded with a fluorescent indicator (Fura-2). We found that Yoda1 at different concentrations (2 and 10 µM) could induce robust intracellular calcium increase in primary microglia, and adding GsMTx4 significantly decreased the Ca^{2+} influx induced by 10 µM Yoda1 (Fig. 1D-F).

We also tested the calcium response patterns in a commonly-used mouse

microglial cell line, BV2. Yoda1 significantly increased intracellular calcium levels in

BV2, and the responses were dose dependent, but GsMTx4 (1µM) inhibited the calcium

increase induced by 10 µM Yoda1 (Fig. S2A-C). Thus, we concluded that Piezo1 was

expressed, and that the channels were functional, in the cellular models of microglia

chosen for our experiments.

Fig. 1. Piezo1 is functional expressed in primary microglia.
A. The relative expression of mechanically- activated ion channel messenger RNA in primary microglia was analyzed by real-time RT-PCR using the ΔΔCT method. Bars represent mean ± SEM. N=3 independent experiments. Two-tailed unpaired t-test. *p < 0.05.
B. Representative traces of Yoda1-induced inward currents in primary microglia

recorded at −60 mV under the indicated conditions.

C. Scatter plots of Yoda1-induced inward currents in the indicated conditions. Bars

represent mean ± SEM. N=4 in Yoda1 group, n=5 in GsMTx4+Yoda1 group. Two-tailed

unpaired t-test. *p < 0.05.

D. Representative live-cell imaging of primary microglia loaded with Fura-2 at rest

state (upper panel) and at the time of maximum fluorescence (lower panel). Scale bar=

20μm.

E. Temporal raster plots of fluorescence changes in primary microglia treated with

different Yoda1 and/or GsMTx4 concentrations. Dashed lines indicate when Yoda1 was

added.

F. Statistical analysis of ΔF/F0 (%) in primary microglia under different Yoda1 and/or

GsMTx4 concentrations. Bars represent the mean ± SD. N= 3 independent experiments;

20-30 cells analyzed per experiment. Two-tailed Mann–Whitney U test. ****p < 0.0001.

Piezo1 deficiency affects expression profiles of genes

Piezo1 has been documented to play key roles in some important physiological

functions, e.g., lymphatic valve formation (Nonomura, Lukacs et al. 2018), neuronal

sensing of blood pressure (Zeng, Marshall et al. 2018), and red blood cell volume

homeostasis (Cahalan, Lukacs et al. 2015). Having confirmed the robust expression of

Piezo1 in primary microglia, we were interested in exploring the role of Piezo1 in

regulating microglial functions. Knockout or knockdown models of genes are a crucial

way to study their roles in various cells. Global deletion of the Piezo1 gene is known to

be lethal to embryos (Li, Hou et al. 2014); thus, we instead generated transgenic mice

using a conditional knockout model. A mouse line having Piezo1 conditionally depleted

from Tmem119-expressing CNS microglia was generated, of the genotype $Piezo1^{flox/flox}$

$Tmem119^{CreERT2/+}$ ($Piezo1^{\Delta Tmem119}$) (Fig. 2A). Mice of the genotype

$Piezo1^{flox/+}Tmem119^{CreERT2/+}$ ($Piezo1^{fl/+}$) were used as a control group for comparison.

Genotyping and Piezo1 knock-out validation results were performed by PCR to confirm

successful deletion (Fig. S3). To evaluate the consequences of Piezo1 loss, primary

microglia were isolated from brains of adult mice by MACS (Fig. 2B). Total RNA was

extracted immediately after sorting and was commercially analyzed for changes in their

transcriptional profile using RNAseq. The analysis revealed that 621 genes were

upregulated and 293 were downregulated in Piezo1$^{\Delta Tmem119}$ compared to the control

(Fig. 2C-D). A gene ontology (GO) pathway analysis was performed, and the top 20

significantly enriched terms were compiled. The analysis revealed several clusters of

dysregulated genes, including genes responsible for extracellular matrix organization

and regulating cellular motility and migration (Fig. 2E). We thus found evidence that

Piezo1 is involved in regulating cell motility, a key aspect of microglial behavior.

Fig. 2. Piezo1 deficiency in microglia affects expression profiles of genes.

A. Schematic illustration of the Piezo1 conditional knock out mice and treatment scheme with tamoxifen.

B. Schematic illustration of RNA-seq preparation procedure.

C. Volcano plot showing Piezo1$^{\Delta \text{Tmem119}}$ vs. Piezo1$^{\text{fl/+}}$ differentially expressed genes by RNA-seq based on primary microglia isolated from Piezo1 conditional knock out mice.

D. Heat map showing the top 50 differentially expressed genes, Piezo1$^{\Delta \text{Tmem119}}$ vs. Piezo1$^{\text{fl/+}}$.

E. The top 20 significantly enriched terms in the gene ontology (GO) enrichment analysis.

Piezo1 regulates migration ability of microglia

Based on the RNAseq data, we were interested in directly investigating the impact

of Piezo1 on microglial migration. We first tested whether the migration of BV2 cells

could be affected by Piezo1 channel activity using an *in vitro* transwell assay. 10% FBS

or 0.5 μM Aβ$_{1-42}$, the main component of plaques in Alzheimer's Disease, were used as

chemo-attractants for BV2 cells seeded in the upper chamber of a membrane insert.

Yoda1 and/or GsMTx-4 were added to the BV2 cells to modify the functioning of

Piezo1. We found that the addition of Yoda1 (23.33 ± 1.202 cells towards 10% FBS,

25.67 ± 0.2404 cells towards 0.5 μM Aβ$_{1-42}$) significantly suppressed the migratory

tendency of BV2 cells compared to an untreated control (168.0 ± 8.544 cells towards

10% FBS, 51.73 ± 0.7688 cells towards 0.5 μM Aβ$_{1-42}$), whereas adding GsMTx-4

(240.3 ± 13.35 cells towards 10% FBS, 67.93 ± 4.476 cells towards 0.5 μM Aβ$_{1-42}$)

enhanced the cells' movement (Fig. 3A-B). The observed migration patterns were

similar with both attractants used. Thus, we found that the activation or blocking of the

Piezo1 channel could indeed affect the ability of BV2 cells to migrate.

We next constructed a BV2-Piezo1 Knock-out (KO) stable cell line using

CRISPR-Cas9 technology. We used a patch clamp setup to confirm that BV2-Piezo1

KO cells were insensitive to Yoda1 stimulation compared to WT cells (Fig. 3C). Inward

current induced by 30 μM Yoda1 in BV2-WT cells was 1176 ± 394.8 pA while it was

much lower (18.67 ± 6.333 pA) in BV2-Piezo1 KO cells (Fig. 3D). The resting

membrane potential of BV2 cells with Piezo1 KO was not obviously different from

wild type cells (Fig. S4) indicating that Piezo1 KO did not itself alter the cells' general

electrical activity and health. Calcium imaging also showed that BV2-Piezo1 KO cells

were insensitive to Yoda1 stimulation (2μM and 10μM) (Fig. 3E-F). Piezo1 depletion

was also found to significantly enhance the cells' migration towards both FBS and $A\beta_{1-42}$

compared to WT (Fig. 3G-H). The cell migration process requires close control and

coordination of actin dynamics and actin signaling dysfunction has been known to lead

to aberrant cell migration (Schaks, Giannone et al. 2019). Hence, we also examined the

effect of Piezo1 KO on the polymerization of actin. Yoda1 was seen to significantly

enhance actin polymerization in WT cells, but this effect was abolished in KO cells (Fig.

S5A-B). Interestingly, Piezo1 KO was found to reduce basal actin polymerization levels.

Thus, knocking out Piezo1 in BV2 cells was found to significantly affect the cells'

electrical responses, calcium dynamics and tendency to migrate.

The role of Piezo1 in mediating the migration ability of microglia was further

examined by a simple chemoattraction assay *in vivo*. The oligomer $A\beta_{1-42}$ HiLyte™

Fluor 555 was injected into the dentate gyrus (DG) of the hippocampus as an attractant.

Counting IBA1-positive cells inside the $A\beta_{1-42}$ -555 signal regions, many more

microglia were found to have migrated to the DG in Piezo1$^{\Delta TMEM119}$ mice than control

Piezo1$^{fl/+}$ mice (Fig. 3I-J) by. We also counted microglia in mice of both genotypes

without any injection of $A\beta_{1-42}$, and found that conditional Piezo1 KO did not affect the

microglia numbers in their basal state (Fig. 3K-L). Thus, we found that Piezo1

expression and activation regulated of the ability of microglia to migrate towards

attractant stimuli *in vitro* and *in vivo*.

228
229 **Fig3. Piezo1 regulates migration ability of microglia both *in vivo* and *in vitro*.**
230 **A.** Representative images of migrated BV2 cells stained by crystal violet treated by
231 Yoda1 (10μM) or GsMTx4 (1μM), attracted by 10% FBS or 0.5 μM Aβ$_{1-42}$ respectively.
232 Scale bar = 100um.
233 **B.** Quantification of migrated cell numbers treated by Yoda1(10μM) or GsMTx4 (1μM)

attracted by 10% FBS or 0.5 μM Aβ$_{1-42}$ respectively. Bars represent mean ± SD. Repeated for 3 times. Two-tailed unpaired t test. *p < 0.05, ****p < 0.0001.

C. Representative traces of 30μM Yoda1 induced inward currents in BV2-WT and BV2- Piezo1 KO cells recorded at −60 mV in the indicated conditions.

D. Quantification of 30 μM Yoda1-induced inward currents in BV2 WT or BV2-Piezo1 KO cells. Bars represent mean ± SD. N=3 cells for each group. Two-tailed unpaired t test. *p < 0.05.

E. Representative time course of calcium fluorescence in BV2-WT and BV2 - Piezo1 KO cells treated with 2μM Yoda1.

F. Statistical analysis of ΔF/F0 (%) in BV2 WT and BV2- Piezo1 KO cells under different Yoda1 concentrations. Bars represent mean ± SEM. N=3 experiments; 30–40 cells were analyzed for each experiment. Two-tailed Mann–Whitney U test. ****p < 0.0001.

G. Representative images of migrated BV2 WT or BV2- Piezo1 KO cells stained by crystal violet, attracted by FBS or Aβ$_{1-42}$ respectively. Scale bar=100um.

H. Quantification of migrated BV2 WT or BV2- Piezo1 KO cell numbers attracted by FBS or Aβ$_{1-42}$ respectively. Bars represent mean ± SD. Repeated for 3 times. Two-tailed unpaired t test. **p < 0.01.

I. Representative images of Aβ$_{1-42}$ injected regions in the dentate gyrus (DG) region of Piezo1$^{fl/+}$ and Piezo1^{ΔTmem119} mouse groups respectively. Scale bar = 50μm.

J. Quantified average values for total Aβ-555/IBA1 positive (IBA1+) cells, normalized to control group (n=6 in Piezo1$^{fl/+}$ group, n=5 in Piezo1^{ΔTmem119} group). Bars represent mean ± SD. Two-tailed unpaired t test. **p < 0.01.

K. Representative images of IBA1 positive cells in DG area of Piezo1$^{fl/+}$ and Piezo1^{ΔTmem119} brain slice respectively (blue: DAPI, green: IBA1). Scale bar= 100μm.

L. Quantification of microglia numbers per mm2 around DG area in Piezo1fl/+ and Piezo1ΔTmem119 mice respectively (n=5 in Piezo1fl/+ group, n=6 in Piezo1ΔTmem119 group). Bars represent mean ± SEM. Two-tailed unpaired t test. n.s., no significance.

Piezo1 regulates the immune response of microglia

Since microglia act as key players in the primary immune response of the CNS,

we also wanted to investigate whether Piezo1 was involved in the microglial immune

response. For this, we used BV2-Piezo1 KO and WT cell lines, stimulated with Yoda1

and/or LPS for 24 hours, a toxin used to trigger an inflammatory response. The

expression levels of some important pro-inflammatory cytokines were then monitored

by RT-PCR. In BV2-WT cells stimulated with LPS, we found that Yoda1 increased IL-

1β, IL-6 and TNF-α mRNA expression levels (Fig. 4A-C). We also found that BV2-

Piezo1 KO cells expressed significantly lower levels of these cytokines even when

stimulated with LPS (Fig. 4D-F). Western blotting for IL-1β further backed up the

finding that loss of Piezo1 expression significantly reduced the ability of microglia to

response to LPS stimulation by producing cytokines (Fig. 4G-H). We further tested the

immune response mediation by Piezo1 *in vivo* using our transgenic mouse model. The

cortex tissues of mice were evaluated for cytokine expression levels after

intraperitoneal LPS injection. Mice with the conditional Piezo1 KO in microglia

showed lower expression of these cytokines when challenged by LPS, but not in the

basal state (Fig. 4I-K). Piezo1 was hence seen to play crucial roles in the LPS-induced

neuroinflammatory response of microglia.

Fig 4. Piezo1 regulates production of pro-inflammatory cytokines by microglia when challenged by LPS both *in vivo* and *in vitro*.

A-C. Relative mRNA expressions of IL-1β, IL-6 and TNF-α in BV2-WT cells, exposed to Yoda1 (10μM) or not, in basal state or stimulated by LPS (100ng/mL), normalized to BV2 basal control state. Bars represent mean ± SD. N=3 independent experiments. *p < 0.05, **p < 0.01.

D-F. Relative mRNA expressions of IL-1β, IL-6 and TNF-α of BV2-WT and BV2-Piezo1 KO cells in basal state or stimulated by LPS (100ng/mL), normalized to BV2-WT basal state. Bars represent mean ± SD. N=3 independent experiments. *p < 0.05, **p < 0.01, ****p < 0.0001.

G. Representative Western blot images for IL-1 β and GAPDH in indicated conditions.

H. Quantification of IL-1 β protein levels normalized to GAPDH compared with BV2-WT treated with LPS group. Bars represent mean ± SD. N=3 independent experiments. Two-tailed unpaired T test. *p < 0.05, **p < 0.01.

I-K. Relative mRNA expressions of IL-1β, IL-6 and TNF-α in cortex of Piezo1$^{fl/+}$ and Piezo1$^{\Delta TMEM119}$ mice in basal state or after administration of LPS (n=3 in Piezo1$^{fl/+}$ -LPS group, n=4 in Piezo1$^{\Delta Tmem119}$ -LPS group, n=3 in Piezo1$^{fl/+}$ +LPS group, n=3 in Piezo1$^{\Delta Tmem119}$ +LPS group). Bars represent mean ± SEM. For panel a-f and i-k, One-way ANOVA with post-hoc Tukey test. *p < 0.05, ***p < 0.001.

Stiffness-mediated microglial migration and activation are mediated by Piezo1

Given that Piezo1 is involved in mechanosensation in many cell types and physiological functions (Li, Hou et al. 2014, Pathak, Nourse et al. 2014, Cahalan, Lukacs et al. 2015, Gudipaty, Lindblom et al. 2017, Romac, Shahid et al. 2018, Solis, Bielecki et al. 2019), we reasoned that Piezo1 would play such a role in microglia as well. To investigate the role of Piezo1 in microglia sensing of stiffness, to regulate migration ability and cytokine production of microglia, BV2- WT and BV2- Piezo1 KO cells were plated onto soft (0.6 kPa) or stiff (35 kPa) polyacrylamide gel substrates (as previously described (Chang, Chen et al. 2017)) for 48 hours. The cells' migration towards 10% FBS and expression of inflammatory cytokines were then compared between groups. We found that enhanced substrate stiffness reduced the migration of BV2- WT cells but not in BV2- Piezo1 KO cells (Fig. 5A-B). We also found that greater substrate stiffness could increase the expression of pro-inflammatory factors (IL-1β, IL-6 and TNF-α) in BV2-WT cells, but did not in IL-1β and TNF-α expression in BV2-Piezo1 KO cells (Fig. 5C-E), indicating Piezo1 is involved in stiffness-induced

microglial migration and activation process.

An unexpected aspect of the soft/stiff substrate culture was that an increase in

stiffness was also found to be correlated with increased Piezo1 expression. BV2 cells

on a stiffer substrate showed significantly increased Piezo1 expression after 48 hours

of culture in both protein levels (Fig. 5F-G) and mRNA levels (Fig. 5H). Stiff-cultured

cells also showed much greater calcium response when stimulated with 10 µM Yoda1

than did soft-cultured cells (Fig. 5I-K), which also points towards an increased

expression of Piezo1. Thus, we found that Piezo1 was both a mediator for the microglial

response to environmental stiffness cues, while also being upregulated in microglia in

response to varying mechanical conditions.

Fig 5. Stiffness-dependent Piezo1 expression/activity modulates microglia pro-inflammatory cytokines production.

A. Representative images of migrated BV2 WT or Piezo1-KO cells pre-cultured for 48h in PA gel of varying stiffness (Soft: 0.6KPa, Stiff: 35KPa) stained by crystal violet. Scale bar = 100um.

B. Quantification of migrated cell numbers of BV2 WT or Piezo1-KO cells pre-cultured for 48h in PA gel of different stiffness (Soft: 0.6KPa, Stiff: 35KPa). Bars represent mean ± SD. N=3 independent experiments. Two-tailed unpaired t test. n.s., no significance, *p < 0.05.

C-E. Relative IL-1β, IL-6 and TNF-α mRNA expression of BV2- WT and BV2- Piezo1 KO seeded on surfaces of indicated stiffness (Soft: 0.6KPa, Stiff: 35KPa). Bars represent mean ± SEM. N=3 independent experiments. Two-tailed unpaired t test. n.s., no significance; **p < 0.01.

F. Immunofluorescence confocal images of Piezo1 (green) in BV2 cell lines seeding on different stiffness of PA gel (Soft: 0.6KPa, Stiff: 35KPa). Scale bar= 20μm.

G. Quantification of Piezo1 mean fluorescent intensity (MFI) of BV2 cell lines seeding on different stiffness of PA gel. Bars represent mean ± SD. Repeated for 3 times, around 30 cells were analyzed for each experiment. Two-tailed unpaired t test. ****p < 0.0001.

H. Relative Piezo1 mRNA levels of BV2 cell lines seeding on different stiffness of PA gel. Bars represent mean ± SEM. N=4 independent experiments. Two-tailed unpaired t test. *p < 0.05.

I. Representative live-cell imaging of BV2 cells loaded with Fura2 calcium indicator treated by Yoda1 (10μM) at time 0 (left panel) and at the time of maximum fluorescence (right panel) seeding on soft and stiff substrate respectively.

J. Representative time course of calcium fluorescence in BV2 cells seeding on soft (black curve) and stiff substrates (red curve) treated with Yoda1 (10μM).

K. Statistical analysis of ΔF/F0 (%) of BV2 cells seeding on soft (black) and stiff substrates (red) induced by Yoda1 (10μM). Bars represent mean ± SD. N=3 independent experiments, 20-30 cells were analyzed for each experiment. Two-tailed Mann–Whitney U test. ****p < 0.0001.

Discussion

In the CNS, microglia constantly encounter mechanically heterogeneous regions of varying cellular density and matrix composition. This implies the presences of a suite of cellular machinery to sense and transduce these cues into cellular signaling. In the present study, we identify a role for the mechanically-activated ion channel Piezo1 in regulating microglial functions, i.e., migration, pro-inflammatory cytokines production and stiffness sensing. We found that Piezo1 is highly expressed in primary cultured microglia. Microglia from mice with conditional Piezo1 knockout exhibited enhanced migration ability towards injected Aβ$_{1-42}$ and reduced pro-inflammatory cytokines production after stimulation by LPS. These phenomena were also observed *in vitro*

using constructed BV2-WT and BV2-Piezo1 KO stable cell lines. I*n vitro*, we found

that activation and blockade of the Piezo1 channel reversely regulated the migration

tendencies of microglia. Furthermore, Yoda1 could enhance pro-inflammatory

cytokines production of microglia stimulated by LPS, but similarly activating Piezo1

in unstimulated microglia had little effect. We also show that substrate stiffness

modulates Piezo1 expressions and stiffness regulated microglia migration and pro-

inflammatory cytokines production (IL-1β, TNF-α) enhancement was affected by

Piezo1 depletion. Together, our study reveals critical roles of Piezo1 in modulating

microglial functions and indicates its significance in maintaining brain health.

Ca^{2+}-mediated signals are a common signal transduction pathway in all living

cells, including microglia (Moller 2002). Ca^{2+} activities of microglia are known to be

altered in some pathological conditions, such as interrupted Ca^{2+} signal transduction

following brain damage (Eichhoff, Brawek et al. 2011) or in Alzheimer's disease (AD)

patients (McLarnon, Choi et al. 2005), though the mechanism is unclear. We found

stiffness-driven variation in Piezo1 expression using our *in vitro* PA gel model. A

previous study using a TgF344-AD transgenic model also found that Piezo1 expression

level in astrocyte increases around stiff amyloid plaques (Velasco-Estevez, Mampay et

al. 2018). Given that mechanical cues are altered in the brain during progression of

several pathological conditions in which microglia are involved, e.g., traumatic brain

injury (Keating and Cullen 2021) or AD (Murphy, Huston et al. 2011, Murphy, Jones et

al. 2016), aberrant Piezo1expression may be a contributor to microglial calcium

signaling dysfunction in such conditions as well. Thus, the role of Piezo1 in microglia,

regulated by changing microenvironment stiffness of the brain, merits further study

using appropriate animal models.

Resting microglia are highly dynamic and can be rapidly activated to participate

in pathological responses, including migrating to affected sites, releasing various

inflammatory molecules, and clearing cellular debris (Kreutzberg 1996, Koizumi,

Shigemoto-Mogami et al. 2007) to maintain brain health. Therefore, migration is a

critical first step for microglia to function after injury or inflammatory stimuli. Our

findings identify a potentially crucial role of Piezo1 in modulating microglial migration.

Numerous studies have shown that microglia tend to migrate towards chemo-attractant

stimuli (Fan, Xie et al. 2017). Based on our observation that Piezo1 serves as a

mechanosensor and plays a part in regulating microglial migration, mechanical cues

exerted on the cells during ageing (Segel, Neumann et al. 2019) and some pathological

conditions (Murphy, Huston et al. 2011, Riek, Millward et al. 2012, Schregel, Wuerfel

et al. 2012, Streitberger, Sack et al. 2012) thus also merit exploration.

Piezo1 also played an important role in microglial production of pro-

inflammatory cytokines (IL-1β, IL-6, TNF-α), which are involved in the pathogenesis

of many neurodegenerative diseases (Smith, Das et al. 2012). Enhanced stiffness,

mediated by Piezo1, was also associated with increased pro-inflammatory cytokine

production in microglia. These cytokines play potentially contradictory roles in the

maintenance of brain homeostasis. On one hand, release of these factors is crucial to

prevent further damage to CNS tissue when, say, under attack by pathogens. However,

on the other hand, the same cytokines may also be toxic to neurons and other glial cells,

and contribute to the progression of neurodegenerative disorders (Smith, Das et al. 2012,

Ogunmokun, Dewanjee et al. 2021, Tansey, Wallings et al. 2022). Interestingly,

substrate stiffness could also regulate Piezo1 expression in microglia and induce

following microglial function changes (Fig5). This phenomenon suggests Piezo1 could

be a key molecule mediating between microenvironment stiffness and microglial

function. Given the mechanical cues generated during the progression of brain diseases,

Piezo1 could be an important subject of study as a participant in signaling pathways

and as a druggable target. Several studies have already described the roles of

mechanosensitive ion channels in regulating brain diseases, i.e., TRPC1 (Wu, Ryskamp

et al. 2018), TRPC3, TRPC6 (Wang, Lu et al. 2015, Chen, Lu et al. 2017), TRPV4

(Wang, Zhou et al. 2019). Given that Piezo1 appears to mediate crucial microglial

functions, further study of its role in the brain could improve our understanding of the

progression of brain diseases and the function of microglia in health and disease.

Materials and methods

Transgenic mouse line generation and genotyping

All animal procedures were approved by the Animal Subjects Ethics Sub-Committee (ASESC) of the Hong Kong Polytechnic University, and were performed in compliance with the guidelines of the Department of Health - Animals (Control of Experiments) of the Hong Kong S.A.R. government. Generation of Piezo1$^{\Delta TMEM119}$ mice was accomplished through breeding Piezo1$^{flox/flox}$ (Jackson Laboratories Stock No. 029213) and Tmem119-2A-Cre^{ERT2} (Jackson Laboratories Stock No: 031820) mice together to generate progeny that were heterozygous for both genes. The generated heterozygotes were then bred with Piezo1$^{flox/flox}$ mice to generate Piezo1$^{flox/flox}$Tmem119$^{Cre/+}$ (Piezo1$^{\Delta Tmem119}$) and Piezo1$^{flox/+}$Tmem119$^{Cre/+}$ (Piezo1$^{fl/+}$) mice. Mice were housed in a 12-h light/12-h dark cycle, with temperatures between 20 °C and 24 °C and 40–70% humidity with food and water available *ad libitum*. Tamoxifen (Sigma-Aldrich, T5648) was dissolved in corn oil containing 5% ethanol at 20 mg/ml under vortex at times for several hours in the dark. At time of administration, the mice weighed 19–26 g and intraperitoneal (i.p.) injection with 100μl of the 20 mg/ml tamoxifen solution for 5 consecutive days. After administration of tamoxifen for at least 2 weeks, mice were used for different *in vivo* assays. Mice tail samples were genotyped using the following primers suggested by Jackson Laboratories: Piezo1-28247- forward: GCC TAG ATT CAC CTG GCT TC; Piezo1-28248- reverse: GCT CTT AAC CAT TGA GCC ATC T; TMEM119-16504: ATC GCA TTC CTT GCA AAA

GT; TMEM119-42648: CAG TAT GTG GGG TCA CTG AAG A; TMEM119-42649:

ACT TGG GGA GAT GTT TCC TG using Phire Tissue Direct PCR Master Mix (F170S,

Thermo Scientific) with the following cycling conditions: Initial denaturation 98°C for

5 min, followed by 10 cycles of 98°C for 5 s, 65°C for 5 s, -0.5 C per cycle decrease,

68°C for 20 s, followed by 30 cycles of 98°C for 5 s, 60°C for 5 s, 72°C for 20 s,

followed by a final hold of 72°C for 1 min. Reactions were separated on 2% agarose

gels yielding the following band sizes: Cre +/-: 280bp and 378bp, Cre -/-: 378bp, Piezo1

fl/fl: 380bp, Piezo1 fl/+: 380bp and 188bp. Successful knock-out Piezo1 were validated

by examining DNA from mice brains using the following primers: Piezo1- 28247-

forward: GCC TAG ATT CAC CTG GCT TC; Piezo1 KO- reverse: AGG TTG CAG

GGT GGC ATG GCT CTT TTT. An obvious Piezo1 KO band was found in 230bp but

none in control mice group. Brain samples were also collected for testing Piezo1 mRNA

levels by qPCR using the following primers: Piezo1-192-forward:

CTCTGGCCTAACTACTGTCT; Piezo1-192-reverse:

GATAAGGTTGGTGGAGTTGG.

Cell culture

The BV2 cell line (RRID: CVCL_0182) was a gift from Dr. Bo Peng's group.

Cells were cultured in DMEM (10566016, Gibco), 10% fetal bovine serum (A3840402,

Gibco), 1% Penicillin-Streptomycin (15140122, Gibco) at 37°C inside a standard cell

culture incubator containing 5% CO_2.

Primary microglia isolation

For primary microglia isolation, P3 pups of C57BL/6J mice were sacrificed, followed by dissociation and isolation using the CD11b isolation kit according to the manufacturer's protocol (130-092-628 and 130-126-725, Miltenyi Biotec). Primary microglia purity was examined by flow cytometry after staining by Cd11b-FITC (130-113-243, Miltenyi Biotec) and Cd45-APC (130-110-660, Miltenyi Biotec). Dishes were pre-coated with 50μg/μl Poly-L-Lysine and cells were then plated on the dishes at a density of 10^5 cells/Ml and cultured in DMEM/F12 (10565018, Gibco), 10% fetal bovine serum (A3840402, Gibco), 1% Penicillin-Streptomycin (15140122, Gibco) at 37°C inside a standard cell culture incubator containing 5% CO_2.

CRISPR knock-out (KO) Piezo1 cell cloning

To generate Piezo1 knock-out (KO) cell lines, single guide RNAs (sgRNAs) ACGCTTCAATGCTCTCTCGC targeting the second exon of mouse piezo1 as reference (Del Marmol, Touhara et al. 2018) was inserted into MLM3636 vector (Addgene Plasmid #43860). SgRNA inserted MLM3636 and Cas9 expression plasmid pX459 v2.0 (Addgene Plasmid #62988) were transfected into BV2 cell line with Lipofectamine™ 3000 transfection reagent (L3000015, Thermo Fisher Scientific). Transfected BV2 cells were selected with 1ug/mL puromycin (A1113803, Life Technologies). After selection, live cells were trypsinized, diluted as 5 cell/mL and then seeded on 96-well plate at an average density of 0.5cell/well for growing to obtain single clones. The successful knock-out clone was verified using Sanger sequencing,

493 patch clamp and calcium imaging.

Intracellular calcium imaging

Cells were plated on confocal dishes (200350, SPL Life Sciences). Prior to

recording, cells were incubated with 2 μM Fura-2 (F1221, Invitrogen) at 37°C for 30

min dispersed in Margo solution with 0.2‰ w/v Pluronic F-127. Cells were then

washed twice and incubated for another 15 min. GsMTx-4 (ab141871, abcam) were

added in this step as needed. Experiments were performed at room temperature using

Nikon Eclipse Ti2-E microscope. Yoda1 (SML1558, Sigma Aldrich) or DMSO

(D12345, Invitrogen) was added during the imaging process. Intracellular calcium

concentration was indicated as the ratio of fura-2 emission (510 nm) intensities for 340

and 380 nm excitation.

Patch clamp

Cells were seeded in culture dishes before patch clamp recording. Borosilicate

glass-made patch pipettes (Vitrex, Modulohm A/S, Herlev, Denmark), were pulled with

micropipette puller (P-97, Sutter Instrument Co., USA) to a resistance of 2 to 5 MΩ

before being filled with KCl pipette solution (in mM): KCl 138, NaCl 10, $MgCl_2$ 1 and

HEPES 10 with D-manitol compensated for osm290. Inward current was recorded with

a data acquisition system (DigiData 1322A, Axon Instruments) and an amplifier

(Axopatch-200B, Axon Instruments, Foster City, CA, USA). The command voltages

were controlled by a computer equipped with pClamp Version 9 software. For cell

resting membrane potential (RMP) test, cells were placed on bath solution (in mM):

NaCl 130, MgCl$_2$ 2, KCl 4.5, Glucose 10, HEPES 20, and CaCl$_2$ 2, pH 7.4. When the

whole-cell Giga seal was formed and the capacitance of cell was measured, the RMP

of cells was recorded when the injection current was set at zero. Afterwards, the inward

currents induced by Yoda1 was recorded by voltage clamp gap free recording mode.

RNA isolation and qPCR

RNA was extracted using the MiniBEST Universal RNA Extraction Kit (Cat No.

9767, TaKaRa). cDNA syntheses were generated using PrimeScript RT Master Mix

(RR036A, TaKaRa). SYBR® Premix Ex Taq™ II (Tli RNaseH Plus) (RR820A,

TaKaRa) were performed with the CFX96 Touch Deep Well Real-Time PCR system

under standard conditions. The $\Delta\Delta$Ct method was used to calculate gene expression.

Primer sequences are provided in the following table.

mouse TRPC1 Forward	AAAGTGGTGGCTCACAACAA
mouse TRPC1 Reverse	CGTTTGCAAAAGCAAAAAGC
mouse TRPC3 Forward	ATTCCTCGCCATCGGCTATTGG
mouse TRPC3 Reverse	GAGGCGTTGAATACAAGCAGACC
mouse TRPC6 Forward	TGGCAGTGAAGTTCCTCGTGGT
mouse TRPC6 Reverse	GGAAAATGGTGAAGGAGGCTGC
mouse TRPV4 Forward	TCACCGCCTACTATCAGCCACT
mouse TRPV4 Reverse	GAACAGGACTCCTGTGAAGAGC
mouse PIEZO1 Forward	GCAGTGGCAGTGAGGAGATT
mouse PIEZO1 Reverse	GATATGCAGGCGCCTATCCA
mouse PIEZO2 Forward	GCACTCTACCTCAGGAAGACTG
mouse PIEZO2 Reverse	CAAAGCTGTGCCACCAGGTTCT
mouse KCNK2 Forward	CACGGTGTTCAAGGCATTGGAG
mouse KCNK2 Reverse	TCCCTGCGTTTATTGCTGCCAC
mouse Asic1a Forward	CGTTTGTGTCTTGCCAGGAGCA
mouse Asic1a Reverse	CGTGATGCTGTAGGAGTCGAAG
mouse Asic3 Forward	TATGTGGCTCGGAAGTGCGGAT
mouse Asic3 Reverse	CAGACACAAGTGTCCTTTCGCAG
mouse Stoml3 Forward	CTTCCTGCTGATGCTCGTTACC

mouse Stoml3 Reverse	CTTGGATGCGTCCCAGTCTGAA
mouse IL-1β Forward	TGCAGCTGGAGAGTGTGGATCCC
mouse IL-1β Reverse	TGTGCTCTGCTTGTGAGGTGCTG
mouse IL-6 Forward	TACCACTTCACAAGTCGGAGGC
mouse IL-6 Reverse	CTGCAAGTGCATCATCGTTGTTC
mouse TNF-α Forward	ACTTCGGGGTGATCGGTCCCC
mouse TNF-α Reverse	TGGTTTGCTACGACGTGGGCTAC
mouse β-actin Forward	GAGATTACTGCCCTGGCTCCTA
mouse β-actin Reverse	TCATCGTACTCCTGCTTGCTGAT

RNA sequencing and data analysis

RNA sequencing and informatics analysis were performed by Novogene Company (Beijing, China). RNA integrity was assessed with the Agilent 2100 Bioanalyzer (Agilent Technologies). Libraries were barcoded, quantified using the NEBNext Ultra RNA Library Prep Kit for Illumina (Cat No. 7530, New England Biolabs (NEB)). Quantified libraries were pooled and sequenced on Illumina platforms, according to effective library concentration and data amount. The clustering of the index-coded samples was performed according to the manufacturer's instructions. After cluster generation, the library preparations were sequenced on an Illumina platform and paired-end reads were generated. Raw data (raw reads) of fastq format were first processed through in-house perl scripts. Clean data (clean reads) were obtained by removing reads containing adapter, reads containing ploy-N and low quality reads from raw data. At the same time, Q20, Q30 and GC content the clean data were calculated. All the downstream analyses were based on the clean data with high quality. Reference genome and gene model annotation files were downloaded from genome website directly. Index of the reference genome was built using Hisat2 v2.0.5 and paired-end clean reads were aligned to the reference genome using Hisat2 v2.0.5. Feature Counts v1.5.0-p3 was

used to count the reads numbers mapped to each gene. Differential expression analysis

of two groups (3 biological replicates per condition) was performed using the DESeq2

R package. Genes with an adjusted P value <=0.05 found by DESeq2 were assigned as

differentially expressed. Gene Ontology (GO) enrichment analysis of differentially

expressed genes was implemented by the cluster Profiler R package, in which gene

length bias was corrected. GO terms with corrected P value less than 0.05 were

considered significantly enriched by differential expressed genes.

Cell migration transwell assays

BV2 cells and primary microglia migration towards 10% FBS were tested in

transwell assays (COSTAR 24-well plate with inserts, 8-μm pore, #3422, Corning, NY).

For the transwell assay, 600 μl DMEM together with the attractant- 10% FBS or 0.5μM

$A\beta_{1-42}$ were added below the membrane inserts. Drugs-1‰ DMSO, 10 μM Yoda1, 1

μM GsMTx4 - were respectively added to 5×10^4 cells in 100 μl media right before

adding the cells to the top wells. Migration was assayed for 20 h at 37°C and 5% CO_2.

Subsequentlly, inserts were removed and washed three times with PBS. BV2 cells were

fixed and stained with 0.5% crystal violet for 20min. Stationary cells were removed

from the top with cotton tips. Five random fields were selected for imaging, and the

migrated cells were quantified using Image J software.

Stereotactic injection of $A\beta_{1-42}$

Beta-Amyloid (1-42), HiLyte™ Fluor 555-labeled (Cat. AS-60480-01, ANASPEC)

were dissolved in DMSO to obtain a 1 mM stock, resuspended in PBS at 100 μM, and

incubated at 37 °C for 1 day to promote $A\beta_{1\text{-}42}$ aggregation. Mice were anesthetized by i.p. injection of a mixture of ketamine (100 mg/kg), xylazine (10 mg/kg), 2 μl of $A\beta_{1\text{-}42}$-555 was distributed into DG area of the hippocampus (AP -2 mm; ML -1.5 mm; DV 2 mm) using stereotaxic equipment (RWD Life Science Co., Ltd). After recovery from surgery, animals were returned to their home cages. Post surgery (16 h), mice were anesthetized and perfused for subsequent experiments.

For histological analyses of microglia migration towards $A\beta_{1\text{-}42}$-555, brains were perfused with ice-cold PBS, fixed in 4% PFA and immersed in 4% PFA for another 4 hours. Brains were then sectioned in 50μM thickness by Vibrating Microtome (Leica VT1200S). Sections were washed, then permeabilized with 0.3% Triton X-100 (Sigma) in PBS with 5% Goat serum (16210072, Gibco) and blocked for 1 hour. The microglia-specific antibody IBA1 (Rabbit mAb 17198S, CST, RRID: AB_2820254) was incubated in blocking buffer overnight at 4°C followed by intensive washing and incubation with a Goat anti-rabbit Adsorbed Secondary Antibody, Alexa Fluor 488 (A-11008, Thermo Fisher Scientific, RRID: AB_143165) for 2 hours. Sections were washed again and mounted with Mounting Medium With DAPI-Aqueous, Fluoroshield (ab104139, Abcam). Data acquisition was performed using a Nikon Eclipse Ti2-E Live-cell Fluorescence Imaging System in Z-stack model.

Western blotting

Cells were rinsed with PBS before being exposed to lysis buffer, a combination of RIPA lysis buffer (89900, Thermo Fisher Scientific) and 1% Halt™ Protease and

Phosphatase Inhibitor Cocktail (78446, Thermo Fisher Scientific). Cells were scraped

to obtain the adhered cells and the lysate was collected. The lysate was put on ice for

another 30 min to achieve thorough lysis. Lysate was spun at 13000 rpm for 20min at

4°C and the supernatant was collected. Samples were quantified using Pierce™ Rapid

Gold BCA Protein Assay Kit (A53225, Thermo Fisher Scientific) and aliquots of 50μg

of protein were prepared.The proteins were denatured through the use of a loading

buffer supplemented with 5% 2-mercaptoethanol at 95°C for 10 min before each sample

was loaded. Gel electrophoresis resulted in the separation of proteins before being

transferred onto nitrocellulose membranes using the Thermo Scientific™ Pierce™ G2

Fast Blotter. Following electroblotting, the membranes were blocked using 5% nonfat

milk in TBST for 1 hour at room temperature. Membranes were then gently washed

with TBST and incubated with Anti-IL-1 beta primary antibody (ab254360, Abcam) at

4°C overnight with continuous rotation. Additional washes, 10 min. 3 times in TBST

followed. Before the membranes were probed with Goat anti-Rabbit IgG (H+L)

Secondary Antibody, HRP (31460, Thermo Fisher Scientific, RRID: AB_228341)

diluted at 1:5000 at room temperature for 1 hour. The membrane was then washed in

TBST and immersed into a chemiluminescent HRP substrate solution (Bio-Rad) and

imaged using a ChemiDoc MP System (Bio-Rad).

Immunofluorescent staining of cells

Cells were washed with PBS 3 times, then fixed with 4% PFA for 15 min. Cells were

then washed with PBS and permeabilized with 0.1% Triton X-100 in PBS for 20 min.

After that, cells were washed with PBS and blocked with 5% normal goat serum

(16210072, Gibco) for 1 hour at room temperature. Cells were then incubated with Piezo1

primary antibody (15939-1-AP, Proteintech, RRID:AB_2231460) diluted 1:200, at 4°C

overnight and washed with PBS 3 times. After that, cells were incubated with Goat anti-

rabbit Adsorbed Secondary Antibody, Alexa Fluor 488 (A-11008, Thermo Fisher

Scientific) at room temperature for 1 hour.

Actin polymerization staining

Green fluorescent phalloidin conjugate (ab112125, Abcam) was used to stain F-actin

according to the manufacturer's recommendation. Cells were then washed with PBS 3

times and sealed with mounting medium containing DAPI. Data acquisition was

performed using a Leica TCS SPE Confocal Microscope.

Quantification and statistical analysis

Statistical tests were performed using GraphPad Prism 8, which was also used to

prepare the graphs. Bars charts represent the mean of the indicated number of

experiments or subjects, unless otherwise indicated. Details of the statistical analyses

performed for each figure are provided in the figure legends. A p value of < 0.05 or

below was considered statistically significant for all experiments.

Resource availability

Lead contact

Further information and request for reagents and resources should be addressed to,

and will be fulfilled by, the Lead Contact, Lei Sun (lei.sun@polyu.edu.hk).

Materials availability

This study did not generate new unique reagents.

Data and code availability

All data reported in this paper will be shared by the lead contact upon request,

unless it is protected by law.

This paper does not report original code.

Any additional information required to reanalyze the data reported in this paper is

available from the lead contact upon request.

Acknowledgements

This work was supported by the Hong Kong Research Grants Council General

Research Fund (15104520, 15102417 and 15326416), Hong Kong Innovation

Technology Fund (MRP/018/18X and MHP/014/19), Key-Area Research and

Development Program of Guangdong Province (2018B030331001), internal funding

from the Hong Kong Polytechnic University (1-ZE1K and 1-ZVW8), and Research

Institute of Smart Ageing (1-CD76). The authors would like to thank the facility and

technical support from University Research Facility in Life Sciences (ULS) and

University Research Facility in Behavioral and Systems Neuroscience (UBSN) of The

Hong Kong Polytechnic University. We thank to Dr. Michael Yuen from ULS for

instructing us in primary microglia isolation. We thank to Mr. Kai Tang for instructing

us in PA gel fabrication.

Author contributions

T.Z. and L.S. designed research. T.Z., S.K., Y.W., H.C., JJ.Z., C.P.C., XY.Z., T.L.

performed research, T.Z., S.K., JH.G., K.F.W., XH.H., MY.Y. analyzed data. T.Z., S.K.,

and L.S. wrote the paper.

Competing interests

The authors declare that they have no competing interest.

References

Aloisi, F. (2001). "Immune function of microglia." Glia **36**(2): 165-179.

Ayata, P. and A. Schaefer (2020). "Innate sensing of mechanical properties of brain tissue by microglia." Curr Opin Immunol **62**: 123-130.

Bae, C., F. Sachs and P. A. Gottlieb (2011). "The mechanosensitive ion channel Piezo1 is inhibited by the peptide GsMTx4." Biochemistry **50**(29): 6295-6300.

Bollmann, L., D. E. Koser, R. Shahapure, H. O. Gautier, G. A. Holzapfel, G. Scarcelli, M. C. Gather, E. Ulbricht and K. Franze (2015). "Microglia mechanics: immune activation alters traction forces and durotaxis." Front Cell Neurosci **9**: 363.

Cahalan, S. M., V. Lukacs, S. S. Ranade, S. Chien, M. Bandell and A. Patapoutian (2015). "Piezo1 links mechanical forces to red blood cell volume." Elife **4**.

Chang, T. Y., C. Chen, M. Lee, Y. C. Chang, C. H. Lu, S. T. Lu, D. Y. Wang, A. Wang, C. L. Guo and P. L. Cheng (2017). "Paxillin facilitates timely neurite initiation on soft-substrate environments by interacting with the endocytic machinery." Elife **6**.

Chen, X., S. Wanggou, A. Bodalia, M. Zhu, W. Dong, J. J. Fan, W. C. Yin, H. K. Min, M. Hu, D. Draghici, W. Dou, F. Li, F. J. Coutinho, H. Whetstone, M. M. Kushida, P. B. Dirks, Y. Song, C. C. Hui, Y. Sun, L. Y. Wang, X. Li and X. Huang (2018). "A Feedforward Mechanism Mediated by Mechanosensitive Ion Channel PIEZO1 and Tissue Mechanics Promotes Glioma Aggression." Neuron **100**(4): 799-815 e797.

Chen, X. Y., M. Lu, X. J. He, L. Ma, L. Birnbaumer and Y. H. Liao (2017). "TRPC3/6/7 Knockdown Protects the Brain from Cerebral Ischemia Injury via Astrocyte Apoptosis Inhibition and Effects on NF-DB Translocation." Molecular Neurobiology **54**(10): 7555-7566.

Cojocaru, A., E. Burada, A. T. Balseanu, A. F. Deftu, B. Catalin, A. Popa-Wagner and E. Osiac (2021). "Roles of Microglial Ion Channel in Neurodegenerative Diseases." J Clin Med **10**(6).

Coste, B., J. Mathur, M. Schmidt, T. J. Earley, S. Ranade, M. J. Petrus, A. E. Dubin and A. Patapoutian (2010). "Piezo1 and Piezo2 are essential components of distinct mechanically activated cation channels." Science **330**(6000): 55-60.

Del Marmol, J. I., K. K. Touhara, G. Croft and R. MacKinnon (2018). "Piezo1 forms a slowly-inactivating mechanosensory channel in mouse embryonic stem cells." Elife **7**.

Echeverry, S., M. J. Rodriguez and Y. P. Torres (2016). "Transient Receptor Potential Channels in Microglia: Roles in Physiology and Disease." Neurotox Res **30**(3): 467-478.

Eichhoff, G., B. Brawek and O. Garaschuk (2011). "Microglial calcium signal acts as a rapid sensor of single neuron damage in vivo." Biochimica Et Biophysica Acta-Molecular Cell Research **1813**(5): 1014-1024.

Elkin, B. S., E. U. Azeloglu, K. D. Costa and B. Morrison, 3rd (2007). "Mechanical heterogeneity of the rat hippocampus measured by atomic force microscope indentation." J Neurotrauma **24**(5): 812-822.

Fan, Y., L. Xie and C. Y. Chung (2017). "Signaling Pathways Controlling Microglia Chemotaxis." Mol Cells **40**(3): 163-168.

Gudipaty, S. A., J. Lindblom, P. D. Loftus, M. J. Redd, K. Edes, C. F. Davey, V. Krishnegowda and J. Rosenblatt (2017). "Mechanical stretch triggers rapid epithelial cell division through Piezo1." Nature **543**(7643): 118-121.

Keating, C. E. and D. K. Cullen (2021). "Mechanosensation in traumatic brain injury." Neurobiol Dis **148**: 105210.

Kigerl, K. A., J. P. de Rivero Vaccari, W. D. Dietrich, P. G. Popovich and R. W. Keane (2014). "Pattern

recognition receptors and central nervous system repair." Exp Neurol **258**: 5-16.

Koizumi, S., Y. Shigemoto-Mogami, K. Nasu-Tada, Y. Shinozaki, K. Ohsawa, M. Tsuda, B. V. Joshi, K.

A. Jacobson, S. Kohsaka and K. Inoue (2007). "UDP acting at P2Y6 receptors is a mediator of microglial

phagocytosis." Nature **446**(7139): 1091-1095.

Konno, M., H. Shirakawa, S. Iida, S. Sakimoto, I. Matsutani, T. Miyake, K. Kageyama, T. Nakagawa, K.

Shibasaki and S. Kaneko (2012). "Stimulation of transient receptor potential vanilloid 4 channel

suppresses abnormal activation of microglia induced by lipopolysaccharide." Glia **60**(5): 761-770.

Kreutzberg, G. W. (1996). "Microglia: a sensor for pathological events in the CNS." Trends Neurosci

19(8): 312-318.

Li, J., B. Hou, S. Tumova, K. Muraki, A. Bruns, M. J. Ludlow, A. Sedo, A. J. Hyman, L. McKeown, R.

S. Young, N. Y. Yuldasheva, Y. Majeed, L. A. Wilson, B. Rode, M. A. Bailey, H. R. Kim, Z. Fu, D. A.

Carter, J. Bilton, H. Imrie, P. Ajuh, T. N. Dear, R. M. Cubbon, M. T. Kearney, R. K. Prasad, P. C. Evans,

J. F. Ainscough and D. J. Beech (2014). "Piezo1 integration of vascular architecture with physiological

force." Nature **515**(7526): 279-282.

Lim, C. G., J. Jang and C. Kim (2018). "Cellular machinery for sensing mechanical force." BMB Rep

51(12): 623-629.

Matcovitch-Natan, O., D. R. Winter, A. Giladi, S. Vargas Aguilar, A. Spinrad, S. Sarrazin, H. Ben-Yehuda,

E. David, F. Zelada Gonzalez, P. Perrin, H. Keren-Shaul, M. Gury, D. Lara-Astaiso, C. A. Thaiss, M.

Cohen, K. Bahar Halpern, K. Baruch, A. Deczkowska, E. Lorenzo-Vivas, S. Itzkovitz, E. Elinav, M. H.

Sieweke, M. Schwartz and I. Amit (2016). "Microglia development follows a stepwise program to

regulate brain homeostasis." Science **353**(6301): aad8670.

McLarnon, J. G., H. B. Choi, L. F. Lue, D. G. Walker and S. U. Kim (2005). "Perturbations in calcium-

mediated signal transduction in microglia from Alzheimer's disease patients." J Neurosci Res **81**(3): 426-

435.

Mobasheri, A., S. D. Carter, P. Martin-Vasallo and M. Shakibaei (2002). "Integrins and stretch activated

ion channels; putative components of functional cell surface mechanoreceptors in articular

chondrocytes." Cell Biol Int **26**(1): 1-18.

Moller, T. (2002). "Calcium signaling in microglial cells." Glia **40**(2): 184-194.

Moshayedi, P., G. Ng, J. C. Kwok, G. S. Yeo, C. E. Bryant, J. W. Fawcett, K. Franze and J. Guck (2014).

"The relationship between glial cell mechanosensitivity and foreign body reactions in the central nervous

system." Biomaterials **35**(13): 3919-3925.

Murphy, M. C., J. Huston, 3rd, C. R. Jack, Jr., K. J. Glaser, A. Manduca, J. P. Felmlee and R. L. Ehman

(2011). "Decreased brain stiffness in Alzheimer's disease determined by magnetic resonance

elastography." J Magn Reson Imaging **34**(3): 494-498.

Murphy, M. C., D. T. Jones, C. R. Jack, Jr., K. J. Glaser, M. L. Senjem, A. Manduca, J. P. Felmlee, R. E.

Carter, R. L. Ehman and J. Huston, 3rd (2016). "Regional brain stiffness changes across the Alzheimer's

disease spectrum." Neuroimage Clin **10**: 283-290.

Nakajima, K., Y. Tohyama, S. Maeda, S. Kohsaka and T. Kurihara (2007). "Neuronal regulation by which

microglia enhance the production of neurotrophic factors for GABAergic, catecholaminergic, and

cholinergic neurons." Neurochem Int **50**(6): 807-820.

Nonomura, K., V. Lukacs, D. T. Sweet, L. M. Goddard, A. Kanie, T. Whitwam, S. S. Ranade, T. Fujimori,

M. L. Kahn and A. Patapoutian (2018). "Mechanically activated ion channel PIEZO1 is required for

lymphatic valve formation." Proceedings of the National Academy of Sciences of the United States of

America **115**(50): 12817-12822.

Ogunmokun, G., S. Dewanjee, P. Chakraborty, C. Valupadas, A. Chaudhary, V. Kolli, U. Anand, J. Vallamkondu, P. Goel, H. P. R. Paluru, K. D. Gill, P. H. Reddy, V. De Feo and R. Kandimalla (2021). "The Potential Role of Cytokines and Growth Factors in the Pathogenesis of Alzheimer's Disease." Cells **10**(10).

Ohana, L., E. W. Newell, E. F. Stanley and L. C. Schlichter (2009). "The Ca2+ release-activated Ca2+ current (I(CRAC)) mediates store-operated Ca2+ entry in rat microglia." Channels (Austin) **3**(2): 129-139.

Ohashi, K., S. Fujiwara and K. Mizuno (2017). "Roles of the cytoskeleton, cell adhesion and rho signalling in mechanosensing and mechanotransduction." J Biochem **161**(3): 245-254.

Pathak, M. M., J. L. Nourse, T. Tran, J. Hwe, J. Arulmoli, D. T. Le, E. Bernardis, L. A. Flanagan and F. Tombola (2014). "Stretch-activated ion channel Piezo1 directs lineage choice in human neural stem cells." Proc Natl Acad Sci U S A **111**(45): 16148-16153.

Riek, K., J. M. Millward, I. Hamann, S. Mueller, C. F. Pfueller, F. Paul, J. Braun, C. Infante-Duarte and I. Sack (2012). "Magnetic resonance elastography reveals altered brain viscoelasticity in experimental autoimmune encephalomyelitis." Neuroimage Clin **1**(1): 81-90.

Romac, J. M. J., R. A. Shahid, S. M. Swain, S. R. Vigna and R. A. Liddle (2018). "Piezo1 is a mechanically activated ion channel and mediates pressure induced pancreatitis." Nature Communications **9**.

Schaks, M., G. Giannone and K. Rottner (2019). "Actin dynamics in cell migration." Essays Biochem **63**(5): 483-495.

Schregel, K., E. Wuerfel, P. Garteiser, I. Gemeinhardt, T. Prozorovski, O. Aktas, H. Merz, D. Petersen, J. Wuerfel and R. Sinkus (2012). "Demyelination reduces brain parenchymal stiffness quantified in vivo by magnetic resonance elastography." Proc Natl Acad Sci U S A **109**(17): 6650-6655.

Segel, M., B. Neumann, M. F. E. Hill, I. P. Weber, C. Viscomi, C. Zhao, A. Young, C. C. Agley, A. J. Thompson, G. A. Gonzalez, A. Sharma, S. Holmqvist, D. H. Rowitch, K. Franze, R. J. M. Franklin and K. J. Chalut (2019). "Niche stiffness underlies the ageing of central nervous system progenitor cells." Nature **573**(7772): 130-134.

Shi, M., F. Du, Y. Liu, L. Li, J. Cai, G. F. Zhang, X. F. Xu, T. Lin, H. R. Cheng, X. D. Liu, L. Z. Xiong and G. Zhao (2013). "Glial cell-expressed mechanosensitive channel TRPV4 mediates infrasound-induced neuronal impairment." Acta Neuropathologica **126**(5): 725-739.

Smith, J. A., A. Das, S. K. Ray and N. L. Banik (2012). "Role of pro-inflammatory cytokines released from microglia in neurodegenerative diseases." Brain Research Bulletin **87**(1): 10-20.

Smith, J. A., A. Das, S. K. Ray and N. L. Banik (2012). "Role of pro-inflammatory cytokines released from microglia in neurodegenerative diseases." Brain Res Bull **87**(1): 10-20.

Solis, A. G., P. Bielecki, H. R. Steach, L. Sharma, C. C. D. Harman, S. Yun, M. R. de Zoete, J. N. Warnock, S. D. F. To, A. G. York, M. Mack, M. A. Schwartz, C. S. Dela Cruz, N. W. Palm, R. Jackson and R. A. Flavell (2019). "Mechanosensation of cyclical force by PIEZO1 is essential for innate immunity." Nature **573**(7772): 69-+.

Streitberger, K. J., I. Sack, D. Krefting, C. Pfuller, J. Braun, F. Paul and J. Wuerfel (2012). "Brain viscoelasticity alteration in chronic-progressive multiple sclerosis." PLoS One **7**(1): e29888.

Syeda, R., M. N. Florendo, C. D. Cox, J. M. Kefauver, J. S. Santos, B. Martinac and A. Patapoutian (2016). "Piezo1 Channels Are Inherently Mechanosensitive." Cell Reports **17**(7): 1739-1746.

Syeda, R., J. Xu, A. E. Dubin, B. Coste, J. Mathur, T. Huynh, J. Matzen, J. M. Lao, D. C. Tully, I. H. Engels, H. M. Petrassi, A. M. Schumacher, M. Montal, M. Bandell and A. Patapoutian (2015). "Chemical

activation of the mechanotransduction channel Piezo1." <u>Elife</u> **4**.

Tansey, M. G., R. L. Wallings, M. C. Houser, M. K. Herrick, C. E. Keating and V. Joers (2022). "Inflammation and immune dysfunction in Parkinson disease." <u>Nat Rev Immunol</u>.

Velasco-Estevez, M., M. Mampay, H. Boutin, A. Chaney, P. Warn, A. Sharp, E. Burgess, E. Moeendarbary, K. K. Dev and G. K. Sheridan (2018). "Infection Augments Expression of Mechanosensing Piezo1 Channels in Amyloid Plaque-Reactive Astrocytes." <u>Front Aging Neurosci</u> **10**: 332.

Wang, J., R. Lu, J. Yang, H. Li, Z. He, N. Jing, X. Wang and Y. Wang (2015). "TRPC6 specifically interacts with APP to inhibit its cleavage by gamma-secretase and reduce Abeta production." <u>Nat Commun</u> **6**: 8876.

Wang, J. H. and B. P. Thampatty (2006). "An introductory review of cell mechanobiology." <u>Biomech Model Mechanobiol</u> **5**(1): 1-16.

Wang, Z. Q., L. Zhou, D. An, W. X. Xu, C. F. Wu, S. Sha, Y. C. Li, Y. C. Zhu, A. D. Chen, Y. M. Du, L. Chen and L. Chen (2019). "TRPV4-induced inflammatory response is involved in neuronal death in pilocarpine model of temporal lobe epilepsy in mice." <u>Cell Death & Disease</u> **10**.

Wu, J., D. Ryskamp, L. Birnbaumer and I. Bezprozvanny (2018). "Inhibition of TRPC1-Dependent Store-Operated Calcium Entry Improves Synaptic Stability and Motor Performance in a Mouse Model of Huntington's Disease." <u>J Huntingtons Dis</u> **7**(1): 35-50.

Zeng, W. Z., K. L. Marshall, S. Min, I. Daou, M. W. Chapleau, F. M. Abboud, S. D. Liberles and A. Patapoutian (2018). "PIEZOs mediate neuronal sensing of blood pressure and the baroreceptor reflex." <u>Science</u> **362**(6413): 464-467.

Zhang, Y., K. Chen, S. A. Sloan, M. L. Bennett, A. R. Scholze, S. O'Keeffe, H. P. Phatnani, P. Guarnieri, C. Caneda, N. Ruderisch, S. Deng, S. A. Liddelow, C. Zhang, R. Daneman, T. Maniatis, B. A. Barres and J. Q. Wu (2014). "An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex." <u>J Neurosci</u> **34**(36): 11929-11947.

Supplementary Figures

Fig S1. Purity of isolated primary microglia.

The purity of isolated primary microglia were determined by flow cytometry with Cd11b-FITC and Cd45-APC staining analyzed by BD Accuri C6 software.

Fig S2. Piezo1 is expressed in the microglial cell line BV2.

A. Representative live-cell imaging of BV2 loaded with Fura-2 calcium indicator at rest state (left panel) and at the time of maximum fluorescence (right panel). Scale bar= 20μm.

B. Representative time course of calcium fluorescence in BV2 cells treated with different concentrations of Yoda1 and if pretreated with GsMTx4.

C. Statistical analysis of ΔF/F0 (%) in BV2 cells treated with different concentrations of Yoda1 and if pretreated with GsMTx4. Bars represent mean ± SEM, N=3 independent experiments; 30–40 cells were analyzed for each experiment. Two-tailed Mann–Whitney U test. ****$p < 0.0001$.

830
Fig S3. Validation of Tmem119- Piezo1 cKO mice.
A. Genotyping results of the transgenic mice. 1: Mouse1, Piezo1 fl/+, Cre-/-. 2: Mouse2, Piezo1 fl/+, Cre+/-. 3: Mouse3, Piezo1 fl/+, Cre-/-. 4: Mouse4, Piezo1 fl/fl, Cre-/-. 5: Mouse5, Piezo1 fl/fl, Cre+/-. M represents for DNA ladder.
B. Piezo1 knock-out validation band. Arrow indicates the Piezo1 knock-out band. KO represents for Piezo1$^{\Delta Tmem119}$ mouse and NC represents for Piezo1$^{fl/+}$ mouse. H_2O is adopted as negative control. M represents the DNA ladder.
C. Piezo1 mRNA expressions (n= 3 in Piezo1$^{fl/+}$ group, n= 4 in Piezo1$^{\Delta Tmem119}$ group) of brain tissues isolated from Control and Tmem119-Piezo1 cKO mice). Transcript levels were calculated using the 2−ΔΔCT method using β-actin as a reference gene and were normalized to the average expression from control samples. Bars represent mean ± SD. Two-tailed unpaired t-test. *$p < 0.05$.

Fig S4. Resting membrane potential of BV2 WT or BV2- Piezo1 KO cells.
N=6 cells for BV2 WT group and N=4 cells for BV2- Piezo1 KO group. Bars represent mean ± SEM. Two-tailed unpaired t-test. n.s., no significance.

Fig S5. Piezo1 mediates actin polymerization of microglia.
A. Representative confocal images of F-actin staining (green) and DAPI (blue) under
Yoda1 (10μM) stimulation in BV2-WT and BV2-Piezo1 KO cells. Scale bar= 20μm.
B. Quantification of F-actin mean fluorescent intensity (MFI) of BV2-WT and BV2-
Piezo1 KO cells stimulated by Yoda1 (10μ M). Repeated for 3 times. 20-35 cells were
analyzed for each group. Bars represent mean ± SD. Two-tailed unpaired t-test. n.s., no
significance; *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.