GCN2 eIF2 kinase promotes prostate cancer by maintaining amino acid homeostasis

Ricardo A. Cordova[1,2], Jagannath Misra[1], Parth H. Amin[1], Angela J. Klunk[1], Nur P. Damayanti[2,3], Kenneth R. Carlson[1], Andrew J. Elmendorf[4], Hyeong-Geug Kim[1], Emily T. Mirek[5], Marcus J. Miller[6], X. Charlie Dong[1], Liang Cheng[2,7,8], Tracy G. Anthony[5], Roberto Pili[9,#], Ronald C. Wek[1,2,#], and Kirk A. Staschke[1,2,#]

[1]Department of Biochemistry and Molecular Biology, Indiana University School of Medicine, Indianapolis, IN 46202, USA.

[2]Indiana University Melvin and Bren Simon Comprehensive Cancer Center, Indianapolis, IN 46202, USA.

[3]Department of Neurological Surgery, Indiana University School of Medicine, Indianapolis, IN 46202, USA.

[4]Indiana Biosciences Research Institute (IBRI), Indianapolis, IN 46202, USA.

[5] Department of Nutritional Sciences, Rutgers University, New Brunswick, NJ 08901, USA.

[6]Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN 46202, USA

[7]Department of Pathology and Laboratory Medicine, Indiana University School of Medicine, Indianapolis, IN 46202, USA

[8]Department of Urology, Indiana University School of Medicine, Indianapolis, IN 46202, USA

[9]Jacobs School of Medicine and Biomedical Sciences, University at Buffalo, Buffalo, NY 14203, USA

#To whom correspondence should be addressed:

Kirk A. Staschke, Department of Biochemistry and Molecular Biology, 635 Barnhill Drive, Indiana University School of Medicine, Indianapolis, Indiana 46202-5122; E-mail:

kastasch@iu.edu

Ronald C. Wek, Department of Biochemistry and Molecular Biology, 635 Barnhill Drive, Indiana University School of Medicine, Indianapolis, Indiana 46202-5122; E-mail: rwek@iu.edu.

Roberto Pili, Department of Medicine, 875 Ellicott Street, University at Buffalo, Buffalo, NY 14203; E-mail: rpili@buffalo.edu.

Running title: GCN2 controls tumor amino acid homeostasis

Keywords: ISR; prostate cancer, GCN2; eIF2; amino acid transport; SLC; solute carrier; CD98; 4F2; 4F2hc; SLC3A2

Conflict of interest statement

R.C.W. is a member of the advisory board and holds equity in HiberCell, Inc. K.A.S. and T.G.A are consultants for HiberCell, Inc. and K.A.S. receives research support from HiberCell, Inc.

Abstract word count: 158
Word Count: 5490
Figures and tables: 7
Supplementary Information:
- Supplementary Figures S1 – S21
- Supplementary Tables S1 – S4
- Supplementary Files S1 – S4
- Supplementary methods

Abstract

A stress adaptation pathway termed the integrated stress response has been suggested to be active in many cancers including prostate cancer (PCa). Here, we demonstrate that the eIF2 kinase GCN2 is required for sustained growth in androgen-sensitive and castration-resistant models of PCa both in vitro and in vivo, and is active in PCa patient samples. Using RNA-seq transcriptome analysis and a CRISPR-based phenotypic screen, GCN2 was shown to regulate expression of over 60 solute-carrier (*SLC*) genes, including those involved in amino acid transport and loss of GCN2 function reduces amino acid import and levels. Addition of essential amino acids or expression of SLC3A2 (4F2) partially restored growth following loss of GCN2, suggesting that GCN2 targeting of SLC transporters is required for amino acid homeostasis needed to sustain tumor growth. A small molecule inhibitor of GCN2 showed robust in vivo efficacy in androgen-sensitive and castration-resistant mouse models of PCa, supporting its therapeutic potential for the treatment of PCa.

Significance: The ISR kinase GCN2 is critical for maintaining tumor amino acid levels to facilitate growth, suggesting a novel therapeutic strategy for the treatment of prostate cancer by inducing starvation for essential amino acids.

Introduction

Prostate cancer (PCa) accounts for 20% of all newly diagnosed cancers and is the second leading cause of cancer-related deaths in men in the United States [1]. Many factors contribute to prostate cancer, including genetics, diet, age, and race [2]. The pathological grade of the PCa and whether it has disseminated into surrounding tissues is prognostic for risk and treatment efficacy [3]. Androgen receptor (AR) is considered the primary oncogenic driver for PCa [4, 5], and therapies targeting AR activity are current standard of care, especially for those

with metastatic PCa [6-9]. Unfortunately, patients develop resistance with time and fail therapy

[8, 10]. In addition, it remains unclear how additional pathways drive PCa through AR-

dependent and AR-independent mechanisms. Therefore, the identification of novel targets for

therapeutic intervention are critically needed and clinically relevant.

The Integrated Stress Response (ISR) is critical for cell adaptation and survival to

environmental stresses. Central to the ISR is phosphorylation of the α subunit of eIF2 (p-eIF2α),

which triggers reductions in global protein synthesis to conserve energy and nutrients, and

concomitantly facilitates adaptive gene expression [11, 12]. Four eIF2α kinases respond to

distinct stress conditions: The eIF2α kinase GCN2 (EIF2AK4) is induced by amino acid

starvation and other conditions that trigger ribosomal collisions or stalling [13-15]. PERK

(EIF2AK3) is responsive to disruptions in the endoplasmic reticulum [16, 17]. PKR (EIF2A2) is

activated by viral infection and inflammation [18, 19], and HRI (EIF2AK1) is enhanced by

restricted heme or iron supply, oxidative stress, and disruptions in mitochondrial function [20-

22]. Activation of these ISR regulators may also occur in response to oncogenic drivers, such as

MYC amplification and loss of the PTEN tumor suppressor, frequently observed in prostate

tumors. Therefore, both extrinsic and intrinsic cell stresses can invoke the ISR in tumors and

pharmacological inhibition of translational control is reported to trigger cytotoxicity in different

cancer models [23-25].

To reprogram gene expression towards recovery from stress damage, the ISR also

induces preferential translation of specific ISR gene transcripts, such as activating transcription

factor 4 (ATF4), a transcriptional regulator of genes involved in amino acid metabolism and

proteostasis control, oxidative stress defenses, along with feedback control of the ISR [26-28].

Therefore, p-eIF2α induces both translational and transcriptional modes of gene expression to

achieve stress adaptation. Targeted inhibition of ATF4 impedes progression of MYC-driven

cancers, emphasizing the important role of this ISR effector in tumor progression [23-25].

Here, we dissect the role of eIF2α kinase(s) induced in PCa and the mechanisms by

which the ISR serves to promote growth of this cancer. Using androgen-sensitive and

castration-resistant human PCa cell lines, primary human PCa tissues, and mouse xenograft

models, we show that the ISR regulator GCN2 is activated in PCa. GCN2 induces expression of

key ISR effector genes involved in nutrient transport that are critical for appropriate maintenance

of free amino acid pools. Genetic or pharmacological inhibition of GCN2 dysregulates

expression of these transporters and inhibits PCa growth. These results point to GCN2 as a

potential therapeutic target for PCa.

Results

GCN2 activity supports prostate cancer growth.

To determine the role of individual eIF2α kinases and their importance for proliferation of

PCa cells in culture, androgen-sensitive LNCaP cells cultured in androgen replete media

conditions were individually depleted of HRI, PKR, PERK, or GCN2 using target-specific

siRNAs. There was robust reduction of protein levels of each of the eIF2α kinases; however,

only cells depleted for GCN2 or PKR showed a significant decrease in proliferation (**Figures**

1A,1B, Supplementary Figure 1A, and Supplementary File 1). Whereas GCN2 depletion

lowered p-eIF2α levels and ATF4 expression, knockdown of PKR had no effect on p-eIF2α and

ATF4 protein levels (**Figure 1B and Supplementary Figure 1A**). These results suggest that

LNCaP cell proliferation requires GCN2 as an ISR regulator.

To broaden our investigation to other PCa cell lines, we knocked down GCN2 in the

androgen-sensitive line LAPC-4 and in the castration-resistant cell lines C4-2B, MR49F, 22Rv1,

and PC-3. MR49F cells express AR, but are resistant to enzalutamide [29], 22Rv1 cells express

the ARv7 splice variant and are also resistant to enzalutamide [30-32], and PC-3 is AR null [33].

In each of these PCa cell lines, depletion of GCN2 reduced proliferation and lowered ATF4

expression (**Figure 1C, Supplementary Figure 1B, and Supplementary File 1**). In 22Rv1 and

PC-3 cells, we were also able to achieve a deletion of *GCN2* using CRISPR/Cas9

(**Supplementary Figure 2A**). In the case of 22Rv1, we determined a similar slow growth

phenotype as observed with siRNA knockdown of GCN2 (**Supplementary Figure 2B and**

Supplementary File 1). Reintroduction of GCN2 into the GCN2 KO cells rescued the slow

growth phenotype, and rescued expression of ATF4 and its target gene ASNS (**Supplementary**

Figure 2B, 2C, and Supplementary File 1). The small molecule GCN2iB is a selective inhibitor

of GCN2 eIF2α kinase activity [34]. Using the inhibitory concentration range of 500 nM to 10 µM

of GCN2iB, we determined that pharmacological inhibition of GCN2 also caused significant

growth inhibition of LNCaP cells, and similarly reduced the growth of C4-2B, 22Rv1, and PC-3

cells **(Figure 1D, Supplementary Figure 3A, and Supplementary File 1).** As expected,

treatment of the LNCaP, 22Rv1, C4-2B, or PC-3 cells with GCN2iB lowered activation of GCN2

as measured by autophosphorylation of the activation loop (p-GCN2-T899), reduced p-eIF2α

levels, and lowered expression of ATF4, and its target genes ASNS and TRIB3. Importantly,

treatment with GCN2iB did not alter protein levels of AR (**Figure 1E and Supplementary**

Figure 3B). To further test the importance of the ISR in PCa, we selectively depleted ATF4

using siRNA in LNCaP, MR49F, C4-2B, and PC-3 cells and observed similar growth inhibition

for each cell line (**Supplementary Figure 2B, Supplementary Figure 4, and Supplementary**

File 1). These results support the hypothesis that the eIF2α kinase activity of GCN2 and the ISR

support PCa proliferation.

We next sought to determine whether GCN2 is activated in human malignant PCa

tissues. Using immunohistochemistry and prostate tumor microarrays (TMAs) featuring normal

and non-malignant prostate tissue (N = 33) and malignant tumor tissues (N = 88), we

determined that there was enhanced GCN2 activity in the tumor samples compared to normal

tissues as measured by p-GCN2-T899 staining (**Figure 1F**). In addition, increased p-GCN2

(Thr-899) staining was also observed in prostate tumors from patient core needle biopsies as

compared to adjacent non-malignant tissue (**Supplementary Figure 5**). Collectively, these

results indicate that GCN2 is active basally in both androgen-sensitive and castration-resistant

PCa models without appreciable extrinsic stress treatments, confers growth advantages, and is

activated in prostate cancer patient tissues.

GCN2 induces ISR transcriptome featuring expression of amino acid transporters.

To address the mechanisms by which GCN2 confers growth advantages in PCa, we

cultured LNCaP cells in the presence or absence of GCN2iB and then measured changes in the

transcriptome by RNA-seq. With 24 hours of GCN2iB treatment, 239 transcripts showed a

significant ≥ 2-fold increase, whereas 374 transcripts were decreased ≥ 2-fold (fdr ≤ 0.05)

compared to vehicle-treated cells (**Figure 2A, Supplementary Table 1**). We carried out a gene

set enrichment analysis (GSEA) of the differentially-expressed transcripts to distinguish the

molecular functions that were altered by GCN2 inhibition. Included among the functional gene

groups showing significant dependence on GCN2 for expression were those linked to amino

acid deprivation and transport (**Figures 2B and 2C)**. These genes are representative of ISR-

targeted gene groups and include genes regulated by ATF4 such as *DDIT3 (CHOP)*, *ASNS*,

and *TRIB3*, those involved in glycine-serine biosynthesis (*PHGDH, PSAT1, PSPH*, and

SHMT2), aminoacyl-tRNA synthetases, and a large number of SLC transporters, many of which

function in the transport of amino acids (**Figure 2B and Supplementary File 2**). As illustrated in

the heat map, expression of the *SLC* genes were significantly lowered after 6 hours of GCN2iB

treatment, with further reductions following 24 hours (**Figure 2C**). It is noteworthy that of the 61

known SLC genes involved in the transport of amino acids, 27 were measurably expressed in

LNCaP cells, and 20 of these expressed transcripts were significantly decreased following

treatment with GCN2iB (**Figure 2B and Supplementary File 2**).

Lowered expression of SLC proteins upon GCN2 inhibition was further validated by

immunoblot analyses. SLC7A11 (xCT), SLC7A1 (CAT-1), SLC1A4 (ASCT1), SLC1A5 (ASCT2),

and SLC3A2 (4F2) protein levels were sharply lowered with GCN2iB treatment of LNCaP cells,

along with reduced expression of ATF4 and its target gene ASNS (**Figure 2D and**

Supplementary Figure 6A). Similarly, treatment of C4-2B, 22Rv1, and PC-3 cells with GCN2iB

also reduced SLC7A5 (LAT1), SLC7A11 (xCT), and SLC3A2 (4F2) (**Supplementary Figure**

3B). We also measured transporter proteins in 22Rv1 and PC-3 cells and showed that deletion

of GCN2 also diminished protein levels of SLC7A11 (xCT), SLC7A5 (LAT1), and SLC3A2 (4F2)

(**Figure 2E, Supplementary Figure 6B**). Furthermore, the reintroduction of GCN2 fully restored

expression of ATF4 and ASNS and partially restored expression of all three SLC proteins in

22Rv1 cells (**Figure 2E**). To more directly address the effects of GCN2 inhibition on amino acid

transport, we measured uptake of leucine, lysine, and arginine in LNCaP and 22Rv1 cells.

Treatment of LNCaP cells with GCN2iB or deletion of GCN2 in 22Rv1 cells significantly lowered

the uptake of each of these amino acids (**Figure 2F and 2G**), suggesting that GCN2-directed

gene expression is important for uptake of amino acids in PCa cells.

GCN2 is critical for maintenance of free amino acid pools.

Given that GCN2 is important for expression of *SLC* genes and amino acid uptake, we

next addressed whether the intracellular levels of free amino acids are disrupted upon inhibition

of GCN2. LNCaP cells were cultured in the presence or absence of GCN2iB for 8 hours and

free amino acid levels were measured. With pharmacological inhibition of GCN2, there was a

sharp reduction in the amounts of essential amino acids (EAA), including histidine, the

branched-chain amino acids, threonine, phenylalanine, and methionine, and nonessential amino

acids (NEAA), such as glutamate and glutamine, aspartate, asparagine, alanine, and tyrosine

(**Figure 3A**). We next considered whether supplementing the culture medium with additional

amino acids would alleviate the growth defect that occurs with the inhibition of GCN2. While

GCN2iB treatment of LNCaP cells cultured in standard medium markedly reduced growth,

supplementation of additional EAAs into the medium reversed this proliferation defect and

increased free amino acid pools in treated cells (**Figure 3B, Supplementary Figure 7,**

Supplementary Figure 8, and Supplementary File 1). By contrast, supplementation with

additional NEAA did not suppress the LNCaP growth defect triggered by GCN2iB treatment

(**Supplementary Figure 8 and Supplementary File 1**).

To address the effect of GCN2 inhibition and EAA supplementation on the cell cycle in

cultured PCa cells, LNCaP cells were treated with GCN2iB or GCN2 expression was reduced

with siRNA. Both methods of GCN2 depletion caused an increase in the G1 phase of the cell

cycle concomitant with a reduction in S phase, indicative of checkpoint control of the cell cycle

that can occur with amino acid depletion [35, 36] (**Figure 3C and Supplemental Figure 9A**).

Supplementation of LNCaP cells with EAA reversed the accumulation of cells in G1 phase

elicited by GCN2iB (**Figure 3C**). These results suggest that inhibition of GCN2 can cause cell

cycle arrest, which was recapitulated with siRNA knockdown of GCN2 or ATF4

(**Supplementary Figure 10A**). Similar findings were observed in 22Rv1 cells deleted for GCN2

or in other PCa cell lines when GCN2 or ATF4 levels were reduced with siRNA

(**Supplementary Figure 9A and 9B**).

Accumulating uncharged tRNA occurs upon severe depletion of cognate amino acids

and is a primary signal activating GCN2 [37]. We cultured LNCaP cells in the presence or

absence of GCN2iB and measured tRNA charging genome-wide using CHARGE-seq [38].

There was a reduction of aminoacylated tRNA^{His} upon GCN2iB treatment that was restored

when additional EAA were supplemented to the growth medium (**Figures 3D, 3E, and**

Supplementary File 3). These results were further supported by measurements of tRNA^{His}

charging levels by qRT-PCR. Addition of EAA to cultured LNCaP cells significantly increased

tRNA^{His} charging levels while treatment with GCN2iB dramatically reduced tRNA^{His} charging

levels which was reversed by the addition of EAA (**Supplementary Figure 10A**). These results

suggest that histidine depletion and the accompanying accumulation of uncharged tRNA^{His} is a

key signal for activation of GCN2. Upon treatment with GCN2iB, which lowered p-eIF2α and

blocked appropriate induction of the ISR, there was severe depletion of histidine accompanied

by accumulation of uncharged tRNA^{His}, culminating in a slow-growth phenotype.

To further address the role of histidine availability in the slow growth phenotype

associated with GCN2 inhibition, we cultured GCN2iB-treated LNCaP cells in medium with

added supplements of individual amino acids. In alignment with our charged tRNA^{His}

measurements, only the addition of histidine to the culture medium restored the growth inhibition

triggered by GCN2iB (**Figure 3F and Supplementary Figure 8**). Removal of histidine from the

growth medium increased G1 accumulation similar to GCN2iB treatment, while supplementation

of additional histidine to the medium alleviated the cell cycle arrest triggered by GCN2iB

treatment (**Figure 3G**). Finally, supplementation with the full complement of EAA or histidine

alone significantly lowered p-GCN2, p-eIF2α, and ATF4 expression, although it is noted that

histidine supplementation rendered only partial suppression of the ISR compared to addition of

EAA (**Figure 3H**). Accompanying this reduction in activation of GCN2, supplementation with

histidine sharply increased aminoacylation of tRNA^{His} (**Figure 3E**) and there was a trend

towards enhanced global protein synthesis as measured by puromycin incorporation. It is

curious that there was a trend towards decreased bulk translation upon GCN2iB treatment that

was restored with the addition of histidine (**Supplementary Figure 10B**). As noted above,

GCN2 inhibition for 48 hours adversely affected the cell cycle and amino acid levels, which

together are suggested to reduce bulk translation upon GCN2iB treatment despite lowered p-

eIF2α levels.

We next determined whether EAA supplementation in other PCa cell lines also

suppressed the slow growth phenotype triggered by loss of GCN2 activity. Addition of EAA or

histidine alone also suppressed the slow growth phenotype of MR49F cells treated with GCN2iB

(**Supplementary Figure 11A**). However, in 22Rv1 cells only lysine alleviated the slow growth

associated with deletion of GCN2 (**Supplementary Figure 11B**). These growth

supplementation studies suggest that while GCN2 is indispensable for maintaining EAA in PCa

cells, the precise processes by which GCN2 carries out these functions can vary among PCa

cell lines presumably due to differences in their unique genetic features and cell-line specific

buffering mechanisms accompanying specific gene expression and metabolic programs.

Depletion of GCN2 in LNCaP and MR49F cells renders them vulnerable to histidine depletion,

whereas deletion of GCN2 sensitizes 22Rv1 cells to lysine depletion.

Select GCN2-regulated SLC transporters are critical for PCa fitness.

Given that GCN2 has a primary role for inducing expression of *SLC* genes, we sought to

determine the contributions of these transporters for PCa growth and viability. To evaluate their

contributions in an unbiased manner, we utilized a previously reported CRISPR/Cas9 library

targeting 394 human SLC genes and pseudogenes [39]. Each of the SLC genes was targeted

with 6 specific single guide RNAs (sgRNAs), along with negative controls for target specificity

and positive controls for assessing cell fitness. The sgRNA library was stably expressed in

LNCaP and 22Rv1 cells using lentivirus delivery. Cells were then subjected to puromycin

selection and then cultured for 3 - 4 weeks (15 - 20 population doublings) as previously reported

[39]. Genomic DNA was prepared and encoded sgRNA sequences were amplified by PCR,

sequenced, and their composition was compared to the starting cells with the sgRNA library at

Day 0. Depletion of the cells expressing specific sgRNAs were then evaluated for statistical

significance, and cell fitness was scored and ranked. A collection of SLC genes showed

significantly diminished growth, including several that require GCN2 for expression, such as

SLC7A5 (LAT1), SLC25A36, SLC25A46, SLC27A1, SLC35A2, and SLC3A2 (4F2) (**Figures 4A**

and Supplementary File 4). *SLC27A1* and *SLC3A2 (4F2)* were required for efficient growth of

both LNCaP and 22Rv1 cells (**Figure 4B and Supplementary Figure 12**). These results

suggest that PCa cells require a specific collection of SLC genes for growth and a subset of

these are dependent on GCN2 for full expression.

Given the importance of SLC3A2 (4F2) in the transport of bulk amino acids [40], we next

focused on the GCN2-dependent regulation of this transport regulator. Treatment of LNCaP

cells with the GCN2iB for 6 or 24 hours resulted in a reduction in GCN2 activation, as shown by

p-GCN2, and lowered ATF4 mRNA and protein. Levels of SLC3A2 (4F2) protein and mRNA

were also lowered with inhibition of GCN2, although this reduction occurred after 24 hours of

treatment (**Figure 4C and Supplementary Figure 13A**). Consistent with our findings that

histidine limitation in part drives GCN2 activation in LNCaP cells, addition of histidine to the

growth media of LNCaP cells in culture reduced GCN2 activation and lowered expression of

ATF4 and SLC3A2 (4F2) protein and mRNA, while removing histidine from the media further

increased GCN2 activation and expression of ATF4 and SLC3A2 (4F2) (**Figure 4D and**

Supplementary Figure 13B). Salubrinal has been reported to increase p-eIF2α by inhibiting its

PP1-directed dephosphorylation [41]. Supporting the idea that p-eIF2α is critical for 4F2

expression, addition of salubrinal alone increased 4F2 expression and also restored 4F2

expression in cells treated with GCN2iB (**Supplementary Figure 14A**). In addition,

overexpression of the PP1-targeting protein GADD34 [42], reduced phosphorylation of eIF2α

and expression of ATF4 and SLC3A2 (4F2) (**Supplementary Figure 14B**). Further supporting

the central role of activation of GCN2 in the induction of SLC3A2 (4F2) expression, treatment of

LNCaP cells with halofuginone, an inhibitor of EPRS and prolyl-tRNA charging [43] resulted in

robust activation of GCN2 and increased expression of ATF4 and SLC3A2 (4F2) protein and

mRNA (**Figure 4E and Supplementary Figure 13C**). These results support the model that

activation of GCN2 during nutrient stress is a major inducer of SLC3A2 (4F2) expression.

We next addressed whether SLC3A2 (4F2) expression is linked to the growth phenotype

following loss of GCN2. First, knock-down of SLC3A2 (4F2) expression using siRNA

significantly reduced growth in LNCaP and 22Rv1 cells and led to activation of GCN2 and

increased ATF4 protein levels (**Figure 4F, 4G, and Supplementary File 1**). As was shown

earlier, loss of GCN2 function using siRNA or GCN2iB treatment in cultured LNCaP cells

reduced growth (**Figure 1A and D**). This growth defect was rescued by overexpression of

SLC3A2 (4F2) (**Figure 4H and Supplementary File 1)**. Importantly, overexpression of SLC3A2

(4F2) reduced GCN2 activity, which lowered p-GCN2 and p-eIF2α levels, along with ATF4

protein expression, and provided a growth advantage in both wild-type LNCaP and 22Rv1 cells

(**Supplementary Figure 14C and 14D**). We next tested the impact of SLC3A2 (4F2) knock-

down on intracellular amino acid levels. As anticipated, knockdown of SLC3A2 reduced a

subset of amino acids as compared to knockdown of GCN2 (**Figure 4I**). These results suggest

that expression of *SLC* genes, especially SLC3A2 (4F2), are important for PCa growth. As

illustrated in the model presented in **Figure 5**, GCN2 plays a critical role ensuring that

appropriate expression of key *SLC* genes and associated amino acid transport are tightly linked

to the ISR**.**

GCN2 supports PCa tumor growth and maintains expression of amino acid transporters

in mouse xenograft models.

To examine the contributions of the GCN2/SLC transporter model for PCa growth in an

in vivo model, we first selected the 22Rv1 cell line as it is castration-resistant. Parental (WT)

22Rv1 cells with functional GCN2 or those with the eIF2α kinase deleted (GCN2 KO) were

injected subcutaneously into the flanks of 8-week-old male NSG mice, and tumor volumes were

measured for up to 12 days. Whereas the parental cells showed sustained tumor growth, loss of

GCN2 significantly reduced tumor growth over time, resulting in a significant reduction in tumor

weight at the end of the study (**Figure 6A**). IHC staining of tumor preparations showed

diminished expression of Ki-67, a marker for cell proliferation in 22Rv1 tumors devoid of GCN2

(**Supplementary Figure 15A**). As expected, the GCN2 protein was not detectable in the GCN2

KO tumors as judged by immunoblot and expression levels of the GCN2-targeted transporters

SLC3A2 (4F2), SLC7A11 (xCT), and SLC7A5 (LAT1) were sharply reduced (**Figure 6B**).

Furthermore, the 22Rv1 tumors maintained expression of the ARv7 variant, which provides for

castration-resistance and resistance to enzalutamide (**Figure 6B**). Similar results were obtained

following deletion of GCN2 in the AR-null castration-resistant cell line PC-3 (**Figure 6C, 6D,**

Supplementary Figure 15B, and 15C). Together, these xenograft experiments using the

22Rv1 and PC-3 models support the hypothesis that GCN2 provides significant growth

advantages to PCa and suggests a mechanism involving enhanced SLC transporter expression

that is independent of AR expression.

To determine the importance of ATF4 in tumor growth *in vivo*, we carried out a similar

xenograft study using 22Rv1 cells deleted for ATF4. While loss of GCN2 led to dramatically

reduced tumor growth, deletion of ATF4 only delayed tumor growth relative to the wild-type

parental cells (**Figure 6E**). Furthermore, deletion of GCN2 led to a further decrease in SLC3A2

(4F2), SLC1A5 (ASCT2), and SLC7A1 (CAT1) protein levels as compared to loss of ATF4

(**Supplementary Figure 16A**). Interestingly, deletion of ATF4 resulted in increased activation of

GCN2, increased levels of p-eIF2α, and in some cases increased amino acids relative to GCN2

KO tumors (**Figure 6F and Supplementary Figure 16A**), indicating that loss of ATF4 can

exacerbate cell stress facilitating further activation of GCN2. Consistent with these *in vivo*

findings, knock-down of GCN2 expression in LNCaP cells in culture reduced amino acid levels

to a greater degree as compared to knock-down of ATF4 (**Supplementary Figure 16B**).

Another key feature of our GCN2/SLC transporter model is that this ISR regulator

functions to ensure appropriate uptake of select EAA. To more directly test this model *in vivo*,

we again utilized the xenograft model featuring 22Rv1 GCN2 KO cells. In this experiment, we

provided additional EAA in the drinking water for the tumor bearing mice. While 22Rv1 cells with

functional GCN2 showed tumor growth within 20 days of injection, loss of GCN2 reduced tumor

growth even over the course of 40 days. However, consumption of EAA-supplemented water by

GCN2 KO-tumor bearing mice provided for increased tumor growth approaching that of wild-

type GCN2 tumors, albeit there was a time delay of 20 days for 22Rv1 with wild-type GCN2 to

about 40 days with EAA-supplemented GCN2 KO (**Figure 6G**). Supplementation of the EAA did

not affect tumor growth of parental 22Rv1 with functional GCN2 (**Figure 6G)**. Similar to previous

reports, supplementation of EAA in the drinking water resulted in a slight decrease in body

weight of mice bearing either parental or GCN2 KO tumors [44] (**Supplementary Figure 17A**).

Measurements of free amino acids in tumors indicated that loss of GCN2 reduced intracellular

arginine and lysine concentrations, with histidine and the branched-chain amino acids trending

lower. Dietary supplementation of EAA partially restored these amino acid levels, although only

histidine achieved statistical significance (**Figure 6H and Supplementary Figure 17B**).

Immunoblot measurements showed that loss of GCN2 lowered key ISR markers, including the

key SLC transporters independent of the supplementation of EAA (**Supplementary Figure**

17C). We also tested the impact of overexpression of SLC3A2 (4F2) on growth of 22Rv1

tumors. Similar to supplementation with EAA, overexpression of SLC3A2 (4F2) in GCN2 KO

tumors partially rescued growth and restored tumor amino acid levels (**Figure 6I,**

Supplementary Figure 18A and 18B). These results further support the notion that GCN2 has

a primary function in maintaining EAA and highlight a key role for GCN2 in the regulation of

SLC3A2 (4F2) expression. In keeping with the cultured PCa cell line experiments, GCN2 played

a vital role supporting amino acid homeostasis in these tumor models.

Pharmacological inhibition of GCN2 reduces tumor growth in xenograft models.

327   We next tested the impact of pharmacological inhibition of GCN2 function in cancer cell-

derived and patient-derived xenograft models. Male NSG mice were injected subcutaneously

with LNCaP or 22Rv1 cells or implanted with tumor fragments from an androgen-sensitive tumor

TM00298 [45]. Alternatively, male castrated NSG mice were implanted with tumor fragments

from LuCaP-35 castration-resistant tumors [46]. Tumors were allowed to grow to 50 -100 mm^3

and mice were randomized to vehicle or GCN2iB treatment groups. Treatment with GCN2iB at

30 mg/kg twice daily for 2 - 6 weeks resulted in robust anti-tumor activity in all four models as

shown by a significant delay in the tumor volume, reductions in tumor mass at endpoint, and

reduced expression of Ki-67 (**Figure 7A – 7D and Supplementary Figure 19A**). Importantly,

GCN2iB had no effect on the body weight of treated mice over the course of treatment

(**Supplementary Figure 19B)**, lowered expression of SLC3A2 (4F2) in both 22Rv1 and

TM00298 tumors, and reduced levels of select amino acids in 22Rv1 tumors from treated mice

(**Figure 7E, 7F, and Supplementary Figure 20A and 20B**). These results suggest that GCN2

is important for tumor growth and expression of SLC3A2 (4F2) in both androgen-sensitive and

castration-resistant tumors.

Our results suggest that GCN2 controls expression of multiple amino acid transporters to

regulate amino acid homeostasis in PCa tumors. We next addressed whether expression of p-

GCN2 or GCN2-dependent mRNA transcripts correlated with expression of SLC3A2 (4F2) in

human tumor samples. Again, using a tumor tissue array containing patient samples, we stained

for p-GCN2 and SLC3A2 (4F2). Similar to recent findings [47], SLC3A2 (4F2) staining was

increased in malignant tissue samples as compared to normal tissue (**Supplementary Figure**

21A). Interestingly, SLC3A2 (4F2) showed a moderate, albeit significant, correlation with

expression of p-GCN2 ($r = 0.27$, $p = 0.025$). This correlation was more striking among tissues

from higher grade tumors ($r = 0.47$, $p = 0.002$, Gleason Score 4 and 5) (**Figure 6G**), which was

further supported by staining of p-GCN2 and SLC3A2 (4F2) of core needle biopsy specimens

from patients with high grade prostate cancers (**Supplementary Figure 21B and 21C**). To

further test the GCN2/SLC3A2 (4F2) model, we derived a gene signature from GCN2-

dependent transcripts from our RNA-seq analysis and measured a significant correlation with

SLC3A2 (4F2) expression ($r = 0.53$, $p < 0.0001$) using a cohort of prostate cancer patients

(PRAD) from the TCGA database (**Figure 7H**). These results further support the idea that

pharmacological inhibition of GCN2 may serve as a potent strategy for the treatment of

androgen-sensitive and castration-resistant PCa by limiting the expression of amino acid

transporters including SLC3A2 (4F2).

Discussion

Tumor cells often demonstrate high demand for amino acids to sustain their increased

metabolism, protein synthesis and proliferation [48], and frequently overexpress amino acid

transporters to meet this increased demand [49, 50]. Altered or increased transport of essential

and non-essential amino acids has been suggested to be a point of vulnerability in several

cancers [51, 52]. In this study, we show that the adaptive ISR pathway driven by GCN2 is active

in PCa tissue and can be genetically and pharmacologically targeted in human models of

androgen-sensitive and castration-resistance disease. Loss of GCN2 function by genetic

deletion or pharmacological inhibition sharply reduced PCa growth *in vitro* and reduces tumor

growth in xenograft models. As illustrated in **Figure 5**, GCN2-directs the expression of a large

number of SLC transporter genes, many of which are critical for PCa growth. Among these

GCN2-targeted genes is SLC3A2 (4F2), a type II integral membrane protein which is known to

form dimers with SLC7 family members including SLC7A5 (LAT1) and SLC7A11 (xCT), each

contributing to uptake of specific amino acids [40]. Loss of GCN2 in PCa reduced free amino

acids both *in vitro* and in tumors derived from xenograft models. Addition of EAAs to the

medium of cultured cells rescued the growth defect associated with GCN2 inhibition;

furthermore, supplementation of EAAs in the drinking water of mice injected subcutaneously

with GCN2 KO PCa cells substantially enhanced tumor growth *in vivo*. While several strategies

to deprive cancer cells of nutrients have been contemplated [51], our results suggest that

inhibition of GCN2 represents a unique treatment paradigm to starve prostate tumors of EAAs.

Further supporting the utility of GCN2 as a therapeutic target for PCa, a small molecule inhibitor

of GCN2 significantly slowed tumor growth in both CDX and PDX models of androgen-sensitive

and castration-resistant disease irrespective of AR-status.

GCN2 is activated by uncharged tRNAs that accumulates during amino acid starvation

or by ribosome collisions [15, 53-55]. As indicated by p-GCN2, this eIF2α kinase is induced in a

variety of different proliferating PCa cell lines and in malignant prostate tissue. While our results

suggest that amino acid limitations and accompanying uncharged tRNA contribute to GCN2

activation in PCa, it is noted that other mechanisms can contribute to regulation of GCN2. For

example, a mitochondrial Rho GTPase 2 (MIRO2) overexpressed in PCa was recently shown to

bind to GCN1, a critical activator of GCN2 [56], and certain mitochondrial defects were reported

to contribute to activation of GCN2 and the ISR [57]. Given that these different PCa models

represent those that are androgen-sensitive, castration-resistant, resistant to enzalutamide, or

lack AR expression altogether, our study suggests that GCN2 functions to control PCa growth

independent of AR function. Supplementation of EAAs reduces p-GCN2 and partially rescued

the growth deficiency of GCN2 KO cells, suggesting that deficiency of at least some of these

critical amino acids are limiting in proliferating PCa cells. This nutrient limitation would trigger

GCN2 induction of the ISR which features induced expression of SLC transporters, including the

central SLC3A2 (4F2). It is noteworthy that SLC3A2 (4F2) has recently been shown to be

overexpressed in PCa and correlated with reduced progression-free survival [47].

Experiments involving supplementation of individual amino acids to cultured LNCaP cells

knocked down for GCN2 suggest that this eIF2α kinase is critical for maintenance of histidine

homeostasis. Histidine is an essential amino acid whose transport is not clearly understood;

although, SLC7A5 (LAT1) has been suggested to facilitate its uptake [58]. Beyond its critical

role in protein synthesis, histidine also functions as an important antioxidant [59] and its

catabolism has been linked to nucleotide synthesis in some cancers [60]. Additional studies will

be required to delineate how histidine contributes to PCa metabolism and growth. Whereas

histidine also rescued the growth defect of GCN2 inhibition in the related MR49F cells, only

lysine alleviated the slow growth of cultured 22Rv1 GCN2 KO cells. These in vitro experiments

suggest a more complex picture where the limiting amino acid is cell-type specific or influenced

by specific genetic alterations which might influence the threshold or mechanism of GCN2

activation. It is acknowledged that amino acids included in the media used in the in vitro

experiments are typically present at high physiological levels so the importance of any given

EAA may be masked by the abundance of these nutrients in the culture medium. Consistent

with this notion, the analysis of 22Rv1 GCN2 KO tumors showed multiple EAAs were reduced.

Activation of GCN2 triggers the ISR by the induction of ATF4, a key transcriptional

activator. ATF4 contributes to PCa growth induced by GCN2, as depletion of ATF4 in multiple

PCa cell lines led to growth inhibition in culture. Previous studies have suggested that ATF4 is

important for growth in multiple cancers [24, 61] including PCa [62]. However, under our

experimental conditions, whereas loss of GCN2 led to a sharp reduction in tumor growth in the

22Rv1 xenograft model, deletion of ATF4 only delayed tumor growth. These results suggest that

while ATF4-directed transcriptional activation is important for PCa proliferation, GCN2 functions

by additional mechanisms, including preferential translation of multiple ISR-target genes.

As noted above activated GCN2 induces a program of genes that are important for

maintenance of amino acid homeostasis. We focused on the critical role of GCN2 in controlling

expression of SLC3A2 (4F2), albeit a collection of SLC genes required GCN2 for full

expression. While SLC3A2 (4F2) is an important contributor to amino acid homeostasis,

deletion of SLC3A2 (4F2) in LNCaP cells led to depletion of only a portion of the free amino

acids that were disrupted by loss of GCN2, consistent with its requirement with a subset of

amino acid transporters [40]. Furthermore, overexpression of SLC3A2 (4F2) only partially

restored growth of 22Rv1 GCN2 KO tumors *in vivo* and did not affect the progression of

parental 22Rv1 tumors. These results indicate that the central role of GCN2 for ensuring

appropriate amino acid homeostasis and enabling PCa proliferation involves induction of

multiple SLC transporters. In addition, the reduction in ATF4 expression that accompanies loss

of GCN2 function reduces expression of several amino acid biosynthetic genes which would

exacerbate the observed effects on uptake of NEAAs. Taken together, these observations

suggest that inhibition of GCN2 may be a more effective therapeutic strategy for starving PCa of

amino acids as compared to therapies that target individual SLC transporters. In summary, our

results indicate that GCN2 may represent a novel therapeutic target for prostate cancer. The

therapeutic landscape for prostate cancer patients is rapidly evolving but AR still remains the

focus. Unfortunately, resistance to androgen deprivation therapies continue to be a major

challenge. The development of new therapeutic strategies that work independently from

inhibition of the AR axis is highly clinically relevant and based on our findings, inhibitors of

GCN2 should be considered further for treatment of androgen-sensitive and castration-resistant

prostate cancers.

Materials and Methods

Cell culture and treatments. LNCaP clone FGC (ATCC Cat. #CRL-1740, RRID:CVCL_1379),

C4-2B (ATCC Cat. #CRL-3315, RRID:CVCL_4784), CW22Rv1 (referred to as 22Rv1, ATCC

Cat. #CRL-2505, RRID:CVCL_1045), PC-3 (ATCC Cat. #CRL-1435, RRID: CVCL_0035), and

HEK-293T (ATCC Cat. #CRL-3216,RRID:CVCL_0063) cells were purchased from the American

Tissue Type Collection (ATCC, Manassas, VA). LAPC-4 (RRID:CVCL_4744) cells were

provided by Dr. Dominic J. Smiraglia (Roswell Park Comprehensive Cancer Center, Buffalo,

NY). MR49F (RRID:CVCL_RW53) cells were obtained from Dr. Timothy Ratliff (Purdue

University, West Lafayette, IN). LNCaP, C4-2B, and 22Rv1 cells were maintained in phenol-free

RPMI-1640 (Gibco, Cat #11835-030) supplemented with 10% FBS (HyClone, Cat #SH30071).

LAPC-4 cells were maintained in phenol-free RPMI-1640 supplemented with 10% FBS as

described above and 10 nM dihydrotestosterone (Sigma-Aldrich, Cat. #D-073). MR49F cells

were maintained in phenol-free RPMI-1640 supplemented with 10% FBS as described above

and 10 µM Enzalutamide (MedChemExpress, Cat. #HY-70002). PC-3 cells were maintained in

Ham's F-12K (Kaighn's) media (Gibco, Cat. #21127-022) or Human Plasma-Like Media (HPLM ,

Gibco Cat. #A4899101) supplemented with 10% FBS. HEK-293T cells were maintained in

DMEM (Corning, Cat. #10-013-CV) supplemented with 10% FBS. All cell lines were maintained

at 37°C in a fully humidified atmosphere containing 5% CO_2. For cell treatments, cells were

plated at 250,000 cells per well in 6-well plates in standard culture media as described above

and allowed to attach overnight. Media was changed to standard culture media containing 500

nM to 10 μM GCN2iB or DMSO as a control or standard media lacking histidine (MyBiosource,

Cat #MBS652918) and incubated for the indicated time period. GCN2iB [34] was synthesized at

Sun-shinechem (Wuhan, China). As indicated, salubrinal (Tocris Biosciences, Cat #405060-95-

9) was added to the culture media at a concentration of 50 μM. To measure bulk protein

translation, 1 μM puromycin (MP Biomedicals Cat #194539) was included in the culture medium

for the last 15 minutes of treatment with GCN2iB, and protein lysates were probed and

quantified by immunoblot analyses using anti-puromycin antibody [63].

Immunoblot analysis. Whole cell extracts and tumor lysates were prepared in 1% SDS lysis

buffer supplemented with 1X Halt Protease and Phosphatase Inhibitor Cocktail (Thermo

Scientific, Cat. #1861281). Tumors were homogenized using a Kinematica Polytron tissue

homogenizer. Lysates were boiled for 10 minutes, sonicated, and centrifuged at 20,000 x g to

remove insoluble material. Protein concentrations were determined by DC Protein Assay (Bio-

Rad, Cat. #5000112) in technical triplicate using BSA as the protein standard. Equal amounts of

protein samples, which ranged from 10 - 20 μg between experiments, were separated on SDS-

PAGE gels, transferred to nitrocellulose membranes, and used for immunoblot analysis. The

primary antibodies used were as follows: phospho-GCN2-T899 (Abcam Cat. #ab75836,

RRID:AB_1310260), total GCN2 (Cell Signaling Technology Cat. #3302, RRID:AB_2277617),

total PERK (Cell Signaling Technology Cat. #3192, RRID:AB_2095847), total HRI (Santa Cruz

Biotechnology Cat. #sc-365239, RRID:AB_10843794), total PKR (Cell Signaling Technology

Cat. #12297, RRID:AB_2665515), phospho-eIF2α-S51 (Abcam Cat. #ab32157,

RRID:AB_732117), total eIF2α (Cell Signaling Technology Cat. #5324, RRID:AB_10692650),

ATF4 (Cell Signaling Technology Cat. #11815, RRID:AB_2616025) or custom rabbit polyclonal

antibody which was prepared against full-length recombinant human ATF4 protein and affinity

purified, ASNS (Cell Signaling Technology, Cat. #20843S), TRIB3 (Abcam Cat. #ab75846,

RRID:AB_1310768), GADD34 (Proteintech Cat. #10449-1-AP, RRID:AB_2168724),

SLC7A5/LAT1 (Cell Signaling Technology Cat. #5347, RRID:AB_10695104), SLC7A11/xCT

(Cell Signaling Technology Cat. #12691, RRID:AB_2687474), SLC3A2/4F2 (Cell Signaling

Technology Cat. #47213, RRID:AB_2799323), SCL7A1/CAT1 (Proteintech Cat. #14195-1-AP,

RRID:AB_2190723), SLC1A4/ASCT1 (Cell Signaling Technology Cat. #8442,

RRID:AB_10828382), SLC1A5/ASCT2 (Cell Signaling Technology Cat. #5345,

RRID:AB_10621427), AR (Cell Signaling Technology Cat. #5153, RRID:AB_10691711),

puromycin (Millipore Cat. #MABE343, RRID:AB_2566826), β-actin (Sigma-Aldrich Cat. #A5441,

RRID:AB_476744), β-tubulin (Cell Signaling Technology Cat. #2146, RRID:AB_2210545).

Immunoblot signals were visualized by enhanced chemiluminescence (ECL) using Clarity

Western ECL Substrate (Bio-Rad, Cat. #170-5060) or SuperSignal West Femto Maximum

Sensitivity Substrate (ThermoFisher Scientific, Cat. #34094) and images were captured using a

ChemiDoc™ Imaging System (Bio-Rad, RRID:SCR_019037). Prestained protein standards

(Bio-Rad Cat. #1610377) were included on each immunoblot, captured as a separate merged

image file, and are indicated in each immunoblot panel.

RNA Interference and proliferation assays. Small Interfering RNAs (siRNAs) were purchased

from Dharmacon and sequences are shown in **Supplementary Table 2**. The siRNAs that

showed the most robust depletion of the targeted genes in the indicated cells were selected for

experiments. For growth assays LAPC-4, C4-2B, MR49F, 22Rv1, or PC-3 cells were seeded

into 96-well plates at 2500 – 5000 cells per well in normal growth media as described above and

allowed to attach overnight. Cells were transfected with 1 pmole of siRNA using Lipofectamine

RNAiMAX Transfection Reagent (Thermo Scientific, Cat. #13778030) or DharmaFECT 3

Transfection Reagent (Horizon, Cat. #T-2003-01) according to manufacturer's protocol and

incubated overnight (considered Day 0). Alternatively, where indicated, growth media was

changed to serum-free Accell Delivery Media (Horizon, Cat. #B-005000) and transfected with 1

pmole of Accell siRNA and incubated 48 hours at which time cells were changed back to normal

growth media (considered Day 0). Cells were incubated for an additional 2 – 6 days and cell

growth was quantified using CellTiter-Glo® 2.0 Reagent (Promega, Cat. #924C). For

immunoblot analysis, PCa cells were seeded into 6-well plates at 250,000 – 350,000 cells per

well and transfected with 30 pmoles of siRNA as described above and incubated for 48 hours

prior to lysate preparation as described above.

***In vitro* amino acid supplementation.** For amino acid supplementation experiments, 1X MEM

Amino Acids Solution (Gibco, Cat. #11130051) was used for essential amino acid (EAA) mix

and 1X MEM Non-Essential Amino Acids Solution (Gibco, Cat. #11140050) was used for non-

essential amino acid (NEAA) mix. Experiments utilizing single amino acids were performed

using the following concentrations: 600 µM L-Arginine hydrochloride, 100 µM L-Cystine

dihydrochloride, 200 µM L-Histidine hydrochloride, 400 µM L-Isoleucine, 400 µM L-Leucine, 395

µM L-Lysine hydrochloride, 100 µM L-Methionine, 200 µM L-Phenylalanine, 400 µM L-

Threonine, 50 µM L-Tryptophan, 200 µM L-Tyrosine, and 400 µM L-Valine. All amino acids were

dissolved in water and were purchased from Sigma-Aldrich (St. Louis, MO).

Amino acid uptake measurements. LNCaP, parental 22Rv1, or 22Rv1 GCN2 KO cells were

plated in 24-well plates (50,000 cells) in standard culture media as described above. LNCaP

cells were treated with vehicle (DMSO) or 2 µM GCN2iB for 24 hours. Cells were washed with

pre-warmed sodium-free Hank's Balanced Salt Solution (HBSS) and treated with 100 nM ^{3}H-

Arginine (5 µCi/ml, Perkin-Elmer), 1 µM ^{14}C-Leucine (0.3 µCi/ml, Perkin-Elmer), or 1 µM ^{14}C-

Lysine (0.3 µCi/ml, Perkin-Elmer) in HBSS containing the appropriate un-labelled amino acid

(100-200 µM Arginine, 1-20 µM Leucine, or 1 µM Lysine) at 25 °C for 2 minutes. Transport was

stopped by adding the appropriate excess cold-amino acid (non-radioactive) in HBSS. Cells

were washed with PBS, lysed with Passive lysis buffer (Promega, Cat #E1941), and a portion

added to scintillation cocktail and counted. The protein concentration of each lysate was

determined as described above and amino acid uptake was represented as nmol/µg of total

protein/min. Each measurement was performed in triplicate.

Plasmid construction. Plasmid pDONR221-SLC3A2_STOP was a gift from RESOLUTE

Consortium & Giulio Superti-Furga (Addgene plasmid # 161379, RRID:Addgene_161379) [39].

Plasmid pENTR223-1 GCN2 (NM_001013703) was purchased from Transomic Technologies

and pLX303 was a gift from David Root (Addgene plasmid # 25897, RRID:Addgene_25897).

GCN2 coding sequence was subcloned into the lentiviral vector pLVX-IRES-Hygromycin

(Clontech, Cat #632185) using an In-Fusion HD Cloning plus kit (TaKaRa, Cat. #638920)

according to manufacturer's protocol yielding plasmids pLVX-GCN2-hygro. The SLC3A2 (4F2)

coding region was subcloned into the lentiviral vector pLX303 by Gateway cloning using

Gateway LR Clonase II Enzyme Mix (Invitrogen, Cat. #11791-020) according to manufacturer's

protocol yielding plasmid pLX303-4F2-blast. Plasmid pMSCV-GADD34-puro contains the

human PPP1R15A (GADD34, NM_014330.5) gene subcloned into the NcoI/EcoRI sites of the

retroviral vector pMSCV-puro (Clontech Laboratories, Mountain View, CA).

Construction of cell lines. GCN2 and ATF4 knock-out cell lines were generated using

CRISPR/Cas9 Human Gene Knockout Kits (Origene, Cat. #KN412459 and #KN402333) using

hGCN2g1 (5'-AATTTAGTTTTGTACCCTCA-3') and hATF4g1 (5'-

CTTCCTGAGCAGCGAGGTGT-3'); respectively, following the manufacturer's protocol. Cell

lines overexpressing GCN2 or SLC3A2 (4F2) were created by lentivirus transduction. Briefly,

HEK-293T cells were cultured in DMEM as described above and were transfected with lentivirus

constructs and ViraPower™ Lentiviral Packaging Mix (Invitrogen, Cat #35-1275) using

Lipofectamine 2000 (ThermoFisher Scientific, Cat # 11668030) according to manufacturer's

protocol and virus containing supernatant was collected over 48 – 72 hours, concentrated using

Lenti-X™ Concentrator (TaKaRa, Cat. #631232), and stored in aliquots at -80 °C. 22Rv1 or PC-

3 cells were transduced with virus-containing culture media supplemented with 10 µg/ml

Polybrene (Millipore, Cat #TR-1003-G) for 24 hours and replaced with fresh growth media. At 72

hours post-transfection, growth media was changed to media supplemented with 250 µg/ml

Hygromycin B (Thermo Cat #10687010) or 10 µg/ml blastocidin S (Sigma, Cat #SBR00022) to

select for stable pools as appropriate. The appropriate empty vector was used as a control for

all experiments.

qRT-PCR analysis. Total RNA was isolated from cultured cells or tumor tissue using TRIzol™

LS Reagent (Invitrogen, Cat. #10296010) and quantified using a Nanodrop Spectrophotometer

(ThermoFisher Scientific). Total RNA (1 µg) was converted to cDNA using a High-Capacity

cDNA Reverse Transcription Kit (ThermoFisher Scientific, Cat. #4368813) according to the

manufacturer's protocol. Following cDNA synthesis, samples were diluted five-fold in molecular

biology grade water and PCR amplification was carried out using PowerUp SYBR Green Master

Mix (ThermoFisher Scientific, Cat. #A25742) using an Applied Biosystems QuantSudio5 PCR

system. The primers used for measuring *ATF4* and *SLC3A2 (4F2hc)* transcript levels are shown

in **Supplementary Table 3**. The relative abundance of each transcript was calculated using the

ΔΔCT method with GAPDH serving as an internal control and data are presented normalized to

each control group.

Cell cycle analysis. Cell cycle analysis on LNCaP, MR49F, C4-2B, 22Rv1, and PC-3 cells was

performed by co-staining with a Click-iT™ EdU Alexa Fluor™ 488 Flow Cytometry Assay Kit

(Invitrogen, Cat. #C10425) and FxCycle Violet Ready Flow Reagent (Invitrogen, Cat. # R37166)

according to manufacturer's protocol. Cell treatments and siRNA knockdowns were carried out

for 48 hours as described above before analysis using an LSR II (LSR4) flow cytometer (BD

Biosystems). A minimum of 10,000 events were analyzed per condition and all experiments

were performed using triplicate wells. Data analysis was performed using FlowJo (FlowJo LLC,

TreeStar) and data are presented as the percentage of cells in G1, S, or G2-M phases of the

cell cycle.

Amino acid analysis. Levels of different amino acids were measured in cultured cells or tumor

lysates by HPLC or LC-MS. Cell pellets or tissues were resuspended in 0.1% formic acid in

methanol, vortexed briefly, and subjected to three cycles of freezing in liquid nitrogen followed

by thawing at 42 °C for 10 minutes. Clarified lysates were analyzed by reverse-phase liquid

chromatography with appropriate standard curves as described previously [64] or using liquid

chromatography tandem mass spectrometry (LC-MS/MS) with stable isotope dilution. For the

MS-based method, lysates were spiked with a mixture containing 37 amino acid isotopic internal

standards and deproteinated using a solution of 90% acetonitrile, 10 mM ammonium formate,

and 0.15% formic acid. Clarified lysates were analyzed using a biphasic LC-MS/MS approach

first with a HILIC separation [65] followed by a mixed-mode chromatographic separation

[66]. Quantification of amino acid levels was determined by fitting response ratios to an eight-

point calibration curve generated using verified reference material for each of the 43 amino

acids quantified.

CRISPR knock-out screen. A human SLC knockout library [39] was purchased from Addgene,

(Cat. #132552). Lentivirus was generated as described above. LNCaP and 22Rv1 cells were

transduced at a multiplicity of infection (MOI) of 0.5 and selected with 1 µg/ml puromycin (MP

Biomedicals Cat #194539) for 7 days. Three independent screens for both LNCaP and 22Rv1

were performed. Cells were then cultured for an additional 4 weeks and genomic DNA (gDNA)

was isolated using DNeasy Blood and Tissue Kit (Qiagen, Cat. # 69506). sgRNA sequences

were PCR amplified from gDNA using Platinum SuperFi II DNA Polymerase (ThermoFisher

Scientific, Cat #12369002) and primers sgRNA_F and sgRNA_R. A second round of PCR was

used to add Illumina adaptor sequences using primers IlluminaAd_F and IlluminaAd_R. PCR

primers are shown in **Supplementary Table 3**. Purified PCR products were sequenced using

next generation sequencing (NGS) at GeneWiz (South Plainfield, NJ). Fastq files were trimmed,

aligned, and analyzed using PinAPL-Py [67] and raw read counts were normalized as counts

per million reads (cpm). The fold-depletion for each sgRNA was calculated for LNCaP or 22Rv1

relative to control human SLC knockout library. sgRNA rankings and significance (Sidak

correction method) were determined according to the negative binomial model implemented in

PinAPL-Py [67]. Gene-depletion significance was determined by a permutation test (10^4

permutations), and *p* values were corrected for multiple testing using the Sidak correction

method.

RNA-seq and network analyses. LNCaP cells were seeded into six-well culture plates and

treated with either vehicle (DMSO) or 2 μM GCN2iB in standard culture media as described

above for 6 or 24 hours as indicated. Total RNA was extracted from cells using a RNeasy Plus

Kit (Qiagen, Cat. #74034) according to the manufacturer's instructions. RNA quality control,

library preparation, and NGS paired-end sequencing on Illumina HiSeq platform was carried out

at GeneWiz (South Plainfield, NJ). Raw fastq data files were trimmed to remove adaptor

sequences using cutadapt [68] and the quality of the trimmed reads was analyzed by FASTQC

[69] available in the Trim Galore package [70]. The trimmed paired sequencing reads were

aligned to human reference genome GRCh38 downloaded from ENSEMBL using HISAT2 [71],

resulting in concordant alignment of >92%. FeatureCounts [72] from the Subread package was

used to count the number of reads mapping to unique exonic sequences and was used for

differential gene expression (DEG) analysis using DESeq2 R-package [73], comparing the

DMSO treated control group to the GCN2iB-treated groups.

The computer cluster Carbonate that was configured for high-performance analyses was

used to perform all the steps of sequencing analysis from trimming to differential analysis.

Volcano plots were created from differentially expressed genes using GraphPad Prism.

Differentially expressed genes with a log2 fold change of $\geq \pm 1$ and adjusted *p*-values of ≤ 0.05

were considered significant for further analysis. Gene set enrichment analysis [74, 75] using the

Hallmark and Curated (C2) gene sets from the Molecular Signatures Database [76, 77] was

used to identify enriched pathways from lists of significant differentially expressed genes. The

GCN2-dependent gene signature was derived from genes showing $\leq$ -2-fold change (fdr ≤ 0.05)

following treatment with GCN2iB for 6 hours in our RNA-seq analysis. The RNA-seq data

generated in this study has been deposited in the NCBI's Gene Expression Omnibus (GEO)

database under the accession code GSE196252 [Reviewer's Token #kpkvcqcunpidhol].

tRNA charging measurements. The levels of charged (aminoacylated) tRNAs genome-wide

were measured using the CHARGE-seq method [38] with some protocol adjustments as

previously described [78]. The CHARGE-seq method relies on the simultaneous sequencing of

both a charged and uncharged tRNA library. A brief description is provided here with more

detailed information provided in the methods supplement. LNCaP cells were treated with vehicle

(DMSO), GCN2iB, or GCN2iB in media supplemented with essential amino acids for 8 hours.

Total RNA was extracted using TRIzol (Life Technologies, Cat. #15596018) and a portion (2 μg)

of total RNA for each sample was oxidized with $NaIO_4$ or left untreated. Following deacylation,

libraries were prepared as described in the supplementary methods and subjected to Illumini

HiSeq paired-end sequencing. The fraction charged for each tRNA isodecoder within each

sample was determined by summing the normalized counts for charged and uncharged reads

for each gene corresponding to that isodecoder, and the mean was then averaged for each of

the four replicates for each treatment. The mean fraction charged +/- SD for each isodecoder

was visualized using Graphpad Prism. Statistical significance in difference of the mean fraction

charged was determined using a Welch's unpaired *t*-test with Benjamini-Hochberg fdr correction

to account for repeated measures. The CHARGE-seq data generated in this study has been

deposited in the NCBI's Gene Expression Omnibus (GEO) database under the accession code

GSE196251 [Reviewer's Token #kpkvcqcunpidhol]. Levels of total and aminoacylated tRNA^{His}

were also carried out by qRT-PCR as previously described [78] using primers specific for the

GTG-tRNA^{His} isodecoder tRNA^{His}-FW and tRNA^{His}-REV (**Supplementary Table 3**).

Animal studies. All animal experiments were approved by the Institutional Animal Care and

Use Committee (IACUC) at Indiana University School of Medicine and comply with all

regulations for ethical conduct of animal research. Male 4 – 6 week old NSG (NOD.Cg-

Prkdcscid Il2rgtm1Wjl/SzJ, RRID:IMSR_JAX:021885) mice were purchased from the In Vivo

Therapeutics Core at Indiana University School of Medicine. GCN2iB used for animal studies

was formulated in 0.5% Methyl Cellulose (Sigma-Aldrich, Cat. #M0262) containing 5% DMSO.

Mice were dosed with vehicle or 30 mg/kg GCN2iB twice daily for 5 days/week by oral gavage

as described in the legends to the figures. For LNCaP, 22Rv1, and PC-3 cell line-derived

xenograft models, 2×10^6 cells in 100 µl of serum-free RPMI-1640 media containing 50%

Matrigel (Corning, Cat. #356237) were injected subcutaneously into the dorsal flank of mice.

Once the tumors reached a palpable stage (~50 – 100 mm^3), the animals were randomized to

different treatment groups using RandoMice v1.1.5 [79]. The LuCaP-35 castration-resistant

(LuCaP-35 CR) PCa PDX model [46, 80] was a generous gift from Dr. Timothy Ratliff (Purdue

University, West Lafayette, IN). LuCaP-35 CR tumors were maintained in male castrated NSG

mice for all experiments to conserve castration-resistant status. TM00298 PCa PDX model [45]

was obtained from Jackson Laboratory and male NSG mice were engrafted with ~1 mm^3 PDX

tumor fragments and randomized once the tumors reached a palpable stage (~50 – 100 mm^3).

For all xenograft studies, tumor volume was monitored twice per week in a blinded fashion using

a digital caliper and the formula: Tumor Volume = (width^2 × length)/2. Body weight was

determined once per week during the course of each study. At the end of the studies, mice were

euthanized using CO_2 and tumors were harvested, weighed, and fixed for

immunohistochemistry staining or flash frozen for immunoblot analyses. For the amino acid

supplementation experiment *in vivo*, mice bearing 22Rv1 WT or 22Rv1 GCN2 KO tumors were

supplemented with essential amino acids (1.5 mg amino acids/ g body weight) in the drinking

water. The control group received drinking water with no amino acid supplementation.

Supplementation of amino acids was started at the same time as tumor cell implantation and

was maintained for the remainder of the study. Amino acids were dissolved in water in

quantities determined by calculating average daily intake of drinking water for mice as

previously reported [44]. Composition, relative percentage, and an estimate of dietary intake of

each amino acid from the drinking water is reported in **Supplementary Table 4**.

Immunohistochemistry. Tumor tissue specimens were fixed for 24 hours in neutral buffered

formalin, paraffin-embedded and sectioned (5 μm). Prostate core needle biopsy specimens

were obtained from the Indiana University Comprehensive Cancer Center Tissue Procurement

and Distribution Core and approval was granted by the Institutional Review Board (IRB) at the

Office of Research Administration at Indiana University. Human tumor tissue microarrays were

purchased from US Biomax (Rockville, MD, Cat. # PR1921b and PR807c). Tissue sections

were deparaffinized and rehydrated through serial graded xylene and alcohol washes. H&E

staining was performed using standard methods. Antigen unmasking was achieved by boiling

the slides in sodium citrate buffer (10 mM Sodium citrate, 0.05% Tween 20, pH 6.0) for 10 – 15

minutes. Sections were further incubated in hydrogen peroxide to reduce endogenous activity.

Tissue blocking was performed with 5% Horse Serum (Vector Laboratories Cat. #PK-7200) in

phosphate-buffered saline (PBS) and incubated with primary antibodies: Ki-67 (1:200, Abcam

Cat. #ab16667, RRID:AB_302459), p-GCN2-T899 (1:100, Abcam Cat. #ab75836,

RRID:AB_1310260), and SLC3A2 (4F2hc/CD98) (1:500, Cell Signaling Technology Cat.

#47213, RRID:AB_2799323). After primary antibody incubation, tissue sections were washed,

incubated in horseradish peroxidase-conjugated anti-rabbit secondary antibody according to the

manufacturer's protocol (Vector Laboratories, Cat #PK-7200), enzymatically developed in

diaminobenzidine (DAB), and counterstained with hematoxylin. Tissue sections were then

dehydrated and mounted with coverslips sealed with Cytoseal™ 60 mounting media

(ThermoFisher Scientific, Cat #23-244257). Stained sections were scanned under brightfield

using a Leica Aperio AT2 slide scanner (Leica Biosystems). Quantification was performed using

Quantitative Pathology & Bioimage Analysis software (QuPath). Representative images were

acquired using an EVOS M5000 cell imaging microscope (Life Technologies).

Statistical Analysis. All statistical tests were carried out using GraphPad Prism version 9.2.0

for Windows as indicated in the legends to the figures or as outlined in **Supplemental File 1**.

Figure 1

Figure 1. GCN2 promotes growth of PCa cells. (A) Expression of the indicated eIF2 kinase was reduced in LNCaP cells using gene specific siRNAs. Two different siRNAs were used for knock-down of each eIF2α kinase and compared to scrambled siRNA control. Cell growth was measured in replicate wells (N = 5) for up to 6 days and is plotted as fold change (mean ± SD) relative to day 0. Statistical analysis is shown in **Supplementary File 1**. **(B)** LNCaP cells were transfected with two different siRNAs targeting GCN2 or scramble siRNA and cell lysates were prepared and immunoblotted for the indicated proteins. Molecular weight markers are shown in kilodaltons. **(C)** Expression of GCN2 was knocked-down in LAPC-4, C4-2B, MR49F, 22Rv1, or PC-3 cells using two different siRNAs and compared to scrambled siRNA control. Cell growth was measured for up to 6 days in replicate wells (N = 5) as described in **A**. Statistical analysis is shown in **Supplementary File 1**. **(D)** LNCaP cells were treated with indicated concentrations of GCN2iB and cell growth was measured for up to 6 days as described in **A**. Statistical analysis is shown in **Supplementary File 1**. **(E)** LNCaP cells were treated with GCN2iB (2 µM) or DMSO control for 24 hours and protein lysates were analyzed by immunoblot using antibodies that recognize total or phosphorylated GCN2-T899, total or phosphorylated eIF2α−S51, ATF4, or actin as indicated. **(F)** Levels of p-GCN2 were measured in prostate tumor microarrays (Biomax PR1921b and PR807c) using immunohistochemistry (IHC). Staining for p-GCN2-T899 from non-malignant (N = 33) and malignant prostate cancer tissue (N = 88) from patients >50 years old was analyzed and quantified using QuPath to determine the histoscore and is represented as a scatterplot. Statistical significance was determined using an unpaired two-tailed *t*-test, *, *p* ≤ 0.05. Representative images showing p-GCN2-T899 staining of normal and malignant prostate tissues are shown. Scale bars shown are 200 µm (main image) and 20 µm (insert).

Figure 2

A

B

Gene Set	NES	p-val
KRIGE_AMINO_ACID_DEPRIVATION	1.75	<0.001
GARGALOVIC_RESPONSE_TO_OXIDIZED_PHOSPHOLIPIDS_RED_UP	1.67	<0.001
REACTOME_FCGR_ACTIVATION	1.67	<0.001
REACTOME_IRE1ALPHA_ACTIVATES_CHAPERONES	1.63	<0.001
REACTOME_AMINO_ACID_TRANSPORT_ACROSS_THE_PLASMA_MEMBRANE	1.62	<0.001
REACTOME_RESPONSE_OF_EIF2AK1_HRI_TO_HEME_DEFICIENCY	1.61	0.001
PACHER_TARGETS_OF_IGF1_AND_IGF2_UP	1.6	<0.001
ZHAN_MULTIPLE_MYELOMA_CD1_VS_CD2_UP	1.58	<0.001
REACTOME_ROLE_OF_PHOSPHOLIPIDS_IN_PHAGOCYTOSIS	1.55	<0.001
ZHAN_MULTIPLE_MYELOMA_CD1_UP	1.54	<0.001
REACTOME_CYTOSOLIC_TRNA_AMINOACYLATION	1.53	0.002
KEGG_GLYCINE_SERINE_AND_THREONINE_METABOLISM	1.53	0.001
KEGG_AMINOACYL_TRNA_BIOSYNTHESIS	1.53	0.001

C

D

E

F

G

Figure 2. GCN2 induces ISR transcriptome featuring expression of amino acid transporters. (A) Volcano plot illustrating $\log_2$ fold change with adjusted *p*-value ($-\log_{10}$) comparing LNCaP cells treated with GCN2iB (2 µM) versus vehicle control (DMSO) for 24 hours. Several amino acid transporters reduced by GCN2iB treatment are highlighted. **(B)** Enrichment plots from Gene Set Enrichment Analysis (GSEA) of gene expression in LNCaP cells treated with GCN2iB (2 µM) for 24 hours versus vehicle control. **(C)** Heat map displaying differential gene expression of *SLC* genes comparing LNCaP cells treated with vehicle (DMSO), or GCN2iB (2 µM) for 6 or 24 hours. *SLC* genes with $\log_2$ fold change ≤ -0.5 and $p ≤ 0.05$ are shown. **(D)** Lysates were prepared from LNCaP cells treated with 2 µM GCN2iB or vehicle control (DMSO) for 6 or 24 hours and immunoblot analysis were carried out using antibodies that recognize ATF4, ASNS, SLC7A11 (xCT), SLC3A2 (4F2), SLC7A1 (CAT1), SLC1A4 (ASCT1), SLC1A5 (ASCT2), or actin. Molecular weight markers are indicated in kilodaltons. **(E)** 22Rv1 WT cells, 22Rv1 GCN2 KO cells, and 22Rv1 GCN2 KO complemented with GCN2 cells were cultured for 24 hours. Lysates were prepared and analyzed by immunoblot for the indicated proteins. **(F)** Amino acid uptake measurements in LNCaP and 22Rv1 cells treated with vehicle (DMSO) or GCN2iB (2 µM) for 24 hours. **(G)** Amino acid uptake measurements for 22Rv1 WT or 22Rv1 GCN2 KO cells cultured for 24 hours. Statistical significance was determined using an unpaired two-tailed *t*-test (N = 4), *$p ≤ 0.05$, **$p ≤ 0.01$, ***$p ≤ 0.001$.

Figure 3

Figure 3. GCN2 is critical for maintenance of free amino acids. (A) Amino acid

measurements of LNCaP cells treated with 2 µM GCN2iB or vehicle control (DMSO) for 8 hours.

Bar graphs in the top panel show high abundance amino acids and the lower panel those with

lower levels. The heatmap on the right shows fold change in amino acid abundance for each

biological replicate of GCN2iB-treated LNCaP cells vs the vehicle with the scale showing the

highest fold change in yellow and lowest in purple. Statistical significance was determined using

an unpaired two-tailed *t*-test. Error bars indicate SD (N = 3); *, $p \leq 0.05$; **, $p \leq 0.01$, ***, $p \leq$

0.001; ****, $p \leq 0.0001$. **(B)** LNCaP cells were treated with vehicle, GCN2iB (2 µM), vehicle +

essential amino acids (EAA), or GCN2iB (2 µM) + EAA, and cell growth was measured for up to

6 days. Error bars indicate SD (N = 5). Statistical analysis is shown in **Supplementary File 1**.

(C) Cell cycle analyses of LNCaP cells treated with vehicle, GCN2iB (2 µM), vehicle + EAA, or

GCN2iB (2 µM) + EAA for 48 hours. Statistical significance was determined using a one-way

ANOVA with Tukey's multiple comparisons. Error bars indicate SD (N=3); ***, $p \leq 0.001$; ****,

$p \leq 0.0001$. **(D)** Genome-wide tRNA charging analysis (CHARGE-seq) of LNCaP cells treated

with vehicle (DMSO), GCN2iB (2 µM), or GCN2iB (2 µM) + EAA for 8 hours. The tRNA charging

ratio is shown as a bar graph with fold change compared to vehicle. Only tRNA isoacceptors

measured in LNCaP cells are shown. Error bars indicate SD (N = 4). **(E)** tRNA charging

percentage for tRNA[His] in LNCaP cells treated with vehicle, GCN2iB, or GCN2iB + EAA.

Statistical significance was determine using a one-way ANOVA with Tukey's multiple

comparisons (N = 4); ***, $p \leq 0.001$; ****, $p \leq 0.0001$. **(F)** LNCaP cells were treated with vehicle,

GCN2iB (2 µM), GCN2iB (2 µM) + EAA, or GCN2iB (2 µM) combined with the indicated

individual amino acids. Cell growth was measured at 4 days in triplicate wells (N = 3). Statistical

significance was determined using a one-way ANOVA with Tukey's multiple comparisons. Error

bars indicate SD; ****, $p \leq 0.0001$. **(G)** Cell cycle analysis of LNCaP cells were treated with

vehicle, GCN2iB (2 µM), GCN2iB (2 µM) + histidine (200 µM), or with media lacking histidine for

48 hours. Statistical significance was determined using a one-way ANOVA with Tukey's multiple

comparisons. Error bars indicate SD (N = 3); ***, $p \leq 0.001$; ****, $p \leq 0.0001$. **(H)** LNCaP cells

were cultured in normal media, media supplemented with EAA mix, or media supplemented with

histidine (200 µM) for 24 hours. Lysates were analyzed by Immunoblot using antibodies that

recognize total or phosphorylated GCN2-T899, total or phosphorylated eIF2α−S51, ATF4, or

actin. Molecular weight markers are presented in kilodaltons for each immunoblot panel.

Figure 4

Figure 4. GCN2 is critical for expression of SLC3A2 and facilitates amino acid transport.

(A) Gene-level depletion for LNCaP and 22Rv1 cells. The average log2 fold-change for the sgRNAs for each gene is shown on the x axis. Significantly depleted genes ($p \leq 0.05$) in LNCaP or 22Rv1 are indicated. Circle size indicates the number of significant sgRNAs. *SLC*-genes in red are dependent on GCN2 for expression. **(B)** Plot of $-Log_{10}$ (p-value) for depleted genes identified in CRISPR screen for LNCaP vs. 22Rv1 cells. Significantly depleted genes ($p \geq 0.05$) in LNCaP, 22Rv1 or both cell lines are indicated. SLC-genes in red are GCN2-dependent. **(C)** Lysates from LNCaP cells were treated with 2 µM GCN2iB for 6 or 24 hours, or with vehicle (DMSO) were analyzed by immunoblot analyses using antibodies that recognize total or phosphorylated GCN2-T899, ATF4, SLC3A2 (4F2), or actin. Molecular weight markers are indicated in kilodaltons for the panels. **(D)** LNCaP cells were cultured in standard culture conditions (NM: normal media), media supplemented with 200 µM histidine (+ His), or media depleted of histidine (- His) for 24 hours. Lysates were analyzed by Immunoblot analyses using antibodies that recognize total or phosphorylated GCN2-T899, ATF4, SLC3A2, or actin. **(E)** LNCaP cells were treated with 100 nM halofuginone (HF) for 2 and 6 hours or vehicle (DMSO). Lysates were analyzed by Immunoblot using antibodies that recognize the indicated proteins. **(F)** SLC3A2 (4F2) expression was reduced in LNCaP or 22Rv1 cells using two different siRNAs or scramble siRNA as a control. Cell growth was measured in replicate wells (N = 5) for up to 6 days and are plotted relative to day 0 (mean ± SD). Statistical analysis is shown in **Supplementary File 1**. **(G)** LNCaP cells transfected with two different siRNAs targeting 4F2 (SLC3A2) or scramble siRNA for 48 hours. Lysate were prepared and analyzed by immunoblot using antibodies that recognize total or phosphorylated GCN2-T899, total or phosphorylated eIF2α−S51, ATF4, SLC3A2, or actin. **(H)** LNCaP cells stably overexpressing SLC3CA2 or vector control were transfected with two different siRNAs targeting GCN2 or scrambled control. Cells were then treated with GCN2iB (2 µM) or vehicle and growth was measured in replicate wells (N = 5) and is plotted relative to day 0 (mean ± SD). Statistical analysis is shown in

Supplementary File 1. **(I)** Amino acid measurements of LNCaP cells transfected siRNA targeting GCN2 (N = 4), 4F2 (SLC3A2, N = 4), or scramble control (N = 8). Two separate bar graphs show high abundance (top) and low abundance (bottom) amino acids. Statistical significance was determined using a one-way ANOVA with Tukey's multiple comparisons Error bars indicate SD; *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$; ****, $p \leq 0.0001$.

Figure 5

Figure 5. Model for GCN2 regulation of amino acid homeostasis in prostate cancer. Model depicting the role of GCN2 in regulating SLC amino acid transporters. Enhanced translation and altered metabolism driven by oncogenes deplete amino acid pools resulting in accumulation of uncharged tRNAs leading to activation of GCN2. Active GCN2 results in increased expression of SLC amino acid transporters, including SLC3A2 (4F2) to increase uptake of amino acids. Loss of GCN2 function disrupts amino acid homeostasis decreasing proliferation of prostate cancer cells.

Figure 6

Figure 6. GCN2 regulates amino acid transporters ensuring sufficient amino acids for tumor growth in mouse PCa xenograft models. (A) WT or GCN2 KO 22Rv1 (clone 7) cells were injected subcutaneously into the dorsal flank of mice as described in the materials and methods section. Tumor volume (TV) was measured on indicated days and is plotted as average TV ± SEM (N = 4). Statistical significance was determined using a two-way ANOVA with Sidak's multiple comparison; *, $p \leq 0.05$; ****, $p \leq 0.0001$. On the right bar graph, the final tumor weight was measured at endpoint and statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD; **, $p \leq 0.01$. **(B)** Protein lysates were prepared from WT and GCN2 KO 22Rv1 tumors and analyzed by immunoblot to measure total GCN2, ATF4, SLC7A5 (LAT1), SLC7A11 (xCT), SLC3A2 (4F2), androgen receptor (AR), AR splice variant 7 (AR-V7), or actin. Molecular weight markers are indicated in kilodaltons for each immunoblot panel. The levels of the SLC proteins normalized to actin are shown in the bar graph (right panels). Statistical significance was determined using an unpaired two-tail t-test. Error bars indicate SD (N = 4); *, $p \leq 0.05$; ***, $p \leq 0.001$. **(C)** Tumor growth of PC-3 WT and PC-3 GCN2 KO (clone 3) cells was analyzed in a mouse xenograft study as in A. Statistical significance was determined using a two-way ANOVA with Sidak's multiple comparison. Error bars indicate SEM (N = 5); ***, $p \leq 0.001$; ****, $p \leq 0.0001$. On the right bar graphs, the final tumor weight was measured at endpoint. Statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD (N = 5); *, $p \leq 0.01$. **(D)** Protein lysates were prepared from the PC-3 WT and PC-3 GCN2 KO tumors and analyzed by immunoblot for the indicated proteins. (Right panels) Quantification of protein levels of SLC7A5 (LAT1), SLC7A11 (xCT), and SLC3A2 (4F2) normalized to actin are shown in the bar graphs. Statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD (N = 4); **, $p \leq 0.01$. **(E)** 22Rv1 WT (N = 4), 22Rv1 GCN2 KO (clone 7, N = 5) and 22Rv1 ATF4 KO (N = 5) were evaluated in the mouse xenograft model. Tumor volumes were measured on the indicated days. Error bars indicated SEM. **(F)** Amino acid measurements of 22Rv1 WT, 22Rv1

GCN2 KO and 22Rv1 ATF4 KO tumors. Two separate bar graphs show high abundance (top) and low abundance (bottom) amino acids. Statistical significance was determined one-way ANOVA with Tukey's multiple comparisons. Error bars indicate SD (N = 4); #, $p \leq 0.1$; *, $p \leq 0.05$.

(G) 22Rv1 WT and 22Rv1 GCN2 KO (clone 7) cells were analyzed in a xenograft model as described for figure **6A**, with or without supplementation of EAA in the drinking water. Tumor volume was measured on indicated days. 22Rv1 WT (N = 4) and 22Rv1 KO (N = 5) are the same tumor growth curves shown in **6E**. 22Rv1 WT + EAA (N = 5), 22Rv1 GCN2 KO + EAA (N = 5) **(H)** Amino acid measurements for 22Rv1 WT, 22Rv1 GCN2, and 22Rv1 GCN2 KO + EAA tumors. Bar graphs show only amino acids present in EAA supplemented water. Statistical significance was determined using a one-way ANOVA with Tukey's multiple comparisons. Error bars indicated SD (N = 4); #, $p \leq 0.1$; *, $p \leq 0.05$; **, $p \leq 0.01$. **(I)** Tumor growth curves for 22Rv1 WT or 22Rv1 GCN2 KO (clone 11) transduced with SLC3A2 (4F2) lentivirus (WT + SLC3A2 and GCN2 KO + SLC3A2) or empty vector (WT + EV and GCN2 KO + EV). Tumor volumes were measured on indicated days. Error bars indicate SEM (N = 5).

Figure 7

Figure 7. Pharmacological inhibition of GCN2 reduces tumor growth in cell line-derived and patient-derived xenograft models. Male NSG mice were injected subcutaneously with LNCaP (N = 5) **(A)** or 22Rv1 (N=4) **(B)** cells, or alternatively implanted with tumor fragments from an androgen-sensitive tumor TM00298 (N = 5) **(C).** Male castrated NSG mice were implanted with tumor fragments from LuCaP-35 CR tumors (N = 5) **(D).** Mice were treated with vehicle or 30 mg/kg GCN2iB twice daily for 5 days per week and tumor volumes were measured on indicated days. Statistical significance was determined using a two-way ANOVA with Sidak's multiple comparison. Error bars indicate SEM; *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$; ****, $p \leq 0.0001$. Final tumor weight was measured at endpoint and is represented in bar graphs (right panels). Statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD; *, $p \leq 0.05$. **(E)** Protein lysates were prepared from 22Rv1 tumors treated with Vehicle or GCN2iB and analyzed by immunoblot for phosphorylated GCN2-T899, ATF4, SLC7A5 (LAT1), SLC7A11 (xCT), SLC3A2 (4F2), and actin. The levels of the SLC proteins normalized to actin are shown. Phosphorylated GCN2-T899 was normalized to total GCN2. Statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD (N = 4); *, $p \leq 0.05$. **(F)** Amino acid measurements of 22Rv1 tumors treated with vehicle or GCN2iB. Bar graphs show high abundance (top) and low abundance (bottom) amino acids. Statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD (N = 4); #, $p \leq 0.1$; *, $p \leq 0.05$; **, $p \leq 0.01$. **(G)** Pearson correlation between p-GCN2-T899 and SLC3A2 (4F2) histoscores calculated from IHC staining from a prostate tumor microarray (Biomax PR807c) containing normal (N = 10), hyperplasia (N = 20) and malignant (N = 50) for all tissues (combined) or Gleason scores 4 and 5. The center lines depict linear regression (95% confidence intervals). Not all samples were analyzed due to damaged/quality of tissue samples. Levels of p-GCN2-T899 and SLC3A2 (4F2) were measured by IHC staining and QuPath was used to determine the histoscore. Two representative cases are shown for high (Case 1) and low (Case 2) p-GCN2-T899 and SLC3A2 (4F2) staining. Scale bar indicates 200

µm (main image) and 20 µm (insert). **(H)** Correlation of expression of SLC3A2 (4F2) and a GCN2-dependent gene signature in prostate adenocarcinoma (PRAD, N = 551) from the Cancer Genome Atlas (TCGA). The GCN2-dependepent gene signature was derived from RNA-seq data as described in the materials and methods.

Data and Materials Availability Statement

The authors declare that all data generated or analyzed in this study are included in the published article and its supplementary information, or are available upon request from the corresponding authors. The CHARGE-seq and RNA-seq datasets generated in this study have been deposited in the NCBI Gene Expression Omnibus (GEO) database under the ascension codes GSE196251 and GSE196252, respectively. Gene expression data from prostate cancer patients (PRAD) in the TCGA database used for correlation analysis is publicly available. Plasmids and cell lines are available upon reasonable request for academic research purposes.

Acknowledgements

This work was supported in part by grants from NIH GM136331 (R.C.W.), DK109714 (T.G.A., R.C.W.), the NCI R21CA221942 (R.P.), the Indiana University Melvin and Bren Simon Comprehensive Cancer Center P30CA082709 (K.A.S.), and the Lilly Endowment, Inc., through its support for the Indiana University Pervasive Technology Institute. We thank the IU Simon Comprehensive Cancer Center for use of the Tissue Procurement and Distribution Core, which provided core needle biopsy specimens for staining, the Preclinical Modeling and Therapeutics Core (PMTC) for assistance with flow cytometry and cell cycle analysis, and the Histology Lab Service Core at the Indiana University School of Medicine for embedding and sectioning of tissue samples. Finally, the authors acknowledge Ludovica Ceci for assistance with slide scanning, Kadir Isidan for assistance with histology, and members of Wek and Pili laboratories for helpful discussions.

Competing Interests

R.C.W. is a member of the advisory board and holds equity in HiberCell, Inc. K.A.S. and T.G.A are consultants for HiberCell, Inc. and K.A.S. receives research support from HiberCell, Inc.

Author Contributions

R.A.C., R.P., R.C.W, and K.A.S. conceived of and designed the experiments, led the project, and coordinated the writing of the manuscript with the assistance of all of the co-authors. N.P.D. provided technical advice and assisted with *in vivo* experiments. A.J.K. and A.E. provided technical assistance with cell culture experiments and immunoblot analyses. H.-G.K. provided technical advice and assistance with IHC staining. L.C. provided pathology interpretation of prostate cancer specimens. E.T.M., M.M., and T.G.A. assisted with amino acid analysis. J.M. carried out the CHARGE-seq and tRNA charging analysis, and P.H.A. and K.C. provided bioinformatics support for the RNA-seq and CHARGE-seq studies.

References

1. Siegel, R.L., K.D. Miller, and A. Jemal, *Cancer statistics, 2019.* CA Cancer J Clin, 2019. **69**(1): p. 7-34.
2. Gann, P.H., *Risk factors for prostate cancer.* Rev Urol, 2002. **4 Suppl 5**: p. S3-S10.
3. Klusa, D., et al., *Metastatic Spread in Prostate Cancer Patients Influencing Radiotherapy Response.* Front Oncol, 2020. **10**: p. 627379.
4. Heinlein, C.A. and C. Chang, *Androgen receptor in prostate cancer.* Endocr Rev, 2004. **25**(2): p. 276-308.
5. Huggins, C. and C.V. Hodges, *Studies on prostatic cancer: I. The effect of castration, of estrogen and of androgen injection on serum phosphatases in metastatic carcinoma of the prostate. 1941.* J Urol, 2002. **168**(1): p. 9-12.
6. Polotti, C.F., et al., *Androgen deprivation therapy for the treatment of prostate cancer: a focus on pharmacokinetics.* Expert Opin Drug Metab Toxicol, 2017. **13**(12): p. 1265-1273.
7. Virgo, K.S., et al., *Second-Line Hormonal Therapy for Men With Chemotherapy-Naive, Castration-Resistant Prostate Cancer: American Society of Clinical Oncology Provisional Clinical Opinion.* J Clin Oncol, 2017. **35**(17): p. 1952-1964.
8. Watson, P.A., V.K. Arora, and C.L. Sawyers, *Emerging mechanisms of resistance to androgen receptor inhibitors in prostate cancer.* Nat Rev Cancer, 2015. **15**(12): p. 701-11.
9. Crawford, E.D., et al., *Androgen-targeted therapy in men with prostate cancer: evolving practice and future considerations.* Prostate Cancer Prostatic Dis, 2019. **22**(1): p. 24-38.
10. Bubley, G.J. and S.P. Balk, *Association Between Androgen Receptor Splice Variants and Prostate Cancer Resistance to Abiraterone and Enzalutamide.* J Clin Oncol, 2017. **35**(19): p. 2103-2105.
11. Baird, T.D. and R.C. Wek, *Eukaryotic initiation factor 2 phosphorylation and translational control in metabolism.* Adv Nutr, 2012. **3**(3): p. 307-21.

12.   Tian, X., et al., *Targeting the Integrated Stress Response in Cancer Therapy.* Front Pharmacol, 2021. **12**: p. 747837.

13.   Dong, J., et al., *Uncharged tRNA activates GCN2 by displacing the protein kinase moiety from a bipartite tRNA-binding domain.* Mol Cell, 2000. **6**(2): p. 269-79.

14.   Inglis, A.J., et al., *Activation of GCN2 by the ribosomal P-stalk.* Proc Natl Acad Sci U S A, 2019. **116**(11): p. 4946-4954.

15.   Wek, S.A., S. Zhu, and R.C. Wek, *The histidyl-tRNA synthetase-related sequence in the eIF-2 alpha protein kinase GCN2 interacts with tRNA and is required for activation in response to starvation for different amino acids.* Mol Cell Biol, 1995. **15**(8): p. 4497-506.

16.   Walter, P. and D. Ron, *The unfolded protein response: from stress pathway to homeostatic regulation.* Science, 2011. **334**(6059): p. 1081-6.

17.   Wek, R.C. and D.R. Cavener, *Translational control and the unfolded protein response.* Antioxid Redox Signal, 2007. **9**(12): p. 2357-71.

18.   Chukwurah, E., et al., *A tale of two proteins: PACT and PKR and their roles in inflammation.* FEBS J, 2021. **288**(22): p. 6365-6391.

19.   Garcia, M.A., et al., *Impact of protein kinase PKR in cell biology: from antiviral to antiproliferative action.* Microbiol Mol Biol Rev, 2006. **70**(4): p. 1032-60.

20.   Fessler, E., et al., *A pathway coordinated by DELE1 relays mitochondrial stress to the cytosol.* Nature, 2020. **579**(7799): p. 433-437.

21.   Guo, X., et al., *Mitochondrial stress is relayed to the cytosol by an OMA1-DELE1-HRI pathway.* Nature, 2020. **579**(7799): p. 427-432.

22.   Suragani, R.N., et al., *Heme-regulated eIF2alpha kinase activated Atf4 signaling pathway in oxidative stress and erythropoiesis.* Blood, 2012. **119**(22): p. 5276-84.

23.   Staschke, K.A. and R.C. Wek, *Adapting to cell stress from inside and out.* Nat Cell Biol, 2019. **21**(7): p. 799-800.

24.   Tameire, F., et al., *ATF4 couples MYC-dependent translational activity to bioenergetic demands during tumour progression.* Nat Cell Biol, 2019. **21**(7): p. 889-899.

25.   Nguyen, H.G., et al., *Development of a stress response therapy targeting aggressive prostate cancer.* Sci Transl Med, 2018. **10**(439).

26.   Harding, H.P., Y. Zhang, and D. Ron, *Protein translation and folding are coupled by an endoplasmic-reticulum-resident kinase.* Nature, 1999. **397**(6716): p. 271-4.

27.   Young, S.K. and R.C. Wek, *Upstream Open Reading Frames Differentially Regulate Gene-specific Translation in the Integrated Stress Response.* J Biol Chem, 2016. **291**(33): p. 16927-35.

28.   Hinnebusch, A.G., I.P. Ivanov, and N. Sonenberg, *Translational control by 5'-untranslated regions of eukaryotic mRNAs.* Science, 2016. **352**(6292): p. 1413-6.

29.   Kuruma, H., et al., *A novel antiandrogen, Compound 30, suppresses castration-resistant and MDV3100-resistant prostate cancer growth in vitro and in vivo.* Mol Cancer Ther, 2013. **12**(5): p. 567-76.

30.   DeSantis, C.E., et al., *Cancer statistics for adults aged 85 years and older, 2019.* CA Cancer J Clin, 2019. **69**(6): p. 452-467.

31.   Li, Y., et al., *Androgen receptor splice variants mediate enzalutamide resistance in castration-resistant prostate cancer cell lines.* Cancer Res, 2013. **73**(2): p. 483-9.

32.   Sramkoski, R.M., et al., *A new human prostate carcinoma cell line, 22Rv1.* In Vitro Cell Dev Biol Anim, 1999. **35**(7): p. 403-9.

33.   Tilley, W.D., et al., *Androgen receptor gene expression in human prostate carcinoma cell lines.* Cancer Res, 1990. **50**(17): p. 5382-6.

34.   Nakamura, A., et al., *Inhibition of GCN2 sensitizes ASNS-low cancer cells to asparaginase by disrupting the amino acid response.* Proc Natl Acad Sci U S A, 2018. **115**(33): p. E7776-E7785.

35.   Lehman, S.L., et al., *Translational Upregulation of an Individual p21Cip1 Transcript Variant by GCN2 Regulates Cell Proliferation and Survival under Nutrient Stress.* PLoS Genet, 2015. **11**(6): p. e1005212.

36.   Leung-Pineda, V., et al., *Induction of p21 and p27 expression by amino acid deprivation of HepG2 human hepatoma cells involves mRNA stabilization.* Biochem J, 2004. **379**(Pt 1): p. 79-88.

37.   Battu, S., et al., *Amino Acid Sensing via General Control Nonderepressible-2 Kinase and Immunological Programming.* Front Immunol, 2017. **8**: p. 1719.

38.   Pavlova, N.N., et al., *Translation in amino-acid-poor environments is limited by tRNA(Gln) charging.* Elife, 2020. **9**.

39.   Girardi, E., et al., *A widespread role for SLC transmembrane transporters in resistance to cytotoxic drugs.* Nat Chem Biol, 2020. **16**(4): p. 469-478.

40.   Pizzagalli, M.D., A. Bensimon, and G. Superti-Furga, *A guide to plasma membrane solute carrier proteins.* FEBS J, 2021. **288**(9): p. 2784-2835.

41.   Boyce, M., et al., *A selective inhibitor of eIF2alpha dephosphorylation protects cells from ER stress.* Science, 2005. **307**(5711): p. 935-9.

42.   Novoa, I., et al., *Feedback inhibition of the unfolded protein response by GADD34-mediated dephosphorylation of eIF2alpha.* J Cell Biol, 2001. **153**(5): p. 1011-22.

43.   Keller, T.L., et al., *Halofuginone and other febrifugine derivatives inhibit prolyl-tRNA synthetase.* Nat Chem Biol, 2012. **8**(3): p. 311-7.

44.   Corsetti, G., et al., *Dietary supplementation with essential amino acids boosts the beneficial effects of rosuvastatin on mouse kidney.* Amino Acids, 2014. **46**(9): p. 2189-203.

45.   Sekhar, K.R., et al., *Radiosensitization by enzalutamide for human prostate cancer is mediated through the DNA damage repair pathway.* PLoS One, 2019. **14**(4): p. e0214670.

46.   Corey, E., et al., *LuCaP 35: a new model of prostate cancer progression to androgen independence.* Prostate, 2003. **55**(4): p. 239-46.

47.   Maimaiti, M., et al., *The heavy chain of 4F2 antigen promote prostate cancer progression via SKP-2.* Sci Rep, 2021. **11**(1): p. 11478.

48.   Lieu, E.L., et al., *Amino acids in cancer.* Exp Mol Med, 2020. **52**(1): p. 15-30.

49.   Broer, S., *Amino Acid Transporters as Targets for Cancer Therapy: Why, Where, When, and How.* Int J Mol Sci, 2020. **21**(17).

50.   Bhutia, Y.D., et al., *Amino Acid transporters in cancer and their relevance to "glutamine addiction": novel targets for the design of a new class of anticancer drugs.* Cancer Res, 2015. **75**(9): p. 1782-8.

51. Pathria, G. and Z.A. Ronai, *Harnessing the Co-vulnerabilities of Amino Acid-Restricted Cancers.* Cell Metab, 2021. **33**(1): p. 9-20.

52. Butler, M., L.T. van der Meer, and F.N. van Leeuwen, *Amino Acid Depletion Therapies: Starving Cancer Cells to Death.* Trends Endocrinol Metab, 2021. **32**(6): p. 367-381.

53. Wu, C.C., et al., *Ribosome Collisions Trigger General Stress Responses to Regulate Cell Fate.* Cell, 2020. **182**(2): p. 404-416 e14.

54. Zaborske, J.M., et al., *Genome-wide analysis of tRNA charging and activation of the eIF2 kinase Gcn2p.* J Biol Chem, 2009. **284**(37): p. 25254-67.

55. Wek, R.C., B.M. Jackson, and A.G. Hinnebusch, *Juxtaposition of domains homologous to protein kinases and histidyl-tRNA synthetases in GCN2 protein suggests a mechanism for coupling GCN4 expression to amino acid availability.* Proc Natl Acad Sci U S A, 1989. **86**(12): p. 4579-83.

56. Furnish, M., et al., *MIRO2 Regulates Prostate Cancer Cell Growth via GCN1-Dependent Stress Signaling.* Mol Cancer Res, 2022.

57. Mick, E., et al., *Distinct mitochondrial defects trigger the integrated stress response depending on the metabolic state of the cell.* Elife, 2020. **9**.

58. Scalise, M., et al., *The Human SLC7A5 (LAT1): The Intriguing Histidine/Large Neutral Amino Acid Transporter and Its Relevance to Human Health.* Front Chem, 2018. **6**: p. 243.

59. Son, D.O., H. Satsu, and M. Shimizu, *Histidine inhibits oxidative stress- and TNF-alpha-induced interleukin-8 secretion in intestinal epithelial cells.* FEBS Lett, 2005. **579**(21): p. 4671-7.

60. Kanarek, N., et al., *Histidine catabolism is a major determinant of methotrexate sensitivity.* Nature, 2018. **559**(7715): p. 632-636.

61. Wortel, I.M.N., et al., *Surviving Stress: Modulation of ATF4-Mediated Stress Responses in Normal and Malignant Cells.* Trends Endocrinol Metab, 2017. **28**(11): p. 794-806.

62. Pallmann, N., et al., *Regulation of the unfolded protein response through ATF4 and FAM129A in prostate cancer.* Oncogene, 2019. **38**(35): p. 6301-6318.

63. Schmidt, E.K., et al., *SUnSET, a nonradioactive method to monitor protein synthesis.* Nat Methods, 2009. **6**(4): p. 275-7.

64. Nikonorova, I.A., et al., *Time-resolved analysis of amino acid stress identifies eIF2 phosphorylation as necessary to inhibit mTORC1 activity in liver.* J Biol Chem, 2018. **293**(14): p. 5005-5015.

65. Prinsen, H., et al., *Rapid quantification of underivatized amino acids in plasma by hydrophilic interaction liquid chromatography (HILIC) coupled with tandem mass-spectrometry.* J Inherit Metab Dis, 2016. **39**(5): p. 651-660.

66. Griffin, C., et al., *Rapid quantification of underivatized alloisoleucine and argininosuccinate using mixed-mode chromatography with tandem mass spectrometry.* J Chromatogr B Analyt Technol Biomed Life Sci, 2019. **1128**: p. 121786.

67. Spahn, P.N., et al., *PinAPL-Py: A comprehensive web-application for the analysis of CRISPR/Cas9 screens.* Sci Rep, 2017. **7**(1): p. 15854.

68. Martin, M., *Cutadapt removes adapter sequences from high-throughput sequencing reads.* 2011, 2011. **17**(1): p. 3.

69. Andrews, S., *FastQC: A quality control tool for high throughput sequence data.* https://www.bioinformatics.babraham.ac.uk/projects/fastqc/, 2010.

70. Krueger, F., *Trim Galore: A wrapper tool around Cutadapt and FastQC to consistently apply quality and adapter trimming to FastQ files.* https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/, 2012.

71. Kim, D., B. Langmead, and S.L. Salzberg, *HISAT: a fast spliced aligner with low memory requirements.* Nat Methods, 2015. **12**(4): p. 357-60.

72. Liao, Y., G.K. Smyth, and W. Shi, *featureCounts: an efficient general purpose program for assigning sequence reads to genomic features.* Bioinformatics, 2014. **30**(7): p. 923-30.

73. Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550.

74. Mootha, V.K., et al., *PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes.* Nat Genet, 2003. **34**(3): p. 267-73.

75. Subramanian, A., et al., *Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles.* Proc Natl Acad Sci U S A, 2005. **102**(43): p. 15545-50.

76. Liberzon, A., et al., *The Molecular Signatures Database (MSigDB) hallmark gene set collection.* Cell Syst, 2015. **1**(6): p. 417-425.

77. Liberzon, A., et al., *Molecular signatures database (MSigDB) 3.0.* Bioinformatics, 2011. **27**(12): p. 1739-40.

78. Misra, J., et al., *Discordant regulation of eIF2 kinase GCN2 and mTORC1 during nutrient stress.* Nucleic Acids Res, 2021. **49**(10): p. 5726-5742.

79. van Eenige, R., et al., *RandoMice, a novel, user-friendly randomization tool in animal research.* PLoS One, 2020. **15**(8): p. e0237096.

80. Montgomery, R.B., et al., *Maintenance of intratumoral androgens in metastatic prostate cancer: a mechanism for castration-resistant tumor growth.* Cancer Res, 2008. **68**(11): p. 4447-54.

81. Langmead, B. and S.L. Salzberg, *Fast gapped-read alignment with Bowtie 2.* Nat Methods, 2012. **9**(4): p. 357-9.

Supplementary Methods.

CHARGE-seq analyses. RNA prepared from each sample was treated with 12.5 mM NaIO$_4$ in sodium acetate solution (pH 4.5) in the dark at room temperature for 20 minutes followed by quenching with 0.3 M glucose. All mature tRNAs end with 3'-CCA and NaIO$_4$ will only oxidize the 3'-A residue of uncharged tRNAs since the 3'-ends of charged tRNAs are protected by linked amino acids. The resulting total RNA was desalted by passing through MicroSpin G-50 column (GE Healthcare, Cat. #27533001) and the tRNA was deacylated through β-elimination in 50 mM Tris–HCl (pH 9.0) at 37 °C for 45 minutes followed by quenching with da solution of 50 mM sodium acetate buffer (pH 4.5) and 100 mM NaCl. Total RNA was then recovered by overnight EtOH precipitation at -20°C. The RNA samples were then treated with polynucleotide kinase (New England BioLabs, Cat. #M0201) to remove the 3'-phosphate group from the uncharged tRNA followed by ligation to 5'-adenylated uniquely barcoded adapters using RNA ligase 2 truncated KQ (New England BioLabs, Cat. # M0351L). The resulting tRNAs were then ligated to a 5'-adaptor, annealed to an RT primer, and converted to cDNA using a using a SuperScript IV RT kit (Invitrogen, Cat. #18090050). The tRNA library was PCR amplified for 10 cycles using Illumina multiplex and barcode PCR primers using xxx polymerase Platinum SuperFi II DNA Polymerase (ThermoFisher Scientific, Cat #12369002). All oligonucleotide primers and adapters utilized in the procedure are listed in **Supplementary Table 3**. The resulting PCR products were electrophoresed on an 8% non-denaturing polyacrylamide TBE gel (Invitrogen, Cat #EC6215) and ~210 bp bands were excised, purified, and subjected to Illumina HiSeq paired-end sequencing. Fastq files from the pooled samples were demultiplexed according to the five-nucleotide sample barcode contained within the ligated adapter using a custom python script, and read pairs without an identifiable barcode were discarded. Illumina sequencing adapters and ligated sample barcode adapters were trimmed using cutadapt [68] in

paired read mode and reads with less <20 nucleotides remaining after trimming, corresponding to empty adapters, were removed.

A custom reference file for human tRNA was constructed from the gtRNAdb GRCh38 high confidence tRNA gene set by removing duplicate sequences and appending 'CCA' to the 3' end of each unique tRNA gene sequence. Trimmed reads were then aligned to the custom tRNA reference file using bowtie2 [81] short read aligner with options "--dovetail -D 20 -R 3 -N 1 -L 20 -i S,1,0.50". The charging status of the aligned tRNA was determined from the 3' end of the aligned portion of each read pair, where reads ending in 'CCA' were determined to be charged and reads ending in 'CC' were determined to be uncharged. This method was able to determine the charging status of ~86% percent of all mapped reads. Counts of charged and uncharged tRNA for each tRNA gene in each sample were then imported to R for further analysis. Differences in read counts between samples were accounted for by multiplying the read counts in each sample by a normalization coefficient equal to the average number of reads across all samples divided by the read count in that sample.