GCN2 eIF2 kinase promotes prostate cancer by maintaining amino acid homeostasis

Ricardo A. Cordova[1,2], Jagannath Misra[1], Parth H. Amin[1], Angela J. Klunk[1], Nur P. Damayanti[2,3], Kenneth R. Carlson[1], Andrew J. Elmendorf[1], Hyeong-Geug Kim[1], Emily T. Mirek[4], Bennet D. Elzey[5,6], Marcus J. Miller[7], X. Charlie Dong[1], Liang Cheng[2,6,8,9], Tracy G. Anthony[4], Roberto Pili[10,#], Ronald C. Wek[1,2,#], and Kirk A. Staschke[1,2,#]

[1]Department of Biochemistry and Molecular Biology, Indiana University School of Medicine, Indianapolis, IN 46202, USA.

[2]Indiana University Melvin and Bren Simon Comprehensive Cancer Center, Indianapolis, IN 46202, USA.

[3]Department of Neurological Surgery, Indiana University School of Medicine, Indianapolis, IN 46202, USA.

[4] Department of Nutritional Sciences, Rutgers University, New Brunswick, NJ 08901, USA.

[5]Department of Comparative Pathology, Purdue University, West Lafayette, IN 47907, USA

[6]Department of Urology, Indiana University School of Medicine, Indianapolis, IN 46202, USA

[7]Department of Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, IN 46202, USA

[8]Department of Pathology and Laboratory Medicine, Indiana University School of Medicine, Indianapolis, IN 46202, USA

[9]Present address: Department of Pathology and Laboratory Medicine, Warren Alpert Medical School of Brown University, Providence, RI 02903, USA

[10]Jacobs School of Medicine and Biomedical Sciences, University at Buffalo, Buffalo, NY 14203, USA

#To whom correspondence should be addressed:

Kirk A. Staschke, Department of Biochemistry and Molecular Biology, 635 Barnhill Drive,

Indiana University School of Medicine, Indianapolis, Indiana 46202-5122; E-mail:

kastasch@iu.edu

Ronald C. Wek, Department of Biochemistry and Molecular Biology, 635 Barnhill Drive, Indiana

University School of Medicine, Indianapolis, Indiana 46202-5122; E-mail: rwek@iu.edu.

Roberto Pili, Department of Medicine, 875 Ellicott Street, University at Buffalo, Buffalo, NY

14203; E-mail: rpili@buffalo.edu.

Running title: GCN2 controls tumor amino acid homeostasis

Keywords: ISR; prostate cancer, GCN2; eIF2; amino acid transport; SLC; solute carrier; CD98; 4F2; 4F2hc; SLC3A2

Conflict of interest statement

R.C.W. is a member of the advisory board and holds equity in HiberCell, Inc. K.A.S. and T.G.A are consultants for HiberCell, Inc. and K.A.S. receives research support from HiberCell, Inc.

Abstract word count: 157
Word Count: 5885
Figures and tables: 7
Supplementary Information:
- Supplementary Figures, 22 total
- Supplementary Tables S1 – S4
- Supplementary Files S1 – S4

Abstract

A stress adaptation pathway termed the integrated stress response has been suggested to be active in many cancers including prostate cancer (PCa). Here, we demonstrate that the eIF2 kinase GCN2 is required for sustained growth in androgen-sensitive and castration-resistant models of PCa both in vitro and in vivo, and is active in PCa patient samples. Using RNA-seq transcriptome analysis and a CRISPR-based phenotypic screen, GCN2 was shown to regulate expression of over 60 solute-carrier (*SLC*) genes, including those involved in amino acid transport and loss of GCN2 function reduces amino acid import and levels. Addition of essential amino acids or expression of 4F2 (SLC3A2) partially restored growth following loss of GCN2, suggesting that GCN2 targeting of SLC transporters is required for amino acid homeostasis needed to sustain tumor growth. A small molecule inhibitor of GCN2 showed robust in vivo efficacy in androgen-sensitive and castration-resistant mouse models of PCa, supporting its therapeutic potential for the treatment of PCa.

Significance: The ISR kinase GCN2 is critical for maintaining tumor amino acid levels to facilitate growth, suggesting a novel therapeutic strategy for the treatment of prostate cancer by inducing starvation for essential amino acids.

Introduction

Prostate cancer (PCa) accounts for 20% of all newly diagnosed cancers and is the second leading cause of cancer-related deaths in men in the United States [1]. Many factors contribute to prostate cancer, including genetics, diet, age, and race [2]. The pathological grade of the PCa and whether it has disseminated into surrounding tissues is prognostic for risk and treatment efficacy [3]. Androgen receptor (AR) is considered the primary oncogenic driver for PCa [4, 5], and therapies targeting AR activity are current standard of care, especially for those

with metastatic PCa [6-9]. Unfortunately, patients develop resistance with time and fail therapy

[8, 10]. In addition, it remains unclear how additional pathways drive PCa through AR-

dependent and AR-independent mechanisms. Therefore, the identification of novel targets for

therapeutic intervention are critically needed and clinically relevant.

The Integrated Stress Response (ISR) is critical for cell adaptation and survival to

environmental stresses. Central to the ISR is phosphorylation of the α subunit of eIF2 (p-eIF2α),

which triggers reductions in global protein synthesis to conserve energy and nutrients, and

concomitantly facilitates adaptive gene expression [11, 12]. Four eIF2α kinases respond to

distinct stress conditions: The eIF2α kinase GCN2 (EIF2AK4) is induced by amino acid

starvation and other conditions that trigger ribosomal collisions or stalling [13-15]. PERK

(EIF2AK3) is responsive to disruptions in the endoplasmic reticulum [16, 17]. PKR (EIF2AK2) is

activated by viral infection and inflammation [18, 19], and HRI (EIF2AK1) is enhanced by

restricted heme or iron supply, oxidative stress, and disruptions in mitochondrial function [20-

22]. Activation of these ISR regulators may also occur in response to oncogenic drivers, such as

MYC amplification and loss of the PTEN tumor suppressor, frequently observed in prostate

tumors. Therefore, both extrinsic and intrinsic cell stresses can invoke the ISR in tumors and

pharmacological inhibition of translational control is reported to trigger cytotoxicity in different

cancer models [23-25].

To reprogram gene expression towards recovery from stress damage, the ISR also

induces preferential translation of specific ISR gene transcripts, such as activating transcription

factor 4 (ATF4), a transcriptional regulator of genes involved in amino acid metabolism and

proteostasis control, oxidative stress defenses, along with feedback control of the ISR [26-28].

Therefore, p-eIF2α induces both translational and transcriptional modes of gene expression to

achieve stress adaptation. Targeted inhibition of ATF4 impedes progression of MYC-driven

cancers, emphasizing the important role of this ISR effector in tumor progression [23-25].

Here, we dissect the role of eIF2α kinase(s) induced in PCa and the mechanisms by

which the ISR serves to promote growth of this cancer. Using androgen-sensitive and

castration-resistant human PCa cell lines, primary human PCa tissues, and mouse xenograft

models, we show that the ISR regulator GCN2 is activated in PCa. GCN2 induces expression of

key ISR effector genes involved in nutrient transport that are critical for appropriate maintenance

of free amino acid pools. Genetic or pharmacological inhibition of GCN2 dysregulates

expression of these transporters and inhibits PCa growth. These results point to GCN2 as a

potential therapeutic target for PCa.

Results

GCN2 activity supports prostate cancer growth.

To determine the role of individual eIF2α kinases and their importance for proliferation of

PCa cells in culture, androgen-sensitive LNCaP cells cultured in androgen replete media

conditions were individually depleted of HRI, PKR, PERK, or GCN2 using target-specific

siRNAs. There was robust reduction of protein levels of each of the eIF2α kinases; however,

only cells depleted for GCN2 or PKR showed a significant decrease in proliferation (**Figures**

1A,1B, Figure 1 - Figure Supplement 1A, and Supplementary File 1). Whereas GCN2

depletion lowered p-eIF2α levels and ATF4 expression, knockdown of PKR, HRI, or PERK had

no effect on p-eIF2α and ATF4 protein levels (**Figure 1B and Figure 1 - Figure Supplement**

1A). These results suggest that LNCaP cell proliferation requires GCN2 as an ISR regulator.

To broaden our investigation to other PCa cell lines, we knocked down GCN2 in the

androgen-sensitive line LAPC-4 and in the castration-resistant cell lines C4-2B, MR49F, 22Rv1,

and PC-3. MR49F cells express AR, but are resistant to enzalutamide [29], 22Rv1 cells express

the ARv7 splice variant and are also resistant to enzalutamide [30-32], PC-3 is AR null [33]. In

each of these PCa cell lines, depletion of GCN2 reduced proliferation and lowered ATF4

expression (**Figure 1C, Figure 1 - Figure Supplement 1B, and Supplementary File 1**). In

22Rv1 and PC-3 cells, we were also able to achieve a deletion of *EIF2AK4* (GCN2) using

CRISPR/Cas9, which reduced p-eIF2α and ATF4 levels (**Figure 1 - Figure Supplement 2A**). In

the case of 22Rv1, we determined a similar slow growth phenotype as observed with siRNA

knockdown of GCN2 (**Figure 1 - Figure Supplement 2B and Supplementary File 1**).

Reintroduction of GCN2 into the GCN2 KO cells rescued the slow growth phenotype, and

rescued expression of ATF4 and its target gene ASNS (**Figure 1 - Figure Supplement 2B,**

Figure 1 – Figure Supplement 2C, and Supplementary File 1). The small molecule GCN2iB

is a selective inhibitor of GCN2 eIF2α kinase activity [34]. Using the inhibitory concentration

range of 500 nM to 10 μM of GCN2iB, we determined that pharmacological inhibition of GCN2

also caused significant growth inhibition of LNCaP cells, and similarly reduced the growth of C4-

2B, 22Rv1, and PC-3 cells **(Figure 1D, Figure 1 - Figure Supplement 3A, and**

Supplementary File 1). Treatment of GCN2 KO cells with up to 10 μM GCN2iB had no effect

on growth demonstrating the specificity of this inhibitor (**Figure 1 - Figure Supplement 3B** and

Supplementary File 1). As expected, treatment of the LNCaP, 22Rv1, C4-2B, or PC-3 cells

with GCN2iB lowered activation of GCN2 as measured by autophosphorylation of the activation

loop (p-GCN2-T899), reduced p-eIF2α levels, and lowered expression of ATF4, and its target

genes ASNS and TRIB3. Importantly, treatment with GCN2iB did not alter protein levels of AR

(**Figure 1E and Figure 1 - Figure Supplement 3C**). To further test the importance of the ISR in

PCa, we selectively depleted ATF4 using siRNA in LNCaP, MR49F, C4-2B, and PC-3 cells and

observed similar growth inhibition for each cell line (**Figure 1 - Figure Supplement 1B, Figure**

1 - Figure Supplement 4, and Supplementary File 1). Importantly, GCN2 is not active in the

non-cancerous prostate epithelial cell line BPH-1 [35], and loss of GCN2 or ATF4, or

pharmacological inhibition of GCN2, did not reduce p-eIF2α or ATF4 levels, or significantly

impact the growth of these cells (**Figure 1 - Figure Supplement 5A, Figure 1 – Figure**

Supplement 5B, Figure 1 – Figure Supplement 5C, and **Supplementary File 1**). These

results support the hypothesis that the eIF2α kinase activity of GCN2 and the ISR support PCa

proliferation.

We next sought to determine whether GCN2 is activated in human malignant PCa

tissues. Using immunohistochemistry and prostate tumor microarrays (TMAs) featuring normal

and non-malignant prostate tissue (N = 33) and malignant tumor tissues (N = 88), we

determined that there was enhanced GCN2 activity in the tumor samples compared to normal

tissues as measured by p-GCN2-T899 staining (**Figure 1F**). In addition, increased p-GCN2

(Thr-899) staining was also observed in prostate tumors from patient core needle biopsies as

compared to adjacent non-malignant tissue (**Figure 1 - Figure Supplement 6**). Collectively,

these results indicate that GCN2 is active basally in both androgen-sensitive and castration-

resistant PCa models without appreciable extrinsic stress treatments, confers growth

advantages, and is activated in prostate cancer patient tissues.

GCN2 induces ISR transcriptome featuring expression of amino acid transporters.

To address the mechanisms by which GCN2 confers growth advantages in PCa, we

cultured LNCaP cells in the presence or absence of GCN2iB and then measured changes in the

transcriptome by RNA-seq. With 24 hours of GCN2iB treatment, 239 transcripts showed a

significant ≥ 2-fold increase, whereas 374 transcripts were decreased ≥ 2-fold (fdr ≤ 0.05)

compared to vehicle-treated cells (**Figure 2A, Supplementary Table 1**). We carried out a gene

set enrichment analysis (GSEA) of the differentially-expressed transcripts to distinguish the

molecular functions that were altered by GCN2 inhibition. Included among the functional gene

groups showing significant dependence on GCN2 for expression were those linked to amino

acid deprivation and transport (**Figures 2B and 2C)**. These genes are representative of ISR-

targeted gene groups and include genes regulated by ATF4 such as *DDIT3 (CHOP), ASNS*,

and *TRIB3*, those involved in glycine-serine biosynthesis (*PHGDH, PSAT1, PSPH*, and

SHMT2), aminoacyl-tRNA synthetases, and a large number of SLC transporters, many of which

function in the transport of amino acids (**Figure 2B and Supplementary File 2**). As illustrated in

the heat map, expression of the *SLC* genes were significantly lowered after 6 hours of GCN2iB

treatment, with further reductions following 24 hours (**Figure 2C**). It is noteworthy that of the 61

known *SLC* genes involved in the transport of amino acids, 27 were measurably expressed in

LNCaP cells, and 20 of these expressed transcripts were significantly decreased following

treatment with GCN2iB (**Figure 2B and Supplementary File 2**).

Lowered expression of SLC proteins upon GCN2 inhibition was further validated by

immunoblot analyses. xCT (SLC7A11), CAT-1 (SLC7A1), ASCT1 (SLC1A4), ASCT2 (SLC1A5),

and 4F2 (SLC3A2) protein levels were sharply lowered with GCN2iB treatment of LNCaP cells,

along with reduced expression of ATF4 and its target gene ASNS (**Figure 2D and Figure 2 -**

Figure Supplement 1A). Similarly, treatment of C4-2B, 22Rv1, and PC-3 cells with GCN2iB

also reduced LAT-1 (SLC7A5), xCT (SLC7A11), and 4F2 (SLC3A2) (**Figure 1 - Figure**

Supplement 3C). We also measured transporter proteins in 22Rv1 and PC-3 cells and showed

that deletion of *EIF2AK4* (GCN2) also diminished protein levels of xCT (SLC7A11), LAT-1

(SLC7A5) (LAT1), and 4F2 (SLC3A2) (**Figure 2E, Figure 2 - Figure Supplement 2B)**.

Furthermore, the reintroduction of GCN2 fully restored expression of ATF4 and ASNS and

partially restored expression of all three SLC proteins in 22Rv1 cells (**Figure 2E**). To more

directly address the effects of GCN2 inhibition on amino acid transport, we measured uptake of

leucine, lysine, and arginine in LNCaP and 22Rv1 cells. Treatment of LNCaP cells with GCN2iB

or deletion of GCN2 in 22Rv1 cells significantly lowered the uptake of each of these amino acids

(**Figure 2F and 2G**), suggesting that GCN2-directed gene expression is important for uptake of

amino acids in PCa cells.

GCN2 is critical for maintenance of free amino acid pools.

Given that GCN2 is important for expression of *SLC* genes and amino acid uptake, we

next addressed whether the intracellular levels of free amino acids are disrupted upon inhibition

of GCN2. LNCaP cells were cultured in the presence or absence of GCN2iB for 8 hours and

free amino acid levels were measured. With pharmacological inhibition of GCN2, there was a

sharp reduction in the amounts of essential amino acids (EAA), including histidine, the

branched-chain amino acids, threonine, phenylalanine, and methionine, and nonessential amino

acids (NEAA), such as glutamate and glutamine, aspartate, asparagine, alanine, and tyrosine

(**Figure 3A)**. We next considered whether supplementing the culture medium with additional

amino acids would alleviate the growth defect that occurs with the inhibition of GCN2. While

GCN2iB treatment of LNCaP cells cultured in standard medium markedly reduced growth,

supplementation of additional EAAs into the medium reversed this proliferation defect and

increased free amino acid pools in treated cells (**Figure 3B, Figure 3 - Figure Supplement 1,**

Figure 3 - Figure Supplement 2, and **Supplementary File 1**). By contrast, supplementation

with additional NEAA did not suppress the LNCaP growth defect triggered by GCN2iB treatment

(**Figure 3 - Figure Supplement 2** and **Supplementary File 1**).

To address the effect of GCN2 inhibition and EAA supplementation on the cell cycle in

cultured PCa cells, LNCaP cells were treated with GCN2iB or GCN2 expression was reduced

with siRNA. Both methods of GCN2 depletion caused an increase in the G1 phase of the cell

cycle concomitant with a reduction in S phase, indicative of checkpoint control of the cell cycle

that can occur with amino acid depletion [36, 37] (**Figure 3C and Figure 3 - Figure**

Supplement 3A). Supplementation of LNCaP cells with EAA reversed the accumulation of cells

in G1 phase elicited by GCN2iB (**Figure 3C**). These results suggest that inhibition of GCN2 can

cause cell cycle arrest, which was recapitulated with siRNA knockdown of GCN2 or ATF4

(**Figure 3 - Figure Supplement 3A**). Similar findings were observed in 22Rv1 cells deleted for

GCN2 or in other PCa cell lines when GCN2 or ATF4 levels were reduced with siRNA (**Figure 3**

- Figure Supplement 3A and **Figure 3 – Figure Supplement 3B**).

Accumulating uncharged tRNA occurs upon severe depletion of cognate amino acids

and is a primary signal activating GCN2 [38]. We cultured LNCaP cells in the presence or

absence of GCN2iB and measured tRNA charging genome-wide using CHARGE-seq [39].

There was a reduction of aminoacylated tRNA^{His} upon GCN2iB treatment that was restored

when additional EAA were supplemented to the growth medium (**Figures 3D, 3E, and**

Supplementary File 3). These results were further supported by measurements of tRNA^{His}

charging levels by qRT-PCR. Addition of EAA to cultured LNCaP cells significantly increased

tRNA^{His} charging levels while treatment with GCN2iB dramatically reduced tRNA^{His} charging

levels which was reversed by the addition of EAA (**Figure 3 - Figure Supplement 4A**). These

results suggest that histidine depletion and the accompanying accumulation of uncharged

tRNA^{His} is a key signal for activation of GCN2. Upon treatment with GCN2iB, which lowered p-

eIF2α and blocked appropriate induction of the ISR, there was severe depletion of histidine

accompanied by accumulation of uncharged tRNA^{His}, culminating in a slow-growth phenotype.

To further address the role of histidine availability in the slow growth phenotype

associated with GCN2 inhibition, we cultured GCN2iB-treated LNCaP cells in medium with

added supplements of individual amino acids. In alignment with our charged tRNA^{His}

measurements, only the addition of histidine to the culture medium reversed the growth

inhibition triggered by GCN2iB (**Figure 3F** and **Figure 3 - Figure Supplement 2**). Removal of

histidine from the growth medium increased G1 accumulation similar to GCN2iB treatment,

while supplementation of additional histidine to the medium alleviated the cell cycle arrest

triggered by GCN2iB treatment (**Figure 3G**). Finally, supplementation with the full complement

of EAA or histidine alone significantly lowered p-GCN2, p-eIF2α, and ATF4 expression,

although it is noted that histidine supplementation rendered only partial suppression of the ISR

compared to addition of EAA (**Figure 3H**). Accompanying this reduction in activation of GCN2,

supplementation with histidine sharply increased aminoacylation of tRNA^{His} (**Figure 3E**) and

there was a trend towards enhanced global protein synthesis as measured by puromycin

incorporation. It is curious that there was a trend towards decreased bulk translation upon

GCN2iB treatment that was restored with the addition of histidine (**Figure 3 - Figure**

Supplement 4B). As noted above, GCN2 inhibition for 48 hours adversely affected the cell

cycle and amino acid levels, which together are suggested to reduce bulk translation upon

GCN2iB treatment despite lowered p-eIF2α levels.

We next determined whether EAA supplementation in other PCa cell lines also

suppressed the slow growth phenotype triggered by loss of GCN2 activity. Addition of EAA or

histidine alone also suppressed the slow growth phenotype of MR49F cells treated with GCN2iB

(**Figure 3 - Figure Supplement 5A**). However, in 22Rv1 cells, EAA or lysine alleviated the slow

growth associated with deletion of *EIF2AK4* (GCN2) (**Figure 3 - Figure Supplement 5B**).

These growth supplementation studies suggest that while GCN2 is indispensable for

maintaining EAA in PCa cells, the precise processes by which GCN2 carries out these functions

can vary among PCa cell lines presumably due to differences in their unique genetic features

and cell-line specific buffering mechanisms accompanying specific gene expression and

metabolic programs. Depletion of GCN2 in LNCaP and MR49F cells renders them vulnerable to

histidine depletion, whereas deletion of GCN2 sensitizes 22Rv1 cells to lysine depletion.

Select GCN2-regulated SLC transporters are critical for PCa fitness.

Given that GCN2 has a primary role for inducing expression of *SLC* genes, we sought to

determine the contributions of these transporters for PCa growth and viability. To evaluate their

contributions in an unbiased manner, we utilized a previously reported CRISPR/Cas9 library

targeting 394 human SLC genes and pseudogenes [40]. Each of the SLC genes was targeted

with 6 specific single guide RNAs (sgRNAs), along with negative controls for target specificity

and positive controls for assessing cell fitness. The sgRNA library was stably expressed in

LNCaP and 22Rv1 cells using lentivirus delivery. Cells were then subjected to puromycin

selection and then cultured for 3 - 4 weeks (15 - 20 population doublings) as previously reported

[40]. Genomic DNA was prepared and encoded sgRNA sequences were amplified by PCR,

sequenced, and their composition was compared to the starting cells with the sgRNA library at

Day 0. Depletion of the cells expressing specific sgRNAs were then evaluated for statistical

significance, and cell fitness was scored and ranked. A collection of SLC genes showed

significantly diminished growth, including several that require GCN2 for expression, such as

LAT1 (SLC7A5), SLC25A36, SLC25A46, SLC27A1, SLC35A2, and 4F2 (SLC3A2) (**Figures 4A**

and **Supplementary File 4**). *SLC27A1* and *4F2* (*SLC3A2)* were required for efficient growth of

both LNCaP and 22Rv1 cells (**Figure 4B** and **Figure 4 - Figure Supplement 1**). These results

suggest that PCa cells require a specific collection of SLC genes for growth and a subset of

these are dependent on GCN2 for full expression.

240   Given the importance of 4F2 (SLC3A2) in the transport of bulk amino acids [41], we next

focused on the GCN2-dependent regulation of this transport regulator. Treatment of LNCaP

cells with the GCN2iB for 6 or 24 hours resulted in a reduction in GCN2 activation, as shown by

p-GCN2, and lowered ATF4 mRNA and protein. Levels of 4F2 (SLC3A2) protein and mRNA

were also lowered with inhibition of GCN2, although this reduction occurred after 24 hours of

treatment (**Figure 4C** and **Figure 4 - Figure Supplement 2A**). Consistent with our findings that

histidine limitation in part drives GCN2 activation in LNCaP cells, addition of histidine to the

growth media of LNCaP cells in culture reduced GCN2 activation and lowered expression of

ATF4 and 4F2 (SLC3A2) protein and mRNA, while removing histidine from the media further

increased GCN2 activation and expression of ATF4 and 4F2 (SLC3A2) (**Figure 4D** and **Figure**

4 - Figure Supplement 2B). Salubrinal has been reported to increase p-eIF2α by inhibiting its

PP1-directed dephosphorylation [42]. Supporting the idea that p-eIF2α is critical for 4F2

(SLC3A2) expression, addition of salubrinal alone increased 4F2 (SLC3A2) expression and also

restored 4F2 (SLC3A2) expression in cells treated with GCN2iB (**Figure 4 - Figure**

Supplement 3A). In addition, overexpression of the PP1-targeting protein GADD34 [43],

reduced phosphorylation of eIF2α and expression of ATF4 and 4F2 (SLC3A2) (**Figure 4 -**

Figure Supplement 3B). Further supporting the central role of activation of GCN2 in the

induction of 4F2 (SLC3A2) expression, treatment of LNCaP cells with halofuginone, an inhibitor

of EPRS and prolyl-tRNA charging [44] resulted in robust activation of GCN2 and increased

expression of ATF4 and 4F2 (SLC3A2) protein and mRNA (**Figure 4E** and **Figure 4 - Figure**

Supplement 2C). These results support the model that activation of GCN2 during nutrient

stress is a major inducer of 4F2 (SLC3A2) expression.

We next addressed whether 4F2 (SLC3A2) expression is linked to the growth phenotype

following loss of GCN2. First, knock-down of 4F2 (SLC3A2) expression using siRNA

significantly reduced growth in LNCaP and 22Rv1 cells and led to activation of GCN2 and

increased ATF4 protein levels (**Figure 4F, 4G,** and **Supplementary File 1**). Importantly, knock-

down of 4F2 (SLC3A2) expression did not impact growth of non-cancerous BPH-1 cells (**Figure**

1 – Figure Supplement 5B and **Figure 1 – Figure Supplement 5C**). As was shown earlier,

loss of GCN2 function using siRNA or GCN2iB treatment in cultured LNCaP cells reduced

growth (**Figure 1A** and **D**). This growth defect was rescued by overexpression of 4F2 (SLC3A2)

(**Figure 4H** and **Supplementary File 1)**. Importantly, overexpression of 4F2 (SLC3A2) reduced

GCN2 activity, which lowered p-GCN2 and p-eIF2α levels, along with ATF4 protein expression,

and provided a growth advantage in both wild-type LNCaP and 22Rv1 cells (**Figure 4 - Figure**

Supplement 3C and **Figure 4 – Figure Supplement 3D**). We next tested the impact of 4F2

(SLC3A2) knock-down on intracellular amino acid levels. As anticipated, knockdown of SLC3A2

reduced a subset of amino acids as compared to knockdown of GCN2 (**Figure 4I**). These

results suggest that expression of *SLC* genes, especially 4F2 (SLC3A2), are important for PCa

growth. As illustrated in the model presented in **Figure 5**, GCN2 plays a critical role ensuring

that appropriate expression of key *SLC* genes and associated amino acid transport are tightly

linked to the ISR**.**

GCN2 supports PCa tumor growth and maintains expression of amino acid transporters

in mouse xenograft models.

To examine the contributions of the GCN2/SLC transporter model for PCa growth in an

in vivo model, we first selected the 22Rv1 cell line as it is castration-resistant. Parental (WT)

22Rv1 cells with functional GCN2 or those with the eIF2α kinase deleted (GCN2 KO) were

injected subcutaneously into the flanks of 8-week-old male NSG mice, and tumor volumes were

measured for up to 12 days. Whereas the parental cells showed sustained tumor growth, loss of

GCN2 significantly reduced tumor growth over time, resulting in a significant reduction in tumor

weight at the end of the study (**Figure 6A**). IHC staining of tumor preparations showed

diminished expression of Ki-67, a marker for cell proliferation in 22Rv1 tumors devoid of GCN2

(**Figure 6 - Figure Supplement 1A**). As expected, the GCN2 protein was not detectable in the

GCN2 KO tumors as judged by immunoblot, p-GCN2-T899 staining was reduced, and

expression levels of the GCN2-targeted transporters 4F2 (SLC3A2), xCT (SLC7A11), and LAT1

(SLC7A5) were sharply reduced (**Figure 6B** and **Figure 6 - Figure Supplement 1A**).

Furthermore, the 22Rv1 tumors maintained expression of the ARv7 variant, which provides for

castration-resistance and resistance to enzalutamide (**Figure 6B**). Similar results were obtained

in the AR-null castration-resistant cell line PC-3 following deletion of GCN2 (**Figure 6C, 6D,**

Figure 6 - Figure Supplement 1B, and **Figure 6 – Figure Supplement 1C**). Together, these

xenograft experiments using the 22Rv1 and PC-3 models support the hypothesis that GCN2

provides significant growth advantages to PCa and suggests a mechanism involving enhanced

SLC transporter expression that is independent of AR expression.

To determine the importance of ATF4 in tumor growth *in vivo*, we carried out a similar

xenograft study using 22Rv1 cells deleted for ATF4. While loss of GCN2 led to dramatically

reduced tumor growth, deletion of ATF4 only delayed tumor growth relative to the wild-type

parental cells (**Figure 6E**). Furthermore, deletion of GCN2 led to a further decrease in 4F2

(SLC3A2), ASCT2 (SLC1A5), and CAT1 (SLC7A1) protein levels as compared to loss of ATF4

(**Figure 6 - Figure Supplement 2A**). It is noted that loss of GCN2 did not reduce p-eIF2α levels

in these end stage tumors, suggesting activation of a secondary eIF2 kinase or reduced

phosphatase-directed dephosphorylation of eIF2α, which has been reported earlier in cells

during extended nutrient limitation or other stress conditions [45, 46]. Interestingly, deletion of

ATF4 resulted in increased activation of GCN2, increased levels of p-eIF2α, and in some cases

increased amino acids relative to GCN2 KO tumors (**Figure 6F** and **Figure 6 - Figure**

Supplement 2A), indicating that loss of ATF4 can exacerbate cell stress facilitating further

activation of GCN2. Consistent with these *in vivo* findings, knock-down of GCN2 expression in

LNCaP cells in culture reduced amino acid levels to a greater degree as compared to knock-

down of ATF4 (**Figure 6 - Figure Supplement 2B**).

Another key feature of our GCN2/SLC transporter model is that this ISR regulator

functions to ensure appropriate uptake of select EAA. To more directly test this model *in vivo*,

we again utilized the xenograft model featuring 22Rv1 GCN2 KO cells. In this experiment, we

provided additional EAA in the drinking water for the tumor bearing mice. While 22Rv1 cells with

functional GCN2 showed tumor growth within 20 days of injection, loss of GCN2 reduced tumor

growth even over the course of 40 days. However, consumption of EAA-supplemented water by

GCN2 KO-tumor bearing mice provided for increased tumor growth approaching that of wild-

type GCN2 tumors, albeit there was a time delay of 20 days for 22Rv1 with wild-type GCN2 to

about 40 days with EAA-supplemented GCN2 KO (**Figure 6G**). Supplementation of the EAA did

not affect tumor growth of parental 22Rv1 with functional GCN2 (**Figure 6G)**. Similar to previous

reports, supplementation of EAA in the drinking water resulted in a slight decrease in body

weight of mice bearing either parental or GCN2 KO tumors [47] (**Figure 6 - Figure Supplement**

3A). Measurements of free amino acids in tumors indicated that loss of GCN2 reduced

intracellular arginine and lysine concentrations, with histidine and the branched-chain amino

acids trending lower. Dietary supplementation of EAA partially restored these amino acid levels,

although only histidine achieved statistical significance (**Figure 6H** and **Figure 6 - Figure**

Supplement 3B). Immunoblot measurements showed that loss of GCN2 lowered key ISR

markers, including the key SLC transporters independent of the supplementation of EAA

(**Figure 6 - Figure Supplement 3C**). We also tested the impact of overexpression of 4F2

(SLC3A2) on growth of 22Rv1 tumors. Similar to supplementation with EAA, overexpression of

4F2 (SLC3A2) in GCN2 KO tumors partially rescued growth and restored tumor amino acid

levels (**Figure 6I, Figure 6 - Figure Supplement 4A,** and **Figure 6 – Figure Supplement 4B**).

These results further support the notion that GCN2 has a primary function in maintaining EAA

and highlight a key role for GCN2 in the regulation of 4F2 (SLC3A2) expression. In keeping with

the cultured PCa cell line experiments, GCN2 played a vital role supporting amino acid

homeostasis in these tumor models.

Pharmacological inhibition of GCN2 reduces tumor growth in xenograft models.

We next tested the impact of pharmacological inhibition of GCN2 function in cancer cell-

derived and patient-derived xenograft models. Male NSG mice were injected subcutaneously

with LNCaP or 22Rv1 cells or implanted with tumor fragments from an androgen-sensitive tumor

TM00298 [48]. Alternatively, male castrated NSG mice were implanted with tumor fragments

from LuCaP-35 castration-resistant tumors [49]. Tumors were allowed to grow to 50 -100 mm^3

and mice were randomized to vehicle or GCN2iB treatment groups. Treatment with GCN2iB at

30 mg/kg twice daily for 2 - 6 weeks resulted in robust anti-tumor activity in all four models as

shown by a significant delay in the tumor volume, reductions in tumor mass at endpoint, and

reduced expression of p-GCN2-T899 and Ki-67 (**Figure 7A – 7E and Figure 7 - Figure**

Supplement 1A). Importantly, GCN2iB had no effect on the body weight of treated mice over

the course of treatment (**Figure 7 - Figure Supplement 1B)**, lowered expression of 4F2

(SLC3A2) in both 22Rv1 and TM00298 tumors, and reduced levels of select amino acids in

22Rv1 tumors from treated mice (**Figure 7E, 7F, Figure 7 - Figure Supplement 2A,** and

Figure 7 – Figure Supplement 2B). These results suggest that GCN2 is important for tumor

growth and expression of 4F2 (SLC3A2) in both androgen-sensitive and castration-resistant

tumors.

Our results suggest that GCN2 controls expression of multiple amino acid transporters to

regulate amino acid homeostasis in PCa tumors. We next addressed whether expression of p-

GCN2 or GCN2-dependent mRNA transcripts correlated with expression of 4F2 (SLC3A2) in

human tumor samples. Again, using a tumor tissue array containing patient samples, we stained

for p-GCN2-T899 and 4F2 (SLC3A2). Similar to recent findings [50], 4F2 (SLC3A2) staining was

increased in malignant tissue samples as compared to normal tissue (**Figure 7 - Figure**

Supplement 3A). Interestingly, SLC3A2 (4F2) showed a moderate, albeit significant,

correlation with expression of p-GCN2-T899 (r = 0.27, p = 0.025). This correlation was more

striking among tissues from higher grade tumors (r = 0.47, p = 0.002, Gleason Score 4 and 5)

(**Figure 7G**), which was further supported by staining of p-GCN2 and 4F2 (SLC3A2) of core

needle biopsy specimens from patients with high grade prostate cancers (**Figure 7 - Figure**

Supplement 3B). To further test the GCN2/SLC3A2 (4F2) model, we derived a gene signature

from GCN2-dependent transcripts from our RNA-seq analysis and measured a significant

correlation with SLC3A2 (4F2) expression (r = 0.53, p < 0.0001) using a cohort of prostate

cancer patients (PRAD) from the TCGA database (**Figure 7H**). These results further support the

idea that pharmacological inhibition of GCN2 may serve as a potent strategy for the treatment of

androgen-sensitive and castration-resistant PCa by limiting the expression of amino acid

transporters including 4F2 (SLC3A2).

Discussion

Tumor cells often demonstrate high demand for amino acids to sustain their increased

metabolism, protein synthesis and proliferation [51], and frequently overexpress amino acid

transporters to meet this increased demand [52, 53]. Altered or increased transport of essential

and non-essential amino acids has been suggested to be a point of vulnerability in several

cancers [54, 55]. In this study, we show that the adaptive ISR pathway driven by GCN2 is active

in PCa tissue and can be genetically and pharmacologically targeted in human models of

androgen-sensitive and castration-resistance disease. Loss of GCN2 function by genetic

deletion or pharmacological inhibition sharply reduced PCa growth *in vitro* and reduces tumor

growth in xenograft models. As illustrated in **Figure 5**, GCN2-directs the expression of a large

number of SLC transporter genes, many of which are critical for PCa growth. Among these

GCN2-targeted genes is 4F2 (SLC3A2), a type II integral membrane protein which is known to

form dimers with SLC7 family members including SLC7A5 (LAT1) and SLC7A11 (xCT), each

contributing to uptake of specific amino acids [41]. Loss of GCN2 in PCa reduced free amino

acids both *in vitro* and in tumors derived from xenograft models. Addition of EAAs to the

medium of cultured cells rescued the growth defect associated with GCN2 inhibition;

furthermore, supplementation of EAAs in the drinking water of mice injected subcutaneously

with GCN2 KO PCa cells substantially enhanced tumor growth *in vivo*. While several strategies

to deprive cancer cells of nutrients have been contemplated [54], our results suggest that

inhibition of GCN2 represents a unique treatment paradigm to starve prostate tumors of EAAs.

Further supporting the utility of GCN2 as a therapeutic target for PCa, a small molecule inhibitor

of GCN2 significantly slowed tumor growth in both CDX and PDX models of androgen-sensitive

and castration-resistant disease irrespective of AR-status.

GCN2 is activated by uncharged tRNAs that accumulates during amino acid starvation

or by ribosome collisions [15, 56-58]. As indicated by p-GCN2, this eIF2α kinase is induced in a

variety of different proliferating PCa cell lines and in malignant prostate tissue. While our results

suggest that amino acid limitations and accompanying uncharged tRNA contribute to GCN2

activation in PCa, it is noted that other mechanisms can contribute to regulation of GCN2. For

example, a mitochondrial Rho GTPase 2 (MIRO2) overexpressed in PCa was recently shown to

bind to GCN1, a critical activator of GCN2 [59], and certain mitochondrial defects were reported

to contribute to activation of GCN2 and the ISR [60]. Given that these different PCa models

represent those that are androgen-sensitive, castration-resistant, resistant to enzalutamide, or

lack AR expression altogether, our study suggests that GCN2 functions to control PCa growth

independent of AR function. Supplementation of EAAs reduces p-GCN2 and partially rescued

the growth deficiency of GCN2 KO cells, suggesting that deficiency of at least some of these

critical amino acids are limiting in proliferating PCa cells. This nutrient limitation would trigger

GCN2 induction of the ISR which features induced expression of SLC transporters, including the

central 4F2 (SLC3A2), and these events are required to sustain tumor growth. It is noteworthy

that 4F2 (SLC3A2) has recently been shown to be overexpressed in PCa and correlated with

reduced progression-free survival [50].

420   Experiments involving supplementation of individual amino acids to cultured LNCaP cells

knocked down for GCN2 suggest that this eIF2α kinase is critical for maintenance of histidine

homeostasis. Histidine is an essential amino acid whose transport is not clearly understood;

although, LAT1 (SLC7A5) has been suggested to facilitate its uptake [61]. Beyond its critical

role in protein synthesis, histidine also functions as an important antioxidant [62] and its

catabolism has been linked to nucleotide synthesis in some cancers [63]. Additional studies will

be required to delineate how histidine contributes to PCa metabolism and growth. Whereas

histidine also rescued the growth defect of GCN2 inhibition in the related MR49F cells, only

lysine alleviated the slow growth of cultured 22Rv1 GCN2 KO cells. These in vitro experiments

suggest a more complex picture where the limiting amino acid is cell-type specific or influenced

by specific genetic alterations which might influence the threshold or mechanism of GCN2

activation. It is acknowledged that amino acids included in the media used in the in vitro

experiments are typically present at high physiological levels so the importance of any given

EAA may be masked by the abundance of these nutrients in the culture medium. Consistent

with this notion, the analysis of 22Rv1 GCN2 KO tumors showed multiple EAAs were reduced.

435   Activation of GCN2 triggers the ISR by the induction of ATF4, a key transcriptional

activator. ATF4 contributes to PCa growth induced by GCN2, as depletion of ATF4 in multiple

PCa cell lines led to growth inhibition in culture. Previous studies have suggested that ATF4 is

important for growth in multiple cancers [24, 64] including PCa [65]. However, under our

experimental conditions, whereas loss of GCN2 led to a sharp reduction in tumor growth in the

22Rv1 xenograft model, deletion of ATF4 only delayed tumor growth. These results suggest that

while ATF4-directed transcriptional activation is important for PCa proliferation, GCN2 functions

by additional mechanisms, including preferential translation of multiple ISR-target genes.

As noted above activated GCN2 induces a program of genes that are important for

maintenance of amino acid homeostasis. It should also be noted that another major nutrient

sensing pathway is regulated by mTORC1 [66] and mTORC1 activity is transiently reduced in

GCN2iB treated cells (data not shown). We focused on the critical role of GCN2 in controlling

expression of 4F2 (SLC3A2), albeit a collection of SLC genes required GCN2 for full

expression. While 4F2 (SLC3A2) is an important contributor to amino acid homeostasis,

deletion of 4F2 (SLC3A2) in LNCaP cells led to depletion of only a portion of the free amino

acids that were disrupted by loss of GCN2, consistent with its requirement with a subset of

amino acid transporters [41]. Furthermore, overexpression of 4F2 (SLC3A2) only partially

restored growth of 22Rv1 GCN2 KO tumors *in vivo* and did not affect the progression of

parental 22Rv1 tumors. These results indicate that the central role of GCN2 for ensuring

appropriate amino acid homeostasis and enabling PCa proliferation involves induction of

multiple SLC transporters. In addition, the reduction in ATF4 expression that accompanies loss

of GCN2 function reduces expression of several amino acid biosynthetic genes which would

exacerbate the observed effects on uptake of NEAAs. Taken together, these observations

suggest that inhibition of GCN2 may be a more effective therapeutic strategy for starving PCa of

amino acids as compared to therapies that target individual SLC transporters. In summary, our

results indicate that GCN2 may represent a novel therapeutic target for prostate cancer. The

therapeutic landscape for prostate cancer patients is rapidly evolving but AR still remains the

focus. Unfortunately, resistance to androgen deprivation therapies continue to be a major

challenge. The development of new therapeutic strategies that work independently from

inhibition of the AR axis is highly clinically relevant and based on our findings, inhibitors of

GCN2 should be considered further as part of a strategy for treatment of androgen-sensitive

and castration-resistant prostate cancers.

Materials and Methods

Cell culture and treatments. LNCaP clone FGC (Cat. #CRL-1740, RRID:CVCL_1379), C4-2B

(Cat. #CRL-3315, RRID:CVCL_4784), CW22Rv1 (referred to as 22Rv1, Cat. #CRL-2505,

RRID:CVCL_1045), PC-3 (Cat. #CRL-1435, RRID: CVCL_0035), and HEK-293T (Cat. #CRL-

3216, RRID:CVCL_0063) cells were purchased from the American Tissue Type Collection

(ATCC, Manassas, VA). BPH-1 (Cat. #SCC256, RRID: CVCL_1091) cells were purchased from

MilliporeSigma (Temecula, CA). LAPC-4 (RRID:CVCL_4744) cells were provided by Dr.

Dominic J. Smiraglia (Roswell Park Comprehensive Cancer Center, Buffalo, NY). MR49F

(RRID:CVCL_RW53) cells were obtained from Dr. Timothy Ratliff (Purdue University, West

Lafayette, IN). LNCaP, C4-2B, 22Rv1, and BPH-1 cells were maintained in phenol-free RPMI-

1640 (Gibco, Cat #11835-030) supplemented with 10% FBS (HyClone, Cat #SH30071). LAPC-

4 cells were maintained in phenol-free RPMI-1640 supplemented with 10% FBS as described

above and 10 nM dihydrotestosterone (Sigma-Aldrich, Cat. #D-073). MR49F cells were

maintained in phenol-free RPMI-1640 supplemented with 10% FBS as described above and 10

µM Enzalutamide (MedChemExpress, Cat. #HY-70002). PC-3 cells were maintained in Ham's

F-12K (Kaighn's) media (Gibco, Cat. #21127-022) or Human Plasma-Like Media (HPLM, Gibco

Cat. #A4899101) supplemented with 10% FBS. HEK-293T cells were maintained in DMEM

(Corning, Cat. #10-013-CV) supplemented with 10% FBS. All cell lines were maintained at 37°C

in a fully humidified atmosphere containing 5% CO_2, used at early passage for experiments, and

tested to be free of mycoplasma contamination. For cell treatments, cells were plated at

250,000 cells per well in 6-well plates in standard culture media as described above and

allowed to attach overnight. Media was changed to standard culture media containing 500 nM to

10 µM GCN2iB or DMSO as a control or standard media lacking histidine (MyBiosource, Cat

#MBS652918) and incubated for the indicated time period. GCN2iB [34] was synthesized at

Sun-shinechem (Wuhan, China). As indicated, salubrinal (Tocris Biosciences, Cat #405060-95-

9) was added to the culture media at a concentration of 50 μM. To measure bulk protein

translation, 1 μM puromycin (MP Biomedicals Cat #194539) was included in the culture medium

for the last 15 minutes of treatment with GCN2iB, and protein lysates were probed and

quantified by immunoblot analyses using anti-puromycin antibody [67].

Immunoblot analysis. Whole cell extracts and tumor lysates were prepared in 1% SDS lysis

buffer supplemented with 1X Halt Protease and Phosphatase Inhibitor Cocktail (Thermo

Scientific, Cat. #1861281). Tumors were homogenized using a Kinematica Polytron tissue

homogenizer. Lysates were boiled for 10 minutes, sonicated, and centrifuged at 20,000 x g to

remove insoluble material. Protein concentrations were determined by DC Protein Assay (Bio-

Rad, Cat. #5000112) in technical triplicate using BSA as the protein standard. Equal amounts of

protein samples, which ranged from 10 - 20 μg between experiments, were separated on SDS-

PAGE gels, transferred to nitrocellulose membranes, and used for immunoblot analysis. The

primary antibodies used were as follows: phospho-GCN2-T899 (Abcam Cat. #ab75836,

RRID:AB_1310260), total GCN2 (Cell Signaling Technology Cat. #3302, RRID:AB_2277617),

total PERK (Cell Signaling Technology Cat. #3192, RRID:AB_2095847), total HRI (Santa Cruz

Biotechnology Cat. #sc-365239, RRID:AB_10843794), total PKR (Cell Signaling Technology

Cat. #12297, RRID:AB_2665515), phospho-eIF2α-S51 (Abcam Cat. #ab32157,

RRID:AB_732117), total eIF2α (Cell Signaling Technology Cat. #5324, RRID:AB_10692650),

ATF4 (Cell Signaling Technology Cat. #11815, RRID:AB_2616025) or custom rabbit polyclonal

antibody which was prepared against full-length recombinant human ATF4 protein and affinity

purified, ASNS (Cell Signaling Technology, Cat. #20843S), TRIB3 (Abcam Cat. #ab75846,

RRID:AB_1310768), GADD34 (Proteintech Cat. #10449-1-AP, RRID:AB_2168724),

SLC7A5/LAT1 (Cell Signaling Technology Cat. #5347, RRID:AB_10695104), SLC7A11/xCT

(Cell Signaling Technology Cat. #12691, RRID:AB_2687474), SLC3A2/4F2 (Cell Signaling

Technology Cat. #47213, RRID:AB_2799323), SCL7A1/CAT1 (Proteintech Cat. #14195-1-AP,

RRID:AB_2190723), SLC1A4/ASCT1 (Cell Signaling Technology Cat. #8442,

RRID:AB_10828382), SLC1A5/ASCT2 (Cell Signaling Technology Cat. #5345,

RRID:AB_10621427), AR (Cell Signaling Technology Cat. #5153, RRID:AB_10691711),

puromycin (Millipore Cat. #MABE343, RRID:AB_2566826), β-actin (Sigma-Aldrich Cat. #A5441,

RRID:AB_476744), β-tubulin (Cell Signaling Technology Cat. #2146, RRID:AB_2210545).

Immunoblot signals were visualized by enhanced chemiluminescence (ECL) using Clarity

Western ECL Substrate (Bio-Rad, Cat. #170-5060) or SuperSignal West Femto Maximum

Sensitivity Substrate (ThermoFisher Scientific, Cat. #34094) and images were captured using a

ChemiDoc™ Imaging System (Bio-Rad, RRID:SCR_019037). Prestained protein standards

(Bio-Rad Cat. #1610377) were included on each immunoblot, captured as a separate merged

image file, and are indicated in each immunoblot panel.

RNA Interference and proliferation assays. Small Interfering RNAs (siRNAs) were purchased

from Dharmacon and sequences are shown in **Supplementary Table 2**. The siRNAs that

showed the most robust depletion of the targeted genes in the indicated cells were selected for

experiments. For growth assays LAPC-4, C4-2B, MR49F, 22Rv1, or PC-3 cells were seeded

into 96-well plates at 2500 – 5000 cells per well in normal growth media as described above and

allowed to attach overnight. Cells were transfected with 1 pmole of siRNA using Lipofectamine

RNAiMAX Transfection Reagent (Thermo Scientific, Cat. #13778030) or DharmaFECT 3

Transfection Reagent (Horizon, Cat. #T-2003-01) according to manufacturer's protocol and

incubated overnight (considered Day 0). Alternatively, where indicated for LAPC-4 cells, growth

media was changed to serum-free Accell Delivery Media (Horizon, Cat. #B-005000) and

transfected with 1 pmole of Accell siRNA and incubated 48 hours at which time cells were

changed back to normal growth media (considered Day 0). Cells were incubated for an

additional 2 – 6 days and cell growth was quantified using CellTiter-Glo® 2.0 Reagent

(Promega, Cat. #924C). For immunoblot analysis, PCa cells were seeded into 6-well plates at

250,000 – 350,000 cells per well and transfected with 30 pmoles of siRNA as described above

and incubated for 48 hours prior to lysate preparation as described above.

***In vitro* amino acid supplementation.** For amino acid supplementation experiments, 1X MEM

Amino Acids Solution (Gibco, Cat. #11130051) was used for essential amino acid (EAA) mix

and 1X MEM Non-Essential Amino Acids Solution (Gibco, Cat. #11140050) was used for non-

essential amino acid (NEAA) mix. Experiments utilizing single amino acids were performed

using the following concentrations: 600 µM L-Arginine hydrochloride, 100 µM L-Cystine

dihydrochloride, 200 µM L-Histidine hydrochloride, 400 µM L-Isoleucine, 400 µM L-Leucine, 395

µM L-Lysine hydrochloride, 100 µM L-Methionine, 200 µM L-Phenylalanine, 400 µM L-

Threonine, 50 µM L-Tryptophan, 200 µM L-Tyrosine, and 400 µM L-Valine. All amino acids were

dissolved in water and were purchased from Sigma-Aldrich (St. Louis, MO).

Amino acid uptake measurements. LNCaP, parental 22Rv1, or 22Rv1 GCN2 KO cells were

plated in 24-well plates (50,000 cells) in standard culture media as described above. LNCaP

cells were treated with vehicle (DMSO) or 2 µM GCN2iB for 24 hours. Cells were washed with

pre-warmed sodium-free Hank's Balanced Salt Solution (HBSS) and treated with 100 nM ^{3}H-

Arginine (5 µCi/ml, Perkin-Elmer), 1 µM ^{14}C-Leucine (0.3 µCi/ml, Perkin-Elmer), or 1 µM ^{14}C-

Lysine (0.3 µCi/ml, Perkin-Elmer) in HBSS containing the appropriate un-labelled amino acid

(100-200 µM Arginine, 1-20 µM Leucine, or 1 µM Lysine) at 25 °C for 2 minutes. Transport was

stopped by adding the appropriate excess cold-amino acid (non-radioactive) in HBSS. Cells

were washed with PBS, lysed with Passive lysis buffer (Promega, Cat #E1941), and a portion

added to scintillation cocktail and counted. The protein concentration of each lysate was

determined as described above and amino acid uptake was represented as nmol/µg of total

protein/min. Each measurement was performed in triplicate.

Plasmid construction. Plasmid pDONR221-SLC3A2_STOP was a gift from RESOLUTE

Consortium & Giulio Superti-Furga (Addgene plasmid # 161379, RRID:Addgene_161379) [40].

Plasmid pENTR223-1 GCN2 (NM_001013703) was purchased from Transomic Technologies

and pLX303 was a gift from David Root (Addgene plasmid # 25897, RRID:Addgene_25897).

GCN2 coding sequence was subcloned into the lentiviral vector pLVX-IRES-Hygromycin

(Clontech, Cat #632185) using an In-Fusion HD Cloning plus kit (TaKaRa, Cat. #638920)

according to manufacturer's protocol yielding plasmids pLVX-GCN2-hygro. The SLC3A2 (4F2)

coding region was subcloned into the lentiviral vector pLX303 by Gateway cloning using

Gateway LR Clonase II Enzyme Mix (Invitrogen, Cat. #11791-020) according to manufacturer's

protocol yielding plasmid pLX303-4F2-blast. Plasmid pMSCV-GADD34-puro contains the

human PPP1R15A (GADD34, NM_014330.5) gene subcloned into the NcoI/EcoRI sites of the

retroviral vector pMSCV-puro (Clontech Laboratories, Mountain View, CA).

Construction of cell lines. GCN2 and ATF4 knock-out cell lines were generated using

CRISPR/Cas9 Human Gene Knockout Kits (Origene, Cat. #KN412459 and #KN402333) using

hGCN2g1 (5'-AATTTAGTTTTGTACCCTCA-3') and hATF4g1 (5'-

CTTCCTGAGCAGCGAGGTGT-3'); respectively, following the manufacturer's protocol. Cell

lines overexpressing GCN2 or SLC3A2 (4F2) were created by lentivirus transduction. Briefly,

HEK-293T cells were cultured in DMEM as described above and were transfected with lentivirus

constructs and ViraPower™ Lentiviral Packaging Mix (Invitrogen, Cat #35-1275) using

Lipofectamine 2000 (ThermoFisher Scientific, Cat # 11668030) according to manufacturer's

protocol and virus containing supernatant was collected over 48 – 72 hours, concentrated using

Lenti-X™ Concentrator (TaKaRa, Cat. #631232), and stored in aliquots at -80 °C. 22Rv1 or PC-

3 cells were transduced with virus-containing culture media supplemented with 10 µg/ml

Polybrene (Millipore, Cat #TR-1003-G) for 24 hours and replaced with fresh growth media. At 72

hours post-transfection, growth media was changed to media supplemented with 250 µg/ml

Hygromycin B (Thermo Cat #10687010) or 10 µg/ml blastocidin S (Sigma, Cat #SBR00022) to

select for stable pools as appropriate. The appropriate empty vector was used as a control for

all experiments.

qRT-PCR analysis. Total RNA was isolated from cultured cells or tumor tissue using TRIzol™

LS Reagent (Invitrogen, Cat. #10296010) and quantified using a Nanodrop Spectrophotometer

(ThermoFisher Scientific). Total RNA (1 µg) was converted to cDNA using a High-Capacity

cDNA Reverse Transcription Kit (ThermoFisher Scientific, Cat. #4368813) according to the

manufacturer's protocol. Following cDNA synthesis, samples were diluted five-fold in molecular

biology grade water and PCR amplification was carried out using PowerUp SYBR Green Master

Mix (ThermoFisher Scientific, Cat. #A25742) using an Applied Biosystems QuantSudio5 PCR

system. The primers used for measuring *ATF4* and *SLC3A2 (4F2hc)* transcript levels are shown

in **Supplementary Table 3**. The relative abundance of each transcript was calculated using the

$\Delta\Delta CT$ method with GAPDH serving as an internal control and data are presented normalized to

each control group.

Cell cycle analysis. Cell cycle analysis on LNCaP, MR49F, C4-2B, 22Rv1, and PC-3 cells was

performed by co-staining with a Click-iT™ EdU Alexa Fluor™ 488 Flow Cytometry Assay Kit

(Invitrogen, Cat. #C10425) and FxCycle Violet Ready Flow Reagent (Invitrogen, Cat. # R37166)

according to manufacturer's protocol. Cell treatments and siRNA knockdowns were carried out

for 48 hours as described above before analysis using an LSR II (LSR4) flow cytometer (BD

Biosystems). A minimum of 10,000 events were analyzed per condition and all experiments

were performed using triplicate wells. Data analysis was performed using FlowJo (FlowJo LLC,

TreeStar) and data are presented as the percentage of cells in G1, S, or G2-M phases of the

cell cycle.

Amino acid analysis. Levels of different amino acids were measured in cultured cells or tumor

lysates by HPLC or LC-MS. Cell pellets or tissues were resuspended in 0.1% formic acid in

methanol, vortexed briefly, and subjected to three cycles of freezing in liquid nitrogen followed

by thawing at 42 °C for 10 minutes. Clarified lysates were analyzed by reverse-phase liquid

chromatography with appropriate standard curves as described previously [68] or using liquid

chromatography tandem mass spectrometry (LC-MS/MS) with stable isotope dilution. For the

MS-based method, lysates were spiked with a mixture containing 37 amino acid isotopic internal

standards and deproteinated using a solution of 90% acetonitrile, 10 mM ammonium formate,

and 0.15% formic acid. Clarified lysates were analyzed using a biphasic LC-MS/MS approach

first with a HILIC separation [69] followed by a mixed-mode chromatographic separation

[70]. Quantification of amino acid levels was determined by fitting response ratios to an eight-

point calibration curve generated using verified reference material for each of the 43 amino

acids quantified.

CRISPR knock-out screen. A human SLC knockout library [40] was purchased from Addgene,

(Cat. #132552). Lentivirus was generated as described above. LNCaP and 22Rv1 cells were

transduced at a multiplicity of infection (MOI) of 0.5 and selected with 1 μg/ml puromycin (MP

Biomedicals Cat #194539) for 7 days. Three independent screens for both LNCaP and 22Rv1

were performed. Cells were then cultured for an additional 4 weeks and genomic DNA (gDNA)

was isolated using DNeasy Blood and Tissue Kit (Qiagen, Cat. # 69506). sgRNA sequences

were PCR amplified from gDNA using Platinum SuperFi II DNA Polymerase (ThermoFisher

Scientific, Cat #12369002) and primers sgRNA_F and sgRNA_R. A second round of PCR was

used to add Illumina adaptor sequences using primers IlluminaAd_F and IlluminaAd_R. PCR

primers are shown in **Supplementary Table 3**. Purified PCR products were sequenced using

next generation sequencing (NGS) at GeneWiz (South Plainfield, NJ). Fastq files were trimmed,

aligned, and analyzed using PinAPL-Py [71] and raw read counts were normalized as counts

per million reads (cpm). The fold-depletion for each sgRNA was calculated for LNCaP or 22Rv1

relative to control human SLC knockout library. sgRNA rankings and significance (Sidak

correction method) were determined according to the negative binomial model implemented in

PinAPL-Py [71]. Gene-depletion significance was determined by a permutation test (10^4

permutations), and *p* values were corrected for multiple testing using the Sidak correction

method.

RNA-seq and network analyses. LNCaP cells were seeded into six-well culture plates and

treated with either vehicle (DMSO) or 2 μM GCN2iB in standard culture media as described

above for 6 or 24 hours as indicated. Total RNA was extracted from cells using a RNeasy Plus

Kit (Qiagen, Cat. #74034) according to the manufacturer's instructions. RNA quality control,

library preparation, and NGS paired-end sequencing on Illumina HiSeq platform was carried out

at GeneWiz (South Plainfield, NJ). Raw fastq data files were trimmed to remove adaptor

sequences using cutadapt [72] and the quality of the trimmed reads was analyzed by FASTQC

[73] available in the Trim Galore package [74]. The trimmed paired sequencing reads were

aligned to human reference genome GRCh38 downloaded from ENSEMBL using HISAT2 [75],

resulting in concordant alignment of >92%. FeatureCounts [76] from the Subread package was

used to count the number of reads mapping to unique exonic sequences and was used for

differential gene expression (DEG) analysis using DESeq2 R-package [77], comparing the

DMSO treated control group to the GCN2iB-treated groups.

The computer cluster Carbonate that was configured for high-performance analyses was

used to perform all the steps of sequencing analysis from trimming to differential analysis.

Volcano plots were created from differentially expressed genes using GraphPad Prism.

Differentially expressed genes with a log2 fold change of ≥ ±1 and adjusted p-values of ≤ 0.05

were considered significant for further analysis. Gene set enrichment analysis [78, 79] using the

Hallmark and Curated (C2) gene sets from the Molecular Signatures Database [80, 81] was

used to identify enriched pathways from lists of significant differentially expressed genes. The

GCN2-dependent gene signature was derived from genes showing ≤ -2-fold change (fdr ≤ 0.05)

following treatment with GCN2iB for 6 hours in our RNA-seq analysis. The RNA-seq data

generated in this study has been deposited in the NCBI's Gene Expression Omnibus (GEO)

database under the accession code GSE196252 [Reviewer's Token #kpkvcqcunpidhol].

tRNA charging measurements. The levels of charged (aminoacylated) tRNAs genome-wide

were measured using the CHARGE-seq method [39] with some protocol adjustments as

previously described [82]. The CHARGE-seq method relies on the simultaneous sequencing of

both a charged and uncharged tRNA library. LNCaP cells were treated with vehicle (DMSO),

GCN2iB, or GCN2iB in media supplemented with essential amino acids for 8 hours. Total RNA

was extracted using TRIzol (Life Technologies, Cat. #15596018) and a portion (2 μg) of total

RNA for each sample was left untreated or oxidized by treatement with 12.5 mM $NaIO_4$ in

sodium acetate solution (pH 4.5) in the dark at room temperature for 20 minutes followed by

quenching with 0.3 M glucose. All mature tRNAs end with 3'-CCA and $NaIO_4$ will only oxidize

the 3'-A residue of uncharged tRNAs since the 3'-ends of charged tRNAs are protected by

linked amino acids. The resulting total RNA was desalted by passing through MicroSpin G-50

column (GE Healthcare, Cat. #27533001) and the tRNA was deacylated through β-elimination in

50 mM Tris–HCl (pH 9.0) at 37 °C for 45 minutes followed by quenching with a solution

containing 50 mM sodium acetate buffer (pH 4.5) and 100 mM NaCl. Total RNA was then

recovered by overnight EtOH precipitation at -20°C. Following deacylation, libraries for NGS

sequencing were prepared as follows, RNA samples were treated with polynucleotide kinase

(New England BioLabs, Cat. #M0201) to remove the 3'-phosphate group from the uncharged

tRNA followed by ligation to 5'-adenylated uniquely barcoded adapters using RNA ligase 2

truncated KQ (New England BioLabs, Cat. # M0351L). The resulting tRNAs were then ligated to

a 5'-adaptor, annealed to an RT primer, and converted to cDNA using a using a SuperScript IV

RT kit (Invitrogen, Cat. #18090050). The tRNA library was PCR amplified for 10 cycles using

Illumina multiplex and barcode PCR primers and Platinum SuperFi II DNA Polymerase

(ThermoFisher Scientific, Cat #12369002). All oligonucleotide primers and adapters utilized in

the procedure are listed in **Supplementary Table 3**. The resulting PCR products were

electrophoresed on an 8% non-denaturing polyacrylamide TBE gel (Invitrogen, Cat #EC6215)

and ~210 bp bands were excised, purified, and subjected to Illumina HiSeq paired-end

sequencing. Fastq files from the pooled samples were demultiplexed according to the five-

nucleotide sample barcode contained within the ligated adapter using a custom python script

[83], and read pairs without an identifiable barcode were discarded. Illumina sequencing

adapters and ligated sample barcode adapters were trimmed using cutadapt [72] in paired read

mode and reads with less <20 nucleotides remaining after trimming, corresponding to empty

adapters, were removed.

A custom reference file for human tRNA was constructed from the gtRNAdb GRCh38

high confidence tRNA gene set by removing duplicate sequences and appending 'CCA' to the 3'

end of each unique tRNA gene sequence. Trimmed reads were then aligned to the custom

tRNA reference file using bowtie2 [84] short read aligner with options "--dovetail -D 20 -R 3 -N 1

-L 20 -i S,1,0.50". The charging status of the aligned tRNA was determined from the 3' end of

the aligned portion of each read pair, where reads ending in 'CCA' were determined to be

charged and reads ending in 'CC' were determined to be uncharged. This method was able to

determine the charging status of ~86% percent of all mapped reads. Counts of charged and

uncharged tRNA for each tRNA gene in each sample were then imported to R for further

analysis. Differences in read counts between samples were accounted for by multiplying the

read counts in each sample by a normalization coefficient equal to the average number of reads

across all samples divided by the read count in that sample. The fraction charged for each tRNA

isodecoder within each sample was determined by summing the normalized counts for charged

and uncharged reads for each gene corresponding to that isodecoder, and the mean was then

averaged for each of the four replicates for each treatment. The mean fraction charged +/- SD

for each isodecoder was visualized using Graphpad Prism. Statistical significance in difference

of the mean fraction charged was determined using a Welch's unpaired *t*-test with Benjamini-

Hochberg fdr correction to account for repeated measures. The CHARGE-seq data generated in

this study has been deposited in the NCBI's Gene Expression Omnibus (GEO) database under

the accession code GSE196251 [Reviewer's Token #kpkvcqcunpidhol].

Levels of total and aminoacylated tRNA^{His} were also carried out by qRT-PCR as

previously described [82] using primers specific for the GTG-tRNA^{His} isodecoder tRNA^{His}-FW

and tRNA^{His}-REV (**Supplementary Table 3**).

Animal studies. All animal experiments were approved by the Institutional Animal Care and

Use Committee (IACUC) at Indiana University School of Medicine (Protocol #21014) and

comply with all regulations for ethical conduct of animal research. Male 4 – 6 week old NSG

(NOD.Cg-Prkdcscid Il2rgtm1Wjl/SzJ, RRID:IMSR_JAX:021885) mice were purchased from the

In Vivo Therapeutics Core at Indiana University School of Medicine. GCN2iB used for animal

studies was formulated in 0.5% Methyl Cellulose (Sigma-Aldrich, Cat. #M0262) containing 5%

DMSO. Mice were dosed with vehicle or 30 mg/kg GCN2iB twice daily for 5 days/week by oral

gavage as described in the legends to the figures. For LNCaP, 22Rv1, and PC-3 cell line-

derived xenograft models, $2x10^6$ cells in 100 µl of serum-free RPMI-1640 media containing 50%

Matrigel (Corning, Cat. #356237) were injected subcutaneously into the dorsal flank of mice.

Once the tumors reached a palpable stage (~50 – 100 mm^3), the animals were randomized to

different treatment groups using RandoMice v1.1.5 [85]. The LuCaP-35 castration-resistant

(LuCaP-35 CR) PCa PDX model [49, 86] was received from Dr. Eva Corey (University of

Washington) and were maintained in male castrated NSG mice for all experiments to conserve

castration-resistant status. TM00298 PCa PDX model [48] was obtained from Jackson

Laboratory and male NSG mice were engrafted with ~1 mm^3 PDX tumor fragments and

randomized once the tumors reached a palpable stage (~50 – 100 mm^3). For all xenograft

studies, tumor volume was monitored twice per week in a blinded fashion using a digital caliper

and the formula: Tumor Volume = (width^2 × length)/2. Body weight was determined once per

week during the course of each study. At the end of the studies, mice were euthanized using

CO_2 and tumors were harvested, weighed, and fixed for immunohistochemistry staining or flash

frozen for immunoblot analyses. For the amino acid supplementation experiment *in vivo*, mice

bearing 22Rv1 WT or 22Rv1 GCN2 KO tumors were supplemented with essential amino acids

(1.5 mg amino acids/ g body weight) in the drinking water. The control group received drinking

water with no amino acid supplementation. Supplementation of amino acids was started at the

same time as tumor cell implantation and was maintained for the remainder of the study. Amino

acids were dissolved in water in quantities determined by calculating average daily intake of

drinking water for mice as previously reported [47]. Composition, relative percentage, and an

estimate of dietary intake of each amino acid from the drinking water is reported in

Supplementary Table 4.

Immunohistochemistry. Tumor tissue specimens were fixed for 24 hours in neutral buffered

formalin, paraffin-embedded and sectioned (5 µm). Prostate core needle biopsy specimens

were obtained from the Indiana University Comprehensive Cancer Center Tissue Procurement

and Distribution Core and approval was granted by the Institutional Review Board (IRB #1796)

at the Office of Research Administration at Indiana University. Human tumor tissue microarrays

were purchased from US Biomax (Rockville, MD, Cat. # PR1921b and PR807c). Tissue

sections were deparaffinized and rehydrated through serial graded xylene and alcohol washes.

H&E staining was performed using standard methods. Antigen unmasking was achieved by

boiling the slides in sodium citrate buffer (10 mM Sodium citrate, 0.05% Tween 20, pH 6.0) for

10 – 15 minutes. Sections were further incubated in hydrogen peroxide to reduce endogenous

activity. Tissue blocking was performed with 5% Horse Serum (Vector Laboratories Cat. #PK-

7200) in phosphate-buffered saline (PBS) and incubated with primary antibodies: Ki-67 (1:200,

Abcam Cat. #ab16667, RRID:AB_302459), p-GCN2-T899 (1:100, Abcam Cat. #ab75836,

RRID:AB_1310260), and SLC3A2 (4F2hc/CD98) (1:500, Cell Signaling Technology Cat.

#47213, RRID:AB_2799323). After primary antibody incubation, tissue sections were washed,

incubated in horseradish peroxidase-conjugated anti-rabbit secondary antibody according to the

manufacturer's protocol (Vector Laboratories, Cat #PK-7200), enzymatically developed in

diaminobenzidine (DAB), and counterstained with hematoxylin. Tissue sections were then

dehydrated and mounted with coverslips sealed with Cytoseal™ 60 mounting media

(ThermoFisher Scientific, Cat #23-244257). Stained sections were scanned under brightfield

using a Leica Aperio AT2 slide scanner (Leica Biosystems). Quantification was performed using

Quantitative Pathology & Bioimage Analysis software (QuPath). Representative images were

acquired using an EVOS M5000 cell imaging microscope (Life Technologies).

Statistical Analysis. All statistical tests were carried out using GraphPad Prism version 9.2.0

for Windows as indicated in the legends to the figures or as outlined in **Supplementary File 1**.

Figure 1

Figure 1. GCN2 promotes growth of PCa cells. (A) Expression of the indicated eIF2α kinase

was reduced in LNCaP cells using gene specific siRNAs. Two different siRNAs were used for

knock-down of each eIF2α kinase and compared to scrambled siRNA control. Cell growth was

measured in replicate wells (N = 5) for up to 6 days and is plotted as fold change (mean ± SD)

relative to day 0. Statistical significance was determined using a two-way ANOVA as described

in **Supplementary File 1**; *, $p \le 0.05$; ***, $p \le 0.001$; ****, $p \le 0.0001$. **(B)** LNCaP cells were

transfected with two different siRNAs targeting GCN2 or a scramble siRNA control and cell

lysates were prepared and immunoblotted for the indicated proteins. Molecular weight markers

are shown in kilodaltons. The relative levels of p-eIF2α normalized to total eIF2α compared to

scramble siRNA control are indicated. **(C)** Expression of GCN2 was knocked-down in LAPC-4,

C4-2B, MR49F, 22Rv1, or PC-3 cells using two different siRNAs and compared to scrambled

siRNA control. Cell growth was measured for up to 6 days in replicate wells (N = 5) as

described in **A**. Statistical significance was determined using a two-way ANOVA as described in

Supplementary File 1; ***, $p \le 0.001$; ****, $p \le 0.0001$. **(D)** LNCaP cells were treated with

indicated concentrations of GCN2iB and cell growth was measured for up to 6 days in replicate

wells (N = 5) as described in **A**. Statistical significance was determined using a two-way ANOVA

as described in **Supplementary File 1**; ****, $p \le 0.0001$. **(E)** LNCaP cells were treated with

GCN2iB (2 μM) or DMSO control for 24 hours and protein lysates were analyzed by immunoblot

using antibodies that recognize total or phosphorylated GCN2-T899, total or phosphorylated

eIF2α−S51, ATF4, or actin as indicated. Relative levels of p-eIF2α normalized to total eIF2α are

shown. **(F)** Levels of p-GCN2 were measured in prostate tumor microarrays (Biomax PR1921b

and PR807c) using immunohistochemistry (IHC). Staining for p-GCN2-T899 from non-malignant

(N = 33) and malignant prostate cancer tissue (N = 88) from patients >50 years old was

analyzed and quantified using QuPath to determine the histoscore and is represented as a

scatterplot. Statistical significance was determined using an unpaired two-tailed *t*-test; *, $p \le$

0.05. Representative images showing p-GCN2-T899 staining of normal and malignant prostate tissues are shown. Scale bars shown are 200 μm (main image) and 20 μm (insert).

Figure 2

A

24 hours GCN2iB/ Vehicle

B

Gene Set	NES	p-val
KRIGE_AMINO_ACID_DEPRIVATION	1.75	<0.001
GARGALOVIC_RESPONSE_TO_OXIDIZED_PHOSPHOLIPIDS_RED_UP	1.67	<0.001
REACTOME_FCGR_ACTIVATION	1.67	<0.001
REACTOME_IRE1ALPHA_ACTIVATES_CHAPERONES	1.63	<0.001
REACTOME_AMINO_ACID_TRANSPORT_ACROSS_THE_PLASMA_MEMBRANE	1.62	<0.001
REACTOME_RESPONSE_OF_EIF2AK1_HRI_TO_HEME_DEFICIENCY	1.61	0.001
PACHER_TARGETS_OF_IGF1_AND_IGF2_UP	1.6	0.001
ZHAN_MULTIPLE_MYELOMA_CD1_VS_CD2_UP	1.58	<0.001
REACTOME_ROLE_OF_PHOSPHOLIPIDS_IN_PHAGOCYTOSIS	1.55	<0.001
ZHAN_MULTIPLE_MYELOMA_CD1_UP	1.54	<0.001
REACTOME_CYTOSOLIC_TRNA_AMINOACYLATION	1.53	0.002
KEGG_GLYCINE_SERINE_AND_THREONINE_METABOLISM	1.53	0.001
KEGG_AMINOACYL_TRNA_BIOSYNTHESIS	1.53	0.001

C

D

E

F

G

Figure 2: GCN2 induces ISR transcriptome featuring expression of amino acid transporters. (A) Volcano plot illustrating $\log_2$ fold change with adjusted *p*-value ($-\log_{10}$) comparing LNCaP cells treated with GCN2iB (2 µM) versus vehicle control (DMSO) for 24 hours. Several amino acid transporters reduced by GCN2iB treatment are highlighted. **(B)** Enrichment plots from Gene Set Enrichment Analysis (GSEA) of gene expression in LNCaP cells treated with GCN2iB (2 µM) for 24 hours versus vehicle control. **(C)** Heat map displaying differential gene expression of *SLC* genes comparing LNCaP cells treated with vehicle (DMSO), or GCN2iB (2 µM) for 6 or 24 hours. *SLC* genes with $\log_2$ fold change $\leq$ -0.5 and $p \leq 0.05$ are shown. **(D)** Lysates were prepared from LNCaP cells treated with 2 µM GCN2iB or vehicle control (DMSO) for 6 or 24 hours and immunoblot analysis were carried out using antibodies that recognize ATF4, ASNS, xCT (SLC7A11), 4F2 (SLC3A2), CAT1 (SLC7A1), ASCT1 (SLC1A4), ASCT2 (SLC1A5), or actin. Molecular weight markers are indicated in kilodaltons. **(E)** 22Rv1 WT cells, 22Rv1 GCN2 KO cells, and 22Rv1 GCN2 KO complemented with GCN2 cells were cultured for 24 hours. Lysates were prepared and analyzed by immunoblot for the indicated proteins. **(F)** Amino acid uptake measurements in LNCaP and 22Rv1 cells treated with vehicle (DMSO) or GCN2iB (2 µM) for 24 hours. **(G)** Amino acid uptake measurements for 22Rv1 WT or 22Rv1 GCN2 KO cells cultured for 24 hours. Statistical significance was determined using an unpaired two-tailed *t*-test (N = 4); $*p \leq 0.05$, $**p \leq 0.01$, $***p \leq 0.001$.

Figure 3

Figure 3. GCN2 is critical for maintenance of free amino acids. (A) Amino acid

measurements of LNCaP cells treated with 2 µM GCN2iB or vehicle control (DMSO) for 8 hours.

Bar graphs in the top panel show high abundance amino acids and the lower panel those with

lower levels. The heatmap on the right shows fold change in amino acid abundance for each

biological replicate of GCN2iB-treated LNCaP cells vs the vehicle with the scale showing the

highest fold change in yellow and lowest in purple. Statistical significance was determined using

an unpaired two-tailed t-test. Error bars indicate SD (N = 3); *, $p \leq 0.05$; **, $p \leq 0.01$, ***, $p \leq$

0.001; ****, $p \leq 0.0001$. **(B)** LNCaP cells were treated with vehicle, GCN2iB (2 µM), vehicle +

essential amino acids (EAA), or GCN2iB (2 µM) + EAA, and cell growth was measured for up to

6 days. Error bars indicate SD (N = 5). Statistical significance was determined using a two-way

ANOVA as described in **Supplementary File 1**; ****, $p \leq 0.0001$. **(C)** Cell cycle analyses of

LNCaP cells treated with vehicle, GCN2iB (2 µM), vehicle + EAA, or GCN2iB (2 µM) + EAA for

48 hours. Statistical significance was determined using a one-way ANOVA with Tukey's multiple

comparisons. Error bars indicate SD (N=3); ***, $p \leq 0.001$; ****, $p \leq 0.0001$. **(D)** Genome-wide

tRNA charging analysis (CHARGE-seq) of LNCaP cells treated with vehicle (DMSO), GCN2iB

(2 µM), or GCN2iB (2 µM) + EAA for 8 hours. The tRNA charging ratio is shown as a bar graph

with fold change compared to vehicle. Only tRNA isoacceptors measured in LNCaP cells are

shown. Error bars indicate SD (N = 4). **(E)** tRNA charging percentage for tRNA^{His} in LNCaP cells

treated with vehicle, GCN2iB, or GCN2iB + EAA. Statistical significance was determine using a

one-way ANOVA with Tukey's multiple comparisons (N = 4); ***, $p \leq 0.001$; ****, $p \leq 0.0001$. **(F)**

LNCaP cells were treated with vehicle, GCN2iB (2 µM), GCN2iB (2 µM) + EAA, or GCN2iB (2

µM) combined with the indicated individual amino acids. Cell growth was measured at 4 days in

triplicate wells (N = 3). Statistical significance was determined using a one-way ANOVA with

Tukey's multiple comparisons. Error bars indicate SD; ****, $p \leq 0.0001$. **(G)** Cell cycle analysis of

LNCaP cells were treated with vehicle, GCN2iB (2 µM), GCN2iB (2 µM) + histidine (200 µM), or

with media lacking histidine for 48 hours. Statistical significance was determined using a one-

way ANOVA with Tukey's multiple comparisons. Error bars indicate SD (N = 3); ***, $p \leq 0.001$;

****, $p \leq 0.0001$. **(H)** LNCaP cells were cultured in normal media, media supplemented with EAA

mix, or media supplemented with histidine (200 µM) for 24 hours. Lysates were analyzed by

Immunoblot using antibodies that recognize total or phosphorylated GCN2-T899, total or

phosphorylated eIF2α−S51, ATF4, or actin. Molecular weight markers are presented in

kilodaltons for each immunoblot panel. The relative levels of p-eIF2α normalized to total eIF2α

compared to normal media (NM) control are indicated.

Figure 4

Figure 4. GCN2 is critical for expression of 4F2 (SLC3A2) and facilitates amino acid transport. (A) Gene-level depletion for LNCaP and 22Rv1 cells. The average log2 fold-change for the sgRNAs for each gene is shown on the x axis. Significantly depleted genes ($p \leq 0.05$) in LNCaP or 22Rv1 are indicated. Circle size indicates the number of significant sgRNAs. *SLC-*genes in red are dependent on GCN2 for expression. **(B)** Plot of -Log_{10} (p-value) for depleted genes identified in CRISPR screen for LNCaP vs. 22Rv1 cells. Significantly depleted genes ($p \geq$ 0.05) in LNCaP, 22Rv1 or both cell lines are indicated. SLC-genes in red are GCN2-dependent. **(C)** Lysates from LNCaP cells were treated with 2 µM GCN2iB for 6 or 24 hours, or with vehicle (DMSO) were analyzed by immunoblot analyses using antibodies that recognize total or phosphorylated GCN2-T899, ATF4, 4F2 (SLC3A2), or actin. Molecular weight markers are indicated in kilodaltons for the panels. **(D)** LNCaP cells were cultured in standard culture conditions (NM: normal media), media supplemented with 200 µM histidine (+ His), or media depleted of histidine (- His) for 24 hours. Lysates were analyzed by Immunoblot analyses using antibodies that recognize total or phosphorylated GCN2-T899, ATF4, 4F2 (SLC3A2), or actin. **(E)** LNCaP cells were treated with 100 nM halofuginone (HF) for 2 and 6 hours or vehicle (DMSO). Lysates were analyzed by Immunoblot using antibodies that recognize the indicated proteins. **(F)** 4F2 (SLC3A2) expression was reduced in LNCaP or 22Rv1 cells using two different siRNAs or scramble siRNA as a control. Cell growth was measured in replicate wells (N = 5) for up to 6 days and are plotted relative to day 0 (mean ± SD). Statistical significance was determined using a two-way ANOVA as described in **Supplementary File 1**; ****, $p \leq 0.0001$. **(G)** LNCaP cells transfected with two different siRNAs targeting 4F2 (SLC3A2) or scramble siRNA for 48 hours. Lysate were prepared and analyzed by immunoblot using antibodies that recognize total or phosphorylated GCN2-T899, total or phosphorylated eIF2α–S51, ATF4, 4F2 (SLC3A2), or actin. **(H)** LNCaP cells stably overexpressing SLC3CA2 or vector control were transfected with two different siRNAs targeting GCN2 or scrambled control. Cells were then treated with GCN2iB (2 µM) or vehicle and growth was measured in

replicate wells (N = 5) and is plotted relative to day 0 (mean ± SD). Statistical significance was determined using a two-way ANOVA as described in **Supplementary File 1**; **, $p \le 0.01$; ****, $p \le 0.0001$. **(I)** Amino acid measurements of LNCaP cells transfected siRNA targeting GCN2 (N = 4), 4F2 (SLC3A2, N = 4), or scramble control (N = 8). Two separate bar graphs show high abundance (top) and low abundance (bottom) amino acids. Statistical significance was determined using a two-way ANOVA as described in **Supplementary File 1**. Error bars indicate SD; *, $p \le 0.05$; **, $p \le 0.01$; ***, $p \le 0.001$; ****, $p \le 0.0001$.

Figure 5

Figure 5. Model for GCN2 regulation of amino acid homeostasis in prostate cancer. Model

depicting the role of GCN2 in regulating *SLC* amino acid transporters. Enhanced translation and

altered metabolism driven by oncogenes deplete amino acid pools resulting in accumulation of

uncharged tRNAs leading to activation of GCN2. Active GCN2 results in increased expression

of SLC amino acid transporters, including 4F2 (SLC3A2), to increase uptake of amino acids.

Loss of GCN2 function disrupts amino acid homeostasis decreasing proliferation of prostate

cancer cells.

Figure 6

Figure 6. GCN2 regulates amino acid transporters ensuring sufficient amino acids for tumor growth in mouse PCa xenograft models. (A) WT or GCN2 KO 22Rv1 (clone 7) cells were injected subcutaneously into the dorsal flank of mice as described in the materials and methods section. Tumor volume (TV) was measured on indicated days and is plotted as average TV ± SEM (N = 4). Statistical significance was determined using a two-way ANOVA with Sidak's multiple comparison; *, $p \leq 0.05$; ****, $p \leq 0.0001$. On the right bar graph, the final tumor weight was measured at endpoint and statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD; **, $p \leq 0.01$. **(B)** Protein lysates were prepared from WT and GCN2 KO 22Rv1 tumors and analyzed by immunoblot to measure total GCN2, ATF4, LAT1 (SLC7A5), xCT (SLC7A11), 4F2 (SLC3A2), androgen receptor (AR), AR splice variant 7 (AR-V7), or actin. Molecular weight markers are indicated in kilodaltons for each immunoblot panel. The levels of the SLC proteins normalized to actin are shown in the bar graph (right panels). Statistical significance was determined using an unpaired two-tail t-test. Error bars indicate SD (N = 4); *, $p \leq 0.05$; ***, $p \leq 0.001$. **(C)** Tumor growth of PC-3 WT and PC-3 GCN2 KO (clone 3) cells was analyzed in a mouse xenograft study as in A. Statistical significance was determined using a two-way ANOVA with Sidak's multiple comparison. Error bars indicate SEM (N = 5); ***, $p \leq 0.001$; ****, $p \leq 0.0001$. On the right bar graph, the final tumor weight was measured at endpoint. Statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD (N = 5); *, $p \leq 0.01$. **(D)** Protein lysates were prepared from the PC-3 WT and PC-3 GCN2 KO tumors and analyzed by immunoblot for the indicated proteins. (Right panels) Quantification of protein levels of LAT1 (SLC7A5), xCT (SLC7A11), and 4F2 (SLC3A2) normalized to actin are shown in the bar graphs. Statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD (N = 4); **, $p \leq 0.01$. **(E)** 22Rv1 WT (N = 4), 22Rv1 GCN2 KO (clone 7, N = 5) and 22Rv1 ATF4 KO (N = 5) were evaluated in the mouse xenograft model. Tumor volumes were measured on the indicated days. Error bars indicated SEM. **(F)** Amino acid measurements of 22Rv1 WT, 22Rv1 GCN2 KO and

22Rv1 ATF4 KO tumors. Two separate bar graphs show high abundance (top) and low abundance (bottom) amino acids. Statistical significance was determined one-way ANOVA with Tukey's multiple comparisons. Error bars indicate SD (N = 4); #, $p \leq 0.1$; *, $p \leq 0.05$. **(G)** 22Rv1 WT and 22Rv1 GCN2 KO (clone 7) cells were analyzed in a xenograft model as described for figure **6A**, with or without supplementation of EAA in the drinking water. Tumor volume was measured on indicated days. 22Rv1 WT (N = 4) and 22Rv1 KO (N = 5) are the same tumor growth curves shown in **6E**. 22Rv1 WT + EAA (N = 5), 22Rv1 GCN2 KO + EAA (N = 5). Statistical significance was determined using a two-way ANOVA with Sidak's multiple comparison; ****, $p \leq 0.0001$. **(H)** Amino acid measurements for 22Rv1 WT, 22Rv1 GCN2, and 22Rv1 GCN2 KO + EAA tumors. Bar graphs show only amino acids present in EAA supplemented water. Statistical significance was determined using a one-way ANOVA with Tukey's multiple comparisons. Error bars indicated SD (N = 4); #, $p \leq 0.1$; *, $p \leq 0.05$; **, $p \leq 0.01$. **(I)** Tumor growth curves for 22Rv1 WT or 22Rv1 GCN2 KO (clone 11) transduced with 4F2 (SLC3A2) lentivirus (WT + 4F2 and GCN2 KO + 4F2) or empty vector (WT + EV and GCN2 KO + EV). Tumor volumes were measured on indicated days. Error bars indicate SEM (N = 5). Statistical significance was determined using a two-way ANOVA with Sidak's multiple comparison; ****, $p \leq 0.0001$.

Figure 7

Figure 7. Pharmacological inhibition of GCN2 reduces tumor growth in cell line-derived and patient-derived xenograft models. Male NSG mice were injected subcutaneously with LNCaP (N = 5) **(A)** or 22Rv1 (N=4) **(B)** cells, or alternatively implanted with tumor fragments from an androgen-sensitive tumor TM00298 (N = 5) **(C).** Male castrated NSG mice were implanted with tumor fragments from LuCaP-35 CR tumors (N = 5) **(D).** Mice were treated with vehicle or 30 mg/kg GCN2iB twice daily for 5 days per week and tumor volumes were measured on indicated days. Statistical significance was determined using a two-way ANOVA with Sidak's multiple comparison. Error bars indicate SEM; *, $p \le 0.05$; **, $p \le 0.01$; ***, $p \le 0.001$; ****, $p \le 0.0001$. Final tumor weight was measured at endpoint and is represented in bar graphs (right panels). Statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD; *, $p \le 0.05$. **(E)** Protein lysates were prepared from 22Rv1 tumors treated with Vehicle or GCN2iB and analyzed by immunoblot for phosphorylated GCN2-T899, ATF4, LAT1 (SLC7A5), xCT (SLC7A11), and 4F2 (SLC3A2), and actin. The levels of the SLC proteins normalized to actin are shown. Phosphorylated GCN2-T899 was normalized to total GCN2. Statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD (N = 4); *, $p \le 0.05$. **(F)** Amino acid measurements of 22Rv1 tumors treated with vehicle or GCN2iB. Bar graphs show high abundance (top) and low abundance (bottom) amino acids. Statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD (N = 4); #, $p \le 0.1$; *, $p \le 0.05$; **, $p \le 0.01$. **(G)** Pearson correlation between p-GCN2-T899 and 4F2 (SLC3A2) histoscores calculated from IHC staining from a prostate tumor microarray (Biomax PR807c) containing normal (N = 10), hyperplasia (N = 20) and malignant (N = 50) for all tissues (combined) or Gleason scores 4 and 5. The center lines depict linear regression (95% confidence intervals). Not all samples were analyzed due to damaged/quality of tissue samples. Levels of p-GCN2-T899 and 4F2 (SLC3A2) were measured by IHC staining and QuPath was used to determine the histoscore. Two representative cases are shown for high (Case 1) and low (Case 2) p-GCN2-T899 and 4F2 (SLC3A2) staining. Scale bar indicates 200

μm (main image) and 20 μm (insert). **(H)** Correlation of expression of 4F2 (SLC3A2) and a GCN2-dependent gene signature in prostate adenocarcinoma (PRAD, N = 551) from the Cancer Genome Atlas (TCGA). The GCN2-dependepent gene signature was derived from RNA-seq data as described in the materials and methods.

Data Availability Statement

The authors declare that all data generated or analyzed in this study are included in the published article, its supplementary information and source files, or are publicly available. The CHARGE-seq and RNA-seq datasets generated in this study have been deposited in the NCBI Gene Expression Omnibus (GEO) database under the ascension codes GSE196251 and GSE196252, respectively. The custom python script used in the analysis of our Charge-seq study is available on GitHub (https://github.com/carlsonkPhD/tRNA_Charge-Seq/). Gene expression data from prostate cancer patients (PRAD) in the TCGA database used for correlation analysis is publicly available.

Acknowledgements

This work was supported in part by grants from NIH GM136331 (R.C.W.), DK109714 (T.G.A., R.C.W.), the NCI R21CA221942 (R.P.), the Indiana University Melvin and Bren Simon Comprehensive Cancer Center P30CA082709 (K.A.S.), and the Lilly Endowment, Inc., through its support for the Indiana University Pervasive Technology Institute. We thank the IU Simon Comprehensive Cancer Center for use of the Tissue Procurement and Distribution Core, which provided core needle biopsy specimens for staining, the Preclinical Modeling and Therapeutics Core (PMTC) for assistance with flow cytometry and cell cycle analysis, and the Histology Lab Service Core at the Indiana University School of Medicine for embedding and sectioning of tissue samples. Finally, the authors acknowledge Ludovica Ceci for assistance with slide scanning, Kadir Isidan for assistance with histology, and members of Wek and Pili laboratories for helpful discussions.

Competing Interests

R.C.W. is a member of the advisory board and holds equity in HiberCell, Inc. K.A.S. and T.G.A are consultants for HiberCell, Inc. and K.A.S. receives research support from HiberCell, Inc.

Author Contributions

R.A.C., R.P., R.C.W, and K.A.S. conceived of and designed the experiments, led the project, and coordinated the writing of the manuscript with the assistance of all of the co-authors. N.P.D. provided technical advice and assisted with *in vivo* experiments. B.D.E. provided technical advice and assistance with the LuCaP-35 CR animal model. A.J.K. and A.E. provided technical assistance with cell culture experiments and immunoblot analyses. H.-G.K. provided technical advice and assistance with IHC staining. L.C. provided pathology interpretation of prostate cancer specimens. E.T.M., M.M., and T.G.A. assisted with amino acid analysis. J.M. carried out the CHARGE-seq and tRNA charging analysis, and P.H.A. and K.C. provided bioinformatics support for the RNA-seq and CHARGE-seq studies.

Supplementary File Descriptions

Supplementary File 1. Statistical analysis for growth assays.

Supplementary File 2. Normalized read counts and analysis of RNA-seq data.

Supplementary File 3. Normalized read counts and analysis of Charge-seq data.

Supplementary File 4. CRISPR screen analysis of depleted genes and sgRNAs.

References

1. Siegel, R.L., K.D. Miller, and A. Jemal, *Cancer statistics, 2019.* CA Cancer J Clin, 2019. **69**(1): p. 7-34.
2. Gann, P.H., *Risk factors for prostate cancer.* Rev Urol, 2002. **4 Suppl 5**: p. S3-S10.
3. Klusa, D., et al., *Metastatic Spread in Prostate Cancer Patients Influencing Radiotherapy Response.* Front Oncol, 2020. **10**: p. 627379.
4. Heinlein, C.A. and C. Chang, *Androgen receptor in prostate cancer.* Endocr Rev, 2004. **25**(2): p. 276-308.
5. Huggins, C. and C.V. Hodges, *Studies on prostatic cancer: I. The effect of castration, of estrogen and of androgen injection on serum phosphatases in metastatic carcinoma of the prostate. 1941.* J Urol, 2002. **168**(1): p. 9-12.
6. Polotti, C.F., et al., *Androgen deprivation therapy for the treatment of prostate cancer: a focus on pharmacokinetics.* Expert Opin Drug Metab Toxicol, 2017. **13**(12): p. 1265-1273.
7. Virgo, K.S., et al., *Second-Line Hormonal Therapy for Men With Chemotherapy-Naive, Castration-Resistant Prostate Cancer: American Society of Clinical Oncology Provisional Clinical Opinion.* J Clin Oncol, 2017. **35**(17): p. 1952-1964.
8. Watson, P.A., V.K. Arora, and C.L. Sawyers, *Emerging mechanisms of resistance to androgen receptor inhibitors in prostate cancer.* Nat Rev Cancer, 2015. **15**(12): p. 701-11.
9. Crawford, E.D., et al., *Androgen-targeted therapy in men with prostate cancer: evolving practice and future considerations.* Prostate Cancer Prostatic Dis, 2019. **22**(1): p. 24-38.
10. Bubley, G.J. and S.P. Balk, *Association Between Androgen Receptor Splice Variants and Prostate Cancer Resistance to Abiraterone and Enzalutamide.* J Clin Oncol, 2017. **35**(19): p. 2103-2105.
11. Baird, T.D. and R.C. Wek, *Eukaryotic initiation factor 2 phosphorylation and translational control in metabolism.* Adv Nutr, 2012. **3**(3): p. 307-21.
12. Tian, X., et al., *Targeting the Integrated Stress Response in Cancer Therapy.* Front Pharmacol, 2021. **12**: p. 747837.
13. Dong, J., et al., *Uncharged tRNA activates GCN2 by displacing the protein kinase moiety from a bipartite tRNA-binding domain.* Mol Cell, 2000. **6**(2): p. 269-79.
14. Inglis, A.J., et al., *Activation of GCN2 by the ribosomal P-stalk.* Proc Natl Acad Sci U S A, 2019. **116**(11): p. 4946-4954.
15. Wek, S.A., S. Zhu, and R.C. Wek, *The histidyl-tRNA synthetase-related sequence in the eIF-2 alpha protein kinase GCN2 interacts with tRNA and is required for activation in response to starvation for different amino acids.* Mol Cell Biol, 1995. **15**(8): p. 4497-506.
16. Walter, P. and D. Ron, *The unfolded protein response: from stress pathway to homeostatic regulation.* Science, 2011. **334**(6059): p. 1081-6.
17. Wek, R.C. and D.R. Cavener, *Translational control and the unfolded protein response.* Antioxid Redox Signal, 2007. **9**(12): p. 2357-71.
18. Chukwurah, E., et al., *A tale of two proteins: PACT and PKR and their roles in inflammation.* FEBS J, 2021. **288**(22): p. 6365-6391.
19. Garcia, M.A., et al., *Impact of protein kinase PKR in cell biology: from antiviral to antiproliferative action.* Microbiol Mol Biol Rev, 2006. **70**(4): p. 1032-60.

20. Fessler, E., et al., *A pathway coordinated by DELE1 relays mitochondrial stress to the cytosol.* Nature, 2020. **579**(7799): p. 433-437.

21. Guo, X., et al., *Mitochondrial stress is relayed to the cytosol by an OMA1-DELE1-HRI pathway.* Nature, 2020. **579**(7799): p. 427-432.

22. Suragani, R.N., et al., *Heme-regulated eIF2alpha kinase activated Atf4 signaling pathway in oxidative stress and erythropoiesis.* Blood, 2012. **119**(22): p. 5276-84.

23. Staschke, K.A. and R.C. Wek, *Adapting to cell stress from inside and out.* Nat Cell Biol, 2019. **21**(7): p. 799-800.

24. Tameire, F., et al., *ATF4 couples MYC-dependent translational activity to bioenergetic demands during tumour progression.* Nat Cell Biol, 2019. **21**(7): p. 889-899.

25. Nguyen, H.G., et al., *Development of a stress response therapy targeting aggressive prostate cancer.* Sci Transl Med, 2018. **10**(439).

26. Harding, H.P., Y. Zhang, and D. Ron, *Protein translation and folding are coupled by an endoplasmic-reticulum-resident kinase.* Nature, 1999. **397**(6716): p. 271-4.

27. Young, S.K. and R.C. Wek, *Upstream Open Reading Frames Differentially Regulate Gene-specific Translation in the Integrated Stress Response.* J Biol Chem, 2016. **291**(33): p. 16927-35.

28. Hinnebusch, A.G., I.P. Ivanov, and N. Sonenberg, *Translational control by 5'-untranslated regions of eukaryotic mRNAs.* Science, 2016. **352**(6292): p. 1413-6.

29. Kuruma, H., et al., *A novel antiandrogen, Compound 30, suppresses castration-resistant and MDV3100-resistant prostate cancer growth in vitro and in vivo.* Mol Cancer Ther, 2013. **12**(5): p. 567-76.

30. DeSantis, C.E., et al., *Cancer statistics for adults aged 85 years and older, 2019.* CA Cancer J Clin, 2019. **69**(6): p. 452-467.

31. Li, Y., et al., *Androgen receptor splice variants mediate enzalutamide resistance in castration-resistant prostate cancer cell lines.* Cancer Res, 2013. **73**(2): p. 483-9.

32. Sramkoski, R.M., et al., *A new human prostate carcinoma cell line, 22Rv1.* In Vitro Cell Dev Biol Anim, 1999. **35**(7): p. 403-9.

33. Tilley, W.D., et al., *Androgen receptor gene expression in human prostate carcinoma cell lines.* Cancer Res, 1990. **50**(17): p. 5382-6.

34. Nakamura, A., et al., *Inhibition of GCN2 sensitizes ASNS-low cancer cells to asparaginase by disrupting the amino acid response.* Proc Natl Acad Sci U S A, 2018. **115**(33): p. E7776-E7785.

35. Hayward, S.W., et al., *Establishment and characterization of an immortalized but non-transformed human prostate epithelial cell line: BPH-1.* In Vitro Cell Dev Biol Anim, 1995. **31**(1): p. 14-24.

36. Lehman, S.L., et al., *Translational Upregulation of an Individual p21Cip1 Transcript Variant by GCN2 Regulates Cell Proliferation and Survival under Nutrient Stress.* PLoS Genet, 2015. **11**(6): p. e1005212.

37. Leung-Pineda, V., et al., *Induction of p21 and p27 expression by amino acid deprivation of HepG2 human hepatoma cells involves mRNA stabilization.* Biochem J, 2004. **379**(Pt 1): p. 79-88.

38. Battu, S., et al., *Amino Acid Sensing via General Control Nonderepressible-2 Kinase and Immunological Programming.* Front Immunol, 2017. **8**: p. 1719.

39.   Pavlova, N.N., et al., *Translation in amino-acid-poor environments is limited by tRNA(Gln) charging.* Elife, 2020. **9**.

40.   Girardi, E., et al., *A widespread role for SLC transmembrane transporters in resistance to cytotoxic drugs.* Nat Chem Biol, 2020. **16**(4): p. 469-478.

41.   Pizzagalli, M.D., A. Bensimon, and G. Superti-Furga, *A guide to plasma membrane solute carrier proteins.* FEBS J, 2021. **288**(9): p. 2784-2835.

42.   Boyce, M., et al., *A selective inhibitor of eIF2alpha dephosphorylation protects cells from ER stress.* Science, 2005. **307**(5711): p. 935-9.

43.   Novoa, I., et al., *Feedback inhibition of the unfolded protein response by GADD34-mediated dephosphorylation of eIF2alpha.* J Cell Biol, 2001. **153**(5): p. 1011-22.

44.   Keller, T.L., et al., *Halofuginone and other febrifugine derivatives inhibit prolyl-tRNA synthetase.* Nat Chem Biol, 2012. **8**(3): p. 311-7.

45.   Jiang, H.Y., et al., *Activating transcription factor 3 is integral to the eukaryotic initiation factor 2 kinase stress response.* Mol Cell Biol, 2004. **24**(3): p. 1365-77.

46.   Zhang, P., et al., *The GCN2 eIF2alpha kinase is required for adaptation to amino acid deprivation in mice.* Mol Cell Biol, 2002. **22**(19): p. 6681-8.

47.   Corsetti, G., et al., *Dietary supplementation with essential amino acids boosts the beneficial effects of rosuvastatin on mouse kidney.* Amino Acids, 2014. **46**(9): p. 2189-203.

48.   Sekhar, K.R., et al., *Radiosensitization by enzalutamide for human prostate cancer is mediated through the DNA damage repair pathway.* PLoS One, 2019. **14**(4): p. e0214670.

49.   Corey, E., et al., *LuCaP 35: a new model of prostate cancer progression to androgen independence.* Prostate, 2003. **55**(4): p. 239-46.

50.   Maimaiti, M., et al., *The heavy chain of 4F2 antigen promote prostate cancer progression via SKP-2.* Sci Rep, 2021. **11**(1): p. 11478.

51.   Lieu, E.L., et al., *Amino acids in cancer.* Exp Mol Med, 2020. **52**(1): p. 15-30.

52.   Broer, S., *Amino Acid Transporters as Targets for Cancer Therapy: Why, Where, When, and How.* Int J Mol Sci, 2020. **21**(17).

53.   Bhutia, Y.D., et al., *Amino Acid transporters in cancer and their relevance to "glutamine addiction": novel targets for the design of a new class of anticancer drugs.* Cancer Res, 2015. **75**(9): p. 1782-8.

54.   Pathria, G. and Z.A. Ronai, *Harnessing the Co-vulnerabilities of Amino Acid-Restricted Cancers.* Cell Metab, 2021. **33**(1): p. 9-20.

55.   Butler, M., L.T. van der Meer, and F.N. van Leeuwen, *Amino Acid Depletion Therapies: Starving Cancer Cells to Death.* Trends Endocrinol Metab, 2021. **32**(6): p. 367-381.

56.   Wu, C.C., et al., *Ribosome Collisions Trigger General Stress Responses to Regulate Cell Fate.* Cell, 2020. **182**(2): p. 404-416 e14.

57.   Zaborske, J.M., et al., *Genome-wide analysis of tRNA charging and activation of the eIF2 kinase Gcn2p.* J Biol Chem, 2009. **284**(37): p. 25254-67.

58.   Wek, R.C., B.M. Jackson, and A.G. Hinnebusch, *Juxtaposition of domains homologous to protein kinases and histidyl-tRNA synthetases in GCN2 protein suggests a mechanism for coupling GCN4 expression to amino acid availability.* Proc Natl Acad Sci U S A, 1989. **86**(12): p. 4579-83.

59.   Furnish, M., et al., *MIRO2 Regulates Prostate Cancer Cell Growth via GCN1-Dependent Stress Signaling.* Mol Cancer Res, 2022.

60.   Mick, E., et al., *Distinct mitochondrial defects trigger the integrated stress response depending on the metabolic state of the cell.* Elife, 2020. **9**.

61.   Scalise, M., et al., *The Human SLC7A5 (LAT1): The Intriguing Histidine/Large Neutral Amino Acid Transporter and Its Relevance to Human Health.* Front Chem, 2018. **6**: p. 243.

62.   Son, D.O., H. Satsu, and M. Shimizu, *Histidine inhibits oxidative stress- and TNF-alpha-induced interleukin-8 secretion in intestinal epithelial cells.* FEBS Lett, 2005. **579**(21): p. 4671-7.

63.   Kanarek, N., et al., *Histidine catabolism is a major determinant of methotrexate sensitivity.* Nature, 2018. **559**(7715): p. 632-636.

64.   Wortel, I.M.N., et al., *Surviving Stress: Modulation of ATF4-Mediated Stress Responses in Normal and Malignant Cells.* Trends Endocrinol Metab, 2017. **28**(11): p. 794-806.

65.   Pallmann, N., et al., *Regulation of the unfolded protein response through ATF4 and FAM129A in prostate cancer.* Oncogene, 2019. **38**(35): p. 6301-6318.

66.   Liu, G.Y. and D.M. Sabatini, *mTOR at the nexus of nutrition, growth, ageing and disease.* Nat Rev Mol Cell Biol, 2020. **21**(4): p. 183-203.

67.   Schmidt, E.K., et al., *SUnSET, a nonradioactive method to monitor protein synthesis.* Nat Methods, 2009. **6**(4): p. 275-7.

68.   Nikonorova, I.A., et al., *Time-resolved analysis of amino acid stress identifies eIF2 phosphorylation as necessary to inhibit mTORC1 activity in liver.* J Biol Chem, 2018. **293**(14): p. 5005-5015.

69.   Prinsen, H., et al., *Rapid quantification of underivatized amino acids in plasma by hydrophilic interaction liquid chromatography (HILIC) coupled with tandem mass-spectrometry.* J Inherit Metab Dis, 2016. **39**(5): p. 651-660.

70.   Griffin, C., et al., *Rapid quantification of underivatized alloisoleucine and argininosuccinate using mixed-mode chromatography with tandem mass spectrometry.* J Chromatogr B Analyt Technol Biomed Life Sci, 2019. **1128**: p. 121786.

71.   Spahn, P.N., et al., *PinAPL-Py: A comprehensive web-application for the analysis of CRISPR/Cas9 screens.* Sci Rep, 2017. **7**(1): p. 15854.

72.   Martin, M., *Cutadapt removes adapter sequences from high-throughput sequencing reads.* 2011, 2011. **17**(1): p. 3.

73.   Andrews, S., *FastQC: A quality control tool for high throughput sequence data.* https://www.bioinformatics.babraham.ac.uk/projects/fastqc/, 2010.

74.   Krueger, F., *Trim Galore: A wrapper tool around Cutadapt and FastQC to consistently apply quality and adapter trimming to FastQ files.* https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/, 2012.

75.   Kim, D., B. Langmead, and S.L. Salzberg, *HISAT: a fast spliced aligner with low memory requirements.* Nat Methods, 2015. **12**(4): p. 357-60.

76.   Liao, Y., G.K. Smyth, and W. Shi, *featureCounts: an efficient general purpose program for assigning sequence reads to genomic features.* Bioinformatics, 2014. **30**(7): p. 923-30.

77.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550.

78. Mootha, V.K., et al., *PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes.* Nat Genet, 2003. **34**(3): p. 267-73.

79. Subramanian, A., et al., *Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles.* Proc Natl Acad Sci U S A, 2005. **102**(43): p. 15545-50.

80. Liberzon, A., et al., *The Molecular Signatures Database (MSigDB) hallmark gene set collection.* Cell Syst, 2015. **1**(6): p. 417-425.

81. Liberzon, A., et al., *Molecular signatures database (MSigDB) 3.0.* Bioinformatics, 2011. **27**(12): p. 1739-40.

82. Misra, J., et al., *Discordant regulation of eIF2 kinase GCN2 and mTORC1 during nutrient stress.* Nucleic Acids Res, 2021. **49**(10): p. 5726-5742.

83. Carlson, K.R., *tRNA_Charge-Seq.* https://github.com/carlsonkPhD/tRNA_Charge-Seq/, 2022.

84. Langmead, B. and S.L. Salzberg, *Fast gapped-read alignment with Bowtie 2.* Nat Methods, 2012. **9**(4): p. 357-9.

85. van Eenige, R., et al., *RandoMice, a novel, user-friendly randomization tool in animal research.* PLoS One, 2020. **15**(8): p. e0237096.

86. Montgomery, R.B., et al., *Maintenance of intratumoral androgens in metastatic prostate cancer: a mechanism for castration-resistant tumor growth.* Cancer Res, 2008. **68**(11): p. 4447-54.

Figure 1- Figure Supplement 1

A

B

LnCaP

MR49F

C4-2B

22Rv1

PC-3

LAPC4

Figure 1 - Figure Supplement 1. Knock-down of eIF2α kinases and ATF4 in PCa cell lines. (A) LNCaP cells were transfected with the indicated gene-specific siRNAs or a scramble control and the protein levels of HRI, PERK, or PKR and p-eIF2α, total eIF2α, and ATF4 were measured by immunoblot. Measurements of actin were used as a protein loading control in the immunoblot experiments. Molecular weight markers are shown in kilodaltons. **(B)** The indicated PCa cell lines were transfected with siRNAs targeting GCN2 or ATF4. Protein lysates were prepared and analyzed by immunoblot to determine the levels of GCN2, ATF4, or actin as indicated. ATF4 was not detected in the LAPC-4 cells.

Figure 1- Figure Supplement 2

Figure 1 - Figure Supplement 2. CRISPR/Cas9 knock-out of GCN2 in 22Rv1 and PC-3 cells. (A) Immunoblot analyses for 22Rv1, PC-3, and GCN2 KO clones. Protein lysates were analyzed by immunoblot to measure the levels of GCN2, ATF4, p-eIF2α, total eIF2α or actin. Molecular weight markers are shown in kilodaltons. The relative levels of p-eIF2α normalized to total eIF2α compared to wild type (WT) parental control are indicated. **(B)** Growth curve of 22Rv1 WT, 22Rv1 GCN2 KO (clone 7), or 22Rv1 GCN2 KO (clone 7) cells overexpressing GCN2. Data from replicate wells (N = 5) are shown as fold-change (mean ± SD) relative to day 0. Statistical significance was determined using a two-way ANOVA as described in **Supplementary File 1**; ****, $p \leq 0.0001$. **(C)** Protein lysates prepared from the 22Rv1 WT, 22Rv1 GCN2 KO (clone 7), or 22Rv1 GCN2 KO (clone 7) cells expressing GCN2 were analyzed by immunoblot for GCN2, ATF4, ASNS, or actin.

Figure 1- Figure Supplement 3

Figure 1 - Figure Supplement 3. Inhibition of the ISR by GCN2iB in PCa cell lines. (A)
C4-2B or 22Rv1 cells, cultured as indicated in the materials and methods, or PC-3 cells cultured
in HPLM media were treated with GCN2iB as indicated for up to 6 days. Cell growth was
measured (N = 5) and plotted as fold change (mean ± SD) relative to day 0. Statistical
significance was determined using a two-way ANOVA as described in **Supplementary File 1**;
****, $p \le 0.0001$. **(B)** 22Rv1 WT and 22Rv1 GCN2 KO (clone 7) cells were treated with GCN2iB
as indicated for up to 6 days. Cell growth was measured (N = 5) and plotted as fold change
(mean ± SD) relative to day 0. Statistical significance was determined using a two-way ANOVA
as described in **Supplementary File 1**; ****, $p \le 0.0001$. **(C)** Lysates were prepared from C4-2B,
22Rv1, or PC-3 cells treated with GCN2iB at the indicated concentrations or vehicle control
(DMSO) for 48 hours and immunoblot analysis was carried out using antibodies that recognize
p-GCN2-T899, total GCN2, p-eIF2α-S51, total eIF2α, ATF4, ASNS, TRIB3, LAT1 (SLC7A5),
xCT (SLC7A11), 4F2 (SLC3A2) AR, or actin. Molecular weight markers are indicated in
kilodaltons.

Figure 1- Figure Supplement 4

Figure 1 - Figure Supplement 4. Knock-down of ATF4 reduces growth in PCa cell lines.

ATF4 expression was reduced in LNCaP, MR49F, C4-2B or PC-3 cells using ATF4-specific siRNAs. Cell growth was measured in replicate wells (N = 5) for up to 6 days and is plotted as fold change (mean ± SD) relative to day 0. Statistical significance was determined using a two-way ANOVA as described in **Supplementary File 1**; ****, $p \leq 0.0001$.

Figure 1- Figure Supplement 5

Figure 1 - Figure Supplement 5. Inhibition of GCN2 in non-cancerous prostate cell line

BPH-1. (A) Lysates were prepared from BPH-1, LNCaP C4-2B, 22Rv1, or PC-3 cells and

immunoblot analysis was carried out using antibodies that recognize p-GCN2-T899, total

GCN2, p-eIF2α-S51, total eIF2α, ATF4, AR, or actin. Molecular weight markers are indicated in

kilodaltons. **(B)** BPH-1 cells were transfected with siRNAs targeting GCN2, ATF4 or 4F2

(SLC3A2). Protein lysates were prepared and analyzed by immunoblot to determine the levels

of GCN2, ATF4, 4F2 (SLC3A2) or actin as indicated. Molecular weight markers are indicated

in kilodaltons. **(C)** Expression of GCN2, ATF4, or 4F2 (SLC3A2) was reduced in BPH-1 cells

using two different gene-specific siRNAs as indicated and compared to a scramble siRNA

control. Cell growth was measured for up to 6 days in replicate wells (N = 5) as described in **A**.

Statistical significance was determined using a two-way ANOVA as described in

Supplementary File 1; *, $p \leq 0.05$; **, $p \leq 0.01$. **(D)** Lysates were prepared from BPH-1 cells

treated with GCN2iB at the indicated concentrations or vehicle control (DMSO) for 48 hours

and immunoblot analysis was carried out using antibodies that recognize p-GCN2-T899, total

GCN2, p-eIF2α-S51, total eIF2α, ATF4, ASNS, TRIB3, LAT1 (SLC7A5), xCT (SLC7A11), 4F2

(SLC3A2), AR, or actin. Molecular weight markers are indicated in kilodaltons. **(E)** BPH-1 cells

were treated with 0.5 – 10 µM GCN2iB or vehicle (DMSO) control as indicated for up to 6 days.

Cell growth was measured (N = 5) and plotted as fold change (mean ± SD) relative to day 0.

Statistical significance was determined using a two-way ANOVA is shown in **Supplementary**

File 1.

Figure 1- Figure Supplement 6

Non-malignant   Malignant

Non-malignant   Malignant

Figure 1 - Figure Supplement 6. p-GCN2 staining of prostate core needle specimens.

H&E and IHC staining using p-GCN2-T899 antibody in prostatic needle biopsy specimens

from patients with prostate cancer with Gleason scores 4+4=8 (top) or 4+5=9 (bottom).

Non-malignant (left) and malignant (right) prostate tissues are highlighted for both cases.

Figure 2 - Figure Supplement 1

A

B

Figure 2 - Figure Supplement 1. Inhibition of GCN2 in PCa cells reduces expression of SLC-genes involved in amino acid transport. (A) LNCaP cells were treated with 2 µM GCN2iB or vehicle control (DMSO) for 6 or 24 hours, protein lysates were prepared, and immunoblotted for the indicated proteins. The bar graphs show the relative levels of the indicated proteins normalized to actin. Statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD (N = 3); *, $p \leq 0.05$; ***, $p \leq 0.001$. **(B)** Immunoblot analysis of PC-3 WT, PC-3 GCN2 KO (clone C-2), and PC-3 GCN2 KO (clone C-3) lysates using antibodies that recognize GCN2, ATF4, ASNS, LAT1 (SLC7A5), xCT (SLC7A11), 4F2 (SLC3A2), ASNS or actin.

Figure 3 - Figure Supplement 1

Figure 3 - Figure Supplement 1. Amino acid measurements of LNCaP cells treated with vehicle (DMSO), 2 µM GCN2iB, or 2 µM GCN2iB + essential amino acid (EAA) for 48 hours. Bar graphs in the top panel show high abundance amino acids and the lower panel highlight those with lower levels. Statistical significance was determined using an unpaired two-tailed *t*-test. Error bars indicate SD (N=4); *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$; ****, $p \leq 0.0001$.

Figure 3 - Figure Supplement 2

Figure 3 - Figure Supplement 2. Effect of AA supplementation on growth of LNCaP cells.
LNCaP cells were treated with GCN2iB (2 µM) or vehicle control (DMSO) in standard growth media, media supplemented with non-essential amino acids (NEAA), supplemented with essential amino acids (EAA), or supplemented with histidine (200 µM), and cultured for up to six days. Cell growth was quantitated using CellTiter-Glo as described in the Material and Methods and is presented as fold-change (mean ± SD) relative to day 0. Statistical significance was determined using a two-way ANOVA as described in Supplementary File 1; ****, $p \le$ 0.0001.

Figure 3 - Figure Supplement 3

A

LNCaP

MR49F

C4-2B

22Rv1

PC-3

B

22Rv1

Figure 3 - Figure Supplement 3. Loss of GCN2 or ATF4 expression in PCa cell lines induces cell cycle arrest. (A) LNCaP, MR49F, C4-2B, 22Rv1, or PC-3 cells were transfected with GCN2 siRNA, ATF4 siRNA, or scrambled siRNA control for 48 hours and subjected to cell cycle analysis as described in the Material and Methods. The percentage of cells in G1, S, and G2-M are shown in the bar graphs. **(B)** 22Rv1 GCN2 KO and the parental WT were analyzed for cell cycle arrest. Statistical significance was determined using a one-way ANOVA with Tukey's multiple comparisons. Error bars indicate SD (N = 3); ***, $p \leq 0.001$; ****, $p \leq 0.0001$.

Figure 3 - Figure Supplement 4

A

B

Puromycin

Figure 3 - Figure Supplement 4. GCN2 inhibition and supplementation with EAA affect charging of tRNA^{HIS} and global translation. (A) LNCaP cells were treated with 2 μM GCN2iB or vehicle (DMSO) control in the absence or presence of EAA for 8 hours and tRNA^{His} charging levels were determined by qRT-PCR as described in the materials and methods. **(B)** LNCaP cells were treated with GCN2iB (2 μM) or vehicle (DMSO) control in the absence or presence of histidine supplementation. Puromycin (1 μM) was added to culture media 15 minutes prior to lysate preparation and puromycin incorporation was measured by immunoblot analysis using anti-puromycin antibody. Puromycin incorporation was quantified from replicate samples and is shown normalized to the vehicle control levels. Statistical significance was determined using a one-way ANOVA with Tukey's multiple comparisons. Error bars indicate SD (N = 3); ns, $p > .5$, **, $p \leq .1$.

Figure 3 - Figure Supplement 5

Figure 3 - Figure Supplement 5. Growth of MR49F and 22Rv1 can be rescued with amino acid supplements. (A) MR49F cells were treated with GCN2iB (2 µM) or vehicle control (DMSO) for 96 hours in normal growth media, growth media supplemented with EAA, or growth media supplemented with individual amino acids as indicated. Cell growth was quantified using CellTiter-Glo as described in the Material and Methods and is presented normalized to the vehicle control group. **(B)** 22Rv1 WT and 22Rv1 GCN2 KO cells were cultured for 96 hours in normal growth media, growth media supplemented with EAA, or growth media supplemented with individual amino acids as indicated and cell growth was similarly measured as described in **A**. Statistical significance in panels A and B was determined using a one-way ANOVA with Tukey's multiple comparisons. Error bars indicate SD (N = 3); ****, $p \leq .1$.

Figure 4 - Figure Supplement 1

LNCaP

22Rv1

Figure 4 - Figure Supplement 1. SLC-specific CRISPR/Cas9 KO library reveals that 4F2 (SLC3A2) is critical for cell fitness in LNCaP and 22Rv1 cells. LNCaP and 22Rv1 cells were transduced with a SLC KO CRISPR/Cas9 library targeting 394 human *SLC* genes and pseudogenes. Transduced cells were cultured for 3 – 4 weeks and sgRNA sequences were amplified by PCR and sequenced to determine the composition of the sgRNA library compared to Control (Day). Scatterplot of sgRNA counts (log1, normalized) for control (N = 3) vs. LNCaP (N = 3) and control (N = 3) vs. 22Rv1 (N = 3). Non-targeting sgRNAs (red), significant depleted sgRNAs (blue) ($p \leq .5$), and 4F2 (SLC3A2) sgRNAs (orange) are indicated.

Figure 4 - Figure Supplement 2

A

4F2 mRNA

ATF4 mRNA

B

4F2 mRNA

ATF4 mRNA

C

4F2 mRNA

ATF4 mRNA

Figure 4 - Figure Supplement 2. GCN2 regulates 4F2 (SLC3A2) mRNA levels in LNCaP cells. (A) 4F2 (SLC3A2) and ATF4 mRNA was measured by qRT-PCR as described in the Materials and Methods in LNCaP cells treated with 2 µM GCN2iB for 6 or 24 hours or vehicle control (DMSO), **(B)** cultured in standard culture conditions (NM: normal media), media supplemented with 200 µM histidine (+ His), or media depleted of histidine (- His) for 24 hours, or **(C)** treated with 100 nM halofuginone (HF) for 2 or 6 hours or untreated (DMSO control). Error bars indicate SD (N = 3). An unpaired two-tailed t-test was used to determine statistical significance; *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$; ****, ≤ 0.0001.

Figure 4- Figure Supplement 3

Figure 4 - Figure Supplement 3. GCN2 and the ISR control expression of 4F2 (SLC3A2) which provides a growth advantage in PCa cell lines. **(A)** LNCaP cells were treated with GCN2iB (2 µM) or vehicle (DMSO) control in the presence or absence of salubrinal (50 µM) for 48 hours. Protein lysates were prepared and analyzed by immunoblot using antibodies that recognize total or phosphorylated GCN2, total or phosphorylated eIF2α, ATF4, 4F2 (SLC3A2), or actin as indicated. **(B)** LNCaP cells transfected with empty vector (EV) control or pMSCV-GADD34-puro expression plasmid encoding the human GADD34 gene were analyzed by immunoblot as indicated in panel A. **(C)** Protein lysates prepared from LNCaP or 22Rv1 stably expressing empty vector (EV) control or 4F2 (SLC3A2) were analyzed by immunoblot using antibodies that recognize total or phosphorylated GCN2, total or phosphorylated eIF2α (S-51), ATF4, or actin as indicated. **(D)** Growth of LNCaP and 22Rv1 cells stably expressing empty vector (EV) control or 4F2 (SLC3A2) were measured in replicate wells (N = 5) for up to 4 days and plotted as fold change (mean ± SD) relative to day 0. Statistical significance was determined using a two-way ANOVA as described in **Supplementary File 1**; **, $p \leq 0.01$; ****, $p \leq 0.0001$.

Figure 6 - Figure Supplement 1

Figure 6 - Figure Supplement 1. Deletion of GCN2 reduces tumor growth and proliferation in 22Rv1 and PC-3 tumors. (A) Representative images for Ki-67, p-GCN2-T899, or 4F2 (SLC3A2) staining of 22Rv1 WT and 22Rv1 GCN2 KO tumors are shown. Bar graph shows quantification for percent positive nuclear staining (Ki67) or Histoscore (p-GCN2-T899 and 4F2). Statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD (N = 4); **, $p \le 0.01$; ****, $p \le 0.0001$. **(B)** PC-3 WT, and PC-3 GCN2 KO clone 2 (C-2), or PC-3 GCN2 KO clone 3 (C-3) cells were injected subcutaneously into the dorsal flank of mice as described in the materials and methods section. Tumor volumes were measured on the indicated days. Statistical significance was determined using a two-way ANOVA with Sidak's multiple comparison. Error bars indicate SEM (N = 5); *, $p \le 0.05$; ***, $p \le 0.001$; ****, $p \le 0.0001$. Bar graph indicates final tumor weight measured at endpoint. Statistical significance was determined using a one-way ANOVA with Tukey's multiple comparisons. Error bars indicate SD (N = 5); **, $p \le 0.01$. **(C)** Representative images for Ki-67, p-GCN2-T899, or 4F2 (SLC3A2) staining of PC-3 WT and PC-3 GCN2 KO tumors are shown. Bar graph shows quantification for percent positive nuclear staining (Ki67) or Histoscore (p-GCN2-T899 and 4F2). Statistical significance was determined using an unpaired two-tailed t-test. Error bars indicate SD (N = 5); *, $p \le 0.05$; **, $p \le 0.01$.

Figure 6 - Figure Supplement 2

Figure 6 - Figure Supplement 2. Deletion of GCN2 or ATF4 reduces the expression of SLC-genes associated with amino acid transport in 22Rv1 tumors. (A) Lysates from 22Rv1 WT, 22Rv1 GCN2 KO and 22Rv1 ATF4 KO tumors were subjected to immunoblot analyses to measure total or phosphorylated GCN2, total or phosphorylated eIF2α, ATF4, LAT1 (SLC7A5), xCT (SLC7A11), 4F2 (SLC3A2), CAT1 (SLC7A1), ASCT1 (SLC1A4), ASCT2 (SLC1A5), androgen receptor (AR), AR splice variant 7 (AR-V7), or actin. Molecular weight markers are indicated in kilodaltons for each immunoblot panel. Levels of the indicated proteins normalized to appropriate control are shown in the bar graph on the right. Statistical significance was determined using a one-way ANOVA with Tukey's multiple comparisons. Error bars indicate SD (N = 4); ns, $p > 0.05$; *, $p \le 0.05$; **, $p \le 0.01$; ***, $p \le 0.001$. **(B)** LNCaP cells were transfected with siRNAs targeting GCN2 (N = 4), ATF4 (N = 4), or scramble control (N = 8) for 48 hours. Amino acid levels were determined as described in the materials and methods sections. Error bars indicate SD. Statistical significance was determined using a one-way ANOVA with Tukey's multiple comparisons; #, $p \le 0.1$; *, $p \le 0.05$; **, $p \le 0.01$; ***, $p \le 0.001$; ****, $p \le 0.0001$. Scramble control and GCN2 knock-down samples are the same as in **Figure 4I**.

Figure 6 - Figure Supplement 3

Figure 6 - Figure Supplement 3. Effect of essential amino acid supplementation on mouse weight, amino acid levels, and expression of SLC-genes associated with amino acid transport in 22Rv1 WT and GCN2 KO tumors. (A) Body weight measurements from mice bearing 22Rv1 WT or 22Rv1 GCN2 KO tumors with or without EAA supplementation in the drinking water as described in **Figure 6G**. Statistical significance was determined using a two-way ANOVA with Sidak's multiple comparison, **, $p \le 0.01$. **(B)** Amino acid measurements for 22Rv1 WT, 22Rv1 GCN2 KO, or 22Rv1 GCN2 KO + EAA tumors. Bar graphs show only amino acids not present in EAA supplemented water. Error bars indicate SD (N = 4). Statistical significance was determined using a one-way ANOVA with Tukey's multiple comparisons; #, $p \le 0.1$; *, $p \le 0.05$. **(C)** Protein lysates were prepared from tumors described in **Figure 6G** and analyzed by immunoblot to measure total GCN2, LAT1 (SLC7A5), xCT (SLC7A11), 4F2 (SLC3A2), CAT1 (SLC7A1), ASCT1 (SLC1A4), ASNS, androgen receptor (AR), AR splice variant 7 (ARv7), or actin. Molecular weight markers are indicated in kilodaltons for each immunoblot panel. The levels of the indicated proteins normalized to actin are shown in the bar graphs. Statistical significance was determined using a one-way ANOVA with Tukey's multiple comparisons. Error bars indicate SD (N = 4); ns, $p > 0.05$; *, $p \le 0.05$; **, $p \le 0.01$; ***, $p \le 0.001$; ****, $p \le 0.0001$.

Figure 6 - Figure Supplement 4

Figure 6 - Figure Supplement 4. Effect of overexpression of 4F2 (SLC3A2) on amino acid levels in 22Rv1 WT and GCN2 KO tumors. (A) Amino acid measurements from 22Rv1 WT + EV, 22Rv1 GCN2 KO + EV, and 22Rv1 GCN2 KO + 4F2 (SLC3A2) tumors. Two separate bar graphs show high abundance (top) and low abundance (bottom) amino acids. Statistical significance was determined using a one-way ANOVA with Tukey's multiple comparisons. Error bars indicate SD (N = 4); #, $p \leq 0.1$; **, $p \leq 0.01$. **(B)** Lysates were prepared from these tumors in A were analyzed by immunoblot to measure total GCN2, 4F2 (SLC3A2), or actin. A long and short exposure for the 4F2 (SLC3A2) immunoblot is shown. Molecular weight markers are indicated in kilodaltons for each immunoblot panel.

Figure 7- Figure Supplement 1

Figure 7 - Figure Supplement 1. GCN2iB reduces proliferation in PCa cell line-derived and patient-derived xenograft models and has no effect on mouse body weight. Male NSG mice were injected subcutaneously with LNCaP (N = 5) or 22Rv1 (N = 4) cells, or implanted with tumor fragments from an androgen-sensitive tumor TM00298 (N = 5) or the castration-resistant LuCaP-35 CR (N = 5) tumor and treated as described in **Figure 7**. **(A)** Representative images showing IHC staining for Ki-67 or p-GCN2-T899 from tumor described above. Bar graphs show quantification for percent positive nuclear Ki-67 cells. Error bars indicate SD. Statistical significance was determined using an unpaired two-tailed *t*-test; *, $p \le 0.05$; **, $p \le 0.01$. **(B)** Mouse body weight was measured on indicated days for mice bearing LNCaP, 22Rv1, TM00298, or LuCaP35 CR tumors treated with vehicle or GCN2iB as described in **Figure 7**.

Figure 7- Figure Supplement 2

A

22Rv1 Tumors

p-GCN2-T899 — 250
GCN2 — 250
ATF4 — 50
LAT1 — 37
xCT — 37
4F2 — 100
Actin — 37

Vehicle   GCN2iB

B

TM00298 Tumors

p-GCN2-T899 — 250
GCN2 — 250
ATF4 — 50
LAT1 — 37
xCT — 37
4F2 — 100
Actin — 37

Vehicle   GCN2iB

ATF4
ATF4/Actin (Fold Change to Vehicle)
Vehicle   GCN2iB

LAT1
LAT1/Actin (Fold Change to Vehicle)
Vehicle   GCN2iB

xCT
xCT/Actin (Fold Change to Vehicle)
Vehicle   GCN2iB

4F2
4F2/Actin (Fold Change to Vehicle)
Vehicle   GCN2iB

Figure 7 - Figure Supplement 2. Treatment with GCN2 inhibitor, GCN2iB, reduces the expression of SLC-genes associated with amino acid transport in 22Rv1 and TM0 0 298 tumors. Immunoblot analysis of 22Rv1 (A) or TM298 (B) tumors from mice treated with GCN2iB as described in **Figure 7.** Molecular weight markers are indicated in kilodaltons for each immunoblot panel. The levels of the indicated proteins from TM298 tumors normalized to appropriate control are shown in the bar graph (N = 4).

Figure 7- Figure Supplement 3

Figure 3 - Figure Supplement 3. 4F2 (SLC3A2) protein levels are higher in malignant prostate cancer tissue compared to normal prostate tissue. (A) The protein levels of 4F2 (SLC3A2) were measured in a prostate tumor microarray (Biomax PR807c) using immunohistochemistry (IHC). Staining for 4F2 (SLC3A2) from normal prostate tissue (N = 10) and malignant prostate cancer tissue (N = 50) was analyzed and quantified using QuPath to determine the histoscore and is represented as a scatter plot. Statistical significance was determined using an unpaired two-tailed *t*-test; *, $p \leq 0.05$. Representative images showing 4F2 (SLC3A2) staining of normal and malignant prostate tissues are shown. Scale bars: 200 µm (main image) and 20 µm (insert). **(B)** H&E and IHC staining using p-GCN2-T899 or 4F2 (SLC3A2) antibody in prostate needle biopsy specimens from patients with high grade prostate cancer with Gleason scores 5+4=9 (top) or 4+5=9 (bottom).