Predation maintains virulence in the opportunistic pathogen Serratia marcescens

Heather A. Hopkins[1] & Kasie Raymann[1]*

- Department of Biology, University of North Carolina at Greensboro, Greensboro NC, 27412

*Kasie Raymann

Email: ktrayman@uncg.edu

Keywords: evolution, virulence, predation, bacteria, opportunistic pathogen

Abstract

Opportunistic pathogens are environmental microbes that are generally harmless and only

occasionally cause disease. Unlike obligate pathogens, the growth and survival of opportunistic

pathogens does not rely on host infection or transmission. Their versatile lifestyles make it challenging

to decipher how and why virulence has evolved in opportunistic pathogens. The Coincidental Evolution

Hypothesis (CEH) postulates that virulence results from exaptation or pleiotropy, i.e., traits evolved for

adaptation to living in one environment that have a different function in another. Several studies have

tested the CEH, but results have been inconclusive. Here we used experimental evolution to determine

how the presence of a predator impacts the virulence and fitness of a ubiquitous environmental

opportunistic bacterial pathogen that has acquired multi-drug resistance: *Serratia marcescens*. We

evolved *S. marcescens* in the presence or absence of a generalist protist predator, *Tetrahymena*

thermophila, and evaluated the genotypic and phenotypic changes that occurred after evolution.

Overall, we found that evolution in the presence of a predator maintained virulence, whereas evolution

in the absence of a predator resulted in attenuated virulence. Strains evolved in the presence or

absence of a predator presented different phenotypic traits, such as increased biofilm formation and

decreased predation resistance, respectively. Whole genome shotgun (WGS) sequencing of the

evolved populations and isolates revealed multiple cases of parallel evolution in genes associated with

virulence. Our results provide evidence that bacterial virulence and virulence related traits are

maintained by selective pressures imposed by protist predators, supporting the CEH.

Introduction

Opportunistic pathogens are environmental bacteria that typically coexist peacefully with hosts and only sometimes cause disease, e.g., in hosts with immunodeficiency or microbiome imbalance [1]. Opportunistic pathogens are intriguing because they have diverse lifestyles and experience strong tradeoffs between adaptation to life in the environment versus life inside of a host. Opportunistic pathogens are also conceptually challenging because, unlike obligate pathogens, their survival does not depend on their ability to infect a host: they are fully capable of reproducing and living in the environment [1,2]. Thus, the factors driving virulence in opportunistic pathogens remain largely enigmatic from an evolutionary standpoint. One hypothesis that has been proposed for opportunistic pathogen virulence is the Coincidental Evolution Hypothesis (CEH) [3,4]. The CEH postulates that virulence evolves indirectly due to selection that occurs in non-host environments [3,4]. Specifically, bacterial defense mechanisms against environmental predators (e.g., amoebae and protists) such as adherence, biofilm formation, digestion resistance, intracellular toxin production, and outer-membrane structures have been shown to play a role in host infection [1,5,6]. Additionally, some studies have supported the CEH by showing that pathogens challenged with microbial predators exhibited increased host virulence [7–12], and genes involved in pathogenesis were upregulated during predation [11].

Experimental evolution represents a powerful approach for evaluating how predation impacts virulence. To date, the limited studies that have experimentally evolved bacteria in the presence of a predator have generally concluded that predation attenuates virulence [13–17]. However, it is difficult to make broad generalizations from these studies as they have used incomparable and often inadequate methods to assess phenotypes (e.g., testing entire evolved populations, mixtures of evolved isolates, or using different individual isolates for various assays). Moreover, to our knowledge, only one bacteria-predator experimental evolution study investigated the genotypes of the evolved bacteria and the authors claim to have identified over 100 mutations within their predator-evolved populations after only

three days of evolution and ~30 mutations (most at low frequency) in individual isolates after 42 days

[17]. This high number of mutations [17] is highly unrealistic and is suggestive of contamination and/or

sequencing error. Indeed, mutation accumulation experiments of bacteria have reported a range of 10^{-8}

to 10^{-10} mutations/bp/generation, meaning that, on average, less than one mutation will arise in one

generation in the absence of negative selection [18] and in the Long-Term Evolution Experiment only

six mutations were observed in an evolved *Escherichia coli* clone (isolate) after 2,000 generations [19].

Therefore, the goal of our study was to experimentally evolve the opportunistic bacterial pathogen

Serratia marcescens in the presence or absence of a ubiquitous environmental protist predator,

Tetrahymena thermophila, and characterize both the genotypes and phenotypes of the evolved

bacteria.

S. marcescens is an opportunistic pathogen that ubiquitously occurs in soil and water [20] and

has been used for several previous bacteria-predator experimental evolution studies [13,14,16]. It is a

host generalist that can colonize many different organisms, where it can be pathogenic and/or

commensal [20]. In humans, *S. marcescens* is a common enteric bacterium that is generally harmless

in the gastrointestinal tract [21]. However, it was recently found to be capable of damaging gut epithelial

cells *in vitro* [22] and is responsible for causing dangerous nosocomial infections worldwide, notably in

the respiratory and urinary tracts [23,24]. In the last few decades, the frequency of nosocomial

infections caused by *S. marcescens* has increased, several hospital-wide outbreaks have been

reported, and most strains have acquired multidrug resistance [23,24]. Additionally, *S. marcescens* has

been recognized as an opportunistic pathogen of many plants and other animals [20]. Thus,

understanding the evolution of virulence in *S. marcescens* is important for the health of many

organisms.

Here, we evaluated whether experimental evolution of *S. marcescens* in the presence and

absence of a predator could be useful for identifying genes involved in virulence and elucidating how

predation impacts opportunistic pathogen evolution. Following 60 days of experimental evolution in the

presence or absence of *T. thermophila*, we sequenced random *S. marcescens* isolates from each

population as well as the entire evolved populations and characterized the mutations that occurred.

Using a single representative isolate from each evolved population we evaluated biofilm production,

growth, and predation resistance to measure fitness and performed survival assays to test virulence in

a natural animal host [25], the honey bee (*Apis mellifera*). Overall, our results revealed that virulence

and grazing resistance is attenuated in the absence of a predator and demonstrated that predation

plays a role in maintaining virulence in opportunistic pathogens, even over short evolutionary

timescales. Additionally, we observed multiple cases of parallel evolution which, when combined with

our phenotypic assays of isolates, allowed us to infer the outcome of some mutations. In conclusion,

our study supports the CEH and provides evidence that experimental evolution can be used to identify

genes involved in virulence and other phenotypes in *S. marcescens*.

Results

82
83   We evolved *S. marcescens* KZ19, isolated from honey bees (23), by serially passaging every

day for 60 days (in Neff media [26]) in the presence or absence of a ubiquitous generalist protist

predator *T. thermophila* (Figure 1). In our study we minimized predator/prey co-evolution by passaging

evolved *S. marcescens* to new axenic *T. thermophila* cultures each day. In total, we created three

media-evolved (ME) and six predator-evolved (PE) replicate lines (populations). The same

concentration of *S. marcescens* was transferred for each passage, and we confirmed successful

passaging via plating on LB agar every day. Viability and approximate population size of *T. thermophila*

was evaluated before every passage via light microscopy. We also confirmed that grazing was

occurring by co-culturing *T. thermophila* with transconjugant *S. marcescens* KZ19 containing an E2

crimson fluorescent protein with spectinomycin resistance [25,27] for 24 hours in Neff media with 180

ug/ml spectinomycin and then visualizing via fluorescent microscopy (Figure 1).

Figure 1. Confirmation of grazing activity. A) Phase contrast image of *T. thermophila* after 24h co-culture with fluorescently labeled *S. marcescens*, B) Cy5 filter image showing the E2 crimson labeled S. marcescens cells inside *T. thermophila*, and C) composite overlay of the phase contrast and fluorescent images. Images were taken at 20X on a Keyence BZ-X700 Series All-In-One Fluorescence Microscope and overlayed in ImageJ.

Following 60 days of evolution, the ancestral strain (n=1), PE populations (n=6), ME populations

(n=3), and isolates from the three ME (n=10) and six PE (n=20) populations were sequenced using

whole genome shotgun (WGS) sequencing. (Supplementary Table 1). We then characterized the

virulence and fitness of one isolate from each evolved population (ME=3 and PE=6).

Mutations in the predator- and media-evolved populations

All evolved populations and isolates were compared to the ancestral genome to identify

mutations. First, the sequencing reads of the ancestral strain were mapped to the reference KZ19

genome (23) to generate a consensus ancestral genome (average of 146X coverage; Supplemental

Table 1). We then corrected for false positives by mapping all the reads of the ancestral strain back to

the new consensus ancestral genome using BreSeq [28]. All mutations predicted via BreSeq [28] when

the ancestral reads were mapped to the ancestral consensus genome were considered false positives

if detected in the evolved populations (i.e., they most likely correspond to sequencing or assembly

issues in the ancestral genome).

We first analyzed the mutation patterns across each population. The total number of mutations

identified in each evolved population (ME n=3; PE n=6) ranged from 76 to 278 with an average of 85

mutations per population (Supplemental Table 1). The number of mutations present in a population was

not correlated with the average genome coverage (P=0.16, R^2=0.26; Figure 2A) or with the average

mutation frequency (P=0.18, R^2=0.25; Figure 2B). A lack of significant correlation between mutations

and genome coverage reveals that our sequencing depth was sufficient to characterize the populations.

Two PE populations that possessed over 230 mutations could be due to a lack of recent sweeps in

these population or the presence of hyper mutators. We did find a significant negative correlation

between average genome coverage and average mutations frequency (P=0.02, R^2=0.55; Figure 2C),

indicating that sequencing depth impacted the number of low frequency variants detected.

Figure 2. Comparison of **A)** the average genome coverage and the number of mutations detected, **B)** the number of mutations and the average mutation frequency, and **C)** the average mutations frequency and the average genome coverage for each evolved population.

In all evolved populations we identified nonsynonymous, synonymous, and intergenic mutations

and a few mutations in non-coding regions (tRNAs) were observed in two ME and two PE populations

(Figure 3A, Supplemental Dataset 1). Nonsynonymous mutations accounted for between 48-61% of the

identified mutations in all populations. To determine the percentage of nonsynonymous, synonymous,

intergenic, and non-coding mutations expected at random we introduced independent mutations into

the ancestral *S. marcescens* KZ19 genome by performing 10,000 independent simulations using a

Jukes & Cantor model. We found that the percentage of nonsynonymous mutations we detected in our

populations was between 11-24% less than the random expectation which is 72.5% (Figure 3A),

indicating that both environments imposed negative (purifying) selective pressure on *S. marcescens*

(i.e., deleterious mutations are being purged).

Figure 3. Characterization of the mutations present in the evolved populations. **A)** Percentage of nonsynonymous (NS), synonymous (S), intergenic (I), and non-coding (NC) mutations detected in each evolved population (PE and ME) and compared to the percentage of each mutation type expected at random (Expected). **B)** Number of parallel (*gene-* and *site-specific*) mutation events identified in only PE populations (n=48), only ME populations (n=5), or in both PE and ME (n=54) and the functional annotations of the genes in which parallel events occurred.

In addition to negative selection, we also found evidence that our populations were undergoing

positive selection based on the identification of numerous mutations either in the exact same position

(*site-specific* parallel evolution) or in the same gene (*gene-specific* parallel evolution) across

populations (Figure 3B, Supplemental Table 2). Parallel evolution events were considered if a mutation

was found in the same position or gene in independently evolved populations (ME, PE, or both). It is

important to note that none of the parallel evolution events occurred across all populations, confirming

that they were not present in the ancestral population. In total, 59 parallel evolution events occurred in

both ME and PE (PE+ME; n=54) or in only ME (ME; n=5) populations, suggesting that these mutations

are conferring adaptation to the Neff media. However, the parallel mutations only found in ME

populations are likely the result of lack of predation rather than adaptation to media. Most strikingly, 48

parallel evolution events were specific to the PE populations, indicating that these mutations/genes are

important for fitness in the presence of a predator. Of the total 107 parallel evolution events identified,

73% could be functionally classified (Figure 3B). Parallel mutations were identified in genes involved in

genetic information and processing (PE=19%, PE+ME=17%), signaling and cellular processes

(PE=15%, ME=40%, PE+ME=12%), environmental Information processing (PE=11%, ME=60%,

PE+ME=4%), and metabolism (PE=32%, PE+ME=33%).

Mutations in the predator- and media-evolved isolates

A total of 10 ME (3-4 per population) and 20 PE (1-5 per population) isolates were sequenced

from the evolved populations (Supplemental Table 3). All isolates were compared to the ancestral

genome to identify mutations using the same procedure applied to the entire populations (described

above and in the methods section).

At least one mutation was detected in each isolate, with a maximum of four and an average of

two mutations per isolate (Supplemental Table 3). We identified a total of 18 different mutations within

or upstream (in promoters) of 11 genes and 78% of the mutations (14/18) were present at a frequency

of >90% in their respective population (Supplemental Table 3). Most isolates from the same population

were clones or only differed by a single mutation, thus we chose one representative clone from each

population for further analysis (ME 1-3, and PE 1-6).

We plotted all the mutations identified in each ME and PE population and their observed

frequencies and found that our isolates possessed all the fixed (100% frequency) mutations present

within the population from which they were isolated (Figure 4A). Only three of the mutations present in

our isolates were detected at a frequency lower than 65% in their respective population (PE5: *yfiN,*

ME2: *barA*, and ME1: *rcsB*), but mutations at the exact same site or within the same gene occurred at

100% frequency in other populations (Figure 4A-B). In fact, we identified three instances of *site-specific*

parallel evolution and two cases of *gene-specific* parallel evolution across the isolates.

Figure 4. Mutations present in evolved isolates. **A)** Plot of all the detected mutations and their frequencies in each population highlighting the mutations present in each representative isolate (circles or triangles). **B)** Mutations present in each isolate grouped by nonsynonymous (NS), synonymous (S), promoter (intergenic mutation in a promotor region) and indicating the mutation type (single nucleotide variant (SNV) or deletion), the gene (or gene upstream) affected, and the functional category of the gene (left color boxes). Gray shading of a row highlights parallel evolution events (*site-* or *gene-specific*) and solid shapes denote cases of *site-specific* parallel evolution. **C)** Pie graph showing the functional classification of the genes (or genes upstream) impacted by mutation events in the individual isolates and the percentage belonging to each category.

The most striking case of *site-specific* parallel evolution was a 66 bp in-frame deletion in the

response regulator *rcsB* [29] present in the ME1 and ME3 isolates (Figure 4B). The two other *site-*

specific parallel evolution events occurred in promoters directly upstream of 1) a methyl-accepting

chemotaxis sensor/transducer protein in the ME2 and ME3 isolates and 2) the outer membrane porin

OmpC protein in the PE4 and PE6 isolates (Figure 4B). One of the *gene-specific* parallel evolution

events occurred in a gene annotated as *barA* (a signal transduction histidine-protein kinase) in the

ME2, ME3, PE2, and PE4 isolates (Figure 4B). The second case of *gene-specific* parallel evolution

arose in a gene annotated as the inner membrane protein *yfiN* in all PE isolates except PE1 and PE3

(Figure 4B). Cases of parallel evolution found across PE and ME lines suggest adaptation to the media,

whereas parallel evolution events unique to either PE or ME lines suggest adaptation to the presence

or to the absence of predation, respectively.

We were able to functionally characterize 73% (8/11) of the genes (or genes adjacent to the

promoter region) in which mutations occurred (Figure 4B-C). The eight characterized genes belonged

to four functional classes: environmental information processing (45%), genetic information processing

(9%), energy metabolism (9%), and signaling and cellular processes (9%) (Figure 4C). Several genes

in which we observed mutations have been associated with virulence factors in *S. marcescens* or

related enterobacteria, such as *yfiN* [30,31], *rcsB* [32,33], and *barA* and its response regulator *uvrY*

[34–36]. Additionally, all intergenic mutations identified in the isolates are located within promoters

upstream of five different genes (Figure 4B), some of which have been associated with biofilm

formation (HmsT protein [37]) and chemotaxis (Methyl-accepting chemotaxis sensor/transducer [38]).

Virulence of predator- and media-evolved strains

S. marcescens KZ19 is pathogenic in honey bees under certain circumstances, e.g., following

antibiotic treatment, when exposed to high doses, or when injected into the hemolymph [25]. Thus, we

determined how predation impacts the virulence of *S. marcescens* by performing survival assays in

honey bees via high dose exposure. We tested the virulence of one isolate from each predator-evolved

population (n=6) and one isolate from each media-evolved population (n=3) by comparing them to the

virulence of the ancestral strain (Figure 5A-B). Three assays (biological replicates) were done with five

technical replicates each, totaling 300 bees per isolate.

Figure 5. Virulence and fitness of evolved isolates. **A-B)** Probability of survival of honey bees exposed to the ancestral strain (black and gray) and the **A)** ME isolates and **B)** PE isolates. Control bees were given sterile sugar water only (white squares). Mortality was recorded each day for 5 days. Graphs are based on three replicate assays per isolate (n=300 bees per isolate). * = P<0.05, *** = P<0.0005; Mantel-Cox Log-rank test with Bonferroni correction. **C-D)** Mean growth rate of each variant based on 600nm OD readings every hour for 24h of **C)** ME and **D)** PE isolates in spent Neff media compared to the ancestral strain (black). **E)** Mean biofilm formation of ME and PE isolates and the ancestral strain based on 560nm OD readings after 36h of growth in Neff media. Error bars show the standard deviation between triplicates. * = P<0.05; *t*-test with Bonferroni correction. **F)** Mean predation resistance of PE and ME isolates and the ancestral strain based on 600nm OD readings of bacterial filtrate after 48h in co-culture with *T. thermophila*. Error bars show the standard deviation between triplicates. * = P<0.05, ** = P<0.005; *t*-test with Bonferroni correction.

All the ME isolates displayed attenuated virulence when compared to the ancestral strain, with

2/3 being significantly less virulent (Figure 5A; $P<0.001$, Mantel-Cox Log-rank test with Bonferroni

correction). Conversely, none of the PE isolates significantly differed from the ancestor in terms of

virulence, except for PE4 which was statistically less virulent than the ancestral strain (Figure 5B;

$P=0.007$, Mantel-Cox Log-rank test with Bonferroni correction). Overall, results show that in the

absence of predation, virulence is attenuated, and predation maintains virulence in *S. marcescens*.

Fitness of predator- and media-evolved strains

To test the fitness of the PE and ME strains, we performed growth, biofilm formation, and predation

resistance (population size after growth in the presence of *T. thermophila*) assays. For growth rate, we

conducted *in vitro* assays using three different media types A) spent Neff media (filtered media that had

supported *T. thermophila* growth for 24h; Figure 5C-D), B) fresh Neff media and C) LB broth. In all

media types, virtually all the ME and PE isolates displayed slightly slower growth than the ancestor over

24h, but in general, the growth of the PE variants was more attenuated than the ME variants (Figure

5C-D, Supplemental Figure S1). This pattern suggests a tradeoff between growth rate and resistance to

predation.

Because biofilm formation is a strategy for resistance to predation and is also involved in

virulence, we measured biofilm production in our ME and PE variants compared to the ancestor (Figure

5E). Overall, the ME isolates did not differ ($P=0.22$, Kruskal-Wallis test) from the ancestor in terms of

biofilm production, whereas the PE isolates produced significantly more biofilms than the ancestor

($P=0.01$, Kruskal-Wallis test). In fact, biofilm production increased between 1.5- to 2.3-fold across PE

isolates relative to the ancestor (Figure 5E), indicating that they are producing more biofilms due to

predation. However, only two of the PE isolates were found to significantly produce more biofilms than

the ancestor when tested individually and corrected for multiple comparisons (PE4 $P=0.03$, PE6

$P=0.02$, *t*-test with Bonferroni correction).

Lastly, we evaluated predation resistance (i.e., growth with predator) of the ME and PE isolates.

Each isolate was grown with *T. thermophila* for 48h and then the population density of the bacteria was

measured (Figure 4F). In general, predation resistance of the PE isolates significantly differed from that

of the ancestor (P=0.01, Kruskal-Wallis). However, predation resistance across PE isolates differed

with some being similar to the ancestor (PE1 P=2.25 and PE2 P=1.8), whereas others were more

resistant (PE3 P= 0.026, PE5 P= 0.0063, and PE6 P= 0.0135) or less resistant (PE4 P= 0.0099) to

predation relative to the ancestor (Figure 4F; t-test with Bonferroni correction). Overall, the ME isolates

were significantly less resistant to predation than the ancestor (Figure 4F; P<0.0001; Kruskal-Wallis

test). All ME isolates were found to be significantly more susceptible to predation than the ancestor

when tested individually and corrected for multiple comparisons (Figure 4F; ME1 P= 0.001, ME2 P=

0.007, ME3 P= 0.001, t-test with Bonferroni correction). Results suggest that lack of predation

decreases grazing resistance while adaptation to predation increases it. Findings also indicate that the

PE lines followed different evolutionary paths over the 60 days of evolution.

Discussion

Evolution of *S. marcescens* in the absence of a predator resulted in attenuated virulence when

compared to the ancestral strain. Conversely, *S. marcescens* that were evolved in the presence of a

predator were not more virulent than the ancestral strain. Thus, our results indicate that virulence is

quickly attenuated in the absence of predation but is maintained in the presence of predation over short

evolutionary time scales. Maintenance rather than increased virulence following evolution with a

predator could be due to the fact that the ancestral strain (KZ19) is a natural opportunistic pathogen of

honey bees [25] and may have limited potential to become more virulent, especially during only 60 days

of evolution and because of the protection provided by the native gut microbes [25,39].

We found that the growth rate of PE and ME isolates did not substantially differ from the

ancestor. Nevertheless, growth rate of the PE isolates was systematically lower than the ME isolates,

likely revealing some fitness tradeoffs involved in adaptation to predation. Biofilm formation was

increased in the PE isolates when compared to the ancestor and the ME isolates. Biofilm production is

a natural defense mechanism against predation and can also play a role in pathogenicity [40]. Only half

of the PE isolates displayed a larger population size than the ancestor after 48h of co-culture with *T.*

thermophila and one PE isolate, PE4, was significantly less resistant to predation than the ancestral

strain, revealing that evolution in the presence of a predator does not always result in increased

resistance to predation. However, the predation-susceptible PE4 isolate also showed attenuated

virulence in honey bees, suggesting a link between predation resistance and virulence. We found that

all ME isolates were significantly less fit in the presence of a predator than the ancestor, indicating that

evolution in the absence of predation reduces fitness in the presence of a predator. Moreover, the ME

isolates that displayed attenuated virulence were also the most susceptible to predation (i.e., had the

smallest population size after co-culture with *T. thermophila*). Together these results imply that

virulence and predation resistance are governed by the same mechanisms. It is important to note that

we observed phenotypic differences after only 60 days of evolution and with only one to four identified

mutations per isolate.

Several different mutations, including some in genes associated with virulence, arose in the ME

and PE isolates. Most mutations observed were non-synonymous or located in promoters, one

synonymous mutation was also identified. Multiple cases of parallel evolution (site-specific and gene-

specific) were observed, indicating that some genes are under strong selection in the presence and/or

absence of a predator. With the exception of one synonymous mutation in *yfiN* in the PE6 isolate, most

instances of parallel evolution involved non-synonymous or intergenic mutations within promoters, both

of which likely impact gene expression and/or function. Though generally thought to be "silent", recent

studies have revealed that synonymous mutations can have an impact on gene expression [41,42].

Therefore, the synonymous mutation that occurred in *yfiN* could have an impact on fitness, especially

because it was fixed in the PE6 population. In fact, virtually all of the parallel mutations observed in the

isolates were fixed within their respective populations.

A gene-specific parallel evolution event occurred in a gene annotated as *barA* in both ME and

PE isolates. In *E. coli*, *barA* encodes a protein part of a two-component system which plays an

important role in the ability to switch between different carbon sources [34]. Additionally, orthologs of

barA have been associated with virulence in several Gram-negative pathogens [34–36]. The mutation

in *barA* present in PE4 most likely resulted in a loss of function since it involved a 1bp frameshift

deletion. In other isolates, it is likely that the mutations altered the function or expression of *barA* since

they consisted of a nonsynonymous SNVs or in-frame deletions. Based on the evidence that *barA* is

involved in a carbon storage regulatory system in *E. coli* [34] and that mutations in this gene arose in

both PE and ME isolates, we predict that they are the result of adaptation to the Neff media, which

consists of proteose peptone, yeast extract, and glucose.

We also identified cases of parallel evolution that were specific to either ME or PE variants,

signifying that they are likely due to adaptation to the presence or to the absence of a predator rather

than the media. For example, in the isolates from ME Lines 1 and 3, the same 66bp in-frame deletion

(site-specific parallel evolution) occurred in a gene annotated as *rcsB*. RcsB is the response regulator

of the Rcs phosphorelay two-component signal transduction system in enterobacteria [32]. RcsB has

been associated with the synthesis of biofilms, capsules, and flagella and antibiotic resistance [32].

Isolates that contained the deletion in *rcsB* displayed significantly attenuated virulence and had much

lower population sizes after co-culture with *T. thermophila* than the ancestral strain, indicating that *rcsB*

plays a role in predation resistance and pathogenicity in *S. marcescens*. One other case of site-specific

parallel evolution found in ME2 and ME3 isolates was an intergenic mutation in the promoter upstream

of a methyl-accepting chemotaxis sensor/transducer protein. Methyl-accepting chemotaxis proteins are

used to sense and respond to environmental stimuli and are involved in the synthesis of biofilms,

flagella, and toxins in many pathogens [38]. The fact that this mutation was in a promoter and found in

isolates from different evolved lines suggests that it confers a fitness benefit in the absence of

predators, potentially by saving resources by reducing the expression or protein complexes.

In most of the PE isolates, a non-synonymous (PE2, PE4, PE5) and synonymous (PE6) SNV

was found in a gene encoding the inner membrane protein YfiN (gene-specific parallel evolution). YfiN

has been implicated in biofilm formation, inhibiting motility, and cell division arrest under stressful

conditions [30,31]. The fact that mutations arose in *yfiN* independently across different PE populations

and were often fixed (100% frequency) implies that this gene is under strong selective pressure in the

presence of predators. Mutations in *yfiN* were present in four of the six PE isolates and we speculate

that they are responsible for the observed increase in biofilm production. The PE variants that did not

acquire a mutation in *yfiN* either contained a mutation in a promoter directly upstream of 1) a

hypothetical protein (unknown function) or 2) the inner membrane protein HmsT. HmsT is associated

with biofilm formation in *Yersinia pestis* [37], one of the closest known relatives of *S. marcescens*.

Although only a speculation, we predict that the intergenic mutations in the promoter regions upstream

of HmsT and the hypothetical protein are involved in increasing biofilm formation in the PE1 and PE3,

respectively, as these were the sole mutations present in these isolates. In addition, PE3 displayed

significantly increased predation resistance, suggesting that the mutation in the promoter region

upstream of the hypothetical protein plays a role in defense against predators.

In conclusion, our results suggest that predation helps maintain virulence in opportunistic

pathogens, thereby supporting the CEH. We identified several instances of parallel evolution across

evolved populations, which, when combined with our phenotypic assays of isolates, allowed us to infer

the outcome of some mutations. For example, we found evidence that *barA* Is involved in carbon

utilization in *S. marcescens*. We also predict that the mutations we observed in *yfiN* and two promoters

(upstream of HmsT and of a hypothetical protein) are responsible for increased biofilm production.

Moreover, our results imply that RscB plays a major role in predation resistance and virulence in *S.*

marcescens. Fundamentally, our study demonstrates the efficacy of using experimental evolution to

identify virulence mechanisms, even over short evolutionary timescales. Surprisingly, we found that

evolution in the absence of a predator yielded the most informative results for identifying factors

associated with virulence. Performing long-term evolutionary experiments under different scenarios with

additional replicate populations will provide a more accurate picture of the forces that drive the

evolution of opportunistic pathogens and the mechanisms responsible for their virulence. Identifying

virulence mechanisms and understanding how virulence evolves can help guide the development of

more effective treatment strategies for combating bacterial infections.

Materials and Methods

Microscopy

To confirm that *T. thermophila* grazes on *S. marcescens* KZ19 we utilized transconjugant KZ19

containing an E2 crimson fluorescent protein with spectinomycin resistance. The transconjugant KZ19

strain was inoculated into 3mL of LB broth containing 180ug/mL spectinomycin and incubated at 30°C

for 24h. After 24h, 1OD of the transconjugant KZ19 was diluted to 10^{-5} in Neff media and 1mL was

pipetted into 3mL of Neff media containing *T. thermophila* and 180ug/mL spectinomycin. The culture

was incubated for 24h at 30°C. One mL of the 24h co-culture was then centrifuged at 2000g for 10m to

collect the *T. thermophila* cells, the supernatant containing bacteria was discarded, and the *T.*

thermophila were resuspended in 1mL of axenic Neff media. The *T. thermophila* cells were then

pipetted onto a microscope slide that was washed with 60% EtOH (which immobilizes the cells without

immediately lysing them), covered with a glass cover slip, and imaged on a Keyence BZ-X700 Series

All-In-One Fluorescence Microscope in brightfield and with a Cy5 filter.

Experimental evolution

Serratia marcescens strain KZ19 was evolved either in media alone (media-evolved) or with the

predator *Tetrahymena thermophila* strain SB210 (predator-evolved). *T. thermophila* strain SB210 was

purchased from the Tetrayhema Stock Center located at Cornell University. The media-evolved (ME)

experiment included three replicate lines and the predator-evolved (PE) experiment included six

replicate lines. First, *S. marcescens* and *T. thermophila* were cultured for 24h at 30°C in Neff media. To

start each evolved line, 1OD of the overnight KZ19 culture was diluted to 10^{-5} in fresh Neff media and

1mL was added to either 3mL of Neff media [26] alone (ME lines) or 3mL Neff media containing equal

concentrations of 24h *T. thermophila* culture (PE lines). The starting ratio of S. marcescens:*T.*

thermophila was based on preliminary trials starting with various concentrations of *S. marcescens* and

T. thermophila co-inoculated in Neff media to determine the ratio that would allow both organisms to

remain in co-culture for 24h.

The experimental cultures were incubated at 30°C without shaking. Every 24h for 60 days the

cultures were vortexed and 10μL of each culture was sampled from the flasks and added to 3mL of

either 1) fresh Neff media or 2) new axenic cultures of *T. thermophila* in Neff media. Every day, prior to

passaging, the cultures were checked for *T. thermophila* viability and density via microscopy and the

successful passaging and growth of *S. marcescens* presence was evaluated by plating on Luria-Bertani

(LB) agar. Every day over the course of the 60-day evolution experiment *T. thermophila* and *S.*

marcescens were always detected and we never observed a noticeable difference in population size

based on plating (*S. marcescens*) and light microscopy (*T. thermophila*).

On day 60, six LB agar plates were streak inoculated from the culture flasks and incubated for

24h at 30°C. Twenty bacterial colonies were randomly selected from the PE lines (1-5 per population)

and ten bacterial colonies were randomly selected from the ME lines (3-4 per population). Pure cultures

of each isolate (30 total) were created by inoculating in 3mL LB broth with 24h incubation at 30°C.

Samples (200μL from each pure culture) were then frozen at -80°C in 20% glycerol (*T. thermophila* do

not survive freezing). These 30 isolates were sequenced, analyzed, and one isolate from each evolved

population (ME n=3, and PE n=6) was used for phenotypic analysis (i.e., virulence, growth, biofilm

production, and predation resistance).

Virulence assays

Pure 1OD cultures of the evolved isolates and the ancestor (10 total) were pelleted via

centrifugation and resuspended in a 1:1 in sterile sugar syrup solution (SSS). Conventional adult honey

bee workers (*Apis mellifera*) were sampled from a single hive located on the Gateway North Research

Campus in Browns Summit, North Carolina. Bees were immobilized at 4C, randomly distributed into

groups, and exposed via the immersion method (23) to ~10μL of one of three treatments: 1) sterile SSS

only, 2) the ancestor in sterile SSS, or 3) each evolved isolate in sterile SSS. Bees were kept in cup

cages (20 bees/cup; 100 bees/treatment) under hive conditions: 35°C and 95% humidity. Mortality

between the treatments was noted every day for 5 days. Assays were performed in triplicate with 300

bees tested in total per isolate. Survival curves (Kaplan-Meier) were created in GraphPad Prism version

9.1.0.

Fitness assays

Growth assays were performed in triplicate in both LB broth, fresh Neff media, and spent Neff media

with *T. thermophila* filtered out (i.e., *T. thermophila* were present in the media for 24h before filtering).

Pure 1OD cultures of the evolved isolates and the ancestor were diluted to 10^{-5} and 5μL of the culture

was pipetted into 200μL of media in a 96-well microplate. Absorbance detection was performed at an

OD of 600nm at 30°C with shaking every hour for 24h on a BioTek Synergy™ 2 plate reader.

Biofilm assays were performed in triplicate for each isolate. Pure 1OD cultures of the evolved

isolates and the ancestor were diluted to 10^{-5} and 5μL of the culture was pipetted into 200μL of media

in a 96-well microplate. The covered plate was incubated for 36h at 30°C without shaking. After 36h,

the supernatant was poured off, the wells were washed twice with 200μL of sterile ddH$_2$O, and the cells

were stained with 150μL of 0.04% crystal violet for 10 minutes. After 10 minutes, the excess crystal

violet was removed, the wells were washed again and stained once more for 10 minutes. The crystal

violet was poured off and cells were solubilized with 150μL of 95% ethanol. OD readings were taken at

absorbance reading 560nm on the BioTek Synergy™ 2 plate reader.

Predation resistance assays were performed in triplicate for each isolate. Pure 1OD cultures of

the evolved isolates and the ancestor were diluted to 10^{-5} and 1mL was added to a culture flask

containing *T. thermophila* in 3mL Neff media. *T. thermophila* alone in 3mL Neff media served as the

negative control. Experimental culture flasks were incubated at 30°C for 48h. After 48h, the cultures

were mixed thoroughly and 1mL was sampled and centrifuged at 5,000 rpm for 10 minutes. The

supernatant was discarded, pellets were resuspended in 1mL 1X PBS, and the solution was filtered

through 5.0 μm MilliporeSigma™ Millex™-SV Sterile PVDF syringe filters to remove the *T. thermophila*

cells. OD readings were taken of filtrates at 600nm after blanking with the negative control filtrate.

DNA Extraction and Sequencing

Cultures of the evolved populations (ME n=3, PE n=6), evolved isolates (ME n=10, PE n=20) and the

ancestor (n=1) were diluted to 1 OD. DNA extractions were performed using the Zymo Quick-DNA

Fungal/Bacterial Miniprep Kit (D6005). The DNA was then prepared for Whole Genome Shotgun

(WGS) sequencing using the Illumina Nextera DNA Flex Library Prep Kit (20018704). Genomes were

sequenced in-house on an Illumina iSeq 100 (2 x 150bp).

Genome Assembly and Analysis

All sequencing reads were demultiplexed using the Illumina BaseSpace Sequence Hub. The ancestral

genome was first mapped to the *S. marcescens* KZ19 (GCA_002915435.1) reference genome [25]. In

brief, the raw sequencing reads were trimmed using BBDuk (sourceforge.net/projects/bbmap/)

implemented in Geneious Prime, using default settings. Trimmed reads were then assembled by

mapping to the ancestral genome in Geneious Prime using Bowtie 2 [43] with default settings.

Assembled contigs were annotated using RAST [44]. We also mapped all the reads of the ancestral

strain back to the consensus ancestral genome using BreSeq [28]. Any mutations predicted by BreSeq

when the ancestral reads were mapped to the ancestral consensus genome were considered

sequencing or assembly issues in the ancestral genome and removed as candidate mutations in the

evolved populations and isolates.

The sequencing reads of the evolved populations and isolates were trimmed using Trimmomatic

(46). The trimmed reads were then mapped to the ancestral genome using BreSeq version 0.37.0 [28]

with a base-quality-cutoff PHRED score of 30 to identify mutations. Promoters were predicted using

iProEP [45]. Functional characterization of genes (Figure 2- Table Supplement 2) was predicted using

KOALA KEGG Orthology And Links Annotation (48). Simulations to determine the percent mutations

expected at random across intergenic, synonymous and non-synonymous positions were conducted

with a custom Python script: we conducted 10,000 independent simulations where one mutation was

introduced at random under a Jukes & Cantor model in the refence genome of *S. marcescens* KZ19

(GCA_002915435.1). The annotations of the genome were used to infer whether the mutation was

intergenic or not. Mutations affecting a coding sequence were inferred as synonymous or non-

synonymous by translating the gene *in silico* before and after introducing the mutation.

Materials availability

All raw sequencing data has been deposited on NCBI under BioProject PRJNA838621.

Acknowledgments

We would like to thank Dr. Louis-Marie Bobay and Dr. Joseph Santin for constructive and helpful

feedback on the manuscript. This work was supported by the National Science Foundation under grant

DEB-1930776 (to K.R.), the University of North Carolina Greensboro (UNCG) College of Arts and

Sciences Faculty First Award (to K.R.), and UNCG Department of Biology Graduate Student Support

grants (to H.A.H.).

Author Contributions

Competing Interest Statement

433   The authors declare no completing interests.

References

1.   Brown SP, Cornforth DM, Mideo N. Evolution of virulence in opportunistic pathogens: generalism,
plasticity, and control. Trends in Microbiology. 2012;20: 336–342. doi:10.1016/j.tim.2012.04.005

2.   Casadevall A, Pirofski L. Accidental Virulence, Cryptic Pathogenesis, Martians, Lost Hosts, and the
Pathogenicity of Environmental Microbes. Eukaryot Cell. 2007;6: 2169–2174. doi:10.1128/EC.00308-07

3.   Levin BR, Edén CS. Selection and evolution of virulence in bacteria: an ecumenical excursion and modest
suggestion. Parasitology. 1990;100: S103–S115. doi:10.1017/S0031182000073054

4.   Levin B. The Evolution and Maintenance of Virulence in Microparasites. Emerg Infect Dis. 1996;2: 93–102.
doi:10.3201/eid0202.960203

5.   Erken M, Lutz C, McDougald D. The Rise of Pathogens: Predation as a Factor Driving the Evolution of
Human Pathogens in the Environment. Microb Ecol. 2013;65: 860–868. doi:10.1007/s00248-013-0189-0

6.   Amaro F, Martín-González A. Microbial warfare in the wild—the impact of protists on the evolution and
virulence of bacterial pathogens. Int Microbiol. 2021;24: 559–571. doi:10.1007/s10123-021-00192-y

7.   Coombes BK, Gilmour MW, Goodman CD. The Evolution of Virulence in Non-O157 Shiga Toxin-Producing
Escherichia Coli. Front Microbio. 2011;2. doi:10.3389/fmicb.2011.00090

8.   Cirillo JD, Cirillo SLG, Yan L, Bermudez LE, Falkow S, Tompkins LS. Intracellular Growth in *Acanthamoeba*
castellanii Affects Monocyte Entry Mechanisms and Enhances Virulence of *Legionella pneumophila*.
Barbieri JT, editor. Infect Immun. 1999;67: 4427–4434. doi:10.1128/IAI.67.9.4427-4434.1999

9.   Hosseinidoust Z, van de Ven TGM, Tufenkji N. Evolution of Pseudomonas aeruginosa Virulence as a Result
of Phage Predation. Appl Environ Microbiol. 2013;79: 6110–6116. doi:10.1128/AEM.01421-13

10. Rasmussen MA, Carlson SA, Franklin SK, McCuddin ZP, Wu MT, Sharma VK. Exposure to Rumen Protozoa
Leads to Enhancement of Pathogenicity of and Invasion by Multiple-Antibiotic-Resistant *Salmonella*
enterica Bearing SGI1. Infect Immun. 2005;73: 4668–4675. doi:10.1128/IAI.73.8.4668-4675.2005

11. Rehfuss MYM, Parker CT, Brandl MT. Salmonella transcriptional signature in Tetrahymena phagosomes and
role of acid tolerance in passage through the protist. ISME J. 2011;5: 262–273. doi:10.1038/ismej.2010.128

12. Adiba S, Nizak C, van Baalen M, Denamur E, Depaulis F. From Grazing Resistance to Pathogenesis: The
Coincidental Evolution of Virulence Factors. Ahmed N, editor. PLoS ONE. 2010;5: e11882.
doi:10.1371/journal.pone.0011882

13. Friman V-P, Lindstedt C, Hiltunen T, Laakso J, Mappes J. Predation on Multiple Trophic Levels Shapes the
Evolution of Pathogen Virulence. Ahmed N, editor. PLoS ONE. 2009;4: e6761.
doi:10.1371/journal.pone.0006761

14. Mikonranta L, Friman V-P, Laakso J. Life History Trade-Offs and Relaxed Selection Can Decrease Bacterial
Virulence in Environmental Reservoirs. Vinatzer BA, editor. PLoS ONE. 2012;7: e43801.
doi:10.1371/journal.pone.0043801

15. Friman V-P, Buckling A. Phages can constrain protist predation-driven attenuation of Pseudomonas
aeruginosa virulence in multienemy communities. ISME J. 2014;8: 1820–1830. doi:10.1038/ismej.2014.40

16. Zhang J, Ketola T, Örmälä-Odegrip A-M, Mappes J, Laakso J. Coincidental Loss of Bacterial Virulence in
Multi-Enemy Microbial Communities. Vinatzer BA, editor. PLoS ONE. 2014;9: e111871.
doi:10.1371/journal.pone.0111871

17. Leong W, Poh WH, Williams J, Lutz C, Hoque MM, Poh YH, et al. Adaptation to an Amoeba Host Leads to
Pseudomonas aeruginosa Isolates with Attenuated Virulence. Nikel PI, editor. Appl Environ Microbiol.
2022;88: e02322-21. doi:10.1128/aem.02322-21

18. Sung W, Ackerman MS, Miller SF, Doak TG, Lynch M. Drift-barrier hypothesis and mutation-rate evolution.
Proc Natl Acad Sci USA. 2012;109: 18488–18492. doi:10.1073/pnas.1216223109

19. Barrick JE, Yu DS, Yoon SH, Jeong H, Oh TK, Schneider D, et al. Genome evolution and adaptation in a long-
term experiment with Escherichia coli. Nature. 2009;461: 1243–1247. doi:10.1038/nature08480

20. Grimont PAD, Grimont F. The Genus Serratia. Annu Rev Microbiol. 1978;32: 221–248.
doi:10.1146/annurev.mi.32.100178.001253

21. Almuneef MA, Baltimore RS, Farrel PA, Reagan-Cirincione P, Dembry LM. Molecular Typing Demonstrating
Transmission of Gram-Negative Rods in a Neonatal Intensive Care Unit in the Absence of a Recognized
Epidemic. Clinical Infectious Diseases. 2001;32: 220–227. doi:10.1086/318477

22. Ochieng JB, Boisen N, Lindsay B, Santiago A, Ouma C, Ombok M, et al. *Serratia marcescens* is injurious to
intestinal epithelial cells. Gut Microbes. 2014;5: 729–736. doi:10.4161/19490976.2014.972223

23. Hejazi A, Falkiner FR. Serratia marcescens. Journal of Medical Microbiology. 1997;46: 903–912.
doi:10.1099/00222615-46-11-903

24. Khanna A. Serratia Marcescens - A Rare Opportunis - tic Nosocomial Pathogen and Measures to Limit its
Spread in Hospitalized Patients. JCDR. 2013 [cited 11 May 2022]. doi:10.7860/JCDR/2013/5010.2737

25. Raymann K, Coon KL, Shaffer Z, Salisbury S, Moran NA. Pathogenicity of Serratia marcescens Strains in
Honey Bees. McFall-Ngai MJ, editor. mBio. 2018;9. doi:10.1128/mBio.01649-18

26. Cassidy-Hanley DM. Tetrahymena in the Laboratory: Strain Resources, Methods for Culture, Maintenance,
and Storage. Methods in Cell Biology. Elsevier; 2012. pp. 237–276. doi:10.1016/B978-0-12-385967-
9.00008-6

27. Leonard SP, Perutka J, Powell JE, Geng P, Richhart DD, Byrom M, et al. Genetic Engineering of Bee Gut
Microbiome Bacteria with a Toolkit for Modular Assembly of Broad-Host-Range Plasmids. ACS Synth Biol.
2018;7: 1279–1290. doi:10.1021/acssynbio.7b00399

28. Deatherage DE, Barrick JE. Identification of Mutations in Laboratory-Evolved Microbes from Next-
Generation Sequencing Data Using breseq. In: Sun L, Shou W, editors. Engineering and Analyzing
Multicellular Systems. New York, NY: Springer New York; 2014. pp. 165–188. doi:10.1007/978-1-4939-
0554-6_12

29. Pan X, Tang M, You J, Liu F, Sun C, Osire T, et al. Regulator RcsB Controls Prodigiosin Synthesis and Various
Cellular Processes in Serratia marcescens JNB5-1. Zhou N-Y, editor. Appl Environ Microbiol. 2021;87:
e02052-20. doi:10.1128/AEM.02052-20

30. Giardina G, Paiardini A, Fernicola S, Franceschini S, Rinaldo S, Stelitano V, et al. Investigating the Allosteric
Regulation of YfiN from Pseudomonas aeruginosa: Clues from the Structure of the Catalytic Domain.
Dobson R, editor. PLoS ONE. 2013;8: e81324. doi:10.1371/journal.pone.0081324

31. Kim HK, Harshey RM. A Diguanylate Cyclase Acts as a Cell Division Inhibitor in a Two-Step Response to
Reductive and Envelope Stresses. Gottesman S, editor. mBio. 2016;7: e00822-16. doi:10.1128/mBio.00822-
16

32. Filippova EV, Zemaitaitis B, Aung T, Wolfe AJ, Anderson WF. Structural Basis for DNA Recognition by the
Two-Component Response Regulator RcsB. Hendrickson WA, editor. mBio. 2018;9: e01993-17.
doi:10.1128/mBio.01993-17

33. Wall E, Majdalani N, Gottesman S. The Complex Rcs Regulatory Cascade. Annu Rev Microbiol. 2018;72:
111–139. doi:10.1146/annurev-micro-090817-062640

34. Pernestig A-K, Georgellis D, Romeo T, Suzuki K, Tomenius H, Normark S, et al. The *Escherichia coli* BarA-
UvrY Two-Component System Is Needed for Efficient Switching between Glycolytic and Gluconeogenic
Carbon Sources. J Bacteriol. 2003;185: 843–853. doi:10.1128/JB.185.3.843-853.2003

35. Herren CD, Mitra A, Palaniyandi SK, Coleman A, Elankumaran S, Mukhopadhyay S. The BarA-UvrY Two-
Component System Regulates Virulence in Avian Pathogenic *Escherichia coli* O78:K80:H9. Infect Immun.
2006;74: 4900–4909. doi:10.1128/IAI.00412-06

36. Teplitski M, Goodier RI, Ahmer BMM. Pathways Leading from BarA/SirA to Motility andVirulence Gene
Expression in *Salmonella*. J Bacteriol. 2003;185: 7257–7265. doi:10.1128/JB.185.24.7257-7265.2003

37. Sun Y-C, Guo X-P, Hinnebusch BJ, Darby C. The Yersinia pestis Rcs Phosphorelay Inhibits Biofilm Formation
by Repressing Transcription of the Diguanylate Cyclase Gene *hmsT*. J Bacteriol. 2012;194: 2020–2026.
doi:10.1128/JB.06243-11

38. Salah Ud-Din AIM, Roujeinikova A. Methyl-accepting chemotaxis proteins: a core sensing element in
prokaryotes and archaea. Cell Mol Life Sci. 2017;74: 3293–3303. doi:10.1007/s00018-017-2514-0

39. Raymann K, Shaffer Z, Moran NA. Antibiotic exposure perturbs the gut microbiota and elevates mortality in
honeybees. Gore J, editor. PLoS Biol. 2017;15: e2001861. doi:10.1371/journal.pbio.2001861

40. Flemming H-C, Wingender J. The biofilm matrix. Nat Rev Microbiol. 2010;8: 623–633.
doi:10.1038/nrmicro2415

41. Bailey SF, Hinz A, Kassen R. Adaptive synonymous mutations in an experimentally evolved Pseudomonas
fluorescens population. Nat Commun. 2014;5: 4076. doi:10.1038/ncomms5076

42. Khademi SMH, Sazinas P, Jelsbak L. Within-Host Adaptation Mediated by Intergenic Evolution in
Pseudomonas aeruginosa. Golding B, editor. Genome Biology and Evolution. 2019;11: 1385–1397.
doi:10.1093/gbe/evz083

43. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9: 357–359.
doi:10.1038/nmeth.1923

44. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST Server: Rapid Annotations using
Subsystems Technology. BMC Genomics. 2008;9: 75. doi:10.1186/1471-2164-9-75

45. Lai H-Y, Zhang Z-Y, Su Z-D, Su W, Ding H, Chen W, et al. iProEP: A Computational Predictor for Predicting
Promoter. Molecular Therapy - Nucleic Acids. 2019;17: 337–346. doi:10.1016/j.omtn.2019.05.028

Supporting Material

Figure S1. Mean growth rate of each variant based on 600nm OD readings every hour for 24h of ME (purple) and PE (green) isolates in LB broth (left) or fresh Neff media (right) compared to the ancestral strain (black).

Table S1. Total reads, number of mapped reads, and average genome coverage for each population and isolate.

Table S2. List of parallel mutations identified across evolved populations.

Table S3. List of all the mutations identified in the evolved isolates and their frequency in the populations.

Dataset S1. List of all the mutations identified in the evolved populations and their frequencies.