An interactive pipeline for multitrait evolutionary analysis in a phylogenetic framework

Dipendra Nath Basu [1,*], Vaishali Bhaumik [1,*], Krushnamegh Kunte

National Centre for Biological Sciences, Tata Institute of Fundamental Research, GKVK

Campus, Bellary Road, Bangalore 560065, India

[1] These authors contributed equally

* Correspondence and email: Dipendra Nath Basu (dipendran@ncbs.res.in, +91 9535-699-

750), Vaishali Bhaumik (vaishali@ncbs.res.in, +91 9731-552-552)

ABSTRACT

Comparative phylogenetic studies based on 2D image data can reveal macroevolutionary patterns and processes by integrating phylogenetic and ecological analyses. The advent of high-throughput imaging facilities has led to the accumulation of large image datasets; however, large-scale data extraction and analysis using these resources can be labour- and computation-intensive. Moreover, there are few standardized methods to analyse multitrait image-based data in a phylogenetic framework. Several packages in the R statistical environment offer utilities for processing and analysing large-scale image data. Techniques for the comparative phylogenetic analysis of multitrait data have also been introduced in other packages. However, most of these approaches are specialized for discrete purposes, and there is a lack of streamlined and efficient methodological pipelines integrating features across packages. Here, we present multEvol, an interactive pipeline for the extraction and phylogenetic analysis of shape and colour pattern data from 2D images in the R statistical computing environment. Using this pipeline, shape outlines and colour pattern data for multiple colours can be extracted, visualised, and compared in a phylogenetic framework. Using various utilities provided in the *phytools*, *RNiftyReg*, *Momocs*, *patternize*, *convevol*, and *geiger* packages, we introduce an interactive interface that allows user inputs for iterative data extraction. This pipeline provides an efficient method for the bulk processing of images of biological specimens from museum collections. As such, it has potential applications in the large-scale morphometric analysis of museum specimens to reveal macroevolutionary patterns in ecology, evolution and integrative biology.

Keywords: Colour pattern, elliptic Fourier analysis, multitrait evolution, phylogenetic analysis, convergence, rate of evolution

INTRODUCTION

Earth's biodiversity is marked by spectacular morphological evolution and diversification

of species. Species diversification typically involves evolution along multiple trait axes, which

helps organisms adapt to their natural environments and occupy new niches. Comprehensive

macroevolutionary analyses of these traits has long posed a challenge to biologists. Quantifying the

degree, direction, and rate of morphological change can help elucidate macro- and micro-

evolutionary processes in biological systems. However, comparative studies in ecology are

frequently limited by a lack of standardized methodology for data extraction and processing [1,2],

especially in the bulk analysis of data within a phylogenetic framework. The extracted data may

also need to be summarized or otherwise re-structured for use in phylogenetic and statistical

analyses, and the methodologies for these are not readily apparent or easily available.

The extraction of shape and colour pattern data from 2D images—such as images of insect

wings or plant leaves—offers a versatile means of data collection using relatively inexpensive

resources. Comparative studies on diverse taxa, such as bats [3] and butterflies [4], have used shape

analysis to examine the evolutionary history and functional constraints of morphological features.

Wing colours and colour patterns are extremely diverse, and play important roles in mimicry [5],

mate recognition [6], and thermoregulation [7], along with other adaptive functions. However,

comparative studies on the evolution of complex traits usually focus on relatively closely related

species [3,8,9], which may limit the utility of these methods in comparisons across study systems.

In addition, standard methods for phylogenetic analysis are typically designed to address specific

questions, and are incompatible with image data. Therefore, it is important to develop a systematic

and interactive pipeline that extracts preliminary data from images and transforms them into

phylogenetically corrected formats that are usable for further comparative analysis across distantly

related taxa.

Here, we present multEvol, a pipeline for data extraction from 2D images and phylogenetic

analysis of morphological data (Fig. 1) in the R statistical computing environment [10]. The

extraction of shape data mostly depends on the *Momocs* package [11], which is a dedicated tool for

shape analysis in R. The extraction of colour pattern data primarily uses the *patternize* package

[12], which allows pattern extraction at the pixel level using a user-defined RGB threshold. In both

cases, we have created functions that allow user interaction, which can help ensure proper

alignment, harmonic calibration (for shape), and pixel-level pattern selection (for colour) between

multiple samples. The outputs can be transformed using phylogenetic principal component analysis

(pPCA, package *phytools* [13]), and the PCs can be used for further statistical or phylogenetic

analysis, such as estimating the degree of convergence (package *convevol* [14]) or rate of evolution
(package *RRphylo* [15]). We demonstrate the utilization of this pipeline using 2D wing images of
butterflies. The pipeline can also be used to analyse any continuous variables—such as
morphometric data—that have been extracted using other methods.
71
METHODS
Data extraction
Outline shape: In geometric morphometric analyses, biological shapes are estimated through two
major approaches: landmark-based analysis, and outline analysis. Although both methods retain
the relative geometric information of the shapes [16,17], coordinate homology is an essential
prerequisite in the landmark method [18]. However, homologous landmarks may not be available
in all biological samples, which restricts the utilization of this approach. Additionally, landmarking
on the homologous points is a time-consuming process. In contrast, elliptical Fourier analysis
(EFA) [17,19,20] is a robust, resource-efficient, and time-efficient method of outline analysis that
accurately quantifies 2D closed contours. Unlike other Fourier-based methods, EFA can operate on
virtually any shape, including ones that lack positions for homologous landmarks [20].
Contemporary studies on geometric morphometrics are increasingly using EFA to estimate 2D
biological shapes [21].
In the EFA approach, the x and y coordinates of closed contours are extracted from the
binary images. These outlines are smoothened to a small factor to account for biological
aberrations, following which they are centred, aligned, and scaled to remove allometric signals
(Fig. 2a). The outlines are size-corrected based on the centroid size. Outline coordinates for each
specimen are resampled to introduce homology, and a Procrustes fit is performed on these
homologous outline coordinates for better alignment. Resetting the first coordinate of each outline
further standardizes the contour estimates. A suitable number of harmonic ranks (a series of
ellipses) are then estimated based on the meanshape of the aligned contours (Fig. 2b). Each contour
is then represented as a series of calibrated elliptical harmonics (Fig. 2c). These ellipses are defined
as the sum of one sine and one cosine curve by decomposing a periodic continuous function
implementing an extension of the Fourier series (Fig. 2d). Fourier transformation calculates four
coefficients for each harmonic, which are eventually normalized based on the first harmonic to
remove the effect of size and rotational variations [17]. These normalized elliptical Fourier
descriptors (EFDs) are used as shape variables for further analysis. We implemented a user-friendly

interactive script 'cleqEFA' based on utilities provided in R package *Momocs* for outline extraction,

alignment, and transformation techniques [11,22].

Colour pattern: Colour pattern is a complex trait with multiple quantifiable dimensions. Colour

pattern data can be quantified in various ways, including relative linear measurements or areas

occupied by a particular colour range [23,24] or estimating the shape variance of homologous

colour patches across multiple surface [25]. These linear and geometric data can be stored in a

presence-absence matrix for each range of colour values (RGB values) on appropriately aligned

surfaces. Therefore, this method of pattern extraction makes the data comparable across variable

and large image datasets, and can be used for further analysis. The accuracy of these data depends

on two components: an efficient registration approach to introduce homology between samples,

and appropriate colour thresholding to extract the spatial and geometric information of homologous

patches.

Images can be superimposed to obtain homology through two main approaches: image

registration and landmark-based homology. Biological specimens often lack homologous

anatomical positions, which restricts the use of landmark-based approaches. Therefore, to obtain

homology in a computationally efficient manner, we have opted for a two-step image

transformation. In the first step, a binary image template is prepared based on either the meanshape

of the 2D surfaces containing colour patterns, or any surface with close resemblance to the

meanshape. Subsequently, one specimen of each taxon or species is affine-transformed onto this

binary template to obtain a taxon- or species-specific template. In the second step, these templates

are used to register all the images using an image registration approach based on similarity of colour

intensity (Fig. 1b). The affine transformations are performed iteratively using the utilities of the R

package *RNiftyReg* [26].

Following image registration, the colour patches that are homologous across multiple

surfaces are isolated through thresholding. Most unsupervised thresholding approaches are limited

to systems with binary or conspicuous colour patterns (that is, systems in which the surfaces have

distinct background and foreground colours), and are not very efficient in systems with multiple

colours or continuous colour boundaries. This pipeline uses a supervised RGB thresholding

approach where colour boundaries are defined by iteratively sampling from a representative

specimen for each species. This approach efficiently extracts colour patterns as raster layer objects

for each selected colour range. These raster layers are then converted to binary presence–absence

matrices and used for subsequent analyses. RGB thresholding and raster layer extraction are

performed using utilities in the R package *patternize* [12].

Phylogenetic analysis

Phylogenetic PCA and size-correction: Most statistical analyses assume that the data have been

collected from independent samples. However, this assumption does not hold during data analysis

in a phylogenetic framework. Comparative phylogenetic analyses that involve statistical

transformation of continuous morphological variables for the estimation of phylogenetic signal or

ancestral state reconstruction may require phylogenetic corrections [27]. Statistical comparisons

without preliminary phylogenetic corrections show higher variance and are prone to type I error.

Hence, this pipeline includes a phylogenetic PCA and offers the option of size correction against

body size, as implemented in the R package *phytools* [13].

Mode of evolution: Several analytical methods have been developed to estimate non-neutral trends

of trait evolution in a phylogenetic framework [9,28]. Estimating these model parameters can reveal

the effects of selection and constraints on trait evolution. The 'pcModels' function uses utilities

provided in the R packages *APE* [29], *phytools* [13], and *geiger* [30] to fit a phylogenetic tree with

a user-defined number of principal components (PCs) and obtain the best fit model. To estimate

the extent of convergence in trait evolution, the *convevol* package is used to calculate the C1 metric

[28], which is a Euclidean distance-based estimate of convergence among groups of species on a

phylomorphospace. Among the various C metrices that can be calculated, C1 denotes the relative

convergence in each trait between a group of species, and is comparable across datasets. The

convevol package also provides a way to simulate data using BM model parameters to evaluate the

significance of the convergence. The simulation results (Fig. 1c) are visualized using the 'c1Plots'

function.

Rate estimation: Comparative phylogenetic analyses often require estimates of the rate of change

in phenotypes across evolutionary time. The rate of trait evolution depends on ancestral state

reconstruction, which can be performed in two ways: 1) ancestral state reconstruction following a

macroevolutionary model (single- or variable-rate models) within a well sampled clade or closely

related clades; and 2) regression-based estimation. A regression-based method is computationally

more efficient than Bayesian approaches, and can accurately predict the ancestral characters of

species belonging to distant clades. The multEvol pipeline implements the ridge regression method

in the *RRphylo* package [15] to estimate the rate of trait evolution of multiple traits using PCs

obtained from the preliminary analysis (Fig. 1c).

OUTPUT

The main *multEvol* functions for shape extraction take images as input; extract, scale, and align the

outlines; and perform elliptical Fourier analysis on those shapes. The output of the 'cleqEFA;

function is a list containing the EFDs as a Coe object, as well as the estimated mean outline for

each species as a Coo object (Momocs). The 'shapeComparison' also extracts the deformation

isolines among the meanshapes of different species for comparative visualization. The 'colData'

function for colour pattern extraction performs RGB thresholding on images using iterative user

inputs in an interactive manner to obtain binary matrices as outputs from each raster object

(*patternize*). These matrices are eventually transformed into a proportional matrix for each colour

range for each species. The 'colData' function also stores the raster layer objects for each colour

range in .png format as confirmation for efficient registration and thresholding, as well as a list of

the user-selected RGB values and threshold for each sample. The 'concat' function concatenates

the linearized proportional matrices of different colour ranges. The 'colourComparison' function

sums the rasterstacks to visualize the heatmaps of intra- and inter-species colour pattern variation

in a comparative manner. The EFDs from outlines, species-level proportional colour pattern data,

and any phylogenetically size-corrected continuous multivariate datasets can be subjected to

phylogenetic PCA using a phylogenetic tree as a phylo object. The PCA results are used to visualise

inter-species variations in trait space, and the PC scores are stored for further analysis. The

'dataPrep' function fits six macroevolutionary models (BM, EB, OU, lambda, kappa, delta) to the

PCs of each trait and stores the AICc values for each model. The 'c1' function estimates and stores

C1 metrics for each user-defined group of species (regimes) and calculates their significance based

on simulations using a Brownian motion (BM) model. The 'c1Plot' function plots the distribution

of C1 values from the BM simulation, along with the actual C1 value for each trait in each regime.

The rate of trait evolution is estimated by the 'rateEvol' function based on the user-defined number

of PCs and a phylo object. This function stores the evolutionary rate for each branch of the

phylogeny in a data frame, which can be used to plot a heatmap cladogram.

EXAMPLE

Study system: In biological systems, mimicry occurs when a palatable prey species evolves

resemblance to a well-defended toxic (aposematic) species to experience reduced predation

pressure and derive a fitness benefit. Aposematic species that resemble each other in a mutualistic interaction are known as Müllerian co-mimics, whereas palatable species that mimic a toxic model in a parasitic interaction are known as Batesian mimics. In this example, we define a mimicry ring as a community that contains at least one well defended aposematic species (Batesian model) and at least one palatable species (Batesian mimic) that resembles the model [31].

Butterflies in a mimicry ring show varying degrees of phenotypic convergence in multiple traits, including colour [32] and colour pattern signals [5,33] as well as locomotor cues [34]. These traits may evolve under different models of selection and at varying rates within and between species [4,23]. The *Tirumala* mimicry ring endemic to the Western Ghats biodiversity hotspot in India consists of five unpalatable models with a shared phenotype and shared ancestry, and three distantly related palatable species (Batesian mimics) that have evolved to resemble the toxic models (Fig. 3) [31,35]. The mimics comprise two species of female-limited mimics (*Pareronia hippia* and *P. ceylanica*), where the males are non-mimetic, and one species of sexually monomorphic mimic (*Papilio clytia* f. *dissimilis*), where both sexes share the mimetic phenotype. Here, we examine the evolution of wing shapes and colour patterns in this mimicry ring using the multEvol pipeline to demonstrate its utility in investigating the evolution of multiple traits under various degrees of mimetic selection. For phylogenetic analyses, we first prepared a time-calibrated phylogeny consisting of the models, mimics, and some of their respective sister species. We then extracted wing shape and colour data from the forewing images of all species and estimated the degree of convergence between models and mimics, as well as the rate of evolution of the two traits, as described above.

Data extraction and preliminary analysis: Using digital SLR cameras and lenses, we photographed one forewing (dorsal surface) of each specimen and extracted the wing shape and colour pattern data. Images were captured under uniform camera flash units because variation in colour tones under different light conditions may produce different results. We extracted shape data from 265 images using the 'cleqEFA' function, and colour data from 200 images using the 'colData' function. Seven harmonic ranks were calibrated to explain the outlines with 99% accuracy (Fig. 2b). The EFDs from these seven harmonics were normalized against those of the first harmonics, resulting in six remaining EFDs. Data were extracted for three colour ranges— melanistic (including black and brown), red/orange, and blue/green—with a resampling factor of three. The resulting proportional matrices of each colour range were linearized into a vector object

and concatenated for each species. The EFD datasets and concatenated colour pattern data were

used for a phylogenetic PCA, and six evolutionary models were fit to the resulting PCs.

RESULTS

The first two PCs explained 96.23% and 98.1% of the variance in shape and colour pattern data,

respectively. Of the six models compared, the lambda model of evolution had the highest support

for the first two of shape (AICc values: PC1, -83.21; PC2, -110.51) and color (AICc values: PC1,

228.76; PC2, 174.03). The meanshapes of each model and mimic species were reconstructed from

their respective EFDs. For each model–mimic pair, the shape changes were visualized as

deformation isolines between the meanshapes of each species (Fig. 4a). In contrast to sexually

monomorphic mimics and mimetic females of female-limited mimics, the non-mimetic males

showed multiple local deformations (red areas in Fig. 4a). The summed raster layers were plotted

as heatmaps to visualise intra-species colour variation (that is, variation in colour patterns between

samples of a single species; plots along the top and left margins in Fig. 5a). Pairwise differences in

the species-level proportional raster layers were plotted as heatmaps to indicate inter-species

differences in colour patterns between each model–mimic pair (5a). Similar to the trend observed

in wing shapes, non-mimetic males showed more differences in colour patches compared to

models, whereas sexually monomorphic and female-limited mimics were similar to models. The

phylomorphospaces for shape and colour pattern analysis indicated an overlap between models and

mimics (Figs. 3b and 4b). In contrast to mimetic females, the non-mimetic males of female-limited

mimetic species were relatively distant from the traitspace occupied by models and mimics. The

convergence indices of various species combinations were plotted using the shape and colour

pattern data to visualize the distribution of C1 values in 100 simulated datasets (Fig. 6a). There was

significant convergence in wing shape and colour patterns within models and between models and

female-limited mimics, as well as in wing shape between models and sexually monomorphic

mimics (star symbols in Fig. 6a). The evolutionary rates of wing shape and colour pattern were

plotted onto heatmap cladograms, which showed that lineages containing models or mimics

generally evolved faster (Fig. 6b).

CONCLUSION AND RECOMMENDATIONS

The *multEvol* pipeline can be used to compare the distribution of colour ranges across species, and

the interactive features allow the iterative selection of appropriate harmonics, RGB values, and

colour pattern matrices across multiple samples. This is especially useful for large-scale supervised

data collection from museum collections or field-collected specimens. Museum collections worldwide represent a vast resource of historical DNA; however, preserved specimens have limited applicability as sources of biological tissue. As such, the systematic photographing and cataloguing of museum specimens can improve their long-term usefulness as sources of morphological data [4,36]. The utilization of museum specimens alleviates several taxonomical and logistical limitations [37,38] and enables the testing of various macroevolutionary hypotheses. The pipeline presented here provides the potential for such large-scale comparative analyses, and we hope that the methods described and pipeline presented here will prove useful to researchers studying shape and colour pattern variation in biological systems.

Author Contributions: DNB and VB conceptualised the analytical pipeline, developed the code, and conceptualised and performed phylogenetic analyses. KK conceived the broad research questions, and supervised the research. DNB and VB wrote the manuscript with inputs from KK.

Competing Interest Statement: Authors declare no competing interests.

Data availability: The functions and their descriptions are available on github (https://github.com/DipendraBasu/multEvol-0.1).

Funding: This research was funded in part by a Ramanujan Fellowship from the Dept. of Science and Technology, Govt. of India, and a research grant from NCBS to KK.

References

1. Koch M, Universitaria C, Federal C, Aires B, Soto IM, Federal C, et al. Zoologica Scripta. 2015; 463–474. doi:10.1111/zsc.12120

2. Watters JL, Cummings ST, Flanagan RL, Siler CD. Review of morphometric measurements used in anuran species descriptions and recommendations for a standardized approach. 2016;4072: 477–495.

3. Pie MR, Tschá MK. Size and shape in the evolution of ant worker morphology. PeerJ. 2013;2013: 1–14. doi:10.7717/peerj.205

4. Owens HL, Lewis DS, Condamine FL, Kawahara AY, Guralnick RP. Comparative phylogenetics of *Papilio* butterfly wing shape and size demonstrates independent hindwing and forewing evolution. Wright A, editor. Syst Biol. 2020;69: 813–819. doi:10.1093/sysbio/syaa029

5. Finkbeiner SD, Briscoe AD, Reed RD. Warning signals are seductive: Relative contributions of color and pattern to predator avoidance and mate attraction in Heliconius

butterflies. Evolution (N Y). 2014;68: 3410–3420. doi:10.1111/evo.12524

6.   Fordyce JA, Nice CC, Forister ML, Shapiro AM. The significance of wing pattern

diversity in the Lycaenidae: mate discrimination by two recently diverged species. J Evol

Biol. 2002;15: 871–879. doi:doi:10.1046/j.1420-9101.2002.00432.x

7.   Gautam S, Kunte K. Adaptive plasticity in wing melanisation of a montane butterfly

across a Himalayan elevational gradient. Ecol Entomol. 2020; een.12911.

doi:10.1111/een.12911

8.   Price-Waldman RM, Shultz AJ, Burns KJ. Speciation rates are correlated with changes in

plumage color complexity in the largest family of songbirds. Evolution (N Y). 2020;74:

1155–1169. doi:10.1111/evo.13982

9.   Mahler DL, Ingram T, Revell LJ, Losos JB. Exceptional convergence on the

macroevolutionary landscape in island lizard radiations. Science. 2013;341: 292–295.

doi:10.1126/science.1232392

10.   R Development Core Team. R: a language and environment for statistical computing.

Vienna: R Foundation for Statistical Computing; 2019.

11.   Bonhomme V, Picq S, Gaucherel C, Claude J. Momocs: outline analysis using R. Journal

of Statistical Software. 2014. doi:10.18637/jss.v056.i13

12.   Van Belleghem SM, Papa R, Ortiz-Zuazaga H, Hendrickx F, Jiggins CD, Owen McMillan

W, et al. patternize: An R package for quantifying colour pattern variation. Vamosi J,

editor. Methods Ecol Evol. 2018;9: 390–398. doi:10.1111/2041-210X.12853

13.   Revell LJ. phytools: an R package for phylogenetic comparative biology (and other

things). Methods Ecol Evol. 2012;3: 217–223. doi:10.1111/j.2041-210X.2011.00169.x

14.   Aguilar-Puntriano C, Avila LJ, De la Riva I, Johnson L, Morando M, Troncoso-Palacios J,

et al. The shadow of the past: convergence of young and old South American desert

lizards as measured by head shape traits. Ecol Evol. 2018;8: 11399–11409.

doi:10.1002/ece3.4548

15.   Castiglione S, Tesone G, Piccolo M, Melchionna M, Mondanaro A, Serio C, et al. A new

method for testing evolutionary rate variation and shifts in phenotypic evolution. Methods

Ecol Evol. 2018;9: 974–983. doi:10.1111/2041-210X.12954

16.   Moellering H, Rayner JN. The harmonic analysis of spatial shapes using dual axis Fourier

shape analysis (DAFSA). Geogr Anal. 1981;13: 64–77. doi:10.1111/j.1538-

4632.1981.tb00715.x

17.   Kuhl FP, Giardina CR. Elliptic Fourier features of a closed contour. Comput Graph Image

Process. 1982;18: 236–258. doi:10.1016/0146-664X(82)90034-X

18.   MacLeod N. Generalizing and extending the Eigenshape method of shape space.

Palaeobiology. 1999;25: 107–138.

19.   Giardina CR, Kuhl FP. Accuracy of curve approximation by harmonically related vectors

with elliptical loci. Comput Graph Image Process. 1977;6: 277–285. doi:10.1016/S0146-

664X(77)80029-4

20.   Ferson S, Rohlf FJ, Koehn RK. Measuring shape variation of two-dimensional outlines.

Syst Biol. 1985;34: 59–68. doi:10.1093/sysbio/34.1.59

21.   Outomuro D, Ángel-Giraldo P, Corral-Lopez A, Realpe E. Multitrait aposematic signal in

Batesian mimicry. Evolution (N Y). 2016;70: 1596–1608. doi:10.1111/evo.12963

22.   Kallal RJ, de Miranda GS, Garcia EL, Wood HM. Patterns in schizomid flagellum shape

from elliptical Fourier analysis. Sci Rep. 2022;12: 3896. doi:10.1038/s41598-022-07823-y

23.   Bijl W, Zeuss D, Chazot N, Tunström K, Wahlberg N, Wiklund C, et al. Butterfly

dichromatism primarily evolved via Darwin's, not Wallace's, model. Evol Lett. 2020;4:

545–555. doi:10.1002/evl3.199

24.   Sezer M, Tekelioglu O. Quantification of Japanese Quail eggshell colour by image

analysis. Biol Res. 2009;42: 99–105. doi:10.4067/S0716-97602009000100010

25.   Le Poul Y, Whibley A, Chouteau M, Prunier F, Llaurens V, Joron M. Evolution of

dominance mechanisms at a butterfly mimicry supergene. Nat Commun. 2014;5: 5644.

doi:10.1038/ncomms6644

26.   Clayden J, Modat M, Presles B, Anthopoulos T, Daga P. RNiftyReg: Image registration

using the "NiftyReg" library. 2021.

27.   Revell LJ. Size-correction and principal components for interspecific comparative studies.

Evolution (N Y). 2009;63: 3258–3268. doi:10.1111/j.1558-5646.2009.00804.x

28.   Stayton CT. The definition, recognition, and interpretation of convergent evolution, and

two new measures for quantifying and assessing the significance of convergence.

Evolution (N Y). 2015;69: 2140–2153. doi:10.1111/evo.12729

29.   Paradis E, Claude J, Strimmer K. APE: analyses of phylogenetics and evolution in R

language. Bioinformatics. 2004;20: 289–290.

doi:https://doi.org/10.1093/bioinformatics/btg412

30.   Harmon LJ, Weir JT, Brock CD, Glor RE, Challenger W. GEIGER: investigating

evolutionary radiations. Bioinformatics. 2008;24: 129–131.

doi:10.1093/bioinformatics/btm538

31. Kunte K, Kizhakke AG, Nawge V. Evolution of mimicry rings as a window into
community dynamics. Annu Rev Ecol Evol Syst. 2021;52: 315–341. doi:10.1146/annurev-
ecolsys-012021-024616
32. Kazemi B, Gamberale-Stille G, Leimar O. Multi-trait mimicry and the relative salience of
individual traits. Proc R Soc B Biol Sci. 2015;282. doi:10.1098/rspb.2015.2127
33. Dittrich W, Gilbert F, Green P, McGregor P, Grewcock D. Imperfect mimicry: a pigeon's
perspective. Proc R Soc B. 1993;251: 195–200. doi:10.1098/rspb.1993.0029
34. Srygley RB, Ellington CP. Discrimination of flying mimetic, passion-vine butterflies
Heliconius. Proc R Soc London Ser B Biol Sci. 1999;266: 2137–2140.
doi:10.1098/rspb.1999.0899
35. Joshi J, Prakash A, Kunte K. Evolutionary assembly of communities in butterfly mimicry
rings. Nuismer SL, Bronstein JL, editors. Am Nat. 2017;189: E58--E76.
doi:10.1086/690907
36. Van Belleghem SM, Alicea Roman PA, Carbia Gutierrez H, Counterman BA, Papa R.
Perfect mimicry between *Heliconius* butterflies is constrained by genetics and
development: mimicry and developmental constraints. Proc R Soc B Biol Sci. 2020;287.
doi:10.1098/rspb.2020.1267rspb20201267
37. Frankino WA, Zwaan BJ, Stern DL, Brakefield PM. Natural selection and developmental
constraints in the evolution of allometries. Science. 2005;307: 718–720.
38. Jantzen B, Eisner T. Hindwings are unnecessary for flight but essential for execution of
normal evasive flight in Lepidoptera. Proc Natl Acad Sci USA. 2008;105: 16636–16640.
doi:10.1073/pnas.0807223105
378
379

 Figure 1: Data extraction, outputs, and potential statistical analyses of shape and colour data

using the multEvol pipeline. Butterfly wings are photographed and converted into binary images

for the extraction of shape (a) or colour pattern (b) data. The 2D shapes are scaled, aligned across

all images, and Fourier transformed to obtain the mean descriptors for each species or group. The

colour data comprise proportional matrices for multiple colours, where each matrix indicates the

pixels that contain the specified colour ($0 = $none of the images sampled contain the specified colour

in that pixel; $1 = $all images sampled contain the specified colour in that pixel). The matrices are

linearized into a single vector onject and used for further analysis, which may include a

phylogenetic principal component analysis (PCA) and estimation of convergence or evolutionary

rate.

Figure 2: Data processing and extraction of elliptical Fourier descriptors for shape outlines

using the 'cleqEFA' function. (a) The right forewings (n=265) of butterflies in the *Tirumala*

mimicry ring are binarized, and their outline coordinates are extracted. These outlines are scaled

and aligned, and the start points of the contours are reset to obtain the final stack of outlines. (b)

Harmonic power plateaus after the seventh harmonic. (c) The harmonic ranks are calibrated to

obtain an accurate representation of the outlines. The coordinates of 12 randomly selected

equidistant points on the wing contours are plotted directionally to observe the periodic

decomposition of the sine and cosine curves from an outline. (d) The harmonic sum of a series of

fitted ellipses (n=7) on the wing outline, where the higher order ellipse role on the periphery of

the lower order ellipse (d) are Fourier transformed to obtain elliptical Fourier descriptors.

Figure 3: Müllerian co-mimics and Batesian mimics in the *Tirumala* mimicry ring of the Western Ghats, India. The five Müllerian co-mimics (red arrows) have shared ancestry and act as Batesian models for the Batesian mimics (blue arrows), which are distantly related. *Pareronia* spp. are female-limited mimics, where the females mimic the models and the males are non-mimetic.

Figure 4: Inter-species variation in wing shape among mimetic butterflies. (a) Pairwise
comparisons of deformation isolines (scale: blue=low deformation, yellow=moderate deformation,
red=high deformation), showing the changes in wing shape in model–mimic species pairs in the
Tirumala mimicry ring. **(b)** A morphospace plot based on the phylogenetic PCA of shape
descriptors, showing the relative positions of models (red), mimics (dark blue), and non-mimetic
males of female-limited mimics (light blue) in the traitspace.

Figure 5: Intra- and inter-species variation in wing colour patterns among mimetic

butterflies. (a) Heatmaps along the top and left margins show intra-species variations in colour

patterns, and the remaining heatmaps exhibit inter-species differences in colour patterns in model–

mimic species pairs in the *Tirumala* mimicry ring. The heatmap scale is shown at the bottom right.

(b) A morphospace plot based on the phylogenetic PCA of colour pattern data, showing the relative

positions of models (red), mimics (dark blue), and non-mimetic males of female-limited mimics

(light blue) in the traitspace.

 Figure 6: Evolutionary characteristics of forewing shape and forewing dorsal colour patterns

in the *Tirumala* mimicry ring. (a) Convergence among Batesian mimics (distantly related

palatable species; includes female-limited dimorphic and sexually monomorphic mimics); among

aposematics (toxic species with a similar colour patterns); between aposematics and female-limited

dimorphic mimics; and between aposematics and sexually monomorphic mimics. C1 values (red:

actual C1; black: mean C1 across 100 simulations) are shown for forewing shape (left) and

forewing dorsal colour patterns (right). A star indicates significant convergence compared to the

convergence estimates from 100 simulations. **(b)** Rates of evolution of forewing shape and

forewing dorsal colour patterns. Colour coding for species names indicates aposematics (red),

Batesian mimics (dark blue), and the males of female-limited Batesian mimics (light blue).

a
Museum specimen photograph → Extracted resized porion of image (resized) → Conversion to binary image → Extracted 2D closed contours → Scaling and aligning

Input: No of harmonics
Fourier transform

Fitting harmonics estimating meanshape outlines

b
2D registration on a binary target (affine transform)

Target

Input:
RGB values for blue →
RGB values for black →
Colour thresholding (each colour, each specimen)

Rasterstack & binary matrix extraction (each colour, each specimen)

Proportional matrix (each colour, each species)

Fourier descriptors for all the shape outlines

Descriptors for mean shape of each species

Any continuous multivariate data

Phylogenetic correction/ standardization

Phylogenetic PCA

c

Convergence

Multi trait comparisons

Comparison of convergence (Mode of trait evolution)

C1 convergence index (Stayton, 2015)

Trait A   Trait B

Heatmapped cladograms (Rate of trait evolution)

Ridged regression

5
0
−5
−5   0   5   10
Phylomorphospace

Phylogeny

Pipelines: ▬ a. Shape trait extraction   ▬ b. Colour pattern trait extraction   ▬ c. Phylogenetic comparisons

Tirumala limniace ♂♀

Tirumala septentrionis ♂♀

♀ *Pareronia ceylanica* ♂

Parantica aglea ♂♀

Parantica nilgiriensis ♂♀

♀ *Pareronia hippia* ♂

Idea malabarica ♂♀

Papilio clytia f. dissimilis ♂♀

Batesian model and Müllerian co-mimic

Batesian mimic

 Müllerian co-mimics

 Aposematic species

Batesian mimics

 Batesian mimics

 Intersexual genetic correlatedness

a   Pairwise comparison forewing shape

Species names	*Idea malabarica*	*Parantica aglea*	*Parantica nilgiriensis*	*Tirumala limniace*	*Tirumala septentrionis*
Papilio clytia ♀					
Papilio clytia ♂					
Pareronia ceylanica ♀					
Pareronia ceylanica ♂					
Pareronia hippia ♀					
Pareronia hippia ♂					

b   Phylomorphospace forewing shape

1. *Parantica aglea*
2. *Parantica nilgiriensis*
3. *Tirumala limniace*
4. *Tirumala septentrionis*
5. *Idea malabarica*
6. *Pareronia hippia* ♀
7. *Pareronia hippia* ♂
8. *Pareronia ceylanica* ♀
9. *Pareronia ceylanica* ♂
10. *Papilio clytia* ♀
11. *Papilio clytia* ♂

a   Pairwise comparison forewing dorsal colour pattern

Species Name	*I. malabarica*	*P. aglea*	*P. nilgiriensis*	*T. limniace*	*T septentrionis*
P. clytia ♀					
P. clytia ♂					
P. ceylanica ♀					
P. ceylanica ♂					
P. hippia ♀					
P. hippia ♂					

0.8 0.4 0.0   -1 0 1

Variation

Intra species   Inter species

b   Phylomorphospace forewing colour pattern

PC2 (25.4%)

PC1 (72.7%)

1. *Parantica aglea*
2. *Parantica nilgiriensis*
3. *Tirumala limniace*
4. *Tirumala septentrionis*
5. *Idea malabarica*
6. *Pareronia hippia* ♀
7. *Pareronia hippia* ♂
8. *Pareronia ceylanica* ♀
9. *Pareronia ceylanica* ♂
10. *Papilio clytia* ♀
11. *Papilio clytia* ♂

a Convergence

b Rate of trait evolution