TITLE

Location-Specific Facilitation in Primate Auditory Cortex

AUTHORS

Chenggang Chen[+], Sheng Xu[+], Yunyan Wang, and Xiaoqin Wang*

AFFILIATIONS

Laboratory of Auditory Neurophysiology, Department of Biomedical Engineering, Johns Hopkins University School of Medicine, Baltimore, Maryland 21205, USA.

[+] Co-first author

CORRESPONDENCE

*Correspondence to: xiaoqin.wang@jhu.edu

SUMMARY

Neural responses to sensory stimuli are markedly influenced by the context in which a stimulus is preceded or embedded. Cortical and subcortical neurons typically exhibit adaptation to repetitive auditory, visual, somatosensory, and olfactory stimulation. Here, we investigated single neuron responses to sequences of sounds either repeatedly delivered from a single spatial location or randomly delivered from multiple spatial locations in the auditory cortex of awake marmosets. Instead of inducing adaptation, repetitive stimulation from a target speaker evoked long-lasting, location-specific facilitation (LSF) in many neurons, irrespective of the visibility of the target speaker. The extent of LSF decreased with decreasing presentation probability of the target speaker. Intracellular recordings showed that repetitive sound stimulation evoked sustained membrane potential depolarization which gave rise to firing rate facilitation. Computational models suggest two distinct neural mechanisms underlying LSF. Our findings revealed a novel form of contextual modulation in the auditory cortex that may play a role in auditory streaming and predictive coding.

INTRODUCTION

It has been well established that responses of neurons in sensory cortical and subcortical areas are influenced by stimulus context. Contextual effects are mainly suppressive when the same stimulus is repeatedly presented. For example, the presence of a preceding sound suppressed a neuron's responses to a succeeding sound of the same feature in ascending auditory pathway (Shore, 1995; Nelson et al., 2009; Xiong et al., 2020; Bartlett and Wang, 2005); visual cortical neurons exhibited repetition suppression to the image that is presented twice (Muller et al., 1999; Williams and Olson, 2022); repetitive whisker stimulation reduced the responses of somatosensory neurons (Ganmor et al., 2010; Heiss et al., 2008); and neural activities in the olfactory pathway habituated to the repeated odor presentation (Wilson, 1998; Kato et al., 2012). In addition to temporal context, spatial context can also suppress neural responses in different modalities. For example, along the ascending visual pathway, neural responses to a stimulus at its receptive field are suppressed by the same surrounding stimulus (Huang et al., 2019; Alitto and Usrey, 2008; Keller et al., 2020); somatosensory cortical neural responses to deflection of its principal whisker are suppressed when its neighboring whiskers are deflected (Mirabella et al., 2001; Petersen et al., 2009); and responses of auditory neurons to a probe sound from one spatial location are suppressed by masker sounds from other locations (Fitzpatrick et al., 1999; Reale and Brugge, 2000; Mickey and Middlebrooks, 2005). In a study of awake marmoset auditory cortex, it was found that a masker placed far away from a neuron's spatial receptive field (SRF) suppressed the response elicited by a probe sound in its SRF, suggesting contextual modulations are widespread and suppressive in the spatial domain (Zhou and Wang, 2012, 2014). The above-cited previous studies suggest that neural adaptation to repetitive stimuli is a general rule in cortical and subcortical neurons across sensory modalities.

An important property of neural adaption in sensory neurons is stimulus-specific

adaptation (SSA) to the stimuli that are presented with a high probability. SSA has been studied extensively in different modalities, including auditory (Ulanovsky et al., 2003; Anderson et al., 2009; Malmierca et al., 2009), visual (Kohn and Movshon, 2003; Movshon and Lennie, 1979), somatosensory (Katz et al., 2006; Liu et al., 2017) and olfactory (Wilson, 1998; Verhagen et al., 2007) systems. SSA has attracted much interest in the past several decades because it was thought to be a potential neuronal correlate of mismatch negativity (MMN) (Ulanovsky et al., 2003; Fishman and Steinschneider, 2012), which has been extensively studied in humans (Näätänen et al., 2007). SSA is also believed to reflect deviance detection (Polterovich et al., 2018; Pérez-González et al., 2021), predictive coding (Parras et al., 2017; Carbajal and Malmierca, 2018) and efficient coding (Wark et al., 2007; Kohn, 2007). Visual and somatosensory neurons exhibit SSA in both spatial (Kohn and Movshon, 2003; Liu et al., 2017) or non-spatial (Movshon and Lennie, 1979; Maravall et al., 2007) domains. In contrast, in the auditory and olfactory systems, SSA has been studied mainly by presenting stimuli with two different frequencies (Natan et al., 2015; Kato et al., 2015; Harpaz et al., 2021) or odorants (Kum et al., 2019).

However, the adaptation alone does not account for the ability of sensory systems to remain vigilant to the environment in spite of repetitive sensory stimulation. In this study, we explored spatial contextual modulations by stimulating neurons in the awake marmoset auditory cortex with sequences of sounds either randomly from various spatial locations (equal-probability mode) or repeatedly from a single location (continuous mode). To our surprise, instead of only inducing adaptation as expected from well-documented SSA literature, repetitive stimulation in the high probability mode from spatial locations away from the center of a neuron's SRF evoked lasting facilitation and adaptation observed by extracellular recordings from single neurons in the auditory cortex. Nearly half of the sampled neuronal population exhibited this spatial facilitation, irrespective of the stimuli type and visibility of the test speaker. The extent of the facilitation decreased with decreasing presentation probability of the test speaker. Intracellular recordings showed that repetitive sound stimulation evoked sustained membrane potential depolarization that was followed by firing rate facilitation. We used computational models to explore neural mechanisms underlying neural facilitation. Taken together, our findings revealed location-specific facilitation (LSF) to repetitively presented sound stimuli in the auditory cortex that has not been observed. This form of spatial contextual modulation may play a role in such functions as detecting the regularity, segregating the sound stream, and solving the cocktail party problem.

RESULTS

Repetitive sound stimulation evoked neural facilitation

We evaluated how extracellularly recorded individual neurons in the auditory cortex responded to broadband sounds played from different speaker locations. Fifteen equally spaced speakers were placed on a semi-spherical surface centered around the animal's head and above the horizontal plane (Figures 1A and 1B). In each test session, we first

Figure 1. Speaker layout and equal-probability sound stimulation

(A) Fifteen equally spaced speakers were placed on a semi-spherical surface centered around the animal's head and above the horizontal plane. View from the front, lateral 35° and elevated 30° (left), and view from directly top (right). Red dots indicate six front speakers, green dots indicate five midline speakers, and brown dots indicate four back speakers.

(B) Three-dimensional front-back space was projected to a two-dimensional plane around midline for displaying purposes.

(C) Spatial receptive field of example Unit M43S-383. White semicircle is the boundary of the front-back space. Black line is the threshold which is defined as the half-maximum firing rate. Black dots indicate fifteen speaker locations.

(D) Spike raster plot of same example neuron at fifteen speaker locations under equal-probability presentation mode. Stimuli at each speaker location were randomly presented ten times.

probed a neuron's spatial selectivity by delivering a frozen wideband noise stimulus from
each speaker location in a randomly shuffled order. We will refer to this stimulation mode
as the *equal-probability presentation mode* for which all locations have the same
occurrence probability of 1/15. Spatial receptive field (SRF) was constructed for each
neuron using the averaged firing rate of the responses to wideband noises in the
equal-probability presentation mode. Figures 1C and 1D show the responses of an
example neuron obtained in the equal-probability presentation mode. This neuron had an
SRF centered around speaker #7 (Figure 1C) and responded to this location with
sustained firing throughout the stimulus duration, and to other locations with onset or
transient firing (Figure 1D). Figure 2A shows firing rate versus speak number for the
equal-probability presentation mode. Speaker #7 evoked the highest firing rate (38.5
spikes/sec) in this neuron, followed by speaker #8 (10.5 spikes/sec). Speakers #14, #13,
#9 and #11 evoked near zero firing rate, and a negative firing rate was observed for
speakers #2, #4, and #5. Because the firing rate for each speaker location is calculated by
subtracting the spontaneous firing rate from the total firing rate, a negative firing rate
indicates inhibition.
17

Figure 2. Repetitive sound stimulation evoked neural facilitation
(A) Firing rate versus speak number of unit M43S-383 recorded under the
equal-probability presentation mode (same data as Figure 1D). Blue dot and orange bar
indicate the target speaker and facilitation threshold (mean + standard deviation),
respectively.
(B), Spike raster of same example neuron tested at speaker #14 under continuous
presentation mode. Red dots indicate spikes belong to the long facilitation phase (i.e., at
least five consecutive trials with firing rates exceeding the facilitation threshold).
(C), Trial-by-trial firing rate of same example neuron. Red dots and line indicate trails
belong to the long facilitation phase. Thick orange line indicates the facilitation threshold.
(D-F), Similar to (A-C), but for Unit M43S-12 that was tested using marmoset vocalization

stimuli.

After the characterization of a neuron's SRF, we further tested each neuron with the

continuous presentation mode in which stimuli were delivered from a speaker location

repeatedly, with each trial separated by an inter-stimulus interval of fixed or variable length

(range: 500 to 5200 ms, see below). Figures 2B and 2C show the responses of the same

neuron depicted in Figures 1C and 1D and Figure 2A to 300 presentations of a 200 ms

frozen wideband noise stimulus delivered from speaker #14 in the continuous

presentation mode. Speaker #14 evoked 0.8 spikes/sec firing rate in the equal

presentation mode (Figure 2A, blue dot) which was considered as the baseline firing rate

of this speaker location. In contrast to the expectation from previous auditory cortex

literature on adaptation, the response of this neuron to the repeated presentations of a

wideband noise stimulus from the same speaker #14 showed epochs consisting of

consecutive trials with substantially higher firing rates than the baseline firing rate (median:

16 spikes/sec, Figure 2C, red dots). Elevated firing rates could be observed in trials long

after the first trial (e.g., between 150th and 250th trials). Also note that during the trials

with elevated firing rates (Figure 2B, red dots), the firing patterns were sustained

throughout the stimulus duration. To further examine the facilitated response in the

continuous presentation mode, we measured the "facilitation phase" to characterize trials

with firing rates exceeding the facilitation threshold which was defined as one standard

deviation above the baseline firing rate (Figures 2A, 2C, orange bar). This neuron

exhibited several facilitation phases lasting up to 42 trials (Figure 2C, red dots). Figures

2D-2F show responses of another example neuron. In the equal-probability presentation

mode, this neuron had weak responses to marmoset vocalization stimuli at speaker #8

(Figure 2D, blue dot, 0.05 spikes/sec). When tested in the continuous presentation mode

with speaker #8 (Figures 2E, 2F), this neuron had weak responses initially, but the

responses gradually built up and eventually led to a facilitation phase (median: 4.8

spikes/sec) lasting 20 trials (Figure 2F, red dots).

Neural facilitation occurred in a variety of stimulus conditions

We tested a total of 104 auditory cortex neurons in four hemispheres of three

marmosets using wideband stimuli, including frozen wideband noises,

amplitude-modulated wideband noises, and marmoset vocalizations. 725 sessions were

tested by both equal-probability and continuous presentation modes. Population statistics

of the facilitation phase are shown in Figure S1A which shows that a facilitation phase

could persist as long as 45 trials. For further analyses, we focused on the sessions that

exhibited facilitation phases lasting at least 5 consecutive trials (129 sessions from 51

neurons). In the majority of sessions, 200 trials were tested (Figure S1B). In most

sessions, it took fewer than 100 trials to achieve the first facilitation phase lasting at least

5 consecutive trials (median: 44 trials) (Figure S1C).

We calculated the proportion of facilitation trials in the continuous presentation mode

for the 129 sessions from 51 neurons that exhibited facilitation phases lasting at least 5

consecutive trials which ranged between 15%-87.5% (median: 32.8%) (Figure 3A, orange

line). As a control, we also calculated the proportion of facilitation trials in the

equal-probability presentation mode for the same group of 51 neurons (Figure 3A, blue

line) which was significantly smaller than that of the continuous presentation mode

Figure 3. Neural facilitation occurred in a variety of stimulus conditions and depends on sound locations

(A) Histogram of the proportion of facilitation trials under the continuous (orange line) and equal-probability (blue line) presentation modes. Only 129 sessions from fifty-one neurons that exhibited facilitation phases lasting at least five consecutive trials were shown.

(B) Histogram of the proportion of facilitation trials under the continuous presentation mode for target speakers located in the front (orange line) and back (blue line). Front speakers: #1, #2, #8, #11, #14, and #15. Back speakers: #3, #4, #5, #6, #7, #9, #10, #12, and #13.

(C) Histogram of the proportion of facilitation trials under the continuous presentation mode using frozen amplitude-modulated wide-band noise or animal vocalization (orange line) and frozen unmodulated wide-band noise (blue line) as stimuli.

(D) Each speaker was assigned a rank number based on its baseline firing rate obtained under the equal-probability presentation mode. Speaker ranked 1st had the highest firing rate. In this example unit M43S-348, speaker #8 ranked 1st and speaker #14 ranked 15th. Orange and blue bars indicate facilitation and adaptation threshold, respectively. Dots and error bars indicate mean ± standard deviation.

(E) Proportion of facilitation (orange dots and lines) and adaptation (blue dots and lines) trials under continuous presentation mode at different speaker ranks for the same example unit. Stimuli at each speaker location were tested 200 times.

(F) Proportion of facilitation (orange dots and lines) and adaptation (blue dots and lines) trials of the population data. All 725 sessions from 104 neurons were shown, regardless of the length of the facilitation phase.

(14.3% vs. 32.8%, p < 0.0001, rank-sum test). The group of 51 neurons scattered across

all recorded cortical areas and did not show any clustering patterns (Figures S2A-S2D).

Most of the neurons (43/51) were recorded at superficial cortical depths (< 1 mm, Figures

S2E, S2F). We attempted to distinguish putative excitatory and inhibitory neurons by their

spike waveform (broad or narrow; Figure S3A) (Mitchell et al., 2007). Spike waveform of

32 neurons had been recorded and 29 neurons had a signal-to-noise ratio higher than 20

dB (Figure S3B). 23 neurons were classified as putative excitatory neurons (yielded 58

sessions) and 6 neurons as putative inhibitory neurons (yielded 12 sessions) (Figure S3C).

The proportions of facilitation trials were similar between putative inhibitory and excitatory

neurons (Figure S3D; 34% vs. 32%, p = 0.2271, rank-sum test). This analysis suggests

that the facilitation can be induced in both putative excitatory and inhibitory neurons.

We conducted control tests to see if the observed facilitation depended on visual

inputs. We found that the facilitation was not limited to speakers located within a

marmoset's visual field (< 90 degrees) (Chaplin et al., 2012) and could be induced from

speaker locations both in front and behind an animal (Figures S4A-S4C). Across all 129

test sessions, the proportions of facilitation trials were similar between front (64 sessions)

and back (65 sessions) speaker locations (Figure 3B; 38% vs. 35%, p = 0.4653, rank-sum

test). The facilitation could still be observed when an animal was tested in the darkness (7

sessions) (Figures S4D-S4F). Interestingly, the proportion of facilitation trials was

significantly higher in the darkness than in the light-on condition (122 sessions) (Figure

S4G; 57% vs. 32%, p < 0.0001, rank-sum test). These results suggest that visual inputs

are not required to induce the facilitation. We also tested the effects of different stimulus

types. Neural facilitation was also observed using unfrozen wideband noises (Figure S4H).

Comparing to frozen wideband noises (83 sessions), slightly larger proportion of

facilitation trials was observed using complex stimuli including amplitude-modulated

frozen wideband noises (37 sessions) or marmoset vocalizations (9 sessions) as stimuli

(Figure 3C; 30% vs. 41%, p = 0.0054, rank-sum test).

Neural facilitation was observed at both short inter-stimulus intervals (ISI) (700 ms,

Figure 2B) and long ISI (1900 ms, Figure 2E). In addition to ISI with a fixed length, we also

tested random ISIs in a subset of sessions. An example is shown in Figure S5A. This

neuron not only showed facilitation at three constant ISIs but also at random ISIs. Across

all 129 test sessions, three groups of ISIs were tested: short (500 ms and 700 ms, 78

sessions), long (>1000 ms, 36 sessions) and random (700 ms to 2200 ms, 15 sessions).

The proportion of facilitation trials was similar between the three ISI groups (Figure S5B;

35%, 41%, and 34%, p = 0.1244, one-way ANOVA).

37 Neural facilitation did not alter a neuron's SRF

Previous studies in both auditory and visual cortices found that after presenting a

stimulus repeatedly, neurons typically exhibited a decrease of response to the stimulus

but an increase of response to other stimuli that are sufficiently different from the repeated

stimulus (Condon and Weinberger, 1991; Dragoi et al., 2000), which suggests changes in

a neuron's receptive field. To investigate whether neural facilitation altered a neuron's SRF,

we compared the SRF measured before and after testing a neuron in the continuous

presentation mode. An example neuron is shown in Figure S6A (same neuron in Figure

1C). After presenting 300 trials of wideband noise at the same location continuously (spike raster shown in Figure 2B), the SRF (Figure S6A, bottom) appeared similar to the SRF measured before (Figure S6A, top). More examples of pre and post SRFs are shown in Figure S6B. To quantitatively characterize SRF changes, we calculated three metrics in 61 neurons in which pre and post SRFs were measured: tuning selectivity, direction selectivity, and correlation coefficient. The average tuning selectivity (Figure S6C, left, 0.153 vs. 0.146) and direction selectivity (Figure S6C, right, 0.477 vs. 0.473) remained similar after tested by the continuous presentation mode. To further quantify the tuning similarity, we computed the pairwise correlation coefficient between each pair of responses to fifteen speaker locations (Figure S6D, left). 75% (197/262) of pair of sessions had a correlation coefficient greater than 0.7 (Figure S6D, right). We also examined if firing rate changed after a neuron was tested by the continuous presentation mode for the highest (1st) and lowest (15th) ranked target speakers (Figure S6E). Both speaker ranks showed similar firing rates before and after being tested by the continuous presentation mode (ranked 1st: 13.8 vs. 13.0 spikes/sec; ranked 15th: 2.9 vs. 2.6 spikes/sec), consistent with the observations on SRF stability (Figures S6A-S6D). These analyses show that the repetitive presentation of stimuli from the same speaker location does not significantly alter the SRF of the neuron being tested.

In the 129 sessions that exhibited facilitation phases lasting for at least 5 consecutive trials when tested by the continuous presentation mode, the average firing rate during the facilitation phases was nearly three times greater than that evoked by the same speaker location under the equal-probability presentation mode (Figure S6F, orange box: 17.5 vs. 6.6 spikes/sec, $p < 0.0001$, one-way ANOVA). The spontaneous firing rates during equal-probability presentation mode, non-facilitation and facilitation phases of the continuous presentation mode were similar (Figure S6F, blue box: 5.1, 5.2, and 6.2 spikes/sec, $p = 0.3924$, one-way ANOVA). Thus, the facilitation phases during the continuous presentation mode were not accompanied by significant changes in spontaneous activities.

Neural facilitation depends on sound locations

Neural facilitation was observed at all tested speaker locations. To investigate the effect of speaker location on the facilitation, each speaker was assigned a rank number based on its baseline firing rate obtained under the equal-probability presentation mode. Speaker ranked 1st had the highest firing rate among all tested speakers and was at or near the center of a neuron's SRF. The lowest ranked speaker usually fell far outside of the SRF and evoked a response not significantly different from the spontaneous activity. The speaker rank of an example neuron is shown in Figure 3D. We found that speakers with lower ranks usually elicited more facilitation phases than speakers with higher ranks when tested under the continuous presentation mode as shown by the example neuron in Figure 3E, orange dots. Note that in this neuron no facilitation was induced by the frontal speaker (speaker #1, rank 5th) and only one facilitation trial (out of 200 trials) was induced by the speaker at the contralateral 90° location (speaker #6, rank 4th). These two locations were commonly used in previous SSA studies (see Discussion). When the test speaker was ranked 15th, 18 of 77 (23.4%) tested sessions had facilitation phases lasting

for at least 5 consecutive trials (Figure S7A) and the average proportion of facilitation trials
was 21.6% (Figure 3F, orange dots). In comparison, when the test speaker was ranked
1st, these statistics dropped to 10.5% (4/38 tested sessions, Figure S7A) and 7.3%
(Figure 3F, orange dots), respectively.
In addition to the facilitation, suppressed responses were also observed in the
continuous presentation mode. We measured the "adaptation phase" to characterize trials
with firing rates lower than the adaptation threshold which was defined as one standard
deviation below the baseline firing rate (Figure 3D, blue bar). At higher ranked speakers,
more adaptation phases were observed than at lower ranked speakers as shown by an
example neuron in Figure 3E, blue dots. Note that 70% of trials (140/200) exhibited
adaptation at the frontal speaker (speaker #1, rank 5th) and 52.5% of trials (105/200)
showed adaptation at the contralateral 90° location (speaker #6, rank 4th). The average
proportion of adaptation trails across all tested sessions was 26.1% when the test speaker
was ranked 1st, whereas that number dropped to 8.9% when the test speaker was ranked
15th (Figure 3F, blue dots). This trend was the opposite of facilitation (Figure 3F, orange
dots).
17
Neural facilitation primarily depends on sound location but not firing rate
We showed above that neural facilitation depends on speaker rank, which was
determined by firing rate in the equal-probability presentation mode at a neuron's
preferred sound level. Sound level was a crucial parameter in influencing the firing rate of
auditory cortex neurons (Sadagopan and Wang, 2008; Wang, 2018). This gives us the
opportunity to determine whether the neural facilitation primarily depends on sound
location or any type of stimuli that could influence the firing rate, including speaker
location and sound level. Therefore, we measured a neuron's responses to wideband
noise delivered from fifteen speaker locations at four sound levels under the
equal-probability presentation mode in a subset of neurons. An example of this analysis is
shown in Figure 4A. We assigned this neuron a speaker rank and sound level rank. In
contrast to previous figures where the speaker rank was determined by firing rate at one
sound level (mostly the best sound level), the speaker rank in this analysis was
determined by the averaged firing rate (Figure 4A, black dots) at four sound levels (Figure
4A, four different color dots). The sound level rank at each speaker location was also
determined by the firing rate in which the 1st ranked sound level had the highest firing rate
among the four tested sound levels. If the neural facilitation was location-dependent, then
the facilitation shall increase from higher ranked speakers to lower ranked speakers, but
not from higher ranked sound levels to lower ranked sound levels. If the neural facilitation
was firing rate-dependent, then the facilitation shall increase from higher ranked speakers
to lower ranked speakers as well as from higher ranked sound levels to lower ranked
sound levels.
We performed the speaker rank and sound level rank analysis in 390 sessions
obtained from 63 neurons. At each sound level rank, the proportion of facilitation trials
tended to be larger at the lower ranked speakers (Figure 4B, fifteen same color dots in x-

Figure 4. Neural facilitation primarily depends on sound location but not firing rate

(A) Example unit M43S-348's averaged responses to wide-band noise played at four sound levels across fifteen speaker locations (violet/brown/green/red dots and lines). Speaker rank was determined by the averaged firing rate at four sound levels (black dots and line).

(B) Proportion of facilitation trials at fifteen speaker ranks (same color dots and lines in x-axis) and four sound level ranks (different color dots and bars in y-axis) were assigned based on the response obtained under the equal-probability mode. Length of upwards arrow is proportional to the proportion of facilitation trials averaged across speakers. Slope of the northeast arrow is proportional to the proportion of facilitation trials averaged across sound levels. Dots and error bars indicate mean - standard deviation of mean.

(C) Similar to (B), but for adaptation. Length of the downwards arrow is proportional to the proportion of adaptation trials averaged across speakers. Slope of the southeast arrow is proportional to the proportion of adaptation trials averaged across sound levels.

axis). When averaged across sound levels, the proportion of facilitation trials was only 3.4%
at the speaker ranked 1st (21 sessions tested) whereas the proportion jumped to 19.1%
when the test speaker was ranked 15th (41 sessions tested) (Figure 4B, slope of
northeast arrow; Figure S7B, orange dots). In contrast, at each speaker rank, the
proportion of facilitation trials did not vary much when the sound level rank changed
(Figure 4B, four different color dots in y-axis). When averaged across speakers, the
proportion of facilitation trials was 12.4% at the sound level ranked 1st (119 sessions
tested) whereas the proportion rose slightly to 13.4% when the sound level was ranked
4th (61 sessions tested) (Figure 4B, length of upwards arrow; Figure S7C, orange dots).

Compared to neural facilitation, the proportion of adaptation trials tended to be larger
at both higher ranked speakers (Figure 4C, fifteen same color dots in x-axis) and higher
ranked sound levels (Figure 4C, four different color dots in the y-axis). When averaged
across sound levels, the proportion of adaptation trials was 18.0% at the speaker ranked
1st and the proportion dropped to 8.9% when the test speaker was ranked 15th (Figure
4C, slope of southeast arrow; Figure S7B, blue dots). When averaged across speakers,
the proportion of adaptation trials was 18.5% at the sound level ranked 1st and the
proportion dropped to 12.3% when the sound level was ranked 4th (Figure 4C, length of
downwards arrow; Figure S7C, blue dots).

These results suggest that neural facilitation is primarily dependent on sound location
but not firing rate. In contrast, neural adaptation is primarily dependent on firing rate which
is expected from previous adaptation literature.

Dependence of neural facilitation on location continuity

As we have shown above, neural facilitation can be induced in the continuous
presentation mode in which the probability of stimuli delivered from a target speaker is
100% (none from other speakers). In the equal-probability presentation mode, the
presentation probability for each speaker is equal to 1/ 15 (number of speakers).
Therefore, the continuous presentation mode provides the location continuity for sound
delivery, whereas the equal-probability presentation mode does not. We further
investigated in a subset of units whether the sound location continuity was necessary to
induce neural facilitation by changing the presentation probability of the target speaker
from 100% to 75%, 50%, 25% and 6.7% (Figure 5A, target speaker: orange square; other
speakers: blue color shapes). An example neuron is shown in Figure 5B. The decrease of
the target speaker's presentation probability to 75%, 50%, and 25% resulted in
increasingly weaker responses of the target speaker (100%: 20, 75%: 20, 50%: 10, and
25%: 6 spikes/sec). Across all 57 test sessions in 12 neurons, the decrease of the
presentation probability of the target speaker from 100% to 75%, 50%, 25% and 6.7%
resulted in weaker response of the target speaker (6.81, 3.43, 1.93, 1.89 and 0.88
spikes/sec, Figure 5C) and the reduction of the proportion of facilitation trials (44%, 23%,
16%, 21% and 14%, Figure 5D). The proportion of facilitation trials lasting for at least 5
consecutive trials showed a similar trend of decrease (Figure 5E, 25%, 12%, 5%, and
0%).

Previous stimulus-specific adaptation (SSA) related studies found that neurons in the
auditory cortex of anesthetized rats were sensitive to statistical regularities: standard and

Figure 5. Dependence of neural facilitation on location continuity

(A) Six stimulus presentation modes were used in our studies. Continuous presentation mode equals to 100% probability presentation mode. 75%, 50% and 25% probabilistic presentation modes play sounds from all fifteen speakers in a randomly shuffled order, while giving the target speaker (orange square) a presentation probability higher than other speakers (blue left-pointing triangle, upward-pointing triangle, hexagram, diamond, downward-pointing triangle, pentagram, right-pointing triangle). For the 50% probability periodic presentation mode (50/p%), the target speaker was interleaved with other speakers. Therefore, the sequence of the target speaker was periodic instead of random. Equal-probability presentation mode equals to 6.7% probability presentation mode.

(B) Firing rate of target speaker (orange dots and lines) and other speakers (blue dots and lines) for example unit M43S-34 under 100% (left), 75% (middle-left), 50% (middle-right), and 25% (right) probability presentation modes. Black dashed line indicates the facilitation threshold.

(C) Firing rate of target speaker for six presentation modes. *, p<0.05, rank-sum test. Dots and error bars indicate mean ± standard deviation.

(D) Proportion of facilitation trials that belong to all facilitation phases at target speaker.

(E) Proportion of facilitation trials that belong to facilitation phases that lasting at least five consecutive trials at target speaker.

deviant tones in random sequences both evoked larger responses than the same tones in
periodic sequences (Yaron et al., 2012; Parras et al. 2017). To investigate whether neural
facilitation is also sensitive to the statical regularity of the target speaker, we played two
different types of sequences with 50% probability of the target speaker. In one sequence,
stimuli from the target speaker and other speakers were randomly arranged, similar to the
75%, 25%, and 6.7% probability mode (Figure 5A, 50). In the other sequence, stimuli from
the target speaker were interleaved with stimuli from other speakers, so that the stimuli
from the target speaker were periodical (Figure 5A, 50/p). In contrast to SSA, we found
that the periodic target speaker sequence evoked significantly stronger responses than
the random target speaker sequence (Figure 5C, 3.91 vs. 1.93 spikes/sec, p = 0.0443,
rank-sum test). The proportions of facilitation trials were similar between random and
periodic target speaker sequence with 50% probability. (Figure 5D, 5E).
13

Repetitive sound stimulation induced sustained membrane potential depolarization

We next asked what are cellular mechanisms underlying the neural facilitation
evoked by repetitive stimuli. In a subset of experiments, we performed intracellular
recordings in awake marmoset (Gao et al. 2016; Gao and Wang, 2019) to examine both
membrane potential and spiking activity during both equal-probability presentation and
continuous presentation modes. Figure 6A shows membrane potential traces of an
example neuron. Figure 6B shows the speaker rank based on firing rate obtained under
the equal-probability presentation mode in this neuron. Our previous analyses of the
spiking activity measured the "facilitation phase" to characterize trials with firing rates
exceeding the facilitation threshold which was defined as one standard deviation above
the baseline firing rate (Figures 2A, 2D, 3D, and 6B, orange bar). Here, we measured the
"depolarization phase" to characterize trials with membrane potential exceeding the
depolarization threshold which was defined as one standard deviation above the baseline
membrane potential (Figure 6C, blue bar). The membrane potential steadily increased
after 40 trials during the continuous presentation (Figure 6D, blue line) which was
accompanied by an increase in spiking activity (Figure 6D, orange line).
We conducted 30 sessions under the continuous presentation mode in 14
intracellularly recorded neurons at different speaker locations. The target speakers were
divided into a low ranked group (10th to 15th) and a high ranked group (1st to 6th). For the
low ranked group, we observed a larger proportion of depolarization trials than
hyperpolarization trials (Figure 6E, gold dots, 43% vs. 19%, p = 0.1810, rank-sum test). In
contrast, the opposite trend was observed for the high ranked group (Figure 6E, violet
dots, 7% vs. 50%, p = 0.0044, rank-sum test). Thus, both the firing rate facilitation and
membrane potential depolarization were influenced by the speaker rank. We further
calculated the membrane potential variation of each trial. Trials were classified into
depolarization, hyperpolarization, or transition groups based on whether they passed the
depolarization threshold, hyperpolarization threshold, or neither. Depolarization trials had
the lowest variation and hyperpolarization trials had the highest variation (Figure 6F, 1.07
vs. 2.97, p < 0.0001, one-way ANOVA).

Figure 6. Repetitive sound stimulation induced sustained membrane potential depolarization

(A) Membrane potential traces of unit M117A-13 recorded under the equal-probability presentation mode at the target speaker location. Each color line indicates one trial.

(B) Firing rate based speaker rank of same neuron obtained under the equal-probability presentation mode. Orange dot and bar indicate the target speaker and its facilitation threshold.

(C) Corresponding membrane potential (spikes removed) at the same speaker rank used in (B). Blue dot and bar indicate the target speaker and its depolarization threshold.

(D) Membrane potential (blue line and dots) and total firing rate (i.e., not minus the spontaneous firing rate, orange line and dots) changes during the continuous presentation mode. Threshold of facilitation (dashed orange line) and depolarization (dashed blue line) were calculated in (B) and (C), respectively.

(E) Target speakers were divided into a low ranked group (10th to 15th, gold dots and bars) and a high ranked group (1st to 6th, violet dots and bars). Dots and error bars indicate mean ± standard deviation of mean.

(F) The standard deviation of membrane potential for depolarization, transition and hyperpolarization trials across all sessions.

(G) Number of trials needed to achieve the first membrane potential depolarization phase and the spiking facilitation phase that both lasting at least five consecutive trials.

(H) Changes in membrane potential magnitude after depolarization (blue dots and line). Black dot indicates the stabilized membrane potential.

For each session, we compared the difference between the number of trials needed
to achieve the first membrane potential depolarization phase and the spiking facilitation
phase that lasted for at least 5 consecutive trials. We found that the depolarization of
membrane potential always preceded the facilitation of spiking activity (Figure 6G, median:
9 trials). Figure 6H shows changes in membrane potential magnitude after depolarization
(seven sessions). The resting membrane potential was -69 mV. After 30 trials, membrane
potential achieved a stable level of -62 mV which was 7 mV depolarized. We will use
these three parameters in our computational models below.
9

Computational models suggest two distinct neural mechanisms underlying location-specific facilitation

Our in vivo extracellular and intracellular recording data showed that both spiking and
subthreshold activity of a neuron could be modulated (facilitation or adaptation of spiking
activity and sustained depolarization or hyperpolarization of membrane potential) by
repetitive stimuli in a location-specific way. We used two computational models to
investigate potential neural mechanisms underlying these observations. In our models,
synaptic inputs were panoramic (i.e., with inputs from every spatial location) based on
observations from previous whole-cell recording studies in rats. Chadderton et al. (2009)
found excitatory postsynaptic potential (EPSP) could be evoked from all spatial locations
in all tested neurons. Kyweriga et al. (2014) found excitatory and inhibitory currents could
be evoked by all interaural level difference (ILD) cues. Panoramic inputs make it possible
for a cortical neuron to generate neural facilitation or sustained depolarization by
amplifying weak responses at unpreferred sound locations.
A conceptual model for location-specific facilitation and adaptation in spiking activity
is shown in Figure S8A. In this model, a neuron receives panoramic excitatory and
inhibitory inputs but with varying strengths according to the speaker ranking as revealed
by the equal probability presentation mode (upper plot). When stimuli are repetitively
presented at the 15[th]-ranked speaker (right plot), stronger inhibitory inputs to the model
neuron would result in larger depression than the depression of weaker excitatory inputs
(Figure S8C). The overall stronger excitation than inhibition would evoke neural facilitation
at this speaker location (Figure S8E). When stimuli are repetitively presented at the
1[st]-ranked speaker (left plot), stronger excitatory inputs to the model neuron would result
in a larger depression than the depression of weaker inhibitory inputs. Then the overall
stronger inhibition than excitation would evoke neural adaptation at this speaker location.
We call this neuron model "EI-LIF model" in which excitatory and inhibitory inputs are
differentially depressed (Figure 7A).
A second conceptual model is shown in Figure S8B for membrane potential. This
model is based on the idea that the brain state is modulated when stimuli switch from
equal-probability presentation mode to continuous presentation mode (see Discussion).
Since neural activity of the network is homeostatically regulated (Pacheco et al. 2019),
sustained depolarization of weaker responses (red line) at the 15[th] ranked speaker will be
accompanied by sustained hyperpolarization of stronger responses (blue line) at the
1[st]-ranked speaker. A sustained depolarization makes it easier to reach the threshold thus
evoke neural facilitation (Figures S8D, S8E). In contrast, a sustained hyperpolarization
45

Figure 7. Computational models suggest two distinct neural mechanisms underlying location-specific facilitation

(A) Inhibitory (blue lines and dots) and excitatory (orange lines and dots) synaptic depression leaky integrate-and-fire model (EI-LIF model). Under the continuous presentation mode, synaptic release probability at five probability presentation modes decreased gradually.

(B) Spike raster plot for one session of EI-model neuron during continuous sound presentation mode. Red dots indicate spikes belong to long facilitation phase (i.e., at least five consecutive trials with firing rates exceeding the facilitation threshold).

(C) Trial-by-trial firing rate from the same session of EI-model neuron. Red dots and line indicate trails belong to the long facilitation phase. Thick orange line indicates the facilitation threshold (i.e., one standard deviation above the baseline firing rate).

(D) Decrease the presentation probability of target speaker resulted in a smaller proportion of facilitation trials for 200 sessions of EI-LIF model neuron (brown dots and bars). Green dots and bars show the data from five probability presentation modes (twelve sessions, same as Figure 5D). Red dots and bars show the data from continuous and equal-probability presentation modes (129 sessions, same as Figure 3A). Dots and error bars indicate mean ± standard deviation.

(E-H), Similar to (A-D) but for membrane potential depolarization leaky integrate-and-fire model (MP-LIF model).

makes it harder to reach the threshold thus evokes neural adaptation. We call this neuron

model "MP-LIF model".

Figures 7B, 7C, 7F, 7G show the neural facilitation simulated with the EI-LIF model

(up) and MP-LIF model (down) in the continuous presentation mode. Notice the elevated

firing rates were observed in tens of trials after the start of continuous sound stimuli

(Figures 7B, 7F, red dots) and epochs consisting of consecutive trials with higher firing

rates than the facilitation threshold (Figures 7C, 7G, red lines). We also simulated neural

adaptation in the EI-LIF model (Figure S8F) and MP-LIF model (Figure S8G). The

threshold of neural facilitation and adaptation were calculated in the equal-probability

presentation mode (gray area). Notice the sparse spikes (Figures S8F, S8G, blue dots)

and epochs consisting of consecutive trials with firing rates lower than the adaptation

threshold (Figures S8F, S8G, blue line).

We further examined whether our models could also simulate the speaker

probability-dependent neural facilitation observed in Figure 5. We hypothesized that the

recovery time of excitatory and inhibitory synaptic release and amplitude of sustained

membrane potential depolarization were proportional to the probability of the target

speaker (Figures 7A, 7E). We found that decreasing the target speaker's presentation

probability reduced the proportion of facilitation trials in the EI-LIF model (Figure 7D, gold

dots) and MP-LIF model (Figure 7H, gold dots). Two models' performances were close to

the experimental results observed in Figure 5D (now shown as Figures 7D, 7H, green

dots). We further compared the proportion of facilitation trials in our models with 129

experimentally tested sessions in the continuous and equal-probability presentation mode

shown in Figure 3A (now shown as Figures 7D, 7H, red dots). The results were also

similar at 100% and 6.7% probabilities (EI-LIF: 39% vs 36% and 18% vs 14%; MP-LIF: 41%

vs 36% and 18% vs 14%). In summary, both EI-LIF and MP-LIF models recapitulated the

neural facilitation and adaptation observed in our experiments and suggest potential

underlying neural mechanisms. Future experimental studies can provide validations of

these suggested mechanisms.

30 DISCUSSION

In this study, we investigated extracellular and intracellular neural responses to

repetitive sound stimulation in the auditory cortex of awake marmoset monkeys. The

major finding of this study is the observation of a novel location-specific facilitation (LSF)

which is dependent on sound location and stimulus presentation mode. LSF raises

questions on the conventional definition of the spatial receptive field as being a static

property of an auditory cortical neuron. LSF is a different phenomenon than the

well-studied stimulus-specific adaptation (SSA). Computational models based on the

synaptic depression or sustained depolarization mechanisms can both reproduce the LSF.

The dependence of facilitation on location continuity and regularity suggests that LSF is a

potential single-neuron substrate of auditory streaming.

42 Implications for the spatial receptive field (SRF) of cortical neurons

The concept of a stable spatial receptive field (SRF) has been a cornerstone of our

understanding of spatial tuning in the central auditory system. Auditory cortex neurons in
anesthetized animals exhibit predominantly broad SRFs that typically increase in size (or
width) as sound level increases (Middlebrooks and Pettigrew 1981; Mrsic-Flogel et al.
2005). In contrast, studies in awake animals have reported restricted SRFs which do not
increase or show less increase in size as sound level increases (Mickey and Middlebrooks
2003; Woods et al. 2006; Zhou and Wang, 2012; Remington and Wang, 2019). It has
been shown that behavior engagement could further decrease the size of SRFs and
therefore sharpen spatial tuning of cortical neurons (Lee and Middlebrooks, 2011; van der
Heijden et al., 2018). The finding of the present study further showed that the spatial
tuning of auditory cortex neurons in awake marmosets is not static in that a non-preferred
spatial location could become responsive under particular conditions. This suggests that
cortical neurons can respond to spatial locations away from the center of SRF dynamically.
When stimuli from other locations were inserted into the repetitively presented sound
sequence from one location, neural facilitation was interrupted and even diminished
(Figure 5). However, neurons still preserve their original SRF after being presented with
repetitive sound stimuli (Figure S6). In contrast, after a repetitive pure tone stimulus is
presented, a neuron changes its spectral receptive field by reducing responses to the
specific tone frequency (Condon and Weinberger, 1991). A non-static SRF could play a
role in spatial and binaural tuning plasticity that is observed in monaural deprived animals
(Popescu and Polley, 2010; Keating et al., 2015).
21
Comparison with stimulus-specific adaptation (SSA) and prediction
Adaptation to repetitive sound stimulation (i.e., a reduction in response to a
high-probability stimulus) by auditory neurons is a commonly observed phenomenon and
has been referred to as stimulus-specific adaptation (SSA) (Harpaz et al., 2021;
Malmierca et al., 2014; Nelken, 2014). In previous studies that demonstrated SSA, both
close field and free field sound stimulation paradigms were used in anesthetized or awake
animals. In close field stimulation, the sound was delivered through a sealed speaker into
the contralateral ear (Condon and Weinberger, 1991; Yaron et al., 2012; Hershenhoren et
al., 2014; Nieto-Diego and Malmierca, 2016) or preferred ear (Ulanovsky et al., 2004). In
free field stimulation, the sound was played from the location contralateral to (Chen et al.,
2015; Kato et al., 2015; Natan et al., 2017), in front of (Natan et al., 2015) or above (Farley
et al., 2010) an animal. Adaptation to repetitive sound stimulation was also observed in
the current study, in particular when stimuli were delivered from preferred sound locations
(i.e., higher ranked speaker locations at or near the center of SRF) (Figures 3E, 3F) or
sound levels (i.e., higher ranked sound levels) (Figure 4C). However, when sound was
delivered from locations away from the center of SRF, our study revealed neural
facilitation to repetitive sound stimulation in the auditory cortex of awake marmosets. To
the best of our knowledge, no previous studies have systematically tested repetitive
sound stimulation across spatial locations.
The most striking difference between the current study and previous SSA studies is
the observation of facilitation instead of adaptation to repetitive stimulation in the auditory
cortex. Although the predominant response in auditory system to high probability stimuli is
adaptation, unadapted and even facilitated responses have been observed in a few

previous studies. Thomas et al. (2012) found that repetitive stimulation did not elicit adaptation in specialized (FM-selective) neurons of bat inferior colliculus (IC). They argued that because in echolocating bats behaviorally relevant sounds are echoes from objects, adaptation to those repetitive echolocation signals that occur with a high probability would be maladaptive during active echolocation. Parras et al. (2017) showed that most neurons on the ascending auditory pathway (IC, auditory thalamus, auditory cortex) of anesthetized rats exhibited repetition suppression. However, repetition enhancement was observed in all three areas. Lesicko et al. 2022 and Kommajosyula et al. 2021 also found repetition enhancement in the IC and auditory thalamus of awake rodents, respectively. This is consistent with our findings that repetitive stimuli evoked facilitation, and periodic target speaker sequences evoked stronger responses than random target speaker sequences (Figure 5C).

In the predictive coding model, the sensory system constantly compares external sensory inputs with internally generated predictions and updates expectations of the environment (Keller and Mrsic-Flogel, 2018). This model requires two classes of neurons: prediction error (PE) neuron and prediction (P) neuron. PE neuron signals a mismatch between predicted and actual sensory inputs, whereas the P neuron signals an expectation of sensory inputs. PE neuron has been observed in the auditory (Eliades and Wang, 2008; Keller and Hahnloser, 2009; Schneider et al., 2018), visual (Keller et al., 2012; Uran et al., 2022) and somatosensory (Ayaz et al., 2019) cortices. In contrast, the P neuron originates from other cortical areas and projects the prediction signal to the PE neuron (Schneider et al., 2014; Leinweber et al., 2017; Garner and Keller, 2022). Instead, our findings suggest that PE (i.e., repetition adaptation) and P (i.e., repetition facilitation) neurons coexist in the auditory cortex. The observation of auditory cortex inactivation reduces or blocks the repetition enhancement in the auditory midbrain and thalamus indicates prediction signals originate from P neurons in the auditory cortex (Lesicko et al., 2022; Kommajosyula et al., 2021).

Together, our finding of neural facilitation to the repetitive sound stimulation provides a complementary contextual modulation effect to the SSA and a new perspective on our current understanding of cortical responses to repetitive stimuli.

Neural mechanisms underlying LSF

We investigated the neural mechanisms underlying LSF with two approaches. Experimentally, we directly recorded the membrane potential from neurons that exhibited LSF in awake marmosets (Figure 6). Computationally, we built a leaky integrate-and-fire (LIF) neuron model and manipulated its excitatory-inhibitory synaptic depression amplitude and recovery time (EI-LIF model) and membrane potential depolarization or hyperpolarization amplitude (MP-LIF model). Both models reproduced LSF and recapitulated the key properties of LSF (Figure 7).

Two mechanisms may account for the LSF observed in this study. One is repetitive stimulus-evoked synaptic depression. Although excitatory and inhibitory synapses are both depressed by repetitive stimuli (Galarreta and Hestrin, 1998), inhibitory synapses may show a larger amplitude of depression than excitatory synapses (Heiss et al., 2008). The imbalance between excitation and inhibition may produce LSF. Two lines of evidence

support this hypothesis. First, the subthreshold activity could be evoked from all tested

sound locations (Chadderton et al., 2009) and strong sound-evoked inhibition is

commonly observed outside of the SRF (Zhou and Wang, 2014; Remington and Wang

2019). Therefore, the panoramic and strong inhibitory inputs are more susceptible to

depression than excitatory inputs. Second, compared to the excitatory inputs, the

inhibitory inputs are more sensitive to context change (Kuchibhotla et al., 2016) and show

stronger depression during the forward masking (Wehr and Zador, 2005), suggesting the

inhibitory inputs are more adjustable than excitatory inputs. Our EI-LIF model could

reproduce the LSF based on the hypothesis that neuron has a stronger depression of their

inhibitory inputs than excitatory inputs, thereby supporting a synaptic depression

mechanism.

Another mechanism is salient stimulus-evoked membrane potential depolarization. A

salient auditory spectrotemporal feature could attract attention automatically (Kayser et al.,

2005; Huang and Elhilali, 2020). A wideband noise used in this study was not salient when

it was presented randomly from different locations to characterize SRF. However, a

wideband noise could become salient when it was repetitively presented from one location

while sounds at all other locations disappeared. This auditory spatial pop-out hypothesis

is similar to visual saliency where a visual item in sharp contrast with its neighboring items

in a simple feature, such as color or orientation, automatically captures attention (Yan et

al., 2018). If a location-specific wideband noise is salient, a more salient sound feature at

the spectrotemporal domain, e.g., amplitude-modulated wideband noise and vocalization,

indeed evoke a larger proportion of facilitation trials than the less salient unmodulated

wideband noise (Figure 3C). In humans, salient auditory stimuli dilate the pupil (Wang et

al., 2014). Pupil dilation is closely correlated with membrane potential depolarization and

a decrease in membrane potential variation (McGinley et al., 2015). Interestingly, we

observed similar changes in membrane potential when neurons exhibit LSF (Figure 6),

suggesting that the ongoing repetitive stimuli are salient to the animals. Importantly, our

MP-LIF model reproduced LSF by incorporating parameters obtained from intracellular

recordings. Together, our intracellular recording data and MP-LIF model simulation

support a salient stimulus-evoked sustained depolarization mechanism underlying the

LSF.

It is not clear whether top-down attention plays a role in LSF. It has been shown that

task engagement could modulate the SRFs (Lee and Middlebrooks, 2011). In the current

study, marmosets passively listened to sound stimuli, though they might have chosen to

pay attention to repeated stimulation from a particular location in the continuous

stimulation mode. However, we did not observe an increase in spontaneous activity when

neural facilitation was observed (Figure S6F), whereas attention tends to increase

spontaneous activity (Luck et al., 1997; Reynolds et al., 2000). Furthermore, we found no

preference of different cell types in exhibiting the LSF (Figure S3D), whereas top-down

attention has stronger modulation over putative inhibitory neurons (Mitchell et al., 2007).

Candidate neural substrate for auditory streaming

43   In a natural environment like at a cocktail party, sounds are often simultaneously and

44 continuously generated by multiple sound sources (Cherry, 1953). One major challenge

for a listener is forming auditory streaming (McDermott, 2009). Streaming requires acoustic cues such as frequency, temporal regularity, and sound location (Shamma and Micheyl, 2010). Over the past decade, a rapidly increasing number of studies have investigated the effect of temporal regularity or repetition for streaming (Bendixen et al., 2010; Andreou et al., 2011). Regular stimulation induces stronger responses than random stimulation when measured with magneto-electro encephalography (M/EEG) and functional MRI (fMRI) in humans (Barascuda et al., 2016; Southwell et al., 2017). The findings that repetitive sound stimulation evoked LSF and regular stimulation evoked a stronger response than random stimulation provide a candidate single-neuron correlate of this perceptual phenomenon. Computational modeling suggests that the change of synaptic efficacy could result in sustained responses to regular stimulation (Auksztulewicz et al., 2017). Interestingly, changing the excitatory and inhibitory synaptic efficacy in our EI-LIF model also generated the LSF. Those two models further suggest that sustained response to regular stimulation and LSF share a similar neural mechanism. Repetition causes the target to pop out from the background and is robust to inattention (McDermott et al., 2011; Masutomi et al., 2016; Mehta et al., 2021). Based on the same pop-out hypothesis, our MP-LIF model could reproduce the LSF. Those similarities suggest that our EI-LIF and MP-LIF models provide a theoretical foundation for both LSF observed in marmosets and enhanced response to regular over random stimulation observed in humans. Together, our findings and models provide valuable new insights into the neural mechanisms of auditory streaming.

METHODS

Animal preparation and experimental setup. Data were collected from five hemispheres of four monkeys (Monkey 1: left, Monkey 2: left and right, Monkey 3: right, Monkey 4: left). All experimental procedures were approved by the Johns Hopkins University Animal Use and Care Committee. These procedures were identical to those described in previous publications from our laboratory (Lu et al. 2001). A typical recording session lasted 3-4 h, during which an animal sat quietly in a specially adapted primate chair with its head immobilized. Throughout the entire recording session, the animal was closely monitored via a video camera by the researcher. The eye position was not controlled, but when the animal closed its eyes for a prolonged period, the experimenter ensured the animal opened its eyes before the next stimulus set was presented.

Experiments were conducted in a double-walled sound-proof chamber (Industrial-Acoustics) with the internal walls and ceiling lined with three-inch acoustic absorption foam (Sonex). Fifteen free-field loudspeakers were placed on a semi-spherical surface centered around the animal's head and above the horizontal plane. The speaker setup was similar to our previous studies (Zhou and Wang, 2012), but with speakers covering the rear sphere. Eight speakers were evenly positioned at 0° elevation, five speakers were evenly spaced at +45° elevation in the frontal hemifield, one speaker was located at +45° elevation in the rear midline, and one speaker was located directly above the animal.

Extracellular and intracellular recordings. Extracellular recording procedures were
identical to those described in our previous publications. A sterile single tungsten
microelectrode (A-M Systems) was held by a micro-manipulator (Narishige) and inserted
nearly perpendicularly into the auditory cortex through a small opening on the skull
(1.0–1.1mm in diameter) and advanced by a hydraulic micro-drive (David Kopf
Instruments). The tip and impedance of the electrode was examined before each
recording session (2-5MΩ impedance). Spikes were detected by a template-based spike
sorter (MSD, Alpha Omega Engineering) and continuously monitored by the experimenter
while data recordings progressed. The raw voltage signal was also recorded. Intracellular
recording procedures were identical to those described in our previous publications (Gao
et al. 2016; Gao and Wang, 2019). The recordings were made in the auditory cortex
through the intact dura using a concentric recording electrode and guide tube assembly.
The sharp recording pipette was made of quartz glass. The guide tube was made of
borosilicate glass. The sharp recording pipette was pulled by a laser puller (P-2000, Sutter
Instrument), and the guide tube was pulled by a conventional puller (P-97, Sutter
Instrument). The electrode assembly was advanced perpendicularly relative to the cortical
surface with a motorized manipulator (DMA1510, Narishige). The electrical signals were
amplified (Axoclamp 2B, Molecular Devices), digitized (RX6, Tucker-Davis Technologies),
and saved using custom programs (MATLAB, Mathworks).
20
Acoustic stimuli. Four different stimulus presentation designs were used. 1) Continuous
(100%): the same stimulus was repeatedly delivered from a fixed speaker location over
many trials. 2) Unequal-probability and random (75%, 50%, and 25%): stimulus was
delivered from multiple speaker locations in a randomly shuffled order, but target speaker
has a higher probability than others. 3) Unequal-probability and periodic (50/p%): stimulus
delivered from the target speaker was interleaved with stimulus delivered from other
speakers, so that the stimulus delivered from the target speaker was periodical. 4) Equal
probability (6.7%): stimulus was delivered from multiple speaker locations in a randomly
shuffled order and all speakers shared the same occurrence probability. For each neuron,
if allowed by the experiment conditions, the equal-probability presentation mode was
tested at multiple separate sessions at different time points and other three presentation
modes were tested between the equal-probability presentation mode.
Stimuli were generated digitally in MATLAB at a sampling rate of 97.7kHz using
custom software, converted to analog signals (RX6, Tucker-Davies Technologies),
attenuated (PA5, Tucker-Davies Technologies), power amplified (Crown Audio), and
played from specified loudspeaker. The sound tokens used included unfrozen wide-band
noise, frozen wide-band noise, amplitude-modulated wide-band noise and
species-specific vocalizations. Sessions collected under continuous unfrozen wide-band
noise stimuli were used only in Figure S4H. Fixed inter-stimulus intervals (ISI) were used
in four presentation modes. A variety of random ISI were used only in continuous
presentation mode. The shortest ISI was 500ms, and the longest ISI was 5200ms.
Rate-level function was used to find the best sound level of tested neurons. Most neurons
were tested using best sound level, except sound level rank experiments where four
different sound levels were used. The same sound level was used when comparing

different presentation modes.

Characterization of spatial receptive fields. Total firing rates were calculated over a

time window beginning 15ms after stimulus onset and 50ms after stimulus offset. Total

firing rates subtracted by the spontaneous rate was the firing rate. SRF characterization

was identical to our previous studies (Remington and Wang, 2019). The threshold was the

half maximal firing rates. Tuning selectivity was defined as the number of areas that have

higher firing rates than the threshold divided by the total number of areas. Direction

selectivity was defined as the product of every area, unit vector and firing rate divided by

the product of every area and firing rate. If a neuron only responded to contralateral and

ipsilateral 90° at horizontal plane and have equal firing rates, then the direction selectivity

was zero. In the plotted SRF, the location of white color dot indicated the preferred sound

location, the dot diameter was proportional to the direction selectivity, the black thick line

was the half-maximum threshold of SRF, area encircled by the threshold was the

reciprocal of tuning selectivity.

Identification of cortical areas, layers and cell types. We used the best frequency (BF)

of neurons to identify the subregions of auditory cortex. For the neurons significantly

responding to at least one tone stimulus played at the front speaker, we specified the

frequency of the tone stimulus that evoked the maximum response rate as the neuron's

BF. Marmoset auditory cortex is situated largely ventral to the lateral sulcus and exhibits a

topographical frequency gradient along the rostral–caudal axis. The boundary between

primary auditory cortex (A1) and the caudal area (caudal-medial and caudal-lateral belt)

can be identified by an abrupt decrease of BF at the high-frequency (caudal) border of A1.

A1 was further divided into the low-frequency (≤8KHz) rostral A1 and high-frequency

(>8KHz) caudal A1 along the rostral-caudal axis. First spike depth was the absolute depth

where the first spike was detected from this electrode, and was depended on thickness of

dura, variations in granulation tissue, proximity to the curvature of the sulcus, and

orthogonality of the electrode penetration to the cortical surface. We used the trough to

peak duration of spike waveforms to identify putative excitatory and inhibitory neurons

(Mitchell et al., 2007). Before analyzing the spike duration, we calculated the signal to

noise ratio of spike waveforms (Sabyasachi and Wang, 2012), which was defined as the

action potential peak to peak height divided by the standard deviation of the background

noise over 1ms preceding all spikes ($20 \times \log10(AP_{peak-peak}/Noise_{SD})$).

The proportion of facilitation, adaptation, depolarization and hyperpolarization

trials. The facilitation phase was defined as the trials whose firing rates were at least one

standard deviation above the mean firing rate evoked under the equal-probability

presentation mode. The sum of facilitation trials divided by the total trial number in each

session was defined as the proportion of facilitation trials. The adaptation phase was

defined as the trials whose firing rates were at least one standard deviation below the

mean firing rate evoked under the equal-probability presentation mode. To analyze the

membrane potential change, spikes were removed from voltage signal with a time window

of 3ms centered around the spike peak. The proportion of depolarization and

hyperpolarization trials were calculated similar to the proportion of facilitation and
adaptation trials.
3
Speaker rank and sound level rank. Speaker tested in each continuous presentation
mode was assigned a rank number based on its firing rate obtained under the
equal-probability presentation mode. Speaker ranked 1st and 15th had the highest and
lowest firing rate among all 15 speakers tested, respectively. When the SRF has a single
peak, speaker ranked 1st and 15th usually had the closest and farthest distance to the
preferred direction, respectively. When the SRF has multiple peaks, low ranked speaker
may next to the preferred direction occasionally. We used speaker firing rate rank instead
of distance rank for two reasons: one, the SRF of some neurons was quite dispersive,
thus it was inaccurate to compute the distance between the target speaker and the SRF
center; and two, the SRF center was usually determined by several high ranked speakers.
The contribution of low ranked speakers was not considered when using the distance rank.
For sound level rank, we measured neurons' responses to 200ms wide-band noise played
at four sound levels under the equal-probability presentation mode. These four sound
levels were a series of fixed values with an interval of 20dB.
18
Computational models of location-specific facilitation. Computational models that
recaptured the location-specific facilitation (LSF) phenomenon were based on two neural
mechanisms (Figures S8A-S8D): excitatory and inhibitory synaptic depression (EI-LIF
model, see equations 4 and 5) and membrane potential depolarization and
hyperpolarization (MP-LIF model, see equation 6 and 7). EI-LIF model dynamically
changed the depression amplitude of synaptic vesicles release and exponential recovery
time constant. MP-LIF model dynamically changed the resting potential and spiking
threshold. In the LSF, the parameters were modulated by the probability of presentation
speaker, e.g., 100% for continuous presentation mode and 6.7% for equal-probability
presentation mode.
The membrane potential V_{t+1} of a leaky integrate-and-fire (LIF) neuron at time step
Δt was:
31
$$V_{t+1} = -\frac{\Delta t}{C}\left[g_{e_t}(V_t - E_e) + g_{i_t}(V_t - E_i) + g_{rest}(V_t - E_{rest})\right] + V_t + \sigma_s \omega_n \sqrt{\Delta t} \quad (1)$$
g_{e_t} and g_{i_t} was the excitatory and inhibitory synaptic conductance (see equations 2 and
3). C, E_e, E_i and g_{rest} was the membrane capacitance, excitatory reversal potential,
inhibitory reversal potential, and leak conductance. Those values were obtained from the
in vivo whole-cell recording in the auditory cortex of anesthetized rats (Wehr and Zador,
2003). Gaussian noise $\sigma_s \omega_n$ was added to generate the spontaneous firing (Lee et al.,
2020). Action potential was evoked when the V_{t+1} reached the spike threshold V_{spike}.
V_{t+1} was reset to E_{rest} after the action potential. E_{rest} was obtained from our
intracellular studies (Figure 6H). It was fixed in the EI-LIF model but was dynamically
modulated in MP-LIF model. V_{spike} is the sum of threshold above resting potential V_{th}
and E_{rest}. It was fixed in the EI-LIF model but was modulated in MP-LIF model.
Excitatory conductance g_{e_t} and inhibitory conductance g_{i_t} were:

$$g_{e_t} = P_{rel\_e} r N_e \Delta t e^{\frac{\Delta t}{\tau}} + \sigma_c \omega_n \quad (2)$$

$$g_{i_t} = P_{rel\_i} N_i \Delta t e^{\frac{\Delta t}{\tau}} + \sigma_c \omega_n \quad (3)$$

P_{rel_e} and P_{rel_i} were the excitatory and inhibitory synaptic release probability, respectively (see equations 4 and 5). P_{rel_e} and P_{rel_i} were modulated in EI-LIF model but fixed to one in MP-LIF model. The inhibitory and excitatory inputs have the same strength and occurred simultaneously, so the inhibitory to excitatory ratio r equal to one and the inhibitory to excitatory delay d (not shown in the equation) equal to zero. The number of excitatory inputs N_e, inhibitory input N_i and time constant τ were fixed and conduction noise $\sigma_c \omega_n$ was added to generate the spontaneous firing (Wehr and Zador, 2003).

For the EI-LIF model, in each trial T, the excitatory and inhibitory synaptic release probability $P_{rel\_e_{t+1}}$ and $P_{rel\_i_{t+1}}$ were:

$$P_{rel\_e_{t+1}} = 1 + \left((1 - A_e) P_{rel\_e_t} - 1 \right) e^{\frac{-\Delta t}{P_s \tau e}} \quad (4)$$

$$P_{rel\_i_{t+1}} = 1 + \left((1 - A_i) P_{rel\_i_t} - 1 \right) e^{\frac{-\Delta t}{P_s \tau_i}} \quad (5)$$

A_e and A_i was the excitatory and inhibitory synaptic depression amplitude, respectively. A_i was larger than A_e because we observed facilitation instead of adaptation in the continuous presentation mode. In addition, Heiss et al., 2008 found that inhibition adapts more than excitation when repetitively stimulating the whisker. A_e and A_i were both fixed in the LSF. Relatively small depression amplitude was chosen due to the slow facilitation processes observed in the recording data.

τ_e and τ_i was the excitatory and inhibitory recovery time constant, respectively. τ_e was longer than τ_i because numerous studies found that inhibitory synapses have a quick recovery than excitatory synapses (Galarreta and Hestrin, 1998; Varela et al., 1999). In the different probability presentation mode, facilitation percent and firing rate decreased when the probability of target speaker decreased. Since the time constant was stimulus frequency dependent (Galarreta and Hestrin, 1998), therefore the τ_e and τ_i were scaled by the presentation probability P_s which resulted in higher synaptic release probability, i.e., less adaptation, when the presentation probability was low. Time constant can across multiple time scales from hundreds of milliseconds to tens of seconds (Varela et al., 1997; Ulanovsky et al., 2004). Since 44 trials were required to reach the first long facilitation phase, τ_e and τ_i were chosen so that the probability of release was stable after 40 trials.

Each session is composed of randomly presented trials T_r for computing the facilitation and adaptation threshold and continuously presented trials T_c for computing the facilitation percent, adaptation percent, and firing rate. The median firing rate in the 100% probability presentation mode was 12 spikes per second. Therefore, the number of stimulus count N_{SC} was chosen so that the average firing rate in the EI-LIF model could match the firing rate in the recording data. Notice that N_{SC} was Poisson distributed and not every stimulus input could evoke a spike output in the LIF neuron. We run EI-LIF model for two hundred sessions for every presentation probability.

For the MP-LIF model, in each trial T, the dynamic resting membrane potential

E_{rest_MP} and spike threshold V_{spike} were:

$$E_{rest\_MP} = \begin{cases} E_{rest}, \ 1 \leq T < T_r \\ E_{rest} : P_s M/(T_{th} - 1) : E_{rest} + P_s M, \ T_r + 1 \leq T < T_r + T_{th} \\ E_{rest} + P_s M, \ T_r + T_{th} + 1 \leq T < T_r + T_c \end{cases} \quad (6)$$

$$V_{spike} = \begin{cases} V_{th} + E_{rest}, \ 1 \leq T < T_r \\ V_{th} + E_{rest} : P_s MS/(T_{th} - 1) : E_{rest} + P_s MS, \ T_r + 1 \leq T < T_r + T_{th} \\ V_{th} + E_{rest} + P_s MS, \ T_r + T_{th} + 1 \leq T < T_r + T_c \end{cases} \quad (7)$$

The depolarization value M and the number of trials to reach the stabilized depolarization

value T_{th} were obtained from our intracellular recordings. Since lower presentation

probability evoked less neural facilitation, therefore M was scaled by the presentation

probability P_s . Spike threshold V_{spike} was further modulated by the spike threshold scale

S . Almost no spike was evoked when S equal to one but excessive spikes were evoked

when S equal to zero. Therefore, S and stimulus count N_{SC} were chosen so that the

average firing rate matched the recording data. We also run MP-LIF model for two

hundred sessions at every presentation probability. Table S1 listed the names of

parameters and corresponding values used in EI-LIF and MP-LIF model neurons

14 DATA AND CODE AVAILABILITY

Source data for generating Figure 1 to Figure 7 is publicly available

(https://github.com/ccg1988/LSF_Figures). Codes (MATLAB R2022a) for generating

Figure 1 to Figure 7 and computational models is publicly available

(https://github.com/ccg1988/LSF_Figures; https://github.com/ccg1988/LSF_Model).

Codes (MATLAB R2012a) for controlling the TDT sound presentation and data acquisition

system are freely available upon request from the corresponding author.

REFERENCES

Alitto, H.J., and Usrey, W.M. (2008). Origin and dynamics of extraclassical suppression in the lateral geniculate nucleus of the macaque monkey. Neuron *57*, 135-146. 10.1016/j.neuron.2007.11.019.

Andreou, L.V., Kashino, M., and Chait, M. (2011). The role of temporal regularity in auditory segregation. Hear Res *280*, 228-235. 10.1016/j.heares.2011.06.001.

Auksztulewicz, R., Barascud, N., Cooray, G., Nobre, A.C., Chait, M., and Friston, K. (2017). The Cumulative Effects of Predictability on Synaptic Gain in the Auditory Processing Stream. J Neurosci *37*, 6751-6760. 10.1523/JNEUROSCI.0291-17.2017.

Ayaz, A., Stauble, A., Hamada, M., Wulf, M.A., Saleem, A.B., and Helmchen, F. (2019). Layer-specific integration of locomotion and sensory information in mouse barrel cortex. Nat Commun *10*, 2585. 10.1038/s41467-019-10564-8.

Barascud, N., Pearce, M.T., Griffiths, T.D., Friston, K.J., and Chait, M. (2016). Brain responses in humans reveal ideal observer-like sensitivity to complex acoustic patterns. Proc Natl Acad Sci U S A *113*, E616-625. 10.1073/pnas.1508523113.

Bartlett, E.L., and Wang, X. (2005). Long-lasting modulation by stimulus context in primate auditory cortex. J Neurophysiol *94*, 83-104. 10.1152/jn.01124.2004.

Bendixen, A., Denham, S.L., Gyimesi, K., and Winkler, I. (2010). Regular patterns stabilize auditory streams. J Acoust Soc Am *128*, 3658-3666. 10.1121/1.3500695.

Carbajal, G.V., and Malmierca, M.S. (2018). The Neuronal Basis of Predictive Coding Along the Auditory Pathway: From the Subcortical Roots to Cortical Deviance Detection. Trends Hear *22*, 2331216518784822. 10.1177/2331216518784822.

Chadderton, P., Agapiou, J.P., McAlpine, D., and Margrie, T.W. (2009). The synaptic representation of sound source location in auditory cortex. J Neurosci *29*, 14127-14135. 10.1523/JNEUROSCI.2061-09.2009.

Chaplin, T.A., Yu, H.H., and Rosa, M.G. (2013). Representation of the visual field in the primary visual area of the marmoset monkey: magnification factors, point-image size, and proportionality to retinal ganglion cell density. J Comp Neurol *521*, 1001-1019. 10.1002/cne.23215.

Chen, I.W., Helmchen, F., and Lutcke, H. (2015). Specific Early and Late Oddball-Evoked Responses in Excitatory and Inhibitory Neurons of Mouse Auditory Cortex. J Neurosci *35*, 12560-12573. 10.1523/JNEUROSCI.2240-15.2015.

Cherry, E.C. (1953). Some experiments on the Recognition of Speech with one or two ears. J Acoust Soc Am.

Condon, C.D., and Weinberger, N.M. (1991). Habituation produces frequency-specific plasticity of receptive fields in the auditory cortex. Behav Neurosci *105*, 416-430. 10.1037//0735-7044.105.3.416.

Dragoi, V., Sharma, J., and Sur, M. (2000). Adaptation-induced plasticity of orientation tuning in adult visual cortex. Neuron *28*, 287-298. 10.1016/s0896-6273(00)00103-3.

Eliades, S.J., and Wang, X. (2008). Neural substrates of vocalization feedback monitoring in primate auditory cortex. Nature *453*, 1102-1106. 10.1038/nature06910.

Farley, B.J., Quirk, M.C., Doherty, J.J., and Christian, E.P. (2010). Stimulus-specific adaptation in auditory cortex is an NMDA-independent process distinct from the sensory novelty encoded

by the mismatch negativity. J Neurosci *30*, 16475-16484.

10.1523/JNEUROSCI.2793-10.2010.

Fishman, Y.I., and Steinschneider, M. (2012). Searching for the mismatch negativity in primary auditory cortex of the awake monkey: deviance detection or stimulus specific adaptation? J Neurosci *32*, 15747-15758. 10.1523/JNEUROSCI.2835-12.2012.

Fitzpatrick, D.C., Kuwada, S., Kim, D.O., Parham, K., and Batra, R. (1999). Responses of neurons to click-pairs as simulated echoes: auditory nerve to auditory cortex. J Acoust Soc Am *106*, 3460-3472. 10.1121/1.428199.

Galarreta, M., and Hestrin, S. (1998). Frequency-dependent synaptic depression and the balance of excitation and inhibition in the neocortex. Nat Neurosci *1*, 587-594. 10.1038/2822.

Ganmor, E., Katz, Y., and Lampl, I. (2010). Intensity-dependent adaptation of cortical and thalamic neurons is controlled by brainstem circuits of the sensory pathway. Neuron *66*, 273-286. 10.1016/j.neuron.2010.03.032.

Gao, L., Kostlan, K., Wang, Y., and Wang, X. (2016). Distinct Subthreshold Mechanisms Underlying Rate-Coding Principles in Primate Auditory Cortex. Neuron *91*, 905-919. 10.1016/j.neuron.2016.07.004.

Gao, L., and Wang, X. (2019). Subthreshold Activity Underlying the Diversity and Selectivity of the Primary Auditory Cortex Studied by Intracellular Recordings in Awake Marmosets. Cereb Cortex *29*, 994-1005. 10.1093/cercor/bhy006.

Garner, A.R., and Keller, G.B. (2022). A cortical circuit for audio-visual predictions. Nat Neurosci *25*, 98-105. 10.1038/s41593-021-00974-7.

Harpaz, M., Jankowski, M.M., Khouri, L., and Nelken, I. (2021). Emergence of abstract sound representations in the ascending auditory system. Prog Neurobiol *202*, 102049. 10.1016/j.pneurobio.2021.102049.

Heiss, J.E., Katz, Y., Ganmor, E., and Lampl, I. (2008). Shift in the balance between excitation and inhibition during sensory adaptation of S1 neurons. J Neurosci *28*, 13320-13330. 10.1523/JNEUROSCI.2646-08.2008.

Hershenhoren, I., Taaseh, N., Antunes, F.M., and Nelken, I. (2014). Intracellular correlates of stimulus-specific adaptation. J Neurosci *34*, 3303-3319. 10.1523/JNEUROSCI.2166-13.2014.

Huang, N., and Elhilali, M. (2020). Push-pull competition between bottom-up and top-down auditory attention to natural soundscapes. Elife *9*. 10.7554/eLife.52984.

Huang, X., Rangel, M., Briggman, K.L., and Wei, W. (2019). Neural mechanisms of contextual modulation in the retinal direction selective circuit. Nat Commun *10*, 2431. 10.1038/s41467-019-10268-z.

Kato, H.K., Chu, M.W., Isaacson, J.S., and Komiyama, T. (2012). Dynamic sensory representations in the olfactory bulb: modulation by wakefulness and experience. Neuron *76*, 962-975. 10.1016/j.neuron.2012.09.037.

Kato, H.K., Gillet, S.N., and Isaacson, J.S. (2015). Flexible Sensory Representations in Auditory Cortex Driven by Behavioral Relevance. Neuron *88*, 1027-1039. 10.1016/j.neuron.2015.10.024.

Katz, Y., Heiss, J.E., and Lampl, I. (2006). Cross-whisker adaptation of neurons in the rat barrel cortex. J Neurosci *26*, 13363-13372. 10.1523/JNEUROSCI.4056-06.2006.

Kayser, C., Petkov, C.I., Lippert, M., and Logothetis, N.K. (2005). Mechanisms for allocating auditory attention: an auditory saliency map. Curr Biol *15*, 1943-1947.

10.1016/j.cub.2005.09.040.

Keating, P., Dahmen, J.C., and King, A.J. (2015). Complementary adaptive processes

contribute to the developmental plasticity of spatial hearing. Nat Neurosci *18*, 185-187.

10.1038/nn.3914.

Keller, A.J., Dipoppa, M., Roth, M.M., Caudill, M.S., Ingrosso, A., Miller, K.D., and Scanziani, M.

(2020). A Disinhibitory Circuit for Contextual Modulation in Primary Visual Cortex. Neuron *108*,

1181-1193 e1188. 10.1016/j.neuron.2020.11.013.

Keller, G.B., Bonhoeffer, T., and Hubener, M. (2012). Sensorimotor mismatch signals in

primary visual cortex of the behaving mouse. Neuron *74*, 809-815.

10.1016/j.neuron.2012.03.040.

Keller, G.B., and Hahnloser, R.H. (2009). Neural processing of auditory feedback during vocal

practice in a songbird. Nature *457*, 187-190. 10.1038/nature07467.

Keller, G.B., and Mrsic-Flogel, T.D. (2018). Predictive Processing: A Canonical Cortical

Computation. Neuron *100*, 424-435. 10.1016/j.neuron.2018.10.003.

Kohn, A. (2007). Visual adaptation: physiology, mechanisms, and functional benefits. J

Neurophysiol *97*, 3155-3164. 10.1152/jn.00086.2007.

Kohn, A., and Movshon, J.A. (2003). Neuronal adaptation to visual motion in area MT of the

macaque. Neuron *39*, 681-691. 10.1016/s0896-6273(03)00438-0.

Kommajosyula, S.P., Bartlett, E.L., Cai, R., Ling, L., and Caspary, D.M. (2021). Corticothalamic

projections deliver enhanced responses to medial geniculate body as a function of the

temporal reliability of the stimulus. J Physiol *599*, 5465-5484. 10.1113/JP282321.

Kuchibhotla, K.V., Gill, J.V., Lindsay, G.W., Papadoyannis, E.S., Field, R.E., Sten, T.A., Miller,

K.D., and Froemke, R.C. (2017). Parallel processing by cortical inhibition enables

context-dependent behavior. Nat Neurosci *20*, 62-71. 10.1038/nn.4436.

Kum, J., Kim, J.W., Braubach, O., Ha, J.G., Cho, H.J., Kim, C.H., Han, H.B., Choi, J.H., and

Yoon, J.H. (2019). Neural Dynamics of Olfactory Perception: Low- and High-Frequency

Modulations of Local Field Potential Spectra in Mice Revealed by an Oddball Stimulus. Front

Neurosci *13*, 478. 10.3389/fnins.2019.00478.

Kyweriga, M., Stewart, W., Cahill, C., and Wehr, M. (2014). Synaptic mechanisms underlying

interaural level difference selectivity in rat auditory cortex. J Neurophysiol *112*, 2561-2571.

10.1152/jn.00389.2014.

Lee, C.C., and Middlebrooks, J.C. (2011). Auditory cortex spatial sensitivity sharpens during

task performance. Nat Neurosci *14*, 108-114. 10.1038/nn.2713.

Leinweber, M., Ward, D.R., Sobczak, J.M., Attinger, A., and Keller, G.B. (2017). A

Sensorimotor Circuit in Mouse Cortex for Visual Flow Predictions. Neuron *96*, 1204.

10.1016/j.neuron.2017.11.009.

Lesicko, A.M.H., Angeloni, C.F., Blackwell, J.M., De Biasi, M., and Geffen, M.N. (2022).

Corticofugal regulation of predictive coding. Elife *11*. 10.7554/eLife.73289.

Liu, C., Foffani, G., Scaglione, A., Aguilar, J., and Moxon, K.A. (2017). Adaptation of Thalamic

Neurons Provides Information about the Spatiotemporal Context of Stimulus History. J

Neurosci *37*, 10012-10021. 10.1523/JNEUROSCI.0637-17.2017.

Luck, S.J., Chelazzi, L., Hillyard, S.A., and Desimone, R. (1997). Neural mechanisms of

spatial selective attention in areas V1, V2, and V4 of macaque visual cortex. J Neurophysiol *77*,

24-42. 10.1152/jn.1997.77.1.24.

Malmierca, M.S., and Auksztulewicz, R. (2021). Stimulus-specific adaptation, MMN and predictive coding. Hear Res *399*, 108076. 10.1016/j.heares.2020.108076.

Malmierca, M.S., Sanchez-Vives, M.V., Escera, C., and Bendixen, A. (2014). Neuronal adaptation, novelty detection and regularity encoding in audition. Front Syst Neurosci *8*, 111. 10.3389/fnsys.2014.00111.

Maravall, M., Petersen, R.S., Fairhall, A.L., Arabzadeh, E., and Diamond, M.E. (2007). Shifts in coding properties and maintenance of information transmission during adaptation in barrel cortex. PLoS Biol *5*, e19. 10.1371/journal.pbio.0050019.

Masutomi, K., Barascud, N., Kashino, M., McDermott, J.H., and Chait, M. (2016). Sound segregation via embedded repetition is robust to inattention. J Exp Psychol Hum Percept Perform *42*, 386-400. 10.1037/xhp0000147.

McDermott, J.H. (2009). The cocktail party problem. Curr Biol *19*, R1024-1027. 10.1016/j.cub.2009.09.005.

McDermott, J.H., Wrobleski, D., and Oxenham, A.J. (2011). Recovering sound sources from embedded repetition. Proc Natl Acad Sci U S A *108*, 1188-1193. 10.1073/pnas.1004765108.

McGinley, M.J., David, S.V., and McCormick, D.A. (2015). Cortical Membrane Potential Signature of Optimal States for Sensory Signal Detection. Neuron *87*, 179-192. 10.1016/j.neuron.2015.05.038.

Mehta, A.H., Feng, L., and Oxenham, A.J. (2021). Neural auditory contrast enhancement in humans. Proc Natl Acad Sci U S A *118*. 10.1073/pnas.2024794118.

Mickey, B.J., and Middlebrooks, J.C. (2003). Representation of auditory space by cortical neurons in awake cats. J Neurosci *23*, 8649-8663.

Mickey, B.J., and Middlebrooks, J.C. (2005). Sensitivity of auditory cortical neurons to the locations of leading and lagging sounds. J Neurophysiol *94*, 979-989. 10.1152/jn.00580.2004.

Middlebrooks, J.C., and Pettigrew, J.D. (1981). Functional classes of neurons in primary auditory cortex of the cat distinguished by sensitivity to sound location. J Neurosci *1*, 107-120.

Mirabella, G., Battiston, S., and Diamond, M.E. (2001). Integration of multiple-whisker inputs in rat somatosensory cortex. Cereb Cortex *11*, 164-170. 10.1093/cercor/11.2.164.

Mitchell, J.F., Sundberg, K.A., and Reynolds, J.H. (2007). Differential attention-dependent response modulation across cell classes in macaque visual area V4. Neuron *55*, 131-141. 10.1016/j.neuron.2007.06.018.

Movshon, J.A., and Lennie, P. (1979). Pattern-selective adaptation in visual cortical neurones. Nature *278*, 850-852. 10.1038/278850a0.

Mrsic-Flogel, T.D., King, A.J., and Schnupp, J.W. (2005). Encoding of virtual acoustic space stimuli by neurons in ferret primary auditory cortex. J Neurophysiol *93*, 3489-3503. 10.1152/jn.00748.2004.

Muller, J.R., Metha, A.B., Krauskopf, J., and Lennie, P. (1999). Rapid adaptation in visual cortex to the structure of images. Science *285*, 1405-1408. 10.1126/science.285.5432.1405.

Naatanen, R., Paavilainen, P., Rinne, T., and Alho, K. (2007). The mismatch negativity (MMN) in basic research of central auditory processing: a review. Clin Neurophysiol *118*, 2544-2590. 10.1016/j.clinph.2007.04.026.

Natan, R.G., Briguglio, J.J., Mwilambwe-Tshilobo, L., Jones, S.I., Aizenberg, M., Goldberg, E.M., and Geffen, M.N. (2015). Complementary control of sensory adaptation by two types of cortical interneurons. Elife *4*. 10.7554/eLife.09868.

Natan, R.G., Rao, W., and Geffen, M.N. (2017). Cortical Interneurons Differentially Shape Frequency Tuning following Adaptation. Cell Rep *21*, 878-890. 10.1016/j.celrep.2017.10.012.

Nelken, I. (2014). Stimulus-specific adaptation and deviance detection in the auditory system: experiments and models. Biol Cybern *108*, 655-663. 10.1007/s00422-014-0585-7.

Nelken, I., Yaron, A., Polterovich, A., and Hershenhoren, I. (2013). Stimulus-specific adaptation beyond pure tones. Adv Exp Med Biol *787*, 411-418. 10.1007/978-1-4614-1590-9_45.

Nelson, P.C., Smith, Z.M., and Young, E.D. (2009). Wide-dynamic-range forward suppression in marmoset inferior colliculus neurons is generated centrally and accounts for perceptual masking. J Neurosci *29*, 2553-2562. 10.1523/JNEUROSCI.5359-08.2009.

Nieto-Diego, J., and Malmierca, M.S. (2016). Topographic Distribution of Stimulus-Specific Adaptation across Auditory Cortical Fields in the Anesthetized Rat. PLoS Biol *14*, e1002397. 10.1371/journal.pbio.1002397.

Parras, G.G., Nieto-Diego, J., Carbajal, G.V., Valdes-Baizabal, C., Escera, C., and Malmierca, M.S. (2017). Neurons along the auditory pathway exhibit a hierarchical organization of prediction error. Nat Commun *8*, 2148. 10.1038/s41467-017-02038-6.

Perez-Gonzalez, D., Parras, G.G., Morado-Diaz, C.J., Aedo-Sanchez, C., Carbajal, G.V., and Malmierca, M.S. (2021). Deviance detection in physiologically identified cell types in the rat auditory cortex. Hear Res *399*, 107997. 10.1016/j.heares.2020.107997.

Petersen, R.S., Panzeri, S., and Maravall, M. (2009). Neural coding and contextual influences in the whisker system. Biol Cybern *100*, 427-446. 10.1007/s00422-008-0290-5.

Polterovich, A., Jankowski, M.M., and Nelken, I. (2018). Deviance sensitivity in the auditory cortex of freely moving rats. PLoS One *13*, e0197678. 10.1371/journal.pone.0197678.

Popescu, M.V., and Polley, D.B. (2010). Monaural deprivation disrupts development of binaural selectivity in auditory midbrain and cortex. Neuron *65*, 718-731. 10.1016/j.neuron.2010.02.019.

Reale, R.A., and Brugge, J.F. (2000). Directional sensitivity of neurons in the primary auditory (AI) cortex of the cat to successive sounds ordered in time and space. J Neurophysiol *84*, 435-450. 10.1152/jn.2000.84.1.435.

Remington, E.D., and Wang, X. (2019). Neural Representations of the Full Spatial Field in Auditory Cortex of Awake Marmoset (Callithrix jacchus). Cereb Cortex *29*, 1199-1216. 10.1093/cercor/bhy025.

Reynolds, J.H., Pasternak, T., and Desimone, R. (2000). Attention increases sensitivity of V4 neurons. Neuron *26*, 703-714. 10.1016/s0896-6273(00)81206-4.

Sadagopan, S., and Wang, X. (2008). Level invariant representation of sounds by populations of neurons in primary auditory cortex. J Neurosci *28*, 3415-3426. 10.1523/JNEUROSCI.2743-07.2008.

Schneider, D.M., Nelson, A., and Mooney, R. (2014). A synaptic and circuit basis for corollary discharge in the auditory cortex. Nature *513*, 189-194. 10.1038/nature13724.

Schneider, D.M., Sundararajan, J., and Mooney, R. (2018). A cortical filter that learns to suppress the acoustic consequences of movement. Nature *561*, 391-395. 10.1038/s41586-018-0520-5.

Shamma, S.A., and Micheyl, C. (2010). Behind the scenes of auditory perception. Curr Opin Neurobiol *20*, 361-366. 10.1016/j.conb.2010.03.009.

Shore, S.E. (1995). Recovery of forward-masked responses in ventral cochlear nucleus neurons. Hear Res *82*, 31-43. 10.1016/0378-5955(94)00160-r.

Southwell, R., Baumann, A., Gal, C., Barascud, N., Friston, K., and Chait, M. (2017). Is predictability salient? A study of attentional capture by auditory patterns. Philos Trans R Soc Lond B Biol Sci *372*. 10.1098/rstb.2016.0105.

Thomas, J.M., Morse, C., Kishline, L., O'Brien-Lambert, A., Simonton, A., Miller, K.E., and Covey, E. (2012). Stimulus-specific adaptation in specialized neurons in the inferior colliculus of the big brown bat, Eptesicus fuscus. Hear Res *291*, 34-40. 10.1016/j.heares.2012.06.004.

Torrado Pacheco, A., Tilden, E.I., Grutzner, S.M., Lane, B.J., Wu, Y., Hengen, K.B., Gjorgjieva, J., and Turrigiano, G.G. (2019). Rapid and active stabilization of visual cortical firing rates across light-dark transitions. Proc Natl Acad Sci U S A *116*, 18068-18077. 10.1073/pnas.1906595116.

Ulanovsky, N., Las, L., Farkas, D., and Nelken, I. (2004). Multiple time scales of adaptation in auditory cortex neurons. J Neurosci *24*, 10440-10453. 10.1523/JNEUROSCI.1905-04.2004.

Ulanovsky, N., Las, L., and Nelken, I. (2003). Processing of low-probability sounds by cortical neurons. Nat Neurosci *6*, 391-398. 10.1038/nn1032.

Uran, C., Peter, A., Lazar, A., Barnes, W., Klon-Lipok, J., Shapcott, K.A., Roese, R., Fries, P., Singer, W., and Vinck, M. (2022). Predictive coding of natural images by V1 firing rates and rhythmic synchronization. Neuron *110*, 1240-1257 e1248. 10.1016/j.neuron.2022.01.002.

van der Heijden, K., Rauschecker, J.P., Formisano, E., Valente, G., and de Gelder, B. (2018). Active Sound Localization Sharpens Spatial Tuning in Human Primary Auditory Cortex. J Neurosci *38*, 8574-8587. 10.1523/JNEUROSCI.0587-18.2018.

Verhagen, J.V., Wesson, D.W., Netoff, T.I., White, J.A., and Wachowiak, M. (2007). Sniffing controls an adaptive filter of sensory input to the olfactory bulb. Nat Neurosci *10*, 631-639. 10.1038/nn1892.

Wang, C.A., Boehnke, S.E., Itti, L., and Munoz, D.P. (2014). Transient pupil response is modulated by contrast-based saliency. J Neurosci *34*, 408-417. 10.1523/JNEUROSCI.3550-13.2014.

Wang, X. (2018). Cortical Coding of Auditory Features. Annu Rev Neurosci *41*, 527-552. 10.1146/annurev-neuro-072116-031302.

Wark, B., Lundstrom, B.N., and Fairhall, A. (2007). Sensory adaptation. Curr Opin Neurobiol *17*, 423-429. 10.1016/j.conb.2007.07.001.

Williams, N.P., and Olson, C.R. (2022). Contribution of individual features to repetition suppression in macaque inferotemporal cortex. J Neurophysiol *128*, 378-394. 10.1152/jn.00475.2021.

Wilson, D.A. (1998). Habituation of odor responses in the rat anterior piriform cortex. J Neurophysiol *79*, 1425-1440. 10.1152/jn.1998.79.3.1425.

Woods, T.M., Lopez, S.E., Long, J.H., Rahman, J.E., and Recanzone, G.H. (2006). Effects of stimulus azimuth and intensity on the single-neuron activity in the auditory cortex of the alert macaque monkey. J Neurophysiol *96*, 3323-3337. 10.1152/jn.00392.2006.

Xiong, C., Liu, X., Kong, L., and Yan, J. (2020). Thalamic gating contributes to forward suppression in the auditory cortex. PLoS One *15*, e0236760. 10.1371/journal.pone.0236760.

Yan, Y., Zhaoping, L., and Li, W. (2018). Bottom-up saliency and top-down learning in the primary visual cortex of monkeys. Proc Natl Acad Sci U S A *115*, 10499-10504.

10.1073/pnas.1803854115.

Yaron, A., Hershenhoren, I., and Nelken, I. (2012). Sensitivity to complex statistical regularities in rat auditory cortex. Neuron *76*, 603-615. 10.1016/j.neuron.2012.08.025.

Zhou, Y., and Wang, X. (2012). Level dependence of spatial processing in the primate auditory cortex. J Neurophysiol *108*, 810-826. 10.1152/jn.00500.2011.

Zhou, Y., and Wang, X. (2014). Spatially extended forward suppression in primate auditory cortex. Eur J Neurosci *39*, 919-933. 10.1111/ejn.12460.

ACKNOWLEDGMENTS

This research was supported by National Institute of Health grants DC003180 and DC005808 to X.W. We thank A. Pistorio, and J. Estes for assistance with animal care, X. Song, X. Liu, and H. Mandal for their comments on the earlier versions of the manuscript.

AUTHOR CONTRIBUTIONS

Conceptualization, S.X., C.C. and X.W.; investigation, S.X., Y.W., and C.C.; visualization, C.C.; funding acquisition, X.W.; project administration, X.W.; supervision, X.W.; data analysis, S.X. and C.C.; modeling, C.C.; writing – original draft, C.C.; writing – review & editing, C.C. and X.W.

DECLARATION OF INTERESTS

The authors declare no completing interests.

Figure S1. Population statistics

(A) Logarithmic histogram of the number of facilitated trials in each facilitation phase. We

chose the sessions that exhibited facilitation phases lasting at least five consecutive trials

as the threshold. Fifty-one neurons passed the threshold.

(B) Histogram of the proportion of tested trials in each session (fifty-one neurons).

(C) Histogram of the proportion of trials that was required to achieve the first facilitation

phase lasting at least five consecutive trials (fifty-one neurons).

Figure S2. Facilitation neurons across cortical areas and layers

(A-D) Best frequency (BF) distribution on the cortical surface from five hemispheres of four monkeys. The BF of recorded neurons (color dots) was examined based on their responses to pure tones stimuli ranging from 2kHz (blue) to 32kHz (red). Fifty-one neurons that exhibited facilitation phases lasting at least five consecutive trials were labeled with asterisks. The black dashed line marked the lateral sulcus line approximated by the bone suture in the temporal lobe. The auditory cortex was divided (gray dashed line) into the rostral primary auditory cortex (A1), caudal A1, and caudal area (caudal lateral and caudal medial) along the rostral-caudal axis based on the BF distribution change. (A) Tonotopy from left hemisphere of Monkey 1. Six facilitated neurons were located in caudal area. (B) Ten facilitated neurons (two overlapped) located in A1 of left hemisphere (top) and twenty-two facilitated neurons (two overlapped) located in A1 and caudal area of right hemisphere (bottom) of Monkey 2. (C) Thirteen facilitated neurons (two overlapped) from right hemisphere of Monkey 3. (D) Fourteen neurons were recorded using the intracellular recording method in the left hemisphere of Monkey 4.

(E) Depth information for all the 104 neurons. Left, first spike depth (x-axis) is the absolute depth where the first spike is detected, unit depth (y-axis) is the absolute depth where the neuron is recorded currently. Right, recording depth equals to the relative depth which is calculated as the unit depth minus first spike depth. Eighty-nine neurons came from supragranular layers (recording depth less than 1mm, orange bars).

(F) Same to (E) but for fifty-one neurons that exhibited facilitation phases lasting at least five consecutive trials. Forty-three neurons came from supragranular layers (orange bars).

Figure S3. Facilitation in putative excitatory and inhibitory neurons

(A) Example putative inhibitory narrow-waveform (NW, left) and putative excitatory

broad-waveform (BW, right) neurons.

(B) Among the fifty-one neurons with facilitation phase lasting at least five consecutive

trials, the waveform of thirty-two neurons has been recorded. Twenty-nine neurons have a

signal-to-noise ratio that was larger than 20dB.

(C) Among the twenty-nine neurons, we used 0.3ms trough to peak duration as the

boundary for classifying putative inhibitory (blue bars) and excitatory (orange bars)

neurons. 21% of neurons were classified as putative inhibitory neurons.

(D) Histogram of the proportion of facilitation trials between putative inhibitory (blue bars)

and excitatory (orange bars) neurons.

1
2
Figure S4. Facilitation occurred in the back, in the darkness, in using
unfrozen wideband noises
(A) Spatial receptive field of example unit M43S-374 obtained under equal-probability
presentation mode. Black arrow indicates the spatial location of continuously presented
stimuli. Six orange circles indicate front speakers, and nine black dots indicate back
speakers.
(B) Firing rate versus speak number of same example neuron under the equal-probability
presentation mode.
(C) Left, spike raster of same example neuron tested at speaker #9 under continuous
sound presentation mode. Gray shaded area indicates the sound presentation period.
Right, trial-by-trial firing rate. Dashed blue line indicates the facilitation threshold. Red
squares include trials belonging to the long facilitation phase (i.e., at least five consecutive
trials with firing rates exceeding the facilitation threshold).
(D-F) Similar to **(A-C)**, but the example unit M43S-34 was tested under the light-off
condition.
(G) Histogram of the proportion of facilitation trials under the continuous presentation
mode for light on (blue bars) and off (orange bars) sessions. Among the seven light-off
sessions, five sessions were tested when light was turned off, and two sessions were
tested when both eyes were blocked by an acoustic drape.
(H) Unfrozen wideband noise stimuli were tested for example unit M43S-374. Four
sessions tested with unfrozen wideband noise stimuli were not included in 129 sessions
mentioned above.

1
2
Figure S5. Facilitation occurred using different ISIs

(A) Example unit M43S-374 showed neural facilitation at fixed length 700ms ISI (top left),
2200ms ISI (top right), 4200ms ISI (bottom left), and random length 700-2200ms ISI
(bottom right). Gray shaded area indicates the sound presentation period.

(B) Across the 129 sessions with facilitation phase lasting at least five consecutive trials,
ISIs were classified into three groups: short (500 and 700ms, 78 sessions), long (1200,
2200, 3200, 4200 and 5200ms, 36 sessions) and random (700-1000 or 700-1200 or
700-1700 or 700-2200 or 1200-2200ms, 15 sessions).

Figure S6. Neural facilitation did not alter SRF

(A) Spatial receptive field under the equal-probability presentation mode before (up) and

after (down) the presentation of continuous sound stimuli (same neuron in Figure 1C).

White arrow indicates the speaker location tested under continuous presentation mode.

Area within the enclosed black line is proportional to the reciprocal of tuning selectivity,

and the size of white dot is proportional to the direction selectivity.

(B) Similar to (A) but for all other three example neurons.

(C) Tuning selectivity (left) and direction selectivity (right) before and after the continuous

presentation mode.

(D) Left, the pairwise correlation coefficient between each pair of responses to fifteen

speaker locations in the example unit M43Q-797. Color dots and lines indicate averaged

firing rate under equal-probability presentation modes. Right, histogram of the correlation

coefficient before and after the continuous presentation mode. Orange bars indicate

paired sessions with a correlation coefficient greater larger than 0.7.

(E) Total firing rates (i.e., without minus the spontaneous firing rate) changes for the

highest (1st, orange circles and line) and lowest (15th, blue circles and line) ranked target

speakers (rank 1st: p = 0.7962, rank 15th: p = 0.9738, rank-sum test). Color dots indicate

the mean value.

(F) Spontaneous (blue dashed boxes) and evoked (orange dashed boxes) total firing

rates of equal-probability presentation mode, non-facilitation, and facilitation phases of the

continuous presentation mode. For equal-probability presentation mode, only trials using

the target speaker were included.

Figure S7. Proportion of facilitation and adaptation trials averaged

across sound levels and speakers

(A) Number of all tested sessions (black bars) and sessions with facilitation phases lasting

at least five consecutive trials (yellow bars) at different speaker rank.

(B) Proportion of facilitation (orange dots and bars) and adaptation (blue dots and bars)

trials averaged across sound levels under the continuous presentation mode. Dots and

error bars indicate mean ± standard deviation of mean.

(C) Proportion of facilitation (orange dots and bars) and adaptation (blue dots and bars)

trials averaged across speakers under the continuous presentation mode.

Figure S8. Explanations of EI-LIF and MP-LIF models.

(A) A conceptual model for location-specific facilitation and adaptation. Orange and blue circles indicate excitatory and inhibitory synaptic inputs, respectively. The width of circles indicates the strength of synaptic inputs. Pentagon and upside-down pentagon shapes indicate 1st and 15th ranked speakers, respectively. Top, under equal-probability presentation mode, EI-LIF model neuron has a lower response at the ipsilateral location due to stronger inhibition (thick blue line) than excitation (thin orange line). Left, under continuous presentation mode, overall stronger inhibition than excitation would evoke neural adaptation at speaker ranked 1st (blue line is thicker than the orange line). Right, under continuous presentation mode, overall stronger excitation than inhibition would evoke neural facilitation at speaker ranked 15th (orange line is thicker than the blue line).

(B) A conceptual model for location-specific sustained depolarization and hyperpolarization. Left, under equal-probability presentation mode, MP-LIF model neuron has panoramic subthreshold activity across all spatial locations but activity at the ipsilateral location is weakest (width of circle). Right, the neural activity of the network is homeostatically regulated: some neurons exhibit sustained depolarization (red circles) while others exhibit sustained hyperpolarization (blue circles). Black lines indicate the connections between two neurons. Bottom, under continuous presentation mode, sustained depolarization of weaker responses (red line) at the 15th-ranked speaker will be accompanied by sustained hyperpolarization of stronger responses (blue line) at the 1st-ranked speaker.

(C) A computational model for location-specific facilitation. Gray shaded area (trial #1 to #20) indicates the equal-probability presentation mode. Under this presentation mode, no

adaptation occurred for both excitatory and inhibitory synaptic inputs, thus the synaptic

release probability is 100% for both inputs. Under the continuous presentation mode (trial

#21 to #100), the inhibitory synapses (blue line) have a stronger adaptation amplitude (i.e.,

lower synaptic release probability) than the excitatory synapses (orange line).

(D) A computational model for location-specific sustained depolarization. Under the

equal-probability presentation mode, the membrane potential is stable at -69mV. Under

the continuous presentation mode, membrane potential reached a stable level of -62mV

which was 7mV depolarized after 30 trials. Those three parameters were obtained from

our intra-cellular recordings.

(E) Leaky integrate-and-fire (LIF) neuron model. LIF neuron integrates multiple excitatory

(orange lines) and inhibitory (blue lines) synaptic inputs and fires a spike (t1, t2) whenever

the membrane potential passes the threshold. Top, location-specific facilitation

mechanism for the EI-LIF model neuron. Bottom, the other mechanism for the MP-LIF

model neuron.

(F) Neural adaptation simulated by EI-LIF model neuron. Left, spike raster plot of one

session under equal-probability (shaded area) and continuous presentation mode. Blue

dots indicate spikes belong to the long adaptation phase (i.e., at least five consecutive

trials with firing rates lower than the adaptation threshold). Right, trial-by-trial firing rate

from the same session. Blue dots and line indicate trails belong to the long adaptation

phase. Thick orange and blue lines indicate the facilitation and adaptation thresholds,

respectively.

(G) Similar to (F) but for MP-LIF model neuron.

Name	Symbol	Source	Value	Range
Leaky integrate-and-fire (LIF, equation 1)				
Time step	Δt	This model	0.0001s	fixed
Capacitance	C	Wehr and Zador 2003	0.25*10⁻⁹F	fixed
Excitatory reversal potential	E_e	Wehr and Zador 2003	0V	fixed
Inhibitory reversal potential	E_i	Wehr and Zador 2003	-0.085V	fixed
Leak conductance	g_{rest}	Wehr and Zador 2003	25*10⁻⁹S	fixed
Resting potential	E_{rest}	Experimental data	-0.069V	fixed (EI-LIF)
Spontaneous noise scale	σ_s	This model	0.01V	fixed
Gaussian noise	ω_n	This model	[-1: 1]	random
Threshold above resting potential	V_{th}	Wehr and Zador 2003	0.02V	fixed
Excitatory and inhibitory conductance (equations 2 and 3)				
Excitatory synaptic release probability	P_{rel_e}	This model	1	fixed (MP-LIF)
Inhibitory to excitatory ratio	r	This model	1	fixed
Excitatory synaptic input number	N_e	Wehr and Zador 2003	10	fixed
Alpha function time constant	τ	Wehr and Zador 2003	0.005S	fixed
Conductance noise scale	σ_c	This model	2.5*10⁻⁸S	fixed
Inhibitory synaptic release probability	P_{rel_i}	This model	1	fixed (MP-LIF)
Inhibitory synaptic input number	N_i	Wehr and Zador 2003	10	fixed
Inhibitory to excitatory delay	d	This model	0S	fixed
Excitatory and inhibitory synaptic depression model (EI-LIF) (equations 4 and 5)				
Excitatory synaptic depression amplitude	A_e	This model	0.003	fixed
Excitatory synaptic time constant	τ_e	This model	20S	fixed
Probability of target speaker	P_s	Experimental data	1/0.75/0.5 0.25/0.07	fixed
Inhibitory synaptic depression amplitude	A_i	This model	0.025	fixed
Inhibitory synaptic time constant	τ_i	This model	10S	fixed
Stimulus count	N_{SC}	Experimental data	36	fixed
Trial step	T	This model	1	fixed
Number of trials in random mode	T_r	This model	20	fixed
Number of trials in continuous mode	T_c	This model	300	fixed
Membrane potential depolarization and hyperpolarization model (MP-LIF) (equations 6 and 7)				
Dynamic resting membrane potential	E_{rest_MP}	Experimental data	-0.062V -0.076V	max min
Membrane potential polarization value	M	Experimental data	±0.007V	max, min
Number of trials to stabilize	T_{th}	Experimental data	30	fixed
Spike threshold scale	S	This model	0.3	fixed
Stimulus count	N_{SC}	Experimental data	30	fixed

1 **Table S1. Parameters and corresponding values used in EI-LIF and**
2 **MP-LIF model neurons.**