Dissection of a complex enhancer cluster at the *Sox2* locus

Ran Brosh[1], Camila Coelho[1,7], André M. Ribeiro-dos-Santos[1,7], Megan S. Hogan[1,5], Hannah J. Ashe[1], Gwen Ellis[1], Nicolette Somogyi[1], Raquel Ordoñez[1], Raven D. Luther[1,6], Emily Huang[1], Jef D. Boeke[1,2,3], Matthew T. Maurano[1,4,*]

[1] Institute for Systems Genetics, NYU School of Medicine, New York, NY 10016, USA.

[2] Department of Biochemistry Molecular Pharmacology, NYU School of Medicine, New York, NY 10016, USA.

[3] Department of Biomedical Engineering, NYU Tandon School of Engineering, Brooklyn NY 11201

[4] Department of Pathology, NYU School of Medicine, New York, NY 10016, USA.

[5] Present address: Neochromosome Inc., Long Island City, NY, 11101, USA.

[6] Present address: BlueRock Therapeutics, New York, NY 10016, USA.

[7] These authors contributed equally to this work.

[*] Corresponding author: maurano@nyu.edu

Abstract

Expression of *Sox2* in mouse embryonic stem cells (mESCs) depends on a distal regulatory cluster of DNase I hypersensitive sites (DHSs), but their individual contributions and degree of independence remain a mystery. Here, we comprehensively analyze the regulatory architecture of *Sox2* at its endogenous locus using Big-IN to scarlessly integrate DNA payloads ranging up to 143 kb. We analyzed 74 payloads incorporating deletions, rearrangements, and inversions affecting single or multiple DHSs, as well as surgical alterations to transcription factor (TF) recognition sequences. Multiple mESC clones were derived for each payload, sequence-verified, and profiled for expression of *Sox2* specifically from the engineered allele. We found that a single DHS comprising a handful of key TF recognition sequences was sufficient to sustain significant expression in mESCs. However, its contribution was modulated synergistically by the presence and relative position of additional DHSs. Our synthetic regulatory genomics approach demonstrates that composite regulatory elements can be reduced to a tractable set of essential sequence features, and is readily scalable to investigate regulatory architecture at other key loci genome-wide.

Main Text

Many developmentally important genes lie near clusters of highly cell-type specific DNase I hypersensitive sites (DHSs), typified by the β-globin locus control region (LCR)[1,2]. More recently, 'super-enhancers' have been proposed to exhibit exceptionally cooperative binding, acting as developmental switches that regulate key TFs[3–5]. General exploration of the broad assumptions involved have been impeded by the challenges of large-scale genomic engineering, leaving fundamental questions unanswered. Individual constituent DHSs demonstrate enhancer activity in transient expression assays[3,6], but it is less clear whether they act independently or synergistically in composite elements[7–10]. Furthermore, nearby enhancers may provide redundancy rather than augmenting gene expression per se[11–14]. Therefore, there is a need for new approaches to investigate the architecture of complex regulatory elements.

Sox2 encodes a key TF that regulates mouse embryonic stem cell (mESC) self-renewal and pluripotency[15]. The *Sox2* gene is surrounded by a proximal enhancer cluster, but its expression in mESCs relies on an LCR comprising multiple DHSs located over 100 kb downstream[16,17]. The

effect of the LCR on *Sox2* expression is influenced by distance and/or intervening CTCF sites[18–20]. Surgical inversion of a single CTCF recognition sequence within the *Sox2* LCR affects chromatin architecture but not *Sox2* expression in mESCs[21], suggesting that proximity and expression might be separate functions[22,23]. The *Sox2* locus therefore provides a natural model for dissecting how the composition and configuration of complex regulatory elements determine transcriptional activity. However, previous studies have focused on transient reporter plasmids and simple deletion analyses, which do not address sufficiency or interaction of distinct elements at the endogenous locus.

We have recently developed the Big-IN platform to enable scarless genome rewriting with payloads of more than 100 kb[24]. Big-IN includes two engineering steps: first, CRISPR/Cas9 is used to target and replace an allele of interest with a landing pad (LP) (**Fig. 1a**). Second, the LP is replaced by a transfected DNA payload using recombinase-mediated cassette exchange (RMCE), and payload-harboring cells are isolated using a positive/negative selection strategy. A comprehensive sequencing verification pipeline accompanies each step to ensure on-target single copy integration and lack of unexpected variants (**Fig. 1b**).

Here we use Big-IN to rewrite the mouse *Sox2* locus with designer payloads incorporating deletions, inversions, translocations, and mutations of single and multiple DHSs. We show that the LCR is a complex element whose constituent DHSs are only partially interchangeable and exhibit unexpected dependencies not previously predicted by reporter assays or deletion analyses. We show that the contribution of individual TF activity depends critically on their local context. Future application of this synthetic regulatory genomics approach to assess regulatory architecture promises to cast new light on the function of key loci genome-wide.

Results

Synthetic regulatory genomics of the *Sox2* locus

To enable interrogation of regulatory element function at the murine *Sox2* locus, we integrated Big-IN LPs into the BL6 allele of C57BL6/6J × CAST/EiJ (BL6xCAST) F1 hybrid mESCs. We targeted independent LPs to replace both the 143-kb *Sox2* locus (LP-Sox2) and the 41-kb region surrounding the *Sox2* LCR (LP-LCR) (**Fig. 1a**). We performed comprehensive quality control (QC) using PCR genotyping and sequencing, which identified clones with single-copy, on-target

LP integration (**Supplementary Fig. S1**, and **Table 1**). Using real-time quantitative reverse transcription PCR (qRT-PCR) with allele-specific primers, we confirmed biallelic *Sox2* expression in parental cells, ablation of BL6 *Sox2* expression in LP-Sox2 mESCs, and near total loss of BL6 *Sox2* expression in LP-LCR mESCs, recapitulating previous reports (**Fig. 1c**)[16,17].

We assembled DNA payloads for Big-IN delivery (size = 177 to 142,667 bp) from a BL6 mouse BAC (bacterial artificial chromosome) encompassing the full *Sox2* locus or from chemically synthesized DNA (**Supplementary Table S1**). We verified the resulting payloads through sequencing (**Supplementary Table S2**), delivered them to LP mESCs, isolated multiple mESC clones for each payload, and comprehensively verified single-copy targeted payload integration using targeted capture sequencing (Capture-seq) (**Table 1, Supplementary Table S3**). Approximately 50% of isolated delivery clones passed PCR genotyping, and the majority of those (~87%) passed subsequent Capture-seq screening. All 301 mESC clones characterized in this study were further confirmed by systematic analyses of sequence coverage depth, allelic ratios, and integration site (**Table 1**). The high proportion of on-target payload delivery attests to the fidelity of RMCE and the efficiency of the positive and negative selection for Big-IN delivery.

To dissect the architecture of the *Sox2* locus, we delineated 28 distinct regulatory regions (**Fig. 2a**) through inspection of genomic annotation including DNA accessibility and TF occupancy data in mESC. We delivered payloads including single and multiple deletions of these regions to LP-Sox2 mESCs (**Fig. 1a**). We measured *Sox2* expression from the engineered allele relative to the unedited CAST allele as an internal control (**Supplementary Fig. S2**). Delivery of the wild-type (*WT*) locus fully rescued *Sox2* expression, whereas payloads lacking the LCR (*ΔLCR*) showed minimal expression (**Fig. 2b**, **Supplementary Table S4**). Deletion of DHSs 10-16 downstream of *Sox2* led to 22% reduction in expression, while a larger deletion of the entire 81-kb intervening region between *Sox2* and its LCR (including DHSs 10-18) showed expression comparable to WT. Given that LCR activation of *Sox2* is distance-dependent[19], it is possible that reduced distance partially compensates for loss of proximal regulatory elements. Deletion of the proximal region upstream of *Sox2* (DHSs 1-8) also reduced expression. Combined deletion of both the upstream and downstream proximal DHSs led to even lower *Sox2* expression than either single deletion. Since a more restricted deletion of DHSs 1-8 with just DHSs 10 & 15 showed similar activity, we conclude that the effect of the downstream deletion is largely attributable to

DHSs 10 & 15. In contrast to these results showing a role for the proximal regulatory regions, single deletion of DHS7 or DHS15 or the combined deletion of DHSs 10-11 have been reported to show no expression difference[16], suggesting that some of these DHSs are redundant and show no effect unless deleted together.

Dissection of the *Sox2* LCR

We next focused on a 41-kb region surrounding the *Sox2* LCR, which contains up to 10 DHSs and CTCF sites active in mESCs (**Fig. 3a**). To interrogate the necessity of each DHS, we delivered a series of LCR payloads to LP-LCR mESCs (**Fig. 1a**) and analyzed their ability to restore the expression of the BL6 *Sox2* allele (**Fig. 3b**). Delivery of the WT LCR completely rescued *Sox2* expression. Analysis of single and multiple DHS deletions delineated a core LCR region comprising DHSs 23-26. Within this region, deletion of DHS24 alone critically ablated *Sox2* expression to just 42% of WT, while deletion of DHS23 or DHS26 showed a lesser but significant impact. Within this core LCR, only deletion of CTCF25 was tolerated with little effect. Deletions outside of the core LCR showed little or no effect.

Examination of the occupancy of key TFs involved in pluripotency using ChIP-seq and ChIP-nexus data[3,25–28] showed that each LCR DHS exhibits a different pattern (**Supplementary Fig. S3**), suggesting that the individual DHSs may be functionally specialized. To investigate whether individual DHSs within the LCR function in isolation or rely on their surrounding DHSs, we delivered minimal payloads containing subsets of LCR DHSs to LP-LCR mESCs and measured their activities (**Fig. 4a** and **Supplementary Fig. S3**). We observed that a minimal payload containing the four core LCR DHSs (*23-26*) recapitulated 88% of the activity of the full WT LCR. Activities of single DHSs varied substantially: DHS24 and DHS26 were able to restore 30% and 14% of *Sox2* expression, respectively, whereas DHS23 was completely inactive. But when DHS23 was linked to either DHS24 or DHS26, it augmented their overall activity nearly twofold. Linking 2 copies of DHS24 (*2x24*) also nearly doubled expression relative to a single copy, demonstrating that increased dosage of DHS24 without addition of any new TF activity can manifest in increased expression. Strikingly, linking DHS19 and DHS20 with CTCF25 and DHS26 yielded significant expression (54% of WT) despite lacking both DHS23 and DHS24. This is particularly surprising given that deletion of DHS19 and DHS20 had no effect provided the rest of the LCR was left intact (**Fig. 3b**), DHS19 by itself had no activity (**Fig. 4a**), and both DHS19 and DHS20 were previously reported as inactive in a reporter assay[16]. We speculate that DHS19 and

DHS20 can augment enhancer activity either due to the increased proximity among each other or to CTCF25 and DHS26. We conclude that the DHSs comprising the LCR are not redundant and many exhibit unexpected function when placed in novel contexts.

Context dependence might further manifest as differences in expression for different configurations of the same set of DHSs. Since certain genomic features have an intrinsic polarity, such as transcription or the interaction between cohesin and CTCF, we investigated the effect of DHS orientation on LCR activity by comparing pairs of payloads that differ in the orientation of single or multiple DHSs (**Fig. 4b**). While inversion of the entire LCR resulted in a 24% decrease in its activity, inversions of smaller LCR regions had a lesser effect. Individual inversions of DHSs 23, 24, 25, and 26 had almost no effect on expression. This is consistent with a prior report showing that surgical inversion of a CTCF recognition sequence within CTCF25 had no effect on *Sox2* expression[21]. To explore whether the effect of larger inversions might be mediated by positional differences rather than the orientation of specific TF or CTCF recognition sites, we profiled multiple permutations of the core DHS order (**Fig. 4c**). Each permutation of DHSs 23, 24, and 25 order reduced overall activity, whereas relocating CTCF25 downstream of DHS26 resulted in increased activity. These results suggest that the conformation of DHSs within the LCR plays a role in their function.

Modeling *Sox2* expression

To coherently summarize the architecture of the *Sox2* locus, we investigated linear regression models that predict *Sox2* expression from surrounding DHS composition and configuration (**Fig. 5a**). We established a baseline model considering only key proximal regions along with the four DHSs in the core LCR. Performance improved substantially upon incorporation of interactions among core LCR DHSs, in particular the contribution of DHS23 in conjunction with DHS24 or DHS26, and CTCF25 in the presence of DHS26 (**Fig. 5b**). Performance was higher when restricting to these three top interaction indicators than when considering all pairwise interactions. Given the potential influence of DHS configuration on function, we extended our top interaction regression model to include the orientation of each core LCR DHS and relative order encoded by indicators for DHS24 preceding DHS23 or CTCF25 preceding DHS24. Inclusion of either orientation or order separately improved model performance, but a model including order alone provided a better fit than orientation alone or a combination of both (**Fig. 5b**). The coefficients of the best fit

model showed strong weights for the presence of proximal regulatory regions and core LCR DHSs, as well as key interactions and DHS order (**Fig. 5c**). While most payloads were predicted with little error, we noted that the contribution of DHS26 as well as inversions and permuted constructs demonstrated consistently higher error than other constructs, suggesting additional context effects not fully captured by our simple model (**Supplementary Fig. S4**).

Given the contribution of interaction terms to model performance, we explored in more detail the extent to which activities of core LCR DHSs depend on their surrounding context. We identified pairs of payloads differing solely by the presence of each core LCR DHS. We calculated the contribution of the focus DHS in that context as the difference in *Sox2* expression between mESC clones for each payload pair (**Fig. 5d**). The synergism between DHS23 and DHS24 was the most evident contextual effect, followed by the higher activity shown by near-complete LCR payloads containing individual DHS deletions vs. minimal elements lacking their surrounding native context. Thus, the function of each of the DHSs comprising the *Sox2* LCR depends strongly on their context.

TF-scale dissection of the core LCR

Given the complex interdependencies among core LCR DHSs when considered as units, we used synthetic DNA to investigate function at the level of individual TF recognition sequences. We designed base payloads covering the core LCR DHSs that avoided or shortened a handful of repetitive regions to facilitate chemical synthesis (**Fig. 6a**). We then designed a series of derivative payloads including surgical deletions, mutations or inversions to assess the individual and collective functions of putative TF or CTCF binding sites identified through analysis of TF motif matches and *in vivo* DNA accessibility and TF occupancy data (**Fig. 6b**).

We started with DHS24, which is the strongest single DHS in the LCR and is capable of reproducing 30% of WT LCR activity (**Fig. 4a**). We identified four distinct TF recognition sequence sites, numbered 24.1-24.4 (**Fig. 6b** and **Supplementary Fig. S5**). These sites were predicted to be occupied by key mESC regulators, including the ESRR, nuclear receptor, SOX, and forkhead TF families. Deletion or mutation of all four sites was sufficient to fully ablate activation of *Sox2* by *syn24* (**Fig. 6c**). All single deletions showed at least some effect, ranging from an 83% (24.2) to 28% (24.4) reduction of base activity (**Fig. 6c**). Double or triple deletions showed a further

negative effect, essentially reaching null when 24.2 and 24.3 were both deleted. A minimal payload containing the most essential site based on the deletion analysis (24.2) with its flanking regions was unable to confer any *Sox2* expression. We analyzed the contribution to *Sox2* expression of each TF site relative to its context, comparing pairs of payloads differing by the presence of a focus TF site (**Fig. 6d**). This analysis identified a clear context dependence for all four TF sites, and the contribution of each TF site increased with the number of other sites present.

As DHS23 alone is incapable of activating *Sox2* expression (**Fig. 4a**), its core DHS peak was analyzed linked to DHS24 (*syn23-24*, **Fig. 6b**). The activity of this payload was slightly higher than the comparable non-synthetic payload including DHSs 23-24 (**Fig. 6e**), possibly due to the decreased distance between the core DHS peaks of DHS23 and DHS24 in the synthetic configuration. Deletion of 7 recognition sequence sites within DHS23 reduced activity by only 30%, and a payload containing a deletion extending halfway through DHS23 reduced activity by 53% (**Supplementary Fig. S6a**). These results suggest that we have identified some, but not all, TF sites responsible for the function of DHS23. This synthetic approach further permitted assessment of two conserved DHSs from the human *SOX2* locus, orthologous to mouse DHS23 and DHS24 (**Supplementary Fig. S6b**). While mouse DHS23 and DHS24 combined are sufficient to confer nearly half of WT *Sox2* expression, the two orthologous human DHSs surprisingly showed no activity whatsoever (**Supplementary Fig. S6c**). While the core of the human DHSs showed sequence conservation with mouse sequence, analysis of TF recognition sequences showed a marked divergence relative to the orthologous mouse sites. This suggests that, despite the sequence orthology, the human LCR has diverged functionally.

Finally, we investigated CTCF25 and DHS26, which delineate a region covered by continuous DNA accessibility that comprises two broad DHS peaks positioned closely together (**Fig. 6a-b**). We identified 9 CTCF recognition sequences in CTCF25 and DHS26, 3 of which are in convergent orientation relative to *Sox2* (**Fig. 6b** and **Supplementary Fig. S5**), including the previously reported CTCF site 25.2[21]. While many of these sites lie outside the primary ChIP-seq peak, they reside within a DHS and demonstrate a trace level of CTCF occupancy in WT cells (**Fig. 6b**), so we reasoned that they might be partially redundant with 25.2. We thus investigated whether inversion or deletion of multiple sites might result in a stronger phenotype than the reported inversion of 25.2 alone. Inversion of the 41-bp CTCF footprint at the 3 sites in convergent orientation to *Sox2* in a baseline payload incorporating DHS23 through DHS27 (*23-27 (Divergent CTCF)*,

Fig. 6b) led to a 20% reduction in expression (**Fig. 6f**). Deletion of 8 CTCF recognition sequences (*23-27 (ΔCTCF)*, **Fig. 6b**) led to an even greater reduction of 36% (**Fig. 6f**), bringing activity down nearly to the baseline of the remaining DHS23 and DHS24, and showing that these sites are collectively required for full function of the CTCF25-DHS26 segment. As DHS26 itself can function as an enhancer (**Fig. 4a**), we investigated the function of TF sites not overlapping with CTCF recognition sequences. A synthetic DHS26 payload demonstrated slightly higher activity than the non-synthetic DHS26 (**Fig. 6f**), likely because it extended into CTCF25 to include a predicted HNF4A/RARB site (25.4) and a CTCF site (25.3) (**Fig. 6b** and **Supplementary Fig. S5**). Deletion of 5 TF recognition sequence sites in this context also completely abrogated activity (**Fig. 6d**). Finally, *Δ26.2-27 & 28*, while lacking all 5 of these TF sites and CTCF sites 26.2-26.6, but containing 26.1 (A CTCF site also occupied by Pbx, and Zic3), showed nearly full expression (**Fig. 6b** and **Supplementary Fig. S6d**). This suggests that, like DHS24, CTCF25-DHS26 is a complex regulatory element harboring multiple essential TF sites whose binding is context dependent and synergistic.

Discussion

Dissection of the *Sox2* LCR shows it to be a highly complex and species-specific element whose function depends on the specific conformation of its constituent DHSs. These context dependencies often are not recapitulated in reporter assays or more limited engineering of the endogenous locus, underscoring the importance of a synthetic regulatory genomics approach for comprehensive analysis of regulatory architecture in context. A key example is our investigation of CTCF25, which previously had been shown to exhibit little or no effect on expression when deleted or inverted[21]. In our study, removal or inversion of additional nearby CTCF recognition sequences yielded a significant difference in *Sox2* expression, yet we found inversion of the full DHS to be less important than its position within the LCR. Notably, redundant CTCF binding is a feature of other insulator elements[18,29,30]. Furthermore, while we did not identify all the specific TFs occupying the deleted recognition sequences, the overlapping motif matches suggests that many of these CTCF and TF sites may exhibit occupancy by multiple distinct TFs.

Our results suggest that TF activity at the *Sox2* LCR demonstrates a high degree of cooperativity, but do not address whether cooperativity arises from direct TF-TF interactions, indirect interaction mediated through the chromatin template, local TF concentrations, or other mechanisms[31,32]. As

Sox2 binds indirectly at many of the LCR DHSs (**Supplementary Fig. S3**), it is possible that this context effect is partially mediated through the trans effect of altered Sox2 levels. Deletions of TF recognition sites had larger effects in more complete contexts, which contrasts with prior results showing that the effect of point variants is buffered by stronger DHSs[33]. We speculate that this discrepancy is related to the size of the perturbation, as TF binding can more readily tolerate point changes without falling below the threshold needed for activity[34]. It is possible that the LCR DHSs might act together differently in other contexts. For example, while *Sox2* expression for a given payload was highly consistent across replicate clones, a single clone lacking DHSs 21-22 showed a nearly complete absence of activity in contrast to the other 8 clones which showed near-WT activity (**Fig. 3b**). This outlier clone passed extensive genotyping and Capture-seq to demonstrate that the locus is intact, suggesting that seemingly redundant sites may influence robustness in certain circumstances.

Our approach investigates function at the endogenous locus, preserving its long-distance architecture and surrounding genomic context. The WT constructs replacing the full locus or just the LCR were sufficient to rescue expression in mESC without any detectable defect, but it is possible their chromatin state might still differ from the endogenous locus. The payloads themselves are epigenetically naïve in that they do not pass through development and their chromatin state is established after delivery based on the surrounding context. Characteristic chromatin and expression patterns are readily reestablished on DNA that passes through the germline[35,36], but it is possible that DNA transfected to cycling cells exhibits only partial functionality[37,38]. For example, the failure of DHS23 to activate *Sox2* expression alone might arise from an inability to independently establish a functional accessible state. Consistent with this, DHS23 augmented the activity of both DHS24 and DHS26, suggesting its function is to some extent flexible. We expect future studies might study the establishment of chromatin state on rewritten genomic loci in more detail.

We have described a general approach to dissect locus architecture in-place from 100-kb scale down to individual bp-level TF binding events, and have shown that analyses of sufficiency and combinatorial rearrangements at the endogenous locus reveal unexpected behavior relative to reporter assays or single deletion analyses. We have already employed Big-IN in multiple cell types and loci[24], suggesting that our approach is readily generalizable. While large-scale multi-

plexed CRISPR screens typically do not verify the effect of each edit, our genotyping and genomics validation has shown that a recombinase-based approach avoiding double stranded breaks yields high efficiency delivery with little or no off-target integration. We expect that engineering of large or complex payloads in yeast will become more widely available. In parallel, Golden Gate-based cloning of shorter payloads from commercial synthetic DNA provides a rapid, scalable, and accessible strategy for high-throughput analyses. These developments suggest that synthetic regulatory genomics approaches will rapidly be deployed to investigate function across additional contexts and loci.

References

1. Grosveld, F., van Assendelft, G. B., Greaves, D. R. & Kollias, G. Position-independent, high-level expression of the human beta-globin gene in transgenic mice. *Cell* **51**, 975–985 (1987).

2. Evans, T., Felsenfeld, G. & Reitman, M. Control of globin gene transcription. *Annual review of cell biology* **6**, 95–124 (1990).

3. Whyte, W. A. *et al.* Master transcription factors and mediator establish super-enhancers at key cell identity genes. *Cell* **153**, 307–319 (2013).

4. Pott, S. & Lieb, J. D. What are super-enhancers? *Nature Genetics* **47**, 8–12 (2014).

5. Blobel, G. A., Higgs, D. R., Mitchell, J. A., Notani, D. & Young, R. A. Testing the super-enhancer concept. *Nat Rev Genet* **22**, 749–755 (2021).

6. Thurman, R. E. *et al.* The accessible chromatin landscape of the human genome. *Nature* **489**, 75–82 (2012).

7. Shin, H. Y. *et al.* Hierarchy within the mammary STAT5-driven Wap super-enhancer. *Nature Genetics* (2016).

8. Hay, D. *et al.* Genetic dissection of the α-globin super-enhancer in vivo. *Nature Genetics* **48**, 895–903 (2016).

9. Dukler, N., Gulko, B., Huang, Y.-F. & Siepel, A. Is a super-enhancer greater than the sum of its parts? *Nat Genet* **49**, 2–3 (2016).

10. Moorthy, S. D. *et al.* Enhancers and super-enhancers have an equivalent regulatory role in embryonic stem cells through regulation of single or multiple genes. *Genome Res* **27**, 246–258 (2017).

11. Hong, J.-W., Hendrix, D. A. & Levine, M. S. Shadow enhancers as a source of evolutionary novelty. *Science* **321**, 1314 (2008).

12. Bothma, J. P. *et al.* Enhancer additivity and non-additivity are determined by enhancer strength in the Drosophila embryo. *eLife* **4**, (2015).

13. Osterwalder, M. *et al.* Enhancer redundancy provides phenotypic robustness in mammalian development. *Nature* **554**, 239–243 (2018).

14. Scholes, C., Biette, K. M., Harden, T. T. & DePace, A. H. Signal Integration by Shadow Enhancers and Enhancer Duplications Varies across the Drosophila Embryo. *Cell Rep* **26**, 2407-2418.e5 (2019).

15. Avilion, A. A. *et al.* Multipotent cell lineages in early mouse development depend on SOX2 function. *Genes & Development* **17**, 126–140 (2003).

16. Zhou, H. Y. *et al.* A Sox2 distal enhancer cluster regulates embryonic stem cell differentiation potential. *Genes & Development* **28**, 2699–2711 (2014).

17. Li, Y. *et al.* CRISPR reveals a distal super-enhancer required for Sox2 expression in mouse embryonic stem cells. *PLoS One* **9**, e114485 (2014).

18. Huang, H. *et al.* CTCF mediates dosage- and sequence-context-dependent transcriptional insulation by forming local chromatin domains. *Nat Genet* **53**, 1064–1074 (2021).

19. Zuin, J. *et al.* Nonlinear control of transcription through enhancer-promoter interactions. *Nature* (2022) doi:10.1038/s41586-022-04570-y.

20. Chakraborty, S. *et al.* High affinity enhancer-promoter interactions can bypass CTCF/cohesin-mediated insulation and contribute to phenotypic robustness. 2021.12.30.474562 Preprint at https://doi.org/10.1101/2021.12.30.474562 (2022).

21. de Wit, E. *et al.* CTCF Binding Polarity Determines Chromatin Looping. *Molecular Cell* **60**, 676–684 (2015).

22. Alexander, J. M. *et al.* Live-cell imaging reveals enhancer-dependent Sox2 transcription in the absence of enhancer proximity. *Elife* **8**, (2019).

23. Taylor, T. *et al.* Transcriptional regulation and chromatin architecture maintenance are decoupled modular functions at the Sox2 locus. 2022.02.09.479674 Preprint at https://doi.org/10.1101/2022.02.09.479674 (2022).

24. Brosh, R. *et al.* A versatile platform for locus-scale genome rewriting and verification. *Proc Natl Acad Sci U S A* **118**, e2023952118 (2021).

25. Kagey, M. H. *et al.* Mediator and cohesin connect gene expression and chromatin architecture. *Nature* **467**, 430–435 (2010).

26. Whyte, W. A. *et al.* Enhancer decommissioning by LSD1 during embryonic stem cell differentiation. *Nature* **482**, 221–225 (2012).

27. Beagan, J. A. *et al.* YY1 and CTCF orchestrate a 3D chromatin looping switch during early neural lineage commitment. *Genome Res* **27**, 1139–1152 (2017).

28. Avsec, Ž. *et al.* Base-resolution models of transcription-factor binding reveal soft motif syntax. *Nat Genet* **53**, 354–366 (2021).

29. Bell, A. C. & Felsenfeld, G. Methylation of a CTCF-dependent boundary controls imprinted expression of the Igf2 gene. *Nature* **405**, 482–485 (2000).

30. Dickson, J. *et al.* VEZF1 elements mediate protection from DNA methylation. *PLoS Genetics* **6**, e1000804 (2010).

31. Arnosti, D. N. & Kulkarni, M. M. Transcriptional enhancers: Intelligent enhanceosomes or flexible billboards? *Journal of cellular biochemistry* **94**, 890–898 (2005).

32. Mirny, L. A. Nucleosome-mediated cooperativity between transcription factors. *Proc Natl Acad Sci U S A* **107**, 22534–22539 (2010).

33. Maurano, M. T., Wang, H., Kutyavin, T. & Stamatoyannopoulos, J. A. Widespread site-dependent buffering of human regulatory polymorphism. *PLoS Genetics* **8**, e1002599 (2012).

34. Singh, G. *et al.* A flexible repertoire of transcription factor binding sites and a diversity threshold determines enhancer activity in embryonic stem cells. *Genome Res* **31**, 564–575 (2021).

35. Peterson, K. R. & Stamatoyannopoulos, G. Role of gene order in developmental control of human gamma- and beta-globin gene expression. *Molecular and Cellular Biology* **13**, 4836–4843 (1993).

36. Wilson, M. D. *et al.* Species-specific transcription in mice carrying human chromosome 21. *Science* **322**, 434–438 (2008).

37. Li, Q. *et al.* Differences of globin transgene expression in stably transfected cell lines and transgenic mice. *Blood* **105**, 3346–3352 (2005).

38. Lienert, F. *et al.* Identification of genetic elements that autonomously determine DNA methylation states. *Nature Genetics* **43**, 1091–1097 (2011).

39. Ribeiro-Dos-Santos, A. M., Hogan, M. S., Luther, R. D., Brosh, R. & Maurano, M. T. Genomic context sensitivity of insulator function. *Genome Res* gr.276449.121 (2022) doi:10.1101/gr.276449.121.

40. Ran, F. A. *et al.* Genome engineering using the CRISPR-Cas9 system. *Nat Protoc* **8**, 2281–2308 (2013).

41. Wild, J., Hradecna, Z. & Szybalski, W. Conditionally amplifiable BACs: switching from single-copy to high-copy vectors and genomic clones. *Genome Res* **12**, 1434–1444 (2002).

42. Zhao, Y. *et al.* Episomal editing of synthetic constructs in yeast using CRISPR. 2022.06.21.496881 Preprint at https://doi.org/10.1101/2022.06.21.496881 (2022).

43. Xie, Z.-X. *et al.* Rapid and Efficient CRISPR/Cas9-Based Mating-Type Switching of Saccharomyces cerevisiae. *G3 (Bethesda)* **8**, 173–183 (2018).

44. Agmon, N. *et al.* Phylogenetic debugging of a complete human biosynthetic pathway transplanted into yeast. *Nucleic Acids Res* **48**, 486–499 (2020).

45. Pinglay, S. *et al.* Synthetic genomic reconstitution reveals principles of mammalian Hox cluster regulation. 2021.07.07.451065 Preprint at https://doi.org/10.1101/2021.07.07.451065 (2021).

46. Gietz, R. D. & Schiestl, R. H. High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method. *Nat Protoc* **2**, 31–34 (2007).

47. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

48. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).

49. Faust, G. G. & Hall, I. M. SAMBLASTER: fast duplicate marking and structural variant read extraction. *Bioinformatics* **30**, 2503–2505 (2014).

50. Neph, S. *et al.* BEDOPS: high-performance genomic feature operations. *Bioinformatics* **28**, 1919–1920 (2012).

51. Bonfield, J. K. *et al.* HTSlib: C library for reading/writing high-throughput sequencing data. *Gigascience* **10**, giab007 (2021).

52. Rausch, T. *et al.* DELLY: structural variant discovery by integrated paired-end and split-read analysis. *Bioinformatics* **28**, i333–i339 (2012).

53. Kent, W. J. *et al.* The human genome browser at UCSC. *Genome Res* **12**, 996–1006 (2002).

54. R Core Team. *R: A Language and Environment for Statistical Computing.* (R Foundation for Statistical Computing, 2018).

55. Halow, J. M. *et al.* Tissue context determines the penetrance of regulatory DNA variation. *Nat Commun* **12**, 2850 (2021).

56. Bonev, B. *et al.* Multiscale 3D Genome Rewiring during Mouse Neural Development. *Cell* **171**, 557-572.e24 (2017).

Online Methods

Cloning landing pad and CRISPR/Cas9 plasmids for genome integration

pLP-PIGA (pLP140) (Addgene #168461) was described previously[24] and harbors a pEF1α-mScarlet-P2A-CreERT2-P2A-PuroR-P2A-hmPIGA-EIF1pA cassette flanked by loxM and loxP sites.

pLP-PIGA2 (pLP300) (Addgene #168462) (**Fig. 1a)** was described previously[24] and harbors a pEF1α-PuroR-P2A-hmPIGA-P2A-mScarlet-EIF1pA cassette flanked by loxM and loxP sites, as well as a pPGK1-ΔTK-SV40pA backbone counter-selectable marker cassette.

To derive LP-PIGA3 (pLP305) (**Fig. 1b**), an intermediate pLP303a was cloned by removing mScarlet-P2A-CreERT2-P2A from pLP-PIGA using NcoI and SalI digest followed by a fill-in reaction with Klenow DNA Polymerase and self-ligation to yield a minimal LP plasmid consisting of pEF1α-PuroR-P2A-hmPIGA-EIF1pA cassette flanked by loxM and loxP sites (**Supplementary Table S5**). Sleeping Beauty inverted terminal repeats (ITRs) were amplified from pMH005[39] using primers oRB_277 + oRB_278 for ITR(L) and oRB_279 + oRB_287 for ITR(R). LP-PIGA3 was cloned using a *Bsa*I Golden Gate reaction in which ITRs were cloned outside the LP region of pLP303a.

To target LPs to specific genomic loci, guide RNAs (gRNAs) (**Supplementary Table S8)** were cloned into pSpCas9(BB)-2A-Puro (pCas9-Puro, Addgene #62988) or pSpCas9(BB)-2A-GFP (pCas9-GFP, Addgene #48138) plasmids using BbsI Golden Gate reactions as described[40]. LP homology arms (HAs) corresponding to the genomic sequence flanking the Cas9 cut sites were amplified from a BAC (**Supplementary Table S6)**. HAs were cloned flanking the LoxM and LoxP sites in pLP-PIGA2 using a BsaI Golden Gate reaction or flanking the ITRs in pLP-PIGA3 using a BsmBI Golden Gate reaction (**Supplementary Table S7**). DNA suitable for transfection was prepped using the ZymoPURE maxiprep kit (Zymo Research D4203) according to the manufacturer's protocol.

Landing pad sequences (excluding HAs and backbone) are provided in **Data S1**.

Payload assembly

All payloads were assembled into pLM1110[24] (Addgene #168460) or pRB051 (a derivative of pLM1110). Both are multifunctional YAC/BAC (yeast and bacterial artificial chromosomes) vectors supporting low-copy DNA propagation, selection (LEU2), and efficient homology-dependent recombination in yeast; low-copy propagation, selection (KanR), and copy number induction in TransforMax EPI300 *E. coli*[41], and transient visualization and selection (eGFP-BSD, enhanced green fluorescent protein - Blasticidin-S deaminase) in mammalian cells[24].

Four different strategies were used to assemble payloads into these YAC/BACs (**Supplementary Table S1, Data S1**), using different primers (**Supplementary Table S5)**, gRNAs (**Supplementary Table S8),** and synthetic DNA fragments (**Supplementary Table S9**):

1) *Yeast CRISPR/Cas9 Assembly*

Desired segments of *Sox2* BAC RP23-144O8 were released by an *in vitro* CRISPR/Cas9 digestion using a pair of synthetic gRNAs and recombinant Cas9 and assembled into BsaI-digested pLM1110 as previously described[24].

2) *Yeast Assembly*

PCR amplicons or synthetic DNA (IDT) tiling the desired payload with >75 bp overlap were assembled into pLM1110. Yeast cells were transformed with 20-50 ng I-SceI-digested pLM1110, 100 ng of each DNA fragment, and 50 ng terminal linker fragments (~400 bp gBlocks, IDT) to enable homologous recombination-dependent assembly followed by selection for LEU+ phenotype.

3) *Yeast CRISPR/Cas9 Editing*

Existing payloads were modified in yeast using CRISPR/Cas9 and synthetic linker fragment(s) (IDT) to mediate up to two deletions, insertions, or inversions simultaneously[42]:

i. A yeast strain carrying a parental payload YAC/BAC was pre-transformed with 100 μg SpCas9 expression vector pNA0519 (carrying a ScHIS3 marker) or pCTC019 (carrying a SpHIS5 marker), and selected for a LEU+/HIS+ phenotype. Yeast were then transformed again with 100 μg single/dual gRNA expression plasmid pNA0304/pNA0308[43] carrying a URA3 marker and with 100 μg synthetic linker fragment(s), and selected for a

LEU+/HIS+/URA+ phenotype. gRNAs (**Supplementary Table S8)** were cloned into the single/dual gRNA yeast expression plasmids pNA304/pNA0308, respectively, using *Hind*III and/or *Not*I Gibson Assembly reactions as described[43]. Yeast Cas9 expression vector pNA0519 pRS413-TEF1p-Cas9 was derived from pNA0306[44] by subcloning the TEF1p-Cas9-CYC1t cassette into a pRS413 (ATCC 87518) in order to replace the LEU2 marker with HIS3[45].

ii. Alternatively, yeast cells were transformed with pYTK-Cas9 plasmids (carrying HIS3 marker) that co-express SpCas9 and a gRNA and with synthetic linker fragment(s), and selected for LEU+/HIS+ phenotype. gRNAs (**Supplementary Table S8)** were cloned into pYTK-Cas9[42] using a BsmBI Golden Gate reaction.

4) Golden Gate Assembly (GGA)

DNA fragments were assembled into pLM1110 using *Bsa*I or into pRB051 using *Esp3*I. pRB051 was cloned by PCR-amplifying the RFP transcriptional unit from pLM1110 using primers oRB_564 + oRB_565, digesting the product with *Esp3*I, and performing a *Bsa*I Golden Gate Assembly into pLM1110. For payload assembly, 100 ng of pLM1110 or pRB051 was mixed with 20 ng of each DNA fragment containing terminal *Bsa*I or *Esp3*I sites, respectively, designed to mediate assembly with neighboring fragments (DNA fragments were sourced either from synthetic fragments (IDT) or from existing payloads by PCR-amplification), and with 0.4 μL *Bsa*I-HF V2 (NEB R3733S) or *Esp3*I (NEB R0734S), 1.5 μL 1 mg/mL BSA, 1 μL T4 Quick Ligase (NEB M2200S) and 1.5 μL T4 Ligase Buffer (NEB) in a total volume of 15 μL. Reactions were cycled 25 times between 37 °C (3 min) and 16 °C (4 min), heat inactivated at 50 °C (5 min) and 80 °C (5 min) before transforming into TransforMax EPI300 cells.

Yeast transformation, payload validation and transfer to *E. coli*

Yeast transformations were performed using the Lithium acetate method[46] with *Saccharomyces cerevisiae* strain BY4741. For screening of correct clones, DNA was isolated from individual yeast colonies by resuspension in 10-40 μl of 20 mM NaOH and boiling for 3 cycles of 95 °C for 3 min and 4°C for 1 min. 2 μl of yeast lysate was used as a template in a 10 μl GoTaq Green reaction (Promega M7123) with 0.25 μM of primers. Clones were screened for the intended payload structure using primers that target newly formed junctions (positive screen), and in some cases using primers that target regions that are present in the parental plasmid but absent in the intended

plasmid (negative screen). YAC/BACs were isolated from candidate yeast clones using the Zymo yeast miniprep I protocol (Zymo Research D2001) and transformed into TransforMax EPI300 *E. Coli* cells by electroporation using the manufacturer's protocol (Lucigen EC300150).

Payload DNA screening and preparation

For initial verification, TransforMax EPI300 *E. coli* colonies were picked into 3-5 mL LB-Kan supplemented with CopyControl Induction Solution (Lucigen CCIS125) and cultured overnight at 30 °C with shaking at 220 RPM. YAC/BACs smaller than 20 kb were isolated using the Zyppy Plasmid Miniprep Kit (Zymo Research D4020).

For larger YAC/BACs, crude DNA extraction was performed as follows: 2-3 mL of induced culture was spun down, resuspended with 300 μL RNase A-supplemented Buffer P1 (QIAGEN 19051), and topped with 300 μL of Buffer P2 (QIAGEN 19052). The suspensions were inverted 10 times, incubated at room temperature for 2 min, topped with 300 μL Buffer P3 (QIAGEN 19053), inverted 10 times and spun down at 12000 RPM for 5 min. Supernatants were transferred to new 2 mL tubes, topped with 900 μL isopropanol, mixed by inverting, and spun down at 12000 RPM for 5 min. Supernatants were discarded, pellets were washed with 500 μl of 70% EtOH, and spun down at 12,000 RPM for 1 min. Supernatants were discarded and the DNA pellets were air-dried and resuspended in 30 μL TE buffer. Tubes were spun down at 12000 RPM for 1 min and clear supernatants transferred to new 1.5 mL tubes.

Payload assembly verification was performed by diagnostic digestion with restriction enzymes, PCR verification of new junctions, and Sanger sequencing, depending on the payload. All YAC/BACs were subsequently sequenced to high coverage depth.

Sequence-verified YAC/BAC clones were grown overnight in 2.5 mL cultures of LB-Kan at 30 °C with shaking, diluted 1:100 in LB-Kan supplemented with CopyControl Induction Solution, and incubated for an additional 8-16 hours at 30 °C with shaking. For transfection, DNA was purified from induced *E. Coli* using the ZymoPURE maxiprep kit (Zymo Research D4203) for YAC/BACs <20 kb or using the Nucleobond XtraBAC kit (Takara 740436) for YAC/BACs >20 kb. YAC/BAC preps were stored at 4 °C.

mESC culture

C57BL6/6J × CAST/EiJ (BL6xCAST) mESCs were cultured as described[24]. Specifically, mESCs were cultured on plates coated with 0.1% gelatin (EMD Millipore ES006-B) in 80/20 medium comprising 80% 2i medium and 20% mESC medium. 2i medium contained a 1:1 mixture of Advanced DMEM/F12 (ThermoFisher 12634010) and Neurobasal-A (ThermoFisher 10888022) supplemented with 1% N2 Supplement (ThermoFisher 17502048), 2% B27 Supplement (ThermoFisher 17504044), 1% GlutaMAX (ThermoFisher 35050061), 1% Pen-Strep (ThermoFisher 15140122), 0.1 mM 2-mercaptoethanol (Sigma M3148), 1,250 U/mL LIF (ESGRO ESG1107l), 3 μM CHIR99021 (R&D Systems 4423), and 1 μM PD0325901 (Sigma PZ0162). mESC medium contained KnockOut DMEM (ThermoFisher 10829018) supplemented with 15% FBS (BenchMark 100- 106), 0.1 mM 2-mercaptoethanol, 1% GlutaMAX, 1% MEM nonessential amino acids (ThermoFisher 11140050), 1% nucleosides (EMD Millipore ES008-D), 1% Pen-Strep, and 1,250 U/mL LIF. Cells were grown at 37 °C in a humidified atmosphere of 5% CO2 and passaged on average twice per week. MK6 C57BL/6J mESCs were provided by the NYU Rodent Genetic Engineering Laboratory.

mESC genome engineering

Landing pad integrations were performed (**Supplementary Table S3**) using the Neon Transfection System as previously described into BL6xCAST Δ*Piga* mESCs, in which the endogenous *Piga* gene was deleted[24]. LP-PIGA A1 mESCs, in which LP-PIGA replaces the BL6 *Sox2* allele were previously described[24].

LP-PIGA2 integration at *Sox2* (replacing a 143-kb genomic region) was performed using 5 μg of pLP-PIGA2 and 2.5 μg of each pCas9-GFP plasmid expressing BL6-specific gRNAs mSox2-5p-1 and mSox2-3p-5, followed by 1 μg/mL puromycin (ThermoFisher A1113803) selection for LP-harboring mESCs and 1 μM ganciclovir (GCV, Sigma PHR1593) selection against LP-PIGA2 backbone's HSV1-ΔTK gene.

LP-PIGA3 integration at the *Sox2* LCR (replacing a 41-kb genomic region) was performed similarly with pCas9-Puro plasmids expressing the non-allele specific gRNA mSox2-DHS18-19 and the BL6-specific gRNA mSox2-3p-5, followed by 1 μg/mL puromycin selection. PCR genotyping and Capture-seq validation were used to identify clone LP-LCR H1 (**Supplementary Fig. S1b** and **Supplementary Fig. S1d**), to which a few LCR payloads were delivered (**Supplementary**

Table S3). DELLY analysis later identified a 7.7 kb deletion (**Supplementary Fig. S1d**) in the CAST allele surrounding the mSox2-DHS18-19 gRNA site in a small number of payload clones derived from LP-LCR H1 cells (not included in this study), indicating a mixed population of cells in the original LP-LCR culture. We therefore sub-cloned LP-LCR H1 to isolate clone LP-LCR C1 and confirmed it did not harbor the CAST allele deletion (**Supplementary Fig. S1 c-d**). LP-LCR C1 was used for all subsequent LCR deliveries.

Payload deliveries were performed as previously described[24] with 1-5 × 10^6 mESCs, 1-10 μg YAC/BAC payload DNA and 2-5 μg pCAG-iCre plasmid (Addgene #89573) per transfection, depending on the payload size (larger payloads required more cells, PL DNA and pCAG-iCre plasmid to obtain sufficient correct mESC clones). Transfected mESCs were selected with 10 μg/mL blasticidin for 2 days starting day 1 post-transfection and with 2 nM proaerolysin for 2 days starting day 6 or 7 post-transfection.

For both LP integration and payload delivery, approximately 9 days post-transfection, individual mESC clones were manually picked into gelatinized 96-well plates prefilled with 100 μL 80/20 media. Two days post-picking, clones were replicated into two gelatinized 96-well plates at 90% and 10% relative densities. Three days later, crude gDNA was extracted from the 90% plate as described[24] and used in PCR genotyping to identify candidate clones, which were then expanded from the 10% density plate for further verification and phenotypic characterization. Genomic DNA was extracted from expanded clones using the DNeasy Blood & Tissue kit (QIAGEN 69506).

mESC genotyping
Genotyping mESC clones was performed either using PCR followed by gel electrophoresis as described[24] or using real-time quantitative PCR (qPCR), which was performed with the KAPA SYBR FAST (Kapa Biosystems KK4610) on a LightCycler 480 Real-Time PCR System (Roche) using either 96-well or 384-well qPCR plates. In most cases loading was performed using an Echo 550 liquid handler (Labcyte): A 384-well qPCR plate was prefilled with 5 μL KAPA SYBR FAST and 4 μL water per well. 100 nL of each 100 μM primer and 0.5 μL of each crude gDNA sample were transferred by the Echo. Genotyping payload clones typically included, in addition to assays designed to detect the newly-formed left and right junctions, assays to detect the loss of the landing pad, the absence of the payload YAC/BAC backbone, and for large (>100 kb) payloads, allele-

specific assays to detect delivered regions of the *Sox2* locus. Genotyping primers are listed in **Supplementary Table S10 and Supplementary Table S11**.

High-throughput sequencing verification

Illumina sequencing libraries were prepared from purified DNA using three methods, as listed in **Supplementary Table S12**:

1) The Illumina dsDNA protocol was previously described in [24]. 1 μg of DNA was sheared in a 96-well microplate using the Covaris LE220 (450 W, 10% Duty Factor, 200 cycles per burst, and 90-s treatment time) to yield fragments between 500 to 900 bp. DNA fragments were end-repaired with T4 DNA polymerase, Klenow DNA polymerase, and T4 polynucleotide kinase (New England Biolabs), and A-tailed using Klenow (3′-5′ exo-; New England Biolabs). Illumina sequencing adapters were then ligated to DNA ends using Quick Ligase (New England Biolabs). The post-ligation product was purified using 18% Sera-Mag Magnetic Beads (Cytiva) in polyethylene glycol. DNA libraries were amplified with KAPA 2× Hi-Fi Hotstart Readymix (Roche) and purified with 18% Sera-Mag Magnetic Beads in polyethylene glycol.

2) One-Pot dsDNA was performed as in the Illumina dsDNA protocol, except that 250 ng of sheared DNA was end-repaired and A-tailed in a single reaction using dATP, dNTPs mix, T4 DNA Polymerase, T4 Polynucleotide Kinase, and Taq Polymerase (New England Biolabs) and incubated in a thermocycler at 12 °C for 10 min, 37 °C for 10 min, and 72 °C for 20 min.

3) The NEBNext Ultra FS II kit (NEB E7805L) was used according to the provided protocol.

Final library concentrations were measured on a Qubit using the dsDNA High Sensitivity Assay Kit (Invitrogen).

Hybridization capture (Capture-seq) for targeted resequencing of engineered regions was performed as previously described[24]. Biotinylated bait was generated using nick translation from BACs covering the *Sox2* locus (RP23-144O8 or RP23-274P9; see **Supplementary Table S6**), landing pad plasmid (LP-PIGA2 or LP-PIGA3), payload plasmid backbone, pCAG-iCre and pSpCas9 plasmids. Bait sets and sequencing statistics are listed in **Supplementary Table S12**.

Sequencing libraries were sequenced in paired-end mode on an Illumina NextSeq 500 operated at the Institute for Systems Genetics. Reads were demultiplexed with Illumina bcl2fastq v2.20 requiring perfect match to the indexing BC sequence. All whole-genome sequencing and Capture-

seq data were processed using a uniform mapping pipeline. Illumina sequencing adapters were trimmed with Trimmomatic v0.39[47]. Reads were aligned using BWA v0.7.17[48] to the appropriate reference genome (GRCm38/mm10 or GRCh38/hg38), including unscaffolded contigs and alternate references, as well as to independent custom references for relevant vectors. PCR duplicates were marked by samblaster v0.1.24[49]. Per-base coverage depth tracks were generated using BEDOPS v2.4.4[50].

Variant calling was performed using a standard pipeline based on bcftools v1.14 [51]:

    bcftools mpileup --excl-flags UNMAP,SECONDARY,DUP --redo-BAQ --adjust-MQ 50 --
    gap-frac 0.05 --max-depth 10000 --max-idepth 200000 -a DP,AD --output-type u |
    bcftools call --keep-alts --ploidy [1|2] --multiallelic-caller -f GQ --output-type u |
    bcftools norm --check-ref w --output-type u |
    bcftools filter -i "INFO/DP>=10 & QUAL>=10 & GQ>=99 & FORMAT/DP>=10" --set-GTs
    . --output-type u |
    bcftools view -i 'GT="alt"' --trim-alt-alleles --output-type z

Bcftools call --ploidy was set to 1 for custom references and 2 for autosomes in mm10.

Large deletions and structural variants were called using DELLY v0.8.7 excluding telomeric and centromeric regions[52]. Variants were required to PASS filters, to have at least 10 paired-end reads and 20% of paired-end reads supporting the variant allele. DELLY results for YAC/BAC and capture sequencing data were inspected manually and used to choose clones for successive analysis.

Data were visualized and explored using the University of California, Santa Cruz Genome Browser[53]. The full processing pipeline is available at https://github.com/mauranolab/mapping.

Payload YAC/BAC sequence verification

To identify potential sample swaps or contamination, mean sequencing coverage was calculated for the genomic regions corresponding to the payload sequence and to the non-payload sequence, defined as the engineered *Sox2* regions that do not overlap the payload. To minimize coverage abnormalities associated with sequence termini, 400 bp were clipped from each end of continuous genomic regions. Mean coverage was normalized by the payload YAC/BAC backbone coverage. We expect a normalized coverage of 1±0.4 for the payload region and <0.1 for the non-

payload region. Human orthologous sequences and regions smaller than 200 bp after clipping were ignored. Coverage analysis results and quality control (QC) calls for each sample are reported in **Supplementary Table S13** and summarized in **Supplementary Table S2**.

To detect payload DNA variants that might have been introduced during assembly or propagation in yeast or bacteria, we analyzed high confidence variant calls relative to the payload custom reference having sequencing depth above 50 and quality score above 100 (DP ≥ 50 & QUAL ≥ 100). Payload variants are reported in **Supplementary Table S14** and QC calls are summarized in **Supplementary Table S2**.

mESC clone sequence verification

To verify correct genomic engineering of mESC clones, mean sequencing coverage was calculated for the genomic regions corresponding to the payload and non-payload (as defined above); and for the payload YAC/BAC backbone, landing pad (LP), LP backbone, and pCAG-iCre custom references. Terminal clipping of 400 bp was performed as described above for all genomic regions. Coverage was normalized by the mean genomic coverage of the engineered region flanks (regions captured by the *Sox2* bait, but unmodified) such that a single copy corresponds to a value of ~0.5. For LP clones we expect normalized coverage of 0.5±0.25 for the non-payload region and <0.1 for LP backbone. For payload clones we expect normalized coverage of 1±0.25 for the payload region, 0.5±0.25 for the non-payload region and <0.1 for the LP, payload YAC/BAC backbone and pCAG-iCre. Human orthologous sequences, and regions smaller than 200 bp after clipping were ignored. Samples where the payload region was ignored were classified as "No call". Coverage analysis results and QC calls for each clone are reported in **Supplementary Table S15** and summarized in **Table 1**.

We further verified the presence of BL6 allele variants, which are lost in regions replaced by landing pads and restored by delivered payloads, by calculating allelic ratios as the mean proportion of reads supporting the reference (BL6) allele (propREF) for the genomic regions described above. Values of 0.5±0.2 and 0±0.2 were expected for payload and non-payload regions, respectively. Regions with 10 or fewer variants were ignored. Samples where the payload region was ignored were classified as "No call". Allelic ratio results and QC calls for each clone are reported in **Supplementary Table S15** and summarized in **Table 1**.

To verify genomic integration sites, we detected sequencing read pairs mapping to two different reference genomes using *bamintersect* as previously described[24] with slight modifications: Same-strand reads mapping within 500 bp were clustered, a minimum width threshold of 125 bp was required for reporting, and junctions with few reads (<1 and 5 reads/10M reads sequenced for LP and payload samples, respectively) were excluded. *Bamintersect* junctions were classified hierarchically based on position (**Supplementary Table S16).** Results and QC calls for each clone are reported in **Supplementary Table S17** and summarized in **Table 1**.

RNA Isolation, cDNA synthesis and mRNA expression analysis by real-time qRT-PCR

RNA was isolated from fresh or frozen cells using the Qiagen RNeasy-mini protocol. Since *Sox2* has no introns, additional steps were taken to ensure that RNA was not contaminated with trace residual genomic DNA. DNase treatment was performed on extracted RNA with the Turbo DNA-free kit (Ambion, Fisher AM1907) using the "rigorous DNase treatment" protocol prescribed by the manufacturer. Following DNase treatment, cDNA was synthesized from 1-2 μg total RNA with the Multiscribe High-Capacity cDNA Reverse Transcription Kit (Fisher 4368814), including a "-RT" no-reverse-transcriptase control for a subset of samples.

Real-time quantitative reverse transcription PCR (qRT-PCR) was performed using KAPA SYBR FAST (Kapa Biosystems KK4610) on a 384-well LightCycler 480 Real-Time PCR System (Roche) and threshold cycle (Ct, also called Cp) values were calculated using Abs Quant/2nd Derivative Max analysis. Primers were designed to detect *Sox2* in an allele-specific manner (**Supplementary Table S18**). An Echo 550 liquid handler was used for loading as described above. Thermal cycling parameters were as follows: 3 min pre-incubation at 95 °C, followed by 40 amplification cycles of 3 sec at 95 °C, 20 sec at 57 °C and 20 sec at 72 °C. For a subset of samples, the -RT controls were verified for lack of amplification and all had Ct>31.

Assays were performed in duplicate and replicate wells on the same plate were averaged after masking wells with no or very low amplification. Raw Ct values are listed in **Data S2**. Ct values for the BL6 *Sox2* allele ranged between 20 (WT) and 30 (ΔSox2), suggesting an accurate quantification range of 2^{10}. ΔCt values for the BL6 and CAST *Sox2* alleles were computed relative to *Gapdh*. Replicate ΔCt measurements of the same clone across different plates were averaged. *Sox2* fold change was defined as the difference between the BL6 and CAST *Sox2* alleles, calculated as $2^{\Delta Ct[CAST-BL6]}$. Fold change was scaled to yield expression values ranging from 0 (ΔSox2)

to 1 (WT) by subtracting the mean fold change calculated for the ΔSox2 samples from all data points, then dividing by the mean fold change of the appropriate WT payload samples (full locus or LCR) (**Supplementary Table S4**).

Modeling

We fitted linear regression models to predict *Sox2* expression based on DHS composition and configuration using R v3.5.2[54]. Models were compared based on Bayesian Information Criterion (BIC) to evaluate the performance of predictor combinations. The composition of proximal (DHSs 1-8 and DHSs 10-15) and core LCR (DHS23, DHS24, CTCF25, and DHS26) regulatory elements was represented by their copy number in the payload. Regulatory element configuration was represented by indicator variables of their relative order (DHS24-DHS23 and CTCF25-DHS24; read as DHS24 before DHS23 and CTCF25 before DHS24) and sequence inversion (inv_DHS23, inv_DHS24, inv_CTCF25, and inv_DHS26). Payloads which did not map clearly onto these features were excluded from the analysis (*Δ26-27 & 28* and *Δ20-23.3*, and the synthetic payloads in **Fig. 6**). Interaction between two elements was represented by indicator variables of both elements' presence (DHS23and24, DHS23and26, and CTCF25andDHS26).

Transcription factor motif analysis

We used motif matches previously derived from scanning the reference genomes using FIMO v4.10.244 with TF motifs as previously described[55].

Data Availability

DNase-seq data were obtained from https://www.encodeproject.org for ES_CJ7 (ENCLB163SYJ, DS13320) and H7 human ESC (ENCLB449ZZZ, DS11909)[6]. CTCF ChIP-seq[27] (GSM2259905), ChIP-seq data[3,25,26] (GSM560343, GSM560345, GSM560350, GSM687280, GSM687282, GSM687285, GSM845236, GSM845238, GSM1082340, GSM1082341, GSM1082342), and ChIP-NEXUS[28] for Zic3 (GSM4087824), Pbx (GSM4087823), Esrrb (GSM4087822), Sox2 (GSM4072777), Oct4 (GSM4072776), Nanog (GSM4072778), and Klf4 (GSM4072779) were obtained from the GEO repository.

Code Availability

The processing pipelines for Capture-seq, ChIP-seq, and DNase-seq data are available on Github at https://github.com/mauranolab/dnase. All code for analyses herein is available upon request.

Acknowledgements

We thank Brendan Camellato, Florrie Zhu, Leslie Mitchell, Sudarshan Pinglay, Weimin Zhang, and Yu Zhao for help with yeast assembly. This work was partially funded by National Institutes of Health (NIH) grants RM1HG009491 (to J.D.B.) and R35GM119703 (to M.T.M.).

Author Contributions

R.B., A.M.R., M.S.H., J.D.B. and M.T.M. designed experiments; R.B., M.S.H., G.E., R.O., and R.D.L. performed experiments; C.C., M.S.H., N.S., R.B., and G.E. assembled DNA payloads; M.S.H., H.J.A., E.H., G.E. and R.D.L. performed sequencing; A.M.R. and M.T.M. performed computational analyses; R.B., A.M.R., and M.T.M wrote the manuscript.

Competing interests

J.D.B. is a Founder and Director of CDI Labs, Inc., a founder of and consultant to Neochromosome, Inc, a founder, SAB member of and consultant to ReOpen Diagnostics, LLC and serves or served on the Scientific Advisory Board of the following: Sangamo, Inc., Modern Meadow, Inc., Rome Therapeutics, Inc., Sample6, Inc., Tessera Therapeutics, Inc. and the Wyss Institute.

Figures

Fig. 1. Engineering the *Sox2* locus.

a. mESC engineering strategy. The BL6 allele of the 143-kb *Sox2* locus (left) or the 41-kb *Sox2* LCR (right) were replaced with LP-PIGA2 or LP-PIGA3, respectively. LP integration was aided by Cas9 and a pair of gRNAs targeting both the replaced allele and the LP plasmid and by short homology arms that facilitate homology-directed repair. LP mESCs were selected with puromycin, while ganciclovir (GCV) selects against pLP-PIGA2's backbone integration. Cre recombinase-mediated cassette exchange (RMCE) enabled replacement of each LP with a series of payloads. Transfected cells were transiently selected with blasticidin, followed by counterselection of LP mESCs cells with proaerolysin.

b. Schematic of DNA assembly, mESC engineering, verification and analysis pipelines.

c. Allele-specific qRT-PCR assay for *Sox2* expression. BL6 and Castaneus (CAST) expression was measured using allele-specific primers in parental BL6xCAST mESCs, in mESCs with deletion of the 143-kb Sox2 locus (ΔSox2) or 41-kb deletion of the LCR (ΔLCR), as well as in BL6 MK6 mESCs.

Fig. 2. Regulatory architecture of the *Sox2* locus.

a. Schematic of the *Sox2* locus. Shown are DNase-seq and CTCF ChIP-seq data in mESCs. DHSs, CTCF binding sites, and transcription start sites (TSSs) were numbered from 1 to 28. Orientation of CTCF recognition sequences is indicated where applicable.

b. *Sox2* expression analysis for payloads delivered to the *Sox2* locus in LP-Sox2 mESCs. Blue rectangles demarcate genomic regions included in each payload. Each point represents the expression of the engineered *Sox2* allele in an independent mESC clone. Bars indicate median. Expression was scaled between 0 (*ΔSox2*) to 1 (*WT LCR*).

Fig. 3. Mapping essential DHSs within the *Sox2* LCR.
a. Schematic of the *Sox2* LCR showing DNase-seq and CTCF ChIP-seq data in mESCs. Regions previously tested in luciferase reporter assays[16] and sites of Hi-C contact with the *Sox2* promoter[56] are indicated.
b. *Sox2* expression analysis for payloads with deletions of single and multiple DHSs within the core LCR delivered to LP-LCR mESCs. Blue rectangles demarcate genomic regions included in each payload. Each point represents the expression of the engineered *Sox2* allele in an independent mESC clone. Bars indicate median. Expression was scaled between 0 (*ΔSox2*) to 1 (*WT LCR*).

Fig. 4. Context-dependent DHS function within the core LCR.
Sox2 expression analysis for selected LCR DHSs. Each point represents the expression of the engineered *Sox2* allele in an independent mESC clone. Bars indicate median. Expression was scaled between 0 (*ΔSox2*) to 1 (*WT LCR*). Payload structures are shown in **Supplementary Fig. S3**. **a**. Sufficiency of minimal payloads. **b-c**. Effect of DHS orientation (**b**) and order (**c**). Expression of corresponding baseline payloads is repeated in blue for each payload. Arrows indicate extent of inversion(s) in **b**.

Fig. 5*. *Modeling the regulatory architecture of the *Sox2* locus.

a-c. Analysis of linear regression models to predict *Sox2* expression.

a. Models include features indicating the presence of selected DHSs. DHSs proximal to the gene were grouped and encoded as two features (DHSs 1-8 and DHSs 10-15).

b. Model performance measured by Bayesian Information Criteria (BIC) and presented as Fit (difference from maximum BIC). Feature sets included in each model are indicated by blue dots. *Interaction* includes all interaction terms for core LCR DHSs or only top interactions (DHS23and24, DHS23and26, and CTCF25andDHS26). *Orientation* includes features indicating inversion of each core LCR DHS. *Order* includes features indicating that DHS24 is positioned *Sox2*-proximal to DHS23 (DHS24-DHS23), and DHS25 *Sox2*-proximal to DHS24 (CTCF25-DHS24). Red indicates the model with best fit.

c. Coefficients for the best model.

d. Context-sensitive activity of DHSs in the core LCR. ΔExpression was computed as the difference in expression between payload pairs differing solely by the presence of each focus DHS. The presence (blue dots) or absence (gray dots) of each DHS is indicated; two copies of DHS24 are shown in red. Points indicate mean and bars indicate SD across all pairwise combinations of clones.

Fig. 6. TF-scale dissection of DHS function.

a. Schematic of *Sox2* LCR DHSs 23-26. Shown are DNase-seq and CTCF ChIP-seq data in mESCs. Selected payloads are illustrated. Gray regions include poly(C) and poly(T) sequences shortened in synthetic (syn) versions of DHSs 23 or 24.

b. Enlargement of engineered regions within DHS23 to DHS24 (left) and CTCF25 to DHS26 (right). DNase-seq windowed density and per-nucleotide cleavages, and ChIP-nexus data for selected TFs in mESCs are shown. Payload schemes indicate surgical deletions of predicted TF recognition sites. Magenta triangles denote CTCF sites and their native orientation. Additional TF sites outside deletions (23.8) or overlapping CTCF sites are shown in black.

c-f. *Sox2* expression analysis for perturbations of TF recognition sequences within the core LCR. Payloads include deletions (Δ) of TF or CTCF recognition sequences shown in **b**. Each point represents the expression of the engineered *Sox2* allele in an independent mESC clone relative to baseline payloads (dashed vertical lines). Bars indicate median. Expression was scaled between 0 (*ΔSox2*) to 1 (*WT LCR*), and vertical gray lines indicate median expression of relevant baseline payloads.

c. *syn24.2* contains only a minimal region surrounding DHS24.2 (see **b**). *syn24 (mut24.1-24.4)* contains point substitutions that ablate TF recognition sequences 24.1-24.4 (**Supplementary Fig. S5**).

d. Context-sensitive activity of TF sites 24.1-24.4. ΔExpression was computed as the difference in expression between payload pairs differing solely by the presence of each focus TF recognition sequence. The presence (blue dots) or absence (gray dots) of each TF site is indicated. Points indicate mean and bars indicate SD across all pairwise combinations of clones.

f. *23-27 (ΔCTCF)* has 8 of 9 identified CTCF sites surgically deleted. *23-27 (Divergent CTCF)* has surgical inversion of 3 CTCF sites (TF sites 25.2, 26.1 and 26.5) so that all 9 CTCF sites lie in divergent orientation relative to *Sox2*. *syn26 (Δ5TFs)* has 5 TF recognition sequences surgically deleted. Deletions and inversions are detailed in panel **b**.

Tables

Table 1. Engineered mESC clones quality control summary.
Landing pad (LP) and payload mESC clones were screened first using PCR genotyping and then Capture-seq. Clones included in this study and profiled for *Sox2* expression were further verified by computational analyses of sequencing coverage depth and allelic ratios (**Supplementary Table S15**) and *bamintersect* integration site analysis (**Supplementary Table S17**). Percentages are relative to the number of clones at each stage.

	LP integration – all clones (n=308)		Payload delivery – all clones (n=1357)		
	Passed	**Failed**	**Passed**	**Failed**	**No call**
PCR genotyping	6 (2%)	302 (98%)	676 (50%)	681 (50%)	0
	LP integration – sequenced clones (n=6)		Payload delivery – sequenced clones (n=392)		
Capture-seq	5 (83%)	1 (17%)	339 (86%)	42 (11%)	11 (3%)
	LP clones used for payload delivery (n=3)		Payload clones selected for characterization (n=301)		
Sequencing coverage QC	3 (100%)	0	207 (69%)	1 (0.3%)[a]	93 (31%)[c]
Allelic ratio QC	3 (100%)	0	192 (64%)	0	109 (36%)[c]
bamintersect QC	3 (100%)	0	297 (99%)	4 (1%)[b]	0

[a] gDNA sample with LP backbone coverage attributed to unrelated plasmid contamination.
[b] Only one of two junctions was verified. No off-target junctions were reported.
[c] Small regions, regions with small number of variants, and human orthologous sites were not analyzed (see **Supplementary Table S15** and **Methods** for details).

Supplementary Information for Brosh et al.

TABLE OF CONTENTS

SUPPLEMENTARY FIGURES

Supplementary Fig. S1. Landing pad integrations.

a. Screening of BL6xCAST LP-Sox2 clones using PCR genotyping primers targeting the novel right junction (R Jx) and left junction (L Jx), as well as allele-specific primers to detect the loss of the targeted *Sox2* BL6 allele. Right, selected clones were revalidated using column gDNA. Clone D4 (orange rectangle) was selected for further verification by Capture-seq. Par, BL6xCAST Δ*Piga* mESCs; col, column gDNA; PC, BL6xCAST LP-Sox2 A1 mESCs[24]. **b**. Screening of BL6xCAST LP-LCR clones using PCR genotyping primers targeting the novel right junction (R Jx) and left junction (L Jx), as well as allele-specific primers to detect the loss of the targeted *Sox2* BL6 allele. Clone H1 (orange rectangle) was selected for further verification by Capture-seq. Par, BL6xCAST Δ*Piga* mESCs; col, column gDNA; PC, BL6xCAST LP-Sox2 A1 mESCs (which do not share the L Jx with LP-LCR clones and should produce a shorter amplicon from the R Jx). **c**. Screening of sub-clones of LP-LCR clone H1 using PCR genotyping primers spanning or internal to the CAST allele deletion. GeneRuler 1 kb Plus DNA Ladder (ThermoFisher Scientific) was used in panels **a-c**. **d.** Capture-seq analysis of parental Δ*Piga* mESCs, LP-Sox2 clone D4, and LP-LCR clones H1 and C1. Reads were mapped to the references indicated above. The PuroR gene in pCas9-Puro cross-maps with the LP-PIGA3 and is shaded gray. Ticks under each coverage track indicate heterozygous (blue) or homozygous non-reference (yellow) variants. ITR, inverted terminal repeats; pEF1α, human EEF1A1 promoter; PuroR, puromycin-resistance gene; hmPIGA, human mini PIGA gene; mScr, mScarlet; pPGK1, human PGK1 promoter.

Supplementary Fig. S2. qRT-PCR results.

Sox2 expression from all mESC clones described in this manuscript. Displayed is the expression of the BL6 allele of *Sox2* relative to the CAST allele, calculated as $2^{\Delta Ct[CAST-BL6]}$. Points represent individual clones and bars indicate median.

Brosh et al SOM 4

Supplementary Fig. S3. *Sox2* **LCR structure and payloads.**
Browser shot for the *Sox2* distal LCR. DHSs and payloads included in **Fig. 3** are demarcated along with DNase-seq, ChIP-seq, and ChIP-nexus tracks.

Supplementary Fig. S4. Linear regression model of *Sox2* regulatory architecture.

a. Feature composition matrix. Gray, absence; blue, presence; red, two copies.

b. Model predictions vs. actual data. Points represent the observed activity (*Sox2* BL6 allele relative expression) for individual mESC clones. Density curves indicate the expected outcome distribution for each regression calculated based on a Student t distribution using the estimated mean and standard error of each PL. Asterisks denote payloads for which measured data diverge from *Proximal+core LCR+Interaction+Position* prediction by >5 standard errors (SE).

Supplementary Fig. S5. Schematic of TF-scale engineering.

Schematic of TF site deletions and mutations in DHS24 (**a**) and TF site deletions and CTCF site inversions in CTCF25-DHS26 (**b-c**). Shown in **a** and **b** are DNase-seq and CTCF ChIP-seq data in mESCs. Selected payloads are shown including TF and CTCF sites, weblogos for matching motifs, and the actual reference (black). Mutated sequence in payload *syn24(mut24.1-24.2)* is in red. **c**. Reference sequences for 9 CTCF sites are shown aligned to the CTCF motif. Orientation in the reference is indicated with triangles. CTCF extended binding modes are represented using three models including the core sequence, an additional upstream sequence, and the up-stream sequence at an additional bp spacing (upstreamP1)[33]. Recognition sequences for other TFs that overlap CTCF recognition sequences are underlined and shown in detail at right and reverse orientation is indicated as (-).

Supplementary Fig. S6. Core LCR and human orthologs analysis.
Sox2 expression analysis. Each point represents the expression of the engineered *Sox2* allele in an independent mESC clone relative to baseline payloads (dashed vertical lines). Bars indicate median. Expression was scaled between 0 (*ΔSox2*) to 1 (*WT LCR*), and vertical gray lines indicate median expression of relevant baseline payloads.

a. Activity of payload *Δ20-23.3* (shown in detail in **Fig. 6b**). Expression data is reproduced from **Fig. 3** and presented here on a different scale in comparison to the activities of payloads *Δ23* and *WT*.

b. Human DHSs orthologous to mouse *Sox2* DHS23 and DHS24. The mouse genomic coordinates of all *Sox2* regulatory regions were broken into 50-bp non-overlapping blocks. These blocks were remapped to the human genome (hg38) using UCSC liftOver with default options and adjacent remapped blocks within 200 bp were merged. Blocks were required to map reciprocally back to the original location in mouse using the same strategy. Shown are DNase-seq in H7 human ESC[6], predicted TF recognition sequences, and phylogenetic conservation. Blue bars demarcate cloned human genomic regions.

c. Activities of payloads comprising human *SOX2* regions orthologous to DHS24 and DHS23-DHS24 delivered to LP-LCR.

d. Activity of payload *Δ26.2-7 & 28* (shown in detail in **Fig. 6b**). Expression data is reproduced from **Fig. 3** and presented here on a different scale in comparison to the activities of payloads *Δ23-25* and *Δ27*.

SUPPLEMENTARY TABLES

Supplementary Table S1. Payloads and assembly details.
Strategies and reagents used for payload YAC/BAC assembly, as well as sequencing data. See **Methods** for details of each assembly strategy, **Supplementary Table S5** for cloning primer sequences, **Supplementary Table S8** for gRNA sequences, **Supplementary Table S9** for synthetic fragments sequences and **Supplementary Table S4** for payload labels used in figures. Sequence-verified unintended payload structures observed in bacteria or in mESCs are indicated in the Assembly strategy field and were assigned n/a values in relevant fields. For two payloads with detected variants (see **Supplementary Table S14**), two different YAC/BAC clones were delivered to mESCs.

This Table is provided as a separate file.

Supplementary Table S2. Summary of YAC/BAC QC.
Summary Table for YAC/BAC QC analyses. See **Supplementary Table S13** and **Supplementary Table S14** for details.

	Assembled payload YAC/BACs (n=71)		
	Passed	**Failed**	**No call**
Sequencing coverage QC	52 (73.2%)	0	19 (26.8%)[a]
Sequencing variants QC	71 (100%)	0	0

[a] Small regions were not analyzed (see **Methods** for details).

Supplementary Table S3. mESC clones.
Summary Table for mESC clones described in this manuscript. Genetic Modification details DNA elements integrated and their integration site in square brackets; transiently transfected plasmids (e.g. pSpCas9 or pCAG-Cre) are indicated by null. See **Supplementary Table S4** for payload labels used in figures. "LP-PIGA A1" corresponds to "Bl6/C LP140(Sox2_1g_5g)_A1", "LP-Sox2 D4" to "BL6/C LP300(Sox2_1g_5g)_201007_D4", "LP-LCR H1" to "Bl6/C LP305(Sox2_6g_7g)_20200915_H1", and "LP-LCR C1" to "Bl6/C LP305(Sox2_6g_7g)_20210326_C01".

This Table is provided as a separate file.

Supplementary Table S4. *Sox2* expression values and activity.
Sox2 expression in engineered mESC clones. The columns Sox2 (CAST) and Sox2 (BL6) list qRT-PCR ΔCt values relative to *Gapdh* for each *Sox2* allele. Fold change indicates the BL6/CAST $2^{\Delta Ct}$ ratio. Activity indicates scaled fold change as detailed in **Methods**. Payload ID, a unique name for each payload; Label, name as in figure panels.

This Table is provided as a separate file.

Supplementary Table S5. Cloning primers.

Primers used for cloning.

Primer name	Primer sequence (5' to 3')
oRB_277	ACGTGGTCTCACCCTCCCTCGAGACGTTATACGCACCGTCTCAAACGTACAGTTGAAGTCGGAAGTTTAC
oRB_278	ACGTGGTCTCACGTTACTTAGTGTATGTAAACTTCTGACC
oRB_279	ACGTGGTCTCGTATGATTGAGTGTATGTAAACTTCTGAC
oRB_287	ACGTGGTCTCAGGATGGATTGAGACGGTGCGTATAACGTCTCGCATATACAGTTGAAGTCGGAAGTTTAC
oRB_413	CTCATTTACATTTCCAATGCTATACCAAAAG
oRB_414	AGGACAGCCAGGGCTACACAG
oRB_419	TGCAGCCCCCTCCTCATCTG
oRB_420	GTTCTAAGCCAGTAGAGAACTATATATCAAGGTCC
oRB_465	TTGAGTACTGGGATGACAAGCAAG
oRB_466	GGGAAGGACCTTGATATATAGTTCTCTACTGGCTTAGAACTGCAGCCCCCTCCTCATCTG
oRB_467	CCCAGCTTTCCGAGCCAGATGAGGAGGGGGCTGCAGTTCTAAGCCAGTAGAGAACTATATATCAAG
oRB_468	GAAGATGGAGCAAGAGGATTACCG
oRB_468	GAAGATGGAGCAAGAGGATTACCG
oRB_470	GGGAAGGACCTTGATATATAGTTCTCTACTGGCTTAGAACGAAGATGGAGCAAGAGGATTACCG
oRB_472	TTGGATATGGTAGTGCTTGCTTGTCATCCCAGTACTCAACTAAATACCTAATAAAAAATGAAAAAAAAATGTACG
oRB_473	GATTTTGAGTTTAAACGTACATTTTTTTTTCATTTTTTATTAGGTATTTAGTTGAGTACTGGGATGACAAGCAAG
oRB_474	TATGGGGGACTTTTGGTATAGCATTGGAAATGTAAATGAGTGCAGCCCCCTCCTCATCTG
oRB_475	CCAGCTTTCCGAGCCAGATGAGGAGGGGGCTGCACTCATTTACATTTCCAATGCTATACC
oRB_476	CTAGCAAAGTTTTGTCAAAGGGCAAGCTAGCCACTGAGAGTTGTCTGAGAAGAAGATGGAGCAAGAGGATTACCG
oRB_477	CTACCTCGAACTCACGGTAATCCTCTTGCTCCATCTTCTTCTCAGACAACTCTCAGTGGC
oRB_478	AACAATCTTGGATATGGTAGTGCTTGCTTGTCATCCCAGTACTCAATCTTTTTTCTATCTTTTAAGGAAGCATCC
oRB_479	TGCGCCACTACGCCCTGCCCTAGGATGCTTCCTTAAAAGATAGAAAAAAGATTGAGTACTGGGATGACAAGCAAG
oRB_483	CTAAATACCTAATAAAAAATGAAAAAAAAATGTACGTTTAAACTC
oRB_484	TCTTTTTTCTATCTTTTAAGGAAGCATCCTAG
oRB_486	TCAGGGTCTCACAGTTCTCATTTACATTTCCAATGCTATACCAAAAG
oRB_487	TCAGGGTCTCTACAGATATAAAATGTCTAAGAGATGGACATGGTGG
oRB_488	TCAGGGTCTCACTGTTCTCATTTACATTTCCAATGCTATACCAAAAG
oRB_489	TCAGGGTCTCTAAAAATATAAAATGTCTAAGAGATGGACATGGTGG
oRB_492	TATGGGGGACTTTTGGTATAGCATTGGAAATGTAAATGAGAAGACTTTAAAATGCATACACATATGTATTCAAGG
oRB_493	TTCAAAATGTGCACCACCATGTCCATCTCTTAGACATTTTATAGTTCTAAGCCAGTAGAGAACTATATATCAAGG
oRB_494	GACGGGGGAAGGACCTTGATATATAGTTCTCTACTGGCTTAGAACTATAAAATGTCTAAGAGATGGACATGGTGG
oRB_495	AAAAAAAATCCTTGAATACATATGTGTATGCATTTTAAAGTCTTCTCATTTACATTTCCAATGCTATACCAAAAG
oRB_496	ACCCAGCTTTCCGAGCCAGATGAGGAGGGGGCTGCAGAAGATGGAGCAAGAGGATTACCG
oRB_497	GCCTACCTCGAACTCACGGTAATCCTCTTGCTCCATCTTCTGCAGCCCCCTCCTCATCTG
oRB_498	CAAAGGGCAAGCTAGCCACTGAGAGTTGTCTGAGAATTGAGTACTGGGATGACAAGCAAG
oRB_499	TGGATATGGTAGTGCTTGCTTGTCATCCCAGTACTCAATTCTCAGACAACTCTCAGTGGC
oRB_500	GCCCTGCCCTAGGATGCTTCCTTAAAAGATAGAAAAAAGATGCAGCCCCCTCCTCATCTG
oRB_501	TTCCGAGCCAGATGAGGAGGGGGCTGCATCTTTTTTCTATCTTTTAAGGAAGCATCCTAG
oRB_502	TTCTCAGACAACTCTCAGTGGCTAG
oRB_504	TGGGATTAAAAGCATGTGCCACC
oRB_505	TTCAATTCCCAGCAACCATTTGACAGGTCACAACTTTCCAGAAGAAATGCAGCAACCAGC
oRB_506	CTTCTGGCTATGGGTGCTGGTTGCTGCATTTCTTCTGGAAAGTTGTGACCTGTCAAATGG
oRB_507	ATGGTAGTGCTTGCTTGTCATCCCAGTACTCAAGCATGCCATACCATCTCTTTTTTGTTG
oRB_508	AAGTTGGTTCAACAAAAAGAGATGGTATGGCATGCTTGAGTACTGGGATGACAAGCAAG
oRB_560	TGTCAAAGGGCAAGCTAGCCACTGAGAGTTGTCTGAGAACTAAATACCTAATAAAAAATGAAAAAAAAATGTACG
oRB_561	ATTTTGAGTTTAAACGTACATTTTTTTTTCATTTTTTATTAGGTATTTAGTTCTCAGACAACTCTCAGTGGCTAG
oRB_562	AAGTTTTGTCAAAGGGCAAGCTAGCCACTGAGAGTTGTCTGAGAATATAAAATGTCTAAGAGATGGACATGGTGG
oRB_563	GCTGGGATTCAAAATGTGCACCACCATGTCCATCTCTTAGACATTTTATATTCTCAGACAACTCTCAGTGGCTAG
oRB_564	TCAGCAGTAGAGACGGCCTGGCTCTAGTAGCGATCTAC
oRB_565	GCCTAAAATGAGACGCAATACGCAAACCGCCTCTCCCCGCGCG
oRB_568	TCAGCGTCTCAGGAATCTCATTTACATTTCCAATGCTATACCAAAAG
oRB_569	TCAGCGTCTCTAAAAATATAAAATGTCTAAGAGATGGACATGGTGGTG

Supplementary Table S6. Genomic coordinates for engineered loci.
Coordinates for payloads, genomic deletions (marked with a "d") and BACs. For some payloads, coordinates include mismatched regions derived from designed sequence mutation. See **Supplementary Table S4** for payload labels used in figures.

This Table is provided as a separate file.

Supplementary Table S7. Homology arm cloning primers.
Primer pairs used to clone left and right homology arms (HAs) using Golden Gate reactions Primer sequences contain *Bsa*I or *BsmB*I sites in lower case, gRNA binding sites (underlined), and PAMs (bold). Length refers to the total size in bp of each HA.

Locus	HA	Length	Primer	Primer sequence (5' to 3')
Sox2	Left	142	Forward	GTGTggtctcACCCT<u>CAAGTCTGAAGTAGTTCAGG</u>**AGG**GTTTGAGGCCAGGAAGGGAT
			Reverse	GTGGTGggtctcACGTTGAACTACTTCAGACTTGGGC
	Right	183	Forward	GTGGTGggtctcGTATGGTTCGGGGACGGTGTTAATATTCTTC
			Reverse	GGTGggtctcAGGAT<u>GAGCTGCAAAGGCTCCCGTT</u>**AGG**AATGAATGCGGATGCCTTGC
Distal LCR	Left	100	Forward	GGTGcgtctcACCC<u>TGTCTTAACCAACCAAAGGTC</u>**AGG**TTATGCTGAGGGTCATCACACAAG
			Reverse	GGTGcgtctcACGTTCTTTGGTTGGTTAAGACGGGTCAC
	Right	138	Forward	GGTGcgtctcGTATGGTTCGGGGACGGTGTTAATATTCTTC
			Reverse	GGTGcgtctcAGGAT<u>GAGCTGCAAAGGCTCCCGTT</u>**AGG**CCAAGCCTTACGGGAACGCC

Supplementary Table S8. gRNAs.

gRNAs used for LP integrations and payload assembly. Allele-specific gRNAs are identified as having CAST variants overlapping the PAM or gRNA seed sequence.

gRNA Names	gRNA Sequence (5' to 3')	Allele specificity
dDHS23	AATTAGTACTCCCCCACGCA	none
invCTCF4.1	GCGTAGAGCTTTCGTCTTTG	none
invCTCF8	TAAAAGCAAGTCCACCAGCA	none
mSox2-3p-5	GAGCTGCAAAGGCTCCCGTT	BL6 (PAM)
mSox2-5p-1	CAAGTCTGAAGTAGTTCAGG	BL6 (PAM)
mSox2-DHS15-16	CATTGGCAGTGTTGTATAGG	none
mSox2-DHS18-19	GTCTTAACCAACCAAAGGTC	none
mSox2-Prom-1	GACAATGAGCGGAGCGTGGC	BL6 (seed)
mSox2_15g	CTGCACCGTCTTAAGATCTC	BL6 (seed)
mSox2_16g	CAGCTGAAAGCGAGCTGCAA	none
mSox2_17g	GTCTAGCGTTTTACAGTCTT	BL6 (seed)
oBRC0453	GTACACCAGAGATCTTAAGA	BL6 (seed)
oBRC0454	TTGGTTAAGACGGGTCACAC	none
oNKS0012	GTATGCATATGCACCACATG	none
oNKS0015	ACGCCGGAACGGTAGCTCAG	none
oNKS0018	TGACTGGCAACCGAGCCGGG	none
oNKS0022	CCTTTACAGAGGAACCCAGG	none
oNKS0023	TGCCAAGTGAAGAAAAGGCA	BL6 (seed)
oNKS0024	CTTGTGTCTATAATCCCAGA	none
oNKS0025	ATGGGGATGCCACTCACTGG	none
oNKS0026	GATCTCATGGTTTGGCAGAG	none
oNKS0028	CTTCACACTGTATCCCAGGA	BL6 (seed)
oNKS0029	GAGTTGAAAAGATGGCTCAG	BL6 (PAM)
oNKS0030	CATGCCTTTAATCCCAGCAG	none
oNKS0031	CTTTAATCTCAACACTTGGG	BL6 (seed)
oNKS0032	GTTTCTATATGTATATCCTG	none
oNKS0033	CACAAAGAAATGAAAGACTC	none
oNKS0034	CTGCTGATACCTGTTCAGCA	none
oNKS0035	GACCCGTCTTAACCAACCAA	none
oNKS0036	GACAGTTCTTGCTCACACAT	none
oNKS0037	GGAGCACTTGTTGGTCTCGG	none
oNKS0038 or dCTCF8	TAAAAGCAAGTCCACCAGCA	none
oNKS0039 or dCTCF14	AGCTAGGAGCCCCTAGGTAC	BL6 (seed)
oNKS0040 or dCTCF17	GAACGCCACTTACTGCTTAC	none
oNKS0041 or dCTCF25	CACAGAAGAAAAAGCGCGGC	none
oNKS0042 or dCTCF1	CTTGAGCAGCGCCCCCACAA	BL6 (seed)
oNKS0043	CCCATGTACTGCAGTTCCTG	none
oNKS0044	TCGTCCGATTTGTCATTAGA	none
oNKS0045	CGCCCCCTATTGGACGCAGT	none
oNKS0048	AGTTCCAGCACACGCGCAGA	BL6 (seed)

Supplementary Table S9. Synthetic fragments.
Sequences of synthetic fragments used for payload assembly.

This Table is provided as a separate file.

Supplementary Table S10. LP and PL integration junction genotyping primers.
Primer pairs used to verify newly formed junctions in engineered cells.

This Table is provided as a separate file.

Supplementary Table S11. LP and PL integration additional genotyping primers.
Primer pairs used to verify or characterize engineered cells.

Assay	Forward primer sequence (5' to 3')	Reverse primer sequence (5' to 3')
Sox2[Bl6]	CTGAAGCATGAGCTTCCTCCCT	GCAGTACATGGGTCGTACACATG
CAST allele deletion spanning	TCCAGCGTCTTTCCCAGTTC	AGGAGGGTTCAAGACAGGGT
CAST allele deletion internal	CTGTTGTCCTAAGCAAGAAGGTAGGC	TTGGGACTCTCAAACCAAAGACTGTAAAAG
Sox2[BL6]_107kb_down	GCCTTGTGAGCTCAGGAGAAGTC	TACTCAAGGTCAGACGTGCG
Sox2[BL6]_7kb_down	GGCCAAGCCATACTGGGCTC	GCTCCAATCAAGCCAGTAGATGTAGAG
Sox2[BL6]_45kb_down	TTGCTCCAGGTCAGTCCAGC	ACTGTACACTCTGGTCGACCT
Sox2[BL6]_80kb_down	GACTTGAATTTGACCCATGAACCAAT	GCAGATGAGGACTTTTCGAGAGT
Sox2[BL6]_15kb_down	GGCAAGTGAAAGAGAGGTGAAAAGATG	GTCACATGAAGAGCACACAGACAC
Loss LP-Sox2	GGCCAGGAAGGGATATTTGAGA	GGGACTGTGGGCGATGTG
Loss of LP-LCR (L Jx)	CTGTTGTCCTAAGCAAGAAGGTAGGT	GGGACTGTGGGCGATGTG
Loss of LP-LCR (R Jx)	TGCGTCCTGCAGGCGCGTTC	CATCACATTCCCAGTGGGTCAG
YAC/BAC backbone (OriV)	CTGTCTATGGCCCTGCAAAC	CAGTGAGGGCCAAGTTTTCC
BSD	TCGCGACGATACAAGTCAGG	GGACCTTGTGCAGAACTCGT

Supplementary Table S12. Sequencing libraries.
Summary Table for DNA and DNA capture sequencing libraries. Sample ID, a unique sequencing library ID; Payload ID, a descriptor of the relevant landing pad or payload (see **Supplementary Table S4** for payload labels used in figures); Clone ID, a unique descriptor of mESC clones. For each sample, the number (#) of sequenced, analyzed and duplicate reads is provided. Bait Set is provided for capture samples.

This Table is provided as a separate file.

Supplementary Table S13. YAC/BAC sequencing coverage QC.
Summary Table for YAC/BAC payload DNA sequencing coverage depth analysis. Length (bp) and normalized (norm) coverage were calculated for regions corresponding to the payload and non-payload, as defined in **Methods**. Quality control (QC) calls are included with comments for unexpected values. See **Supplementary Table S4** for payload labels used in figures.

This Table is provided as a separate file.

Supplementary Table S14. YAC/BAC variant calling.

Summary Table for YAC/BAC payload variants relative to custom references. Ref, reference sequence; Alt, variant sequence; Qual, quality score; DP, sequencing depth; Feature, the custom reference feature overlapping the variant; RefSeq, variant and surrounding sequence; Comments, annotation and interpretation of variant location and relevance. None of the detected variants are expected to confound the measured activities of integrated payloads. See **Supplementary Table S4** for payload labels used in figures.

This Table is provided as a separate file.

Supplementary Table S15. mESC clone capture sequencing QC.

Summary of QC analyses of Capture-seq samples derived from engineered mESC clones, including sequencing coverage, allelic ratios, and integration site analysis (*bamintersect*). See **Methods** for more detail. Sample ID, a unique sequencing library identifier; Payload ID, LIMS name for relevant payload or LP; Clone ID, a unique descriptor of the mESC clone. Analyses were performed separately for: payload (genomic regions covered by the delivered payload), non-payload (genomic regions overlapping engineered locus and absent from payload), flanks (regions captured by the *Sox2* bait, but not engineered), PL[BB] (YAC/BAC backbone), LP (landing pad), LP[BB] (landing pad plasmid backbone), and iCre (pCAG-iCre plasmid). norm coverage, normalized mean coverage; propREF, proportion of reference to non-reference allele reads. QC calls for the coverage and variants analyses are included, as well as for the *bamintersect* analysis (see **Supplementary Table S16**). R/L Jx, right/left junction. See **Supplementary Table S4** for payload labels used in figures.

This Table is provided as a separate file.

Supplementary Table S16. *bamintersect* junction categories.

Hierarchical criteria used to classify *bamintersect* junctions. bam1 and bam2 are the two references to which junctional reads were mapped. LP, Landing Pad; PL, payload; BB, backbone; Jx, Junction; Chr, chromosome.

Clone Type	Priority	Jx class	bam1 Chr	bam1 Position	bam1 Strand	bam2 Chr	bam2 Position	bam2 Strand
LP	1	L Jx	chr3	0-1000 bp upstream of HAL's 5'	+	LP	First 1000 bp	-
LP	2	R Jx	chr3	0-1000 bp downstream of HAR's 3'	-	LP	Last 1000 bp	+
LP	3	LP-BB	bam1 or bam2 maps to LP backbone					
LP	4	off-target	None of the above					
PL	1	L Jx[a]	chr3	0-1000 bp upstream of HAL 3'	+	PL	First 1000 bp	-
PL	2	R Jx[a]	chr3	0-1000 bp downstream of HAR's 5'	-	PL	Last 1000 bp	+
PL	3	L ITR-PL[b]	LP	First 1000 bp	+	PL	First 1000 bp	-
PL	4	R ITR-PL[b]	LP	Last 1000 bp	-	PL	Last 1000 bp	+
Any	5	Non-informative[c]	chr3	Engineered region	Any	Any	Any	Any
Any	6	PL-BB[d]	bam1 or bam2 maps to YAC/BAC PL backbone					
Any	7	off-target	None of the above					

[a] For payloads with wild type termini, CAST allele reads are indistinguishable from BL6 allele reads.
[b] Only applicable to payloads delivered to LP-PIGA3 (LP305) as the LP's ITRs remain flanking the integration site after PL delivery (see **Fig. 1**).
[c] Non-informative junctions often originate from the CAST allele within the engineered locus
[d] LP-BB and PL-BB junctions are often a result of contamination of gDNA samples by LP/PL plasmids.

Supplementary Table S17. mESC *bamintersect* results.

Summary of *bamintersect* integration site analysis for mESC clones capture sequencing samples. Each line describes a junction (Jx) detected by *bamintersect* (see **Methods**). Chrom (chromosome), chromStart, chromEnd, and Strand for bam1 and bam2 refer to the mapped reads location for the first and second reference genome, respectively. NearestGene indicates the nearest UCSC gene, including distance in bp (negative value indicates an upstream gene) for mm10 or nearest feature for custom references. ReadsPer10M is the number of read pairs supporting the junction per 10 million sequencing reads. Sample ID, a unique sequencing library identifier; Payload ID, LIMS name for relevant payload or LP; Clone ID, a unique descriptor of the mESC clone; Genomes indicate the reference genomes compared. Junc_type indicates the junction classification based on criteria detailed in **Supplementary Table S16**. See **Supplementary Table S4** for payload labels used in figures.

This Table is provided as a separate file.

Supplementary Table S18. qRT-PCR primers.

Primer pairs used for qRT-PCR.

Assay	Forward primer sequence (5' to 3')	Reverse primer sequence (5' to 3')
Sox2[BL6]	AACCGATGCACCGC	GAGCATTATCAGATTTTTCC
Sox2[Cast]	AGCCGATGCACCGA	TGAGCATTATCAGATTTTTCT
Gapdh	AGAACATCATCCCTGCATCC	CACATTGGGGGTAGGAACAC

SUPPLEMENTARY DATA

Data S1. Landing pad and payload sequences.
Sequences for LPs and payloads described in this manuscript, provided in fasta format. For LPs, sequences include only the integrated region (excluding homology arms). For payloads, sequences include only the payload region of the YAC/BAC (from loxM to loxP) and correspond to the designed assembly (sequence variants are not included). See **Supplementary Table S4** for payload labels used in figures.

Data S2. Raw qRT-PCR Ct values.
Raw qRT-PCR Ct values for all mESC clones and assays. Source, the date when assay was performed; RT, indicates whether cDNA sample was prepared with a reverse transcriptase (TRUE) or without ("-RT" control, FALSE); Ct, threshold cycle. See **Supplementary Table S4** for payload labels used in figures.