TITLE:

Non-random distribution of mitochondrial m.3243A>G heteroplasmy in human retina and its impact on cellular phenotype

AUTHORS:

Nathaniel K. Mullin[1,2,3], Andrew P. Voigt[1,2,3], Miles J. Flamme-Wiese[1,2], Xiuying Liu[1,2], Megan J. Riker[1,2], Katayoun Varzavand[1,2], Edwin M. Stone[1,2], Budd A. Tucker[1,2], Robert F. Mullins[1,2]

[1] University of Iowa Institute for Vision Research, Iowa City, IA, USA

[2] Department of Ophthalmology and Visual Sciences, University of Iowa, IA, USA

[3] Medical Scientist Training Program, University of Iowa, Iowa City, IA, USA

Address correspondence to:

Robert F. Mullins

4130 MERF

375 Newton Rd

Iowa City, IA 52242 USA

Phone: (319) 335-8222

Email: robert-mullins@uiowa.edu

CONFLICT-OF-INTEREST STATEMENT:

The authors have declared that no conflict of interest exists.

Abstract word count: 199

Total word count: 9,370

Figure count: 9

Supplemental Figure count: 3

Tables count: 0

Reference count: 70

ABSTRACT

Variants within the mitochondrial genome (mtDNA) can disrupt organelle function and lead to severe multi-system disease. The wide range of disease manifestations observed in heteroplasmic patients results from varying fractions of abnormal mtDNA molecules in different cells and tissues. Here, we profiled the transcriptome, chromatin accessibility state, and heteroplasmy of the most common pathogenic mitochondrial mutation, m.3243A>G, in single cells isolated from the retina and choroid of a patient with retinal and systemic disease. We found that the proportion of the pathogenic m.3243G allele was neither evenly nor randomly distributed across ocular cell types. All classes of cells in the neural retina exhibited a high percentage of the mutant variant. However, endothelial and lymphocyte populations of the choroid were almost homoplasmic for the wildtype allele. Gene expression analysis revealed that the presence of m.3243G causes upregulation of *MTRNR2L1*, a pseudogene of humanin, known to be protective against oxidative damage. We further found by joint multimodal sequencing of the retinal pigment epithelium that a high proportion of the 3243G variant was associated with transcriptionally and morphologically abnormal cells. Together, these findings show the non-random nature of mitochondrial variant partitioning and demonstrate that m.3243A>G causes non-cell autonomous dysfunction in the eye.

BRIEF SUMMARY

Transcriptome and ATAC sequencing in single cells from a human donor with MELAS show that m.3243A>G variant segregation follows a cell type-specific pattern.

GRAPHICAL ABSTRACT

MELAS Patient Eye

% m.3243A>G

scRNAseq

mt-scATACseq

RPE-initiated retinal dysfunction

Sparing of the choriocapillaris

↑ % m.3243G in RPE

↓ % m.3243G in choriocapillaris

INTRODUCTION

Several single nucleotide variants on the mitochondrial genome are known to cause retinal dystrophy and multi-system disease (1). A notable feature of these variants is that they exhibit a high degree of phenotypic heterogeneity (2). This heterogeneity manifests in terms of timing of disease onset, organ systems involved, and severity of tissue dysfunction (3, 4). The heterogenous nature of genetic mitochondrial disease is likely due, at least in part, to the fact that the mitochondrial genome exists at a high copy number in relation to the nuclear genome (5). Since every cell possesses hundreds to thousands of mitochondrial genomes (mtDNA) (6), individual cells may contain a mixture of mutant and wildtype alleles, a phenomenon known as heteroplasmy (7). Mutant mitochondrial alleles are known to be unevenly distributed among organs (8, 9) and cells within the same organ (10, 11). How the distribution of mutant and wildtype mitochondrial molecules is established during organogenesis and maintained throughout life, as well as influence of this distribution on the resulting disease phenotype, are areas of active interest in the study of mitochondrial disease (3, 5, 12, 13).

Here, we focus on the most common pathogenic mitochondrial variant: m.3243A>G, a transition mutation found in the mitochondrial leucine tRNA gene *MT-TL1* (14). m.3243A>G is thought to disrupt electron transport chain function through mistranslation of the thirteen protein-coding genes found on the mtDNA (15-20). Depending on the age of onset and organ system involvement, patients with this variant are typically diagnosed with one of three syndromic conditions: maternally inherited diabetes and deafness (MIDD, OMIM 520000), mitochondrial encephalopathy, lactic acidosis, and stroke-like episodes (MELAS, OMIM 540000) or Leigh syndrome (OMIM 256000) (20, 21). Symptoms can include vision loss (22-24), sensorineural hearing loss (25), diabetes mellitus, gastrointestinal disturbances, myopathy, stroke-like episodes (20), and premature death (26). Notably, the affected organs in these syndromes arise from all three primordial germ layers. In addition to causing dysfunction in a broad range of organ systems, the phenotypic expression of m.3243A>G is also highly variable, even within affected families (27). The heterogenous pattern of organ involvement between patients with the m.3243G variant has led to speculation that disease may be explained by the random partitioning of the pathogenic allele during early development, such that affected organs are simply those that randomly contain a higher proportion of m.3243G. However, emerging evidence has called some of the assumptions underpinning this model of m.3243A>G pathogenesis into question. Global mutant allele

burden is not sufficient to explain the heterogeneity observed in m.3243A>G associated disease phenotype (28, 29), implying a role for variable intra-tissue heteroplasmy. Recent work in circulating blood suggests that allelic segregation within tissues can be non-random (11). The concept of controlled mitochondrial DNA distribution builds on previous observations that mutant allele proportion followed predictable patterns between tissues (21, 30) and that such patterns may even be heritable, implying the involvement of nuclear genetic factors in governing the segregation or propagation of mutant alleles (3, 29).

In the current study, our goal was to investigate the relationship between the distribution of the mitochondrial m.3243A>G variant across the retina to the clinical retinal phenotype in MELAS. We sought to test whether the m.3243G variant was evenly distributed among the cell types of the human retina and to examine the effect of pathogenic variant dosing in specific ocular cell types on cellular phenotype. To accomplish this, we utilized primary post-mortem human ocular tissue from a patient diagnosed with m.3243A>G associated MELAS to better understand the cell type-specific features that underly disease pathogenesis. From this tissue, we collected neural retina, retinal pigment epithelium (RPE), and choroid (31). The retina, RPE, and choroid are an ideal system for the study of the cell type-specific dynamics that govern heteroplasmy and cellular response to a mitochondrial variant in a complex tissue. While the neural retina and RPE arise from neural ectoderm, the cells of the choroid arise from an independent pool of mesenchymal progenitors (32, 33). The endothelial cells of the choriocapillaris develop largely via hemo-vasculogenesis from choroidal mesoderm progenitors (32-34). The circulating and resident blood cells in the retina and choroid provide an opportunity to study an extraocular embryological lineage (bone marrow). We provide evidence that the ratio m.3243A to m.3243G is not randomly distributed across the cell types of the neural retina and choroid. A high proportion of m.3243G in the RPE and neural retinal cells causes dysregulation of gene expression. Together, combining intra-organ heteroplasmy data with prior knowledge of retinal metabolism sheds new light on m.3243A>G pathogenesis in the eye.

RESULTS

Clinical ophthalmologic history of m.3243A>G disease.

Although m.3243A>G is known to be sufficient to cause retinal disease (14, 35, 36), the sequence of events that connect the presence of the mutation in cells to the widespread retinal atrophy is not fully understood (3, 29). Fundus examination of the proband revealed macular atrophy typical of the m.3243A>G variant (**Figure 1A, B**) (22). Optical coherence tomography (OCT) of the retina showed outer retinal tubulations (ORTs) (**Figure 1C**), a clinical finding known to be associated with m.3243A>G associated disease and with RPE/choroid dysfunction leading to retinal dystrophy more broadly (37-39). The proband had been previously confirmed to carry the m.3243A>G variant by sequencing and digital PCR (**Figure S1A**), and the pedigree structure was compatible with mitochondrial inheritance. This patient was shown previously to harbor a relatively high global burden of the m.3243A>G variant compared to patients with less severe disease manifestations (30). Intra-tissue heteroplasmy distribution has not previously been explored in the eye.

Figure 1. Macular retinopathy caused by the m.3243A>G variant. (A) Color fundus photograph of the right eye of a healthy control patient with 20/20 acuity. **(B)** Color fundus photograph of the proband. Macular atrophy of the RPE and choriocapillaris typical of MELAS is appreciable (white arrowheads). **(C)** Optical coherence tomography (OCT) of the proband's retina demonstrates outer retinal tubulations (white arrowhead) indicative of Müller cell activation following RPE cell death (GCL; ganglion cell layer, INL; inner nuclear layer, ONL; outer nuclear layer, RPE; retinal pigment epithelium/Bruch's complex).

Single cell profiling of MELAS retina and choroid.

To better understand how the proportion of m.3243G varies within tissues and how such variability affects cellular dysfunction, we utilized a microfluidic-based single-cell sequencing approach to profile the

Figure 2. Transcriptome and chromatin accessibility profiling in single cells isolated from a MELAS patient's eye. (A) Experimental schematic of scRNAseq and mt-scATACseq studies. **(B)** Two-dimensional UMAP embedding of neural retinal cells based on scRNAseq data from the proband and control donor. **(C)** Two-dimensional UMAP embedding of RPE, and choroidal cells based on scRNAseq data from the proband and control donor. **(D)** Dot plot indicating the magnitude of expression (shade of red) and proportion of cells (size of dot) expressing known marker genes in the neural retinal clusters annotated in (F). **(E)** Dot plot of curated marker genes in the RPE/choroid clusters annotated in (G). **(F)** Two-dimensional UMAP embedding of neural retinal cells based on mt-scATACseq data from the proband and control donor. **(G)** Two-dimensional UMAP embedding of RPE and choroidal cells based on mt-scATACseq data from the proband and control donor (EC; endothelial cell, BC; bipolar cell, RGC; retinal ganglion cell, SMC; smooth muscle cell, RPE; retinal pigment epithelium).

transcriptome, nuclear chromatin accessibility, and m.3243 heteroplasmy of single cells isolated from the neural retina and underlying RPE and choroid. Cells were isolated from the right eye (OD) of a patient with clinically and genetically diagnosed MELAS. As m.3243A>G is known to cause region-specific retinal atrophy (22), samples were recovered from both the macular (M) and peripheral (P) retina, as well as from that of an age and sex-matched control individual. The neural retina and underlying RPE & choroid were each dissected, dissociated, and profiled with scRNAseq or mt-scATACseq (40) (**Figure 2A**). Cells profiled by single-cell RNA sequencing (scRNAseq) were projected in a low dimensional space based on gene expression (**Figure 2B, C**), and resulting clusters were manually annotated using previously described marker genes for human ocular cell types (41, 42) (**Figure 2D, E**). Due to the sparseness inherent in the single cell assay of transposase-accessible chromatin by sequencing (scATAC-seq) data, we annotated mt-scATAC-seq clusters via unsupervised label transfer from the corresponding scRNA-seq datasets (**Figure 2F, G**) using previously described methods (43). Clusters were examined to ensure proper integration of both control and proband

samples and between cells taken from the macula or periphery (**Figure S1E-H**). No major region-specific changes in gene expression were observed and cells from these regions were computationally pooled for subsequent analyses. Since the scRNA-seq and mt-scATAC-seq datasets were derived from the same samples, a high correlation between cell type proportion was observed (**Figure S1B**) lending confidence to the automated label transfer approach. Confidence scores for predicted cell type classification were generated (**Figure S1C, D**), and only cells with a prediction score greater than 0.6 were used in downstream analysis. Cells not reaching this prediction score are colored in dark grey in **Figure 2F and G**.

Figure 3. Non-random distribution of m.3243G in the retina and choroid measured by mt-scATACseq. (A) Schematic of the neural retina, RPE, and choroid indicating the cell classes captured by scRNAseq, mt-scATACseq, and LCM-dPCR. **(B)** Two-dimensional UMAP embeddings of cells recovered from the proband neural retina. Individual cells are colored based on proportion of m.3243A>G mutant allele (red indicating 100% mutant allele and blue indicating 100% wildtype allele). **(C)** Two-dimensional UMAP embeddings of cells recovered from the proband neural RPE and choroid colored based on proportion of m.3243A>G mutant allele. **(D)** Violin plot demonstrating the proportion of m.3243A>G mutant allele in individual cells isolated from the neural retina sample. Cells are grouped based on cell type cluster identity. **(E)** Violin plot demonstrating the proportion of m.3243A>G mutant allele in individual cells isolated from the RPE/choroid sample. A large fraction of endothelial cells and T cells are homoplasmic for wildtype m.3243A.

Non-random m.3243G partitioning in the retina and choroid.

Since the partitioning of the m.3243A>G variant within tissues would likely affect the resulting clinical phenotype, we asked whether heteroplasmy of the m.3243A>G variant differed among cells isolated from the

neural retina and RPE/choroid (**Figure 3A**). mt-scATACseq data were used, and only cells with sufficient coverage of the m.3243 locus were included in the analysis (see **Methods**). The proportion of pathogenic m.3243G was computed on a per-cell basis and overlayed onto cells projected in a low dimensional space (**Figure 3B, C**). The three major cell types of the neural retina (photoreceptor cells, interneurons, and glia) were near homoplasmic for the pathogenic m.3243G allele (**Figure 3B**), all having median m.3243G proportions of 1. This finding was corroborated by bulk dPCR analysis of the same tissue sample (**Figure S1A**). The RPE cells and fibroblasts of the RPE/choroid sample also had median m.3243G proportions of 1 (**Figure 3C**). However, endothelial and T cell populations were near homoplasmic for wildtype m.3243A, all having median m.3243G proportions beneath 0.05 (**Figure 3C**). The proportion of the pathogenic m.3243G allele within individual cells grouped by cell type is plotted in **Figure 3D** and **E**. The melanocytes and pericytes from the RPE/choroid sample were heterogenous in terms of proportion m.3243G (**Figure 3E**).

Figure 4. m.3243G partitioning in the retina and choroid measured by digital PCR. (A-C) Color photomicrographs of neural retina, RPE, and choroidal regions of fixed ocular sections. Regions of interest are shown before laser capture microdissection (*Pre-LCM*). **(D-F)** Color photomicrographs of fixed ocular sections after isolation of photoreceptors (PRC) **(D)**, RPE **(E)**, and choriocapillaris (CC) **(F)** by LCM (*Post-LCM)* Regions isolated for subsequent digital PCR are indicated with dotted lines. **(G)** Proportion of mutant m.3243G in each LCM-captured region as measured by digital PCR.

To validate the finding that choroidal endothelial cells have a reduced proportion of m.3243G compared to other choroidal and retinal cell types, we performed laser capture microdissection followed by digital PCR (LCM-dPCR) on independent m.3243A>G patient samples. Photoreceptor (PRC), RPE, and choriocapillaris (CC) layers (see **Figure 4A**) were separately captured from the contralateral eye of the proband analyzed by mt-scATAC-seq (MELAS 1*) as well as from an unrelated MELAS patient known to carry the m.3243A>G variant (MELAS 2) (**Figure 4A-F**). The CC showed routinely lower m.3243A>G than the RPE and PRC layers

(**Figure 4G**), consistent with the results of the mt-scATAC-seq approach (**Figure 3B-E**). Due to the highly

complex and dense nature of the choroid, isolation of pure endothelial cells is not possible with LCM-dPCR,

explaining the elevated heteroplasmy compared to the mt-scATACseq approach.

We used transmission electron microscopy (TEM) to evaluate the effect of m.3243A>G heteroplasmy on the

ultrastructural phenotype in retinal cells. We found the mitochondria in MELAS neural retina (**Figure 5A**) and

RPE (**Figure 5C**) to be swollen and hypertrophied compared to those in a control sample (**Figure 5B, D**).

Figure 5. m.3243G partitioning in the retina and choroid affects mitochondria ultrastructure. (A)
Transmission electron micrographs of photoreceptor cell inner segments from MELAS patient samples reveal
abnormal mitochondrial structure compared to control tissue **(B)**. **(C)** Transmission electron micrograph of RPE
cells from MELAS patient samples also show abnormal mitochondrial structure compared to control (**D**).
Mitochondria are indicated with asterisks. Scale bar represents 1μm.

Cytoprotective gene upregulation in the presence of m.3243A>G.

After determining the non-random partitioning of m.3243A>G across the cell types of the neural retina and

RPE/choroid, we next asked how the presence of pathogenic m.3243G impacts gene expression in different

cell types. We performed differential expression analysis of cell types from the neural retina and RPE/choroid obtained from the MELAS proband and a normal control individual. The nuclear encoded gene *MTRNR2L1* was upregulated in nearly all cell types in the MELAS samples compared to controls (**Figure 6A-D, E-F**). This gene was upregulated in cell types found to have low levels of m.3243G, such as endothelial cells (**Figure 6D, white arrowhead**), as well as those cell types found to have high levels m.3243G, such as fibroblasts (**Figure 6D, black arrowhead**). The expression of *MTRNR2L1* in individual cell types across samples is shown (**Figure 6E, F**). *MTRNR2L1* encodes the peptide humanin-like 1, a peptide that is highly homologous to humanin, which is encoded by the mitochondrial gene *MT-RNR2*. Humanin is known to be cytoprotective against oxidative stress and has been shown to rescue RPE cells from oxidative damage *in vitro* (44).

Figure 6. Upregulation of *MTRNR2L1* in MELAS retina and choroid. (A, B) *MTRNR2L1* is upregulated in all classes of neural retinal cells in the MELAS sample versus control. Individual cells (dots) are shaded based on expression, with darker shading indicating higher expression. **(C, D)** *MTRNR2L1* is upregulated in all classes of choroid cells including endothelial cells, which contain a low level of the mutant m.3243 allele (white arrowhead). **(E, F)** Visualization of the distribution of *MTRNR2L1* expression across cell types in MELAS and control samples.

Cell type-specific chromatin accessibility changes in MELAS.

Chromatin remodeling has previously been implicated in the cellular phenotype caused by m.3243A>G (45). We sought to use our mt-scATACseq data to understand how the m.3243G variant alters transcription factor binding and expression in cells isolated from the retina and RPE/choroid. We used chromVAR (46) to identify

Figure 7. Cell type and disease state-specific transcription factor binding motif enrichment. **(A)** Heatmap of scaled chromVAR enrichment scores for top cell type-specific motifs. A sample of no more than 100 cells are shown for each cell type. JASPAR matrix names corresponding to marker motifs are shown. **(B, C)** UMAP of neural retinal cells colored by chromVAR enrichment score for the MA0476.1/FOS motif. Enrichment is shown in the control Müller cell cluster but not MELAS. **(D)** Violin plot of MA0476.1/FOS motif enrichment in the Müller cell cluster. **(E, F)** scRNA-seq drive UMAP of neural retinal cells colored by *FOS* transcript abundance. Downregulation is shown in the MELAS Müller cell cluster compared to control. **(G)** Downregulation of *FOS* expression in MELAS Müller cells.

transcription factor-binding motifs that were statistically enriched in these cells (**Figures 7 and S2**). The most highly enriched, cell type-specific motifs from the neural retina are shown in **Figure 7A**. One motif, MA0476.1, was enriched only in a subset of Müller cells (**Figure 7A, box and motif logo**). This motif is known to be bound by FOS (47), a leucine zipper protein and AP-1 transcription factor complex component associated with cell proliferation and apoptosis. We overlaid the chromVAR enrichment score for the FOS motif onto the neural retina UMAP projection, split by disease status (**Figure 7B, C**) such that cells with higher motif availability are shaded darker green. This visualization showed that the FOS motif was enriched in the control Müller cells compared to those from the MELAS patient. Differential enrichment was further visualized in the Müller cluster in a violin plot (**Figure 7D**).

Figure 8. Multimodal sequencing of MELAS and control RPE/choroid. (A) Schematic of multimodal sequencing approach to measure gene expression, nuclear chromatin accessibility, and heteroplasmy in individual cells isolated from the RPE/choroid. **(B)** Two-dimensional UMAP embeddings of cells from the MELAS proband and two controls based on weighted nearest neighbor (WNN) analysis of scRNAseq and mt-scATACseq data. Diverging embeddings of RPE cluster between MELAS and control samples indicate abnormal gene expression (inset). **(C)** Gene expression of the RPE marker gene *BEST1*. Cells are colored based on magnitude of *BEST1* mRNA expression. **(D)** Two-dimensional UMAP embeddings of cells recovered from the proband samples. Individual cells are colored based on proportion of m.3243A>G mutant allele (red indicating 100% mutant allele and blue indicating 100% wildtype allele). **(E)** Violin plot demonstrating the proportion of m.3243A>G mutant allele in individual cells isolated from the RPE/choroid. RPE cells are homoplasmic for the mutant m.3243G allele.

We next asked whether FOS transcript abundance was also lower in MELAS Müller cells, which would suggest bona fide pathway downregulation in which transcription factor abundance correlated with motif availability. We overlaid FOS expression onto the scRNA-seq driven UMAP of control (**Figure 7E**) and MELAS **(Figure 7F)** neural retina cells and again observed a higher level specifically in control Müller cells. The differential FOS expression is further shown in **Figure 7G**. Notably, while several neural retina cell classes had a high proportion of m.3243G (**Figure 3B, D**) FOS motif enrichment is Müller cell-specific, indicating that m.3243G-driven nuclear chromatin remodeling may be cell type-dependent.

Figure 9. m.3243A>G-induced dysfunction in the RPE. (A) Hematoxylin and eosin staining of the RPE and choroid (CC) of a healthy control donor. RPE monolayer shows typical cuboidal epithelial morphology. (**B**) RPE dysplasia is observed in the proband, including the thinning that is associated with epithelial to mesenchymal transition. **(C)** RPE morphology is also abnormal in an unrelated MELAS patient with the m.3243A>G variant. **(D)** Differential gene expression analysis of multimodal sequencing data between MELAS and control RPE cells shows several differentially expressed genes known to be involved in normal RPE function. Scale bar represents 50μm.

Multimodal sequencing reveals aberrant gene expression profile in MELAS RPE.

RPE dysfunction has been implicated in the pathogenesis of m.3243A>G-associated vision loss (23, 48) and the outer retinal tubulations observed in the proband (**Figure 1C**) are indicative of RPE cell death (49). Since the primary functional cell type of the choroid (endothelial cells) were homoplasmic for wildtype m.3243, we hypothesized that the RPE is the site of the initial tissue injury in the eye. To confirm the high proportion of mutant m.3243G in the RPE and investigating its impact on gene expression, we jointly profiled heteroplasmy, gene expression, and chromatin accessibility in the same single cells from choroidal and purified RPE samples of the MELAS proband and controls (**Figure 8A**). Due to a low RPE cell count in unselected RPE/choroid samples (**Figure 2B, C, F, G**), manually purified RPE samples were included in this experiment to guarantee

sufficent cell number. Cell clustering was performed using WNN-analysis (50), which combines gene expression and chromatin accessibility modalities in a weighted manner to drive and refine cluster identification (**Figure 8B**) beyond that derived from single modalities (**Figure S3A-C**). The relative modality weighting behind each cluster is shown (**Figure S3D**). While MELAS and control samples generally clustered together across most cell types (**Figure S3E**), the RPE cluster fragmented (**Figure 8B, inset**) revealing a unique MELAS RPE sub-cluster. The RPE identity of each sub-cluster was confirmed by expression of the marker gene *BEST1* (**Figure 8C**). Confirming prior mt-scATACseq analysis, the MELAS proband RPE was homoplasmic for mutant m.3243G (**Figure 8D, E**). We next sought to understand how this high proportion of m.3243G in the RPE related to cellular phenotype and overall retinal function.

We performed H&E staining on two unrelated MELAS retinal samples. Both donor sections exhibited signs of RPE dysfunction, specifically an epithelial to mesenchymal transition (EMT) that has been previously recognized in models of MELAS *in vitro* (48). MELAS donors exhibited RPE thinning (**Figure 9A-C**), a recognized signed of RPE EMT (51). Additionally, when differential gene expression between MELAS and control RPE cells were analyzed, several genes known to be RPE-specific (*BEST1*, *RLBP1*) and involved in lysosomal function (*ATP10B*) were found to be dysregulated (**Figure 9D**).

DISCUSSION

In this study, we sought to probe the nature and consequences of m.3243A>G heteroplasmy in the human retina. We profiled heteroplasmy, gene expression, and chromatin accessibility in single cells and found that m.3243G is non-randomly distributed, being high in neural retina and RPE and lower in the choroidal endothelium. We also found that the presence of m.3243G results in elevated expression of the pseudogene of a known protective peptide, *MTRNR2L1*. Upregulation of this gene was found in both cell types with high and low levels of m.3243G, implying a response to global environment of oxidative stress in the retina. We showed that m.3243G dysregulates transcription factor binding motif enrichment in a cell type-specific manner. Finally, by combining previous ophthalmologic clinical observations (24, 52) and multimodal sequencing of the RPE, we showed that RPE dysfunction associated with high proportions of m.3243G is likely an early pathogenic step in the retinal dystrophy of MELAS and related mitochondrial conditions (i.e., MIDD and Leigh Syndrome).

m.3243G is non-randomly distributed between ocular cell types.

Mitochondria harboring pathogenic variants have long been known to be subject to purifying selection during oogenesis and early embryonic development (53). More recently, evidence has emerged that such selection may occur within tissues on a cell type-specific basis (10, 11, 54). Currently, the mechanism that leads to uneven mitochondrial DNA partitioning between cells of the same individual is not clear. One hypothesis is that mitochondrial genomes are randomly partitioned into early embryonic progenitors during the first zygotic divisions. In such a model, high proportions of a mutant mtDNA in the central nervous system would arise from mutant mtDNA genomes being randomly partitioned into a neuroectoderm-fated cell in early embryonic development. However, reproducible patterns of mtDNA distribution in specific tissues of unrelated individuals (30) and specific cell types within tissues of unrelated individuals (11) suggest that there is cell type-specific control over mtDNA partitioning. Our data show that within the environment of the retina and choroid, certain cell types develop lower levels of m.3243G despite a background of globally elevated m.3243G. We find that choroidal T cells possess low levels of m.3243G, building on previous observations of lower m.3243G in circulating T cells (11) and observe a similar phenomenon in the endothelial cells of the choroid.

The m.3243A>G causes macular retinopathy featuring outer retinal tubulations.

Patterns of retinal atrophy in inherited retinal disease can be traced to specific cell types. We and others (55) have previously demonstrated that different retinal and choroidal cell types express unique gene expression profiles, and such profiles can vary regionally even within a single cell type. The m.3243A>G mutation causes a characteristic macular atrophy *(*Figure 1A*)*. It has been hypothesized that the unique macular specificity in this disease is caused by region-specific dysfunctional RPE cells (56). Another notable anatomical feature of retinal dystrophy caused by m.3243A>G is the presence of outer retinal tubulations (ORT) (24, 37, 38), indicating outer retinal dysfunction and Müller cell activation (49). ORTs are known to form in response to RPE or choroidal malfunction, such as in the age related macular degeneration (AMD) (57) and the monogenetic disease choroideremia (58). Clinical examination of the retina of MELAS patients suggests a disease process that begins in the RPE and/or choroid, and eventually leads to outer retinal damage. Our data showing a lack

of the mutant allele in the choroidal endothelium suggest that the RPE is the key player in m.3243A>G-associated retinal disease.

Choroidal m.3243A>G heteroplasmy in the context of development.

The distribution of m.3243A>G we observed among the cell types of the choroid can be understood in the context of choroidal development during embryonic and fetal life. The choroid is home to cell types found in many other organs (e.g., endothelial cells), yet is a unique microenvironment that confers choroid-specific gene expression signatures to such cells (e.g., *CA4* expression in the choriocapillaris (59)). We hypothesize that the lower proportion of m.3243G in choroidal endothelial cells is one such choroid-specific trait.

The endothelial cells of the choriocapillaris form in early development from a pool of hemanogioblasts that also give rise to primitive blood cells (32-34). This process is known to begin in human development as early as gestational week 6, when lumenized vessels can be observed expressing markers of proliferation such as CD34 and VEGFR (34). The differentiation of mesenchymal progenitors that gives rise to the choriocapillaris is distinct from the angiogenic process that later connects the choroidal circulation to that of the rest of the body (34). The unique developmental origin of choroidal endothelium may explain the low proportion of m.3243G observed in this study as compared to a previous study of m.3243A>G in other endothelial cell populations (60).

Retinal m.3243A>G heteroplasmy in the context of metabolism.

The metabolic program of retinal cells varies greatly by cell type (61, 62). Photoreceptor cells have the highest energy needs in the retina, primarily utilizing ATP to drive ion pumps that are essential for light modulated changes in membrane potential (62). However, photoreceptors generate most of their ATP through glycolysis as opposed to the more efficient process of oxidative phosphorylation (63). While photoreceptors contain numerous mitochondria, their primary function may be to generate biosynthetic intermediates via the TCA cycle instead of ATP via the electron transport chain. Because of this, mistranslation of electron transport chain components may photoreceptors less than the RPE.

Lactate generated by photoreceptor glycolysis is exported and utilized by glial and RPE cells as a substrate for oxidative phosphorylation in their mitochondria (61) in a reversed process reminiscent of the neuron/astrocyte metabolic linking elsewhere in the central nervous system. The reliance of the RPE on oxidative phosphorylation is necessary to support photoreceptor cell glycolysis, given their physical arrangement with respect to the choroid. Glucose arrives to the outer retina via choroidal blood flow and must be passed through the RPE layer to reach the photoreceptors. Because of this arrangement, RPE preference for oxidative phosphorylation using lactate and other fuels allows for minimal glucose loss during transport to the photoreceptors (63). Additionally, RPE energy demand is supplemented by oxidation of ingested photoreceptor outer segment fatty acids (64).

Cell type-specific metabolic pathway preferences such as these appear to underlie the observed distribution of m.3243A>G burden across the retina and choroid. The m.3243A>G variant is thought to impair tRNA charging by cognate aminotransferases (15, 65) and as a result, likely disrupts electron transport chain protein translation (18). These changes in turn disrupt the process of oxidative phosphorylation in cells with high levels of m.3243A>G (20). Photoreceptor cells do not largely rely on the electron transport chain for energy production, and as such may be more tolerant of the accumulation of mtDNA carrying the pathogenic allele. However, a high proportion of m.3243G in the RPE, as seen in the current study, likely causes decreased electron transport chain function and leads to derangement of the RPE-photoreceptor cooperative metabolic ecosystem, eventually causing photoreceptor dysfunction and vision loss.

Limitations of the current study

The current study has limitations that necessitate future experimental work. Challenges facing the study of m.3243A>G and other mitochondrial variants include the lack of animal models and the technical infeasibility of mitochondrial genome editing. The lack of animal models of genetic mitochondrial disease particularly impairs understanding of pathogenic variant segregation during complex organ development. We addressed this challenge by studying primary ocular tissue from clinically diagnosed MELAS patients in this study. However, this approach limits the population size available. The current study is also limited by only having single timepoints available. This means that any understanding of the developmental processes underlying

mitochondrial genome segregation must be inferred by genome distribution at a single moment in time. Further work in models amenable to sampling over a time course and larger study populations is necessary to confirm these inferences and validate the observation of non-random mtDNA genome partitioning.

Finally, in the current study we focused only on the most common pathogenic mitochondrial variant. While our results imply a generalizable phenomenon for the establishment of mitochondrial variant segregation in the retina and choroid, further study is needed to clarify whether variants segregate in a generalized or variant-specific manner. Such knowledge may also clarify the underlying mechanism of mtDNA segregation within tissues.

Conclusions and Future Directions

In this study, we used a single-cell approach to measure m.3243G heteroplasmy in single cells of a developmentally complex tissue. We found that m.3243G was lowest in the endothelial cells of the choriocapillaris of two MELAS patients and highest in the neural retina and RPE. We also found that in cells with high m.3243G chromatin accessibility and gene expression is altered. Together, these data support the hypothesis that high m.3243G in the RPE initiates MELAS-related retinal dystrophy.

Observation of non-random m.3243G partitioning in adult samples implies selection against heteroplasmy in certain cell types during development. Further work is needed to identify the mechanism by which endothelial cells come to possess only wildtype m.3243 in a patient with globally high m.3243G. *In vitro* models of human genetic mitochondrial disease have proven useful in understanding mtDNA segregation **(66, 67)**. Particularly, human induced pluripotent stem cells (iPSC) can be produced from patient samples harboring mutant mtDNA genomes **(68)**. When combined with contemporary differentiation protocols, such systems may be useful for modeling mtDNA segregation during development. Ultimately, elucidation of how certain cells can regulate m.3243A>G heteroplasmy and other variants may lead to new therapeutic approaches for MELAS and other mitochondrial diseases.

METHODS

Ocular tissue acquisition

Human donor eyes used in single cell studies from a patient affected with MELAS and an unaffected donor were acquired from the Iowa Lions Eye Bank. A second pair of eyes from a donor affected with MELAS was obtained from the Minnesota Lions Eye Bank. All eyes used in the study were obtained or preserved within 8 hours of death and all experiments were performed in compliance with the Declaration of Helsinki and following full consent of the donor's next of kin. Retinal and choroidal tissues were isolated from 8-mm trephine punch biopsies of the macular and inferotemporal peripheral regions and processed as previously described (41). Briefly, the neural retina was dissected away from the underlying retinal pigment epithelium and choroid prior to tissue dissociation. Neural retina samples were dissociated with 20 units/mL of papain (Worthington Biochemical Corporation) with 0.0005% DNase I (Worthington Biochemical Corporation) on a shaker at 37°C for 75 minutes. RPE/choroid samples were dissociated mechanically using razor blades into one-millimeter pieces then dissociated enzymatically using collagenase on a shaker at 37°C for 60 minutes. Following dissociation, cells were cryopreserved in DMSO-based Recovery Cell Cryopreservation Media (Gibco). After overnight cooling at -80°C, samples were transferred to liquid nitrogen (vapor phase) for long-term storage.

Single-cell gene expression (scRNAseq) library construction

Cryopreserved cells were rapidly thawed at 37°C and resuspended in dPBS-/- (Gibco) with 0.04% non-acetylated bovine serum albumin (New England Biolabs). Cells were filtered through a 70μm filter and diluted to target 8,000 cells per run. Single cells were then partitioned and barcoded with the Chromium Controller instrument (10X Genomics) and Single Cell 3' Reagent (v3.1 chemistry) kit (10X Genomics) according to the manufacturer's specifications with no modification (Rev C). Final libraries were quantified using the Qubit dsDNA HS Assay Kit (Life Technologies) and diluted to 3ng/μL in buffer EB (Qiagen). Library quality and concentration was confirmed using the Bioanalyzer High Sensitivity DNA Assay (Agilent) prior to sequencing.

Mitochondrial single-cell ATAC (mt-scATACseq) library construction

Cells were prepared for mt-scATAC sequencing using the protocol described previously by Lareau et al. (40). Briefly, cells were thawed as above and filtered through a 70μm cell strainer. Cells were then washed twice with dPBS-/- (Gibco). Cells were fixed with 1% formaldehyde (Sigma) in PBS-/- for 10 minutes at room

temperature. Fixation was quenched with the addition of glycine (Research Products International) to a final concentration of 0.125M for 5 minutes at room temperature. Following fixation, cells were washed twice with ice-cold PBS-/-. Following the second wash, cells were resuspended in 100µL ice-cold Lysis Buffer (10 mM Tris-HCl pH 7.4 (Millipore-Sigma), 10 mM NaCl (Ambion/Invitrogen), 3 mM $MgCl_2$ (Ambion/Invitrogen), 0.1% Nonidet-P40 substitute (Research Products International), 1% Fraction V BSA (Research Products International)) and incubated on ice for 2.5 minutes. Following incubation, 1mL of ice-cold Wash Buffer (10 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM $MgCl_2$, 1% BSA) was added, and cells were pelleted at 500 x g for 5 minutes at 4˚C. Cells were resuspended in 10µL diluted Nuclei Buffer (10X Genomics) and counted using Trypan Blue (Gibco) with the Countess automated cell counter (Invitrogen). Based on these counts, cell suspensions were diluted in diluted Nuclei Buffer to a final target concentration of 6,000 cells per µL. Cells were then processed following the 10X Chromium Next GEM Single Cell ATAC Reagent Kit v1.1 User Guide (Rev. D) without modification. Final libraries were quantified using the Qubit dsDNA HS Assay Kit (Life Technologies) and diluted to 3ng/µL. Library quality was checked using the Bioanalyzer instrument (Agilent), wherein periodicity of fragment length was observed, consistent with high quality ATAC-seq library construction.

Multimodal mt-scATAC and scRNA library preparation

Cryopreserved RPE & choroid samples were thawed as described above. Cells were prepared for multimodal profiling using the sample preparation modifications described previously (69). Briefly, cells were thawed and filtered through a 70µm cell strainer. Cells were then washed twice with dPBS-/- (Gibco). Cells were fixed with 1% formaldehyde (Sigma) in PBS-/- for 10 minutes at room temperature. Fixation was quenched with the addition of glycine (Research Products International) to a final concentration of 0.125M for 5 minutes at room temperature. Following fixation, cells were washed twice with ice-cold PBS-/-. Following the second wash, cells were resuspended in 100µL ice-cold Lysis Buffer (10 mM Tris-HCl pH 7.4 (Millipore-Sigma), 10 mM NaCl (Ambion/Invitrogen), 3 mM MgCl2 (Ambion/Invitrogen), 0.1% Nonidet-P40 substitute (Research Products International), 1% Fraction V BSA (Research Products International)) and incubated on ice for 2.5 minutes. Following incubation, 1mL of ice-cold Wash Buffer (10 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM MgCl2, 1% BSA) was added, and cells were pelleted at 500 x g for 5 minutes at 4˚C. Cells were resuspended in 10µL

diluted Nuclei Buffer (10X Genomics) and counted using Trypan Blue (Gibco) with the Countess automated cell counter (Invitrogen). Cells were then processing following the 10X Genomics Chromium Next GEM Single Cell Multiome ATAC + Gene Expression (Rev. E) without modification. Final libraries were quantified using the Qubit dsDNA HS Assay Kit and diluted to 3ng/µL. Library quality was checked using the Bioanalyzer system.

scRNA-seq sequencing, preprocessing, and mapping

scRNA libraries were pooled and sequenced using the NovaSeq 6000 instrument (Illumina) generating 100-bp paired end reads. Sequencing was performed by the Genomics Division of the Iowa Institute of Human Genetics. FASTQ files were generated from base calls with the bcl2fastq software (Illumina), and reads were mapped to the pre-built GRCh38 reference with Cell Ranger v3.0.1 (10X Genomics) using the 'count' function. Only cells passing the default Cell Ranger call were analyzed further. Neural retina and RPE/choroid samples were integrated separately using canonical correlation analysis (CCA) in Seurat v3.1 (43). Only cells with between 500 and 7,000 unique genes (features) were included in the analysis. For RPE/choroid samples, only cells with < 65% of reads mapping to mtDNA-encoded genes and < 40% of reading mapping to ribosomal RNA genes were included. For neural retina samples, only cells with < 40% of reads mapping to mtDNA-encoded genes and < 20% of reading mapping to ribosomal RNA genes were included.

mtscATAC-seq sequencing, preprocessing, and mapping

mt-ATAC libraries were pooled and sequenced using the Illumina NovaSeq 6000 instrument generating 100-bp paired end reads. Sequencing was carried out by the Genomics Division of the Iowa Institute of Human Genetics. FASTQ files were generated from base calls with the bcl2fastq software (Illumina). mtscATAC-seq reads were mapped to a modified GRCh38 reference genome that was generated by hard-masking nuclear regions that share homology with mitochondrial sequences, as described previously (40). Mapping and quantification were carried out by Cell Ranger ATAC (v2.0.0, 10X Genomics) using the 'count' function, while the mt-ATAC portion of the multimodal library reads were mapped using Cell Ranger ARC (v.2.0, 10X Genomics). Only cells passing the default Cell Ranger call were analyzed further. Single-cell heteroplasmy of mitochondrial variants was determined using the mgatk package (v0.6.1).

Neural retina and RPE/choroid samples were integrated separately using Signac v1.4.0 (43). Briefly, integration anchors were identified with reciprocal LSI projection. LSI embeddings were then integrated using the 'IntegrateEmbeddings' function, and a UMAP was generated. Only cells with peak region fragments between 1,000 and 75,000 and average mtDNA depth between 10x and 1,000x were used for integration and in downstream analyses. Manually annotated cluster labels were transferred from the scRNA-seq data to the integrated mt-scATAC-seq data. First, transfer anchors between the integrated scRNA-seq and mt-scATAC-seq objects were identified using the 'FindTransferAnchors' function. Next, predicted cluster labels were generated using the TransferData function. Only cells annotated with a prediction.score > 0.6 were used in subsequent analysis.

Multimodal sequencing, preprocessing, and mapping

FASTQ files were generated from base calls with the bcl2fastq software (Illumina). Sequencing reads were mapped to a modified GRCh38 reference genome that was generated by hard-masking nuclear regions that share homology with mitochondrial sequences, as described previously (40). Mapping and quantification were carried out by Cell Ranger ARC (v2.0.0, 10X Genomics) using the 'count' function.

For gene expression libraries, cells were filtered with the Seurat (v3.1) subset function. Cells with nUMIs less than 100 (to remove cells with poor read quality) or greater than 4000 (to remove cells likely to be doublets) were removed. RPE/choroid cells with greater than 40% of reads originating from mitochondrial genes were also removed. Cells with greater than 40% of reads originating from ribosomal RNA genes were also removed. For each of the gene expression libraries, reads were normalized with the Seurat (v3.1) NormalizeData function, and variable features were found with the FindVariableFeatures command. Integration anchors from the first 25 dimensions of the canonical correlation analysis (CCA) were used to integrate gene expression data. Data scaling, principal component analysis, and clustering of the gene expression data were performed using Seurat (v3.1) (43).

For ATAC libraries, cells were filtered with the Signac (v1.4.0) subset function. Cells with fewer than 1000 peak region fragments or greater than 30000 peak region fragments were removed. Cells with greater than 1000x

mtDNA depth were also removed. Mitochondrial DNA sequencing depth (mtDNA_depth) was calculated using mgatk. Term frequency inverse document frequency (TF-IDF) normalization and partial singular value decomposition were performed on each ATAC library using RunTFIDF() and RunSVD() (Signac) with default settings. Gene activities were computed using GeneActivity().

Integration anchors between ATAC datasets were identified using reciprocal LSI projection using FindIntegrationAnchors() in Signac. Dimensions 2-30 from the CCA were used. LSI embeddings were integrated using IntegrateEmbeddings based on these anchors and using dimensions 1-30. A UMAP was generated using the integrated embeddings. Cell identity within the gene expression portion of the experiment was inferred using the TransferData() function in Seurat from the annotated RPE/choroid scRNAseq data obtained in this study (Figure 2C, E). Transfer anchors between were identified using FindTransferAnchors with the following parameters: reduction = 'pca', dims = 1:30. Cells were annotated with a prediction score > 0.8.

Gene expression and ATAC portions of the experiment were combined using Weighted Nearest Neighbor (WNN) analysis (Seurat). Integrated ATAC and gene expression objects were subset to include only common cells (i.e., cells that passed both ATAC and gene expression quality control thresholds). UMAP and cluster identification were performed on the resulting object. Cluster identification used the SLM algorithm with a resolution of 1 (Seurat). Following multimodal integration, cluster identities (i.e., cell type annotation) were refined manually based on canonical marker gene expression.

Sequencing data availability

All raw and processed data are deposited in the Gene Expression Omnibus under the accession number GSE202747. In addition, fully processed single-cell scRNAseq, mt-scATACseq, and multimodal sequencing data are hosted online on Spectacle (70) at https://singlecell-eye.org for interactive visualization.

Single cell heteroplasmy analysis

Per-cell m.3243A>G heteroplasmy was determined as described by Walker et al. (11). mt-scATACseq data

were subset to retain only cells with coverage of the m.3243 locus greater than 10x and less than 1.5 times the upper boundary of the interquartile range. Multimodal sequencing data were subset to retain only cells with coverage of the m.3243 locus greater than 5x and less than 1.5 times the upper boundary of the interquartile range. Only the proband (i.e., not control) patient was included in m.3243A>G heteroplasmy analysis. Of these subsets, m.3243A>G proportion was plotted in violin plots by cell type or as a color gradient on UMAP plots.

Differential gene expression analysis

Differential gene expression was calculated between MELAS and control samples for each cell type identified by scRNAseq data. Differential gene expression analysis was performed on the multimodal sequencing data for the RPE cluster only.

Transcription factor binding motif enrichment analysis

Transcription factor binding motif enrichment was computed using ChromVAR run in Signac (46). The JASPAR 2020 database of motif position frequencies was used (47). The top marker motifs enriched in each cell type were identified based on average fold change ChromVAR score.

Transmission electron microscopy

Donor samples were fixed in ½ strength Karnovsky's fixative and processed for TEM as described previously (34). Sections were cut at 85 nm with an ultra-microtome (Leica) and collected on copper slot grids coated with Formvar (0.5%) and images collected on a Hitachi HT7800 transmission electron microscope.

Laser Capture Microdissection

Fixed frozen blocks containing neural retina and choroid from both MELAS donors and an unaffected control donor were sectioned at 7μm thickness onto membrane slides (Zeiss). Prior to sectioning, slides were exposed to UV light for 30 minutes in a tissue culture hood, per manufacturer's suggestion. Mounted sections were rinsed once with cold deionized water prior to laser capture microdissection (LCM). LCM was carried out using the PALM Combi LCM system (Zeiss). Regions of interest (i.e., photoreceptors, RPE, or choriocapillaris) were traced and captured into adhesive cap tubes following the manufacturer's recommended protocol. DNA was

isolated using the Arcturus PicoPure DNA Extraction Kit (Applied Biosystems). Briefly, samples were digested overnight in a Proteinase K solution at 65˚C, followed by a 10-minute inactivation at 95˚C. DNA was stored undiluted at -20˚C.

Genotyping by digital PCR

Custom probe-based assays were designed against either variant at the m.3243 locus as previously described (30). Probes were designed incorporating several locked nucleic acids (indicated with '+' below) to attain adequate binding specificity to provide allelic specificity and manufactured by Integrated DNA Technologies (Coralville, IA). Digital PCR was carried out using the 3D PCR platform (Applied Biosystems). For the m.3243A/G assay, the following mixture was used (per reaction): 8.7µL 3D Master Mix V2 (Applied Biosystems), 0.85µL A-FAM Probe (5µM), 0.85µL G-HEX Probe (5µM), 1.67µL F Primer (5µM), 1.67 R Primer (5µM), 1.66µL H2O, 1µL DNA lysate from LCM. 14.5µL of the mixture was loaded into a 3D Digital PCR chip using the QuantStudio Digital PCR Chip Loaded following the manufacturer's instruction.

Probe Name	Color	F Primer	R Primer	Probe
MT-TL 3243 A	FAM	ACACCCACCCAAGAACA	AGGAATTGAACCTCTGACTGTAAA	AT+GG+CA+G+A+GC
MT-TL 3243 G	HEX	ACACCCACCCAAGAACA	AGGAATTGAACCTCTGACTGTAAA	ATGGCA+G+G+GC

The thermal cycler program was run as follows for the m.3243A/G assay: Step 1: 95˚C – 10 minutes; Step 2: 56˚C – 2 minutes; Step 3: 98˚C – 30 seconds; Step 4: 60˚C – 2 minutes, with steps 2 and 3 cycled 45 times. After amplification, chips were allowed to equilibrate to room temperature for approximately 10 minutes and were then read. Data were analyzed using the 3D Quantstudio cloud-based web application, and the automatically generated thresholds were used. The proportion of G allele for the *MT-TL1* A/G heteroplasmy assay was calculated as:

$$m.3243G \text{ proportion} = \frac{\text{Copies VIC per µL}}{\text{Copies VIC per µL} + \text{Copies FAM per µL}}$$

Isolation of Primary Retina Pigmented Epithelium (RPE)

Donor eyes were dissected as previously described (42). Following removal of the vitreous and neural retina, the RPE was manually scraped away from the underlying Bruch's membrane and choroid. Cells were

transferred directly to cryopreservation media and cryopreserved in the same manner as described for the dissociated neural retina and RPE/choroid above.

Human donor samples

Donor ID	Age	Sex	PMI (h:m)	Experiment	Genotype	Region
MELAS 1	41	Male	8:09	1, 2, 3, 4, 5, 7	m.3243A>G	Macula + Periphery
MELAS 2	70	Female	7:38	5, 6, 7	m.3243A>G	Macula
Control 1	40	Male	5:40	1, 2	n/a	Macula + Periphery
Control 2	59	Male	6:59	3	n/a	Periphery
Control 3	60	Male	6:14	4	n/a	Periphery
Control 4	70	Female	5:22	5	n/a	Macula
Control 5	83	Male	8:30	6	n/a	Macula
Control 6	77	Male	8:00	6	n/a	Macula
Control 7	41	Female	7:38	7	n/a	Macula

1. scRNAseq
2. mt-scATACseq
3. Multimodal profiling (RPE/Choroid)
4. Multimodal profiling (Purified RPE)
5. LCM-dPCR
6. Transmission electron microscopy
7. Hematoxylin and eosin staining

Statistics

Differential gene expression analysis was performed using the 'FindMarkers' of Seurat. P values for the differential expression of genes between condition were generated based on the Wilcoxon rank sum test statistic (43). P values were adjusted for multiple testing using the Bonferroni correction based on all features (genes) present in the dataset.

Study approval

Human donor eyes used in this study were acquired from the Iowa Lions Eye Bank or the Minnesota Lions Eye Bank in accordance with the Declaration of Helsinki following consent of the donor's next of kin. The human subjects aspects of the study were approved by the Institutional Review Board of the University of Iowa. The patients for whom clinical data is available were seen at the University of Iowa Hospitals and Clinics and provided written informed consent for inclusion in this study.

AUTHOR CONTRIBUTIONS

N.K.M., A.P.V., B.A.T., and R.F.M. designed the study. N.K.M., A.P.V., M.J.F.-W., X.L., and K.V. conducted experiments and acquired data. E.M.S. examined the proband and provided clinical data. N.K.M. and A.P.V. analyzed the data. N.K.M. wrote the original draft of the manuscript. A.P.V., E.M.S., B.A.T., and R.F.M. edited the manuscript. All authors reviewed and approved the final manuscript.

ACKNOWLEDGEMENTS

This work was supported by the National Institutes of Health (T32GM008629, T32GM139776, P30EY025580, F30EY031923, EY033308), Research to Prevent Blindness, the Elmer and Sylvia Sramek Charitable Trust, and The Edward N. & Della L. Thome Memorial Foundation. The authors would like to acknowledge the extraordinary generosity of the tissue donors and their families that made this study possible. We would also like to thank K. Mulfaul, E. Burnight and the other members of the Tucker and Mullins laboratories for their helpful conversation and advice. Data presented herein were obtained at the Genomics Division of the Iowa Institute of Human Genetics which is supported, in part, by the University of Iowa Carver College of Medicine. We would like to acknowledge specifically the advice and support of M. Boes, G. Hauser, K. Knudtson, A. Sheehan, and E. Snir. Colors used in plots were based on www.colorbrewer2.org by Cynthia A. Brewer, Department of Geography, Pennsylvania State University.

REFERENCES

1.   Hahn A, and Zuryn S. The Cellular Mitochondrial Genome Landscape in Disease. *Trends Cell Biol.* 2019;29(3):227-40.

2.   Aryaman J, Johnston IG, and Jones NS. Mitochondrial Heterogeneity. *Front Genet.* 2018;9:718.

3.   Boggan RM, Lim A, Taylor RW, McFarland R, and Pickett SJ. Resolving complexity in mitochondrial disease: Towards precision medicine. *Mol Genet Metab.* 2019;128(1-2):19-29.

4.   Moggio M, Colombo I, Peverelli L, Villa L, Xhani R, Testolin S, et al. Mitochondrial disease heterogeneity: a prognostic challenge. *Acta Myol.* 2014;33(2):86-93.

5.   Stewart JB, and Chinnery PF. The dynamics of mitochondrial DNA heteroplasmy: implications for human health and disease. *Nat Rev Genet.* 2015;16(9):530-42.

6.   Filograna R, Mennuni M, Alsina D, and Larsson NG. Mitochondrial DNA copy number in human disease: the more the better? *FEBS Lett.* 2021;595(8):976-1002.

7.   Lightowlers RN, Chinnery PF, Turnbull DM, and Howell N. Mammalian mitochondrial genetics: heredity, heteroplasmy and disease. *Trends in Genetics.* 1997;13(11):450-5.

8.   Scholle LM, Zierz S, Mawrin C, Wickenhauser C, and Urban DL. Heteroplasmy and Copy Number in the Common m.3243A>G Mutation-A Post-Mortem Genotype-Phenotype Analysis. *Genes (Basel).* 2020;11(2).

9.   Li M, Schroder R, Ni S, Madea B, and Stoneking M. Extensive tissue-related and allele-related mtDNA heteroplasmy suggests positive selection for somatic mutations. *Proc Natl Acad Sci U S A.* 2015;112(8):2491-6.

10. Hubner A, Wachsmuth M, Schroder R, Li M, Eis-Hubinger AM, Madea B, et al. Sharing of heteroplasmies between human liver lobes varies across the mtDNA genome. *Sci Rep.* 2019;9(1):11219.

11. Walker MA, Lareau CA, Ludwig LS, Karaa A, Sankaran VG, Regev A, et al. Purifying Selection against Pathogenic Mitochondrial DNA in Human T Cells. *N Engl J Med.* 2020;383(16):1556-63.

12. Stewart JB, and Chinnery PF. Extreme heterogeneity of human mitochondrial DNA from organelles to populations. *Nat Rev Genet.* 2021;22(2):106-18.

13. Parakatselaki ME, and Ladoukakis ED. mtDNA Heteroplasmy: Origin, Detection, Significance, and Evolutionary Consequences. *Life (Basel).* 2021;11(7).

14. Goto Y, Nonaka I, and Horai S. A mutation in the tRNA(Leu)(UUR) gene associated with the MELAS subgroup of mitochondrial encephalomyopathies. *Nature.* 1990;348(6302):651-3.

15. Flierl A, Reichmann H, and Seibel P. Pathophysiology of the MELAS 3243 transition mutation. *J Biol Chem.* 1997;272(43):27189-96.

16. Kirino Y, Yasukawa T, Ohta S, Akira S, Ishihara K, Watanabe K, et al. Codon-specific translational defect caused by a wobble modification deficiency in mutant tRNA from a human mitochondrial disease. *Proc Natl Acad Sci U S A.* 2004;101(42):15070-5.

17. Janssen GM, Hensbergen PJ, van Bussel FJ, Balog CI, Maassen JA, Deelder AM, et al. The A3243G tRNALeu(UUR) mutation induces mitochondrial dysfunction and variable disease expression without dominant negative acting translational defects in complex IV subunits at UUR codons. *Hum Mol Genet.* 2007;16(20):2472-81.

18. Sasarman F, Antonicka H, and Shoubridge EA. The A3243G tRNALeu(UUR) MELAS mutation causes amino acid misincorporation and a combined respiratory chain assembly defect partially suppressed by overexpression of EFTu and EFG2. *Hum Mol Genet.* 2008;17(23):3697-707.

19. Li R, and Guan MX. Human mitochondrial leucyl-tRNA synthetase corrects mitochondrial dysfunctions due to the tRNALeu(UUR) A3243G mutation, associated with mitochondrial encephalomyopathy, lactic acidosis, and stroke-like symptoms and diabetes. *Mol Cell Biol.* 2010;30(9):2147-54.

20. Richter U, McFarland R, Taylor RW, and Pickett SJ. The molecular pathology of pathogenic mitochondrial tRNA variants. *FEBS Lett.* 2021;595(8):1003-24.

21. de Laat P, Koene S, van den Heuvel LP, Rodenburg RJ, Janssen MC, and Smeitink JA. Clinical features and heteroplasmy in blood, urine and saliva in 34 Dutch families carrying the m.3243A > G mutation. *J Inherit Metab Dis.* 2012;35(6):1059-69.

22. Coussa RG, Parikh S, and Traboulsi EI. Mitochondrial DNA A3243G variant-associated retinopathy: Current perspectives and clinical implications. *Surv Ophthalmol.* 2021;66(5):838-55.

23. Smith PR, Bain SC, Good PA, Hattersley AT, Barnett AH, Gibson JM, et al. Pigmentary retinal dystrophy and the syndrome of maternally inherited diabetes and deafness caused by the mitochondrial DNA 3243 tRNALeu A to G mutation11The authors have no proprietary interest in the development or marketing of any device or medications mentioned in the article or any competing device. *Ophthalmology.* 1999;106(6):1101-8.

24. Oh JK, Lima de Carvalho JR, Jr., Nuzbrokh Y, Ryu J, Chemudupati T, Mahajan VB, et al. Retinal Manifestations of Mitochondrial Oxidative Phosphorylation Disorders. *Invest Ophthalmol Vis Sci.* 2020;61(12):12.

25. Iwanicka-Pronicka K, Pollak A, Skorka A, Lechowicz U, Pajdowska M, Furmanek M, et al. Postlingual hearing loss as a mitochondrial 3243A>G mutation phenotype. *PLoS One.* 2012;7(10):e44054.

26. Ciafaloni E, Ricci E, Shanske S, Moraes CT, Silvestri G, Hirano M, et al. MELAS: clinical features, biochemistry, and molecular genetics. *Ann Neurol.* 1992;31(4):391-8.

27. Finsterer J, and Laccone F. Phenotypic Heterogeneity in 5 Family Members with the Mitochondrial Variant m.3243A>G. *Am J Case Rep.* 2020;21:e927938.

28. Grady JP, Pickett SJ, Ng YS, Alston CL, Blakely EL, Hardy SA, et al. mtDNA heteroplasmy level and copy number indicate disease burden in m.3243A>G mitochondrial disease. *EMBO Mol Med.* 2018;10(6).

29. Pickett SJ, Grady JP, Ng YS, Gorman GS, Schaefer AM, Wilson IJ, et al. Phenotypic heterogeneity in m.3243A>G mitochondrial disease: The role of nuclear factors. *Ann Clin Transl Neurol.* 2018;5(3):333-45.

30. Mullin NK, Anfinson KR, Riker MJ, Wieland KL, Tatro NJ, Scheetz TE, et al. Sensitive quantification of m.3243A>G mutational proportion in non-retinal tissues and its relationship with visual symptoms. *Hum Mol Genet.* 2021.

31. DOLLERY CT, BULPITT CJ, and KOHNER EM. Oxygen supply to the retina from the retinal and choroidal circulations at normal and increased arterial oxygen tensions. *Investigative Ophthalmology & Visual Science.* 1969;8(6):588-94.

32. Lutty GA, and McLeod DS. Development of the hyaloid, choroidal and retinal vasculatures in the fetal human eye. *Prog Retin Eye Res.* 2018;62:58-76.

33. Lutty GA, Hasegawa T, Baba T, Grebe R, Bhutto I, and McLeod DS. Development of the human choriocapillaris. *Eye (Lond).* 2010;24(3):408-15.

34. Hasegawa T, McLeod DS, Bhutto IA, Prow T, Merges CA, Grebe R, et al. The embryonic human choriocapillaris develops by hemo-vasculogenesis. *Dev Dyn.* 2007;236(8):2089-100.

35. Sakuta R, Goto Y-i, Horai S, and Nonaka I. Mitochondrial DNA mutations at nucleotide positions 3243 and 3271 in mitochondrial myopathy, encephalopathy, lactic acidosis, and stroke-like episodes: A comparative study. *Journal of the Neurological Sciences.* 1993;115(2):158-60.

36. Latkany P, Ciulla TA, Cucchillo P, and Malkoff MD. Mitochondrial maculopathy: geographic atrophy of the macula in the MELAS associated A to G 3243 mitochondrial DNA point mutation. *American Journal of Ophthalmology.* 1999;128(1):112-4.

37. Tripathy K, Sarma B, and Mazumdar S. Outer retinal tubulation and inner retinal pseudocysts in a patient with maternally inherited diabetes and deafness evaluated with optical coherence tomography angiogram. *Indian J Ophthalmol.* 2020;68(1):250-3.

38. Raja MS, Goldsmith C, and Burton BJ. Outer retinal tubulations in maternally inherited diabetes and deafness (MIDD)-associated macular dystrophy. *Graefes Arch Clin Exp Ophthalmol.* 2013;251(9):2265-7.

39. Han IC, Critser DB, and Stone EM. Subliminal Message Outer Retinal Tubulations Resembling Mitochondria in Maternally Inherited Diabetes and Deafness. *Ophthalmol Retina.* 2020;4(11):1102.

40. Lareau CA, Ludwig LS, Muus C, Gohil SH, Zhao T, Chiang Z, et al. Massively parallel single-cell mitochondrial DNA genotyping and chromatin profiling. *Nat Biotechnol.* 2021;39(4):451-61.

41. Voigt AP, Whitmore SS, Flamme-Wiese MJ, Riker MJ, Wiley LA, Tucker BA, et al. Molecular characterization of foveal versus peripheral human retina by single-cell RNA sequencing. *Exp Eye Res.* 2019;184:234-42.

42. Voigt AP, Mulfaul K, Mullin NK, Flamme-Wiese MJ, Giacalone JC, Stone EM, et al. Single-cell transcriptomics of the human retinal pigment epithelium and choroid in health and macular degeneration. *Proc Natl Acad Sci U S A.* 2019;116(48):24100-7.

43. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al. Comprehensive Integration of Single-Cell Data. *Cell.* 2019;177(7):1888-902 e21.

44. Sreekumar PG, Ishikawa K, Spee C, Mehta HH, Wan J, Yen K, et al. The Mitochondrial-Derived Peptide Humanin Protects RPE Cells From Oxidative Stress, Senescence, and Mitochondrial Dysfunction. *Invest Ophthalmol Vis Sci.* 2016;57(3):1238-53.

45.   Kopinski PK, Janssen KA, Schaefer PM, Trefely S, Perry CE, Potluri P, et al. Regulation of nuclear epigenome by mitochondrial DNA heteroplasmy. *Proc Natl Acad Sci U S A.* 2019;116(32):16028-35.

46.   Schep AN, Wu B, Buenrostro JD, and Greenleaf WJ. chromVAR: inferring transcription-factor-associated accessibility from single-cell epigenomic data. *Nat Methods.* 2017;14(10):975-8.

47.   Castro-Mondragon JA, Riudavets-Puig R, Rauluseviciute I, Berhanu Lemma R, Turchi L, Blanc-Mathieu R, et al. JASPAR 2022: the 9th release of the open-access database of transcription factor binding profiles. *Nucleic Acids Res.* 2022;50(D1):D165-D73.

48.   Homma K, Toda E, Osada H, Nagai N, Era T, Tsubota K, et al. Taurine rescues mitochondria-related metabolic impairments in the patient-derived induced pluripotent stem cells and epithelial-mesenchymal transition in the retinal pigment epithelium. *Redox Biol.* 2021;41:101921.

49.   Dolz-Marco R, Litts KM, Tan ACS, Freund KB, and Curcio CA. The Evolution of Outer Retinal Tubulation, a Neurodegeneration and Gliosis Prominent in Macular Diseases. *Ophthalmology.* 2017;124(9):1353-67.

50.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, et al. Integrated analysis of multimodal single-cell data. *Cell.* 2021;184(13):3573-87 e29.

51.   Zhou M, Geathers JS, Grillo SL, Weber SR, Wang W, Zhao Y, et al. Role of Epithelial-Mesenchymal Transition in Retinal Pigment Epithelium Dysfunction. *Front Cell Dev Biol.* 2020;8:501.

52.   Muller PL, Treis T, Pfau M, Esposti SD, Alsaedi A, Maloca P, et al. Progression of Retinopathy Secondary to Maternally Inherited Diabetes and Deafness - Evaluation of Predicting Parameters. *Am J Ophthalmol.* 2020;213:134-44.

53.   Monnot S, Gigarel N, Samuels DC, Burlet P, Hesters L, Frydman N, et al. Segregation of mtDNA throughout human embryofetal development: m.3243A>G as a model system. *Hum Mutat.* 2011;32(1):116-25.

54.   Barrett A, Arbeithuber B, Zaidi A, Wilton P, Paul IM, Nielsen R, et al. Pronounced somatic bottleneck in mitochondrial DNA of human hair. *Philos Trans R Soc Lond B Biol Sci.* 2020;375(1790):20190175.

55.   Voigt AP, Mullin NK, Whitmore SS, DeLuca AP, Burnight ER, Liu X, et al. Human photoreceptor cells from different macular subregions have distinct transcriptional profiles. *Hum Mol Genet.* 2021;30(16):1543-58.

56. Ortolan D, Sharma R, Volkov A, Maminishkis A, Hotaling NA, Huryn LA, et al. Single-cell-resolution map of human retinal pigment epithelium helps discover subpopulations with differential disease sensitivity. *Proc Natl Acad Sci U S A.* 2022;119(19):e2117553119.

57. Litts KM, Ach T, Hammack KM, Sloan KR, Zhang Y, Freund KB, et al. Quantitative Analysis of Outer Retinal Tubulation in Age-Related Macular Degeneration From Spectral-Domain Optical Coherence Tomography and Histology. *Invest Ophthalmol Vis Sci.* 2016;57(6):2647-56.

58. Xue K, Oldani M, Jolly JK, Edwards TL, Groppe M, Downes SM, et al. Correlation of Optical Coherence Tomography and Autofluorescence in the Outer Retina and Choroid of Patients With Choroideremia. *Invest Ophthalmol Vis Sci.* 2016;57(8):3674-84.

59. Hageman GS, Zhu XL, Waheed A, and Sly WS. Localization of carbonic anhydrase IV in a specific capillary bed of the human eye. *Proc Natl Acad Sci U S A.* 1991;88(7):2716-20.

60. Pek NMQ, Phua QH, Ho BX, Pang JKS, Hor JH, An O, et al. Mitochondrial 3243A > G mutation confers pro-atherogenic and pro-inflammatory properties in MELAS iPS derived endothelial cells. *Cell Death Dis.* 2019;10(11):802.

61. Viegas FO, and Neuhauss SCF. A Metabolic Landscape for Maintaining Retina Integrity and Function. *Front Mol Neurosci.* 2021;14:656000.

62. Pan WW, Wubben TJ, and Besirli CG. Photoreceptor metabolic reprogramming: current understanding and therapeutic implications. *Commun Biol.* 2021;4(1):245.

63. Kanow MA, Giarmarco MM, Jankowski CS, Tsantilas K, Engel AL, Du J, et al. Biochemical adaptations of the retina and retinal pigment epithelium support a metabolic ecosystem in the vertebrate eye. *Elife.* 2017;6.

64. Adijanto J, Du J, Moffat C, Seifert EL, Hurle JB, and Philp NJ. The retinal pigment epithelium utilizes fatty acids for ketogenesis. *J Biol Chem.* 2014;289(30):20570-82.

65. Chomyn A, Enriquez JA, Micol V, Fernandez-Silva P, and Attardi G. The mitochondrial myopathy, encephalopathy, lactic acidosis, and stroke-like episode syndrome-associated human mitochondrial tRNALeu(UUR) mutation causes aminoacylation deficiency and concomitant reduced association of mRNA with ribosomes. *J Biol Chem.* 2000;275(25):19198-209.

66.	Hamalainen RH, Manninen T, Koivumaki H, Kislin M, Otonkoski T, and Suomalainen A. Tissue- and cell-type-specific manifestations of heteroplasmic mtDNA 3243A>G mutation in human induced pluripotent stem cell-derived disease model. *Proc Natl Acad Sci U S A.* 2013;110(38):E3622-30.

67.	Latorre-Pellicer A, Lechuga-Vieco AV, Johnston IG, Hamalainen RH, Pellico J, Justo-Mendez R, et al. Regulation of Mother-to-Offspring Transmission of mtDNA Heteroplasmy. *Cell Metab.* 2019;30(6):1120-30 e5.

68.	Moreira JD, Gopal DM, Kotton DN, and Fetterman JL. Gaining Insight into Mitochondrial Genetic Variation and Downstream Pathophysiology: What Can i(PSCs) Do? *Genes (Basel).* 2021;12(11).

69.	Mimitou EP, Lareau CA, Chen KY, Zorzetto-Fernandes AL, Hao Y, Takeshima Y, et al. Scalable, multimodal profiling of chromatin accessibility, gene expression and protein levels in single cells. *Nat Biotechnol.* 2021.

70.	Voigt AP, Whitmore SS, Lessing ND, DeLuca AP, Tucker BA, Stone EM, et al. Spectacle: An interactive resource for ocular single-cell RNA sequencing data analysis. *Exp Eye Res.* 2020;200:108204.