Chrna5 and lynx prototoxins identify acetylcholine super-responder subplate neurons

Sridevi Venkatesan[1], Tianhui Chen[1], Yupeng Liu[1], Eric E Turner[2], Shreejoy Tripathy[1,3,4,5], *Evelyn K Lambe[1,4,6]

1. Department of Physiology, Temerty Faculty of Medicine, University of Toronto, Toronto ON, Canada
2. Center for Integrative Brain Research, Seattle Children's Research Institute, Seattle WA, USA
3. Krembil Centre for Neuroinformatics, Centre for Addiction and Mental Health, Toronto ON, Canada
4. Department of Psychiatry, University of Toronto, Toronto ON, Canada
5. Institute of Medical Science, University of Toronto, Toronto ON, Canada
6. Department of Obstetrics and Gynecology, University of Toronto, Toronto ON, Canada

Corresponding author:
E.K. Lambe, Ph.D.
1 King's College Circle
Toronto ON
Canada M5S 1A8
(416) 946-0910
Email: evelyn.lambe@utoronto.ca

Summary (150 words)
Attention depends on cholinergic excitation of prefrontal neurons but is sensitive to perturbation of α5-containing nicotinic receptors encoded by *Chrna5*. However, *Chrna5*-expressing (Chrna5+) neurons remain enigmatic, despite their potential as a target to improve attention. Here, we generate complex transgenic mice to probe Chrna5+ neurons and their sensitivity to endogenous acetylcholine. Through opto-physiological experiments, we discover that Chrna5+ neurons contain a distinct population of acetylcholine super-responders. Leveraging single-cell transcriptomics, we discover molecular markers conferring subplate identity on this subset. We determine that Chrna5+ super-responders express a unique complement of GPI-anchored lynx prototoxin genes (*Lypd1*, *Ly6g6e,* and *Lypd6b*), predicting distinct nicotinic receptor regulation. To manipulate lynx regulation of endogenous nicotinic responses, we developed a pharmacological strategy guided by transcriptomic predictions. Overall, we reveal *Chrna5*-Cre mice as a transgenic tool to target the diversity of subplate neurons in adulthood, yielding new molecular strategies to manipulate their cholinergic activation relevant to attention disorders.

Keywords
Acetylcholine, Chrna5, nicotinic receptor, alpha5, prefrontal cortex, subplate, lynx prototoxins, transcriptomics, optogenetics, electrophysiology

Introduction

Cholinergic modulation of the medial prefrontal cortex (mPFC) is essential for attention and detection of sensory cues [1–4]. Deep-layer pyramidal neurons in the PFC are critically involved in such executive function [5–7] and are robustly excited by acetylcholine [8,9], via nicotinic and muscarinic receptor activation [10]. The α5 nicotinic receptor subunit encoded by *Chrna5* is specifically expressed in deep-layer pyramidal neurons [11,12], forming high-affinity nicotinic receptors in combination with α4 and β2 subunits. Electrophysiological, behavioural, and genetic evidence in both rodents and humans point to an important role of *Chrna5* expression for nicotinic receptor function, attention, and executive function.

Constitutive deletion of *Chrna5* in mice, or knockdown in the adult rat PFC disrupt attention and reduce nicotinic receptor activation by exogenous acetylcholine stimulation in layer 6 neurons [13–15]. Optogenetic experiments in *Chrna5-/-* mice show that *Chrna5* is required for rapid onset of postsynaptic cholinergic activation and prevents desensitization of the endogenous cholinergic response during prolonged stimulation [16]. In humans, the non-synonymous rs16969968 (D398N) polymorphism in *Chrna5* is associated with nicotine dependence, schizophrenia and cognitive impairment [17–19]. Nicotinic α4β2α5 receptors with the D398N polymorphism have partial loss of function, attributed to changes in receptor desensitization, calcium permeability, or membrane trafficking [20–22].

Despite α5 nicotinic receptor involvement in attention and prefrontal cholinergic activation, systematic characterization of *Chrna5*-expressing neurons using modern genetic tools is lacking. Deep-layer neurons include diverse corticothalamic (L6CT), corticocortical, and layer 6b (L6b) populations [23–25]. *Chrna5* is predicted to be expressed in L6CT neurons [8,26], which are usually identified by their expression of *Syt6*, a conserved L6CT neuronal marker [27,28,29]. *Syt6*-Cre mice have been widely used to characterize prefrontal L6CT neurons and their cholinergic properties [30–32]. However, it is unclear whether these are the same neurons expressing *Chrna5*. Characterization of *Chrna5*-expressing neurons has been limited by the lack of verified antibodies for the α5 subunit that could be used for post-hoc immunostaining. Previous BAC-transgenic mice labeling *Chrna5*-expressing neurons had altered expression of other genes in the tightly linked *Chrna5/a3/b4* gene cluster, limiting their use for functional examination [33]. This issue was circumvented by disrupting the open reading frames of *Chrna3/b4* in the BAC transgene to generate a *Chrna5*-Cre mouse without misexpression artifacts [34].

Here, we create compound transgenic mice to investigate how prefrontal *Chrna5*-expressing (Chrna5+) neurons respond to optogenetic release of endogenous acetylcholine. The fast and strong response prompted a multi-approach examination of Chrna5+ neurons together with a control population of *Syt6*-expressing (Syt6+) neurons, a more traditional molecular marker of deep-layer prefrontal cortex. We demonstrate a large fraction of Chrna5+ neurons have distinctive high-affinity cholinergic responses. Single-cell RNAseq analysis reveals the expression of several subplate markers (*Cplx3*, *Ctgf*, and *Lpar1*) in this Chrna5+ subset, identifying them as subplate neurons born early in development that are critical for establishing thalamocortical connectivity [35–37]. Chrna5+ subplate neurons show distinct expression pattern of GPI-anchored lynx prototoxins (*Ly6g6e*, *Lypd1*, and *Lypd6b*) capable of exerting complex

modulation of nicotinic receptor properties [38,39]. As predicted from the transcriptomic analysis,
our pharmacological manipulations targeting lynx prototoxins successfully alter endogeneous
nicotinic responses, revealing cell-type specific lynx regulation of nicotinic receptors.

Recent studies have examined cell-type specific modulation of nicotinic receptors by
different lynx prototoxins and the consequences for cortical development and cognition [40–43].
Here, we discovered endogenous lynx modulation of nicotinic properties relevant for attention in
subplate neurons expressing *Chrna5*. Our work highlights *Chrna5*-Cre mice as a transgenic tool
to target acetylcholine super-responder subplate neurons and identifies strategies to fine-tune
their cholinergic activation by manipulating GPI-anchored lynx prototoxins.

Results

Chrna5 expression identifies acetylcholine 'super-responders'

To characterize prefrontal *Chrna5*-expressing (Chrna5+) neurons, we generated triple transgenic
Chrna5-Cre$^{/+}$Ai14$^{/+}$ChAT-ChR2$^{/+}$ mice (**Fig 1A**) and examined optogenetic cholinergic
responses in labeled Chrna5+ neurons. These mice expressed tdTomato in Chrna5+ neurons and
channelrhodopsin-2 in cholinergic axons as seen by two-photon imaging (**Fig 1B**). The location
of reporter labeled Chrna5+ neurons across the brain in *Chrna5*-Cre mice is specific to layer 6
and found across the entire cortical mantle (**Fig S1**). We recorded endogenous cholinergic
responses in labeled Chrna5+ neurons with optophysiology in mPFC slices, using unlabeled
pyramidal neurons as a control (**Fig 1C**). Optogenetic Chrna5+ neurons clearly possessed larger-
amplitude cholinergic responses with significantly faster onset compared to control neurons (**Fig
1D-F**). The rising slope was significantly larger in Chrna5+ neurons (220 ± 32 pA/s, 24 cells
from 4 mice) compared to control (123 ± 22 pA/s, 15 cells; $t_{(37)} = 2.19$, $P = 0.02$, unpaired t-test;
Cohen's d: 0.72). Similarly, peak current evoked by optogenetic acetylcholine release was
greater in Chrna5+ neurons (15 ± 2 pA) compared to control (8 ± 1 pA, $t_{(37)} = 2.52$, $P = 0.02$;
Cohen's d: 0.83), as well as the area of the cholinergic response ($t_{(37)} = 2.06$, $P = 0.046$). Intrinsic
electrophysiological properties, however, did not differ between Chrna5+ and unlabeled neurons
(**Table S1**). Additionally, we determined that Chrna5+ neurons are under presynaptic muscarinic
autoinhibitory control, which can be relieved by atropine, resulting in even larger responses (post
atropine response: $194 \pm 42\%$ of baseline, n = 6 cells). Our application of the nicotinic antagonist
DHBE eliminated the optogenetic cholinergic response (post DHBE response : 3 ± 2 % of
baseline, n = 4 cells), indicating the relevant nicotinic receptors contain β2 subunits. This
characterization of Chrna5+ neurons revealed them to be acetylcholine 'super-responders' with
stronger and faster onset cholinergic responses distinct from other deep-layer pyramidal neurons.
We next examined whether Chrna5+ neurons in other cortical regions are also acetylcholine
super-responders. Chrna5+ labeled neurons deep in layer 6 of primary somatosensory cortex
(SSp) were also found to show significantly stronger and faster onset nicotinic responses to
optogenetic acetylcholine release (**Fig S2)** confirming that *Chrna5*-labeling identifies
acetylcholine super-responders in multiple cortical areas.

To further probe the distinct cholinergic responses of prefrontal Chrna5+ neurons, we
examined factors differentiating these cells from neurons labeled by *Syt6* expression (Syt6+),
which has hereto been the predominant marker of layer 6 corticothalamic neurons used to
characterize their function during PFC-dependent tasks [28,30,31,44]. Since the extent of overlap
between *Chrna5* and *Syt6* expressing deep-layer pyramidal neuron populations was unclear, we
adopted an imaging strategy to visualize the distribution of Chrna5+ and Syt6+ neurons in the
PFC and determine the exact proportion of distinctive non-overlapping Chrna5+ neurons. We
generated a compound transgenic *Chrna5*-Cre$^{/+}$Ai14$^{/+}$*Syt6*-EGFP$^{/+}$ mouse to simultaneously
express tdTomato in Chrna5+ neurons and EGFP in Syt6+ neurons and performed confocal and
two-photon imaging of the endogenous fluorescence of these reporters in mPFC brain slices.
Chrna5+ and Syt6+ neurons were both present primarily in layer 6, with a few Chrna5+ neurons
in layer 5 (**Fig 1G, I**). Fig 1J (left) shows the proportion of all cells expressing tdTomato
(*Chrna5*), GFP (*Syt6),* or both, as a percentage of the total number of fluorescent cells in that
region. Closer investigation confirmed the existence of a substantial proportion of exclusively
Chrna5+ neurons (37% of all labeled cells) which do not express *Syt6*, in addition to overlapping
Chrna5+Syt6+ neurons (39%) which express both markers, and exclusively Syt6+ neurons
(24%) which do not express *Chrna5* (N = 4 mice, **Fig 1H, J**). Thus, nearly half of all *Chrna5*-
expressing neurons are not labeled by *Syt6*-expression and would have been excluded in previous
studies using *Syt6*-Cre mice. Furthermore, *Chrna5*-expressing neurons are found even in primary
visual cortex, where *Syt6*-labeling is greatly reduced [31,45] (**Fig S3**).

Distinct Chrna5+ neurons with highly-resilient nicotinic receptor responses

To elucidate the distinct subset of Chrna5+ neurons not found with Syt6-labeling approaches, we
measured acetylcholine-evoked signals in multiple neurons simultaneously with *ex vivo*
GCaMP6s Ca^{2+} imaging. We generated transgenic mice expressing GCaMP6s in either Chrna5+
(*Chrna5*-Cre$^{/+}$/Ai96$^{/+}$) or Syt6+ (*Syt6*-Cre$^{/+}$/Ai96$^{/+}$) neurons and performed two-photon imaging
of mPFC layer 6 (**Fig 2A**). We measured Ca^{2+} signals evoked by acetylcholine (1 mM, 15 s) in
Chrna5+ (**Video S1**) and Syt6+ neurons (**Video S2**). To pharmacologically interrogate these
populations, we measured changes in the Ca^{2+} signal and proportions of acetylcholine-responsive
neurons (**Fig 2A, B**). Application of the competitive nicotinic antagonist DHBE (10 µM, 10 min)
left residual acetylcholine-evoked Ca^{2+} signals in Chrna5+ neurons (35 ± 3 % of baseline, n = 71
cells, 6 mice; **Fig 2C i**) that were greater to those in Syt6+ neurons (21 ± 3% of baseline, n = 112
cells, 7 mice; Mann Whitney U = 2400, $P < 10^{-4}$). The cumulative distribution of responses
remaining after DHBE was significantly right-shifted in Chrna5+ neurons compared to Syt6+
neurons (**Fig 2C ii**, Kolmogorov Smirnov D = 0.37, $P < 10^{-4}$). A majority of Chrna5+ neurons
(~83%) still showed acetylcholine-evoked responses after DHBE, whereas fewer Syt6+ neurons
(~50%) retained their responses, with a complete elimination of acetylcholine-evoked responses
in the rest (**Fig 2C iii**, Fisher's exact test: $P < 10^{-4}$). Yet, the addition of muscarinic antagonist
atropine did not attenuate the striking differences between Chrna5+ and Syt6+ neurons, raising
the possibility of an underlying nicotinic mechanism (**Fig 2D**). In the presence of DHBE +
atropine, a subset of Chrna5+ neurons still showed substantial acetylcholine-evoked Ca^{2+} signals
(6 ± 1 % of baseline; **Fig 2D i**), whereas almost all Syt6+ neurons' responses were completely
blocked (0.2 ± 0.1 % of baseline; Mann Whitney U = 2083, $P < 10^{-4}$) as seen from the

cumulative distribution (**Fig 2D ii**, Kolmogorov Smirnov D = 0.36, $P < 10^{-4}$). The proportion of
Chrna5+ and Syt6+ neurons showing acetylcholine-evoked responses after DHBE+ Atropine
was significantly different (41% Chrna5+ vs 6% Syt6+, Fisher's exact test: $P < 10^{-4}$, **Fig 2D iii**).
Chrna5+ neurons resilient to DHBE+ Atropine within a slice appear to be localized to layer 6b
(**Fig S4**). Since DHBE is a competitive antagonist, it can be outcompeted by exogenous
acetylcholine at high affinity nicotinic receptors. Therefore, we hypothesized that the resilient
exogenous acetylcholine-evoked responses would require non-competitive antagonist.

To address the hypothesis of higher-affinity nicotinic binding and its consequences for
spiking in these neurons, we switched to whole-cell recording in individual Chrna5+ and Syt6+
neurons. We recorded current clamp responses to acetylcholine (1 mM, 15 s) in labeled Chrna5+
and Syt6+ neurons from *Chrna5*-Cre$^{/+}$Ai14$^{/+}$ and *Syt6*-Cre$^{/+}$Ai14$^{/+}$ or *Syt6*-EGFP mice
respectively (**Fig 2E**). Chrna5+ neurons showed stronger acetylcholine-evoked firing: attaining
significantly higher peak firing frequency (29 ± 6 Hz, n=12 cells, 6 mice; $t_{(24)}$ = 2.74, P = 0.01,
unpaired t-test; Cohen's d: 1.08) compared to Syt6+ neurons (13 ± 2 Hz, n = 14 cells, 5 mice;
Fig 2E). The intrinsic electrophysiological properties of Chrna5+ and Syt6+ neurons did not
show statistically significant differences (**Table S2**). We next examined the sensitivity of
acetylcholine-evoked firing to competitive nicotinic receptor block by DHBE in the presence of
atropine. Acetylcholine-evoked firing was completely eliminated in all Syt6+ neurons (**Fig 2F**)
whereas a large subset of Chrna5+ neurons (7/11) retained their ability to respond to
acetylcholine (Average peak firing rate in Chrna5+ neurons: 6 ± 2 Hz; $t_{(19)}$ = 3.22, P = 0.004,
unpaired t-test) demonstrating similar resilience to competitive nicotinic receptor block as
observed with Ca^{2+} imaging. We used the non-competitive nicotinic receptor blocker
mecamylamine [46,47] to test our hypothesis that nicotinic receptors in this Chrna5+ subset were
higher affinity and therefore allowed exogenous acetylcholine to outcompete DHBE. The
addition of 5 μM mecamylamine was sufficient to eliminate acetylcholine-evoked firing in all the
Chrna5+ neurons that were resilient to competitive nicotinic block ($t_{(4)}$ = 5.14, P = 0.007, paired
t-test; **Fig 2F**). Together, our Ca^{2+} imaging and electrophysiology experiments revealed the
existence of a distinct subset of Chrna5+ neurons dissimilar to Syt6+ neurons, with high affinity
nicotinic responses resilient to competitive nicotinic antagonism.

Single-cell transcriptomics identifies Chrna5+ subplate neurons with Lynx genes

To determine the molecular identity of distinct Chrna5+ neurons with enhanced cholinergic
responses, we pursued single-cell RNAseq. We extracted gene expression data of L5-6
glutamatergic neurons (n = 2422 cells) in the mouse anterior cingulate cortex from the Allen
Institute single cell RNAseq databases [48–50]. We classified these deep-layer pyramidal neurons
into 3 groups: those expressing only *Chrna5* (Chrna5+, n= 243), both *Chrna5* and *Syt6* (Chrna5+
Syt6+, n = 834), or only *Syt6* (Syt6+, n = 564) **(Fig 3A)**. 781 cells showed no expression of
Chrna5 or *Syt6* and consisted primarily of L6 Intratelencephalic cells which have been
previously shown to have purely muscarinic M2/M4 mediated hyperpolarizing cholinergic
responses [16,51]. We focused on the Chrna5+, Syt6+ and Chrna5+Syt6+ groups to examine their
transcriptomic differences. Prior work has shown that deep cortical neurons can be separated into
L6b, L5, and L6CT subclasses by the expression of characteristic markers and hierarchical

clustering [50]. Single-cell analysis revealed that the Chrna5+ group primarily included L6b (44%), L5 near-projecting (L5NP, 19%), and L6CT neurons (30%), whereas the Chrna5+Syt6+ and Syt6+ groups were predominantly composed of L6CT neurons (>90%) (**Fig 3B**). We further examined the expression of marker genes in these respective groups to validate our cell-classification. Chrna5+ neurons showed distinctive expression of several marker genes, *Ctgf, Cplx3, Kcnab1,* and *Lpar1* (**Fig 3B,C**) associated with subplate neurons [35,52]. Subplate neurons are early-born and vital for brain development, leaving L6b neurons as descendants in adulthood [36,37,53]. Notably, the highest fold enrichment among all differentially expressed genes in Chrna5+ neurons was found for subplate markers *Ctgf* (Fold change, 5.69) and *Cplx3* (3.81) (**Table 1**). All differentially expressed genes are listed in Table S4. Overall, Chrna5+ neurons including both L5NP and L6b subpopulations highly express subplate marker genes. In contrast, *Syt6*-expressing Chrna5+Syt6+ and Syt6+ groups are only enriched in the corticothalamic markers *Foxp2* and *Syt6*, consistent with their corticothalamic subtype. These results support our imaging, electrophysiological, and pharmacological results suggesting the exclusive Chrna5+ population is a distinct cell type from typical L6CT *Syt6*-expressing neurons.

To identify molecular changes predictive of Chrna5+ nicotinic 'super-responders', we examined differential expression of genes that exert effects on postsynaptic cholinergic responses (**Fig 3D-E**). We selected cholinergic receptor genes (nicotinic *Chrna5-2, Chrna7, Chrnb2-4,* and muscarinic subunits *Chrm1-4*), acetylcholinesterase (*Ache*), and members of the family of genes that encode lynx proteins (*Ly6e, Ly6h, Ly6g6e, Lynx1, Lypd1, Lypd6, Lypd6b*) known to allosterically modulate nicotinic receptor responses [38]. We found substantial and highly significant changes in the expression of three lynx prototoxins *Lypd1*, *Ly6g6e* and *Lypd6b* (**Fig 3D**). While both Chrna5+ and Chrna5+Syt6+ populations express *Chrna5,* there was slightly higher expression of *Chrna5* (25% increase) as well as lower expression of the inhibitory muscarinic receptor *Chrm2* (20% decrease) in Chrna5+Syt6+ neurons. There were no significant differences in other nicotinic and muscarinic subunit expression between the two groups. Acetylcholinesterase, the enzyme that breaks down acetylcholine was also highly expressed (50% increase) in Chrna5+ neurons, which may benefit their nicotinic responses by protecting receptors from overactivation and desensitization. Of note, *Chrnb2* expression does seem to be in fewer cells than expected (Fig 3E), which can be attributed to sparse detection of *Chrnb2* levels by scRNAseq approaches [54,56]. The fold-change of the genes in **Fig 3E** between Chrna5+ and Chrna5+Syt6+ neurons is shown in **Table S3**. Notably, the top three genes with highest fold change in Chrna5+ neurons were the GPI-anchored lynx prototoxins: Lynx2 encoded by *Lypd1* (Fold change: 2.55), *Ly6g6e* (2.03), and *Lypd6b* (1.51). The distinct pattern of expression of specific lynx proteins in Chrna5+ neurons suggests unexpectedly complex endogenous control of nicotinic responses in these prefrontal subplate neurons.

Transcriptomic examination of *Chrna5*-expressing neurons in other cortical regions including primary motor (MOp), somatosensory (SSp), and visual cortices (VISp) also confirmed that Chrna5+ neurons consist of L6b and L5/6NP neurons expressing subplate marker genes, in addition to the lynx modulatory proteins as described above in the PFC (**Fig S5**). Therefore, *Chrna5*-expression identifies a conserved population of deep-layer 6 neurons with subplate

identity across multiple cortical regions that display specialized expression of lynx proteins to
modulate nicotinic receptors.

Perturbing native prefrontal cortical lynx-modulation of optogenetic nicotinic responses

To examine whether the molecular machinery of deep layer prefrontal neurons endows them
with greater dynamic range in responding to acetylcholine, we sought to experimentally perturb
endogenous lynx modulation. Members of the lynx-family are GPI-anchored (**Fig 4A**), and work
in cell expression systems [39] suggests these anchors can be cleaved via activation of
phospholipase C (PLC). These experiments are important because the potential impact of GPI
cleavage on nicotinic responses in a native system is not well understood. We hypothesized that
perturbing lynx-mediated control could affect endogenous nicotinic properties in a complex
manner (**Fig 4A**) since both positive (eg. Ly6g6e) and negative modulatory lynx proteins (eg.
Lynx1) are expressed. To cleave GPI-anchored proteins, we used the PLC activator compound
m-3M3FBS [55,57–59]. Nicotinic responses of deep layer pyramidal neurons from ChAT-ChR2 mice
to optogenetic acetylcholine release were recorded in the continuous presence of atropine before
and after treatment with m-3M3FBS (25μM, 5 min; **Fig 4B**). The rising slope of the nicotinic
responses showed a significant increase after m-3M3FBS treatment ($23 \pm 17\%$; Paired Cohen's d
$= 0.83$; $P = 0.008$, Wilcoxon matched-pairs test), compared to the baseline change observed in
the same cells prior to PLC activation ($-6 \pm 4\%$; 8 cells, 6 mice; **Fig 4C, D**). This increase was
not observed with the inactive ortholog o-3M3FBS that does not activate PLC (Paired Cohen's d
$= 0.09$, $P = 0.625$, Wilcoxon matched-pairs test, data not shown). The area under the nicotinic
response known as charge transfer also showed a statistically significant increase following PLC
activation ($22 \pm 7\%$; Cohen's d $= 1.68$; $P = 0.016$, Wilcoxon matched-pairs test; **Fig 4D**),
compared to baseline change ($-9 \pm 6\%$). Thus, PLC activation causes a specific increase in
nicotinic receptor responses, presumably due to cleavage of inhibitory GPI-anchored prototoxins
such as Lynx1.

To test the transcriptomic prediction that cell-type specific differences in lynx modulation
lead to different cholinergic properties, we obtained purified water-soluble recombinant Ly6g6e
protein and examined its effects on optogenetic nicotinic responses in labeled Chrna5+ and
Syt6+ neurons (**Fig 4F**). These experiments were conducted in *Chrna5*-Cre$^{/+}$Ai14$^{/+}$ChAT-
ChR2$^{/+}$ and *Syt6*-Cre$^{/+}$Ai14$^{/+}$ChAT-ChR2$^{/+}$ mice. We hypothesized that the modulation of
Chrna5+ neuronal nicotinic receptors by endogenous Ly6g6e would occlude the effect of
exogenous soluble Ly6g6e, whereas Syt6+ neurons would be altered by exposure to the
exogenous Ly6g6e (Fig 4G). Consistent with this hypothesis, we found that 10 minute
application of soluble Ly6g6e did not significantly alter the amplitude of optogenetically evoked
nicotinic responses in labeled Chrna5+ neurons (Change in peak = -2.1 ± 1.2 pA, $t_{(6)} = 1.79$, $P =$
0.12, paired t-test). However, in labeled Syt6+ neurons lacking endogenous expression of
Ly6g6e, exogenous application of soluble Ly6g6e caused a significant decrease in the amplitude
(Change in peak = -10 ± 1.8 pA, $t_{(8)} = 5.60$, $P < 0.001$, paired t-test; **Fig 4H-I**). The change in
peak and area of the nicotinic responses caused by solube Ly6g6e was significantly different
between Chrna5+ and Syt6+ neurons (change in peak : $t_{(14)} = 3.43$, $P = 0.004$; Change in area:
$t_{(14)} = 2.53$, $P = 0.024$, Unpaired t test; **Fig 4I - J**). Of note, soluble and endogenous GPI-
anchored prototoxins are known to have opposite effects on nicotinic receptors and the exact
direction of endogenous modulation of nicotinic receptors by different lynx proteins is still
debated [41,60,61]. The key outcome of this experiment is the difference in the Ly6g6e modulation

of Chrna5+ and Syt6+ neurons, not the direction. We conclude that Chrna5+ neurons exert
specialized molecular control over their nicotinic receptors, shaping their fate as acetylcholine
super-responders.

Discussion

Our work examines the effects of GPI-anchored lynx prototoxins on native nicotinic receptor-
mediated optogenetic responses, advancing from work in heterologous expression systems.
These results are a first step in showing how endogenous lynx regulation of nicotinic responses
can act in a complex cell-type specific fashion leading to specialized cholinergic properties in a
subset of neurons. Overall, our study reveals a previously uncharacterized population of *Chrna5*-
expressing subplate neurons in the prefrontal cortex that are exquisitely sensitive to
acetylcholine, with differential expression of several lynx prototoxin genes that allow flexible
tuning of their high-affinity nicotinic responses (**Fig 5**).

Specialized cholinergic properties of Chrna5-expressing neurons
While important for attention, the neurophysiological impact of the auxiliary α5 nicotinic subunit
in its native neuronal environment has remained beyond the reach of previous work. The
contributions of α5 to high-affinity nicotinic receptors have been extrapolated based on results of
cell system experiments and work in rodents deleted for *Chrna5* [13,16,22,62,63]. Here, *Chrna5*-Cre
mice allowed us to affirmatively demonstrate that neurons expressing the α5 nicotinic subunit
respond faster and more strongly to endogenous acetylcholine. This cholinergic heterogeneity
among layer 6 neurons prompted a larger scale comparison between Chrna5+ neurons and a
well-defined layer 6 population labeled by *Syt6* [29]. These experiments revealed a subset of
Chrna5+ acetylcholine 'super-responders' with high affinity nicotinic responses that were not
found in Syt6+ neurons.

Heterogeneity of cell types expressing Chrna5
Previously, the deep-layer cell types expressing *Chrna5* were uncharacterized, and generally
thought to include L6CT neurons [8]. Investigation of L6CT neurons have relied on *Syt6*-Cre and
Ntsr1-Cre mouse lines that label similar sets of neurons [27–29], with only *Syt6*-Cre mice
successfully labeling this population in prefrontal cortex [31]. L6CT neurons labeled by *Syt6* or
Ntsr1 expression are excited by acetylcholine [32,64], but the degree to which their nicotinic
response relied on *Chrna5* expression was unclear. Strikingly, we reveal that acetylcholine
super-responders with high affinity nicotinic receptors are from the population of Chrna5+
neurons without *Syt6*-expression. Transcriptomic analysis demonstrates that majority of these
likely Chrna5+ 'super-responders' arise from L5 Near-Projecting and L6b neurons, populations
that express multiple markers linking them to the developmental subplate. These enigmatic cells
are remnants of earliest-born cortical neurons that serve as a relay for establishing connections
between cortex and thalamus [65,66]. Subplate neurons receive cholinergic inputs at birth [67],
highlighting their role in developmental cholinergic modulation.
324
325 *Advantages of Chrna5 as a marker for subplate cells*
326 In contrast to L6CT neurons, subplate neurons remain relatively uncharacterized due to the lack
327 of transgenic mice to definitively label the diverse subtypes, and inaccessibility of the available

lines for *in vivo* targeting. Transcriptomic analysis (Fig 5, Table 1) suggests that the Chrna5+ population is enriched for known subplate markers *Ctgf* (Connective tissue growth factor), *Cplx3* (Complexin 3), *Kcnab1,*and *Lpar1* [35,68,69]. Significantly, the lynx prototoxin and nicotinic receptor modulator *Ly6g6e*, which is highly expressed in Chrna5+ neurons, is also a marker of subplate neurons [70]. Our study is the first to identify enhanced cholinergic activation regulated by *Chrna5* and lynx-gene expression in subplate/L6b neurons. Subplate neurons have recently been found to strongly regulate cortical output through their intracortical connections [71,72]. Enhanced cholinergic activation in these neurons will have different consequences for prefrontal processing, challenging the popular conception that cholinergic modulation of attention occurs only through top-down control of thalamic input by L6CT neurons.

Molecular determinants of nicotinic receptor properties in Chrna5+ neurons

Our transcriptomic analysis revealed enhanced expression of GPI-anchored lynx prototoxin genes *Ly6g6e*, *Lypd1*, and *Lypd6b* in Chrna5+ neurons (Fig 5). Lynx proteins are well known modulators of nicotinic receptor properties and trafficking [41,43], but most of the insight into their actions comes from heterologous cell systems, deletion, and overexpression experiments. Their effects on nicotinic receptors in their native environment is unclear. In expression systems, Ly6g6e potentiates α4β2 nicotinic responses, slowing their desensitization [39], predicting cholinergic responses in Chrna5+ neurons would be resistant to desensitization as has been implied by *Chrna5* deletion work [16]. In contrast, Lynx2 (*Lypd1*) is a predicted negative modulator that can increase desensitization of α4β2 nicotinic receptors [73]. Lynx2 acts intracellularly to reduce surface expression of α4β2 nicotinic receptors [39], but may preferentially act on lower affinity $(\alpha 4)_3(\beta 2)_2$ receptors [62,74] and indirectly promote expression of high affinity $(\alpha 4)_2(\beta 2)_2\alpha 5$ nicotinic receptors. The effect of Lypd6b on $(\alpha 4)_2(\beta 2)_2\alpha 5$ nicotinic receptors is yet to be determined and may further contribute to the complex control of Chrna5+ nicotinic responses [75]. In addition, Lynx1, a well-known negative modulator of α4β2 nicotinic receptors [42,76–78] is also expressed in Chrna5+ neurons. Consistent with such complex lynx regulation, our experiments confirmed that removing GPI-anchored lynx proteins increases nicotinic response onset and amplitude potentially due to removal of Lynx1. In contrast, exogenous application of recombinant Ly6g6e had different effects in Chrna5+ and Syt6+ neurons, consistent with cell-type specific lynx modulation in Chrna5+ neurons predicted by transcriptomics.

Functional consequences

The effects of lynx proteins on nicotinic receptor function have so far been determined by heterologous expression systems [39], knockout studies [73,79], exogenous application of recombinant water-soluble lynx proteins [80,81], and viral manipulation of expression in the brain [42,82]. We advance this field by revealing, in native tissue, complex endogenous regulation of optogenetic nicotinic responses by multiple GPI-anchored lynx proteins. Inhibitory lynx expression and high levels of acetylcholinesterase in Chrna5+ neurons suggest that their responses are restrained and our experiments likely underestimated their nicotinic receptor function. These responses could be dramatically enhanced when acetylcholinesterase and inhibitory lynx modulation is reduced through other signaling mechanisms. Such flexible tuning of nicotinic responses by lynx prototoxins in Chrna5+ neurons can provide greater dynamic range and poises them to be key

players during attentional processing (Fig 5). A recent study found that preventing developmental increase in Lynx1 expression in corticocortical neurons by viral knockdown led to altered cortical connectivity and impaired attention [42]. Thus cell-type specific changes in lynx expression during development are critical for maturation of attention circuits. It is of interest to examine such changes during development in Chrna5+ neurons and how they differ from Syt6+ neurons.

Caveats and limitations of study

Subplate marker genes and localization of subplate cells can vary across cortical regions, and they have been better studied in sensory cortical regions [83–85]. Our optophysiological, anatomical and transcriptomic data show Chrna5+ acetylcholine super-responder neurons to be found in layer 6b both in the primary somatosensory cortex and PFC (Fig S2-S3). However, an outstanding question is whether transcriptomic cell classes such as L5/6NP and L6b, which express multiple subplate marker genes, correlate with early developmental subplate origin. Anatomical evidence alone is insufficient for this purpose. For example, while the distinct transcriptomic subclasses L6b and L5/6NP show overlapping spatial distribution in deep-layers of the motor cortex [86,87], these subclasses possess distinct epigenetic signatures [88] whose implications are yet to be understood. It will be necessary for future work to examine whether transcriptomic /epigenetically defined cell classes (L5/6NP, L6b) expressing subplate markers also show distinct early developmental origin. Lack of labeling strategies has limited efforts to characterize these neuronal subclasses and verify their developmental subplate identity. *Chrna5-Cre* mice will be an important tool to rectify this as they consistently label these conserved subclasses across multiple cortical regions. Another caveat of our study is the perturbation of lynx prototoxins using chemical strategies. While we have demonstrated cell-type specific effects of recombinant ly6g6e, it remains to be seen whether cell-type specific expression of ly6g6e or other lynx prototoxins is necessary for nicotinic modulation. Future work is needed to genetically manipulate lynx prototoxins in specified cell populations.

Summary of advances

Our study reveals a distinct group of 'acetylcholine super-responder' neurons in the prefrontal cortex identified by *Chrna5*-expression that constitute subplate neurons vital for cortical development. We identify that their high affinity α5 subunit-containing nicotinic receptors are under complex regulation by several lynx prototoxins and acetylcholinesterase. *Chrna5*-Cre mice are a valuable tool for future studies examining the in vivo role of these specialized neurons.

Acknowledgements: We acknowledge the generous support of the Canadian Institutes of Health Research (CIHR MOP 89825, EKL; CIHR PJT-153101, EKL), the Canada Research Chair in Developmental Cortical Physiology (EKL), Ontario Graduate Scholarships (SV). We thank Ms. Janice McNabb and Mr. Ha-Seul Jeoung for expert technical assistance. Earlier presentations of this work received valuable feedback from Dr. Junchul Kim, Dr. Beverly Orser, and Dr. Steve Prescott of the University of Toronto.

Conflicts of interest: None

Figure legends

Figure 1. *Chrna5* expression identifies a distinct population of prefrontal neurons with
stronger and faster-onset optogenetic cholinergic responses. A, Breeding scheme to obtain
triple transgenic *Chrna5*-Cre$^{/+}$Ai14$^{/+}$ChAT-ChR2$^{/+}$ mice expressing tdTomato in *Chrna5*-
expressing (Chrna5+) neurons and Channelrhodopsin 2 in cholinergic axons. **B,** (*Top*) Schematic
of coronal mPFC slice with region of interest, adapted from [89]. (*Bottom*) two photon imaging (3D
projection) of tdTomato-labeled Chrna5+ neurons and EYFP-labeled cholinergic axons in layer 6
of mPFC slices. **C,** IRDIC (*left*) and widefield fluorescence (TRITC, *right*) images of tdTomato
labeled Chrna5+ and unlabeled layer 6 neurons during whole cell patch clamp electrophysiology.
Clearing induced by the pipette is visible. **D,** Average light-evoked endogenous cholinergic
response of labeled Chrna5+ vs neighbouring unlabeled Chrna5- neurons. Dotted lines are the
slope of the response onset. (*Inset*) Individual responses are zoomed in to show the onset. **E-F,**
Bar graph comparing (E) Rising slope and (F) Peak current of endogenous cholinergic responses
between labeled Chrna5+ neurons (n = 24 cells) and unlabeled Chrna5- neurons (n = 15 cells, 4
mice). *$P < 0.05$, Unpaired t-test. **G,** (*Top*) Breeding scheme to obtain triple transgenic *Chrna5*-
Cre$^{/+}$Ai14$^{/+}$*Syt6*-EGFP$^{/+}$ mice expressing tdTomato in Chrna5+ neurons and EGFP in Syt6+
neurons. (*Bottom*) Confocal imaging in mPFC slices shows Chrna5+ and Syt6+ neurons
distributed in layer 6. **H-I,** Confocal (H) and two-photon imaging (I) reveal three populations of
neurons: exclusively Chrna5+ neurons which do not express *Syt6*, overlapping Chrna5+Syt6+
neurons which express both markers, and exclusively Syt6+ neurons which do not express
Chrna5. **J,** (*left*) Graph quantifies the percentage of each cell type with respect to all labeled cells
per sample. (*Right*) Average proportions of Chrna5+, Chrna5+Syt6+ and Syt6+ neurons. (See
Figs S1-3 for additional information on distribution of Chrna5+ cells across the cortex, and
additional recordings showing stronger and faster opto-cholinergic responses in Chrna5+ neurons
in primary somatosensory cortex.)

Figure 2. Calcium imaging in Chrna5+ and Syt6+ populations reveals a distinct subset of
Chrna5+ neurons with resilient nicotinic responses. A, (*Top*) Two photon Ca^{2+} imaging in
prefrontal brain slices from *Chrna5*-Cre$^{/+}$Ai96$^{/+}$ (*left*) and *Syt6*-Cre$^{/+}$Ai96$^{/+}$ mice (*right*) showing
acetylcholine-evoked GCaMP6s responses in Chrna5+ and Syt6+ neurons respectively (scale 50
μm). (*Bottom*) Acetylcholine-evoked GCaMP6s signals were sequentially recorded after
application of competitive nicotinic antagonist DHBE and addition of muscarinic antagonist
atropine (scale 10 μm). **B,** Normalized fluorescence signal **(**ΔF by F) evoked by acetylcholine in
individual Chrna5+ and Syt6+ neurons in a brain slice at baseline (*left*), after DHBE (*middle*),
and after DHBE + Atropine (*right*). (Inset, average response and standard deviation. Scale: same
as main figure). **C-D i,** Boxplot shows the percentage of response remaining after the application
of (C) DHBE and (D) DHBE + Atropine (Inset shows the same boxplot with a restricted y-axis,
'+' symbols denote respective means). Responses were quantified by the area under the ΔF/F

curve (n = 71 neurons, 6 mice for Chrna5+, and 112 neurons, 7 mice for Syt6+, ****$P < 10^{-4}$,
Mann-Whitney test). **C-D ii,** Cumulative frequency distribution of the percentage response
remaining after (C) DHBE and (D) DHBE + Atropine ($P < 10^{-4}$, Kolmogorov-Smirnov test). **C-**
D iii, Proportion of cells showing zero and non-zero responses after (C) DHBE and (D) DHBE +
Atropine ($P < 10^{-4}$ for both C & D, Fisher's exact test). **E,** Current clamp responses evoked by
1mM acetylcholine (15s) in fluorescently labeled Chrna5+ and Syt6+ layer 6 neurons patched in
mPFC slices from mice *Chrna5*-Cre$^{/+}$ Ai14$^{/+}$ and *Syt6*-Cre$^{/+}$Ai14$^{/+}$ or *Syt6*-EGFP mice
respectively. **F,** Peak spike frequency in Chrna5+ and Syt6+ neurons evoked by acetylcholine
(*left*), and in the presence of competitive nicotinic antagonist DHBE and atropine (*middle; **$P <$*
0.01,$P < 0.05$*, unpaired t-test). Residual response remaining after DHBE + Atropine in
Chrna5+ neurons is blocked by non-competitive nicotinic antagonist mecamylamine (*right, **P*
< 0.01, paired t-test). (See **Fig S4** for localization of Chrna5+ neurons in PFC with resilient
nicotinic responses in layer 6b.)

Figure 3. Single cell transcriptomic analysis reveals Chrna5+ subset to span subplate
neuron populations with differential expression of lynx prototoxin genes A, Single cell-
RNAseq data for 2422 L5-6 glutamatergic neurons in the anterior cingulate cortex (ACA, shown
in schematic on the left) was obtained from publicly available datasets (Allen Institute,
SMARTSeq ACA and MOP (2018)). (*Right*) Scatter plot showing *Chrna5* vs *Syt6* expression in
$\log_{10}$ (Copies per million +1) for each neuron, with the frequency distribution shown on the
corresponding axes. Neurons were classified into Chrna5+, Chrna5+Syt6+ and Syt6+ groups
based on their expression of *Chrna5* and *Syt6* genes. Cells which expressed neither gene were
excluded from subsequent analyses. **B,** The major neuronal subclasses within Chrna5+,
Chrna5+Syt6+ and Syt6+ groups is indicated by the colorbar on top. NP- Near projecting, CT-
Corticothalamic. Heatmap shows expression of subplate and corticothalamic marker genes in
each cell in all 3 groups. **C,** Dotplot shows summary of subplate and corticothalamic marker
expression in Chrna5+, Chrna5+Syt6+ and Syt6+ groups. Dot size indicates the percentage of
cells within each group expressing that gene, color of the dot indicates average expression level
relative to other groups. Chrna5+ neurons highly express multiple subplate marker genes, but not
corticothalamic markers. **D,** Violin plots show expression of Lynx prototoxin genes *Ly6g6e*,
Lypd1 (Lynx2) and *Lypd6b* which show highest fold-change between Chrna5+ and
Chrna5+Syt6+ neurons. **E,** Dotplot shows expression of major genes known to modulate
cholinergic function, including nicotinic, muscarinic subunits, acetylcholinesterase, and lynx
prototoxins in Chrna5+, Chrna5+Syt6+ and Syt6+ neurons. Genes are ordered by decreasing fold
change in expression. Dot size indicates the percentage of cells within each group expressing that
gene, color of the dot indicates average expression level relative to other groups. Fold change of
all the genes shown in this dotplot are listed in Table S3. (See **Fig S5** for transcriptomic analysis
of Chrna5+ neurons in other cortical areas showing conserved subplate markers and lynx
prototoxin gene expression across cortical regions)

Figure 4. Regulation of optogenetic nicotinic responses by endogenous GPI-anchored lynx
proteins and cell type specific effects of recombinant Ly6g6e. A, Schematic of nicotinic
receptor environment showing endogenous GPI-anchored lynx proteins exerting positive and
negative modulation of nicotinic receptors. The compound m-3M3FBS activates PLC, cleaving

the GPI anchor and perturbing lynx-mediated modulation of nicotinic responses. **B,** Optogenetic nicotinic responses in prefrontal deep-layer pyramidal neurons from ChAT-ChR2 mice before and after treatment with m-3M3FBS (5 min). **C,** PLC activation significantly increased the rising slope of optogenetic nicotinic responses. Percent change in **D,** Rising slope and **E,** Area of nicotinic response in control and after PLC activation. (*$P < 0.05$, **$P < 0.01$, Wilcoxon matched-pairs test). **F,** IRDIC (*left*) and widefield fluorescence (*right*) images of tdTomato labeled Chrna5+ (*top*) and Syt6+ (*bottom*) neurons during whole-cell patch clamp electrophysiology in *Chrna5*-Cre$^{/+}$Ai14$^{/+}$ChAT-ChR2$^{/+}$ and *Syt6*-Cre$^{/+}$Ai14$^{/+}$ChAT-ChR2$^{/+}$ mouse brain slices respectively. **G,** Schematic summarising predicted and observed effects of recombinant water-soluble ly6g6e on Chrna5+ and Syt6+ neuronal nicotinic receptors. **G,** Optogenetic nicotinic responses are reduced in amplitude following 10 minute application of soluble ly6g6e in Syt6+ but not Chrna5+ neurons. Change in peak current (**I**) and area of the nicotinic response (**J**) of Chrna5+ vs Syt6+ neurons (*$P < 0.05$, ** $P < 0.01$, Unpaired t-test)

Figure 5. Graphical summary. Deep-layer pyramidal neurons can be divided into three groups (Chrna5+, Chrna5+Syt6+, Syt6+) by their expression of *Chrna5* and *Syt6* genes. The subset of *Chrna5*-expressing neurons without *Syt6* expression are molecularly distinct and comprise of subplate neurons, whereas *Syt6*-expressing neurons are of the corticothalamic subtype. Nicotinic receptors in these neurons are under complex regulation by endogenous lynx prototoxins. Inhibitory prototoxin gene *Lynx1* is expressed uniformly in all neurons, whereas Chrna5+ subplate neurons additionally have specific expression of *Ly6g6e*, *Lypd1* and *Lypd6b* prototoxin genes. These Chrna5+ subplate neurons in layer 6b show enhanced α5 subunit nicotinic receptor-mediated cholinergic responses that are differently modulated by specific lynx prototoxins.

 Tables

Genes	P value	Chrna5+ cells %	Chrna5+Syt6+ cells %	Adjusted P value	Fold change (Chrna5+/ Chrna5+Syt6)
Cplx3	1.16E-43	0.535	0.149	3.37E-39	5.69
Ctgf	1.60E-28	0.77	0.568	4.65E-24	3.81
Ptn	3.68E-28	0.864	0.758	1.07E-23	3.60
Tmem163	2.12E-44	0.51	0.129	6.16E-40	2.58
Lypd1	3.40E-23	0.593	0.381	9.88E-19	2.55
Kcnab1	3.81E-74	0.741	0.191	1.11E-69	2.51
Serpini1	1.54E-50	1	0.993	4.49E-46	2.50
Hpcal1	2.48E-27	0.658	0.39	7.20E-23	2.47
Nrsn2	1.89E-30	0.864	0.767	5.50E-26	2.28
Ntm	1.83E-56	0.889	0.59	5.31E-52	2.21
Tshz2	2.82E-16	0.535	0.327	8.20E-12	2.17
Etv1	5.70E-16	0.395	0.186	1.66E-11	2.16
Rcn2	5.18E-46	0.996	0.984	1.51E-41	2.14
Olfm3	1.67E-44	0.897	0.716	4.86E-40	2.11
Crtac1	5.50E-46	0.728	0.347	1.60E-41	2.04
Ly6g6e	2.13E-25	0.671	0.434	6.20E-21	2.03
Adk	6.31E-29	0.922	0.922	1.83E-24	2.00
Trp53i11	2.73E-60	0.724	0.222	7.94E-56	1.93
Cd164	6.72E-18	0.897	0.882	1.95E-13	1.92
Dner	2.35E-45	0.712	0.326	6.82E-41	1.85

513 **Table 1. Top 20 differentially expressed genes between Chrna5+ and Chrna5+Syt6+**
514 **neurons.** Top 20 differentially expressed genes determined by the FindMarkers function on
515 Seurat ordered by decreasing fold change. Several known subplate neuron markers (highlighted
516 by *****) are the highest enriched genes in Chrna5+ neurons. All differentially expressed genes are
517 listed in supplementary table S4.

STAR Methods

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to Dr. Evelyn Lambe (evelyn.lambe@utoronto.ca) or Dr. Eric Turner (eeturner@uw.edu).

Materials availability

This study did not generate new unique reagents.

Data and code availability

This paper analyzes existing, publicly available single cell RNAseq data. The links for the Allen Institute scRNAseq database are provided in the key resources table. The paper does not report original code. Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Experimental model and subject details

Syt6-Cre$^{/+}$GCaMP6s$^{/+}$ and *Chrna5*-Cre$^{/+}$GCaMP6s$^{/+}$ mice used for Ca^{2+} imaging were obtained by crossing *Chrna5*-Cre (Gift from Dr. Eric Turner) and *Syt6*-Cre mice (*Syt6*-Cre KI148, RRID:MMRRC 037416-UCD, [90]) respectively, with Ai96 mice (JAX: 024106). For electrophysiological recordings of labeled Chrna5+ and Syt6+ neurons, we used *Chrna5*-Cre$^{/+}$Ai14$^{/+}$, and *Syt6*-Cre$^{/+}$Ai14$^{/+}$ mice respectively. *Syt6*-EGFP$^{/+}$ mice were additionally used for few experiments (*Syt6*-EGFP EL71, RRID:MMRRC 010557-UCD, [91]).

Triple transgenic mice labeling both *Chrna5* and *Syt6*-expressing neurons with EGFP in Syt6+ neurons and tdTomato in Chrna5+ neurons were used to examine the overlap between the two cell types. *Syt6*-EGFP and Ai14 mice [92] were bred together and the offspring were crossed with *Chrna5*-Cre mice to generate *Chrna5*-Cre$^{/+}$Ai14$^{/+}$*Syt6*-EGFP$^{/+}$ mice used for these experiments. A set of experiments measuring optogenetic cholinergic responses was also performed in ChAT-ChR2 (ChAT$^{/+}$) mice (JAX: 014546). To examine optogenetic cholinergic responses in labeled Chrna5 and Syt6 cell populations, the respective Cre lines were crossed with ChAT$^{/+}$Ai14$^{/+}$ mice to generate *Chrna5*-Cre$^{/+}$Ai14$^{/+}$ChAT$^{/+}$ and *Syt6*-Cre$^{/+}$Ai14$^{/+}$ChAT$^{/+}$ mice.

All animals were bred on a C57BL/6 background, except *Syt6*-EGFP which were Black Swiss. Adult male and female animals age >P60 were used in the study. Mice were separated based on sex after weaning at P21 and group-housed (2-4 mice per cage). Animals had ad libitum access to food and water and were on a 12-h light/dark cycle with lights on at 7 AM. Guidelines of the Canadian Council on Animal Care were followed, and all experimental procedures were approved by the Faculty of Medicine Animal Care Committee at the University of Toronto. 42 mice were used for the entire study, with similar numbers of males and females.

Method details

Brain slicing and electrophysiology

Slicing and electrophysiology followed procedures described previously [16]. An intraperitoneal injection of chloral hydrate (400 mg/kg) was given to anesthetize mice prior to decapitation. The brain was rapidly extracted in ice cold sucrose ACSF (254 mM sucrose, 10 mM D-glucose, 26 mM $NaHCO_3$, 2 mM $CaCl_2$, 2 mM $MgSO_4$, 3 mM KCl and 1.25 mM NaH_2PO_4). 400 μm thick coronal slices of prefrontal cortex (Bregma 2.2 - 1.1) were obtained on a Dosaka linear slicer (SciMedia, Costa Mesa, CA, USA). Slices were left to recover for at least 2 hours in oxygenated (95% O_2, 5% CO_2) ACSF (128 mM NaCl, 10 mM D-glucose, 26 Mm $NaHCO_3$, 2 mM $CaCl_2$, 2 mM $MgSO_4$, 3 Mm KCl, and 1.25 mM NaH_2PO_4) at 30°C before being used for electrophysiology or two-photon Ca^{2+} imaging. Brain slices were transferred to the stage of a BX51WI microscope (Olympus, Tokyo, Japan) and perfused with oxygenated ACSF at 30°C. Recording electrodes (2 - 4 MΩ) containing patch solution (120 mM potassium gluconate, 5 mM KCl, 10 mM HEPES, 2 mM $MgCl_2$, 4 mM K_2-ATP, 0.4 mM Na_2-GTP and 10 mM sodium phosphocreatine, pH adjusted to 7.3 using KOH) were used to patch pyramidal neurons in layer 6 - 6b based on morphology and proximity to white matter. Only regular spiking neurons were included. Multiclamp 700B amplifier at 20 kHz with Digidata 1440A and pClamp 10.7 software (Molecular devices) were used for data acquisition. All recordings were compensated for liquid junction potential (14 mV). Voltage-clamp responses were examined at -75 mV and in current-clamp at rest or starting from -70 mV.

Optogenetics

5 ms pulses of blue light (473 nm) were delivered through the 60X objective lens with an LED (Thorlabs, 2 mW) to excite channelrhodopsin containing cholinergic fibers. Pattern of stimulation was as in a previous study, with 8 pulses of light delivered in a frequency accommodating manner [16].

Pharmacology

Acetylcholine (1mM, Sigma) was used to exogenously stimulate cholinergic receptors. Atropine (200 nM, Sigma) and Dihydro-β-erythroidine (DHBE, 10 μM, Tocris) were used to competitively block muscarinic receptors and β2 subunit-containing nicotinic receptors respectively. Mecamylamine (5 μM, Tocris) was used to further non-competitively block nicotinic receptors. Phospholipase C activator m-3M3FBS (25 μM, Tocris) was used to cleave GPI-anchored Lynx prototoxins and the inactive ortholog o-3M3FBS (25 μM, Tocris) was used as a control [55]. Cyclodextrin (1 mM, Tocris) was included in a small subset of experiments to improve solubility of 3M3FBS compounds, but no further improvement in efficacy was observed. Water soluble recombinant Ly6g6e (0.5 mg/ml) was obtained by custom purification (Creative Biomart) and used for exogenous application at 1: 1000 and 3:1000 dilution. Effects on nicotinic receptors were not distinguishable between the two different protein concentrations. Only freshly thawed protein aliquots were used for experiments.

Two-photon imaging

Two-photon imaging of GCaMP6s Ca^{2+} signals in L6 neurons was performed using a 60× water-immersion objective with 0.90 numerical aperture using an Olympus Fluoview FV1000

microscope and a Titanium-Sapphire laser sapphire laser (Newport) at 930nm. Images were
sampled at 512 x 512 pixels (2.4 pixels/µm) at a frame rate of 0.9 Hz. Following a 2-minute
washout period for this initial application, GCaMP6s Ca^{2+} signals were measured in response to
acetylcholine (1 mM, 15 s). The cellular responses to acetylcholine were measured at baseline,
then after application of competitive nicotinic receptor antagonist DHBE (10 µM, 10 min), and
again after the addition of muscarinic antagonist atropine (200 nM, 10 min).

Dual color two-photon imaging (910 nm excitation, using 570 nm dichroic mirror with
green (540-595 nm) and red (570-620 nm) filters) was performed in brain slices of triple
transgenic *Chrna5*-Cre$^{/+}$Ai14$^{/+}$*Syt6*-EGFP$^{/+}$ mice to examine overlap in fluorescent reporter
expression between Chrna5+ and Syt6+ neurons. Z-stacks of 30 frames acquired in 1-µm steps
were taken in layer 6 of mPFC slices and the maximum projection used to count cells with the
cell counter feature in Fiji. A set of mPFC brain slices from *Chrna5*-Cre$^{/+}$Ai14$^{/+}$*Syt6*-EGFP$^{/+}$
mice were also fixed and mounted for confocal imaging with LSM880 (Leica) microscope.

Single cell RNAseq analysis
Single cell RNAseq data for Anterior Cingulate Cortex (ACA) of adult mice was taken from the
ACA and MOP Smart-Seq (2018) database, with cell-type annotations from Whole cortex &
Hippocampus Smart-Seq (2019) database from the Allen Institute for Brain Science at
https://portal.brain-map.org/atlases-and-data/rnaseq [48,50]. Single cell analysis was performed
using the R package Seurat (v 4.04) [93–95] . Layer 5 and 6 glutamatergic neurons were selected
and sorted into three cell classes based on their expression of *Chrna5* and *Syt6* genes: those
expressing only *Chrna5* (Chrna5+, n = 243), only *Syt6* (Syt6+, n = 834), or both *Chrna5* and
Syt6 (Chrna5+Syt6+, n = 564). Expression (copies per million) greater than zero was used as the
threshold. 781 cells did not express either *Chrna5* or *Syt6* and were not used in subsequent
analyses. Rare cell types with fewer than 10 cells per group are not shown in the heatmap in
figure 5 but are included for the differential expression analysis. The FindMarkers function in
Seurat was used to identify genes differentially expressed between the Chrna5+ and
Chrna5+Syt6+ populations. Adjusted p value < 0.05 was used as the cutoff for identifying
differentialy expressed genes. Similar approaches were used to perform the same analyses in
multiple cortical regions (Fig S5). L5-6 glutamatergic cells cells in ACA, MOp, SSp, and VISp
were taken from the Whole cortex & Hippocampus Smart-Seq (2019) dataset.

Quantification and Statistical Analysis
Analysis of electrophysiological data was conducted on Clampfit 10.7 and Axograph. Rising
slope was measured by fitting a line to the first 50ms of the optogenetic cholinergic responses.
Cholinergic response magnitude in voltage clamp was determined by peak current (picoamperes)
and charge transfer (picocoulombs) measured by the area under the current response for 1
second.

GCaMP6s imaging data were extracted using the multi-measure feature in Fiji. Maximum
projections across time for each experiment were first used to identify acetylcholine-responsive
cells and add them to the ROI manager, then fluorescence intensity at all timepoints for each cell
was measured. Fluorescence was normalized to the background fluorescence averaged over the
first 10 frames. Area under the peak (AUP) of the signal after baseline correction was used to

quantify the magnitude of cells' response to acetylcholine. Percentage response remaining after DHBE and DHBE+Atropine was calculated from the cell's AUP before and after the blockers. The statistical analysis for electrophysiological recordings and calcium imaging treated each cell as independent with several cells recorded from each slice.

GraphPad Prism 8 was used for statistical analysis and plotting graphs. Bar graphs depicting mean with standard error and boxplots with median and quartiles are shown. Effect sizes are reported as Cohen's d for major results [96]. Unpaired t-tests or Mann-Whitney tests were used when comparing response properties between cell types, and paired t-test or Wilcoxon test to quantify effect of pharmacological manipulations within cells. Kolmogorov-Smirnov and Fisher's exact tests were used to compare cumulative distributions and proportions of cells respectively.

References

1. Dalley, J.W., Cardinal, R.N., and Robbins, T.W. (2004). Prefrontal executive and cognitive
functions in rodents: neural and neurochemical substrates. Neurosci Biobehav Rev *28*, 771–784.
10.1016/j.neubiorev.2004.09.006.

2. McGaughy, J., Dalley, J.W., Morrison, C.H., Everitt, B.J., Robbins, T.W., and Robbins, TW.
(2002). Selective behavioral and neurochemical effects of cholinergic lesions produced by
intrabasalis infusions of 192 IgG-saporin on attentional performance in a five-choice serial
reaction time task. J Neurosci *22*, 1905–1913. 10.1523/jneurosci.5112-07.2008.

3. Dalley, J.W., Theobald, D.E., Bouger, P., Chudasama, Y., Cardinal, R.N., and Robbins, T.W.
(2004). Cortical Cholinergic Function and Deficits in Visual Attentional Performance in Rats
Following 192 IgG-Saporin-induced Lesions of the Medial Prefrontal Cortex. Cerebral Cortex *14*,
922–932. 10.1093/cercor/bhh052.

4. Gritton, H.J., Howe, W.M., Mallory, C.S., Hetrick, V.L., Berke, J.D., and Sarter, M. (2016).
Cortical cholinergic signaling controls the detection of cues. Proc Natl Acad Sci U S A *113*,
E1089-97. 10.1073/pnas.1516134113.

5. Briggs, F., and Usrey, W.M. (2008). Emerging views of corticothalamic function. Curr Opin
Neurobiol *18*, 403–407. 10.1016/j.conb.2008.09.002.

6. Voigts, J., Deister, C.A., and Moore, C.I. (2020). Layer 6 ensembles can selectively regulate the
behavioral impact and layer-specific representation of sensory deviants. Elife *9*.
10.7554/eLife.48957.

7. Spellman, T., Svei, M., Kaminsky, J., Manzano-Nieves, G., and Liston, C. (2021). Prefrontal deep
projection neurons enable cognitive flexibility via persistent feedback monitoring. Cell *184*, 2750-
2766.e17. 10.1016/J.CELL.2021.03.047.

8. Kassam, S.M., Herman, P.M., Goodfellow, N.M., Alves, N.C., and Lambe, E.K. (2008).
Developmental excitation of corticothalamic neurons by nicotinic acetylcholine receptors. J
Neurosci *28*, 8756–8764. 10.1523/JNEUROSCI.2645-08.2008.

9. Sparks, D.W., Tian, M.K., Sargin, D., Venkatesan, S., Intson, K., and Lambe, E.K. (2018).
Opposing cholinergic and serotonergic modulation of layer 6 in prefrontal cortex. Front Neural
Circuits *11*. 10.3389/fncir.2017.00107.

10. Venkatesan, S., Jeoung, H.-S., Chen, T., Power, S.K., Liu, Y., and Lambe, E.K. (2020).
Endogenous Acetylcholine and Its Modulation of Cortical Microcircuits to Enhance Cognition. In
Behavioral Pharmacology of the Cholinergic System (Springer, Berlin, Heidelberg), pp. 47–69.
10.1007/7854_2020_138.

11. Winzer-Serhan, U.H., and Leslie, F.M. (2005). Expression of alpha5 nicotinic acetylcholine
receptor subunit mRNA during hippocampal and cortical development. J Comp Neurol *481*, 19–
30. 10.1002/cne.20357.

12. Wada, E., McKinnon, D., Heinemann, S., Patrick, J., and Swanson, L.W. (1990). The distribution
of mRNA encoded by a new member of the neuronal nicotinic acetylcholine receptor gene family
(α5) in the rat central nervous system. Brain Res *526*, 45–53. 10.1016/0006-8993(90)90248-A.

13. Bailey, C.D.C., De Biasi, M., Fletcher, P.J., and Lambe, E.K. (2010). The nicotinic acetylcholine
receptor alpha5 subunit plays a key role in attention circuitry and accuracy. J Neurosci *30*, 9241–
9252. 10.1523/JNEUROSCI.2258-10.2010.

14. Tian, M.K., Bailey, C.D.C., De Biasi, M., Picciotto, M.R., and Lambe, E.K. (2011). Plasticity of
prefrontal attention circuitry: upregulated muscarinic excitability in response to decreased
nicotinic signaling following deletion of α5 or β2 subunits. J Neurosci *31*, 16458–16463.
10.1523/JNEUROSCI.3600-11.2011.

15. Howe, W.M., Brooks, J.L., Tierney, P.L., Pang, J., Rossi, A., Young, D., Dlugolenski, K.,
Guillmette, E., Roy, M., Hales, K., et al. (2018). α5 nAChR modulation of the prefrontal cortex
makes attention resilient. Brain Struct Funct *223*, 1035–1047. 10.1007/s00429-017-1601-1.

16. Venkatesan, S., and Lambe, E.K. (2020). Chrna5 is essential for a rapid and protected response to
optogenetic release of endogenous acetylcholine in prefrontal cortex. The Journal of Neuroscience
40, 7255–7268. 10.1523/jneurosci.1128-20.2020.

17. Han, W., Zhang, T., Ni, T., Zhu, L., Liu, D., Chen, G., Lin, H., Chen, T., and Guan, F. (2019).
Relationship of common variants in CHRNA5 with early-onset schizophrenia and executive
function. Schizophr Res *206*, 407–412. 10.1016/j.schres.2018.10.011.

18. Bierut, L.J., Stitzel, J.A., Wang, J.C., Hinrichs, A.L., Grucza, R.A., Xuei, X., Saccone, N.L.,
Saccone, S.F., Bertelsen, S., Fox, L., et al. (2008). Variants in the Nicotinic Receptors Alter the
Risk for Nicotine Dependence. Am J Psychiatry *165*, 1163. 10.1176/APPI.AJP.2008.07111711.

19. Schuch, J.B., Polina, E.R., Rovaris, D.L., Kappel, D.B., Mota, N.R., Cupertino, R.B., Silva, K.L.,
Guimarães-da-Silva, P.O., Karam, R.G., Salgado, C.A.I., et al. (2016). Pleiotropic effects of
Chr15q25 nicotinic gene cluster and the relationship between smoking, cognition and ADHD. J
Psychiatr Res *80*, 73–78. 10.1016/J.JPSYCHIRES.2016.06.002.

20. Maskos, U. (2020). The nicotinic receptor alpha5 coding polymorphism rs16969968 as a major
target in disease: Functional dissection and remaining challenges. J Neurochem *154*, 241–250.
10.1111/JNC.14989.

21. Scholze, P., and Huck, S. (2020). The α5 Nicotinic Acetylcholine Receptor Subunit Differentially
Modulates α4β2* and α3β4* Receptors. Front Synaptic Neurosci *12*, 607959.
10.3389/fnsyn.2020.607959.

22. Prevost, M.S., Bouchenaki, H., Barilone, N., Gielen, M., and Corringer, P.J. (2020). Concatemers
to re-investigate the role of α5 in α4β2 nicotinic receptors. Cellular and Molecular Life Sciences
2020 78:3 *78*, 1051–1064. 10.1007/S00018-020-03558-Z.

23. Thomson, A.M. (2010). Neocortical layer 6, a review. Front Neuroanat *4*, 13.
10.3389/fnana.2010.00013.

24. Briggs, F. (2010). Organizing principles of cortical layer 6. Front Neural Circuits *4*, 3.
10.3389/neuro.04.003.2010.

25.   Sorensen, S.A., Bernard, A., Menon, V., Royall, J.J., Glattfelder, K.J., Desta, T., Hirokawa, K.,
721   Mortrud, M., Miller, J.A., Zeng, H., et al. (2015). Correlated Gene Expression and Target
722   Specificity Demonstrate Excitatory Projection Neuron Diversity. Cerebral Cortex *25*, 433–449.
723   10.1093/cercor/bht243.

724   26.   Heath, C.J., King, S.L., Gotti, C., Marks, M.J., and Picciotto, M.R. (2010). Cortico-Thalamic
725   Connectivity is Vulnerable to Nicotine Exposure During Early Postnatal Development through
726   α4/β2/α5 Nicotinic Acetylcholine Receptors. Neuropsychopharmacology *35*, 2324–2338.
727   10.1038/npp.2010.130.

728   27.   Harris, J.A., Hirokawa, K.E., Sorensen, S.A., Gu, H., Mills, M., Ng, L.L., Bohn, P., Mortrud, M.,
729   Ouellette, B., Kidney, J., et al. (2014). Anatomical characterization of Cre driver mice for neural
730   circuit mapping and manipulation. Front Neural Circuits *8*, 76. 10.3389/fncir.2014.00076.

731   28.   Harris, J.A., Mihalas, S., Hirokawa, K.E., Whitesell, J.D., Choi, H., Bernard, A., Bohn, P.,
732   Caldejon, S., Casal, L., Cho, A., et al. (2019). Hierarchical organization of cortical and thalamic
733   connectivity. Nature *575*, 195–202. 10.1038/s41586-019-1716-z.

734   29.   Nectow, A.R., Moya, M. V., Ekstrand, M.I., Mousa, A., McGuire, K.L., Sferrazza, C.E., Field,
735   B.C., Rabinowitz, G.S., Sawicka, K., Liang, Y., et al. (2017). Rapid Molecular Profiling of
736   Defined Cell Types Using Viral TRAP. Cell Rep *19*, 655–667. 10.1016/j.celrep.2017.03.048.

737   30.   Nakayama, H., Ibañez-Tallon, I., and Heintz, N. (2018). Cell-type-specific contributions of medial
738   prefrontal neurons to flexible behaviors. Journal of Neuroscience *38*, 4490–4504.
739   10.1523/JNEUROSCI.3537-17.2018.

740   31.   Vaasjo, L.O., Han, X., Thurmon, A.N., Tiemroth, A.S., Berndt, H., Korn, M., Figueroa, A., Reyes,
741   R., Feliciano□Ramos, P.A., and Galazo, M.J. (2022). Characterization and manipulation of
742   Corticothalamic neurons in associative cortices using Syt6□Cre transgenic mice. Journal of
743   Comparative Neurology *530*, 1020–1048. 10.1002/cne.25256.

744   32.   Tian, M.K., Bailey, C.D.C., and Lambe, E.K. (2014). Cholinergic excitation in mouse primary vs.
745   associative cortex: region-specific magnitude and receptor balance. Eur J Neurosci *40*, 2608–2618.
746   10.1111/ejn.12622.

747   33.   Ables, J.L., Görlich, A., Antolin-Fontes, B., Wang, C., Lipford, S.M., Riad, M.H., Ren, J., Hu, F.,
748   Luo, M., Kenny, P.J., et al. (2017). Retrograde inhibition by a specific subset of interpeduncular
749   α5 nicotinic neurons regulates nicotine preference. Proc Natl Acad Sci U S A *114*, 13012–13017.
750   10.1073/pnas.1717506114.

751   34.   Morton, G., Nasirova, N., Sparks, D.W., Brodsky, M., Sivakumaran, S., Lambe, E.K., and Turner,
752   E.E. (2018). Chrna5-expressing neurons in the interpeduncular nucleus mediate aversion primed
753   by prior stimulation or nicotine exposure. J Neurosci, 0023–18. 10.1523/JNEUROSCI.0023-
754   18.2018.

755   35.   Hoerder-Suabedissen, A., and Molnár, Z. (2013). Molecular Diversity of Early-Born Subplate
756   Neurons. Cerebral Cortex *23*, 1473–1483. 10.1093/CERCOR/BHS137.

757   36.   Luhmann, H.J., Kirischuk, S., and Kilb, W. (2018). The superior function of the subplate in early
758   neocortical development. Front Neuroanat *12*, 97. 10.3389/FNANA.2018.00097/BIBTEX.

759   37.   Kanold, P.O., and Luhmann, H.J. (2010). The Subplate and Early Cortical Circuits. Annu Rev
760   Neurosci *33*, 23–48. 10.1146/ANNUREV-NEURO-060909-153244.

761   38.   Miwa, J.M., Anderson, K.R., and Hoffman, K.M. (2019). Lynx Prototoxins: Roles of Endogenous
762   Mammalian Neurotoxin-Like Proteins in Modulating Nicotinic Acetylcholine Receptor Function
763   to Influence Complex Biological Processes. Front Pharmacol *10*, 343. 10.3389/fphar.2019.00343.

764   39.   Wu, M., Puddifoot, C.A., Taylor, P., and Joiner, W.J. (2015). Mechanisms of inhibition and
765   potentiation of α4β2 nicotinic acetylcholine receptors by members of the Ly6 protein family. J
766   Biol Chem *290*, 24509–24518. 10.1074/jbc.M115.647248.

767   40.   Demars, M.P., and Morishita, H. (2014). Cortical parvalbumin and somatostatin GABA neurons
768   express distinct endogenous modulators of nicotinic acetylcholine receptors. Mol Brain *7*, 75.
769   10.1186/s13041-014-0075-9.

770   41.   Miwa, J.M. (2021). Lynx1 prototoxins: critical accessory proteins of neuronal nicotinic
771   acetylcholine receptors. Curr Opin Pharmacol *56*, 46–51. 10.1016/J.COPH.2020.09.016.

772   42.   Falk, E.N., Norman, K.J., Garkun, Y., Demars, M.P., Im, S., Taccheri, G., Short, J., Caro, K.,
773   Mccraney, S.E., Cho, C., et al. (2021). Nicotinic regulation of local and long-range input balance
774   drives top-down attentional circuit maturation. Sci. Adv *7*.

775   43.   Anderson, K.R., Hoffman, K.M., and Miwa, J.M. (2020). Modulation of cholinergic activity
776   through lynx prototoxins: Implications for cognition and anxiety regulation. Neuropharmacology
777   *174*, 108071. 10.1016/j.neuropharm.2020.108071.

778   44.   Nectow, A.R., Moya, M. V., Ekstrand, M.I., Mousa, A., McGuire, K.L., Sferrazza, C.E., Field,
779   B.C., Rabinowitz, G.S., Sawicka, K., Liang, Y., et al. (2017). Rapid Molecular Profiling of
780   Defined Cell Types Using Viral TRAP. Cell Rep *19*, 655–667. 10.1016/j.celrep.2017.03.048.

781   45.   Heintz, N. (2004). Gene Expression Nervous System Atlas (GENSAT). Nature Neuroscience 2004
782   7:5 *7*, 483–483. 10.1038/nn0504-483.

783   46.   Webster, J.C., Francis, M.M., Porter, J.K., Robinson, G., Stokes, C., Horenstein, B., and Papke,
784   R.L. (1999). Antagonist activities of mecamylamine and nicotine show reciprocal dependence on
785   beta subunit sequence in the second transmembrane domain. Br J Pharmacol *127*, 1337–1348.
786   10.1038/SJ.BJP.0702686.

787   47.   Papke, R.L., Stokes, C., Muldoon, P., and Imad Damaj, M. (2013). Similar activity of
788   mecamylamine stereoisomers in vitro and in vivo. Eur J Pharmacol *720*, 264.
789   10.1016/J.EJPHAR.2013.10.018.

790   48.   Yao, Z., van Velthoven, C.T.J., Nguyen, T.N., Goldy, J., Sedeno-Cortes, A.E., Baftizadeh, F.,
791   Bertagnolli, D., Casper, T., Chiang, M., Crichton, K., et al. (2021). A taxonomy of transcriptomic
792   cell types across the isocortex and hippocampal formation. Cell. 10.1016/j.cell.2021.04.021.

793   49.   Tasic, B., Menon, V., Nguyen, T.N., Kim, T.K., Jarsky, T., Yao, Z., Levi, B., Gray, L.T.,
794   Sorensen, S.A., Dolbeare, T., et al. (2016). Adult mouse cortical cell taxonomy revealed by single
795   cell transcriptomics. Nat Neurosci *19*, 335–346. 10.1038/nn.4216.

50.   Tasic, B., Yao, Z., Graybuck, L.T., Smith, K.A., Nguyen, T.N., Bertagnolli, D., Goldy, J., Garren,
E., Economo, M.N., Viswanathan, S., et al. (2018). Shared and distinct transcriptomic cell types
across neocortical areas. Nature *563*, 72–78. 10.1038/s41586-018-0654-5.

51.   Yang, D., Günter, R., Qi, G., Radnikow, G., and Feldmeyer, D. (2020). Muscarinic and Nicotinic
Modulation of Neocortical Layer 6A Synaptic Microcircuits Is Cooperative and Cell-Specific.
Cerebral Cortex *30*, 3528–3542. 10.1093/cercor/bhz324.

52.   Ghezzi, F., Marques-Smith, A., Anastasiades, P.G., Lyngholm, D., Vagnoni, C., Rowett, A.,
Parameswaran, G., Hoerder-Suabedissen, A., Nakagawa, Y., Molnar, Z., et al. (2021). Non-
canonical role for lpar1-egfp subplate neurons in early postnatal mouse somatosensory cortex.
Elife *10*. 10.7554/ELIFE.60810.

53.   Wess, J.M., Isaiah, A., Watkins, P. v, and Kanold, P.O. (2017). Subplate neurons are the first
cortical neurons to respond to sensory stimuli. Proc Natl Acad Sci U S A *114*, 12602–12607.
10.1073/pnas.1710793114.

54.   Hansen, J.Y., Markello, R.D., Tuominen, L., Nørgaard, M., Kuzmin, E., Palomero-Gallagher, N.,
Dagher, A., and Misic, B. (2022). Correspondence between gene expression and neurotransmitter
receptor and transporter density in the human brain. Neuroimage *264*, 119671.
10.1016/J.NEUROIMAGE.2022.119671.

55.   Bae, Y.S., Lee, T.G., Park, J.C., Hur, J.H., Kim, Y., Heo, K., Kwak, J.Y., Suh, P.G., and Ryu, S.H.
(2003). Identification of a Compound That Directly Stimulates Phospholipase C Activity. Mol
Pharmacol *63*, 1043–1050. 10.1124/MOL.63.5.1043.

56.   Abbondanza, A., Bas, I.R., Modrak, M., Capek, M., Minich, J., Tyshkevich, A., Naser, S.,
Rangotis, R., Houdek, P., Sumova, A., et al. (2022). Nicotinic Acetylcholine Receptors Expressed
by Striatal Interneurons Inhibit Striatal Activity and Control Striatal-Dependent Behaviors. The
Journal of Neuroscience *42*, 2786. 10.1523/JNEUROSCI.1627-21.2022.

57.   Krjukova, J., Holmqvist, T., Danis, A.S., Åkerman, K.E.O., and Kukkonen, J.P. (2004).
Phospholipase C activator m-3M3FBS affects Ca2+ homeostasis independently of phospholipase
C activation. Br J Pharmacol *143*, 3–7. 10.1038/SJ.BJP.0705911.

58.   Sturgeon, R.M., and Magoski, N.S. (2018). A Closely Associated Phospholipase C Regulates
Cation Channel Function through Phosphoinositide Hydrolysis. Journal of Neuroscience *38*,
7622–7634. 10.1523/JNEUROSCI.0586-18.2018.

59.   Horowitz, L.F., Hirdes, W., Suh, B.-C., Hilgemann, D.W., Mackie, K., and Hille, B. (2005).
Phospholipase C in Living Cells Activation, Inhibition, Ca2+ Requirement, and Regulation of M
Current. Journal of General Physiology *126*, 243–262. 10.1085/JGP.200509309.

60.   Kulbatskii, D., Shenkarev, Z., Bychkov, M., Loktyushov, E., Shulepko, M., Koshelev, S.,
Povarov, I., Popov, A., Peigneur, S., Chugunov, A., et al. (2021). Human Three-Finger Protein
Lypd6 Is a Negative Modulator of the Cholinergic System in the Brain. Front Cell Dev Biol *9*,
2593. 10.3389/FCELL.2021.662227/BIBTEX.

61.   Arvaniti, M., Jensen, M.M., Soni, N., Wang, H., Klein, A.B., Thiriet, N., Pinborg, L.H., Muldoon,
P.P., Wienecke, J., Damaj, M.I., et al. (2016). Functional interaction between Lypd6 and nicotinic
acetylcholine receptors. J Neurochem *138*, 806–820. 10.1111/JNC.13718.

62.   Kuryatov, A., Onksen, J., and Lindstrom, J. (2008). Roles of accessory subunits in α4β2* nicotinic
837   receptors. Mol Pharmacol. 10.1124/mol.108.046789.

838   63.   Howe, W.M., Brooks, J.L., Tierney, P.L., Pang, J., Rossi, A., Young, D., Dlugolenski, K.,
839   Guillmette, E., Roy, M., Hales, K., et al. (2018). α5 nAChR modulation of the prefrontal cortex
840   makes attention resilient. Brain Struct Funct *223*, 1035–1047. 10.1007/s00429-017-1601-1.

841   64.   Sundberg, S.C., Lindström, S.H., Sanchez, G.M., and Granseth, B. (2018). Cre-expressing neurons
842   in visual cortex of Ntsr1-Cre GN220 mice are corticothalamic and are depolarized by
843   acetylcholine. Journal of Comparative Neurology *526*, 120–132. 10.1002/cne.24323.

844   65.   Molnár, Z., Luhmann, H.J., and Kanold, P.O. (2020). Transient cortical circuits match
845   spontaneous and sensory-driven activity during development. Science *370*.
846   10.1126/science.abb2153.

847   66.   Marx, M., Qi, G., Hanganu-Opatz, I.L., Kilb, W., Luhmann, H.J., and Feldmeyer, D. (2017).
848   Neocortical Layer 6B as a Remnant of the Subplate - A Morphological Comparison. Cerebral
849   Cortex *27*, 1011–1026. 10.1093/CERCOR/BHV279.

850   67.   Mechawar, N., and Descarries, L. (2001). The cholinergic innervation develops early and rapidly
851   in the rat cerebral cortex: a quantitative immunocytochemical study. Neuroscience *108*, 555–567.
852   10.1016/S0306-4522(01)00389-X.

853   68.   Hoerder-Suabedissen, A., Wang, W.Z., Lee, S., Davies, K.E., Goffinet, A.M., Rakić, S.,
854   Parnavelas, J., Reim, K., Nicolić, M., Paulsen, O., et al. (2009). Novel Markers Reveal
855   Subpopulations of Subplate Neurons in the Murine Cerebral Cortex. Cerebral Cortex *19*, 1738–
856   1750. 10.1093/CERCOR/BHN195.

857   69.   Tiong, S.Y.X., Oka, Y., Sasaki, T., Taniguchi, M., Doi, M., Akiyama, H., and Sato, M. (2019).
858   Kcnab1 is expressed in subplate neurons with unilateral long-range inter-areal projections. Front
859   Neuroanat *13*, 39. 10.3389/FNANA.2019.00039/BIBTEX.

860   70.   Hoerder-Suabedissen, A., Oeschger, F.M., Krishnan, M.L., Belgard, T.G., Wang, W.Z., Lee, S.,
861   Webber, C., Petretto, E., Edwards, A.D., and Molnár, Z. (2013). Expression profiling of mouse
862   subplate reveals a dynamic gene network and disease association with autism and schizophrenia.
863   Proc Natl Acad Sci U S A *110*, 3555–3560. 10.1073/pnas.1218510110.

864   71.   Zolnik, T.A., Ledderose, J., Toumazou, M., Trimbuch, T., Oram, T., Rosenmund, C., Eickholt,
865   B.J., Sachdev, R.N.S., and Larkum, M.E. (2020). Layer 6b Is Driven by Intracortical Long-Range
866   Projection Neurons. Cell Rep *30*, 3492-3505.e5. 10.1016/J.CELREP.2020.02.044.

867   72.   Egger, R., Narayanan, R.T., Guest, J.M., Bast, A., Udvary, D., Messore, L.F., Das, S., de Kock,
868   C.P.J., and Oberlaender, M. (2020). Cortical Output Is Gated by Horizontally Projecting Neurons
869   in the Deep Layers. Neuron *105*, 122-137.e8. 10.1016/J.NEURON.2019.10.011.

870   73.   Tekinay, A.B., Nong, Y., Miwa, J.M., Lieberam, I., Ibanez-Tallon, I., Greengard, P., and Heintz,
871   N. (2009). A role for LYNX2 in anxiety-related behavior. Proc Natl Acad Sci U S A *106*, 4477.
872   10.1073/PNAS.0813109106.

873   74.   Nichols, W.A., Henderson, B.J., Yu, C., Parker, R.L., Richards, C.I., Lester, H.A., and Miwa, J.M.
874   (2014). Lynx1 Shifts α4β2 Nicotinic Receptor Subunit Stoichiometry by Affecting Assembly in

the Endoplasmic Reticulum *. Journal of Biological Chemistry *289*, 31423–31432.
10.1074/JBC.M114.573667.

75. Ochoa, V., George, A.A., Nishi, R., and Whiteaker, P. (2016). The prototoxin LYPD6B modulates
heteromeric a3β4-containing nicotinic acetylcholine receptors, but not α7 homomers. The FASEB
Journal *30*, 1109–1119. 10.1096/FJ.15-274548.

76. Miwa, J.M., Ibaňez-Tallon, I., Crabtree, G.W., Sánchez, R., Šali, A., Role, L.W., and Heintz, N.
(1999). lynx1, an Endogenous Toxin-like Modulator of Nicotinic Acetylcholine Receptors in the
Mammalian CNS. Neuron *23*, 105–114. 10.1016/S0896-6273(00)80757-6.

77. Ibañez-Tallon, I., Miwa, J.M., Wang, H.L., Adams, N.C., Crabtree, G.W., Sine, S.M., and Heintz,
N. (2002). Novel modulation of neuronal nicotinic acetylcholine receptors by association with the
endogenous prototoxin lynx1. Neuron *33*, 893–903. 10.1016/S0896-6273(02)00632-3.

78. Morishita, H., Miwa, J.M., Heintz, N., and Hensch, T.K. (2010). Lynx1, a Cholinergic Brake,
Limits Plasticity in Adult Visual Cortex. Science (1979) *330*, 1238–1240.
10.1126/science.1195320.

79. Sherafat, Y., Chen, E., Lallai, V., Bautista, M., Fowler, J.P., Chen, Y.C., Miwa, J., and Fowler,
C.D. (2021). Differential Expression Patterns of Lynx Proteins and Involvement of Lynx1 in
Prepulse Inhibition. Front Behav Neurosci *15*. 10.3389/FNBEH.2021.703748.

80. Thomsen, M.S., Arvaniti, M., Jensen, M.M., Shulepko, M.A., Dolgikh, D.A., Pinborg, L.H.,
Härtig, W., Lyukmanova, E.N., and Mikkelsen, J.D. (2016). Lynx1 and Aβ1–42 bind
competitively to multiple nicotinic acetylcholine receptor subtypes. Neurobiol Aging *46*, 13–21.
10.1016/J.NEUROBIOLAGING.2016.06.009.

81. Shenkarev, Z.O., Shulepko, M.A., Bychkov, M.L., Kulbatskii, D.S., Shlepova, O. v., Vasilyeva,
N.A., Andreev-Andrievskiy, A.A., Popova, A.S., Lagereva, E.A., Loktyushov, E. v., et al. (2020).
Water-soluble variant of human Lynx1 positively modulates synaptic plasticity and ameliorates
cognitive impairment associated with α7-nAChR dysfunction. J Neurochem *155*, 45–61.
10.1111/JNC.15018.

82. Sadahiro, M., Demars, M.P., Burman, P., Yevoo, P., Zimmer, A., and Morishita, H. (2020).
Activation of Somatostatin Interneurons by Nicotinic Modulator Lypd6 Enhances Plasticity and
Functional Recovery in the Adult Mouse Visual Cortex. Journal of Neuroscience *40*, 5214–5227.
10.1523/JNEUROSCI.1373-19.2020.

83. Viswanathan, S., Bandyopadhyay, S., Kao, J.P.Y., and Kanold, P.O. (2012). Changing
Microcircuits in the Subplate of the Developing Cortex. Journal of Neuroscience *32*, 1589–1601.
10.1523/JNEUROSCI.4748-11.2012.

84. Viswanathan, S., Sheikh, A., Looger, L.L., and Kanold, P.O. (2017). Molecularly Defined
Subplate Neurons Project Both to Thalamocortical Recipient Layers and Thalamus. Cereb Cortex
27, 4759–4768. 10.1093/CERCOR/BHW271.

85. Hoerder-Suabedissen, A., Hayashi, S., Upton, L., Nolan, Z., Casas-Torremocha, D., Grant, E.,
Viswanathan, S., Kanold, P.O., Clasca, F., Kim, Y., et al. (2018). Subset of Cortical Layer 6b
Neurons Selectively Innervates Higher Order Thalamic Nuclei in Mice. Cerebral Cortex *28*, 1882–
1897. 10.1093/CERCOR/BHY036.

86.   Zhang, M., Eichhorn, S.W., Zingg, B., Yao, Z., Cotter, K., Zeng, H., Dong, H., and Zhuang, X.
(2021). Spatially resolved cell atlas of the mouse primary motor cortex by MERFISH. Nature
2021 598:7879 *598*, 137–143. 10.1038/s41586-021-03705-x.

87.   Muñoz-Castañeda, R., Zingg, B., Matho, K.S., Chen, X., Wang, Q., Foster, N.N., Li, A.,
Narasimhan, A., Hirokawa, K.E., Huo, B., et al. (2021). Cellular anatomy of the mouse primary
motor cortex. Nature 2021 598:7879 *598*, 159–166. 10.1038/s41586-021-03970-w.

88.   Yao, Z., Liu, H., Xie, F., Fischer, S., Adkins, R.S., Aldridge, A.I., Ament, S.A., Bartlett, A.,
Behrens, M.M., van den Berge, K., et al. (2021). A transcriptomic and epigenomic cell atlas of the
mouse primary motor cortex. Nature 2021 598:7879 *598*, 103–110. 10.1038/s41586-021-03500-8.

89.   Paxinos, G., and Franklin, K. (2004). Paxinos and Franklin's the Mouse Brain in Stereotaxic
Coordinates.

90.   Gong, S., Doughty, M., Harbaugh, C.R., Cummins, A., Hatten, M.E., Heintz, N., and Gerfen, C.R.
(2007). Targeting Cre recombinase to specific neuron populations with bacterial artificial
chromosome constructs. Journal of Neuroscience *27*, 9817–9823. 10.1523/JNEUROSCI.2707-
07.2007.

91.   Gong, S., Zheng, C., Doughty, M.L., Losos, K., Didkovsky, N., Schambra, U.B., Nowak, N.J.,
Joyner, A., Leblanc, G., Hatten, M.E., et al. (2003). A gene expression atlas of the central nervous
system based on bacterial artificial chromosomes. Nature *425*, 917–925. 10.1038/NATURE02033.

92.   Madisen, L., Zwingman, T.A., Sunkin, S.M., Wook Oh, S., Zariwala, H.A., Gu, H., Ng, L.L.,
Palmiter, R.D., Hawrylycz, M.J., Jones, A.R., et al. (2010). A robust and high-throughput Cre
reporting and characterization system for the whole mouse brain. 10.1038/nn.2467.

93.   Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., Hao, Y.,
Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data.
Cell *177*, 1888-1902.e21. 10.1016/j.cell.2019.05.031.

94.   Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-cell
transcriptomic data across different conditions, technologies, and species. Nat Biotechnol *36*, 411–
420. 10.1038/nbt.4096.

95.   Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J., Wilk, A.J.,
Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-cell data. Cell *184*,
3573-3587.e29. 10.1016/j.cell.2021.04.048.

96.   Ho, J., Tumkaya, T., Aryal, S., Choi, H., and Claridge-Chang, A. (2019). Moving beyond P
values: data analysis with estimation graphics. Nat Methods *16*, 565–566. 10.1038/s41592-019-
0470-3.

 Supplementary Material Legends

Video S1: Video shows Chrna5+ neurons in mPFC slices from *Chrna5*-Cre$^{/+}$Ai96$^{/+}$ mice
responding to exogenous application of 1mM acetylcholine with an increase in GCaMP6s
fluorescence signal

Video S2: Video shows Syt6+ neurons from *Syt6*-Cre$^{/+}$Ai96$^{/+}$ mice responding to exogenous
application of 1mM acetylcholine with an increase in GCaMP6s fluorescence signal.

(Note: Figures S1-5 and Tables S1-3 are in one PDF file. Table S4 is attached as a separate
excel file.)

Figure S1. *Chrna5*-Cre reporter expression in different brain regions.

Figure S2. Chrna5+ neurons in primary somatosensory cortex are super-responders to
optogenetic acetylcholine release.

Figure S3. Chrna5+ cells in primary visual cortex from *Chrna5*-Cre$^{/+}$ Ai14$^{/+}$ Syt6-EGFP$^{/+}$ mice.

Figure S4. Chrna5+ acetycholine super-responders in layer 6b.

Figure S5. Transcriptomic characterization of Chrna5+ super-responders in other cortical
regions.

Table S1. Intrinsic electrophysiological properties of Chrna5+ and Chrna5- unlabeled deep-layer
neurons in *Chrna5*-Cre$^{/+}$Ai14$^{/+}$ChAT-ChR2$^{/+}$ mice.

Table S2. Intrinsic electrophysiological properties of Chrna5+ and Syt6+ deep-layer neurons.

Table S3. Comparing expression of major genes modulating postsynaptic cholinergic responses
in Chrna5+ and Chrna5+Syt6+ neurons.

Table S4. Excel file containing all differentially expressed genes between Chrna5+ and
Chrna5+Syt6+ neurons with adjusted p value < 0.05

A

Chrna5-Cre$^{/+}$
ChAT-ChR2$^{/+}$ ×   Ai14   →   *Chrna5*-Cre$^{/+}$ Ai14$^{/+}$
                                    ChAT-ChR2$^{/+}$

B

Chrna5+ cells   ChAT axons

C

Chrna5+ labeled L6 pyramidal neuron

10 µm

Chrna5- unlabeled L6 pyramidal neuron

50 µm

D

– – Chrna5+ labeled
– – Chrna5- unlabeled

0.5 s

5 pA

20 ms

5 pA

E

Rising slope (pA/s)

Chrna5+ labeled   Chrna5- unlabeled

*

F

Peak current (pA)

Chrna5+ labeled   Chrna5- unlabeled

*

G

Chrna5-Cre$^{/+}$ ×   *Syt6*-EGFP$^{/+}$ →   *Chrna5*-Cre$^{/+}$ Ai14$^{/+}$
                            Ai14$^{/+}$ *Syt6*-EGFP$^{/+}$

Chrna5+   Syt6+   Overlay

L6   L6   L6

500 µm

H

100 µm

I

L6   L5   L2/3   L1

Chrna5+   Syt6+   Chrna5+Syt6+

100 µm

J

% of total cells

Chrna5+   Syt6+   Chrna5+
                    Syt6+

Proportions of cells

37%   Chrna5+
39%   Chrna5+Syt6+
24%   Syt6+

A GCaMP6s Ca²⁺ imaging

1mM ACh, 15s | DHBE | DHBE+Atropine

Chrna5-Cre^/+ Ai96^/+

Syt6-Cre^/+ Ai96^/+

DHBE   DHBE+Atropine

DHBE   DHBE+Atropine

B Chrna5+ cells

1mM ACh   DHBE   DHBE + Atropine

1 ΔF/F

30 s

Syt6+ cells

0.5 ΔF/F

30 s

C DHBE

i   ****

% response remaining

Chrna5+   Syt6+

+Mean

ii KS test p < 10⁻⁴

Cumulative fraction

% response remaining

iii   Non-zero   Zero

83%   52%

n = 71   n = 112

D DHBE+Atropine

i   ****

% response remaining

Chrna5+   Syt6+

ii KS test p < 10⁻⁴

Cumulative fraction

% response remaining

iii   Non-zero   Zero

41%   6%

n = 59   n = 112

E

Chrna5-Cre^/+Ai14^/+

Syt6-Cre^/+Ai14^/+ or *Syt6*-EGFP

20 mV
10 s

1 mM ACh

F Acetylcholine

Peak frequency (Hz)

*   Chrna5+   Syt6+

+Competitive block

**   Chrna5+   Syt6+

+Non-competitive block

**   Chrna5+   Chrna5 +MEC

Acetylcholine

Opto Nicotinic response

LY6G6E
LYPD1
LYPD6B

Cell-type specific
Lynx modulation of
nicotinic receptors

CT *CT* *CT*

SP

SP

Layer 6

Layer 6b

CT- Corticothalamic
SP- Subplate

Syt6+

Chrna5+Syt6+

Chrna5+

Cholinergic afferents

 α4β2α5 nicotinic receptor

 α4β2 nicotinic receptor

Lynx prprototoxins

 LY6G6E

 LYNX1

 LYPD6B

 LYPD1
(LYNX2)