Structural basis of LRPPRC-SLIRP-dependent translation by the mitoribosome

Vivek Singh[1,7], J. Conor Moran[2,7], Yuzuru Itoh[1,6,*], Iliana C. Soto[3], Flavia Fontanesi[2], Mary Couvillion[3], Martijn A. Huynen[4], Stirling Churchman[3,*], Antoni Barrientos[5,*], Alexey Amunts[1,*]

[1] Science for Life Laboratory, Department of Biochemistry and Biophysics, Stockholm University, Solna 17165, Sweden.

[2] Department of Biochemistry and Molecular Biology, University of Miami Miller School of Medicine, Miami FL 33136, USA

[3] Blavatnik Institute, Department of Genetics, Harvard Medical School, Boston MA 02115, USA.

[4] Medical BioSciences, Radboud University Medical Center, Nijmegen 6525, the Netherlands.

[5] Department of Neurology, University of Miami Miller School of Medicine, Miami FL 33136, USA.

[6] Current address: Department of Biological Sciences, Graduate School of Science, University of Tokyo, 113-0033 Tokyo, Japan.

[7] These authors contributed equally

* correspondence: yuzuru.itoh@bs.s.u-tokyo.ac.jp, churchman@genetics.med.harvard.edu, abarrientos@med.miami.edu, amunts@scilifelab.se

In mammalian mitochondria, mRNAs are co-transcriptionally stabilized by the protein factor LRPPRC. Here, we characterize LRPPRC as an mRNA delivery factor and report its cryo-EM structure in complex with SLIRP, mRNA and the mitoribosome. The structure shows that LRPPRC associates with the mitoribosomal proteins mS39 and the N-terminus of mS31 through recognition of the LRPPRC helical repeats. Together, the proteins form a corridor for hand-off the mRNA. The mRNA is directly bound to SLIRP, which also has a stabilizing function for LRPPRC. To delineate the effect of LRPPRC on individual mitochondrial transcripts, we used an RNAseq approach, metabolic labeling and mitoribosome profiling that showed a major influence on ND1, ND2, ATP6, COX1, COX2, and COX3 mRNA translation efficiency. Our data suggest that LRPPRC-SLIRP acts in recruitment of mitochondrial mRNAs to modulate their translation. Collectively, the data define LRPPRC-SLIRP as a regulator of the mitochondrial gene expression system.

The mitoribosome is organised in a small and large subunit (SSU and LSU) that are assembled from multiple components in a coordinated manner and through regulated sequential mechanisms [1-5]. The SSU formation is accomplished by the association of the mitoribosomal protein mS37 and the initiation factor mtIF3, leading to a mature state that is ready for translation of the mRNA [3,6]. In mammals, mitochondrial transcription is polycistronic and gives rise to two long transcripts, corresponding to almost the entire heavy and light mtDNA strands. The individual mRNAs are available for translation only after they are liberated from the original polycistronic transcripts and polyadenylated [7]. In *Escherichia coli*, a functional transcription-translation coupling mechanism has been characterised involving a physical association of the RNA polymerase with the SSU, termed the expressome [8-10]. In contrast, in mammalian mitochondria, nucleoids are not compartmented with protein synthesis; mitoribosomes are independently tethered to the membrane [11,12], and no coupling with the RNA polymerase has been reported. In addition, human mitochondrial mRNAs and the mitoribosome do not have the Shine–Dalgarno (SD) and anti-SD sequences that are used to recruit mRNA to SSU in bacteria [13]. Mitochondrial mRNAs also lack cap 5′ modifications, which is a hallmark of eukaryotic cytosolic mRNAs translation initiation. In the cytosol, mRNA is recruited to a pre-initiation complex, consisting of the SSU and translation initiation factors, which then scans along the 5′ untranslated region to find the start codon [14,15]. No equivalent mechanism has been found in

mitochondria, and thus, how mRNAs are delivered for translation in mitochondria remained
unknown.

The 130-kDa protein factor LRPPRC (leucine-rich pentatricopeptide repeat-containing protein),
a member of a Metazoa-specific pentatricopeptide repeat family, was reported to act as a global
mitochondrial mRNA chaperone that binds co-transcriptionally [16-18]. LRPPRC is an integral part
of the post-transcriptional processing machinery required for mRNA stability, polyadenylation,
and translation [16-19]. Mutations in the gene encoding for LRPPRC lead to French-Canadian type
Leigh syndrome (LSFC) an untreatable paediatric neurodegenerative disorder caused by
ultimately impaired mitochondrial energy conversion [20].

LRPPRC has been reported to interact with a small 11-kDa protein cofactor SLIRP (SRA stem-
loop-interacting RNA-binding protein) [21,22] that plays roles in LRPPRC stability and maintaining
steady-state mRNA levels [23]. *SLIRP* silencing results in the destabilization of respiratory
complexes, loss of enzymatic activity, and reduction in mRNA levels, implicating a role in
mRNA homeostasis [24]. *SLIRP* variants cause a respiratory deficiency that leads to mitochondrial
encephalomyopathy [25]. In addition, *SLIRP* knockdown results in increased turnover of LRPPRC
[23,25,26], and *in vivo* co-stabilisation suggests that the two entities have interdependent functions
[23,27]. The interaction of LRPPRC and SLIRP *in vitro* has been previously studied [28].

LRPPRC has also been implicated in coordinating mitochondrial mRNA stability and translation
[18,29]. Previous analysis showed a correlation between presence of LRPPRC and mRNA on the
mitoribosome [30]. However, there are no structures available for LRPPRC, SLIRP, or any
complexes containing them, and *in vitro* reconstitution could not provide meaningful
information, in part because not all the components of the mitochondrial gene expression system
have been characterised. Thus, although isolated mitoribosomal models have been determined [31-
33], the molecular mechanisms of mRNA delivery to the SSU for activation of translation and the
potential involvement of LRPPRC-SLIRP in this process remained unknown.

Results

Structure determination of LRPPRC-SLIRP bound to the mitoribosome

To explore the molecular basis for translation activation in human mitochondria, we used low
salt conditions to isolate a mitoribosome:LRPPRC-SLIRP-mRNA complex for cryo-EM. We

merged particles containing tRNAs in the A- or P-site, as well as an extra density in the vicinity

of the mRNA entry channel and applied iterative local-masked refinement and classification with

signal subtraction (Extended Data Fig. 1a). It resulted in a 2.9 Å resolution map of the

mitoribosome during mRNA delivery to the SSU, with the local resolution for the LRPPRC

binding region of ~3.4 Å (Extended Data Fig. 1b,c). The reconstruction showed a clear density

only for the LRPPRC N-terminal domains (residues 64-644, average local resolution ~4.5 Å)

bound to the SSU head, which is consistent with a previous mass-spectrometry analysis

(Extended Data Fig. 1d,e) [30]. It allowed us to model 34 α-helices, 17 of which (α2-18) form a

ring-like architecture, while the rest form an extended tail that adopts a 90° curvature and

projects 110 Å from the SSU body in parallel to the L7/L12 stalk (Fig. 1a,b). The C-terminal

domains (residues 645-1394) were not resolved. The complete LRPPRC model obtained with

AlphaFold2 [34] combined with Translation/Liberation/Screw Motion Determination (TLSMD)

analysis [35,36] defined the C-terminal domains as individual segments, indicating potential

flexibility (Extended Data Fig. 2).

When LRPPRC-SLIRP is bound to the mitoribosome, a previously disordered density of mS31

that extends from the core also becomes ordered, revealing its N-terminal region (Fig. 2a). This

region is arranged in two helix-turn-helix motifs, offering a 1930 $Å^2$ surface area for direct

interactions with LRPPRC (Fig. 1c, Fig. 2). The position of the LRPPRC residue 354, in which

the mutation A354V leads to LSFC with a clinically distinct cytochrome *c* oxidase deficiency

and acute fatal acidotic crises is in a buried area of helix 17, close to the mRNA binding region

(Fig. 1c, Extended Data Fig. 2a). A previous study demonstrated that the mutation is

abolishing the interaction with the protein SLIRP [37]. Consistent with mass spectrometry analysis

[28] and the interaction interface previously determined [37], the remaining associated density was

assigned as SLIRP, found to be located close to the Epsin N-terminal homology (ENTH) domain

of LRPPRC (Fig. 2a). Finally, SLIRP is connected to an elongated density on the LRPPRC

surface that is also associated with six of the mitoribosomal proteins and corresponds to the

endogenous mRNA (Fig. 2).

Figure 1. Structure of mitoribosome with LRPPRC-SLIRP bound to mRNA.
a, Overview of the mitoribosome:LRPPRC-SLIRP model. Right panel, top view of the model colored by atomic *B*-factor (Å^2), tRNAs in surface (red, orange, brown), mRNA path (light green) is highlighted. **b,** A close-up view of the mitoribosome:LRPPRC-SLIRP-mRNA

interactions. LRPPRC associates with mS31-m39 via a ring-like structure (α2-18) that together form a corridor for the hand-off the mRNA from SLIRP. **c,** Contact sites between LRPPRC and mS31-mS39 (within 4 Å distance), view from the interface. Right panel, schematic diagram showing the topology of LRPPRC consisting of 34 helices. Colours represent engagement in interactions with mS39 (light green), mS31 (purple), SLIRP (orange), mRNA (blue). The position of LSFC variant (A354V) is indicated with an asterisk on helix 17.

Figure 2. Overview of density for LRPPRC, SLIRP and mRNA and their interactions with SSU proteins.
a, The density map for LRPPRC (dark green), SLIRP (orange), mRNA (blue) on the SSU is shown in the centre. The model and map for mS39-LRPPRC-SLIRP and corresponding bound mRNA residues are shown in closeup views on the left, and arginines involved in mRNA binding are indicated. The bottom closeup views show SLIRP with its associated densities for LRPPRC and mRNA. For clarity, the map for SLIRP has been low-pass filtered to 6 Å resolution. **b,** Schematic of protein-protein interactions, where node size corresponds to relative molecular mass. Nodes of proteins involved in mRNA binding are encircled in blue. **c,** RRM containing proteins: SLIRP, hnRNPA1/B2 (PDBID 5WWE), PolyA binding protein (PABP, PDBID 1CVJ) and Nucleolin (PDBID 1RKJ) are shown in complex with RNA with RNP1 and RNP2 sub-motifs colored blue.

SLIRP is stably associated with mRNA and LRPPRC on the SSU

The binding of SLIRP in our model is enabled via LRPPRC helices α20 and 22, which is consistent with crosslinking mass spectrometry data and mutational analysis [28]. The structure reveals that SLIRP links the nuclear export signal (NES) domain with the curved region of the ENTH domain of LRPPRC (Fig. 1b, Fig. 1c). This binding of SLIRP contributes to a corridor for the mRNA that extends to mS31 and mS39 (Fig. 1b, Supplementary Video 1). Through this corridor, the mRNA extends over ~180 Å all the way to the decoding center (Fig. 2a). In our structure, SLIRP is oriented such that the conserved RNA recognition motif (RRM), including its submotifs RNP1 (residues 21-26) and RNP2 (residues 60-67) [38-40] form an interface with modeled mRNA (Fig. 2). The arrangement of RNP1 and RNP2 with respect to the mRNA is similar to that observed in previously reported structures of other RRM proteins [41-43] (Fig. 2c, Extended Data Fig. 3). Moreover, residues R24, R25 of RNP2 and L62 of RNP1 motifs previously implicated to be required for RNA binding by SLIRP [37] are positioned within an interacting distance of the mRNA (Fig. 2a). Thus, SLIRP contributes to the LRPPRC specific scaffold, and accounts for a role in binding the mRNA.

The *B*-factor distribution of SLIRP in our model is similar to that of LRPPRC, while still lower than some of the more mobile components of the mitoribosome, such as the acceptor arm of the CP-tRNA^{Val} (Fig. 1a). This indicates a functionally relevant association with LRPPRC in terms of stability of binding. Our finding that SLIRP is involved in hand-off the mRNA to the mitoribosome provides a mechanistic explanation for the previous results from biochemical studies showing that SLIRP affects LRPPRC properties *in vitro* [27,28], and the presentation of the mRNA to the mitoribosome *in vivo* [23].

Since in *E. coli*, the expressome-mediating protein NusG was proposed to regulate mRNA unwinding [8], and SSU proteins uS3 and uS4 have an intrinsic RNA helicase activity [44], we searched for known helicase signature motifs [45] in the LRPPRC sequence, but no such motifs were present. In the mitoribosome, where the mRNA channel entry site is located, a bacteria-like ring-shaped entrance is missing, the entrance itself has shifted, and its diameter expanded [31]. The mRNA extends all the way into the head/beak of the SSU stabilised by mitoribosome-specific components: mS39 helical repeats, mS35 N-terminus that extends from the side of the SSU head, and N-terminal extension of uS9m that contacts the mRNA nucleotide at position 15 (numbered from the E-site).

LRPPRC-SLIRP hands-off the mRNA to mS31-mS39, channeling it for translation

Next, we analysed the structural basis for the complex formation. The association of LRPPRC with the mitoribosome involves the helices α1, 2, 6-11, that form a mitoribosome-binding surface (Fig. 1b, Fig. 1c). The binding is mediated by four distinct contacting regions (Extended Data Fig. 6): 1) α1-2 (residues 64-95) is flanked by a region of mS39, a PPR domain-containing protein, that consists of four bundled helices (α11-14); 2) α7 and 9 form a shared bundle with two N-terminal helices of mS31 (residues 175-208, stabilized by C-terminal region of mS39) that encircle the NES-rich domain to complement the PPR domain; 3) α10-11 are capped by a pronounced turn of mS31 (residues 209-232) acting as a lid that marks the LRPPRC boundary, and it is sandwiched by the mS39 helix α19 and C-terminus from the opposite side; 4) in addition, α11 is also positioned directly against the helix α23 of mS39. Thus, LRPPRC docks onto the surface of the mitoribosome via mS31-mS39, which are tightly associated with each other, and each provides two contact patches to contribute to stable binding.

Based on the structural analysis, the hand-off of the mRNA for translation is mediated by four of the LRPPRC helices: α1, 2, 16, and 18 (Fig. 1b, Fig. 1c, Supplementary Video 1). The mRNA nucleotides 33-35 (numbered from the E-site) are stabilized in a cleft formed by α1-2 on one side and α16, 18 on the other. Nucleotides 31 and 32 contact residues R332, R333 from LRPPRC, as well as R344 from mS39 (Fig. 2a). This region is within 120-130 Å from the P-site. The involvement of the NES-rich domain of LRPPRC in mRNA binding in our structure is consistent with a biochemical analysis of recombinant LRPPRC where the N-terminal PPR segments were systematically removed, which showed a reduced formation of protein-RNA complexes [28]. The rest of the mRNA is situated too far from LRPPRC to interact with it. Here, the mRNA is handed to mS31-mS39, consistently with a translation initiation complex [46].

In the structure, mS31, mS39, and LRPPRC together form a 60-Å long corridor that channels the mRNA from SLIRP toward the mitoribosomal core (Fig. 1b, Supplementary Video 1). The binding of LRPPRC is mediated by four distinct contact regions (Extended Data Fig. 6): contact-1 is formed by α1-2 (residues 64-95) flanked by the mS39 PPR domain of four bundled helices (α11-14); contact-2 is formed by α7, 9 that generate a shared bundle with two N-terminal helices of mS31 (residues 175-208, stabilized by C-terminal region of mS39) that encircle the head to complement the PPR domain; contact-3 is formed by α10-11 that are capped by a pronounced turn of mS31 (residues 209-232) acting as a lid that marks the LRPPRC boundary, and it is

sandwiched by the mS39 helix α21 and C-terminus from the opposite side; contact-4 is formed

by α11 that is positioned directly against the mS39 α23.

With respect to mRNA binding, nucleotides 26-30 bind mS39 PPR domain 5, and nucleotide 26

connects to contact-2 (Fig. 2a, Extended Data Fig. 6). Thus, the mRNA hand-off is achieved

through functional cooperation between LRPPRC-SLIRP and mS31-mS39. Therefore, in the

mitoribosome:LRPPRC-SLIRP with mRNA model, LRPPRC performs three functions:

coordination of SLIRP, which plays a key role in the process of mRNA recruitment, association

with the SSU, and hand-off of the mRNA for translation (Fig. 1, Extended Data Fig. 6).

Supplementary Video 1).

LRPPRC is recruited for translation of mRNAs

Next, we asked whether LRPRRC-SLIRP delivers all mRNAs to the mitoribosome or is

selective. We generated an *LRPPRC*-knockout cell line [29] that was rescued with either a wild-

type *LRPPRC* or a variant carrying the LSFC founder mutation A354V [20] (Extended Data Fig.

4). The steady-state levels of the LSFC variant were reduced by 60%, suggesting protein

instability as reported in patients [22], and the levels of SLIRP were equally decreased (Extended

Data Fig. 4). We then implemented an RNAseq approach that confirmed a substantially depleted

mitochondrial transcriptome [17,47,48] (Fig. 3a). In the LRPPRC-knockout, transcripts from the

heavy strand were lowered by 1.5-4-fold, except for ND3, which remained stable, consistent

with protein synthesis data (Fig. 3b), while the single light strand-encoded *ND6* mRNA was not

affected as reported [16], and the effect of the LSFC mutation on RNA stability was limited to six

transcripts (Fig. 3a), indicating LRPPRC's role in heavy strand mRNA stability is non-specific.

Metabolic labeling assays using [^{35}S]- methionine indicated that incorporation of the

radiolabeled amino acid into most newly synthesized mitochondrial proteins is severely

decreased in *LRPPRC*-KO cells (Fig. 3b). However, there were differential effects among

transcripts; synthesis of ND3, ND4L, and ATP8 remained above 50% of the WT, but translation

of other transcripts proceeded at a lower rate (*e.g.*, *ND1*, *ND2*, or *ATP6*) or was virtually blocked

(*e.g.*, *COX1, COX2, or COX3)* (Fig. 3b). The translational defect results in a decrease in the

steady-state levels of the four OXPHOS complexes that contain mtDNA-encoded subunits

(Extended Data Fig. 5).

To assess whether the mitochondrial translation efficiency (TE) is decreased in an mRNA-specific manner in the *LRPPRC*-knockout, we performed mitoribosome profiling (Fig. 4). These data show a decreased TE, calculated by dividing spike-in normalized ribosome footprint reads by the spike-in normalized RNA sequencing reads (in other words, how well a particular transcript is translated) [49,50]. In *LRPPRC*-knockout cells, TE was attenuated for *COX1* and *COX2* transcripts and the bicistronic *ATP8/ATP6* transcript, intriguingly more for ATP6 than ATP8 (Fig. 4a). Thus, our data suggest that LRPPRC-SLIRP is required for the translation of some transcripts, in agreement with the metabolic labeling of newly synthesized mitochondrial products (Fig. 3b).

To support the role of LRPPRC in mRNA binding, we determined an average length of the mitoribosome-protected fragments using mitoribosome profiling (Fig 4b). In the *LRPPRC*-knockout cells, we observed a decrease in the average protected fragment length, compared to the wild type (Fig 4b). This observation is consistent with the structural data showing the association of LRPPRC with mRNA and the mitoribosome. Previous studies also showed that LRPPRC–SLIRP relaxes structures of mRNAs [17], potentially to expose it to initiate translation [16]. The average protected fragment length in LSFC cells were smaller than in wild-type cells, similar to the *LRPPRC*-knockout (Fig 4b), suggesting that whereas the mutant protein participates in translation, it does so differently than the wild-type protein

Figure 3. Mechanism of LRPPRC-SLIRP mediated mRNA binding and stabilization.

a, Whole-cell RNAseq normalized by read depth, comparing *LRPPRC*-KO cells (KO) with KO

cells reconstituted with WT *LRPPRC* (KOR) or the LSFC variant (A354V). The results are the

average of two biological replicates. The differentially expressed mitochondrial transcripts are

color-coded: coding for subunits of cytochrome *c* oxidase (CIV) in red, NADH dehydrogenase

(CI) in yellow, coenzyme Q-cytochrome *c* oxidoreductase (CIII) in blue, and ATP synthase (CV)

in green. **b,** Metabolic labeling with [^{35}S]-labeled methionine of newly-synthesized

mitochondrial polypeptides for the indicated times, in the presence of emetine to inhibit cytosolic

protein synthesis, in whole HEK-293T WT, *LRPPRC*-KO cells, and KO cells reconstituted with

LRPPRC (KO+WT) or the LSFC variant (KO+LSFC). Bottom panel, representative plots of

[^{35}S]-labeled methionine incorporation into specific polypeptides in WT or *LRPPRC*-KO cells.

The images were quantified in two independent experiments.

Figure 4. Mitochondrial translation efficiency is decreased in *LRPPRC*-KO cells. a, Change in TE and RNA abundance in *LRPPRC*-KO cells compared to LRPPRC-reconstituted cells ("rescue"). Mitoribosome profiling data and RNA-seq data were normalized using a mouse lysate spike-in control (RPKS) [29], then TE was calculated from these normalized values (mitoribosome profiling / RNA-seq). Biological replicates are shown as individual points. The mitochondrial transcripts are color-coded as in Fig. 3. **b**, Heat map showing the length distribution for reads mapping to mitochondrial mRNAs [29].

The mitoribosome:LRPPRC-SLIRP complex is specific to Metazoa

To place the structural data into an evolutionary context, we performed comparative phylogenetic analysis of the proteins involved in the mRNA hand-off process. Since the mitochondrial rRNA has been generally reduced in Metazoa [51], we examined whether this loss might coincide with the origin of LRPPRC and its interactors. The orthology database EGGNOGG [52] and previous analysis [53] indicated that LRPPRC and mS31 are only present in Bilateria, while mS39 only occurs in Metazoa. We then confirmed the results with more sensitive homology detection [54] followed by manual sequence analysis examining domain composition, which put the origin of LRPPRC and mS31 at the root of the Metazoa. Thus, the appearance of these proteins coincides with the loss of parts of the rRNA (Fig. 5a). SLIRP appears to originate slightly later than LRPPRC, but its small size makes determining its phylogenetic origin less conclusive.

The correlation between rRNA reduction and protein acquisition is important, because the rRNA regions h16 (410-432) and h33-37 (997-1118) that bridge the mRNA to the channel entrance in bacteria [8] are either absent or reduced in the metazoan mitoribosome. However, a superposition of the mitoribosome:LRPPRC-SLIRP-mRNA complex with *E. coli* expressome [8] shows not only

that the nascent mRNA follows a comparable path in both systems, but also the mRNA

delivering complexes bind in a similar location with respect to their ribosomes (Fig. 5b). To test

whether protein-protein interactions can explain the conservation, we compared the interface

with the *E. coli* expressome [8]. Indeed, in the expressome, NusG binds to uS10 and restrains

RNAP motions [8], and in our structure uS10m has a related interface between its α2 helix with

mS31-mS39, which induces association of these two proteins (Fig. 5c, Extended Data Fig. 7).

Yet, most of the interactions rely on a mitochondria-specific N-terminal extension of uS10m

where it shares a sheet with mS39 via the strand β1, and helices α1,16 and 18 are further

involved in the binding (Extended Data Fig. 7). A similar conclusion can be reached from

comparison with the *M. pneumoniae* expressome [10]. Together, this analysis suggests that a

specific protein-based mechanism must have evolved in the evolution of the metazoan

mitoribosome for mRNA recognition and protection.

Figure 5. Formation of mitoribosome:LRPPRC-SLIRP and 70S:RNAP complexes.
a, Phylogenetic analysis shows correlation between acquisition of LRPPRC, SLIRP, mS31, mS39 and reduction of rRNA in Metazoa. Black rectangles indicate the presence of proteins, grey indicates uncertainty about the presence of an ortholog. *Hs*: *Homo sapiens, Ds: Drosophila melanogaster, Nv*: *Nematostella vectensis, Ta: Trichoplax adhaerens, Aq*: *Amphimedon queenslandica* (sponge)*, Mb*: *Monosiga brevicollis* (unicellular choanoflagellate)*, Co*: *Capsaspora owczarzaki* (protist)*, Sc*: *Saccharomyces cerevisiae* (fungi)*, Rp*: *Rickettsia prowazekii* (alpha-proteobacterium). Dating in million years ago (mya) is based on previously described method [55]. **b,** Model of the mitoribosome:LRPPRC-SLIRP complex compared with the uncoupled model of the expressome from *E. coli* [8]. **c,** Schematic representation indicating association of mRNA-delivering proteins in the mitoribosome compared to NusG-coupled expressome [8].

Discussion

LRPPRC is an mRNA chaperone that regulates human mitochondrial transcription and translation and is involved in a neurodegenerative disorder. In this study, we report the cryo-EM structure of the LRPPRC-SLIRP in complex with the mitoribosome and characterize its function with respect to the mRNA delivery. We identified that LRPPRC, in complex with SLIRP, binds to mRNAs to hand-off transcripts to the mitoribosome for translation. The docking of LRPPRC is realized through the mitoribosomal proteins mS39 and the N-terminus of mS31, that together recognize eight of the LRPPRC helical repeats. The structural comparison with the unbound state uncovers that the N-terminus of mS31 adopts a stable conformation upon LRPPRC association.

Our structure also shows that SLIRP is directly involved in interactions with mRNA. Those interactions are supported by the comparison with other RNA-binding proteins that contain RNP domains, similar to SLIRP. SLIRP further stabilizes the architecture of LRPPRC, and both are required for mRNA binding. The mRNA is then channeled through a corridor formed with mS39 towards the decoding center.

Although *LRPPRC* knockout results in an overall decrease in the steady-state levels of the four OXPHOS complexes that contain mtDNA-encoded subunits, by implementing an RNAseq approach and metabolic labeling assays, we show that beyond its role in mRNA stabilization, LRPPRC has differential effects on the translational efficiency of mitochondrial transcripts. Specifically, the synthesis of ND1, ND2, ATP6, COX1, COX2, and COX3 are particularly affected. Furthermore, our mitoribosome profiling data together with the structural analysis show that LRPPRC-SLIRP does not preexist on the mitoribosome as a structural element. Thus, the LRPPRC-SLIRP-dependent translation is not the sole regulatory pathway, and other mechanisms involving mRNA binding are likely to co-exist.

Since mS39 and mS31 are specific to Metazoa, as well as LRPPRC-SLIRP, the proposed mechanism in which some of the mitochondrial mRNAs are recruited for translation has developed in a co-evolutionary manner in Metazoa. However, also the presence of large RNA-binding moieties was also reported in association with mitoribosomes in other species [56-60]. Therefore, the principle of regulation by facilitation of molecular coupling might be a general feature, while unique molecular connectors involved in different species.

Overall, these findings define LRPPRC-SLIRP as a regulator of mitochondrial gene expression
and explain how its components modulate the function of translation via mRNA binding. Given
the challenge of studying mitochondrial translation due to the lack of an *in vitro* system, the
native structures are crucial for explaining fundamental mechanisms. The identification of the
components involved enhances our understanding of mitochondrial translation. Together, these
studies provide the structural basis for translation regulation and activation in mitochondria.
350

References

1.   Brown, A., Rathore, S., Kimanius, D., Aibara, S., Bai, X.-chen, Rorbach, J., Amunts, A., &
Ramakrishnan, V. Structures of the human mitochondrial ribosome in native states of
Assembly. *Nature Structural & Molecular Biology*, **24**(10), 866–869 (2017).

2.   Sissler, M., & Hashem, Y. Mitoribosome assembly comes into view. *Nature Structural &*
Molecular Biology, **28**(8), 631–633 (2021).

3.   Itoh, Y. *et al.* Mechanism of mitoribosomal small subunit biogenesis and preinitiation.
Nature **606,** 603–608 (2022).

4.   Lavdovskaia, E., Hillen, H. S., & Richter-Dennerlein, R. Hierarchical folding of the catalytic
core during mitochondrial ribosome biogenesis. *Trends in Cell Biology*, **32**(3), 182-185
(2022).

5.   Harper, N.J., Burnside, C. & Klinge, S. Principles of mitoribosomal small subunit assembly
in eukaryotes. *Nature* **614**, 175–181 (2023).

6.   Khawaja, A. *et al.* Distinct pre-initiation steps in human mitochondrial translation. *Nat*
Commun **11**, 2932, (2020).

7.   Rackham, O., Filipovska, A. Organization and expression of the mammalian mitochondrial
genome. *Nat Rev Genet*. **10**, 606-623 (2022).

8.   Webster, M. W. *et al.* Structural basis of transcription-translation coupling and collision in
bacteria. *Science* **369,** 1355–1359 (2020).

9.   Wang, C. *et al.* Structural basis of transcription-translation coupling. *Science* **369,** 1359–
1365 (2020).

10. OʼReilly, F. J. *et al.* In-cell architecture of an actively transcribing-translating expressome.
Science **369,** 554–557 (2020).

11. Itoh, Y. *et al.* Mechanism of membrane-tethered mitochondrial protein synthesis. *Science*
371, 846–849 (2021).

12. Ott, M., Amunts, A. & Brown, A. Organization and regulation of mitochondrial protein
synthesis. *Annual Review of Biochemistry* **85,** 77–101 (2016).

13. Christian, B. and Spremulli, L. Preferential selection of the 5′-terminal start codon on
leaderless mRNAs by mammalian mitochondrial ribosomes *J Biol Chem* **285**, 28379-28386
(2010).

14. Yi, SH. *et al.* Conformational rearrangements upon start codon recognition in human 48S
translation initiation complex. *Nucleic Acid Research* **50**, 5282–5298 (2022).

15. Lapointe, C.P. *et al.* eIF5B and eIF1A reorient initiator tRNA to allow ribosomal subunit
joining. *Nature*, **607** (7917) 185-190 (2022).

16. Ruzzenente, B. *et al.* LRPPRC is necessary for polyadenylation and coordination of
translation of mitochondrial mrnas. *The EMBO Journal* **31,** 443–456 (2012).

17. Siira, S. J. *et al.* LRPPRC-mediated folding of the mitochondrial transcriptome. *Nature*
Communications **8,** (2017).

18. Chujo, T. *et al.* LRPPRC/SLIRP suppresses pnpase-mediated mrna decay and promotes
polyadenylation in human mitochondria. *Nucleic Acids Research* **40,** 8033–8047 (2012).

19. Sasarman, F., Brunel-Guitton, C., Antonicka, H., Wai, T. & Shoubridge, E. A. LRPPRC and
SLIRP interact in a ribonucleoprotein complex that regulates posttranscriptional gene
expression in mitochondria. *Molecular Biology of the Cell* **21,** 1315–1323 (2010).

20. Mootha, V. K. *et al.* Identification of a gene causing human cytochrome *c* oxidase
deficiency by Integrative Genomics. *Proceedings of the National Academy of Sciences* **100,**
605–610 (2003).

21. Antonicka, H. *et al*. A high-density human mitochondrial proximity interaction network.
Cell Metab **32**, 479-497 (2020).

22. Sasarman, F., Brunel-Guitton, C., Antonicka, H., Wai, T. & Shoubridge, E.A. LRPPRC and
SLIRP interact in a ribonucleoprotein complex that regulates posttranscriptional gene
expression in mitochondria. *Mol. Biol. Cell* **21**, 1315-1323. (2010).

23. Lagouge, M. *et al.* SLIRP regulates the rate of mitochondrial protein synthesis and protects
LRPPRC from degradation. *PLOS Genetics* **11,** (2015).

24. Baughman, J. M. *et al.* A computational screen for regulators of oxidative phosphorylation
implicates SLIRP in mitochondrial RNA homeostasis. *PLoS Genetics* **5,** (2009).

25. Guo, L. *et al.* Pathogenic SLIRP variants as a novel cause of autosomal recessive
mitochondrial encephalomyopathy with complex I and IV deficiency. *European Journal of*
Human Genetics **29,** 1789–1795 (2021).

26. Sasarman, F. *et al.* Tissue-specific responses to the LRPPRC founder mutation in French
Canadian leigh syndrome. *Human Molecular Genetics* **24,** 480–491 (2014).

27. Xu, F., Addis, J. B. L., Cameron, J. M. & Robinson, B. H. LRPPRC mutation suppresses
cytochrome oxidase activity by altering mitochondrial RNA transcript stability in a mouse
model. *Biochemical Journal* **441,** 275–283 (2011).

28. Spåhr, H. *et al.* SLIRP stabilizes LRPPRC via an RRM–PPR protein interface. *Nucleic*
Acids Research **44,** 6868–6882 (2016).

29. Soto, I. *et al.* Balanced mitochondrial and cytosolic translatomes underlie the biogenesis of
human respiratory complexes. *Genome Biol.* **23,** 170 (2022).

30. Aibara, S., Singh, V., Modelska, A. & Amunts, A. Structural basis of mitochondrial
translation. *eLife* **9,** (2020).

31. Amunts, A., Brown, A., Toots, J., Scheres, S. H. & Ramakrishnan, V. The structure of the
human mitochondrial ribosome. *Science* **348,** 95–98 (2015).

32. Greber, B. J. *et al.* The complete structure of the 55S mammalian mitochondrial ribosome.
Science **348,** 303–308 (2015).

33. Itoh, Y., *et al.* Structure of the mitoribosomal small subunit with streptomycin reveals Fe-S
clusters and physiological molecules *eLife* 11:e77460 (2022).

34. Jumper J, *et al.* Highly accurate protein structure prediction with Alphafold. *Nature* **596,**
583–589 (2021).

35. Painter, Jay, and Ethan A. Merritt. "Optimal description of a protein structure in terms of
multiple groups undergoing TLS motion." *Acta Crystallographica Section D: Biological*
Crystallography **62,** 439-450 (2006).

36. Painter, Jay, and Ethan A. Merritt. "TLSMD web server for the generation of multi-group
TLS models." *Journal of applied crystallography* **39**, 109-111 (2006).

37. Coquille, S. & Thore, S. Leigh syndrome-inducing mutations affect LRPPRC / SLIRP
complex formation. *bioRxiv* https://doi.org/10.1101/2020.04.16.044412 (2020).

38. Burd, C. G., & Dreyfuss, G. Conserved structures and diversity of functions of RNA-
binding proteins. *Science*, **265,** 615-621 (1994).

39. Hatchell, E. C., Colley, S. M., Beveridge, D. J., Epis, M. R., Stuart, L. M., Giles, K. M., ... &
Leedman, P. J. SLIRP, a small SRA binding protein, is a nuclear receptor corepressor.
Molecular cell, **22,** 657-668 (2006).

40. Johansson, C., Finger, L. D., Trantirek, L., Mueller, T. D., Kim, S., Laird-Offringa, I. A., &
Feigon, J. Solution structure of the complex formed by the two N-terminal RNA-binding
domains of nucleolin and a pre-rRNA target. *Journal of molecular biology*, **337,** 799-816
(2004).

41. Williams, P., Li, L., Dong, X., & Wang, Y. Identification of SLIRP as a G quadruplex-
binding protein. *Journal of the American Chemical Society*, **139,** 12426-12429 (2017).

42. Deo, R. C., Bonanno, J. B., Sonenberg, N., & Burley, S. K. Recognition of polyadenylate
RNA by the poly (A)-binding protein. *Cell*, **98,** 835-845 (1999).

43. Wu, B., Su, S., Patil, D. P., Liu, H., Gan, J., Jaffrey, S. R., & Ma, J. Molecular basis for the
specific and multivariant recognitions of RNA substrates by human hnRNP A2/B1. *Nature*
communications, **9,** 1-12 (2018).

44. Takyar, S., Hickerson, R. P., & Noller, H. F. mRNA helicase activity of the ribosome. *Cell*,
120, 49–58 (2005).

45. Umate, P., Tuteja, N., & Tuteja, R. Genome-wide comprehensive analysis of human
helicases. *Communicative & Integrative Biology*, **4**, 118–137 (2011).

46. Kummer E, Leibundgut M, Rackham O, Lee RG, Boehringer D, Filipovska A & Ban N.
Unique features of mammalian mitochondrial translation initiation revealed by cryo-EM
Nature **560**, 263-267 (2018).

47. Xu, F., Morin, C., Mitchell, G., Ackerley, C. & Robinson, B.H. The role of the *LRPPRC*
(leucine-rich pentatricopeptide repeat cassette) gene in cytochrome oxidase assembly:
mutation causes lowered levels of COX (cytochrome c oxidase) I and *COX III* mRNA.
Biochem J. **382**, 331-336. (2004).

48. Ruzzenente, B. *et al.* LRPPRC is necessary for polyadenylation and coordination of
translation of mitochondrial mRNAs. *EMBO J.* **31**, 443-456 (2012).

49. Couvillion, M.T., Soto, I.C., Shipkovenska, G. & Churchman, L.S. Synchronized
mitochondrial and cytosolic translation programs. *Nature* **533**, 499-503 (2016).

50. Ingolia, N.T., Ghaemmaghami, S., Newman, J.R. & Weissman, J.S. Genome-wide analysis
in vivo of translation with nucleotide resolution using ribosome profiling. *Science* **324**, 218-
223 (2009).

51. Petrov, A. S. *et al.* Structural patching fosters divergence of mitochondrial ribosomes.
Molecular Biology and Evolution **36,** 207–219 (2018).

52. Huerta-Cepas, J. *et al.* Eggnog 5.0: A hierarchical, functionally and phylogenetically
annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids*
Research **47,** (2018).

53. Sterky, F. H., Ruzzenente, B., Gustafsson, C. M., Samuelsson, T. & Larsson, N.-G.
LRPPRC is a mitochondrial matrix protein that is conserved in metazoans. *Biochemical and*
Biophysical Research Communications **398,** 759–764 (2010).

54. Eddy, S. R. Accelerated profile HMM searches. *PLoS Computational Biology* **7,** (2011).

55. Erwin, D. H. Early metazoan life: divergence, environment and ecology. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, **370,** 1-7 (2015).

56. Amunts, A. *et al.* Structure of the yeast mitochondrial large ribosomal subunit. *Science* **343,**
1485–1489 (2014).

57. Tobiasson, V. & Amunts, A. Ciliate mitoribosome illuminates evolutionary steps of
mitochondrial translation. *eLife* **9,** (2020).

58. Waltz, F., Soufari, H., Bochler, A., Giegé, P. & Hashem, Y. Cryo-EM structure of the RNA-
rich plant mitochondrial ribosome. *Nature Plants* **6,** 377–383 (2020).

59. Waltz, F. *et al.* How to build a ribosome from RNA fragments in Chlamydomonas
mitochondria. *Nature Communications* **12,** (2021).

60. Tobiasson, V., Berzina, I. & Amunts, A. Structure of a mitochondrial ribosome with
fragmented rRNA in complex with membrane-targeting elements. *Nature Communications*
13, 6132 (2022).

Methods

Experimental model and culturing

HEK293S-derived cells (T501) were grown in Freestyle 293 Expression Medium containing 5% tetracycline-free fetal bovine serum (FBS) in vented shaking flasks at 37°C, 5% CO_2 and 120 rpm (550 x g). Culture was scaled up sequentially, by inoculating at 1.5 x 10^6 cells/mL and subsequently splitting at a cell density of 3.0 x 10^6 cells/mL. Finally, a final volume of 2 L of cell culture at a cell density of 4.5 x 10^6 cells/mL was used for mitochondria isolation, as previously described [61].

Mitoribosome purification

HEK293S-derived cells were harvested from the 2 L culture when the cell density was 4.2 x 10^6 cells/mL by centrifugation at 1,000 g for 7 min, 4 °C. The pellet was washed and resuspended in 200 mL Phosphate Buffered Saline (PBS). The washed cells were pelleted at 1,000 g for 10 min at 4 °C. The resulting pellet was resuspended in 120 mL of MIB buffer (50 mM HEPES-KOH, pH 7.5, 10 mM KCl, 1.5 mM $MgCl_2$, 1 mM EDTA, 1 mM EGTA, 1 mM dithiothreitol, cOmplete EDTA-free protease inhibitor cocktail (Roche) and allowed to swell in the buffer for 15 min in the cold room by gentle stirring. About 45 mL of SM4 buffer (840 mM mannitol, 280 mM sucrose, 50 mM HEPES-KOH, pH 7.5, 10 mM KCl, 1.5 mM $MgCl_2$, 1 mM EDTA, 1 mM EGTA, 1 mM DTT, 1X cOmplete EDTA-free protease inhibitor cocktail (Roche) was added to the cells in being stirred in MIB buffer and poured into a nitrogen cavitation device kept on ice. The cells were subjected to a pressure of 500 psi for 20 min before releasing the nitrogen from the chamber and collecting the lysate. The lysate was clarified by centrifugation at 800 x g and 4 °C, for 15 min, to separate the cell debris and nuclei. The supernatant was passed through a cheesecloth into a beaker kept on ice. The pellet was resuspended in half the previous volume of MIBSM buffer (3 volumes MIB buffer + 1 volume SM4 buffer) and homogenized with a Teflon/glass Dounce homogenizer. After clarification as described before, the resulting lysate was pooled with the previous batch of the lysate and subjected to centrifugation at 1,000 x g, 4 °C for 15 minutes to ensure complete removal of cell debris. The clarified and filtered supernatant was centrifuged at 10,000 x g and 4 °C for 15 min to pellet crude mitochondria. Crude mitochondria were resuspended in 10 mL MIBSM buffer and treated with 200 units of

Rnase-free Dnase (Sigma-Aldrich) for 20 min in the cold room to remove contaminating genomic DNA. Crude mitochondria were again recovered by centrifugation at 10,000 g, 4 °C for 15 min and gently resuspended in 2 mL SEM buffer (250 mM sucrose, 20 mM HEPES-KOH, pH 7.5, 1 mM EDTA). Resuspended mitochondria were subjected to a sucrose density step-gradient (1.5 mL of 60% sucrose; 4 mL of 32% sucrose; 1.5 mL of 23% sucrose and 1.5 mL of 15% sucrose in 20 mM HEPES-KOH, pH 7.5, 1 mM EDTA) centrifugation in a Beckmann Coulter SW40 rotor at 28,000 rpm (139,000 x g) for 60 min. Mitochondria seen as a brown band at the interface of 32% and 60% sucrose layers were collected and snap-frozen using liquid nitrogen and transferred to −80 °C.

Frozen mitochondria were transferred on ice and allowed to thaw slowly. Lysis buffer (25 mM HEPES-KOH, pH 7.5, 50 mM KCl, 10 mM Mg(Oac)$_2$, 2% polyethylene glycol octylphenyl ether, 2 mM DTT, 1 mg/mL EDTA-free protease inhibitors (Sigma-Aldrich) was added to mitochondria and the tube was inverted several times to ensure mixing. A small Teflon/glass Dounce homogenizer was used to homogenize mitochondria for efficient lysis. After incubation on ice for 5-10 min, the lysate was clarified by centrifugation at 30,000 x g for 20 min, 4 °C. The clarified lysate was carefully collected. Centrifugation was repeated to ensure complete clarification. A volume of 1 mL of the mitochondrial lysate was applied on top of 0.4 mL of 1 M sucrose (v/v ratio of 2.5:1) in thick-walled TLS55 tubes. Centrifugation was carried out at 231,500 x g for 45 min in a TLA120.2 rotor at 4 °C. The pellets thus obtained were washed and sequentially resuspended in a total volume of 100 μl resuspension buffer (20 mM HEPES-KOH, pH 7.5, 50 mM KCl, 10 mM Mg(Oac)$_2$, 1% Triton X-100, 2 mM DTT). The sample was clarified twice by centrifugation at 18,000 g for 10 min at 4 °C. The sample was applied on to a linear 15-30% sucrose (20 mM HEPES-KOH, pH 7.5, 50 mM KCl, 10 mM Mg(Oac)$_2$, 0.05% n-dodecyl-β-D-maltopyranoside, 2 mM DTT) gradient and centrifuged in a TLS55 rotor at 213,600 x g for 120 min at 4 °C. The gradient was fractionated into 50 μL volume aliquots. The absorption for each aliquot at 260 nm was measured and fractions corresponding to the monosome peak were collected. The pooled fractions were subjected to buffer exchange with the resuspension buffer.

Cryo-EM data acquisition

3 μL of ~120 nM mitoribosome was applied onto a glow-discharged (20 mA for 30 sec) holey-

carbon grid (Quantifoil R2/2, copper, mesh 300) coated with continuous carbon (of ~3 nm

thickness) and incubated for 30 sec in a controlled environment of 100% humidity and 4 ºC. The

grids were blotted for 3 sec, followed by plunge-freezing in liquid ethane, using a Vitrobot

MKIV (ThermoFischer). The data were collected on FEI Titan Krios (ThermoFischer)

transmission electron microscope operated at 300 keV, using C2 aperture of 70 μm and a slit

width of 20 eV on a GIF quantum energy filter (Gatan). A K2 Summit detector (Gatan) was used

at a pixel size of 0.83 Å (magnification of 165,000X) with a dose of 29-32 electrons/$Å^2$

fractionated over 20 frames.

Cryo-EM data processing

The beam-induced motion correction and per-frame B-factor weighting were performed using

RELION-3.0.2 [62,63]. Motion-corrected micrographs were used for contrast transfer function

(CTF) estimation with gctf [64]. Unusable micrographs were removed by manual inspection of the

micrographs and their respective calculated CTF parameters. Particles were picked in RELION-

3.0.2, using reference-free followed by reference-aided particle picking procedures. Reference-

free 2D classification was carried out to sort useful particles from falsely picked objects, which

were then subjected to 3D classification. 3D classes corresponding to unaligned particles and

LSU were discarded, and monosome particles were pooled and used for 3D auto-refinement

yielding a map with an overall resolution of 2.9-3.4 Å for the five datasets. Resolution was

estimated using a Fourier Shell Correlation cut-off of 0.143 between the two reconstructed half

maps. Finally, the selected particles were subjected to per-particle defocus estimation, beam-tilt

correction, and per-particle astigmatism correction followed by Bayesian polishing. Bayesian

polished particles were subjected to a second round per-particle defocus correction. A total of

994,919 particles were pooled and separated into 86 optics groups in RELION-3.1 [65], based on

acquisition areas and date of data collection. Beam-tilt, magnification anisotropy, and higher-

order (trefoil and fourth-order) aberrations were corrected in RELION-3.1 [65]. Particles with A-

site and/or P-site occupied with tRNAs which showed comparatively higher occupancy for the

unmodeled density potentially corresponding to the LRPPRC-SLIRP module, were pooled and

re-extracted in a larger box size of 640 Å. The re-extracted particles were subjected to 3D

autorefinement in RELION3.1 [65]. This was followed by sequential signal subtraction to remove

signal from the LSU, and all of the SSU except the region around mS39 and the unmodeled

density, in that order. The subtracted data was subjected to masked 3D classification (T=200) to

enrich for particles carrying the unmodeled density. Using a binary mask covering mS39 and all

of the unmodeled density, we performed local-masked refinement on resulting 41,812 particles

within an extracted sub-volume of 240 Å box size leading to 3.37 Å resolution map.

<u>Model building and refinement</u>

At the mRNA channel entrance, a more accurate and complete model of mS39 could be built

with 29 residues added to the structure. Improved local resolution enabled unambiguous

assignment of residues to the density which allowed us to address errors in the previous model.

A total of 28 α-helices could be modeled in their correct register and orientation. Further, a 28-

residue long N-terminal loop of mS31 (residues 247–275) along mS39 and mito-specific N-

terminal extension of uS9m (residues 53–70) approaching mRNA were modeled by fitting the

loops in to the density maps.

For building LRPPRC-SLIRP module, the initial model of the full length LRPPRC was obtained

from *AlphaFold2* Protein Structure Database (Uniprot ID P42704). Based on the analysis, three

stable domains were identified that are connected by flexible linkers (673-983, 1035-1390). We

then systematically assessed the domains against the map, and the N-terminal region (77-660)

could be fitted into the density. The initial model was real space refined into the 3.37 Å

resolution map of mS39-LRPPRC-SLIRP region obtained after partial signal subtraction using

reference-restraints in *Coot* v0.9 [66]. The N-terminal region covering residues 64-76 was

identified in the density map and allowed us to model 34 helices of LRPPRC (residues 64-644).

Helices α1-29 could be confidently modeled. An additional five helices, as predicted by

Alphafold2 [34], could be accommodated into the remaining density. After modelling LRPPRC into

the map, there was an unaccounted density that fits SLIRP. The initial model of SLIRP was

obtained from *Alphafold2* Protein Structure Database (Uniprot ID Q9GZT3). The unmodeled

density agreed with the secondary structure of SLIRP. The model was real space refined into the

density using reference-restraints as was done for LRPPRC in *Coot* v0.9 [66]. Five additional RNA

residues could be added to the 3' terminal of mRNA to account for tubular density extending

from it along the mRNA binding platform. The A/A P/P E/E state model was rigid body fitted

into the corresponding 2.85 Å resolution consensus map. Modeled LRPPRC was merged with

the rigid body fitted monosome model to obtain a single model of the mitoribosome bound to

LRPPRC and SLIRP. The model was then refined against the composite map using PHENIX

v1.18 [67] (Supplementary Information Table 1).

<u>Phylogenetic analysis</u>

Phylogenetic distribution of proteins was determined by examining phylogeny databases [55],

followed by sensitive homology detection to detect homologs outside of the bilateria. Orthologs

were required to have identical domain compositions, and Dollo parsimony was used to infer the

evolutionary origin of a protein from its phylogenetic distribution. When multiple homologs of a

protein were detected in a species, a neighbor-joining phylogeny was constructed to assess

monophyly of putative orthologs to the human protein. The short length of the SLIRP candidate

protein from *T. adhaerens* (B3SAC0_TRIAD), that is part of the large RRM family, precludes

obtaining a reliable phylogeny to confidently assess its orthology to human SLIRP is therefore

tentative.

<u>TLSMD analysis</u>

The TLSMD analysis [35,36] was performed with full length LRPPRC model obtained from

AlphaFold Database (AF-P42704-F1), and mitochondrial targeting sequence (residues 1-59) was

removed. The model was divided into TLS segments (N), and single chain TLSMD is performed

on all atoms using the isotropic analysis model. Instead of using atomic B-factors, the values for

a per residue confidence score of AlphaFold called predicted local distance difference test

(pLDDT) were used as reference to calculate the least squared residuals against the

corresponding values calculated by TLSMD analysis. This is based on the assumption that local

mobility of the model should be inversely correlated with the pLDDT score. AlphaFold pLDDT

values and the corresponding calculated values were plotted for every iteration to monitor

improvement in prediction and across the length of LRPPRC. The data in Extended Data Fig. 2

is presented for N=4, where segments 1 and 2 (residues 60-373 and 374-649) correspond to the

modeled region, whereas segments 3 and 4 correspond to the remaining domains that could not

be modeled.

Helicase sequence analysis

To address the possibility that LRPPRC may serve as a helicase, we inspected the sequence of full-length LRPPRC (Uniprot ID P42704). First, we checked the sequence for matches with consensus motifs characteristic of helicases using regular expression search. The following motifs were searched, GFxxPxxIQ, AxxGxGKT, PTRELA, TPGR, DExD, SAT, FVxT, RgxD (DDX helicases); GxxGxGKT, TQPRRV, TDGML, DExH, SAT, FLTG, TNIAET, QrxGRAGR (DHX helicases); AHTSAGKT, TSPIKALSNQ, MTTEIL (others). Next, we carried out multiple sequence analysis against representative member helicases of the DHX and DDX family to verify the results of the regular-expression sequence search and to find potentially valid weaker matches.

Human cell lines and cell culture conditions

Human HEK293T embryonic kidney cells (CRL-3216, RRID: CVCL-0063) were obtained from ATCC. The HEK293T *LRPPRC* knock-out (KO) cell line was engineered in-house and previously reported [27]. The LRPPRC-KO cell line was reconstituted with either the wild-type *LRPPRC* gene [27], or a variant causing Leigh syndrome, French-Canadian type (LSFC). The LSFC variant carries a single base change (nucleotide C1119T transition), predicting a missense A354V change at a conserved protein residue [46].

Cells were cultured in high-glucose Dulbecco's modified Eagle's medium (DMEM, Thermo Fisher Scientific, CAT 11965092), supplemented with 10% FBS (Thermo Fisher Scientific, CAT A3160402), 100 μg/mL of uridine (Sigma, CAT U3750), 3 mM sodium formate (Sigma CAT 247596), and 1 mM sodium pyruvate (Thermo Fisher Scientific, CAT 11360070) at 37 °C under 5% CO_2. Cell lines were routinely tested for mycoplasma contamination.

To generate an *LRPPRC*-KO cell line reconstituted with the LSFC variant of the gene, a Myc-DDK tagged *LRPPRC* ORF plasmid was obtained from OriGene (CAT: RC216747). This ORF was then subcloned into a hygromycin resistance-containing pCMV6 entry vector (OriGene, CAT PS100024) and used to generate an *LRPPRC*-KO cell line reconstituted with a wild-type *LRPPRC* gene as reported [27]. To generate the *LRPPRC*-LSFC variant carrying the C1119T mutation, we used the Q5® Site-Directed Mutagenesis Kit from NEB. ~ 10 pg of template pCMV6-A-Myc-DDK-Hygro-*LRPPRC* vector were used, along with the primers LSFC-Q5-F 5'

GGAAGATGTAGTGTTGCAGATTTTAC and LSFC-Q5-R 5'

AATTTTTCAGTGACTAAAAGTAAAATG, designed to include the codon to be mutated.

After exponential amplification and treatment with kinase and ligase, 2.5 µl of the reaction were

transformed into competent *Escherichia coli* cells. Several transformants were selected, their

plasmid DNA purified, and then sequenced to select the correct pCMV6-A-Myc-DDK-Hygro-

LRPPRC-LSFC construct.

For transfection of the construct into *LRPPRC*-KO cells, we used 5 µl of EndoFectin mixed with

2 µg of vector DNA in OptiMEM-I media according to the manufacturer's instructions. Media

was supplemented with 200 µg/ml of hygromycin after 48 h, and drug selection was maintained

for at least one month.

<u>Whole-Cell extracts and Mitochondria isolation</u>

For SDS-PAGE electrophoresis, pelleted cells were solubilized in RIPA buffer (25 mM Tris-HCl

pH 7.6, 150 mM NaCl, 1% NP-40, 1% sodium deoxycholate, and 0.1% SDS) with 1 mM PMSF

(phenylmethylsulfonyl fluoride) and mammalian protease inhibitor cocktail (Sigma). Whole-cell

extracts were cleared by 5 min centrifugation at 20,000 x g at 4 °C.

Mitochondria-enriched fractions were isolated from at least ten 80% confluent 15-cm plates as

described previously [68-70]. Briefly, the cells were resuspended in ice-cold T-K-Mg buffer (10 mM

Tris–HCl, 10 mM KCl, 0.15 mM $MgCl_2$, pH 7.0) and disrupted with 10 strokes in a homogenizer

(Kimble/Kontes, Vineland, NJ). Using a 1 M sucrose solution, the homogenate was brought to a

final concentration of 0.25 M sucrose. A post-nuclear supernatant was obtained by centrifugation

of the samples twice for 5 min at 1,000 x g. Mitochondria were pelleted by centrifugation for 10

min at 10,000 x g and resuspended in a solution of 0.25 M sucrose, 20 mM Tris-HCl, 40 mM

KCl, and 10 mM $MgCl_2$, pH 7.4.

<u>Denaturing and native electrophoresis, followed by immunoblotting</u>

Protein concentration was measured by the Lowry method [71]. 40–80 µg of mitochondrial protein

extract was separated by denaturing SDS–PAGE in the Laemmli buffer system [72]. Then, proteins

were transferred to nitrocellulose membranes and probed with specific primary antibodies

against the following proteins: β-ACTIN (dilution 1:2,000; Proteintech; Rosemont, IL; 60008-1-

Ig), ATP5A (1:1000; Abcam; Cambridge, MA; ab14748), CORE2 (1:1,000; Abcam; Cambridge,

MA; ab14745), COX1 (dilution 1:2,000; Abcam; Cambridge, MA; ab14705), LRPPRC (dilution

1:1,000; Proteintech; Rosemont, IL; 21175-1-AP), NDUFA9 (1:1000; Proteintech; Rosemont,

IL; 20312-1-AP), SDHA (1:1,000; Proteintech; Rosemont, IL; 14865-1-AP) or SLIRP (1:1000;

Abcam; Cambridge, MA; ab51523). Horseradish peroxidase-conjugated anti-mouse or anti-

rabbit IgGs were used as secondary antibodies (dilution 1:10,000; Rockland; Limerick, PA). β-

ACTIN was used as a loading control. Signals were detected by chemiluminescence incubation

and exposure to X-ray film.

Blue-native polyacrylamide gel electrophoresis (BN-PAGE) analysis of mitochondrial OXPHOS

complexes in native conditions was performed as described previously [73,74]. To extract

mitochondrial proteins in native conditions, we pelleted and solubilized 400 μg mitochondria in

100 μl buffer containing 1.5 M aminocaproic acid and 50 mM Bis-Tris (pH 7.0) with 1% n-

dodecyl b-D-maltoside (DDM). Solubilized samples were incubated on ice for 10 min in ice and

pelleted at 20,000 x g for 30 min at 4 °C. The supernatant was supplemented with 10 μl of

sample buffer 10X (750 mM aminocaproic acid, 50 mM Bis-Tris, 0.5 mM EDTA

(ethylenediaminetetraacetic acid), 5% Serva Blue G-250). Native PAGE™ Novex® 3-12% Bis-

Tris Protein Gels (Thermo Fisher) gels were loaded with 40 μg of mitochondrial proteins. After

electrophoresis, the gel was stained with 0.25% Coomassie brilliant blue R250, or proteins were

transferred to PVDF membranes using an eBlot L1 protein transfer system (GenScript,

Piscataway, NJ) and used for immunoblotting.

<u>Pulse Labeling of Mitochondrial Translation Products</u>

To determine mitochondrial protein synthesis, 6-well plates were pre-coated at 5 μg/cm^2 with 50

μg/mL collagen in 20 mM acetic acid and seeded with WT or LRPPRC cell lines (two wells per

sample per timepoint). 70% confluent cell cultures were incubated for 30 min in DMEM without

methionine and then supplemented with 100 μl/ml emetine for 10 min to inhibit cytoplasmic

protein synthesis as described [68]. 100 μCi of [^{35}S]-methionine was added and allowed to

incorporate to newly synthesized mitochondrial proteins for increasing times from 15- to 60-

minute pulses. Subsequently, whole-cell extracts were prepared by solubilization in RIPA buffer,

and equal amounts of total cellular protein were loaded in each lane and separated by SDS-

PAGE on a 17.5% polyacrylamide gel. Gels were transferred to a nitrocellulose membrane and

exposed to a Kodak X-OMAT X-ray film. The membranes were then probed with a primary

antibody against β-ACTIN as a loading control. Optical densities of the immunoreactive bands

were measured using the Histogram function of the Adobe Photoshop software in digitalized

images.

<u>Whole-cell transcriptomics</u>

Cells were grown to 80% confluency in a 10 cm plate (two plates per sample) and were collected

by trypsinization and washed once with PBS before resuspending in one mL of Trizol

(ThermoFisher Scientific). RNA was extracted following the Trizol manufacturer's

specifications. The aqueous phase was transferred to a new tube, and an equal volume of 100%

isopropanol and 3 μL of glycogen were added to precipitate the RNA. The sample was incubated

at -80 °C overnight and centrifuged at 15,000 xg for 45 min at 4°C. RNA was resuspended in 50

μL of RNAse-free water and quantified by measuring absorbance at a wavelength of 260 nm. 2

μg of RNA was sent to Novogene (Sacramento, CA) for further processing. Novogen services

included library preparation, RNA sequencing (RNAseq) on an Illumina HiSeq platform

according to the Illumina Tru-Seq protocol, and bioinformatics analysis. The raw data was

cleaned to remove low-quality reads and adapters using Novogen in-house Perl scripts in

Cutadapt [75]. The reads were mapped to the reference genome using the HISAT2 software [76]. The

transcripts were assembled and merged to obtain an mRNA expression profile with the StringTie

algorithm [77], the RNA-seq data was then normalized to account for the total reads sequenced for

each sample (the read depth), and differentially expressed mRNAs were identified by using the

Ballgown suite [78] and the DESeq2 R package [79]. GraphPad Prism version 9.0 software

(GraphPad Software, San Diego, CA, USA) was used to prepare the volcano plots.

<u>Mitoribosome profiling</u>

Mitoribosome profiling, matched RNA-seq, and data analysis were performed as described [29].

Briefly, human and mouse cell lysates were prepared and mixed 95:5 human:mouse. For

mitoribosome profiling, the combined lysates were subjected to RNaseI treatment and

fractionated across a linear sucrose gradient. Sequencing libraries were prepared from the

monosome fraction after phenol/chloroform extraction. For RNA-seq, RNA was extracted from

the undigested combined lysate, fragmented by alkaline hydrolysis, and sequencing libraries

prepared. Reads were cleaned of adapters, filtered of rRNA fragments, and PCR duplicates were

removed. Read counts were summed across features (coding sequences) using Rsubread feature

Counts [80], then normalized by feature length and mouse spike-in read counts. TE was calculated

by dividing spike-in normalized mitoRPF reads per kilobase by spike-in normalized RNA-seq

reads per kilobase. Values are expressed as $\log_2$-fold change in the *LRPPRC*-KO cells compared

to the *LRPPRC* rescue cells. Mitoribosome profiling and RNA-seq data for *LRPPRC*-KO and

LRPPRC-reconstituted cell lines are deposited in GEO under the accession number GSE173283.

Mitoribosome profiling and RNA-seq data for the LSFC-reconstituted is deposited in GEO under

the accession number GSEXXXXXX.

The mitoRPF length distribution was determined from mitochondrial mRNA-aligned reads. First,

soft-clipped bases were removed using jvarkit [81], then frequency for each length was output

using Samtools stats [82].

Data availability statement

The atomic coordinates were deposited in the RCSB Protein Data Bank, and EM maps have been

deposited in the Electron Microscopy Data bank under accession numbers 8ANY and EMD-

15544.

The atomic coordinates that were used in this study: 6ZTJ (*E.coli* 70S-RNAP expressome

complex in NusG); 6ZTN (*E.coli* 70S-RNAP expressome complex in NusG); 1RKJ (human

Nucleolin); 5WWE (human hnRNPA2/B1); 1CVJ (Poly-adenylate binding protein, PABP)

Additional References

61. Aibara, S., Andréll, J., Singh, V. & Amunts, A. Rapid isolation of the mitoribosome from
Hek Cells. *Journal of Visualized Experiments* (2018). doi:10.3791/57877.
62. Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure determination
in relion-3. *eLife* **7,** (2018).
63. Zivanov, J., Nakane, T. & Scheres, S. H. A bayesian approach to beam-induced motion
correction in Cryo-EM single-particle analysis. *IUCrJ* **6,** 5–17 (2019).
64. Zhang, K. GCTF: Real-time CTF determination and correction. *Journal of Structural
Biology* **193,** 1–12 (2016).
65. Zivanov, J., Nakane, T. & Scheres, S. H. Estimation of high-order aberrations and
anisotropic magnification from Cryo-EM data sets in *relion*-3.1. *IUCrJ* **7,** 253–267 (2020).
66. Emsley, P. & Cowtan, K. *Coot*: Model-building tools for Molecular Graphics. *Acta
Crystallographica Section D Biological Crystallography* **60,** 2126–2132 (2004).
67. Afonine, P. V. *et al.* Real-space refinement in *phenix* for cryo-EM and Crystallography. *Acta
Crystallographica Section D Structural Biology* **74,** 531–544 (2018).

68. Bourens, M., Boulet, A., Leary, S.C. & Barrientos, A. Human COX20 cooperates with
SCO1 and SCO2 to mature COX2 and promote the assembly of cytochrome *c* oxidase. *Hum.*
Mol. Genet. **23**, 2901-2913 (2014).
69. Fernandez-Vizarra, E. *et al.* Isolation of mitochondria for biogenetical studies: An update.
Mitochondrion **10**, 253-262 (2010).
70. Moreno-Lastres, D. *et al.* Mitochondrial Complex I Plays an Essential Role in Human
Respirasome Assembly. *Cell Metab.* **15**, 324-335 (2012).
71. Lowry, O.H., Rosebrough, N.J., Farr, A.L. & Randall, R.J. Protein measurement with the
Folin phenol reagent. *J. Biol. Chem.* **193**, 265-275 (1951).
72. Laemmli, U.K. Cleavage of structural proteins during the assembly of the head of
bacteriophage T4. *Nature* **227**, 680-685 (1970).
73. Diaz, F., Barrientos, A. & Fontanesi, F. Evaluation of the mitochondrial respiratory chain
and oxidative phosphorylation system using blue native gel electrophoresis. *Curr. Protoc.*
Hum. Genet. Chapter, Unit19.14. (2009).
74. Timón-Gómez, A. *et al.* Protocol for the analysis of yeast and human mitochondrial
respiratory chain complexes and supercomplexes by Blue Native-PAGE. *STAR Protocols* **1**,
100119 (2020).
75. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet J. **17**, 10-12 (2021).
76. Kim, D., Paggi, J.M., Park, C., Bennett, C. & Salzberg, S.L. Graph-based genome alignment
and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907-915 (2019).
77. Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from RNA-
seq reads. *Nat. Biotechnol.* **33**, 290-295 (2015).
78. Frazee, A.C. *et al.* Ballgown bridges the gap between transcriptome assembly and
expression analysis. *Nat. Biotechnol.* **33**, 243-246 (2015).
79. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).
80. Liao, Y., Smyth, G.K. & Shi, W. The R package Rsubread is easier, faster, cheaper and
better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Res.* **47**, e47
(2019).
81. Lindenbaum, P. & Redon, R. bioalcidae, samjs and vcffilterjs: object-oriented formatters
and filters for bioinformatics files. *Bioinformatics* **34**, 1224-1225 (2018).
82. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10** (2021).
835

Acknowledgments

We thank S. Aibara and J. Andrell for help with data collection. The research was funded by the

Swedish Foundation for Strategic Research (FFL15:0325), Ragnar Söderberg Foundation

(M44/16), European Research Council (ERC-2018-StG-805230) , Knut and Alice Wallenberg

Foundation (2018.0080) , NIH R01 (R01-GM123002) to SC, NIH R35 (R35-GM118141) to AB.

V.S. was supported by the Horizon 2020 - Marie Sklodowska-Curie Innovative Training

Network (721757), Y.I. was supported by H2020-MSCA-IF-2017 (799399-Itohribo), and C.M

was supported by the Eunice Kennedy Shriver National Institute Of Child Health & Human

Development of the National Institutes of Health under Award Number F30HD107939. The

SciLifeLab cryo-EM facility is funded by the Knut and Alice Wallenberg, Family Erling

Persson, and Kempe foundations. The content is solely the responsibility of the authors and does

not necessarily represent the official views of the National Institutes of Health.

849 Author contributions

850 V.S. collected cryo-EM data, processed the data, and built the models. V.S., Y.I. and A.A.

851 performed structural analysis. C.M., F.F. and A.B. performed mitochondrial translation,

852 OXPHOS, and RNAseq analysis. I.S., M.C. and S.C. performed mitoribosome profiling and

853 RNAseq analysis. V.S., M.H. and A.A. performed evolutionary analysis. A.A. wrote the

854 manuscript. All authors contributed to data interpretation and manuscript writing.

856 Competing Interests Statement

857 The authors declare no competing interests.

Extended Data Fig. 1. Cryo-EM data processing and map for mS39-LRPPRC-SLIRP region.
a. Focused 3D-classification with signal subtraction using mask around mS39-LRPPRC-SLIRP
region (transparent orange) of mitoribosome particles to identify LRPPRC-SLIRP containing
monosome particles (2.86 Å overall resolution), followed by, masked refinement with signal
subtraction on mS39-LRPPRC-SLIRP region to improve the local resolution. **b.** The mS39-
LRPPRC-SLIRP map is shown colored by local resolution (top left) and by proteins assigned to the
density (top right). The consensus map (bottom left) and the masked refined maps shown as a single
composite map colored by local resolution (bottom right). **c.** Fourier shell correlation curves for the
post-subtraction masked refined mS39-LRPPRC-SLIRP map (top) and individual masked refined
maps. **d.** Map comparison for LRPPRC region between our work and EMD-11397. The map has
been Gaussian filtered for better visibility. **e.** Density shown as mesh around helices α5-6, 8 and 11-
12. Corresponding regions are indicated with arrows in panel (d).

Extended Data Fig. 2. *AlphaFold* **model and TLSMD analysis of LRPPRC.**

a. The modeled region of LRPPRC (residues 64-644) is compared with the AlphaFold model (AF-P42704-F1) of full length (right). The modeled region is green, the unmodeled is white. The position of LSFC variant (A354V) is indicated. **b.** TLSMD analysis of the AlphaFold model of LRPPRC up to 20 TLS segments (N). Graph plots least-square residuals assigned per-residue confidence score values (pLDDT) versus those calculated by TLS analysis. **c.** Model colored by TLS segments for N=4. Regions between the segments with high pLDDT values correspond to loop regions and are shown as spheres **d.** Comparison of AlphaFold assigned versus calculated pLDDT values at N=4.

Extended Data Fig. 3. Multiple sequence alignment between SLIRP and representative RRM containing proteins. Alignment of SLIRP with representative RRM family proteins, heterogeneous nuclear ribnucleoproteins (hnRNPA2/B1), poly-A binding protein (PABP), and nucleolin shows conservation of submotifs RNP1 and RNP2 highlighted and indicated by corresponding residue numbers in SLIRP. Individual sequences are marked by residue numbers in the beginning and end and residues are colored by present identity.

Extended Data Fig. 4. Reconstitution of the *LRPPRC*-KO with wild-type and LSCF variants of LRPPRC. Immunoblot analysis to estimate the steady-state levels of LRPPRC and SLIRP in the indicated cell lines. β-ACTIN was used as a loading control. The images were digitized, and the specific signals were quantified using the histogram function of Adobe Photoshop from three independent repetitions.

Extended Data Fig. 5. Mitochondrial protein synthesis is altered in *LRPPRC*-KO cells. Blue-native PAGE analyses in WT, *LRPPRC*-KO, and KO+WT cell lines. Intact respiratory complexes were extracted from purified mitochondria using 1% n-dodecyl β-D-maltoside. An asterisk indicates the ATPase **(**CV) F_1 module that accumulates due to the low levels of the mitochondrion-encoded F_O module subunits ATP6 and ATP8.

Extended Data Fig. 6. LRPPRC-SLIRP contacts with the SSU head. a. Comparison of SSU from mitoriboome:LRPPRC-SLIRP complex with SSU from E-site tRNA bound monosome. Zoom-in shows N-terminal region of mS31 and C-terminal loop of mS39 (in surface) stabilized by LRPPRC. **b.** Contact regions of LRPPRC with mS31 and mS39 shown in cartoon and surface representations.

Extended Data Fig. 7. Close-up view of uS10m interactions with mS31-mS39.

Interface between uS10m with mS31-mS39 that serve as the platform for LRPPRC-SLIRP is similar to that formed between uS10 and NusG that binds RNA polymerase in bacterial expressome [8].

Structural basis of LRPPRC-SLIRP-dependent translation by the mitoribosome

Vivek Singh[1,7], J. Conor Moran[2,7], Yuzuru Itoh[1,6,*], Iliana C. Soto[3], Flavia Fontanesi[2], Mary Couvillion[3], Martijn A. Huynen[4], Stirling Churchman[3,*], Antoni Barrientos[5,*], Alexey Amunts[1,*]

[1] Science for Life Laboratory, Department of Biochemistry and Biophysics, Stockholm University, 17165 Solna, Sweden.

[2] Department of Biochemistry and Molecular Biology, University of Miami Miller School of Medicine, Miami, FL, 33136, USA

[3] Blavatnik Institute, Department of Genetics, Harvard Medical School, Boston, MA, 02115, USA.

[4] Centre for Molecular and Biomolecular Informatics, Radboud Institute for Molecular Life Sciences, Radboud University Medical Center, Nijmegen, the Netherlands.

[5] Department of Neurology, University of Miami Miller School of Medicine, Miami, FL, 33136, USA.

[6] Current position: Department of Biological Sciences, Graduate School of Science, The University of Tokyo, 113-0033 Tokyo, Japan.

[7] These authors contributed equally

* correspondence: yuzuru.itoh@bs.s.u-tokyo.ac.jp, churchman@genetics.med.harvard.edu, abarrientos@med.miami.edu, amunts@scilifelab.se

SUPPLEMENTARY INFORMATION

Supplementary table 1. Data collection and model statistics.

	Monosome with LRPPRC:SLIRP PDB: 8ANY EMD-15544
Data collection and processing	
Electron microscope	Titan Krios
Camera	K2 Summit (counting mode)
Magnification	165,000
Voltage (kV)	300
Electron exposure ($e^-/\text{Å}^2$)	29-32
No. of frames	20
Defocus range (μm)	-0.6 to -2.8
Pixel size (Å)	0.83
Symmetry imposed	C_1
Final particle number (no.)	41,815
Map resolution (Å) (Overall/ LSU-body/ CP/ L10-L12-stalk/ L1-stalk/ SSU-body/ SSU-head/ mS39/ SSU-tail/ mS39-LRPPRC-SLIRP)	2.85/2.69/ 3.07/ 3.07/ −/ 2.89/ 2.84/−/ 3.02/3.37
FSC threshold	0.143
Map resolution range (Å)	2.5-8.0
Refinement	
Initial model used (PDB code)	6ZSG, 6RW4
Model resolution (Å)	2.7
Model to map CC (CC_{volume})	0.84
FSC threshold	0.5
Map-sharpening B factor (Å^2) (Overall/ LSU-body/ CP/ L10-L12-stalk/ L1-stalk/ SSU-body/ SSU-head/ mS39/	-33/ -28/ -58/ -49/−/ -39/ -39/−/-55/ -60
Model composition	
Non-hydrogen atoms	356138
Hydrogen atoms	160529
Protein chains	90
RNA chains	7
Protein residues (non-modified/ N-acetylAla/ N-acetylSer/ N-acetylThr O^1-methylisoAsp)	15624/3/1/1/1
RNA residues (non-modified/ mG/ mU/ m^1A/ m^2G/ψ /m^4C/ m^5C/ m^5U/ m^6_2A)	2822/2/ 1/ 2/ 1/ 2/ 1/ 1/ 1/ 2
Ligands (ATP/ GDP/ NAD/ 2Fe-2S/ spermine/ spermidine/ putrescine)	1/ 1/ 1/ 3 /1 /4/ 1
Ions (Zn^{2+}/ K^+/ Mg^{2+})	3/ 49/ 206
Waters	6,926
Mean atomic B-factor (Å^2)	
Protein	38.28
RNA	36.42
Ligand	18.47
Water	15.21
Validation	
Ramanchandran plot (%)	
Outliers	0.05
Allowed	1.85
Favored	98.11
Clash score	2.64
RMSD	
Bonds (Å)	0.002
Angles (°)	0.432
Rotamer outliers (%)	0.00
C_β outliers (%)	0.00
CaBLAM outliers (%)	0.90

Supplementary Video 1. Structure of LRPPRC-SLIRP module bound to monosome.

The video shows the structure of LRPPRC-SLIRP determined in this work and how docking of mRNA on SSU is achieved by LRPPRC-SLIRP together with mito-specific proteins mS31 and mS39.