Past, present, and future of the Living Planet Index

Sophie E H Ledger[1*], Louise McRae[1*], Jonathan Loh[2], Rosamunde Almond[3], Monika Böhm[4], Christopher F Clements[5], Jessica Currie[6], Stefanie Deinet[1], Thomas Galewski[7], Monique Grooten[3], Martin Jenkins, Valentina Marconi[1], Brett Painter[8], Kate Scott-Gatty[1], Lucy Young[9], Michael Hoffmann[10] and Robin Freeman[1]

[1] Institute of Zoology, Zoological Society of London (ZSL), London, UK

[2] School of Anthropology and Conservation, University of Kent, Canterbury, UK

[3] WWF Netherlands – World Wide Fund for Nature, Zeist, Netherlands

[4] Global Center for Species Survival, Indianapolis Zoo, Indianapolis, USA

[5] School of Biological Sciences, University of Bristol, Bristol, UK

[6] WWF Canada – World Wildlife Fund, Toronto, Canada

[7] Institut de recherche pour la conservation des zones humides méditerranéennes, Tour du Valat, Arles, France

[8] Environment and Climate Change Canada (ECCC), Government of Canada, Gatineau, Canada

[9] WWF UK – World Wide Fund for Nature, Cambridge, UK

[10] Conservation and Policy, Zoological Society of London (ZSL), London, UK

* *Joint first and corresponding authors:* sophie.ledger@ioz.ac.uk; louise.mcrae@ioz.ac.uk

Abstract

As we enter the next phase of international policy commitments to halt biodiversity loss (e.g. Post-2020 Biodiversity Framework), biodiversity indicators will play an important role forming the robust basis upon which targeted, and time sensitive conservation actions are developed. Population trend indicators are perhaps the most powerful tool in biodiversity monitoring due to their responsiveness to changes over short timescales and their ability to aggregate species trends from global down to at a sub-national or even local scale. We consider how the project behind the foremost population level indicator - the Living Planet Index - has evolved over the last 25 years, its value to the field of biodiversity monitoring, and how its components have portrayed a compelling account of the changing status of global biodiversity through its application at policy, research and practice levels. We explore ways the project can develop to enhance our understanding of the state of biodiversity and share lessons learned to inform indicator development and mobilise action.

Box 1. The Living Planet Index Project

The Living Planet Index project (the index, methodology, and database) and its secondary outputs (methods papers and R code, database and website, global index, and subset indices) have had wide-ranging applications within the fields of biodiversity monitoring and research, as well as across policy, education, and outreach.

The Living Planet Index (LPI) is a biodiversity indicator which tracks trends in the relative abundance of wild vertebrate populations (where population is defined as to a single species in a defined location rather than the biological definition). Relative abundance captures how populations are changing over time on average in comparison to a reference point, or "baseline" (the LPI uses 1970). It is often described as analogous to a stock market index for species. The index is comprised of thousands of population time-series for vertebrate species from locations around the world; the trends from these populations are averaged to produce terrestrial, freshwater, and marine indices, which are further aggregated to a global LPI. The latest global LPI shows a decline of 68% between 1970 and 2016 globally [1]. This is an average trend based on time-series data from 20,811 populations of 4,392 species of mammals, birds, reptiles, amphibians, and fish.

The LPI database (LPD) can include population data for any species for which time-series population data could be found, regardless of threat status, or whether they show increasing or declining trends. These population time-series are sourced from scientific papers, online databases, government, and expert led published reports. They can be searched and downloaded from the project website (www.livingplanetindex.org). More technical information is available on the LPI stats website (http://stats.livingplanetindex.org/).

Introduction

The Living Planet Index (LPI) (Box 1) was first proposed as a means of evaluating environmental change, particularly by tracking trends in global biodiversity, a quarter of a century ago [2]. At that time, although there was mounting evidence of anthropogenic impacts on nature [3], there were very few indicators of the state of biodiversity or ecosystems at a global, or even regional scale. The initial version of the LPI, based on trends in vertebrate populations and forest cover, indicated that biodiversity was in decline globally [2]. A successful response to what is now widely recognised as a global biodiversity crisis [4-7] will involve transformative changes in the way humans use the planet's resources,[8-10] widespread intergovernmental action [11] and ambitious targets [9,10] (intergovernmental agreements such as the Convention on Biological Diversity (CBD) [12] and the United Nations Sustainable Development Goals (SDGs) [13]). To this end we need meaningful and reliable biodiversity indicators, generated from high quality and large-scale data to track progress towards targets down to the national level [10,14]. As such, the development of biodiversity indicators has become an increasing focus in conservation science [15-17], particularly to ensure they are fit for purpose as tools for management and policy, as well as to improve the representation of the underlying data beyond well-studied taxa and regions.

Within this review we chart the history, progression, and applications of the LPI project (Box 1). We review the LPI as a tool for public engagement and outreach, policy, and to drive further research and, analyse citation data to explore other applications of the LPI. We discuss challenges faced in maintaining a large biodiversity dataset and in current uses of the LPI. Finally, we look to the future and propose how the LPI project could evolve by enabling global collaboration to strengthen the indicator, harnessing new technologies for collecting population data, and developing new analysis to better understand the relationships between drivers and wildlife population trends.

The origins and development of the Living Planet Index

The Living Planet Index was conceived in 1997 by the World Wildlife Fund for Nature (WWF International). The primary aim was to "*develop a measure of the changing state of the world's biodiversity over time*" [18] using aggregate time-series population trends for a large sample of species from across the world. As very

little data were available on plants, fungi or invertebrate species, the pragmatic approach was taken to
restrict the initial LPI taxonomically to vertebrates. There was also geographic unevenness in the
distribution of the available data: long-term monitoring studies dating back decades were located mainly in
Europe and North America. To address the biases in data coverage, a benchmark of 1970 was set, and the
data were divided up into three broad biomes – terrestrial, freshwater and marine – and then further into
regional groupings. The source data and LPI outputs were at first collaboratively managed by WWF and the
World Conservation Monitoring Centre (now UN Environment Programme WCMC) for use within WWF's
flagship publication, the Living Planet Report (LPR). First published in 1998, the LPR used the initial
iteration of the LPI as a communications tool to convey biodiversity trends into a singular message on the
health of the planet for a broad audience, alongside measures of humanity's impact on the planet [2].
Calculated as -32% between 1970 and 1995 (Loh, et al. [2]), the downward trend of the LPI was already
apparent.
70
In the early 2000's, as the LPI dataset and methods were developed further [18], their potential for use in
advocacy, research, and as an indicator for monitoring biodiversity were recognised more widely. In 2002,
the Parties to the CBD committed to achieve a significant reduction of the rate of biodiversity loss at the
global, regional and national level by 2010 and required a framework of biodiversity indicators to monitor
their progress [19]. The first national LPI, the 'Living Uganda Index', was published with the National
Biodiversity Data Bank recording scheme at Makerere University, Uganda in 2004 [20,21] and was presented
as a case study for country-level applications of species population indices at CBD COP 7 [22]. A Discussion
Meeting held at the Royal Society in 2004 brought together leading academic and NGO researchers
working on biodiversity indicators, and the resulting papers, including one on the LPI, were published in a
special issue of Philosophical Transactions B [23]. This meeting laid much of the groundwork for subsequent
indicator development in the context of the CBD and other international biodiversity monitoring processes
[24]. In 2005, the Convention's scientific advisory body adopted the LPI metric as part of a suite of biodiversity
indicators, deployed to monitor progress towards that target [25]. In 2010, the CBD Parties agreed a further
set of biodiversity targets, the Aichi Targets, for the period 2011 to 2020 [4] and the LPI was identified as an
indicator for several of these.
86
To strengthen the LPI's scientific foundations and improve its capacity as an indicator for tracking progress
towards international biodiversity policy targets, an in-depth peer-reviewed paper on the methodology was
published [18] and the current partnership between WWF and the Zoological Society of London (ZSL) was
subsequently formed in 2006. Since then, two updates to the methodology behind the global index have
been published [26,27] and the research potential of the LPI data has expanded by incorporating metadata on
ecology, geography, threats and management into the database, the core data of which were made openly
accessible online in 2013 (18% of the data set is not available due to a confidentiality clause in the data
sharing agreement, often for rare or threatened species).
95

96   **Applications of the LPI**
97   Here we provide an overview of the uses of the different LPI project elements (see Box 1) and outputs,
98   grouped into three themes: public engagement and advocacy, policy and research.

99   1. *The LPI as a communication tool for public engagement and advocacy.*
100   From its inception, the LPI was seen as a powerful tool and WWF communications found that it
101   resonated with the public better than any other conservation messages at that time. The LPI helps
102   to set the scene for the state of global biodiversity by conveying a complex topic as a singular
103   takeaway message for a broad audience. The key conduit for the global LPI has been as the
104   headline biodiversity indicator within the (LPR). The LPR is an open access, biennial publication of
105   the latest research and insights into global biodiversity trends, the human drivers behind them, and
106   proposed solutions to halt biodiversity loss and "bend the curve" [10] back towards restoration. Its
107   widespread distribution and WWF's communications expertise have provided a regular global media
108   platform and, emphasizing opportunities for awareness raising and advocacy regarding the

biodiversity crisis. The 13[th] edition, published in 2020, was translated into 16 languages and
circulated around the world, with over 290 million social media views and 3,560 mentions from
monitored global news outlets within the first month of its launch [28]. The consistent use and media
exposure within the LPR has accorded the LPI with familiarity within the public realm (see
Communication and interpretation of the LPI). An analysis of online posts and articles (in English)
containing the LPR 2020's keywords or hashtags showed that 51% mentioned the 2020 global LPI
statistic [28]. Apart from global LPI figures, analysis of subset indices such as those featured in the
LPR 2020 (LPI by The Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem
Services (IPBES) regions, taxonomic focus (e.g. reptiles) and ecological biome (e.g. forests and
freshwater)) have been used to draw focus towards trends within different species groups [5,29,30].

Both the underlying data in the LPI and the global results have been used in several educational
formats, in schools and higher education. As part of the latest LPR outreach campaign, a youth
edition including the LPI trends was prepared [31] and adapted by WWF country offices to enable
young people to learn from the report's key messages and promote engagement of schools globally
in biodiversity issues.

Nature documentaries provide another medium for large-scale biodiversity outreach [32]. The 2019
Netflix series "Our Planet," narrated by Sir David Attenborough, used the global LPI statistic from
LPR 2018 to set the scene for its narrative alongside other headline biodiversity indicators and,
within the first month of the launch, was viewed by 45 million accounts across the world [33].

National scale LPI analysis and LPRs such as those undertaken by WWF offices in Belgium [34], the
Netherlands [35] and Canada [36], and regional approaches like the 2013 "Wildlife Comeback in
Europe" report [37] have used LPI figures to illustrate species trends and raise public awareness to
what is happening to status and trends of the biodiversity on their doorstep. The Wildlife Comeback
report reached 138 million people across Europe and worldwide [38].
133

2.   *The use of the LPI project within policy*
Analyses of the LPI dataset and trends within a geopolitical, ecological or taxonomic focus have
been used to provide evidence of biodiversity change for policy makers, fed into policy and target
development, and monitored progress towards those targets. The LPI is part of a suite of
biodiversity indicators adopted by the CBD, measuring trends in relative abundance of vertebrates
and deployed to monitor progress towards the 2010 Biodiversity Target [19], subsequent 2020 Aichi
targets [4], and is one of the indicators within Goal A of the Post-2020 Biodiversity Monitoring
Framework [39]. As a measure of population trends compiled at annual intervals, the LPI is sensitive
enough to detect annual changes, which is of value for informing policy [15] and evaluating the impact
of conservation interventions [40].[19]

ZSL and WWF joined the Biodiversity Indicators Partnership (BIP) in 2007 to further develop the LPI
and make it available for use under the CBD strategic plan. This resulted in the use of the LPI as
evidence of biodiversity decline in international policy documents (Table 1): global and regional
assessments (Millennium Ecosystem Assessment (2005) [41], IPBES global, regional and thematic
assessments [7,42-45] and successive updates of UN Global Environment Outlook [25,46-49] and UN
Global Biodiversity Outlook [50-53]) as well as thematic assessments (Ramsar Convention on
Wetlands, (2018) [54], Mediterranean Wetlands Outlooks (2012 and 2018) [55,56], the Convention on
Migratory Species reports (CMS) (2008 and in 2019) [57,58] and Arctic Biodiversity Assessment (2013)
[59]). More recently, the global and regional indices were used to illustrate the state of nature and how
this varies geographically as part of the evidence base for the Dasgupta review, an independent
report on the economics of biodiversity [60].

LPIs have been used as a scientific basis and in their scene setting capacity, to influence policy
development when advocating for transformative change and setting ambitious biodiversity targets
[9,10]. The global LPI statistic has featured in high-level biodiversity discussions, for example within
Volkan Bozkir's (President of the UN General Assembly) speech to heads of state at the 75[th] UN

Summit on Biodiversity in 2020 and within UK parliament in 2016 to support an Early Day Motion on
Global Biodiversity [61].

The LPI dataset and guidance on applying the method at a sub-global scale [62] have allowed for
regional, national and in some areas, sub-national scale analysis (Table 1). This 'scalability' is a key
requirement for indicators to be effective at tracking progress of signatory parties towards larger
intergovernmental targets [62,63]. CBD parties, for example, can develop national LPIs to fulfil part of
their progress reporting requirements within their National Biodiversity Strategy and Action Plans
(NBSAP) [3]. Several members, including the Netherlands, Uganda, Canada, and China have
provided LPI analysis of species trends within their NBSAP reports. In France, this process has
been scaled down even further and; provinces such as Provence-Alpes-Côte d'Azur have used LPI
analysis to track progress towards their National Biodiversity Strategy [64]. In Australia, a new
application of the LPI method focussed on threatened species to monitor their national progress
towards Aichi Target 12 (extinction prevented) [65].

Aside from tracking CBD commitments, nations have adapted the LPI method and applied it to suit
their state biodiversity indicator needs such as the "Canadian Species Index," developed by ZSL in
partnership with Environment and Climate Change Canada (ECCC) [66,67]. The package in the
programming language "R" for calculating the LPI (rlpi), is freely available via GitHub [68], and has
been used by collaborators from around the world to produce their own regional and national
indices e.g. national and scientific agencies within Brazil use it within a national bird and mammal
monitoring programme [69].

Application of the LPI	Corresponding biodiversity and sustainable development targets and other multilateral environmental agreements (MEAs)		
Disaggregation and reference	*CBD*	*SDG*	*MEAs*
Sub national			
LPI-Cat, State of Nature in Catalonia 2020 report for Catalunya, Spain [70]	CBD 12	SDG 14, SDG 15	ipbes
Indice Région Vivante (IRV), province of Provence-Alpes-Côte d'Azur, France [64]	CBD 12	SDG 14, SDG 15	ipbes
Indice Région Vivante (IRV), bird indicator for the province of Franche-Comté, France [71]	CBD 12	SDG 15	CMS ipbes
National			
Living Uganda Index (LUI), Uganda [21,72-75]	CBD 12	SDG 15	ipbes
Living Planet Index or Naturindeks for Norge, Norway [76]	CBD 12	SDG 14, SDG 15	ipbes
Canadian Species Index (CSI), one of a suite of Canadian Environmental Sustainability Indicators, Canada. [67,77] The Canadian Living Planet Index (C-LPI) [36]	CBD 12	SDG 14, SDG 15	ipbes
Living Planet Index Netherlands, the Netherlands [78,79]	CBD 12	SDG 14, SDG 15	ipbes
Living Planet Index, China [80 81]	CBD 12	SDG 15	ipbes
Belgian Living Planet Index, Belgium [34]	CBD 12	SDG 14, SDG 15	ipbes
Threatened Species Index (TSX), for birds, Australia [65]	CBD 12	SDG 15	ipbes
The Austrian Living Planet Index, Austria [82]	CBD 12	SDG 15	ipbes
Regional			
Arctic Species Trend Index (ASTI) for vertebrates across the Arctic [59,83,84]	CBD 10, CBD 12	SDG 14, SDG 15	ipbes Ramsar
ASTI for Arctic marine mammals, birds and fish [85]	CBD 10, CBD 12	SDG 14	ipbes

Application	Aichi Targets	SDG Goals	Conventions/MEAs
Arctic Migratory Birds Index [86]	10, 12	14, 15	CMS, ipbes
Mediterranean wetlands Living Planet Index [55,56,87]	5	6	ipbes, Ramsar
European marine vertebrates Living Planet Index, European Environment Agency (EEA) [88]	6, 12	14	ipbes
Ecological			
Living Planet Index for global estuarine systems [89]	6, 7, 10, 12	6, 14, 15	ipbes, Ramsar
Living Planet Index for migratory species [58,90]	12	14, 15	CMS, ipbes
Living Planet Index by marine, freshwater and terrestrial biomes [54,91]	5	6	ipbes, Ramsar
Living Planet Index for Reptiles [92]	12	14, 15	ipbes
Living Planet Index for freshwater megafauna [93]	7, 10, 12	6, 15	CMS, ipbes, Ramsar
Living Planet Index for migratory freshwater fish [94]	6, 10, 12	6, 15	CMS, ipbes, Ramsar
Forest Specialists Index [95]	12	6, 15	ipbes
Conservation management and species utilisation			
Protected areas and protected area management [96-99]	11	15	
Impacts of conservation management on species [40] and threatened species [36,100]	12	15	
Living Planet Index for recovering populations of European mammals and birds [37]	5, 7, 11, 15	15	ipbes
Living Planet Index for utilized species [101,102]	4, 12	8, 12, 14, 15	CITES, ipbes
Trends in target and bycatch species (oceanic sharks and rays) [103]	6, 12	14	ipbes
Other influences of the LPI			
Index of Linguistic Diversity [104,105]	18	1, 16	ipbes
The Wetland Extent Trends Index [106,107]	5	6	ipbes, Ramsar
Sustainability Policy Transparency Toolkit (SPOTT) Index [108]	4, 5, 7	12	
The Species Awareness Index (SAI) [109]	1		

Table 1. Selected applications of the LPI data and/or method and the corresponding and suggested uses for tracking global conventions on biodiversity, sustainable development, and other multilateral environmental agreements (MEAs). Sourced from: UNEP (United Nations Environment Programme) [12], UN [13], UNEP-WCMC (UN Environment Programme World Conservation Monitoring Centre) [110]. The Post-2020 Biodiversity Framework targets were not finalised at the point of submission and are not included.

3. *The LPI project as a tool for research*
The LPI methods, dataset and metrics have been used either individually or in unison for numerous
research projects around the world (Table 1 and Figure 1). Within a random sample of 341 citations
containing the term "Living Planet Index," 90% of author and document affiliation was classed as
research (academic institution or university); of the outputs themselves, 53% were within academic
journals (Supplementary Materials A and B).

The Living Planet Database (LPD) (except for about 18% of the data marked as confidential – see
the origins and development of the Living Planet Index) has been publicly available since 2013
when the LPI website was created to facilitate viewing and downloading the data. Prior to this,
subsets of the database were shared upon request. The LPD is now the largest repository of
vertebrate population trend data (containing over 38,000 populations of more than 5,000 species at
the time of writing), adding to a wealth of available biodiversity data for species occurrence (GBIF
[111]), species extinction risk (IUCN Red List [112]) and ecological community data (PREDICTS [113],
BioTIME [114]). To date, www.livingplanetindex.org has had over 6,000 registered users from 145
countries around the world.

Within the LPD, the population and ancillary data (Supplementary Materials D Figure 5) have
facilitated a wide range of research topics (Table 1). In particular, the threat and management data
at population-level allows for more fine-grained analysis compared with using species-level data.
Recent applications of the data include: measuring the effectiveness of protected areas [96-98];
evaluating the correlates of abundance trends in subsets of species such as mammals, reptiles,
forest specialists, freshwater megafauna and migratory species [90,92,93,95,115]; the nature of population
dynamics in response to threats or management [99-101,116,117]; the effects of land use and climate on
species [118] and exploring linkages between human development variables and wildlife population
trends [119].

The LPD has been incorporated into an open access repository at the University of Edinburgh,
dedicated to providing free online courses in statistics for ecology and environmental scientists [120].
In a more informal setting, LPI data have been used to present challenges for data visualisation or
analysis as part of Hackathons, one of which led to the development of a tool to automatically
identify papers containing abundance data [121].

The framework used to calculate the LPI has been applied to produce other metrics and not just for
biodiversity. Conceptually, relative change, as calculated by the geometric mean, can be applied to
other units of measurement that have been collected consistently over time. Using the code for
calculating the LPI, new indicators have been developed for wetland areas [106,107], linguistic diversity
[104], monitoring environmental, social and governance transparency in palm oil production [108] and
biodiversity awareness [109]. The first two of these are part of the ongoing suite of indicators for the
CBD (Table 1).

222
223 **Fig. 1. The number of publications per year citing the Living Planet Index between 1998-2020**. The secondary Y-axis shows
224 the cumulative total of publications. These 2,152 citations are from academic and grey literature in English and non-English
225 languages between the years 1998-2020 (as of 18th of January 2021). See Supplementary Materials A for details on the methods.
226

Challenges and opportunities

Along with other high-profile biodiversity indicators and reports [122,123], the underlying data, methods, and
interpretation of the LPI have come repeatedly under scrutiny, which has been a positive catalyst for new
research, collaborations and ameliorations on the scientific rigour of the index. Here we provide an outline
of the challenges faced by the LPI and aim to provide clarity on common misconceptions that have arisen
within recent years.

1) The dataset underpinning the LPI
One of the strengths of the LPD is that it is not static: data are continually added and updated to
provide the most complete and accurate picture possible of relative trends in population sizes
(Figure 2). To ensure data are comparable, only species-level time-series which fulfil the following
criteria are added: they are a measure of population abundance (or proxy, such as number of
breeding pairs), with two or more years of data, collected within a specified geographic location
under consistent methods (or explicitly corrected for) [26]. Supplementary metadata (Supplementary
Materials D) are continually updated for both new, and existing time-series, adding a further step in
the data extraction process [26]. The rigorous evaluation of data sources and data extraction not only
limits the amount of applicable data that can be included, but it is also time consuming and labour
intensive, and affects the volume of data that can be processed for each update. Storing these data
in suitable infrastructure and the financial support required to maintain it are a further limitation
common to other biodiversity databases [124]. The costs of running the entire project can be complex
to calculate as the source data are often already published and there are many stakeholders
including researchers and policymakers to consider.

Long-term, population, abundance studies at a species population level are a limited resource in
themselves, particularly for highly speciose taxa such as invertebrates and plants which have not
been included in the LPD to date (see The future). Studies which include population data may not
have been designed for long-term population monitoring but to assess population size and so their
methods and survey effort might change with advances in population estimate approaches (e.g.
revised Orangutan estimates in Sabah [125], this renders these data incompatible for inclusion in the

LPD). This issue is amplified for regions and taxa which are recognised as underrepresented within the dataset such as tropical regions and fish, reptiles and amphibians (see Supplementary Materials E) [27,126].

Subsequently, the composition of the LPD is likely to reflect bias inherent in species monitoring schemes which tend to favour certain taxa (e.g. birds), or regions (e.g. high income countries) [27,127,128]. This is a challenge shared by biodiversity indicators and databases in general [122,129]. In addition, attempts to source data from grey literature or offline databases is often dependent on the time and expertise available from researchers and field contacts within chronically neglected and underfunded areas [130]. To counteract bias in the resulting LPI, two approaches are taken. At the data inputting stage, a gap analysis of the taxonomic and geographic representation of the LPD is used to prioritise taxa and regions for targeted data searches (Supplementary Materials E). However, focussed searches are not always fruitful: within the 2020 LPI, only 4 populations of African amphibians were included despite targeted efforts [131]. The second step for overcoming bias in the LPI is in the adoption of the diversity-weighted method (see The LPI methods).

Language is a further constraint to collating representative data for the LPI and can exacerbate existing geographic biases [132]. The dominance of English-language data sources is partly a reflection of the LPI project being hosted in an English-speaking country but also of English as a globally used language for science [133]. However, over a third of biodiversity documents from a single year were published in languages other than English [134], so there are likely to be data that have not been captured because language barriers have not yet been adequately addressed.

Collating and storing a continually increasing repository of LPI data, that aligns with FAIR (Findable, Accessible, Interoperable, and Reusable) Data Principles, requires ongoing investment in the data infrastructure and management [124,135]. Coupled with this is the importance of promoting data sharing in a way that alleviates concerns over data ownership and provides appropriate credit to data providers. Unless a system is in place whereby data providers maintain ownership and control of their data, there is likely to be a barrier to mobilising data.

Figure 2. a-d. Growth in number of populations and species in the Living Planet Database (LPD) by region and taxa. The cumulative number of new populations (panel a) and species (b) entered by region, and the cumulative number of populations (c) and species (c) entered by taxon. Please note 2b adds up to more than the individual number of species as some species occur in more than one region.

 2) The LPI methods

The key methodological challenges for the LPI project are to generate a robust LPI indicator of
biodiversity and to model the time-series data in the LPD, which vary in length and scale, in a way
that allows exploration of underlying patterns in population trends. A further challenge that
underpins both issues, is addressing the taxonomic and geographic gaps in the underlying data
(Supplementary Materials E).

The basic formula for calculating the LPI has remained consistent: each logged population trend is
averaged within a single species and the species trends are aggregated to produce a single index
[18]. This aggregation is produced using a geometric mean, an approach used to generate other
indices of relative abundance from species abundance data [136-138]. Further levels of aggregation are
often used for global, national, and local contexts (see Supplementary Materials F for the global
example).

A challenge in the use of a geometric mean of abundance for the calculation of indicators is that it
can be sensitive to outliers in the data which may impact the precision of the long-term trend if not
addressed [139-141]. While this method is still considered to be a more suitable and sensitive metric to
assess changes in biodiversity [142,143], understanding the impact of outliers is important. To tackle
this, each new iteration of the global LPI analysis includes sensitivity tests on the influence of single
species on the trends and of the effect of short time-series on the LPI, as these are more commonly
associated with highly variable or extreme trends [131]. These tests are published in the
supplementary information, blog or website for transparency and to demonstrate the robustness of
any index [131].

The modelling of the time-series data in the LPD has been periodically improved. In early iterations
of the LPI, the chain method was implemented, which involved linearly interpolating the rate of
change between 5-year intervals, (following Loh, et al. [18]). As this approach was sensitive to abrupt
changes in population trends, generalised additive modelling (GAM) was adopted to better capture
long-term nonlinear trends in populations [26]. National variations of modelling have been tailored to
the type of species monitoring data in the country in question, for example the use of linear
regression for short-term trends in the Canadian Species Index [66,79].

More recently, Bayesian approaches such as state-space models have been applied to model the
population time-series whilst incorporating observation error into the estimation of trends [144],
something that the GAM framework does not account for. This has allowed for new ways of
analysing the LPD, which lend themselves to uncovering the correlates of vertebrate population
trends [145] and the taxonomic and geographic patterns of population trends globally [141].

A significant challenge remains in tackling the underrepresentation in the LPI database of particular
taxa and regions in the LPD so an adaptation to the LPI method was made to mitigate the impacts
of this bias on the index. This diversity-weighted approach was developed and subsequently
adopted for calculating global and regionals LPIs [27]. This method places greater weight on species
trends from regions and taxa that are more species-rich but tend to be disproportionately under-
represented in the LPD e.g. the Neotropics. This provides a more representative picture of global
vertebrate trends in lieu of a more complete data set. One drawback is that weight is often placed
on species and regions with the lowest data availability so if the sample of data from a region is not
representative, this could cause an over- or under- estimation of trends. Efforts are also underway
to address gaps in the data set through targeted data collection and to develop models to predict
trends in locations and for taxa which are data deficient, as has been done for extinction risk [146].
329

3) Communication and interpretation of the LPI
Key attributes of biodiversity indicators are that they should be simplified and easily understood [138].
The LPI was developed with these criteria in mind and, by aggregating trends from different
ecological realms and geographic regions, it can provide a useful overview and communication tool
for broad audiences. However, the index has been critiqued as oversimplifying the state of
biodiversity [147] and masking important trends [141]. Arguably, there is need for a balance between

providing a simple, clear message about global biodiversity trends whilst supporting it with more in-
depth analysis [123]. To explore this variation, disaggregations of the LPI have been developed (Table
1), for example, for forest specialists [95].

The limited availability of quality, ecological data prior to the 1970s is a common limitation to many
biodiversity indicators [129,148]. The LPI is benchmarked at a temporal baseline of 1970, and this raises
the importance of interpreting the index in context, as geopolitical regions have been impacted by
anthropogenic pressure at different points in time and varying intensity. In Europe, for example, a
significant amount of habitat destruction and overexploitation of some species had occurred prior to
the 1970s and therefore the LPI baseline is set at a significantly depleted reference point [37]. The
year chosen as a baseline can affect the interpretation of the state of biodiversity in a particular
region [149]. Without taking this into consideration, it is possible to underestimate the gravity of the
decline in biodiversity or overestimate a recovery within any given landscape.

Communications around biodiversity indicators and biodiversity loss have often centred on species,
and species extinctions respectively, rather than attempting to explain the multi-faceted nature of
biodiversity change and how we measure it [123,150]. Miscommunication and oversimplification of
biodiversity and biodiversity loss, or decline, across the science-society and science-policy
interface, are a challenge shared by biodiversity indicators in general [123]. The impact of substituting
a single word for another in press and media communications, "loss" vs "decline", has sometimes
led to misinterpretation of the global LPI statistic. A negative trend in the LPI depicts a relative
decline in population sizes, on average, since 1970. The use of the word "loss" in some media
articles can imply that a negative LPI trend is analogous with the disappearance of populations and
even the extinctions of species, which can prove challenging to correct. Media headlines have
referred to large percentages of populations being "wiped out" [151], which could mislead the public
about the severity of biodiversity decline and, it has been argued, such negative statements about
environmental issues may be counterproductive in trying to stimulate action [152].

Efforts to minimise misinterpretation have been made with each iteration of the LPI, by engaging
with journalists directly through press briefings, providing background information to
communications teams and publicising the supporting information available in technical
supplements to the LPR [131], websites (http://stats.livingplanetindex.org/) and blogs [153]. These efforts
have also reinforced the LPI as a measure of "relative abundance" rather than "abundance" to help
avoid misinterpretations [154]. We have already seen an uptake in the use of the LPR 2020 technical
supplement in recent publications and blogs exploring the LPI [155,156]. The analogy of a FTSE index
for biodiversity is most commonly used to describe the LPI, but a focus in the future should be on
finding other ways to communicate the index that mitigate the use of dramatic narratives, whilst
retaining the simple message of the LPI that can be broadly understood.

The future

The LPI project has grown significantly over the last 25 years and provides an important dataset to
communicate the trends in vertebrate populations and investigate the factors that influence them. We
identify four key priorities for the immediate future.

1. Increasing representation in the LPI
The composition of the LPI needs to be improved, crucially by increasing the taxonomic and geographic
representation of the data particularly for aquatic species. Incorporating invertebrate and plant species into
the LPI is likely to be challenging given the paucity of monitoring compared to some vertebrate groups [157]
but is key to attaining an indicator of broader biodiversity in addition to and providing a fuller data set for
macro-ecological research. Many national LPIs have already been developed, and maintaining this focus
on increasing the representation of species within countries will provide nations with a tool to track progress
towards future CBD and SDG targets. Indicators also need to be ecologically relevant [138], so ensuring that
different functional attributes of species within an ecosystem are reflected will be the focus of new research.
These developments in the data set will be realised through the use of emerging techniques to incorporate
unstructured data, such as that collected through citizen science initiatives [158-160], and capitalising on

growing technology for monitoring biodiversity such as eDNA, satellite monitoring and AI-assisted counting of species, provided they can be transformed into usable metrics of abundance.

2. Streamlining data collation and data access
Sourcing and extracting data continue to be significant bottlenecks for the development of the LPD. Data searches can be automated to some degree using predictive models based upon titles and abstracts [121], but extracting data automatically remains a challenge. Working with publishers, data holders, government institutions and research funding bodies to automate the process of identifying and extracting data from articles would be beneficial particularly if a standardised workflow is developed (e.g. Cardoso, et al. [161]), and systematic review tools may advance data collation in a community-driven way [162]. To address language barriers, which in turn could help to fill taxonomic and regional data gaps [163], a protocol for conducting data searches in multiple languages is under development. This should be part of a broader strategy to build a sustainable data network for the LPI, which provides accessibility to a global database (both for data download and upload, e.g. from new national LPI datasets) whilst retaining data quality and ownership, and assuring appropriate credit to data gatherers and providers. It is also important that the LPD is made as accessible as possible, both through simple, downloadable, tidy data formats [164] and the development of Application Programming Interfaces (API) to allow the data to interoperate with other resources such as the IUCN Red List [112], Protected Planet [165] and GBIF [111].

3. Better models to link population trends with drivers
The LPI continues to highlight that global biodiversity is in trouble and understanding (and predicting) which regions and species are likely to decline most in the future is useful. As such, models to better predict wildlife abundance trends for species and regions where we have poorer data is critical. Understanding the quality and utility of these models will allow us to make concrete and valuable predictions. The varied response of some populations to their changing environment highlights an important question – are some populations useful 'canaries' of pending ecosystem collapse and how might we best identify them?

Models that combine LPI data with drivers such as land-use and climate-change data have demonstrated that both are important drivers of population trends [118]. Developing these models further allows us to make predictions about how biodiversity might change under future scenarios and management interventions [9], highlighting one evolving use of biodiversity datasets like the LPD.

Whilst incorporating data on drivers from other global data sets can inform explanatory analysis for species trend data [118], population-scale information can also provide a powerful set of variables, for example in understanding the effect of different direct drivers [101] or to pave the way for counterfactual analysis of different management types (e.g. Jellesmark, et al. [166]). However, the current coding for threats and conservation action in the LPD lacks alignment with established frameworks [167], so transferring the ancillary information into these classification schemes and maintaining the recording of population drivers will improve the utility of models and ground-truthing of broad scale datasets in the future.

4. Increasing the utility of the LPI for policy
From a policy perspective, an emphasis on developing LPIs at the national level is needed to expand its use as a communication and reporting tool. With reporting requirements at a national level for the SDGs and the CBD, national LPIs would serve a dual purpose of providing countries with a sensitive indicator for reporting while boosting data representation for the global index. Disaggregations of the LPI on themes such as use, trade, migration and wetlands should continue to be developed, so that these are available for reporting against other multilateral environmental agreements such as the Ramsar Convention on Wetlands, CITES and the CMS.

The LPI performed well in an evaluation of biodiversity indicators using decision science [17], although gaps were identified in the practice of regular tests of the index and in assessing the cost- effectiveness of the LPI relative to other indicators. Creating a better understanding of how the LPI fits within the growing suite of biodiversity indicators such as the Red List Index [168] and the Biodiversity Intactness Index [169], and clearly presenting the complementarity of these indicators with each other, will be key to developing a clear and consistent narrative of global biodiversity change [14] and to ensure the suitability of the LPI within any multi-dimensional indicator framework [170,171].

Conclusion

The LPI has evolved from a simple communications tool to a large and growing database, policy tool and foundation for research. The open-access dataset and method are globally important resources for the scientific community and beyond, but improvements are still needed to enhance the representation of biodiversity in the underlying data and produce clear and meaningful outputs. Collaboration and engagement within the fields of science, policy, conservation and communication — some of which have fuelled much of the development to date, will continue to be important for ensuring the LPI project remains fit for purpose.

Acknowledgements

This research was partly funded by Research England, SL was funded by WWF-NL; LM, SD, VM were funded by WWF-UK.

We acknowledge the following individuals who were instrumental in the initial development and/or funding of the LPI project; Georgina Mace, Ben Collen, Jonathan Baillie, and Raj Amin. We also thank the LPI volunteers, collaborators, and contributors to the LPD past, and present for their essential support to the LPI project.

Competing interests

SL, LM, RF, KSG, VM, MF and SD are employed by ZSL (which hosts the LPD and partners with WWF to deliver the global LPI) and work directly on the management of the LPD and global LPI. MG, RA, LY and JC are employed by WWF offices (the umbrella organisation of which founded the LPI). MG and RA are Editors of the Living Planet Report. WWF-UK and WWF-NL have provided funding for project and research support and supported the research in kind.

Supporting information

Supplementary information is available for this paper at: DOI XXX

These include the methods for the citation search in academic and grey literature, and metadata coding (Supplementary materials A), the results of citation search and metadata coding (Supplementary materials B), the results of the Altmetric data analysis for key LPI papers (Supplementary materials C), an infographic of the underlying data within the LPD and summary of the growth in populations and species in the LPD over time (Supplementary materials D), a summary of LPD data diagnostics for underrepresented taxa and realms (Supplementary materials E) and a visualisation of the global aggregation LPI method (Supplementary materials F).

References

1   WWF/ZSL. *The Living Planet Index Database (LPD)*, <www.livingplanetindex.org> (2020).

2   Loh, J. *et al.* Living planet report: 1998. (WWF, Gland, Switzerland, 1998).

3   UN. Convention on biological diversity. 5th June 1992. (United Nations Conference on Environment and Development, Rio de Janeiro, 1992).

4   Tittensor, D. P. *et al.* A mid-term analysis of progress toward international biodiversity targets. *Science* **346**, 241-244, doi:10.1126/science.1257484 (2014).

5   WWF. Living planet report 2020 - Bending the curve of biodiversity loss. (WWF, Gland, Switzerland, 2020).

6   Diaz, S. *et al.* Assessing nature's contributions to people. *Science* **359**, 270-272,
doi:10.1126/science.aap8826 (2018).

7   IPBES. Global assessment report on biodiversity and ecosystem services of the Intergovernmental
Science-Policy Platform on Biodiversity and Ecosystem Services. (IPBES Secretariat, Bonn,
Germany, 2019).

8   Diaz, S. *et al.* Pervasive human-driven decline of life on Earth points to the need for transformative
change. *Science* **366**, doi:10.1126/science.aax3100 (2019).

9   Leclère, D. *et al.* Bending the curve of terrestrial biodiversity needs an integrated strategy. *Nature*
585, 551-556, doi:10.1038/s41586-020-2705-y (2020).

10 Mace, G. M. *et al.* Aiming higher to bend the curve of biodiversity loss. *Nature Sustainability* **1**, 448-
451, doi:10.1038/s41893-018-0130-0 (2018).

11 Xu, H. *et al.* Ensuring effective implementation of the post-2020 global biodiversity targets. *Nat Ecol*
Evol **5**, 411-418, doi:10.1038/s41559-020-01375-y (2021).

12 UNEP (United Nations Environment Programme). Decision X/2: The strategic plan for biodiversity
2011–2020 and the Aichi Biodiversity Targets. Adopted at the 10th Conference Of The Parties
(COP) to the Convention On Biological Diversity (CBD). (UNEP, Montreal, Canada, 2010).

13 UN. Transforming our world: The 2030 agenda for sustainable development. (United Nations (UN),
2015).

14 Hill, S. L. L. *et al.* Reconciling Biodiversity Indicators to Guide Understanding and Action.
Conservation Letters **9**, 405-412, doi:10.1111/conl.12291 (2016).

15 Jones, J. P. *et al.* The why, what, and how of global biodiversity indicators beyond the 2010 target.
Conserv Biol **25**, 450-457, doi:10.1111/j.1523-1739.2010.01605.x (2011).

16 Nicholson, E. *et al.* Making robust policy decisions using global biodiversity indicators. *PLoS ONE* **7**,
e41128, doi:10.1371/journal.pone.0041128 (2012).

17 Watermeyer, K. E. *et al.* Using decision science to evaluate global biodiversity indices. *Conserv Biol*
35, 492-501, doi:10.1111/cobi.13574 (2021).

18 Loh, J. *et al.* The Living Planet Index: Using species population time series to track trends in
biodiversity. *Philosophical Transactions of the Royal Society B: Biological Sciences* **360**, 289-295,
doi:10.1098/rstb.2004.1584 (2005).

19 Butchart, S. H. *et al.* Global biodiversity: Indicators of recent declines. *Science* **328**, 1164-1168,
doi:10.1126/science.1187512 (2010).

20 Arinaitwe, H., Pomeroy, D. E. & Tushabe, H. The State of Uganda's Biodiversity: 2000. 56 (National
Biodiversity Data Bank, Makerere University Institute of Environment and Natural Resources,
Kampala, Uganda, 2000).

21 Pomeroy, D. & Tushabe, H. The State of Uganda's Biodiversity 2004. (National Biodiversity
DataBank (NBDB). Makerere University Institute of Environment and Natural Resources (MUIENR),
Kampala, Uganda, 2004).

22 Jenkins, M., Kapos, V. & Loh, J. Rising to the biodiversity challenge. The role of species population
trend indices like the Living Planet Index in tracking progress towards global and national
biodiversity targets. (World Bank, Washington, DC. USA, 2004).

23 Balmford, A., Crane, P. R., Green, R. E. & Mace, G. M. Discussion Meeting Issue 'Beyond
extinction rates: monitoring wild nature for the 2010 target'. *Philosophical Transactions of the Royal*
Society B: Biological Sciences **360**, 219-477 (2005).

24 UNEP (United Nations Environment Programme). Decision VIII/15: Framework for monitoring
implementation of the achievement of the 2010 target and integration of targets into the thematic
programmes of work. Adopted at the 8th Conference Of The Parties (COP) to the Convention On
Biological Diversity (CBD). (UNEP, Curitiba, Brazil, 2006).

25 UNEP (United Nations Environment Programme). Report on the eighth meeting of the Conference
of the Parties to the Convention on Biological Diversity, CBD. (UNEP, Nairobi, 2006).

26 Collen, B. *et al.* Monitoring change in vertebrate abundance: The Living Planet Index. *Conservation
Biology* **23**, 317-327, doi:10.1111/j.1523-1739.2008.01117.x (2009).

27 McRae, L., Deinet, S. & Freeman, R. The diversity-weighted Living Planet Index: Controlling for
taxonomic bias in a global biodiversity indicator. *PLOS ONE* **12**, e0169156,
doi:10.1371/journal.pone.0169156 (2017).

28 WWF. Living Planet Report 2020 - Network results. (WWF, Internal report, 2021).

29 WWF. Living planet Report 2020. Bending the curve of biodiversity loss: a deep dive into climate
and biodiversity. (WWF, Gland, Switzerland, 2020).

30 WWF. Living planet Report 2020. Bending the curve of biodiversity loss: a deep dive into
freshwater. (WWF, Gland, Switzerland, 2020).

31 WWF. Living Planet Report 2020 youth edition: A guide for our future. (WWF, Gland, Switzerland,
2020).

32 Jones, J. P. G., Thomas-Walters, L., Rust, N. A., Veríssimo, D. & Januchowski-Hartley, S. Nature
documentaries and saving nature: Reflections on the new Netflix series Our Planet. *People and
Nature* **1**, 420-425, doi:10.1002/pan3.10052 (2019).

33 WWF/tve. Our Planet: Our impact - The first year of the Our Planet Project. (WWF-UK, Woking, UK,
2020).

34 WWF. Rapport Planète Vivante - La Nature en Belgique. (WWF, Brussels, Belgium, 2020).

35 Wereld Natuur Fonds. Living Planet Report Nederland. Natuur en landbouw verbonden. (WWF-NL,
Zeist, 2020).

36 WWF-Canada. Living Planet Report Canada: Wildlife At Risk. (World Wildlife Fund Canada,
Toronto, Canada, 2020).

37 Deinet, S. *et al.* Wildlife comeback in Europe: The recovery of selected mammal and bird species.
Final report to Rewilding Europe by ZSL, BirdLife International and the European Bird Census
Council. (ZSL, London, UK, 2013).

38 Rewilding Europe. Annual review 2013. (Rewilding Europe, The Netherlands, 2013).

39 UNEP (United Nations Environment Programme). in *24th Meeting of the Subsidiary Body On
Scientific, Technical And Technological Advice (SBSTTA)*   (Convention on Biological Diversity
(CBD),, 2020).

40 Jellesmark, S. *et al.* Assessing the global impact of targeted conservation actions on species
abundance. *bioRxiv*, 2022.2001.2014.476374, doi:10.1101/2022.01.14.476374 (2022).

41 Millennium Ecosystem Assessment. Ecosystems and Human Well-being: Biodiversity Synthesis.
(World Resources Institute, Washington, DC., 2005).

42 IPBES. The IPBES regional assessment report on biodiversity and ecosystem services for the
Americas of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem

Services. (Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.
IPBES Secretariat, Bonn, Germany, 2018).

43 IPBES. The IPBES regional assessment report on biodiversity and ecosystem services for Europe
and Central Asia of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem
Services. (Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.
IPBES Secretariat, Bonn, Germany, 2018).

44 IPBES. The IPBES regional assessment report on biodiversity and ecosystem services for Africa of
the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.
(Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. IPBES
Secretariat, Bonn, Germany, 2018).

45 IPBES. The IPBES regional assessment report on biodiversity and ecosystem services for Asia and
the Pacific of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem
Services. (Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.
IPBES Secretariat, Bonn, Germany, 2018).

46 UNEP (United Nations Environment Programme). Global Environment Outlook 3: Past, present and
future perspectives. (United Nations Environment Programme, Nairobi, Kenya, 2002).

47 UNEP (United Nations Environment Programme). Global Environment Outlook 4: Environment for
development. (United Nations Environment Programme, Nairobi, Kenya, 2007).

48 UNEP (United Nations Environment Programme). Global Environment Outlook 5: Environment for
the future we want. (United Nations Environment Programme, Nairobi, Kenya, 2012).

49 UNEP (United Nations Environment Programme). *Global Environment Outlook – GEO-6: Healthy*
planet, healthy people. (United Nations Environment Programme. Cambridge University Press,
2019).

50 Secretariat of the Convention on Biological Diversity. Global Biodiversity Outlook 2. 81 + vii
(Montréal, Canada, 2006).

51 Secretariat of the Convention on Biological Diversity. Global Biodiversity Outlook 3. 94 (Montréal,
Canada, 2010).

52 Secretariat of the Convention on Biological Diversity. Global Biodiversity Outlook 4. 155 (Montréal,
Canada, 2014).

53 Secretariat of the Convention on Biological Diversity. Global Biodiversity Outlook 5. (Montréal,
Canada, 2020).

54 Ramsar Convention on Wetlands. Global wetland outlook: State of the world's wetlands and their
services to people. (Ramsar Convention Secretariat, Gland, Switzerland, 2018).

55 MWO (Mediterranean Wetlands Observatory). Mediterranean Wetlands Outlook. First
Mediterranean Wetlands Observatory report - Technical report. 128 pages (Tour du Valat, France,
2012).

56 MWO (Mediterranean Wetlands Observatory). Mediterranean Wetlands Outlook 2: Solutions for
Sustainable Mediterranean Wetlands. (Tour du Valat, France, 2018).

57 Deinet, S. The Living Planet Index (LPI) for species listed on the CMS Appendices. Technical
summary submitted to UNEP-WCMC and the CMS Secretariat. 11 (ZSL,
UNEP/CMS/COP13/Doc.24/Annex 5, 2019).

58 Latham, J., Collen, B., McRae, L. & Loh, J. The Living Planet Index for migratory species: An index
of change in population abundance. 22 (ZSL/WWF, 2008).

59   CAFF (Conservation of Arctic Flora and Fauna). *Arctic biodiversity assessment. Status and trends*
in Arctic biodiversity. (Conservation of Arctic Flora and Fauna, 2013).

60   Dasgupta, P. The economics of biodiversity: The Dasgupta review. (HM Treasury, London, 2021).

61   UK Parliament. *Early Day Motions. Number 624: Global biodiversity. Tabled 31st October, 2016*
(2016-17 Session) (2016).

62   McRae, L. *et al.* Living Planet Index Guidance for national and regional use Version 1.1. 11
(Cambridge, UK., 2008).

63   Brooks, T. M. *et al.* Analysing biodiversity and conservation knowledge products to support regional
environmental assessments. *Scientific Data* **3**, 160007, doi:10.1038/sdata.2016.7 (2016).

64   PACA (Observatoire Régional de la Biodiversité en Provence-Alpes-Côte d'Azur). Indice Région
Vivante. Comment évolue la biodiversité en Provence-Alpes-Côte d'Azur? , (Observatoire Régional
de la Biodiversité en Provence-Alpes-Côte d'Azur, 2018).

65   Bayraktarov, E. *et al.* A threatened species index for Australian birds. *Conservation Science and*
Practice **3**, doi:10.1111/csp2.322 (2020).

66   Marconi, V. *et al.* Population declines among Canadian vertebrates: But data of different quality
show diverging trends. *Ecological Indicators* **130**, 108022, doi:10.1016/j.ecolind.2021.108022
(2021).

67   Environment and Climate Change Canada (ECCC). Canadian Environmental Sustainability
Indicators: Canadian species index. (2019).

68   Freeman, R., McRae, L., Deinet, S., Amin, R. & Collen, B. *rlpi: Tools for calculating indices using*
the Living Planet Index method. R Package, <https://github.com/Zoological-Society-of-London/rlpi>
(2017).

69   ICMBio-CENAP/Programa-Monitora-Florestal-Global. *Analise de dados Mastoaves do protocolo*
florestal global do programa Monitora, <https://github.com/ICMBio-CENAP/Programa-Monitora-
Florestal-Global> (2021).

70   Brotons, L. *et al.* Estat de la Natura a Catalunya 2020. (Departament de Territori i Sostenibilitat.
Generalitat de Catalunya, Barcelona, 2020).

71   Maas, S. & Giroud, I. Indice Région Vivante (IRV) : indicateurs oiseaux de Franche-Comté. 11p
(LPO Franche-Comté, DREAL Bourgogne Franche-Comté et Conseil Régional Bourgogne Franche-
Comté, 2016).

72   Pomeroy, D., Tushabe, H. & Loh, J. The State of Uganda's Biodiversity 2017. (National Biodiversity
Data Bank. Makerere University, Kampala, 2017).

73   Pomeroy, D. & Tushabe, H. The State of Uganda's Biodiversity 2006. (Makerere Institute of
Environment and Natural Resources/National Biodiversity Data Bank, 2006).

74   NEMA (National Environment Management Authority). State of Environment Report for Uganda.
332 (NEMA, Kampala, Uganda, 2006/7).

75   Pomeroy, D. & Tushabe, H. The state of Uganda's biodiversity report: Sixth biennial report.
(National Biodiversity Data Bank (NBDB), Makerere University Institute of Environment and Natural
Resources (MUIENR), 2008).

76   WWF-Norge. Naturindeks for Norge 2005. Utfor bakke med norsk natur. (WW-Norge, Oslo, Norway,
2005).

77   Muller, H. *et al.* The Canadian Species Index. (ZSL/Environment Canada, 2016).

78 CBS, PBL, RIVM & WUR. *Trend fauna - all species monitored - Living Planet Index Netherlands,*
1990-2018 (indicator 1569, version 05 , 30 March 2020), <https://www.clo.nl/en/indicators/en1569-
living-planet-index-for-the-netherlands> (2021).

79 van Strien, A. J. *et al.* Modest recovery of biodiversity in a western European country: The Living
Planet Index for the Netherlands. *Biological Conservation* **200**, 44-50,
doi:10.1016/j.biocon.2016.05.031 (2016).

80 WWF China. Living planet report China 2015: Development, species and ecological civilization.
(WWF China in partnership with China Council for International Cooperation on Environment and
Development (CCICED), Institute of Geographic Sciences and Natural Resources Research
(IGSNRR) and Institute of Zoology of Chinese Academy of Sciences (CAS), and the Global
Footprint Network, 2015).

81 The Government of China. Sixth national report to the Conventional on Biological Diversity.
(Secretariat of the Convention on Biological Diversity (SCBD), The Clearing-House Mechanism of
the Convention on Biological Diversity (CHM), 2019).

82 Semmelmayer, K. & Hackländer, K. Monitoring vertebrate abundance in Austria: Developments over
30 years. *Die Bodenkultur: Journal of Land Management, Food and Environment* **71**, 19-30,
doi:10.2478/boku-2020-0003 (2020).

83 McRae, L., Böhm, M., Deinet, S., Gill, M. & Collen, B. The Arctic Species Trend Index: using
vertebrate population trends to monitor the health of a rapidly changing ecosystem. *Biodiversity* **13**,
144-156, doi:10.1080/14888386.2012.705085 (2012).

84 McRae, L. *et al.* Arctic Species Trend Index 2010. Tracking Trends in Arctic Wildlife. (CAFF
International Secretariat, 2010).

85 McRae, L., Deinet, S., Gill, M. & Collen, B. The Arctic Species Trend Index: Tracking trends in Arctic
marine populations. (Conservation of Arctic Flora and Fauna (CAFF), Iceland, 2012).

86 Deinet, S. *et al.* Arctic Species Trend Index: Migratory Birds Index. (Conservation of Arctic Flora and
Fauna (CAFF), Akureyri, Iceland, 2015).

87 Galewski, T., Segura, L., Biquet, J., Saccon, E. & Boutry, N. Living Mediterranean Report:
Monitoring species trends to secure one of the major biodiversity hotspots. (Tour du Valat (TdV),
France, 2021).

88 EEA (European Environment Agency). Marine messages II: Navigating the course towards clean,
healthy and productive seas through implementation of an ecosystem-based approach. (European
Environment Agency, Luxembourg, 2019).

89 Deinet, S. *et al.* The Living Planet Index for Global Estuarine Systems: Technical report. (WWF/ZSL,
2010).

90 Hardesty-Moore, M. *et al.* Migration in the Anthropocene: how collective navigation, environmental
system and taxonomy shape the vulnerability of migratory species. *Philos Trans R Soc Lond B Biol*
Sci **373**, doi:10.1098/rstb.2017.0017 (2018).

91 WWF. Living planet report 2016. Risk and resilience in a new era. Report No. 978-2-940529-40-7,
(WWF, Gland, Switzerland, 2016).

92 Saha, A. *et al.* Tracking Global Population Trends: Population Time-Series Data and a Living Planet
Index for Reptiles. *Journal of Herpetology* **52**, doi:10.1670/17-076 (2018).

93 He, F. *et al.* The global decline of freshwater megafauna. *Glob Chang Biol* **25**, 3883-3892,
doi:10.1111/gcb.14753 (2019).

94 Deinet, S. *et al.* The Living Planet Index (LPI) for migratory freshwater fish - Technical Report.
(World Fish Migration Foundation, The Netherlands, 2020).

95 Green, E. J. *et al.* Below the canopy: global trends in forest vertebrate populations and their drivers.
Proc Biol Sci **287**, 20200533, doi:10.1098/rspb.2020.0533 (2020).

96 Geldmann, J. *et al.* A global analysis of management capacity and ecological outcomes in terrestrial
protected areas. *Conservation Letters* **11**, e12434, doi:10.1111/conl.12434 (2018).

97 Craigie, I. D. *et al.* Large mammal population declines in Africa's protected areas. *Biological
Conservation* **143**, 2221-2228, doi:10.1016/j.biocon.2010.06.007 (2010).

98 Barnes, M. D. *et al.* Wildlife population trends in protected areas predicted by national socio-
economic metrics and body size. *Nature Communications* **7**, 12747, doi:10.1038/ncomms12747
(2016).

99 Costelloe, B. *et al.* Global biodiversity indicators reflect the modeled impacts of protected area policy
change: Biodiversity indicators and protected areas. *Conservation Letters* **9**, 14-20,
doi:10.1111/conl.12163 (2016).

100 Currie, J., Marconi, V. & Kerr, J. An analysis of threats and factors that predict trends in Canadian
vertebrates designated as at-risk. *Facets* **5**, 49-66, doi:10.1139/facets-2019-0017 (2020).

101 McRae, L. *et al.* A global indicator of utilised wildlife populations: regional trends and the impact of
management. *bioRxiv*, doi:10.1101/2020.11.02.365031 (2021).

102 Tierney, M. *et al.* Use it or lose it: Measuring trends in wild species subject to substantial use. *Oryx
48, 420-429, doi:10.1017/S0030605313000653 (2014).

103 Pacoureau, N. *et al.* Half a century of global decline in oceanic sharks and rays. *Nature* **589**, 567-
571, doi:10.1038/s41586-020-03173-9 (2021).

104 Harmon, D. & Loh, J. The index of linguistic diversity: A new quantitative measure of trends in the
status of the world's languages. *Language Documentation & Conservation* **4**, 97-151,
doi:http://hdl.handle.net/10125/4474 (2010).

105 Loh, J. & Harmon, D. Biocultural Diversity: threatened species, endangered languages. (WWF
Netherlands, Zeist, The Netherlands, 2014).

106 Dixon, M. J. R. *et al.* Tracking global change in ecosystem area: The Wetland Extent Trends index.
Biological Conservation **193**, 27-35, doi:10.1016/j.biocon.2015.10.023 (2016).

107 Darrah, S. E. *et al.* Improvements to the Wetland Extent Trends (WET) index as a tool for
monitoring natural and human-made wetlands. *Ecological Indicators* **99**, 294-298,
doi:10.1016/j.ecolind.2018.12.032 (2019).

108 Oppenheimer, P. *et al.* The SPOTT index: A proof-of-concept measure for tracking public disclosure
in the palm oil industry. *Current Research in Environmental Sustainability* **3**,
doi:10.1016/j.crsust.2021.100042 (2021).

109 Millard, J. W., Gregory, R. D., Jones, K. & Freeman, R. The Species Awareness Index (SAI): a
Wikipedia-derived conservation culturomics metric for public biodiversity awareness. *bioRxiv*,
doi:10.1101/2020.08.17.254177 (2020).

110 UNEP-WCMC (UN Environment Programme World Conservation Monitoring Centre). *The
Biodiversity Indicators Partnership (BIP)* <https://www.bipindicators.net/> (2021).

111 GBIF. *GBIF: The Global Biodiversity Information Facility*, <https://www.gbif.org/> (2021).

112 IUCN. *The IUCN Red List of Threatened Species. Version 2021-3.*, <https://www.iucnredlist.org>
(2021).

113 Hudson, L. N. *et al.* The database of the PREDICTS (Projecting Responses of Ecological Diversity
In Changing Terrestrial Systems) project. *Ecology and Evolution* **7**, 145-188, doi:10.1002/ece3.2579
(2017).

114 Dornelas, M. *et al.* BioTIME: A database of biodiversity time series for the Anthropocene. *Glob Ecol*
Biogeogr **27**, 760-786, doi:10.1111/geb.12729 (2018).

115 Collen, B. *et al.* Predicting how populations decline to extinction. *Philosophical Transactions of the*
Royal Society B: Biological Sciences **366**, 2577-2586, doi:10.1098/rstb.2011.0015 (2011).

116 Di Fonzo, M. D., Collen, B. & Mace, G. M. A new method for identifying rapid decline dynamics in
wild vertebrate populations. *Ecol Evol* **3**, 2378-2391, doi:10.1002/ece3.596 (2013).

117 Capdevila, P., Noviello, N., McRae, L., Freeman, R. & Clements, C. F. Global patterns of resilience
decline in vertebrate populations. *Ecology Letters* **25**, 240-251, doi:10.1111/ele.13927 (2022).

118 Spooner, F. E. B., Pearson, R. G. & Freeman, R. Rapid warming is associated with population
decline among terrestrial birds and mammals globally. *Glob Chang Biol* **24**, 4521-4531,
doi:10.1111/gcb.14361 (2018).

119 Ament, J. M. *et al.* Compatibility between agendas for improving human development and wildlife
conservation outside protected areas: Insights from 20 years of data. *People and Nature* **1**, 305-
316, doi:10.1002/pan3.10041 (2019).

120 University of Edinburgh. *Our Coding Club*, <https://ourcodingclub.github.io/> (2021).

121 Cornford, R. *et al.* Fast, scalable, and automated identification of articles for biodiversity and
macroecological datasets. *Global Ecology and Biogeography* **30**, 339-347, doi:10.1111/geb.13219
(2020).

122 Rodrigues, A. S., Pilgrim, J. D., Lamoreux, J. F., Hoffmann, M. & Brooks, T. M. The value of the
IUCN Red List for conservation. *Trends Ecol Evol* **21**, 71-76, doi:10.1016/j.tree.2005.10.010 (2006).

123 Turnhout, E. & Purvis, A. Biodiversity and species extinction: categorisation, calculation, and
communication. *Griffith Law Review* **29**, 669-685, doi:10.1080/10383441.2020.1925204 (2021).

124 Kays, R., McShea, W. J., Wikelski, M. & Zurell, D. Born-digital biodiversity data: Millions and billions.
Diversity and Distributions **26**, 644-648, doi:10.1111/ddi.12993 (2020).

125 Ancrenaz, M. *et al.* Aerial surveys give new estimates for orangutans in Sabah, Malaysia. *PLoS Biol*
3, e3, doi:10.1371/journal.pbio.0030003 (2005).

126 Collen, B., Ram, M., Zamin, T. & McRae, L. The tropical biodiversity data gap: Addressing disparity
in global monitoring. *Tropical Conservation Science* **1**, 75-88, doi:10.1177/194008290800100202
(2008).

127 Moussy, C. *et al.* A quantitative global review of species population monitoring. *Conserv Biol*,
doi:10.1111/cobi.13721 (2021).

128 Proença, V. *et al.* Global biodiversity monitoring: From data sources to Essential Biodiversity
Variables. *Biological Conservation* **213**, 256-263, doi:10.1016/j.biocon.2016.07.014 (2017).

129 Hoffmann, M., Brooks, T. M., Butchart, S. H. M., Gregory, R. D. & McRae, L. in *Encyclopedia of the*
Anthropocene 175-184 (2018).

130 Stephenson, P. J. *et al.* Priorities for big biodiversity data. *Frontiers in Ecology and the Environment*
15, 124-125, doi:10.1002/fee.1473 (2017).

131 WWF. Living planet Report 2020. Bending the curve of biodiversity loss: a deep dive into the Living
Planet Index. Marconi, V., McRae, L., Deinet, S., Ledger, S. and Freeman, F. in (WWF, Gland,
Switzerland, 2020).

132 Amano, T. & Sutherland, W. J. Four barriers to the global understanding of biodiversity
conservation: wealth, language, geographical location and security. *Proc Biol Sci* **280**, 20122649,
doi:10.1098/rspb.2012.2649 (2013).

133 Montgomery, S. L. *Does science need a global language?* , (The University of Chicago Press,
2013).

134 Amano, T., Gonzalez-Varo, J. P. & Sutherland, W. J. Languages Are Still a Major Barrier to Global
Science. *PLoS Biol* **14**, e2000933, doi:10.1371/journal.pbio.2000933 (2016).

135 Wilkinson, M. D. *et al.* The FAIR Guiding Principles for scientific data management and stewardship.
Sci Data **3**, 160018, doi:10.1038/sdata.2016.18 (2016).

136 Buckland, S. T., Magurran, A. E., Green, R. E. & Fewster, R. M. Monitoring change in biodiversity
through composite indices. *Philosophical Transactions of the Royal Society B: Biological Sciences*
360, 243-254, doi:10.1098/rstb.2004.1589 (2005).

137 Buckland, S. T., Marsden, S. J. & Green, R. E. Estimating bird abundance: making methods work.
Bird Conservation International **18**, S91-S108, doi:10.1017/s0959270908000294 (2008).

138 Gregory, R. D. *et al.* Developing indicators for European birds. *Philos Trans R Soc Lond B Biol Sci*
360, 269-288, doi:10.1098/rstb.2004.1602 (2005).

139 Buckland, S. T., Studeny, A. C., Magurran, A. E., Illian, J. B. & Newson, S. E. The geometric mean
of relative abundance indices: a biodiversity measure with a difference. *Ecosphere* **2**,
doi:10.1890/es11-00186.1 (2011).

140 Gregory, R. D., Skorpilova, J., Vorisek, P. & Butler, S. An analysis of trends, uncertainty and
species selection shows contrasting trends of widespread forest and farmland birds in Europe.
Ecological Indicators **103**, 676-687, doi:10.1016/j.ecolind.2019.04.064 (2019).

141 Leung, B. *et al.* Clustered versus catastrophic global vertebrate declines. *Nature* **588**, 267-271,
doi:10.1038/s41586-020-2920-6 (2020).

142 Santini, L. *et al.* Assessing the suitability of diversity metrics to detect biodiversity change. *Biological
Conservation* **213**, 341-350, doi:10.1016/j.biocon.2016.08.024 (2017).

143 van Strien, A. J., Soldaat, L. L. & Gregory, R. D. Desirable mathematical properties of indicators for
biodiversity change. *Ecological Indicators* **14**, 202-208, doi:10.1016/j.ecolind.2011.07.007 (2012).

144 Auger-Méthé, M. An introduction to state-space modeling of ecological time series. *arXiv preprint*,
doi:arXiv:2002.02001 (2020).

145 Daskalova, G. N., Myers-Smith, I. H. & Godlee, J. L. Rare and common vertebrates span a wide
spectrum of population trends. *Nat Commun* **11**, 4394, doi:10.1038/s41467-020-17779-0 (2020).

146 Bland, L. M., Collen, B., Orme, C. D. & Bielby, J. Predicting the conservation status of data-deficient
species. *Conserv Biol* **29**, 250-259, doi:10.1111/cobi.12372 (2015).

147 Jaspers, A. Can a single index track the state of global biodiversity? *Biological Conservation* **246**,
108524, doi:10.1016/j.biocon.2020.108524 (2020).

148 Martin, L. J., Blossey, B. & Ellis, E. Mapping where ecologists work: biases in the global distribution
of terrestrial ecological observations. *Frontiers in Ecology and the Environment* **10**, 195-201,
doi:10.1890/110154 (2012).

149 Collins, A. C., Böhm, M. & Collen, B. Choice of baseline affects historical population trends in
hunted mammals of North America. *Biological Conservation* **242**, doi:10.1016/j.biocon.2020.108421
(2020).

150 Navarro, M. & Tidball, K. G. Challenges of biodiversity education: A review of education strategies
for conserving biodiversity. *International Electronic Journal of Environmental Education* **2**, 13–30
(2012).

151 Carrington, D. *Humanity has wiped out 60% of animal populations since 1970, report finds*,
<https://www.theguardian.com/environment/2018/oct/30/humanity-wiped-out-animals-since-1970-
major-report-finds> (2018).

152 O'Neill, S. & Nicholson-Cole, S. "Fear won't do it": Promoting positive engagement with climate
change through visual and iconic representations. *Science Communication* **30**, 355-379,
doi:10.1177/1075547008329201 (2009).

153 Freeman, R. *The Living Planet Index – data analysis, clusters and biodiversity loss*,
<https://www.zsl.org/blogs/science/the-living-planet-index--data-analysis-clusters-and-biodiversity-
loss> (2020).

154 Puurtinen, M., Elo, M. & Kotiaho, J. S. The Living Planet Index does not measure abundance.
Nature **601**, E14-E15, doi:10.1038/s41586-021-03708-8 (2022).

155 Buschke, F. T., Hagan, J. G., Santini, L. & Coetzee, B. W. T. Random population fluctuations bias
the Living Planet Index. *Nature Ecology & Evolution* **5**, 1145-1152, doi:10.1038/s41559-021-01494-
0 (2021).

156 Ritchie, H. *Living Planet Index*, <https://ourworldindata.org/living-planet-index> (2021).

157 Hochkirch, A. *et al.* A strategy for the next decade to address data deficiency in neglected
biodiversity. *Conserv Biol* **35**, 502-509, doi:10.1111/cobi.13589 (2021).

158 Outhwaite, C. L., Gregory, R. D., Chandler, R. E., Collen, B. & Isaac, N. J. B. Complex long-term
biodiversity change among invertebrates, bryophytes and lichens. *Nature Ecology & Evolution* **4**,
384-392, doi:10.1038/s41559-020-1111-z (2020).

159 Pocock, M. J. O., Logie, M. W., Isaac, N. J. B., Outhwaite, C. L. & August, T. Rapid assessment of
the suitability of multi-species citizen science datasets for occupancy trend analysis. *bioRxiv*,
813626, doi:10.1101/813626 (2019).

160 van Strien, A. J., van Swaay, C. A. M., van Strien-van Liempt, W. T. F. H., Poot, M. J. M. &
WallisDeVries, M. F. Over a century of data reveal more than 80% decline in butterflies in the
Netherlands. *Biological Conservation* **234**, 116-122, doi:10.1016/j.biocon.2019.03.023 (2019).

161 Cardoso, P., Stoev, P., Georgiev, T., Senderov, V. & Penev, L. Species Conservation Profiles
compliant with the IUCN Red List of Threatened Species. *Biodivers Data J*, e10356,
doi:10.3897/BDJ.4.e10356 (2016).

162 Grames, E. *et al.* Trends in global insect abundance and biodiversity: A community-driven
systematic map protocol. *Open Science Framework (osf.io/uxk4a)* doi:10.17605/OSF.IO/Q63UY
(2019).

163 Amano, T. *et al.* Tapping into non-English-language science for the conservation of global
biodiversity. *bioRxiv*, doi:10.1101/2021.05.24.445520 (2021).

164 McRae, L., Deinet, S. & Freeman, R. LPI_LPR2016data_public.csv. *Figshare. Dataset*,
doi:https://doi.org/10.6084/m9.figshare.4300022.v1 (2016).

165 UNEP-WCMC & IUCN. *Protected Planet*, <https://www.protectedplanet.net/en> (2021).

166   Jellesmark, S. *et al.* A counterfactual approach to measure the impact of wet grassland
861   conservation on U.K. breeding bird populations. *Conserv Biol* **35**, 1575-1585,
862   doi:10.1111/cobi.13692 (2021).

863   167   Salafsky, N. *et al.* A standard lexicon for biodiversity conservation: unified classifications of threats
864   and actions. *Conserv Biol* **22**, 897-911, doi:10.1111/j.1523-1739.2008.00937.x (2008).

865   168   Butchart, S. H. *et al.* Improvements to the Red List Index. *PLoS One* **2**, e140,
866   doi:10.1371/journal.pone.0000140 (2007).

867   169   Hill, S. L. L. *et al.* Worldwide impacts of past and projected future land-use change on local species
868   richness and the Biodiversity Intactness Index. *bioRxiv*, doi:10.1101/311787 (2018).

869   170   Soto-Navarro, C. A. *et al.* Building a Multidimensional Biodiversity Index – A scorecard for
870   biodiversity health. Project report. (UN Environment Programme World Conservation Monitoring
871   Centre (UNEP-WCMC), Cambridge, UK and Luc Hoffmann Institute (LHI), Gland, Switzerland.,
872   2020).

873   171   Soto-Navarro, C. A. *et al.* Towards a multidimensional biodiversity index for national application.
874   *Nature Sustainability* **4**, 933-942, doi:10.1038/s41893-021-00753-z (2021).