High-throughput phenotyping of infection by diverse microsporidia species reveals a wild

***C. elegans* strain with opposing resistance and susceptibility traits**

Calvin Mok[1#], Meng A. Xiao[1*], Yin C. Wan[1*], Winnie Zhao[1], Shanzeh M. Ahmed[1], Robert J.

Luallen[2] and Aaron W. Reinke[1#]

[1] Department of Molecular Genetics, University of Toronto, Toronto, ON, Canada

[2] Department of Biology, San Diego State University, San Diego, CA 92182, USA

*These authors contributed equally

#Corresponding authors

calvin.mok@utoronto.ca

aaron.reinke@utoronto.ca

Abstract

Animals are under constant selective pressure from a myriad of diverse pathogens. Microsporidia

are ubiquitous animal parasites, but the influence they exert on shaping animal genomes is mostly

unknown. Using multiplexed competition assays, we measured the impact of four different species

of microsporidia on 22 wild isolates of *Caenorhabditis elegans*. This resulted in the identification

and confirmation of 13 strains with significantly altered population fitness profiles under infection

conditions. One of these identified strains, JU1400, is sensitive to an epidermal-infecting species

by lacking tolerance to infection. JU1400 is also resistant to an intestinal-infecting species and

can specifically recognize and destroy this pathogen. Genetic mapping of JU1400 demonstrates

that these two opposing phenotypes are caused by separate loci. Transcriptional analysis reveals

the JU1400 sensitivity to epidermal microsporidia infection results in a response pattern that

shares similarity to toxin-induced responses. In contrast, we do not observe JU1400 intestinal

resistance being regulated at the transcriptional level. The transcriptional response to these four

microsporidia species is conserved, with *C. elegans* strain-specific differences in potential

immune genes. Together, our results show that phenotypic differences to microsporidia infection

amongst *C. elegans* are common and that animals can evolve species-specific genetic

interactions.

Introduction

Obligate intracellular parasites only propagate inside of host cells, which places extreme pressure

on the host, resulting in genomic changes [1–3]. Animals have coevolved with pathogens and

genetic variation to pathogen susceptibility is common in animal populations [4,5]. Much of this

genetic and phenotypic diversity to infection is specific to individual species of pathogens or even

to different strains [6,7]. Hosts can evolve several ways to improve their fitness in the presence

of pathogens. They can become more resistant by preventing pathogen invasion or by limiting

pathogen growth [8]. Alternatively, hosts can evolve to better tolerate the infection, by limiting the

negative effects of infection without reducing the amount of pathogen [9,10].

Microsporidia are a large phylum of eukaryotic obligate intracellular pathogens with a wide host

range including humans and agriculturally important animals such as honey bees and shrimp [11–

14]. Microsporidia infections cause developmental delays, a reduction in progeny, and increased

mortality [15–17]. Several studies have shown that microsporidia can affect host population

dynamics and evolution. Microsporidia have been suggested to influence the composition of

Caenorhabditis elegans genotypes and resistant strains can outcompete sensitive ones in just a

few generations [18–20]. In competition experiments with over a dozen isolates of *Daphnia*

magna, different microsporidia species have been shown to select for specific strains [21,22].

Using genetic mapping, several genomic regions contributing to microsporidia resistance in *C.*

elegans, *D. magna*, and honey bees have been identified [19,23–26].

C. elegans has become a powerful model for studying host-pathogen coevolution [27–30]. In the

wild, *C. elegans* contend with bacterial, fungal, viral, and microsporidian pathogens and these

coevolving species represent strong selective pressures on the animal's genome [31,32]. To

facilitate the study of natural diversity, hundreds of isolates of *C. elegans* have been collected

from around the world [33]. These wild isolates have different levels of susceptibility to pathogens

and natural variants have been discovered that impact traits such as bacterial avoidance and viral

susceptibility [7,19,34–37]. Although studying phenotypic variation in wild isolates is a powerful

approach, measuring each strain individually is time consuming and laborious. To circumvent this

issue, strategies have been developed to perform multiplex competitive fitness assays under

selective conditions such as starvation stress or different bacterial diets [38–40]. Several

sequencing-based approaches, termed either PhenoMIP or MIP-seq, determine the identity of

these individual strains by using molecular inversion probes (MIPs) to enrich for their unique

genomic signatures [39,40].

A common microbial infection found in wild *C. elegans* is from species of microsporidia in the

Nematocida genus. Infection of *C. elegans* by microsporidia begins with ingestion of a chitin-

coated spore [41]. The spore then fires its unique infectious apparatus, known as a polar tube,

and deposits its sporoplasm within the host [42]. The sporoplasm then replicates to form

membrane-bound, multinucleated cells know as meronts which then differentiate into spores that

are shed by the host [16]. Some *Nematocida* species, such as *N. parisii*, *N. ausubeli*, and *N.*

ironsii, infect only the intestine, while others, such as *N. displodere*, *N. ferruginous*, and *N. cider*,

can proliferate in several tissues including the muscle and epidermis [15,19,41,43,44]. *C. elegans*

possesses several mechanisms to defend itself against microsporidia [45,46]. Infection with

microsporidia induces transcriptional upregulation of genes known as the Intracellular Pathogen

Response (IPR) [47]. Mutations in several negative regulators of the IPR cause elevated

expression of the IPR and resistance to microsporidia infection [48–50]. *C. elegans* also possess

a separate form of inherited immunity that is activated by infection of parents and is transferred

to progeny [49].

Here we present the results of a high-throughput fitness competition of 22 pooled *C. elegans* wild

isolates infected by four species of microsporidia at multiple doses. We identified 16 candidate

interactions of either host resistance or sensitivity. We demonstrate that the wild isolate JU1400

has distinct and opposing responses to infection by different microsporidia species. This strain

has a defect in tolerance to the epidermally-infecting *N. ferruginous* but is resistant to the

intestinal-infecting *N. ironsii* through specific elimination of infection by this species. Through

mapping experiments and the generation of recombinant strains, we demonstrate that the

opposing phenotypes of JU1400 are caused by separate loci. In contrast to the species-specific

phenotypes we observed from fitness competitions, transcriptional analysis showed a host

response to infection that is mostly independent of the microsporidia species, but dependent upon

the *C. elegans* strain. Notably, we identified a signature of down-regulated genes that has overlap

with the *C. elegans* response to toxins, and this downregulated response was most prominent in

the tolerance-defective strain JU1400 infected with *N. ferruginous*. Our results demonstrate that

PhenoMIP can efficiently identify resistant and sensitive wild isolates and that microsporidia

genotype-specific interactions are common in *C. elegans*.

Results

A PhenoMIP screen of wild isolate strains reveals diverse responses to microsporidia infection

To identify *C. elegans* strains that vary in their resistance or susceptibility to microsporidia infection, we carried out a modified PhenoMIP screen. We chose 22 *C. elegans* wild isolate strains that are genetically diverse and isolated from a range of geographic regions (**Supplementary Figure S1**). We selected four pathogenic microsporidia species that differ in their impact on *C. elegans* development and the tissues they infect (*N. parisii*, *N. ausubeli*, and *N. ironsii* infect the intestine and *N. ferruginous* infects the muscle and epidermis) (**Figure 1A and Supplementary Table S1**) [15,16,19,43,51]. To synchronize infections, we treated adult populations with a sodium hydroxide and sodium hypochlorite mixture (known as bleach synchronizing) and allowed the recovered embryos to hatch in the absence of food. The process of bleach synchronizing animals for microsporidia infection has been shown to have similar levels of infection between bleached versus laid embryos, and short-term starvation is reported to have minimal impact on wild isolates [19,20,40]. We pooled equal numbers of synchronized first larval stage (L1) wild isolates and a laboratory reference-derived strain (VC20019) into a single population. VC20019, was chosen as a reference because it has many genomic variants that can be used to distinguish it from N2 while having developmental timing and fecundity similar to that of N2 [39,52]. We then plated these animals along with their food source of *Escherichia coli* bacteria mixed with one of four doses of each microsporidia species (high, medium, low, and none) (**Supplementary Table S2**). We hypothesized that a high-dose exposure could identify *C. elegans* strains that were resistant or tolerant to microsporidia infection while medium- and low-dose exposures might identify strains exhibiting mild to strong susceptibility to microsporidia infection. Animals were exposed to microsporidia spores for 72 hours at 21°C, before collecting and bleach synchronizing the population. This mixture inhibits microsporidia from infecting the next generation of animals and synchronizes the population [49]. This microsporidia-free population of L1s was then grown in the

absence of infection for 72 hours at 21°C to allow intergenerational immunity to reset [49]. Except

for the *N. ausubeli* medium-dose infection replicates (**see Methods**), worms in each condition

were infected for an additional three cycles for a total of eight generations. Adult animals were

collected every two generations at the end of the uninfected phase and genomic DNA was isolated

for further analysis (**Figure 1B**).

To determine how population composition changes due to exposure to microsporidia, we

calculated the relative abundance of each strain within each condition. We used MIPs to target

strain-specific variants for sequencing at high depth. In total we sequenced 127 samples, which

is the equivalent of 2162 infections and 736 mock/control infections. From each of the eight mock-

infection controls and twenty-four infection replicates for every strain we calculated a mean fold-

change rate (FCR) from across all timepoints. The distribution of FCRs for each strain was then

converted to a modified z-score distribution to identify candidate interactions (**See Methods**).

From this dataset, we identified 16 nematode/microsporidia pairings where the modified z-score

for both high-infection replicates was consistently outside the range of +/- 1.5 (**Figure 2,**

Supplementary Figure S2 and Table 1). We also generated mean growth profiles across all

timepoints for each worm strain in environments that were either pathogen-free or infected by

each of the microsporidia species (**Figure 3A and B**). From our analyses we observed ten

interaction pairs with reduced population fitness (low-fitness strains), suggesting potential

susceptibility to infection. In addition, we observed six interactions with increased population

fitness (high-fitness strains), suggesting a potential resistance to infection conditions. Our high-

fitness interaction set included the previously reported resistance interaction of CB4856 to *N.*

ironsii (**Supplementary Figure S3**) [19,51] which further suggested our scoring threshold was

appropriate for the detection of interactions in this assay format. Of the 15 novel candidate

interactions, we identified five interactions with *N. ironsii* (ERTm5), eight with *N. ferruginous*

(LUAm1), two with *N. ausubeli* (ERTm2), and no strong interactions with *N. parisii* (ERTm1). In

total, our screen yielded 13 *C. elegans* wild isolate strains with significant interactions, including

the strains JU360, JU1400, and MY2 that had interactions with more than one species of

microsporidia (**Figure 2 and Table 1**).

JU1400 fails to tolerate *N. ferruginous* infection

To validate a subset of our large-scale multiplex identification of microsporidia-strain phenotypes,

we performed isogenic population infection experiments. We infected individual strains with doses

of microsporidia spores equivalent to those used in the PhenoMIP experiment (**Figure 3C and D**

and Supplementary Figures S4-S6). We examined a combination of phenotypes including

morphology of adult populations, formation of microsporidia meronts or spores, and embryo

production. Most high-fitness strains tended to experience a smaller reduction in embryo

production ratios compared to changes in lab reference N2 in identical infection conditions. Low-

fitness strains, on the other hand, experienced developmental delay phenotypes and/or a

reduction in fecundity–both of which would contribute negatively to the fitness readout observed

via PhenoMIP. These phenotypes were most evident in JU1400 and MY1 which both had severe,

dose-dependent responses to infection by the epidermal microsporidia species *N. ferruginous*

(**Figure 3C and D and Supplementary Figure S7**). In total, of the 12 pairs that we observed

(**Table 1**), nine had altered relative fecundity profiles that could account for the fitness changes

observed from our PhenoMIP screen. Additionally, of the seven pairs we retested that had a z-

score below -1.5, six had a significant reduction in embryos when infected, compared to N2.

To examine how a *C. elegans* strain responds to different microsporidia species, we focused on

JU1400, in which we observed reduced fitness to infection by both *N. ferruginous* and *N. ironsii*.

To determine why JU1400 appeared sensitive to *N. ferruginous*, we performed pulse-chase

experiments by infecting animals for 3 hours, washing away free spores, and replating animals

for an additional 69 hours [19]. We observed that the percentage of JU1400 and N2 animals

infected with *N. ferruginous* was similar (**Figure 3E and Supplementary Figure S8A and B**).

Additionally, the amount of meronts formed during infection did not significantly differ between N2

and JU1400 (**Figure 3F and Supplementary Figure S8C and 8D**). Taken together, these results

demonstrate that although JU1400 is more sensitive to the effects of *N. ferruginous*, JU1400 is

not significantly more infected, suggesting a defect in tolerance.

To determine the specificity of JU1400 to *N. ferruginous* infection, we infected JU1400 with other

related microsporidia species and strains. We infected JU1400 with a second isolate of *N.*

ferruginous, LUAm3, as well as a separate muscle and epidermal-infecting microsporidian, *N.*

cider [44]. Infection by LUAm3 caused the same severe phenotype as LUAm1 (**Supplementary**

Figure S9A and B). JU1400 is infected by *N. cider* to a similar extent as N2, but infection by this

species resulted in a dose-dependent reduction in embryo numbers in JU1400 that was not

observed in N2 (**Supplementary Figure S9C and D**).

JU1400 specifically clears *N. ironsii* infection

In addition to the verified susceptibility to *N. ferruginous,* our PhenoMIP screen also identified a

more moderate fitness disadvantage for JU1400 in high-dose exposures to *N. ironsii* (**Figure 3A**).

We infected N2 and JU1400 with *N. ironsii* for 72 hours and compared the relative change in

embryos for each strain versus mock-infected controls. We observed no significant difference in

this normalized number of embryos between the N2 and JU1400 strains (**Figure 4A**). However,

JU1400 is significantly less infected, suggesting that JU1400 is resistant to *N. ironsii* infection

(Figure 4C and Supplementary Figure S10A and C). To determine whether the observed

resistance conferred a fitness advantage at a later time point, we exposed JU1400 animals to *N.*

ironsii for 96 hours and observed significantly more embryos and significantly less infection

compared to N2 (**Figure 4B and D and Supplementary Figure S10B and D**). Together our data

suggest that although JU1400 appears to have a short-term disadvantage in our pooled

competition assays, this strain has a long-term advantage due to enhanced resistance towards

N. ironsii.

Several mechanisms of *C. elegans* resistance to microsporidia infection have been described, including a block of invasion or clearance of infected parasite [19,20,49,51]. To determine how JU1400 is resistant to *N. ironsii* infection, we performed a pulse-chase experiment by infecting animals for 3 hours, washing away any external microsporidia spores, and examining animals at 3, 24, and 72 hours post infection (hpi). Both N2 and JU1400 had similar numbers of sporoplasms at 3 hpi, but JU1400 had decreased levels of pathogen at both 24 and 72 hpi (**Figure 5A**). This experiment revealed that JU1400 does not block microsporidia invasion, but instead has the capability to actively eliminate *N. ironsii* infection. To determine if the resistance in JU1400 was dependent upon development stage, we performed pulse-chase infections at the L4 stage and observed that JU1400 animals could no longer clear the infection (**Figure 5B**).

Resistance to *N. ironsii* has previously been observed in the *C. elegans* wild isolate, CB4856 [19]. This resistance was shown to occur through a modest decrease in invasion coupled with the ability to clear the infection at the L1 stage. To compare the properties of these two resistant strains, we infected both strains with *N. ironsii* for 72 hours. This experiment shows that JU1400 is more resistant to infection **(Figure 5C)**. To determine the specificity of JU1400 resistance, we tested infection with either *N. parisii* or *N. ausubeli* for 72 hours. CB4856 was slightly, but not significantly less infected by *N. parisii* and *N. ausubeli,* whereas JU1400 was infected by these species to the same extent as N2 (**Figure 5D**). We also tested the ability of JU1400 and CB4856 to clear microsporidia infection, and both strains could only eliminate *N. ironsii* (**Supplementary Figure S10E and F**). Together, these results demonstrate that JU1400 resistance is specific to *N. ironsii*.

The immunity of JU1400 against *N. ironsii* could be a specific recognition of *N. ironsii* infection that induces broad immunity towards microsporidia or the immunity itself could be specific for *N. ironsii*. To distinguish between these possibilities, we set up coinfection experiments using species-specific FISH probes and then coinfected JU1400 and N2 animals with *N. ausubeli* and

either *N. ironsii* or *N. parisii*. Under these conditions, *N. ironsii* sporoplasms are eliminated

regardless of the presence of *N. ausubeli*. There is no clearance of *N. ausubeli* observed, either

by itself or in the presence of *N. ironsii*. (**Figure 5E and Supplementary figure S11A and B**). No

clearance was observed in N2 regardless of the infection condition (**Supplementary Figure**

S11C). These results further demonstrate that JU1400 has specific immunity towards *N. ironsii*.

Genetic mapping reveals multiple loci are responsible for the interactions of JU1400 with

microsporidia

The opposing phenotypes of JU1400 in response to microsporidia infection could be caused by

the same or different genetic loci. To determine the genetic basis for JU1400 being sensitive to

N. ferruginous and resistant to *N. ironsii*, we mapped the causative alleles using a pooled

competitive fitness assay [52]. We crossed JU1400 to the mapping strain VC20019 and infected

F2 animals with either *N. ferruginous* for 72 hpi or *N. ironsii* for 96 hpi (**Supplementary Table S3**

and Methods). After two rounds of selection, genomic DNA was isolated from each sample and

MIP-MAP sequencing libraries were generated [52]. As a control, we used this approach to map

the resistance of CB4856 to *N. ironsii.* We then compared our results to a previous study that

determined regions responsible for resistance [19]. Although the previous mapping study was

based on pathogen load as opposed to our mapping of competitive fitness advantage, several

regions mapped concordantly between the two approaches on chromosome II and the right arm

of chromosome V (**Supplementary Figure S12)**.

Our mapping results of JU1400 with *N. ferruginous* identified a single peak of VC20019 marker

fixation on the left-hand side of chromosome I (**Figure 6A**). This mapping was repeated with MY1,

another strain that exhibited a fitness disadvantage with *N. ferruginous* (**Figure 3B-D)**. The

mapped region under negative selection was the same in MY1. The MY1 strain also carries the

zeel-1/peel-1 incompatibility locus at chromosome I, which accounts for the specific signal seen

at the I:2,726,021 marker in both the infected and uninfected samples (**Supplementary Figure**

S13A) [53]. To determine if the same locus was responsible for *N. ferruginous* in MY1 and JU1400, we tested their ability to complement. First, we determined that both MY1 and JU1400 display a recessive phenotype when crossed to our control strain. We then mated MY1 and JU1400 and the resulting cross progeny were susceptible to *N. ferruginous* infection, indicating that the two strains do not complement (**Supplementary Figure S14A and B**). We then infected MY1 with *N. ironsii*. There were fewer MY1 animals containing spores compared to N2, and less JU1400 animals had spores or meronts compared to MY1 animals (**Supplementary Figure S14C and D**). Together this data suggests a region on the left side of chromosome I, is responsible for sensitivity to *N. ferruginous* in multiple strain*s*.

Our mapping of JU1400 with *N. ironsii* at 96 hours identified three regions that are linked with resistance to infection, one on the left-hand side of chromosome II from 1,639,865-4,426,687, a second in a broad area of chromosome V from 11,035,658-18,400,066, and a third on the right-hand side of chromosome X from 15,180,772-17,676,467 (**Figure 6B**). A similar pattern was observed when these mapping experiments were performed at 72 hours, with the exception that the region on chromosome X was no longer observed (**Supplementary Figure S13B**). We also observed a region on the left side of chromosome I where JU1400 variants underwent weak negative selection upon over multiple *N. ironsii* infection rounds. Together, our data suggests that JU1400 resistance to *N. ironsii* is multigenic and may be influenced by infection time.

To confirm that the region on chromosome I was responsible for sensitivity to *N. ferruginous* infection, we generated Near Isogenic Lines (NILs). These NILs were constructed by crossing a version of VC20019 that contained a CRISPR-integrated single-copy GFP on chromosome I (AWR133) (**Supplementary Figure S15A**). We tested ten independent NILs that contained at least some JU1400 sequence to the left of I:2,851,000. All ten of these NILs were sensitive to infection by *N. ferruginous* (**Supplementary Figure S15B).** We used molecular markers to genotype these NILs, mapping the susceptibility locus to I:1-1,051,810 (**Supplementary Table**

S4). We then outcrossed one of these NILs, AWR140 back to N2 six times. During this outcrossing

process, we recovered two lines, AWR144 and AWR145, that contained different responses to

N. ferruginous infection (**Figure 6D**). We performed whole-genome sequencing of AWR138

(another independently isolated NIL), AWR140, AWR144, and AWR145 which narrowed the

critical region of association for our phenotype to I:1-701,078 (**Figure 6C**). Within this region, we

identified 12 genes shared between JU1400 and MY1 (same complementation group) but absent

in ED3042 and JU360 which tested into separate complementation groups **(Supplementary**

Table S5). The NIL AWR145 is sensitive to *N. ferruginous* infection but is only partially resistant

to infection by *N. ironsii* (**Figure 6D and E).** Together, our data demonstrate that the *N.*

ferruginous sensitivity and *N. ironsii* resistance to JU1400 are determined by distinct genetic loci.

Transcriptional analysis of the JU1400 response to microsporidia infection

To determine how the JU1400 strain differs in its response to microsporidia infection, we

performed transcriptional analysis. We grew three biological replicates of N2 and JU1400 L1

stage animals either without microsporidia or infected with one of *N. parisii*, *N. ironsii*, *N.*

ferruginous, or *N. ausubeli* for 48 hours. We then extracted and sequenced the *C. elegans* and

microsporidia mRNA. First, we mapped the microsporidia mRNA in each of the eight infected

samples. For the three intestinal infecting species, between ~1-9% of reads were mapped for

each sample, with the exception of *N. ironsii* in JU1400, where few reads were identified, which

is consistent with this strain being resistant to this species. For *N. ferruginous*, fewer reads were

observed in either strain compared to the intestinal species, which is consistent to what we

observed by microscopy (**Supplementary Figure S16 and Supplementary Table S6**).

We then sought to determine how the two *C. elegans* strains responded to the four different

microsporidia species (**Supplementary Table S7)**. First, we used principal component analysis

(PCA), and projected samples back onto the two principal components which accounted for more

than 80% of the variance in our samples. PCA analysis revealed a clear clustering between the

two strains (**Figure 7A)** and a secondary analysis by strain clustered the uninfected samples

against pathogen infected samples (**Supplementary Figure S17**). Second, we clustered the

differential expression profiles of the eight infected samples. This approach revealed several large

groups of genes that were differentially expressed either across most samples or within the same

strain **(Figure 7B** and **Supplementary Table S8).** Although the largest difference in the samples

is between the two strains, we observe significant correlation between strains infected with each

microsporidia species (**Supplementary Figure S18**).

We then determined the properties of genes that are differentially regulated in each sample**.** The

number of differentially upregulated or downregulated genes ranged from 6-819, with the most

being in JU1400 infected with *N. ferruginous* **(Figure 7C).** We then classified differentially

expressed genes into either conserved, microsporidia species (pathogen) specific, *C. elegans*

strain specific, or strain-pathogen specific categories **(Figure 7D)**. We identified 97 genes

upregulated in all samples, and no genes that were broadly downregulated. We identified ~30-50

differentially expressed genes specific to either *N. ausubeli* or *N. ferruginous* infection, one gene

specific to *N. parisii* infection, and no genes specific to *N. ironsii* infection. We also observed 45

or more differentially upregulated genes that were specific to either N2 or JU1400. To determine

the types of differentially expressed genes in these samples, we annotated them using a set of

domains previously observed to be regulated during microsporidia infection [47,48,54]

(**Supplementary Table S9)**. About 35% of the conserved response were genes containing these

domains, with the most frequent being PALS, F-box, and MATH/BATH domains. About 25% of

the strain-specific genes also contained these domains, with each strain having at least one

PALS, F-box, and MATH/BATH, c-type lectin, and skr domain-containing gene expressed that is

specific to that strain **(Figure 7D and Supplementary Table S10)**. Skp-related proteins (*skr*) in

C. elegans were shown to bind to CUL-6 to ubiquitylate specific proteins [55]. *skr-4* is upregulated

in both host strains infected by all microsporidia species. Among the upregulated genes, we also

observe strain-specific *skr* genes such as JU1400 specific *skr-6*, and N2 specific *skr-5*

(**Supplementary Table S8)**.

To understand how transcriptional responses correlated with resistant and susceptible

phenotypes, we analyzed genes that were specifically regulated in JU1400 when infected with

either *N. ironsii* or *N. ferruginous* (**Figure 7D)**. Surprisingly, we only observed three differentially

expressed genes specific to *N. ironsii* infection of JU1400. In contrast, we saw 181 upregulated

and 266 down-regulated genes specific to *N. ferruginous* infection of JU1400. Many of the

upregulated genes contained domains enriched in the IPR. The down-regulated genes contained

a few genes with IPR domains, as well as 23 genes with cuticle collagen domains. These proteins

with collagen domains were especially striking, as we never identified more than one of these

genes having altered expression in any of the other gene sets.

To determine if other stresses had similarities to the response of JU1400 to *N. ferruginous*, we

compared the set of genes specifically differentially expressed in this sample to previously

published *C. elegans* transcriptional datasets [56]. We identified the top ten chemical or microbial

treatment data sets that had the most significant overlap with JU1400 infected by *N. ferruginous*

(**Supplementary Table S11)**. We then determined the overlap of these data sets to all our

samples (**Figure 7E-F and Supplementary Table S12).** This revealed that genes differentially

downregulated in JU1400 in response to *N. ferruginous* infection overlapped significantly with

treatments of chemical poisons such as pesticides (fenamiphos and dichlorvos), organic

compounds (acrylamide and tert-butyl hydrogen peroxide), and heavy metals (mercury and

cadmium). All of these treatments were previously shown to inhibit development or the production

of progeny in *C. elegans* [57–60]. We also observed significant overlap of the response to these

chemicals with JU1400 infected with *N. ironsii*, and both strains infected with *N. ausubeli*. These

three infected samples along with JU1400 infected with *N. ferruginous* also contained the largest

number of significantly down-regulated genes **(Figure 7C and Supplementary Figure S19).**

Additionally, we observed significant overlap of these four microsporidia infected samples to

pathogenic bacteria or toxins including *Staphylococcus aureus*, *Serratia marcescens*, *Bacillus*

thuringiensis, *Xenorhabdus nematophila*, and *Pseudomonas aeruginosa* (**Figure 7E-F**). These

patterns of transcriptional overlap were the most striking with differentially downregulated genes,

although some overlap with upregulated genes was also observed (**Supplementary Figure S20**).

We also performed enrichment analysis of our transcriptional data sets, observing that the

upregulated genes are commonly enriched for metabolic processes and signalling pathways**.** In

contrast, down regulated genes are enriched for proteolysis, fatty-acid related metabolic

processes and drug metabolism (**Supplementary Tables S13 and S14).** We also analyzed the

12 candidate genes within the mapped chromosome I region that contain genetic variants in both

JU1400 and MY1. None of these genes are differentially regulated in JU1400 upon *N. ferruginous*

infection.

Discussion

Animal populations often have genetic diversity to bacterial, viral, and fungal pathogens, but less

is known about how microsporidia infection influences host variation [5,6]. To investigate how

different microsporidia species impact genetic diversity, we measured the phenotypic responses

to infection in a collection of genetically diverse *C. elegans* wild isolates. A total of 13 out of 22

wild isolate strains displayed a significant difference in population fitness after microsporidia

infection when compared to the population as a whole. We detected these interactions among

three of the four microsporidia species we measured, with *N. ironsii* and *N. ferruginous* accounting

for the bulk of the fitness interactions observed. We attributed these fitness changes as signatures

of susceptibility (fitness decreases) or resistance (fitness increases) with respect to microsporidia

infection.

So far, the causative gene for any host variants that influence microsporidia interactions has not

been identified. Previous genetic mapping in *C. elegans*, *D. magna*, and honey bees have

revealed 3-6 loci involved in resistance and sensitivity to microsporidia [19,23–26]. Additionally,

some loci in *D. magna* appear to be specific to different microsporidia species, which we show to

be the case for JU1400 [24]. Our genetic analysis of JU1400 identified that susceptibility to *N.*

ferruginous in both JU1400 and MY1 mapped to a single region on the left arm of chromosome I

while the resistance phenotype of JU1400 to *N. ironsii* appears to be caused by at least three

genomic regions. Taken together, our results add to the increasing evidence that suggests that

microsporidia have had a major impact on the evolution of animal genomes.

Microsporidia infect a wide diversity of animal phyla and infections are highly prevalent [12,13,61].

Nine species of microsporidia have been reported to infect *C. elegans*, with four having been

found in wild isolates and five capable of infecting *C. elegans* after isolation of microsporidia from

other nematode species [15,19,32,41]. The most common are the intestinal-infecting *N. parisii*

and *N. ausubeli,* which have been found in Europe, the United States, India, and Africa. The

geographical relationships between the strongest interactions we detected is not clear; JU1400

was isolated in Spain, *N. ironsii* in Hawaii, and *N. ferruginous* in France. Although host genetic

factors were shown to be responsible for the geographic range of a microsporidian infecting *D.*

magna, further work is needed to understand if there is a geographic basis to microsporidia

resistance and susceptibility in *C. elegans* [62].

Pathogen tolerance is being increasingly appreciated as a valuable way for hosts to improve

fitness in the presence of pathogens, although these mechanisms are less understood [63].

Tolerance is generally defined as the ability of a host to limit the consequences of a given level of

infection [64]. Here we show that JU1400 is very sensitive to infection by two different epidermal

infecting microsporidian species, but this strain does not display significantly higher levels of *N.*

ferruginous infection than non-sensitive control strains. RNA-seq analysis reveals that JU1400

has a unique response to microsporidia when infected with *N. ferruginous*. This response shares

similarities to several chemical toxins and pathogens that have been shown to damage *C.*

elegans. Although the mechanistic basis of this transcriptional similarity is likely different between

the various conditions, this result suggests that JU1400 infected with *N. ferruginous* has a

response similar to worms being exposed to toxic conditions. This signature is also detected to a

lesser extent in other strain-microsporidia pairings that cause developmental delay, including *N.*

ausubeli with either strain or *N. ironsii* with JU1400 [16]. Thus, one potential consequence of the

lack of tolerance to *N. ferruginous* is a developmental delay. Consistent with this result is the

observation of many genes containing collagen domains being down regulated in JU1400 infected

with *N. ferruginous*. Further work uncovering the genetic variant responsible for this aberrant

response will help elucidate the mechanistic basis of tolerance to microsporidia infection. None

of the differentially regulated genes are among the list of 12 candidate genes in the critical region

of chromosome I for *N. ferruginous* sensitivity, suggesting that causative gene or variant is not

differentially regulated during infection.

Our initial observation from our PhenoMIP experiment is that JU1400 has severely reduced

fitness when infected with *N. ferruginous* and a moderate reduction of fitness when infected with

N. ironsii (**Figure 3A**). In our follow up experiments using embryo counts as a proxy for fitness,

we observed that at 72 hours there was no relative difference in reduction of embryos during

infection with *N. ironsii* (**Figure 4A**) and significantly less reduction of embryos in JU1400 at 96

hours (**Figure 4B**). However, we also observed a decrease in embryos at 72 hours when JU1400

was infected with *N. ironsii* compared to N2 or VC20019 (**Supplementary Figure S4B**). We also

performed MIP-MAP experiments using similar conditions to those in our initial PhenoMIP

experiment. These experiments showed that JU1400 regions were selected for at both 72 hours

(**Supplementary Figure S13B**) and at 96 hours (**Figure 6B**). Regions on both chromosome II

and chromosome V were selected for at both times, but only the region on the X chromosome

was selected for at 96 hours. In addition to our experiments on host fitness, we also provide strong

evidence that there is resistance to *N. ironsii* in JU1400 due to its ability to clear the parasite.

Thus, our evidence is supportive of a model where JU1400 is resistant to *N. ironsii* infection and

has increased fitness compared to control strains at a 96-hour time point. Our data is inconclusive

about the impact of *N. ironsii* on JU1400 at 72 hours as we have conflicting results depending on

the experiment. Our results suggest that a limitation of our PhenoMIP experiment is that

resistance to microsporidia infection doesn't always result in a fitness advantage under certain

conditions. Our data also suggests that future competitive selection experiments using

microsporidia could be improved using 96-hour selections to identify resistant strains.

JU1400 is resistant to *N. ironsii* infection by clearing the invaded parasite. This immune

mechanism is similar to previous results seen in *N. ironsii* infection of CB4856. However, the

immunity in JU1400 results in less infection than in CB4856. The JU1400 resistance mechanism

is specific, as the clearance we observe does not occur in even the closely related sister species

N. parisii, which shares 92% identical DNA [43]. A strain of mosquito that is resistant to infection

by a parasitic nematode was shown to rely on activation of an upregulated immune response [65].

However, we see no evidence for an increased transcriptional response of JU1400 in response

to *N. ironsii*. A potential caveat to these experiments is that we infected JU1400 at the L1 stage,

but measured the transcriptional response at the L4 stage, a stage where if infections are initiated,

parasite clearance is not observed. We do, however, observe additional parasite clearance

between 24 and 72 hours (**Figure 5A**), which corresponds to between the L2/L3 stages and

adults, suggesting that if there was transcriptional signature of resistance, it would be observed

under our experimental RNA-seq conditions. Instead of a specific transcriptional response of

JU1400 to *N. ironsii*, we observe a *C. elegans* strain-specific response of immune genes including

upregulation of several CHIL genes, which have been observed to be upregulated in response to

epidermally infecting oomycetes, but have not previously been observed in response to

microsporidia infection [66,67]. This strain-specific-response of *C. elegans* has also been

observed with several bacterial pathogens, suggesting that there exists a diversity of innate

immune responses within the *C. elegans* population [68]. We also observe more strain-specific

regulated genes than pathogen-specific regulated genes, suggesting that *C. elegans* responds

similarly to infection by different species of *Nematocida* [69].

Here we demonstrate that PhenoMIP can be used to identify variants of pathogen infection in a

high-throughput manner. We compressed the equivalent of 2898 individual infection experiments

into 127 pooled population experiments, allowing a large savings in time and reagents required

for this study. The genomic loci responsible for these phenotypes can then be identified using

MIP-MAP [52]. We only tested a small fraction of the current *C. elegans* isotypes identified,

suggesting there is much variation to microsporidia within *C. elegans*. Our pooled selection

approach could also be applied to other *C. elegans*-related microbes such as those isolated from

the *C. elegans* microbiome or known pathogens and viruses [70,71]. Here we studied the animal's

reproductive fitness, but our approach could be modified, for instance, to select for pathogen load

using fluorescently labelled pathogens. The wild isolate collections of *C. elegans* provide a unique

opportunity to study how animals evolve in response to infection. The *Caenorhabditis elegans*

Natural Diversity Resource (CeNDR) collection, hosts more than 1500 wild isolate strains across

550 isotypes [33]. Based on our initial results, many of these strains are likely to have fitness

interactions with microsporidia infection. Identifying additional sensitive or resistant strains would

drive the discovery and further elucidation of evolved mechanisms of host-responses in

coevolving pathogens.

Acknowledgements

We thank Hala Tamim El Jarkass and Keir Balla for providing helpful comments on the

manuscript. We thank John Calarco for the kind gift of plasmids to perform CRISPR and Robert

Waterston and Donald Moerman for the kind gift of reagents to perform PhenoMIP and MIP-MAP.

This work was supported by Canadian Institutes of Health Research grants no. 400784 and

461807 and an Alfred P. Sloan Research Fellowship FG2019-12040 to A.W.R. and by NIH grant

R35 GM146836 to R.J.L. Some strains were provided by the CGC, which is funded by NIH Office

of Research Infrastructure Programs (P40 OD010440) and we thank WormBase.

Author contributions

C. M. and A. W. R. conceived of the project. C. M., M. A. X., Y. C. W., and A.R.W. designed and

analyzed the experiments and cowrote the paper, with consolation from all the authors. C. M., M.

A. X., W.Z., and S. M.A. conducted experiments. R. J. L. provided novel reagents. A.W.R.

provided funding and mentorship.

Competing Interests

The authors declare that they have no competing interests.

Methods

Nematode maintenance and worm pooling

Nematode strains were maintained at 21°C on a modified version of nematode growth media

NGM) with a 1:1 ratio of Agar:Agarose to reduce burrowing by wild isolate strains. NGM plates

were seeded with OP50-1 bacteria grown to saturation overnight in lysogeny broth (LB) at 37°C

and concentrated to 10X. All wild isolate strains used (**Supplementary Data S1**) are available

from the CGC. The AWR133 mapping strain was generated from VC20019 by inserting a GFP at

I:2,851,135 using CRISPR-based homologous recombination [72]. Nematode strains were grown

continuously without any starvation or contamination for a minimum of two weeks before seeding

individual strains to two 15-cm NGM plates seeded with 1.5 ml of 40X concentrated OP50-1. After

96 hours, gravid adult populations were bleach-synchronized [73] and allowed to hatch into axenic

M9 media for ~16 hours at 20°C. Synchronized L1 samples were counted and equal numbers of

animals were mixed into a single population before redistributing ~13,000 L1s to each 15-cm

NGM experimental plate with a mixture of 1.5 ml of 40X OP50-1 and the appropriate amount of

spores as needed (**Supplementary Table S2**).

PhenoMIP competitive assays

PhenoMIP competitive assays were based on a previously described protocol [39]. We modified

this method to use bleach-synchronization between generations to directly measure egg-

production as an output of population fitness. At each generation, samples were bleach-

synchronized as gravid adults ~72 hours after plating L1s and incubating at 21°C. Bleached

embryos were then allowed to hatch over 18-22 hours while incubating on a shaker at 21°C in

axenic M9 media. Each sample was gently pelleted and 3 – 7 µl of each pellet was used to seed

the experimental plate for the next generation of animals. Overall, the competitive assay consisted

of four total cycles, with one infection generation and one uninfected (resting) generation per

cycle. For each infection cycle, 3 ml of 40X OP50-1 culture was mixed in 15-ml falcon tubes with

twice the indicated dose of spores used on each infection experiment before splitting the mixture

to each duplicate plate. Worm populations for each specific replicate were then added to the plate

and samples were distributed evenly across the NGM surface before drying. For each rest cycle,

worm pellets were plated onto 15-cm NGM plates previously seeded with 1.5 ml of 40X OP50-1

culture. Prior to bleach-synchronizing the 72h-resting generation, a subsample of the population

was collected and frozen as a genomic DNA source. Microsporidia doses were adjusted after the

first infection round in order to ensure high enough selection pressure throughout the population

without causing the populations to become completely infertile or developmentally arrested

(**Supplementary Table 2**) [74]. The initial infection round with *N. ausubeli* caused animals on the

high-dose plates to produce no progeny. Therefore, the medium-dose infection series became

the high-dose series at 53,000 spores/cm^2. A new medium-dose series was started using an

uninfected control replicate population from the first rest phase. This new medium-dose series

(39,800 spores/cm^2) therefore only experienced three rounds of infection selection of *N. ausubeli*.

PhenoMIP sequencing and analysis

PhenoMIP sequencing libraries were generated using previously designed MIPs for the wild

isolate strains from the Million Mutation Project [39]. Individual MIPs were normalized to a

concentration of 100 μM and pooled to a maximum volume of 85 μl. 10 μl of 10X Polynucleotide

Kinase (PNK) Buffer and 5 μl of PNK were added to a volume of 85 μl pooled MIPs before

incubating for 45 minutes at 37°C and 20 minutes at 80°C. This pool was then diluted to a working

concentration of 330 nM. Libraries were generated using 160 ng of genomic DNA from each

sample and generated as previously described [52]. Libraries were sequenced on an Illumina

NextSeq system with each PhenoMIP library ranging between 0.0882×10^6 and 2.723×10^6 reads

and a mean of 9566 ± 7994 reads per probe.

For each sequencing library, reads were initially analysed as previously described [52] with the

exclusion of the normalization step for each MIP. After the abundance of each MIP was calculated,

an average abundance was calculated for each strain as well as a standard deviation across this

average. These values were used in downstream analysis of population structure across multiple

timepoints.

Population structure and fold-change analysis was calculated across each experiment using the

amalgamated data from above. Strains with a starting abundance value below 5×10^{-3} (CB4854

and JU312) were eliminated from downstream population analysis. Remaining data were further

transformed with any values below 1.0×10^{-3} being converted to this value to accommodate $\log_2$

growth analysis. Total fold-change and mean fold-change were calculated based on starting and

end-point changes in abundance versus total generations (two generations per infection cycle).

In samples with negative trajectories, however, the final generation of growth was calculated as

the first instance of abundance at or below the lower limit of 1.0×10^{-3}. Mean fold-change rate was

calculated based on the total fold-change abundance in the final generation of growth divided by

the expected number of generations passed. Each experimental condition may have had multiple

biological replicates (referred to in text as replicates), each with a time series of samples. Each

fold-change rate calculated for a single time series represented a single replicate. Mean fold-

change rate for each strain was therefore a combination of multiple replicates across multiple

experimental conditions (**Supplementary Data S1**).

Given the nature of the smaller datasets and the expectation of potential outliers stretching out

the tails of the mean FCR distribution, modified z-scores were calculated using the following

formula where $\tilde{Y}$ is the median of the data and $MAD = median(|Y_i - \tilde{Y}|)$

554
$$M_i = \frac{0.6745(x_i - \tilde{x})}{MAD}$$

To identify outliers and potential interaction candidates, we used a minimum z-score of ± 1.5 as

our threshold where both replicate samples were each required to meet the threshold.

Microsporidia infection assays

Continuous microsporidia infection assays were conducted using ~1,000 bleach-synchronized

L1s obtained from cultures rotating at 21°C for 18-22 hours. Worm populations were pre-mixed

in a 1.5 ml microfuge tube with 400 μl of 10X OP50-1 and microsporidia spores (**Supplementary**

Table S2). Samples were then spread onto individual unseeded 6-cm NGM plates and allowed

to   dry   before   incubating   at   21°C.   Plate   populations   were   imaged   using   a

Leica S9i Stereomicroscope before fixing samples for microscopy. Animals were washed in

M9/0.1% Tween-20 at least three times before fixing with acetone. Chitin staining for

microsporidia and embryos was performed using Direct Yellow 96 [75]. Meronts and sporoplasms

were visualized using FISH probes conjugated to CAL Fluor Red 610 (LGC Biosearch

Technologies). *N. ferruginous* was stained using 3 ng/ul of a mix of three CAL Fluor Red 610

probes   designed   to   target   the   18S   rRNA: GTTGCCCGCCCTCTGCC,

CTCTGTCCGTCCTCGGCAA, and CGGTCTCTAATCGTCTTC. For pathogen-specific staining,

MicroF-CAL Fluor Red 610 (AGACAAATCAGTCCACGAATT) was used for *N. parisii* and

Microsp1A-FAM (CAGGTCACCCCACGTGCT) was used for *N. ausubeli*. Unless otherwise

noted, all remaining FISH probes used 5 ng/µl of MicroB, a *Nematocida* microsporidia 18S rRNA

sequence: CTCTCGGCACTCCTTCCTG [15]. Stained samples were mixed with Biotium

EverBrite mounting medium with DAPI before mounting on slides for quantitation.

Embryo analysis and normalization

Unless otherwise noted, embryo analyses were completed using normalized values which were

calculated in the following manner. For each replicate experiment, the uninfected control for each

strain was used to establish a baseline mean embryo value ($\bar{Y}$). Embryo counts for individual

animals (Y) of the same strain within the same replicate experiment were normalized ($Y/\bar{Y}$) before

combining with the normalized values from matching replicate experiments. This normalization

step facilitated the analysis of relative changes to embryo production, rather than absolute

changes in embryo counts which could vary both between nematode strains and replicate

experiments. The combined normalized populations were compared using two-way ANOVA with

a Tukey HSD post hoc.

Microscopy and image quantification

All imaging was done using an Axio Imager.M2 (Zeiss) and captured via ZEN software under

identical exposure times per experiment. Animals containing clumps of spores visible in the body

after 72 hours by DY96 straining were considered as 'new spore formation'. Animals with one

sporoplasm or more in intestinal cells stained with FISH were considered infected. Gravidity of

worms was determined by counting the number of embryos per hermaphrodite animal.

Pathogen burden and body size was determined using ImageJ/FIJI software [76]. Individual

worms were outlined as "region of interest" with selection tools. Body size of worms were

determined using the "measure area" function. To quantify fluorescence (pathogen burden),

signal from FISH staining was subjected to the "threshold" function followed by "measure percent

area" tool.

Pulse-chase assays

Pulse-chase assays were performed on 6-cm NGM plates with roughly 6,000 bleach-

synchronized L1s obtained from cultures rotating at 21°C for 18-22 hours. Worm populations were

first premixed in a 1.5 mL microcentrifuge tube with 5 µL of 10X OP50-1 and 1.25 million spores

per microsporidia strain. This dose was chosen to produce N2 controls with 85-90% infection

rates at 3 hpi. Samples were then spread onto individual 6-cm NGM plates and allowed to dry

before incubating at 21°C for 3 hours. To subsample the population at 3 hpi, plates were washed

at least two times with M9 in a 1.5 mL microcentrifuge tube to remove OP50-1 and residual

spores. Half the population was then fixed in acetone and stored at -20°C. The other half was

mixed with 50 µL of 10X OP50-1 and plated onto 6-cm NGM plates. Samples were then incubated

at 21°C for 21 hours (24 hpi), washed, subsampled, fixed, and replated (with 300 µL of 10X OP50-

1) in the same manner as described above. The plates were then incubated at 21°C for 48hours

(72hpi) before washing and fixing in acetone. Pulse-chase co-infection assays were performed

as described above, except samples were infected with either one or two species of microsporidia

for 24 hours total. Pulse-chase assays initiated at the L4 stage were performed by using ~3000

bleach-synchronized L1s mixed with 200 µL of 10X OP50-1 and plated on unseeded 6-cm NGM

plates for 48 hours. Worm populations were then washed two times with M9 in 1.5 mL

microcentrifuge tubes. 50 µL of 10X OP50-1 and spores were added to each worm population

and then plated on unseeded 6-cm NGM plate for 3 hours at 21°C. Samples were then

subsampled, washed, and replated as described above. Chitin and FISH staining were performed

as described above for all pulse-infection experiments.

MIP-MAP and variant analysis

Wild isolates were mapped using a competitive fitness approach after crossing with males of the

mapping strain AWR133, a CRISPR-inserted Pmyo-2::GFP version of VC20019. Mapping was

completed using 2-4 competitive selection cycles on 10-cm NGM plates. Each cycle consisted of

2 generations of recombinant populations, each added to plates as bleach-synchronized L1

animals. In the first generation ~10,000 L1 cross-progeny animals were exposed to continuous

microsporidia infection (See **Supplementary Table S3**) for ~72 hours before bleach-

synchronizing. Next, ~8,000 L1s were plated within 24 hours to "rest" on 10-cm OP50-seeded

NGM plates and grown for ~72 hours before being bleach-synchronized. L1 progeny of the "rest"

phase were then plated onto both an OP50-1-seeded plate with spores to begin the next cycle

and a secondary plate without spores to generate a sample to collect genomic DNA. Genomic

samples were prepared with 250 ng of genomic DNA and VC20019 MIP-MAP library probes as

previously described [52]. Sequencing libraries were run on an Illumina NextSeq, with 161,054-

801,840 reads per library and a mean of 4005 ± 2760 reads per MIP across the entire data set.

Near isogenic lines construction

Near isogenic lines were tracked across chromosome I using four markers located at four

locations matching sequenced variants identified in JU1400: cewivar00066519 (491,763),

cewivar00067570 (1,051,810), cewivar00069114 (1,567,450), and cewirvar00070489

(2,012,869). These markers were used for restriction fragment length polymorphism analysis with

the enzyme DraI. JU1400 hermaphrodites were crossed with AWR133 males. F1 GFP+ animals

at the L4 stage were chosen to produce F2 animals. 48 GFP+ animals were chosen and screened

at marker cewivar00066519 for JU1400 homozygosity and further screened at cewirvar00070489

for JU1400 homozygosity or heterozygosity. Seven F2 lines were chosen based on their

genotyping and four of these gave rise to five main progenitor NILs, each of which was used to

create a separate GFP+ and GFP- NIL strain. In total 10 NILs were generated and tested for

susceptibility against LUAm1. One line was chosen for outcrossing (AWR140) which gave rise to

the AWR144 and AWR145 NILs.

C. elegans **Whole Genome Sequencing Analysis**

Worm strains used for whole genome sequencing were grown on 10-cm plates with OP50-1 for

3 days and allowed to starve. Plates were then washed at least two times with M9 in

microcentrifuge tubes before freezing in liquid freezing solution and then stored at -80C. gDNA

was then extracted and quantified with Qubit™ HS system. gDNA quality was determined by

agarose gel electrophoresis and photometric OD260/280 of 1.8-2.0. gDNA was submitted to The

Centre for Applied Genomics (TCAG, Hospital for Sick Children, Toronto, ON, Canada) for library

preparation and whole genome sequencing on an Illumina HiSeq 2500 SR50 Rapid Run flowcell

in pools of six samples per lane. Raw sequencing reads of each NIL were aligned to the N2

reference of the WBcel215 release (accession: PRJNA13758) using bowtie2 [77]. Then, samtools

was used to convert it into .bam format and freebayes was used to perform variant calling with

the default settings [78,79]. The breakpoints between VC20019 and JU1400 in chromosome I of

each NIL were computed and visualised using vcf-kit [80].

RNA extraction and mRNA sequencing

Infection samples for mRNA sequencing were prepared by mixing ~10,000 bleach-synchronized

L1 stage animals with 1 mL of 20X OP50-1 and 10 million *N. parisii* spores, 10 million *N. ausubeli*

spores, 10 million *N. ironsii* spores, and 30 million *N. ferruginous* spores in four separate

microcentrifuge tubes. Samples were then spread onto individual 10-cm NGM plates and allowed

to dry before incubating at 21°C for 48 hours. Samples for RNA extraction were washed in

M9/0.1% Tween-20 in 15 mL conical tubes at least three times before adding 1 mL of TRI-

Reagent® (Sigma-Aldrich). Total RNA extraction and ethanol precipitation protocols was

performed as per Tri-Reagent® manufacturers instruction. Libraries were prepared using

NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina® kit as per manufacturer's

instruction. RNA sequencing was carried out using Illumina NovaSeq-SP flow cell by The Centre

for Applied Genomics (TCAG).

Microsporidia RNA-seq analysis

Using Bowtie2 v2.4.4, indexes were built from *Nematocida parisii* strain ERTm1 (Genbank:

GCF_000250985.1), *Nematocida ausubeli* strain ERTm2 (Genbank: GCA_000738915.1),

Nematocida parisii strain ERTm5 (Genbank: GCA_001642415.1) and *Nematocida ferruginous*

strain LUAm1 (Accession: JALPMW000000000) The genome annotation file of each *Nematocida*

species was converted into gtf format using Gffread v0.12.3. Paired-end reads were aligned to

corresponding index and genome annotation files using TopHat v2.1.2. Cufflinks v2.2.1 was used

to generate a transcriptome assembly for each sample, which was then combined with the

corresponding reference genome annotation using Cuffmerge of the Cufflinks 2.2.1 package to

produce a single annotation file.

***C. elegans* mRNA-seq analysis**

The paired end reads generated for each strain were submitted as separate projects to Alaska

v1.7.2 (http://alaska.caltech.edu), which is an online automated RNA-seq analysis tool in

collaboration with Wormbase. Briefly, Bowtie2 [77], Samtools [78], RseQC [81], FastQC, and

MultiQC [82] were used for quality control of the input files. Kallisto [83] was used for the read

alignment and quantification, followed by differential analysis by Sleuth. For both N2 and JU1400

samples, the reads were aligned to the N2 reference of the WS268 release (accession:

PRJNA13758).

Principal component analysis

689   The N2 and JU1400 normalized abundance measurements generated by Alaska, were read by

690   the readRDS() function of the R package base v4.1.1. Normalized counts for each gene in each

691   strain were generated via sleuth_to_matrix() with the "obs_norm" data and "tpm" units. PCA of

692   gene expression was performed using samples from each strain, and merged samples, via the R

693   package pcaexplorer v2.20.2 [84].

Differentially expressed genes of N2 and JU1400 hosts

Log_2 fold change values of duplicated genes in each of the eight samples were averaged using

the R package dplyr v1.0.8. Genes with FDR-adjusted p-value of <0.01 were regarded as

significant. Significant genes expressed in at least three out of the eight samples were used for

hierarchical clustering via the function hclust() from the R package stats v4.1.1, then plotted using

R packages ggdendrogram and ggplot2 v3.3.5. Differentially upregulated genes were defined as

those with an FDR-adjusted p-value<0.01 and log_2 fold change >0 (infected vs. control);

differentially downregulated genes were defined as those with an FDR-adjusted p-value <0.01

and log_2 fold change <0 (infected vs. control).

Strain-specific, pathogen-specific and strain-pathogen pair specific genes

Strain-specific genes were defined as genes that are significantly upregulated or downregulated

in at least three out of the four JU1400 or N2 samples. Pathogen-specific genes were defined as

those significantly upregulated or downregulated in both JU1400 and N2 upon infection by the

same microsporidia species, but not observed when infected by other microsporidia species.

Moreover, strain-pathogen pair specific genes were defined as genes that are significantly

upregulated or downregulated in a particular strain-pathogen sample, but not observed in the

other seven strain-pathogen samples.

Domain enrichment analysis

We chose to look at F-box, MATH/BATH, PALS, DUF684 and DUF713 domains containing

genes, implicated from differentially regulated genes of *N. parisii* infected *C. elegans* in previous

publications [54,85]. Lists of gene names from respective gene classes were downloaded

separately from Wormbase (http://wormbase.org/) and the corresponding protein-coding

sequences were extracted from Wormbase ParaSite (https://parasite.wormbase.org/).

Alignments of other domains or classes were downloaded directly from Pfam

(http://pfam.xfam.org/) separately. After deleting duplicated sequences, protein sequences of

these classes of genes were converted to Stockholm format using Clustal Omega Multiple

Sequence Alignment tool (https://www.ebi.ac.uk/Tools/msa/clustalo/). Hmmbuild of HMMER

v3.3.2 [86] used the Stockholm format files to build respective profile HMMs, which was then

searched against *C. elegans* (accession: PRJNA13758) proteome of WS268 release using

hmmsearch. Output genes with E-value <1e-5 were used as the gene list for the enrichment

analyses. In the output genes, MATH and BATH genes were combined into one list; fbxa, fbxb

and fbxc genes were merged into the gene list for F-box (**Supplementary Table S10**). Using the

R package gplots v3.1.1, the number of genes overlapping between the lists and samples were

computed, then plotted using R package ggplot2 v3.3.5. Genes that do not fall into any of these

gene classes or have any of these domains were categorised as "other".

Gene Set Enrichment Analysis

The WormExp [56] dataset, which contains published expression data for *C. elegans*, was

downloaded from the WormExp v1.0 website (https://wormexp.zoologie.uni-kiel.de/wormexp/).

The overlap between published datasets of *C. elegans* exposed to different chemicals or

pathogens (**Supplementary Table S12**) and our samples, as well as the strain specific,

pathogen-specific, and strain-pathogen pair specific genes in our dataset were computed using

the R package gplots v3.1.1. For the statistical evaluation of the overlap, the p-values were

calculated using the adjusted Fisher exact test method from the program EASE, which removes

one gene in the set of intersected genes. Correction for multiple testing was implemented using

the Bonferroni method via the p.adjust() function in the R package stats v4.1.1.

GO Biological Process, GO Molecular Function and KEGG pathway enrichment analyses

Using the R package enrichR v3.0 [87], the connection to WormEnrichr, a gene list enrichment

analysis tool for *C. elegans*, was set through the setEnrichrSite ("WormEnrichr") function. The

"GO_Molecular_Function_2018", "GO_Biological_Process_2018" and "KEGG_2019" databases

were selected then used to query enrichr with the enrichr() function, before being plotted as

Enrichr GO-BP output **(Supplementary Tables S13 and S14)**.

Volcano plots

Values of duplicated genes in each of the four host-microsporidia samples were averaged using

the R package dplyr v1.0.8. Genes with FDR<0.05 and $\log_2$ Fold Change >0 or $\log_2$ Fold Changes

<0 were categorised as differentially upregulated or downregulated respectively. The volcano

plots were generated by ggplot2 v3.3.5 in R.

Data and code availability

All samples were deposited under NCBI BioProject PRJNA841603 and the sequence reads for

all samples were submitted to the NCBI Sequence Read Archive. Data from all the infection

experiments is in **Supplementary Data S1**. PhenoMIP analysis scripts can be found at

https://github.com/camok/PhenoMIP.

Figures

Figure 1. A PhenoMIP high-throughput screen to identify wild strains that are either resistant or susceptible to microsporidia infection. (A) N2 L1 animals that were uninfected (i) or exposed to either 3.5 million *N. parisii*, *N. ironsii*, or *N. ausubeli* spores or 1.25 million *N. ferruginous* spores for 72 hours, were fixed and stained with the following 18S FISH probes (i, ii,

iv; MicroB-CAL Fluor Red 610, iii; Microsp1A-FAM and v; a combination of three Fluor Red 610

probes described in the Methods). Scale bars, 100 μm. (B) PhenoMIP screening protocol for

microsporidia requires that separate unstarved populations for each of the 22 wild isolate strains

as well as VC20019 (lab reference-derived strain) be sodium hyplochlorite-synchronized (i) before

pooling in equimolar amounts (ii) and infecting (iii) with different microsporidia species at doses

that can impose fitness challenges to the lab reference strain. Populations are grown for 72 hours

post-infection (iv) before sodium hypochlorite-synchronizing (v) and growing free of infection for

an additional 72 hours (vi). Uninfected populations are harvested, subsampled for genomic DNA,

and sodium hypochlorite-synchronized (vii) before repeating the infection and rest phases for an

additional three rounds.

Figure 2. Analysis of PhenoMIP mean fold-change rate identifies 16 candidate

nematode/microsporidia interactions. Violin plots of *C. elegans* wild isolate strains infected

by specific microsporidia species. Violins are derived from modified z-scores of mean fold-

change rate (FCR) per generation using the eight mock-infection replicates of that strain.

Overlayed are the six infection replicates of low (green circles), medium (blue triangles), and

high (purple diamonds) doses for the microsporidia species indicated in each subplot title. Mock

infection replicates are also overlayed (yellow squares). Red dotted lines represent the

candidate hit threshold for modified z-scores of ±1.5.

Figure 3. JU1400 lacks tolerance to *N. ferruginous* infection. (A-B) Line graph of fold change

across multiple time points and conditions for JU1400 (A) and MY1 (B). Microsporidia strains are

denoted by point shape and doses of microsporidia are denoted by line type. The eight uninfected

control replicates are bounded within the greyed area and the mean of the uninfected control is

denoted by a solid red line. All other lines represent mean FCR of two biological replicate samples.

(C-D) Strains were infected for 72 hours with *N. ferruginous*. (C) Dissecting microscope images

of live animals growing on plates. (D) Dot plot of normalized embryo counts of animals fixed and

stained with DY96 with mean and SEM bars. Data is combined from three biological replicates

with n={41,50} worms per replicate sample. (E-F) N2 and JU1400 animals were infected for three

hours with 1.25 million spores of *N. ferruginous* before washing away uneaten spores from each

population. Animals were fixed after 3 hours (E) or after 72 hours (F). (E) Dot plot of sporoplasms

counted in N2 and JU1400 animals. Data is combined from three biological replicates with n=40

worms per replicate sample. (F) Bar plot with SEM bars depicting mean parasite burden as a

function of percent fluorescent area per animal by FISH staining in N2 and JU1400 populations.

Data is combined from three biological replicates with n={29, 53} worms per replicate sample. p-

values were determined by two-way ANOVA with Tukey post-hoc. Significance was defined as p

≤ 0.001 (***) and not significant as $p > 0.05$ (ns).

Figure 4. JU1400 is resistant to *N. ironsii* infection. (A-D) N2 and JU1400 were infected with
N. ironsii for 72 hours (A and C) or 96 hours (B and D), fixed, and stained with DY96. (A-B) Dot
plots of normalized embryo counts of animals fixed and stained with DY96 with mean and SEM
bars. Data is combined from three biological replicates with n={47,50} worms per replicate
sample. (C-D) Bar plot with SEM bars of percent population with new spore formation from three
combined replicate experiments. p-values were determined by two-way ANOVA with Tukey post-
hoc. Significance was defined as $p \leq 0.001$ (***) and not significant as $p > 0.05$ (ns).

Figure 5. JU1400 specifically clears *N. ironsii* infection at the earliest larval stage. (A-B) N2

and JU1400 animals were pulse infected for 3 hours with *N. ironsii* before washing away excess

spores. Animals were pulse infected at either the L1 stage and fixed at 3-, 24-, and 72 hpi (A) or

animals were pulse infected at the L4 stage before fixing at 3- and 24 hpi (B). Samples were then

stained with an *N. parisii* 18S RNA FISH probe and animals containing either sporoplasms or

meronts were counted as infected. Bar plot with SEM bars show percent of animals infected per

replicate from three biological replicates with n=20 worms per replicate. (C) CB4856 and JU1400

were infected at the L1 stage with either medium or high doses of *N. ironsii* for 72 hours and then

fixed and stained with DY96. (D) N2, JU1400, and CB4856 were infected at the L1 stage with

either *N. parisii*, *N. ausubeli*, or *N. ironsii* for 72 hours and then fixed and stained with DY96. (C-

D) Bar plot with SEM bars show percent of animals generating spores per replicate from three

biological replicates with n=20 worms per replicate. (E) JU1400 animals were pulse infected for 3

hours with *N. parisii*, *N. ausubeli*, *N. ironsii, N. parisii + N. ausubeli*, or *N. ausubeli + N. ironsii*

before washing away excess spores. Animals were either fixed immediately at 3 hpi or at 24 hpi.

Samples were then stained with species-specific 18S rRNA FISH probes and animals containing

either sporoplasms or meronts were counted as infected. Bar plot with SEM bars show percent

of animals infected per replicate from three biological replicates with n={78, 147} worms per

replicate. p-values were determined by two-way ANOVA with Tukey post-hoc. Significance was

defined as $p \leq 0.01$ (**), $p \leq 0.001$ (***), and not significant as $p > 0.05$ (ns).

Figure 6. Genetic mapping of JU1400 host-response phenotypes identifies distinct loci.
Genetic mapping of JU1400 susceptibility to *N. ferruginous* (A) and JU1400 resistance to *N. ironsii*
(B) was performed with MIP-MAP using the AWR133 mapping strain. Green square-points and
dotted lines represent a mock-infection replicate with the solid blue and orange lines representing
two biological replicates exposed to specific infection conditions. Red arrows indicate candidate

regions of interest based on the expected genome fixation direction during phenotype selection.

(C) Whole-genome sequencing haplotype map of chromosome I NILs generated from crosses

between JU1400 and AWR133. Regions of chromosome I for specific NILs are coloured by their

inferred haplotype. Strain names are indicated to the left of each with AWR138, AWR140, and

AWR145 sharing a susceptibility phenotype (Red text) while AWR144 is not susceptible (Blue

text) to *N. ferruginous* infection. (D) N2, JU1400, AWR144, and AWR145 L1 stage animals were

infected with *N. ferruginous* before being fixed at 72 hpi and stained with DY96 to visualize

embryos. Dot plot of normalized embryo counts of with mean and SEM bars. Data is combined

from three biological replicates with n=50 worms per replicate sample. (E) N2, JU1400, AWR144,

and AWR145 L1 stage animals were infected with *N. ironsii* before being fixed at 72 hpi and

stained with DY96 to visualize mature spores. Bar plot with SEM bars show percent of animals

generating new spores per replicate from three biological replicates with n=50 worms per

replicate. p-values were determined by two-way ANOVA with Tukey post-hoc. Significance was

defined as $p \leq 0.001$ (***) and not significant as $p > 0.05$ (ns).

Figure 7. RNA-seq analysis of N2 and JU1400 infected with four species of microsporidia.

N2 and JU1400 animals were infected with one of four species of microsporidia (see Methods for

doses) for 48 hours and RNA was extracted and sequenced. (A) Principal component analysis of

uninfected and infected N2 and JU1400 samples. Circles represent confidence ellipses around

each strain at 95% confidence interval. (B) Heatmap of transcriptional profiles of differentially

regulated genes in N2 and JU1400 samples. Rows represent genes clustered hierarchically.

Scale is of differential regulation of infected compared to uninfected samples. Grey tile indicates

that the gene is not significantly differentially regulated in the sample. (C-D) Domain enrichment

analysis of significantly upregulated and downregulated genes in each sample (C) and of specific

gene sets (strain, pathogen, and strain-pathogen pair) (D). (E-F) Published datasets of

differentially expressed *C. elegans* genes upon exposure to chemicals or bacterial infection were

compared for overlap with the downregulated genes from each sample (E) or for overlap between

of differentially expressed genes in strain specific, pathogen specific, or strain-pathogen sets (F).

(A-F) uni (uninfected), npa (*N. parisii*), nau (*N. ausubeli*), nir (*N. ironsii),* and nfe (*N. ferruginous)*.

Upregulated samples are represented with orange text; downregulated samples are represented

with pink text.

Figure S14. JU1400 and MY1 susceptibility to *N. ferruginous* is recessive and MY1 is less

resistant than JU1400 to *N. ironsii*.

Figure S15. Generation of NILs carrying the chromosome I region that is responsible for

susceptibility to *N. ferruginous*.

Figure S16. Microsporidia mRNA reads per sample demonstrates JU1400 resistance to *N.*

***ironsii*.**

Figure S17. Principal component analysis of N2 and JU1400 samples.

Figure S18. Comparison of correlation between N2 and JU1400 strains after infection with

different microsporidia species.

Figure S19. Volcano plots of differentially regulated genes.

Figure S20. Overlap between the significant upregulated genes upon microsporidia

infection and differentially expressed datasets of *C. elegans* genes upon exposure to

chemicals or bacterial infection.

Table S1. Microsporidia species information.

Table S2. PhenoMIP microsporidia infection doses.

Table S3. Wild isolate mapping microsporidia infection doses.

Table S4. NIL strain chromosome I region composition.

Table S5. Candidate genes found in chromosome I region for *N. ferruginous* sensitivity.

Table S6. Microsporidia gene FPKM values in N2 and JU1400.

Table S7: Normalized *C. elegans* gene counts for all RNA-seq replicates from this study.

Table S8: Results of *C. elegans* differential expression analyses for all RNA-seq
comparisons performed in this study.

Table S9. Gene classes and domains used for enrichment analyses.

Table S10. The number of differentially regulated genes that contain enriched domains.

Table S11. Published data sets from WormExp used in this study.

Table S12. List of gene set overlaps from this study to published data sets.

Table S13. GO Biological Processes, GO Molecular Functions, and KEGG Pathways
enriched in N2 genes upregulated or downregulated upon microsporidia infection.

Table S14. GO Biological Processes, GO Molecular Functions, and KEGG Pathways
enriched in JU1400 genes upregulated or downregulated upon microsporidia infection.

Data S1. Data of all infection experiments.

A

937 **Figure S1. Geographical locations of *C. elegans* and microsporidia used in this study.** (A)

938 World map highlighting the locations where the *C. elegans* wild isolate strains (orange circles),

939 *Nematocida* species (green triangles), and the VC20019 reference strain (purple star) used in this

940 study were originally isolated.

Figure S2. Analysis of PhenoMIP mean fold-change rate identifies candidate nematode/microsporidia interactions. Violin plots of 20 wild isolates and lab reference-derived control strain VC20019 showing modified z-scores of mean fold-change rate (FCR) per generation. Violin shapes are derived from the eight mock-infection replicates. Overlayed are the eight mock infections replicates and six infection replicates for each of *N. parisii* (A), *N. ausubeli* (B), *N. ironsii* (C), and *N. ferruginous* (D). Replicates are coloured by mock infection (yellow), low dose (green), medium dose (blue), and high dose (purple) with increasing point size associated with increasing dose. Dotted lines represent the z-score candidate threshold of ±1.5. The strains CB4854 and JU312 were excluded from analysis due to low initial population abundance.

Figure S3. The Hawaiian (CB4856) strain shows resistance to infection by high-dose *N. ironsii*. Line graph of fold change across multiple time points and conditions for CB4856 demonstrates potential fitness increase relative to mock-infection conditions when infected with a high dose of *N. ironsii*. Microsporidia strains are denoted by point shape and doses of microsporidia are denoted by line type. The eight uninfected control replicates are bounded within the greyed area with the mean of the uninfected controls denoted by a solid red line. All other lines represent mean FCR of two biological replicate samples.

Figure S4. Single-strain testing of PhenoMIP interaction candidates with *N. ferruginous*

infection. (A-C) Strains of *C. elegans* were infected with medium and high doses of *N. ferruginous*

for 72 hours, imaged, fixed, and stained with DY96 and a *N. ferruginous* 18S rRNA FISH probe.

(A) images of N2 and VC20019 controls versus AB1, JU397, ED3042, and JU360 at 72 hpi. (B)

Dot plot with mean and SEM bars of normalized embryo counts from infected populations with

n={41,70} worms per sample. (C) Bar plot depicting the percent of animals with detectable meront

967   staining. p-values denote comparison to N2 controls and were determined by two-way ANOVA

968   with Tukey post-hoc. Significance was defined as $p \leq 0.001$ (***) and not significant as $p > 0.05$

969   (ns).

Figure S5. Single-strain testing of PhenoMIP interaction candidates with *N. ironsii* infection. (A-C) Strains of *C. elegans* were infected with medium and high doses of *N. ironsii* for 72 hours, imaged, fixed, and stained with DY96. (A) images of N2 and VC20019 controls versus CB4856, JU300, JU1400, and MY2 at 72 hpi. (B) Dot plot with mean and SEM bars of normalized embryo counts from infected populations with n={45,115} worms per sample. (C) Bar plot depicting the percent of animals with newly generated spores. p-values denote comparison to N2

978   controls and were determined by two-way ANOVA with Tukey post-hoc. Significance was defined

979   as $p \leq 0.001$ (***), $p \leq 0.01$ (**), $p \leq 0.05$ (*) and not significant as $p > 0.05$ (ns).

Figure S6. Single-strain testing of PhenoMIP interaction candidates with *N. ausubeli* infection. (A-C) Strains of *C. elegans* were infected with medium and high doses of *N. ausubeli* for 72 hours, imaged, fixed, and stained with DY96. (A) images of N2 and VC20019 controls versus JU360 and MY2 at 72 hpi. (B) Dot plot with mean and SEM bars of normalized embryo counts from infected populations with n={54,71} worms per sample. (C) Bar plot depicting the

987   percent of animals with newly generated spores. p-values denote comparison to N2 controls and

988   were determined by two-way ANOVA with Tukey post-hoc. Significance was defined as $p \leq 0.05$

989   (*) and not significant as $p > 0.05$ (ns).

Figure S7. JU1400 and MY1 share similar response phenotypes to infection by _N._

**ferruginous****.** JU1400, MY1, and N2 L1s were infected with a medium dose of _N. ferruginous_ for

72 hours, fixed, and stained with _N. ferruginous_ 18S rRNA FISH probes and DY96. (A) Dot plot

with mean and SEM bars of embryo counts from infected populations. (B) Bar plot with SEM bars

depicting the percent of animals with visible meronts by FISH staining. Data is combined from

three biological replicates with n={41,50} worms per replicate sample. p-values denote

comparison to N2 controls and were determined by two-way ANOVA with Tukey post-hoc.

Significance was defined as p ≤ 0.01 (**), p ≤ 0.05 (*) and not significant as p > 0.05 (ns).

1001

1002

Figure S8. JU1400 and N2 display similar levels of *N. ferruginous* infection. (A-D) JU1400

and N2 L1s were infected with 1.25 million spores of *N. ferruginous* for 3 h, washed to remove

spores, and replated for an additional 69 h. Animals were fixed at 3 hpi and 72 hpi and stained

with *N. ferruginous* 18S rRNA FISH probes. (A) Representative images of infected animals at 72

hpi. Scale bars are 100 μm. (B) Bar plot with SEM bars depicting the percent of animals displaying

either sporoplasm (3 hpi) or meronts (72 hpi) by FISH staining. (C-D) quantitation of FISH stain

at 72 hpi. (C) Dot plot with mean and SEM bars of parasite burden by area in infected populations

with a minimum of n={41,50} worms per sample. (D) Bar plot with SEM bars depicting mean

parasite burden as a function of percent fluorescent area per animal by FISH staining. Data in (C,

D) is combined from three biological replicates with n=40 worms per replicate sample. p-values

denote comparison to N2 or JU1400 controls and were determined by two-way ANOVA with

Tukey post-hoc. Not significant was defined as p > 0.05 (ns).

1015

1016

Figure S9. JU1400 displays a similar response to *N. ferruginous* strains and sensitivity to

N. cider. (A-D) N2 and JU1400 L1 stage animals were infected with medium doses of *N.*

ferruginous strain LUAm1 or LUAm3 (A-B) or a medium dose of *N. cider* (C-D) for 72 hours, fixed,

and stained with DY96 and 18S rRNA FISH probe. (A, C) Dot plot of normalized embryo counts

of animals with mean and SEM bars. (B, D) Bar plot with SEM bars of percent population with

meronts visible. Data are combined from three biological replicates with n=50 worms per replicate.

p-values were determined by two-way ANOVA with Tukey post-hoc. Significance was defined as

p ≤ 0.05 (*), p ≤ 0.001 (***), and not significant as p > 0.05 (ns).

1025

1026

1027

1028

1029

Figure S10. JU1400 is resistant to infection by *N. ironsii*. N2 and JU1400 L1 stage animals were infected with mock, medium, and high doses of *N. ironsii* for 72 (A and C) and 96 (B and D) hours and fixed. Embryos and mature spores are visualized by staining with DY96 (green), meronts are stained by a 18S rRNA FISH probe. (A-B) Representative images of high dose

infection from each experiment. Scale bars are 100 µm. (C-D) Bar plots with SEM bars of percent

population with meronts visible. Data in (C, D) is combined from three biological replicates each

with n={47,50} worms per replicate. (E-F) N2, JU1400, and CB4856 animals at the L1 stage were

infected for 3 hours with 1.25 million spores of either *N. parisii*, *N. ausubeli*, or *N. ironsii* before

washing away excess spores. Animals were either fixed at 3 hours (E) or replated for an additional

21 hours (F) and then fixed. Shown are bar plots with SEM bars of percent population with

meronts visible. Data in (E, F) are combined from three biological replicates each with n={59,179}

worms per replicate. p-values were determined by two-way ANOVA with Tukey post-hoc.

Significance was defined as $p \leq 0.05$ (*), $p \leq 0.001$ (***), and not significant as $p > 0.05$ (ns).

1043

1044

1045

1046

Figure S11. JU1400, but not N2, can specifically eliminate *N. ironsii* infection. N2 and JU1400 animals at the L1 stage were pulse infected for 3 hours with *N. parisii*, *N. ausubeli*, *N. ironsii, N. parisii + N. ausubeli, or N. ausubeli + N. ironsii* before washing away excess spores. Animals were either fixed immediately at 3 hpi or at 24 hpi. Samples were then stained with species-specific 18S RNA FISH probes (*N. ausubeli* = green; *N. parisii* and *N. ironsii* = magenta) and animals containing either sporoplasms or meronts were counted as infected. (A) Representative images of N2 and JU1400 infected with *N. parisii* and *N. ausubeli* (A) and *N. ironsii* and *N. ausubeli* (B). Scale bar 10 μm for 3 hpi and 25 μm for 24 hpi. Insets at top right for each

1055   image show an enlarged image of the outlined boxes. (C) Bar plot with SEM bars show percent

1056   of N2 animals infected per replicate from three biological replicates with n={59, 179} worms per

1057   replicate. p-values were determined by two-way ANOVA with Tukey post-hoc. Non-significance

1058   was defined as $p > 0.05$ (ns).

1059

1060

Figure S12. Genetic mapping of CB4856 resistance to *N. ironsii*. Genetic mapping of CB4856

resistance to *N. ironsii* was performed with MIP-MAP using the AWR133 mapping strain. Green

square-points and dotted lines represent a mock-infection replicate with the solid blue and orange

lines representing two biological replicates exposed to high dose infection conditions for 72-hours

per selection cycle. A red arrow on chromosome V indicates a strong signal on the right arm.

Additional red arrows on chromosomes II and IV indicate broad regions of interest across the

chromosomes that may contribute to CB4856 resistance phenotypes. Golden arrow indicates the

z*eel-1/peel-1* incompatibility locus at chromosome I.

1069

Figure S13. Genetic mapping of interaction phenotypes of JU1400 to *N. ironsii* and MY1 to

***N. ferruginous* using 72-hour selections.** Genetic mapping of (A) MY1 susceptibility to *N.*

ferruginous and (B) JU1400 resistance to *N. ironsii* was performed with MIP-MAP using the

AWR133 mapping strain. Green square-points and dotted lines represent a mock-infection

replicate with the solid blue and orange lines representing two biological replicates exposed to

specific infection conditions. Red arrows indicate candidate regions of interest based on the

expected genome fixation direction during phenotype selection. Golden arrow indicates the *zeel-*

1/peel-1 incompatibility locus at chromosome I.

1078

1079

1080

1081

1082

1083

Figure S14. JU1400 and MY1 susceptibility to *N. ferruginous* is recessive and MY1 is less resistant than JU1400 to *N. ironsii*. (A-B) Analysis of F1 progeny from crosses between

AWR133 (VC20019 with integrated pmyo-2::GFP) and the wild isolates JU1400 or MY1 as well

as F1 progeny from crosses between JU1400 hermaphrodites and MY1 males. F1s were

examined 72 hours post-infection by *N. ferruginous* for body size (A) and normalized embryo

counts (B). (B) Dot plot with mean and SEM bars of normalized embryo counts from infected

populations with a minimum of n={47,51} worms per sample. (C-D) L1 stage animals were

infected for 72 hours with *N. ironsii*, fixed, and stained with DY96 and an 18S rRNA FISH probe.

Bar plots depict the percent of animals with newly generated spores (C) or meront signal (D). p-

values were determined by two-way ANOVA with Tukey post-hoc. Significance was defined as p

≤ 0.001 (***), p ≤ 0.05 (*) and not significant as p > 0.05 (ns).

1095

1096

1097

1098

1099

1100

1101

1102

Figure S15. Generation of NILs carrying the chromosome I region that is responsible for susceptibility to *N. ferruginous*. A schematic (A) of 10 separate NILs generated through a cross between AWR133 and JU1400. These lines were infected with a medium dose of *N. ferruginous*

1107   for 72 hours (B) and their resulting body size phenotype was assessed in comparison to N2 and

1108   JU1400 control strains.

1109

1110

1111   **Figure S16. Microsporidia mRNA reads per sample demonstrates JU1400 resistance to *N.***

1112   ***ironsii*.** Scatterplot of overall read mapping rate of *Nematocida* in both strains with mean and

1113   standard deviation. uni (uninfected), npa (*N. parisii*), nau (*N. ausubeli*), nir (*N. ironsii),* and nfe (*N.*

1114   *ferruginous)*.

A

B

Infection: Infected Uninfected

1116 **Figure S17. Principal component analysis of N2 and JU1400 samples.** Principal component

1117 plots of expression data from only (A) N2 samples or (B) JU1400 samples. Circles represent

1118 confidence ellipses around each strain at 95% confidence interval. uni (uninfected), npa (*N.*

1119 *parisii)*, nau (*N. ausubeli*), nir (*N. ironsii),* and nfe (*N. ferruginous)*.

1120

1121

Figure S18. Comparison of correlation between N2 and JU1400 strains after infection with

different microsporidia species. Correlation between $\log_2$ fold changes in genes of N2 and

JU1400 animals infected with (A) *N. parisii* (B) *N. ausubeli* (C) *N. ironsii* (D) *N. ferruginous*.

Pearson's correlation value and p-value for each cluster (pink, green, blue, purple) are presented

in the upper left corner. Pearson's correlation value and p-value for all the genes are presented

in the bottom left corner. A gene with FDR adjusted p-value of <0.01 is deemed significant in each

strain.

1129

1130

1131

1132

1133

1134

1135

1137

Figure S19. Volcano plots of differentially regulated genes. (A) *N. parisii*-infected N2. (B) *N.*

parisii-infected JU1400. (C) *N. ausubeli*-infected N2. (D) *N. ausubeli*-infected JU1400. (E) *N.*

ironsii-infected N2. (F) *N. ironsii*-infected JU1400. (G) *N. ferruginous*-infected N2. (H) *N.*

ferruginous-infected JU1400. Each point represents a gene, orange points indicate differentially

upregulated genes, pink points are differentially downregulated genes while grey points represent

genes with FDR-adjusted p-value >0.01 (red horizontal line). The top five differentially

upregulated and downregulated genes with FDR-adjusted p-value <0.01 are labelled with their

respective gene names.

1146

1147

A

B

Bonferroni corrected p-values | 0.05~1 | 1e-10~0.05 | 1e-50~1e-10 | 1e-100~1e-50

1149 **Figure S20. Overlap between the significant upregulated genes upon microsporidia**

1150 **infection and differentially expressed datasets of** *C. elegans* **genes upon exposure to**

1151 **chemicals or bacterial infection.** (A) upregulated genes from microsporidia-infected samples.

1152 (B) upregulated gens from strain specific, pathogen specific, and strain-pathogen specific gene

1153 sets. uni (uninfected), npa (*N. parisii)*, nau (*N. ausubeli*), nir (*N. ironsii),* and nfe (*N. ferruginous)*.

1154

1155

1156

1157

1158

1159 **Table 1. PhenoMIP screen interaction hits**

Strain	Site of isolation	Microsporidia species	Relative Impact on host fitness	Modified z-score	Single infection embryo phenotype?
AB1	Adelaide, Australia	*N. ferruginous*	Increase	1.65, 1.64	Yes
JU397	Ouistreham, France	*N. ferruginous*	Increase	2.58, 2.59	Yes
LKC34	Antananarivo, Madagascar	*N. ferruginous*	Increase	2.14, 2.40	Untested
ED3040	Johannesburg, South Africa	*N. ferruginous*	Decrease	-1.90, -1.90	Untested
ED3042	Capetown, South Africa	*N. ferruginous*	Decrease	-2.53, -1.57	Yes
JU360	Franconville, France	*N. ferruginous*	Decrease	-2.68, -2.74	Yes
MY1	Munster, Germany	*N. ferruginous*	Decrease	-6.05, -6.05 -4.52[M], -4.25[M] -2.17[L] -1.85[L]	Yes
JU1400	Sevilla, Spain	*N. ferruginous*	Decrease	-6.43, -6.43 -3.33[M], -3.33[M]	Yes
CB4856	Hawaii, USA	*N. ironsii*	Increase	2.48, 2.88	Yes
JU300	Le Blanc, France	*N. ironsii*	Increase	2.19, 2.70	No
MY2	Munster, Germany	*N. ironsii*	Decrease	-2.83, -3.50	Yes
JU533	Primel-Tregastel, France	*N. ironsii*	Decrease	-2.00, -2.00	Untested
JU642	Paris, France	*N. ironsii*	Decrease	-2.44, -3.06	Untested
JU1400	Sevilla, Spain	*N. ironsii*	Decrease	-1.78, -1.78	Yes
JU360	Franconville, France	*N. ausubeli*	Increase	2.03, 1.88	No
MY2	Munster, Germany	*N. ausubeli*	Decrease	-2.07, -2.42	No

1160 All modified z-scores are identified from high dose infections unless otherwise noted.

1161 [M] Medium dose infection modified z-score

1162 [L] Low dose infection modified z-score

1163

1164

 Table S1. Microsporidia species information.

Microsporidia species	Original host	Infected tissue	Geographical origin
Nematocida parisii [15]	*C. elegans*	Intestine	France
Nematocida ausubeli [15,32]	*C. briggsae*	Intestine	India
Nematocida ironsii [19,43]	*C. briggsae*	Intestine	Hawaii
Nematocida ferruginous [44]	*C. elegans*	Epidermis/muscle	France

1166

1167 **Table S2. PhenoMIP microsporidia infection doses.**

Microsporidia species	Microsporidia strain	Infection round	Spores per cm^2 media		
			Low	Medium	High
Nematocida parisii	ERTm1	1	8,000	16,000	32,000
		2-4	13,269	26,539	53,078
		Follow-up infections		26,539	53,078
Nematocida ausubeli	ERTm2	1	26,539	NA	53,078
		2-4	26,539	39,808	53,078
		Follow-up infections		17,692	53,078
Nematocida ironsii	ERTm5	1	22,116	44,232	88,464
		2-4	30,692	61,924	123,849
		Follow-up infections		61,924	123,849
Nematocida ferruginous	LUAm1	1	88,464	176,928	353,857
		2-3	159,235	318,471	636,942
		4	159,235	318,471	751,592
		Follow-up infections		318,471	751,592
	LUAm3	**Follow-up infections**		353,857	
Nematocida cider	AWRm77	**Follow-up infections**		141,542	

1168 * effective dose of spores used in retests and additional infections

1169

1170 **Table S3. Wild isolate mapping microsporidia infection doses.**

Wild isolate strain(s)	Microsporidia strain	Infection time (hours)	Infection round	Spores per cm^2 media
JU1400, MY1	*Nematocida ferruginous*	72	1-2	350,318
			3	445,859
JU1400	*Nematocida ironsii*	72	1	178,343
			2-3	254,777
JU1400	*Nematocida ironsii*	96	1-2	509,554

| CB4856 | *Nematocida ironsii* | 72 | 1 | 178,343 |
| | | | 2-3 | 254,777 |

Table S4. NIL strain chromosome I region composition.

Strain	I:491,763	I:1,051,810	I:1,567,450	I:2,012,869	GFP Status I-2,851,135
AWR134	JU1400	JU1400	JU1400	--	+
AWR135	JU1400	JU1400	JU1400	JU1400	-
AWR136	JU1400	JU1400	JU1400	JU1400	+
AWR137	JU1400	JU1400	JU1400	JU1400	-
AWR138	JU1400	--	--	--	+
AWR139	JU1400	JU1400	JU1400	JU1400	-
AWR140	JU1400	JU1400	--	--	+
AWR141	JU1400	JU1400	JU1400	JU1400	-
AWR142	JU1400	JU1400	--	--	+
AWR143	JU1400	JU1400	JU1400	JU1400	-

Table S9. Gene classes and domains used for enrichment analyses.

Domain/ Gene Family	Databse	Wormbase gene class OR Pfam ID
F-box	Wormbase	fbxa, fbxb, fbxc
MATH (meprin or Traf homology) or BATH (BTB and MATH domain-containing)	Wormbase	math, bath
PALS (protein containing ALS2CR12 signature)	Wormbase	pals
C-type lectins	Wormbase	clec
DUF713	Pfam	PF015218
DUF684	Pfam	PF05075
chil	Wormbase	chil
Histone	Pfam	PF00125
Nematode cuticle collagen N-terminal domain	Pfam	PF01484
skr (Skp1-related)	Wormbase	Skr
Other (Any genes not in the domain or gene family above)	-	-

Table S11. Published data sets from WormExp used in this study.

Chemical/Pathogen	WormExp Term
Cadmium	• down by Cadmium (Huffman)

	• UP by Cadmium (Huffman)
Dichlorvos	• Down by high dichlorvos • UP by high dichlorvos
Fenamiphos	• Down by fenamiphos high • UP by fenamiphos high
Mercury	• Down by mercury high 8H • UP by mercury high 8H
Ivermectin (IVM)	• Down by 1ng/ml ivermectin • UP by 1ng/ml ivermectin
Mefloquine	• Down by mefloquine high • UP by mefloquine high
Heat Shock F3	• Down HS F3 vs. CTL F3 • UP HS F3 vs. CTL F3
Acrylamide	• Down by high acrylamide 8H • UP by high acrylamide 8H
Multi-walled carbon nanotubes (MWCNT)	• Down by MWCNT 24H vs. 0H • UP by MWCNT 24H vs. 0H
Tert-butyl hydrogen peroxide	• down by tert-butyl hydrogen peroxide (Oliveira) • UP by tert-butyl hydrogen peroxide (Oliveira)
Bacillus thuringiensis strain BT247	• down by B. thuringiensis at 6h (BT247, 1:2) (Yang) • down by B. thuringiensis at 6h (BT247, 1:2) (Yang)
Bacillus licheniformis strain 141	• down by B. licheniformis 141 • UP by B. licheniformis 141
Drechmeria coniospora	• Down D. coniospora 12h, RNASeq • UP D. coniospora 12h, RNASeq
Pseudomonas aeruginosa strain PA14	• down by PA14, 24h • UP by PA14, 24h
Cry5B toxin	• down by Bt toxin,Cry5B • UP by Bt toxin,Cry5B
Candida albicans	• down by C. albicans (Pukkila-Worley) • UP by C. albicans (Pukkila-Worley)
Lactobacillus rhamnosus strain CNCM I-3690	• Down fed with L. rhamnosus CNCM I-3690 vs. OP50,3 days • UP fed with L. rhamnosus CNCM I-3690 vs. OP50,3 days
Staphylococcus aureus	• down by S. aureus (Bond) • UP by S. aureus (Bond)
Serratia marcescens	• Down infected by S. Marcescens • UP on S. marcescens
Xenorhabdus nematophila	• Down infected by X. Nematophila

	• UP on X. nematophila

1179

References

1. Band G, Rockett KA, Spencer CCA, Kwiatkowski DP. A novel locus of resistance to severe malaria in a region of ancient balancing selection. Nature. 2015;526: 253–257. doi:10.1038/nature15390

2. Samson M, Libert F, Doranz BJ, Rucker J, Liesnard C, Farber CM, et al. Resistance to HIV-1 infection in caucasian individuals bearing mutant alleles of the CCR-5 chemokine receptor gene. Nature. 1996;382: 722–725. doi:10.1038/382722a0

3. Sabeti PC, Schaffner SF, Fry B, Lohmueller J, Varilly P, Shamovsky O, et al. Positive natural selection in the human lineage. Science. 2006;312: 1614–1620. doi:10.1126/science.1124309

4. Lazzaro BP, Sceurman BK, Clark AG. Genetic Basis of Natural Variation in D. melanogaster Antibacterial Immunity. Science. 2004;303: 1873–1876. doi:10.1126/science.1092447

5. Duxbury EM, Day JP, Maria Vespasiani D, Thüringer Y, Tolosana I, Smith SC, et al. Host-pathogen coevolution increases genetic variation in susceptibility to infection. eLife. 2019;8: 1–21. doi:10.7554/eLife.46440

6. Lazzaro BP, Sackton TB, Clark AG. Genetic variation in Drosophila melanogaster resistance to infection: A comparison across bacteria. Genetics. 2006;174: 1539–1554. doi:10.1534/genetics.105.054593

7. Schulenburg H, Ewbank JJ. Diversity and specificity in the interaction between Caenorhabditis elegans and the pathogen Serratia marcescens. BMC Evolutionary Biology. 2004;4: 49. doi:10.1186/1471-2148-4-49

8. Marquet S. Overview of human genetic susceptibility to malaria: From parasitemia control to severe disease. Infect Genet Evol. 2018;66: 399–409. doi:10.1016/j.meegid.2017.06.001

9. Raberg L, Sim D, Read AF. Disentangling genetic variation for resistance and tolerance to infectious diseases in animals. Science. 2007;318: 812–814. doi:10.1126/science.1148526

10. Medzhitov R, Schneider DS, Soares MP. Disease tolerance as a defense strategy. Science. 2012;335: 936–941. doi:10.1126/science.1214935

11. Wadi L, Reinke AW. Evolution of microsporidia: An extremely successful group of eukaryotic intracellular parasites. PLoS Pathog. 2020;16: e1008276. doi:10.1371/journal.ppat.1008276

12. Stentiford GD, Feist SW, Stone DM, Bateman KS, Dunn AM. Microsporidia: diverse,
dynamic, and emergent pathogens in aquatic systems. Trends Parasitol. 2013;29: 567–578.
doi:10.1016/j.pt.2013.08.005

13. Murareanu BM, Sukhdeo R, Qu R, Jiang J, Reinke AW, Weiss LM. Generation of a
Microsporidia Species Attribute Database and Analysis of the Extensive Ecological and
Phenotypic Diversity of Microsporidia. mBio. 2021. pp. e01490-21.

14. Willis AR, Reinke AW. Factors That Determine Microsporidia Infection and Host
Specificity. Exp Suppl. 2022;114: 91–114. doi:10.1007/978-3-030-93306-7_4

15. Troemel ER, Felix M-A, Whiteman NK, Barriere A, Ausubel FM. Microsporidia are
natural intracellular parasites of the nematode Caenorhabditis elegans. PLoS Biol. 2008;6:
2736–2752. doi:10.1371/journal.pbio.0060309

16. Balla KM, Luallen RJ, Bakowski MA, Troemel ER. Cell-to-cell spread of microsporidia
causes Caenorhabditis elegans organs to form syncytia. Nature Microbiology. 2016;1:
16144.

17. Higes M, Martín-Hernández R, Botías C, Bailón EG, González-Porto AV, Barrios L, et al.
How natural infection by Nosema ceranae causes honeybee colony collapse. Environ
Microbiol. 2008;10: 2659–2669. doi:10.1111/j.1462-2920.2008.01687.x

18. Richaud A, Zhang G, Lee D, Lee J, Félix M-A. The Local Coexistence Pattern of Selfing
Genotypes in Caenorhabditis elegans Natural Metapopulations. Genetics. 2018;208: 807–
821. doi:10.1534/genetics.117.300564

19. Balla KM, Andersen EC, Kruglyak L, Troemel ER. A Wild C. Elegans Strain Has
Enhanced Epithelial Immunity to a Natural Microsporidian Parasite. PLoS Pathogens.
2015;11: e1004583. doi:10.1371/journal.ppat.1004583

20. Tamim El Jarkass H, Mok C, Schertzberg MR, Fraser AG, Troemel ER, Reinke AW. An
intestinally secreted host factor promotes microsporidia invasion of C. elegans. Lourido S,
Sengupta P, editors. eLife. 2022;11: e72458. doi:10.7554/eLife.72458

21. Capaul M, Ebert D. Parasite-Mediated Selection in Experimental Daphnia Magna
Populations. Evolution. 2003;57: 249–260. doi:10.1111/j.0014-3820.2003.tb00260.x

22. Haag CR, Ebert D. Parasite–mediated selection in experimental metapopulations of
Daphnia magna. Proceedings of the Royal Society of London Series B: Biological Sciences.
2004;271: 2149–2155. doi:10.1098/rspb.2004.2841

23. Huang Q, Lattorff HMG, Kryger P, Conte YL, Moritz RFA. A selective sweep in a
microsporidian parasite Nosema-tolerant honeybee population, Apis mellifera. Animal
Genetics. 2014;45: 267–273. doi:10.1111/age.12114

24.   Keller D, Kirk D, Luijckx P. Four QTL underlie resistance to a microsporidian parasite that
1246   may drive genome evolution in its Daphnia host. bioRxiv. 2019; 847194.
1247   doi:10.1101/847194

1248   25.   Krebs M, Routtu J, Ebert D. QTL mapping of a natural genetic polymorphism for long-term
1249   parasite persistence in Daphnia populations. Parasitology. 2017;144: 1686–1694.
1250   doi:10.1017/S0031182017001032

1251   26.   Routtu J, Ebert D. Genetic architecture of resistance in Daphnia hosts against two species of
1252   host-specific parasites. Heredity (Edinb). 2015;114: 241–248. doi:10.1038/hdy.2014.97

1253   27.   King KC, Brockhurst MA, Vasieva O, Paterson S, Betts A, Ford SA, et al. Rapid evolution
1254   of microbe-mediated protection against pathogens in a worm host. The ISME Journal.
1255   2016;10: 1915–1924. doi:10.1038/ismej.2015.259

1256   28.   Morran LT, Schmidt OG, Gelarden IA, Parrish RC, Lively CM. Running with the Red
1257   Queen: Host-Parasite Coevolution Selects for Biparental Sex. Science. 2011;333: 216–218.
1258   doi:10.1126/science.1206360

1259   29.   Papkou A, Guzella T, Yang W, Koepper S, Pees B, Schalkowski R, et al. The genomic
1260   basis of Red Queen dynamics during rapid reciprocal host–pathogen coevolution. PNAS.
1261   2019;116: 923–928. doi:10.1073/pnas.1810402116

1262   30.   Masri L, Branca A, Sheppard AE, Papkou A, Laehnemann D, Guenther PS, et al. Host–
1263   Pathogen Coevolution: The Selective Advantage of Bacillus thuringiensis Virulence and Its
1264   Cry Toxin Genes. PLOS Biology. 2015;13: e1002169. doi:10.1371/journal.pbio.1002169

1265   31.   Schulenburg H, Félix M-A. The Natural Biotic Environment of . Genetics. 2017;206: 55–
1266   86.

1267   32.   Zhang G, Sachse M, Prevost M-C, Luallen RJ, Troemel ER, Felix M-A. A Large Collection
1268   of Novel Nematode-Infecting Microsporidia and Their Diverse Interactions with
1269   Caenorhabditis elegans and Other Related Nematodes. PLoS Pathog. 2016;12: e1006093.
1270   doi:10.1371/journal.ppat.1006093

1271   33.   Cook DE, Zdraljevic S, Roberts JP, Andersen EC. CeNDR, the Caenorhabditis elegans
1272   natural diversity resource. Nucleic Acids Research. 2017. doi:10.1093/nar/gkw893

1273   34.   Ashe A, Bélicard T, Le Pen J, Sarkies P, Frézal L, Lehrbach NJ, et al. A deletion
1274   polymorphism in the Caenorhabditis elegans RIG-I homolog disables viral RNA dicing and
1275   antiviral immunity. eLife. 2013;2013: 1–21. doi:10.7554/eLife.00994.001

1276   35.   Sterken MG, van Sluijs L, Wang YA, Ritmahan W, Gultom ML, Riksen JAG, et al.
1277   Punctuated Loci on Chromosome IV Determine Natural Variation in Orsay Virus
1278   Susceptibility of Caenorhabditis elegans Strains Bristol N2 and Hawaiian CB4856. Journal
1279   of Virology. 2021;95: e02430-20. doi:10.1128/JVI.02430-20

1280   36.   Reddy KC, Andersen EC, Kruglyak L, Kim DH. A Polymorphism in npr-1 Is a Behavioral
1281   Determinant of Pathogen Susceptibility in C. elegans. Science. 2009;323: 382–384.
1282   doi:10.1126/science.1166527

1283   37.   Huang Y, Kammenga JE. Genetic Variation in Caenorhabditis elegans Responses to
1284   Pathogenic Microbiota. Microorganisms. 2020;8: 618.
1285   doi:10.3390/microorganisms8040618

1286   38.   Webster AK, Hung A, Moore BT, Guzman R, Jordan JM, Kaplan REW, et al. Population
1287   Selection and Sequencing of Caenorhabditis elegans Wild Isolates Identifies a Region on
1288   Chromosome III Affecting Starvation Resistance. G3: Genes, Genomes, Genetics. 2019;9:
1289   3477–3488. doi:10.1534/g3.119.400617

1290   39.   Mok C, Belmarez G, Edgley ML, Moerman DG, Waterston RH. PhenoMIP: High-
1291   Throughput Phenotyping of Diverse Caenorhabditis elegans Populations via Molecular
1292   Inversion Probes. G3: Genes, Genomes, Genetics. 2020;10: 3977–3990.
1293   doi:10.1534/g3.120.401656

1294   40.   Webster AK, Chitrakar R, Powell M, Chen J, Fisher K, Tanny RE, et al. Using population
1295   selection and sequencing to characterize natural variation of starvation resistance in
1296   Caenorhabditis elegans. Hobert O, James DE, McGrath P, editors. eLife. 2022;11: e80204.
1297   doi:10.7554/eLife.80204

1298   41.   Luallen RJ, Reinke AW, Tong L, Botts MR, Felix M-A, Troemel ER. Discovery of a
1299   Natural Microsporidian Pathogen with a Broad Tissue Tropism in Caenorhabditis elegans.
1300   PLoS Pathog. 2016;12: e1005724. doi:10.1371/journal.ppat.1005724

1301   42.   Murareanu BM, Antao NV, Zhao W, Dubuffet A, El Alaoui H, Knox J, et al. High-
1302   throughput small molecule screen identifies inhibitors of microsporidia invasion and
1303   proliferation in C. elegans. Nat Commun. 2022;13: 5653. doi:10.1038/s41467-022-33400-y

1304   43.   Reinke AW, Balla KM, Bennett EJ, Troemel ER. Identification of microsporidia host-
1305   exposed proteins reveals a repertoire of rapidly evolving proteins. Nat Commun. 2017;8:
1306   14023. doi:10.1038/ncomms14023

1307   44.   Wadi L, El Jarkass HT, Tran TD, Islah N, Luallen RJ, Reinke AW. Genomic and
1308   phenotypic evolution of nematode-infecting microsporidia. bioRxiv. 2022;
1309   2022.08.28.505597. doi:10.1101/2022.08.28.505597

1310   45.   Jarkass HTE, Reinke AW. The ins and outs of host-microsporidia interactions during
1311   invasion, proliferation and exit. Cellular Microbiology. 2020;22: e13247.
1312   doi:10.1111/cmi.13247

1313   46.   Tecle E, Troemel ER. Insights from C. elegans into Microsporidia Biology and Host-
1314   Pathogen Relationships. Exp Suppl. 2022;114: 115–136. doi:10.1007/978-3-030-93306-7_5

47. Reddy KC, Dror T, Sowa JN, Panek J, Chen K, Lim ES, et al. An Intracellular Pathogen
Response Pathway Promotes Proteostasis in C. elegans. Curr Biol. 2017;27: 3544-3553.e5.
doi:10.1016/j.cub.2017.10.009

48. Reddy KC, Dror T, Underwood RS, Osman GA, Elder CR, Desjardins CA, et al.
Antagonistic paralogs control a switch between growth and pathogen resistance in C.
elegans. PLoS Pathog. 2019;15: e1007528. doi:10.1371/journal.ppat.1007528

49. Willis AR, Zhao W, Sukhdeo R, Wadi L, Jarkass HTE, Claycomb JM, et al. A parental
transcriptional response to microsporidia infection induces inherited immunity in offspring.
Science Advances. 2021;7: eabf3114. doi:10.1126/sciadv.abf3114

50. Tecle E, Chhan CB, Franklin L, Underwood RS, Hanna-Rose W, Troemel ER. The purine
nucleoside phosphorylase pnp-1 regulates epithelial cell resistance to infection in C.
elegans. PLOS Pathogens. 2021;17: e1009350. doi:10.1371/journal.ppat.1009350

51. Balla KM, Lažetić V, Troemel ER. Natural variation in the roles of C. elegans autophagy
components during microsporidia infection. PLOS ONE. 2019;14: e0216011.
doi:10.1371/journal.pone.0216011

52. Mok CA, Au V, Thompson OA, Edgley ML, Gevirtzman L, Yochem J, et al. MIP-MAP:
High-Throughput Mapping of Caenorhabditis elegans Temperature-Sensitive Mutants via
Molecular Inversion Probes. Genetics. 2017;207: 447–463.
doi:10.1534/genetics.117.300179

53. Seidel HS, Ailion M, Li J, Oudenaarden A van, Rockman MV, Kruglyak L. A Novel
Sperm-Delivered Toxin Causes Late-Stage Embryo Lethality and Transmission Ratio
Distortion in C. elegans. PLOS Biology. 2011;9: e1001115.
doi:10.1371/journal.pbio.1001115

54. Bakowski MA, Desjardins CA, Smelkinson MG, Dunbar TL, Lopez-Moyado IF, Rifkin
SA, et al. Ubiquitin-mediated response to microsporidia and virus infection in C. elegans.
PLoS Pathog. 2014;10: e1004200. doi:10.1371/journal.ppat.1004200

55. Panek J, Gang SS, Reddy KC, Luallen RJ, Fulzele A, Bennett EJ, et al. A cullin-RING
ubiquitin ligase promotes thermotolerance as part of the intracellular pathogen response in
Caenorhabditis elegans. Proceedings of the National Academy of Sciences. 2020;117:
7950–7960. doi:10.1073/pnas.1918417117

56. Yang W, Dierking K, Schulenburg H. WormExp: a web-based application for a
Caenorhabditis elegans-specific gene expression enrichment analysis. Bioinformatics.
2016;32: 943–945. doi:10.1093/bioinformatics/btv667

57. Lewis JA, Szilagyi M, Gehman E, Dennis WE, Jackson DA. Distinct patterns of gene and
protein expression elicited by organophosphorus pesticides in Caenorhabditis elegans.
BMC Genomics. 2009;10: 202. doi:10.1186/1471-2164-10-202

58. Helmcke KJ, Syversen T, Miller DM, Aschner M. Characterization of the effects of
methylmercury on Caenorhabditis elegans. Toxicology and Applied Pharmacology.
2009;240: 265–272. doi:10.1016/j.taap.2009.03.013

59. Eom H-J, Kim H, Kim B-M, Chon T-S, Choi J. Integrative Assessment of Benzene
Exposure to Caenorhabditis elegans Using Computational Behavior and Toxicogenomic
Analyses. Environ Sci Technol. 2014;48: 8143–8151. doi:10.1021/es500608e

60. Menzel R, Menzel S, Swain S, Pietsch K, Tiedt S, Witczak J, et al. The Nematode
Caenorhabditis elegans, Stress and Aging: Identifying the Complex Interplay of Genetic
Pathways Following the Treatment with Humic Substances. Frontiers in Genetics. 2012;3.
Available: https://www.frontiersin.org/article/10.3389/fgene.2012.00050

61. Trzebny A, Slodkowicz-Kowalska A, Becnel JJ, Sanscrainte N, Dabert M. A new method
of metabarcoding Microsporidia and their hosts reveals high levels of microsporidian
infections in mosquitoes (Culicidae). Molecular Ecology Resources. 2020;20: 1486–1504.
doi:https://doi.org/10.1111/1755-0998.13205

62. Lange B, Kaufmann AP, Ebert D. Genetic, ecological and geographic covariables
explaining host range and specificity of a microsporidian parasite. Journal of Animal
Ecology. 2015;84: 1711–1719. doi:10.1111/1365-2656.12421

63. Bonneaud C, Tardy L, Giraudeau M, Hill GE, McGraw KJ, Wilson AJ. Evolution of both
host resistance and tolerance to an emerging bacterial pathogen. Evolution Letters. 2019;3:
544–554. doi:https://doi.org/10.1002/evl3.133

64. Ayres JS, Schneider DS. Tolerance of Infections. Annual Review of Immunology. 2012;30:
271–294. doi:10.1146/annurev-immunol-020711-075030

65. Edgerton EB, McCrea AR, Berry CT, Kwok JY, Thompson LK, Watson B, et al.
Activation of mosquito immunity blocks the development of transmission-stage filarial
nematodes. PNAS. 2020;117: 3711–3717. doi:10.1073/pnas.1909369117

66. Grover M, Fasseas MK, Essmann C, Liu K, Braendle C, Félix M-A, et al. Infection of C.
elegans by Haptoglossa Species Reveals Shared Features in the Host Response to
Oomycete Detection. Frontiers in Cellular and Infection Microbiology. 2021;11: 973.
doi:10.3389/fcimb.2021.733094

67. Osman GA, Fasseas MK, Koneru SL, Essmann CL, Kyrou K, Srinivasan MA, et al. Natural
Infection of C. elegans by an Oomycete Reveals a New Pathogen-Specific Immune
Response. Current Biology. 2018;28: 640-648.e5. doi:10.1016/j.cub.2018.01.029

68. Lansdon P, Carlson M, Ackley BD. Wild-type Caenorhabditis elegans isolates exhibit
distinct gene expression profiles in response to microbial infection. BMC Genomics.
2022;23: 229. doi:10.1186/s12864-022-08455-2

69. Wan YC, Troemel ER, Reinke AW. Conservation of Nematocida microsporidia gene
expression and host response in Caenorhabditis nematodes. PLOS ONE. 2022;17:
e0279103. doi:10.1371/journal.pone.0279103

70. Dirksen P, Assié A, Zimmermann J, Zhang F, Tietje A-M, Marsh SA, et al. CeMbio - The
Caenorhabditis elegans Microbiome Resource. G3: Genes, Genomes, Genetics. 2020;10:
3025–3039. doi:10.1534/g3.120.401309

71. Powell JR, Ausubel FM. Models of Caenorhabditis elegans Infection by Bacterial and
Fungal Pathogens. In: Ewbank J, Vivier E, editors. Innate Immunity. Totowa, NJ: Humana
Press; 2008. pp. 403–427. doi:10.1007/978-1-59745-570-1_24

72. Norris AD, Kim H-M, Colaiácovo MP, Calarco JA. Efficient Genome Editing in
Caenorhabditis elegans with a Toolkit of Dual-Marker Selection Cassettes. Genetics.
2015;201: 449–458. doi:10.1534/genetics.115.180679

73. Stiernagle T. Maintenance of C. elegans. WormBook. 2006; 1–11.
doi:10.1895/wormbook.1.101.1

74. Luallen RJ, Bakowski MA, Troemel ER. Characterization of Microsporidia-Induced
Developmental Arrest and a Transmembrane Leucine-Rich Repeat Protein in
Caenorhabditis elegans. PLOS ONE. 2015;10: e0124065.
doi:10.1371/journal.pone.0124065

75. Willis AR, Jarkass HTE, Reinke AW. Studying Inherited Immunity in a Caenorhabditis
elegans Model of Microsporidia Infection. JoVE (Journal of Visualized Experiments).
2022; e63636. doi:10.3791/63636

76. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nature Methods. 2012;9: 676–682.
doi:10.1038/nmeth.2019

77. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012;9: 357–359. doi:10.1038/nmeth.1923

78. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25: 2078–2079.
doi:10.1093/bioinformatics/btp352

79. Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing. 2012
[cited 4 Jun 2022]. doi:10.48550/ARXIV.1207.3907

80. Cook DE, Andersen EC. VCF-kit: Assorted utilities for the variant call format.
Bioinformatics. 2017; btx011. doi:10.1093/bioinformatics/btx011

81. Wang L, Wang S, Li W. RSeQC: quality control of RNA-seq experiments. Bioinformatics.
2012;28: 2184–2185. doi:10.1093/bioinformatics/bts356

82.   Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for
1422   multiple tools and samples in a single report. Bioinformatics. 2016;32: 3047–3048.
1423   doi:10.1093/bioinformatics/btw354

1424   83.   Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq
1425   quantification. Nat Biotechnol. 2016;34: 525–527. doi:10.1038/nbt.3519

1426   84.   Marini F, Binder H. pcaExplorer: an R/Bioconductor package for interacting with RNA-seq
1427   principal components. BMC Bioinformatics. 2019;20: 331. doi:10.1186/s12859-019-2879-1

1428   85.   Chen K, Franz CJ, Jiang H, Jiang Y, Wang D. An evolutionarily conserved transcriptional
1429   response to viral infection in Caenorhabditis nematodes. BMC Genomics. 2017;18: 303.
1430   doi:10.1186/s12864-017-3689-3

1431   86.   Eddy SR. Accelerated Profile HMM Searches. PLOS Computational Biology. 2011;7:
1432   e1002195. doi:10.1371/journal.pcbi.1002195

1433   87.   Kuleshov MV, Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, et al. Enrichr: a
1434   comprehensive gene set enrichment analysis web server 2016 update. Nucleic Acids Res.
1435   2016;44: W90–W97. doi:10.1093/nar/gkw377

1436