The effects of conscious movement processing on the neuromuscular control of posture

Li-Juan Jie[1,2], Elmar Kal[3,4], Toby Ellmers[5], Joëlle Rosier[1], Kenneth Meijer[1], Tjeerd Boonstra[6]

[1] Department of Nutrition and Movement Sciences, NUTRIM School of Nutrition and Translational Research in
Metabolism, Maastricht University, the Netherlands
[2] Research Centre for Nutrition, Lifestyle and Exercise, Zuyd University of Applied Sciences, the Netherlands
[3] College of Health, Medicine and Life Sciences, Brunel University London, UK
[4] Centre for Cognitive Neuroscience, Brunel University London, UK
[5] Centre for Vestibular Neurology, Imperial College London, UK
[6] Department of Neuropsychology and Psychopharmacology, Faculty of Psychology and Neuroscience,
Maastricht University, the Netherlands

Correspondence concerning this article should be addressed to:

Dr. Li-Juan Jie

Department of Nutrition and Movement Sciences

Maastricht University

Universiteitssingel 50, 6200 MD Maastricht the Netherlands

Email address: li-juan.jie@maastrichtuniversity.nl

Word count: 4849

Abstract

Maintaining balance typically occurs sub-consciously. Occasionally, however, individuals will direct conscious attention towards balance, e.g., in response to a postural threat. Such conscious movement processing (CMP) increases the reliance on attentional resources and may disrupt control of balance, but the underlying changes in neuromuscular control remain poorly understood. We investigated the effects of CMP (manipulated using verbal instructions) on neural control of posture in twenty-five adults (11 females, mean age = 23.9, range = 18–33). Participants performed 90-second, narrow-stance balance trials in high- and low-CMP conditions, and in stable (solid surface) and unstable (foam) task conditions. Postural sway amplitude, frequency and complexity were used to assess postural control. Surface EMG was recorded bilaterally from lower leg muscles (Soleus, Tibialis Anterior, Gastrocnemius Medialis, Peroneus Longus) and intermuscular coherence (IMC) was assessed for 12 muscle pairs across four frequency bands. We found significantly increased sway amplitude, and decreased sway frequency and complexity in the high- compared to the low-CMP condition. All sway variables (frequency, amplitude and complexity) increased in the unstable compared to the stable condition. Results suggested reduced beta band IMC between several muscles pairs during high- compared to low-CMP, but these findings did not remain significant after controlling for multiple comparisons. Finally, IMC significantly increased in the unstable conditions for most muscle combinations and frequency bands. In all, results tentatively suggest that CMP-induced changes in sway outcomes may be underpinned by reduced beta-band IMC, but these findings need to be replicated before they can be interpreted more conclusively.

Keywords: Attention; Postural Control; Balance; Intermuscular Coherence; Movement Reinvestment; Internal Focus;

1. Introduction

Humans typically control their posture in a relatively automatic manner, without much awareness of how they maintain balance. However, in some situations, e.g., when anxious about falling (Adkin & Carpenter, 2018; Young & Williams, 2015) or when instructed to do so when (re-)learning a skill, or within some populations such as stroke or Parkinson's' disease (Masters et al, 2007; Orrell, Masters, & Eves, 2009), individuals will direct attention towards controlling their balance. This increased reliance on attentional resources may alter the neuromuscular control of balance. For instance, researchers have suggested that conscious control may underpin the frequently observed 'postural stiffening' strategy individuals adopt when their balance is threatened, e.g., when standing at the edge of a raised platform (Adkin & Carpenter, 2018; Young & Williams, 2015).

Postural stiffening is characterised by a concurrent reduction in sway amplitude and increase in sway frequency, and is thought to result from enhanced co-contraction of the ankle muscles (Adkin & Carpenter, 2018; Stins et al., 2011). These changes may be driven in part by fear-related changes in attention (Adkin & Carpenter, 2018), as we typically observe concomitant increases in conscious (cortical) control of balance when individuals are fearful of falling (see Adkin & Carpenter, 2018). However, only few studies have directly investigated the effects of conscious balance processing on ankle stiffening in the absence of other fear-related physiological responses (Richer et al., 2020; Richer et al, 2017), but their results suggest that conscious processing may have little direct effect on neuromuscular outcomes of postural stiffening. Recent research has also presented findings that suggest that conscious balance processing actually *drives* behaviours that are opposite to postural stiffening responses (e.g., reduced sway frequency and increased sway amplitude (Ellmers et al., 2021; Kal et al., 2022). However, the work by Ellmers et al. (2021) and Kal et al. (2022) restricted their analysis to postural sway outcomes and did not study neuromuscular control outcomes.

A more detailed understanding of the effects of conscious processing on neuromuscular control of posture may be obtained using intermuscular coherence (IMC) analysis. IMC, a measure of linear correlation between two EMG signals in the frequency domain, can be used to estimate common spinal input to different muscles (Boonstra, 2013; Boonstra et al., 2016; Farina & Negro, 2015). IMC is observed in different frequency bands and common input at these frequencies may originate from different cortical or sub-cortical processes (Grosse et al., 2002; Kerkman et al., 2018). During quiet standing, lower frequency bands (0-5 Hz and 6-15 Hz) are thought to reflect afferent and supraspinal input during quiet standing (Boonstra et al., 2008; Nandi et al., 2019; Obata et al., 2014), whereas IMC in the beta band (15-30Hz) is considered to reflect cortical input (Farmer et al, 1993; Grosse et al., 2002). Conscious control is expected to be reliant on cortical processes, for example some studies have shown that increased conscious movement processing is associated with increased activity of the verbal-analytical and motor-

planning regions of the cortex (Ellmers et al., 2016; Zhu et al., 2011). Therefore, corticospinal drive is expected to change with increasing conscious movement processing, as manifested by increased IMC at higher frequencies.

There are currently no studies that have directly aimed to isolate the effects of (experimentally-induced) conscious movement processing on IMC during balance. Preliminary insight can be gathered from several studies which have indirectly modulated conscious processing during a variety of postural control tasks. For example, Boonstra et al. (2015) evaluated IMC while manipulating attention during different task conditions likely to alter conscious processing of balance, such as counting backwards (steps of sevens), holding a cup, and raising platform height. IMC was dependent on the type of standing task: when counting (when there is *less* opportunity for conscious processing of balance), coherence increased between agonist and antagonist lower leg muscles at 10Hz. Yet during the height condition (when we typically observe an *increase* in conscious processing) a pronounced increase in bilateral Tibialis Anterior IMC was found at 16Hz (Boonstra et al., 2015). During this same height condition, coherence values were generally higher across many muscle combinations and frequencies when compared to the other conditions. Zaback et al. (2022) also reported increased IMC in this frequency range (i.e., in the 5-19Hz frequency band) for bilateral Soleus muscles when participants were standing at a raised platform. Boonstra et al. (2015) and Zaback et al. (2022) both suggest such postural threat to increase IMC in the beta band (20-35 Hz), plantar flexor-dorsal flexor pairs (Boonstra et al., 2015) and bilateral Soleus muscles (Zaback et al., 2022) respectively. This suggests an increased corticospinal drive due to anxiety-induced conscious processing (Zaback et al., 2022). As such, we would expect that promoting conscious movement processing in isolation (i.e., using specific verbal instructions) may yield similar effects (i.e., increases in beta band IMC). However, the direct effects of conscious balance processing on IMC have not yet been studied.

The current study investigates how conditions that either promote or minimise conscious movement processing influence the neuromuscular control of posture in young healthy adults. In this study, postural control was evaluated using metrics of i) *performance,* i.e., sway amplitude (root mean square) ii) *movement automaticity,* i.e., sway frequency (mean power frequency) and sway complexity (sample entropy) and iii) *muscle coordination,* i.e., intermuscular coherence. We assessed IMC in the following lower limb muscles relevant to postural control: Soleus, Tibialis Anterior, Gastrocnemius Medialis and Peroneus Longus. We expect that during high-conscious movement processing i) performance would worsen, as evidenced by increased sway amplitude (e.g., Kal et al., 2022), ii) movement automaticity would be reduced, as evidenced by reduced sway frequency and complexity (e.g., Kal, et al., 2022), iii) and that this would be accompanied by increased beta-band IMC in both agonist and antagonist muscle pairs (Boonstra et al., 2015; Zaback et al., 2022). Finally, for a more comprehensive assessment, participants

performed the standing balance task in stable (solid surface) and unstable (foam) conditions, to assess if the above outcomes would be influenced by task difficulty. By exploring the effects of CMP on IMC the current study may ultimately help improve our understanding of how elevated CMP – as often observed in anxious individuals, and clinical populations - contributes to (mal)adaptive changes in postural control.

2. Experimental Procedures

2.1. Participants

Healthy young adults were included if they were aged between 18 and 40 years, had a BMI between 18 and 27 kg/m2 (to optimise reliability of EMG data), and were free from any (self-reported) neurological or musculoskeletal conditions that could influence balance performance. We aimed for a pragmatic sample size of 20 to 25 young healthy adults. The heuristic sample size was chosen based on previous studies with similar research set-ups, tasks and populations (Boonstra et al., 2015; Zaback et al., 2022). IMC analyses in the present study are largely exploratory and are performed to better understand the muscle pairings, frequency bands and experimental tasks to focus on in future work. Yet, results of the study allow us to draw some preliminary interpretations on how CMP (in absence of fall-related anxiety) affects IMC. Recruitment took place via the local network of the researchers (online adverts and by word of mouth) and from the university population for course credits. Each participant received an information sheet and provided written consent prior to the measurement session. The Ethics Review Committee Psychology and Neuroscience from Maastricht University approved the study protocol (ERCPN-230_137_11_2020).

2.2. Postural task

Participants performed a 90-second narrow, bipedal stance with one foot on each force plate (feet 10 cm apart; exact positions were marked on the floor to ensure consistency of positioning across trials). Participants were instructed to stand straight and keep their hands by their side looking straight ahead at a fixation cross displayed on a screen (at eye level) in front of them (see Figure 1). A full factorial (2×2) design was used in which participants performed this task in 'low' and 'high' Conscious Movement Processing (CMP) conditions and in 'stable' and 'unstable' task conditions. The high-CMP condition aimed to induce conscious movement processing through previously validated instructions (Kal et al., 2022), whereas the low-CMP condition aimed to distract people from focusing on their movement by using a cognitively less demanding distraction task (see Table 1). To investigate whether the effects of CMP on balance control were influenced by task difficulty (as suggested by Young & Williams, 2015), the task was performed with different levels of surface stability. In the 'stable' condition, participants performed the balance task whilst standing on a flat, solid surface, while in the 'unstable' condition, the task was performed while standing on foam (Balance pad Dittmann; 50 x 40 x 6 cm). Therefore, participants

172   performed the task four times in total (a single 90-second trial for each of the four specific conditions) and

173   the order of conditions was counterbalanced across participants (balanced Latin-square).

175   **Figure 1**. Schematic illustration of the experimental set up. Height of the screen was adjustable and set to
176   eye level of the participant. Distance between the standing position and the screen was approximately 2
177   meters. The grey squares on the floor indicate two independent force plates.
178
179   **Table 1.** Standardised instructions

Condition	Instructions

High-CMP[1]	Please stand straight with your feet on **the marked positions** with your **hands by your side** and **look straight ahead**. At a certain point during this trial, you will hear a 'beep'. When **you hear a 'beep'** we will ask you
	    1) Where your weight was currently distributed beneath your feet, and
	    2) How your weight distribution changed in the 10 seconds prior to the 'beep'.
	You can **look at the cross** X that will be displayed on this screen.

Low-CMP	Please stand straight with your feet on **the marked positions** with your **hands by your side** and look **straight ahead**. A number of colours will be presented. At the end, you have to tell how many times the colour '**red**'[2] has passed. Count the relevant colour **in your head**. So **not out loud**!

[1]*The high-CMP instructions have been validated in a study by Kal et al. (2022) This manipulation has been found to reliably induce increases*
in self-reported conscious movement processing. Although participants were told that the beep could occur at any point during the trial
(after the initial 10 seconds had passed), in fact the beep always occurred at the end of the trial, once data had stopped recording.
[2] *In total 5 different colours (the whole screen displayed either the colour: red, purple, blue, green, or yellow) within a 90 second time frame*
were presented at random to the participant. Each colour was presented to them for 2 seconds. Between each colour a fixation cross was
presented on the screen (e.g., green, fixation cross, red, fixation cross, yellow, fixation cross etc.) for the duration of 1 second. Participants
were asked to either count the colour "RED" or "BLUE.
187

2.3. Experimental procedures

Measurements took place in the Human Performance Lab at Maastricht University. To describe the

characteristics of the population we first collected demographic information related to age, gender, leg

dominance (i.e., front leg in tandem stance), and BMI. In addition, propensity for reinvestment was

measuring using the Movement Specific Reinvestment Scale (MSRS) (Masters et al., 2005). Attentional

capacity was measured using the D2 (Brickenkamp and Zillmer, 1998; this test measures sustained

attention, specifically). If desired, there was the possibility to have short (5 min) resting breaks in between

the tests.

Prior to performance of the postural task, sixteen EMG sensors were placed bilaterally on the Tibialis

Anterior (TA), Soleus (SOL), Gastrocnemius Medialis (GM), Peroneus Longus (PL), Rectus Femoris (RF),

Biceps Femoris (BF), Gluteus Maximus (GLM), and the Erector Spinae (ES) (Fig. 2). Electrode placement

and skin preparation (i.e., the electrode position was shaved and disinfected with alcohol) were

performed in accordance with the SENIAM recommendations (www.seniam.org). Furthermore, as the

current study was part of a larger project, participants completed the task whilst also wearing an EEG cap.

Please note, however, that we only used the force plate data and the EMG data for the SOL, TA, GM, and

PL for the current analyses as these are the primary ankle plantar and dorsiflexors during stance. After a

familiarisation trial, i.e., performance of a 90-second narrow bipedal stance without any CMP-related

instructions, the experiment started and instructions were given to the participants (Table 1).

2.4. Data acquisition

Force signals were recorded using two force platforms (600mmx400mm; AMTI OR6-Series, Watertown,

USA; 1000Hz). Muscle activity was recorded bilaterally with sixteen EMG sensors (PicoEMG, Cometa

Systems, Newburg, USA; 1000Hz). All signals (EMG, EEG, force plates) were concurrently recorded and

temporally aligned with triggers generated using Presentation software (Version 22.1, Neurobehavioral

Systems Inc, Albany, CA, USA; www.neurobs.com). A 23-inch wide-screen monitor (HP Pavilion 2309m,

Hewlett-Packard, Palo Alto, CA, USA) was used to present the instructions (Table 1).

2.5. Data analyses

2.5.1.   Postural stability

Analyses were performed using customised Matlab scripts (The Mathworks, Natick, MA, USA) and are

freely available through the OSF repository (https://osf.io/j3e6a/). Force data was low-pass filtered (5 Hz,

2^{nd} order Butterworth), after which COP coordinates were determined. The COP data was used to

estimate the amplitude (RMS; Root Mean Square), frequency (MPF; Mean Power Frequency), and

complexity (SEn; Sample Entropy) of postural adjustments for each condition in Medio Lateral (ML) and

Anterior Posterior (AP) directions. Higher MPF values have been shown to reflect a more automatic mode

of postural control (Ellmers, Kal, & Young, 2021; Richer et al., 2017). SEn provides a measure of

complexity of sway. Higher SEn values reflect more unpredictable, irregular sway, which has been

associated with more automatic postural control (Richer & Lajoie, 2020; Roerdink et al., 2011). Optimized

parameter settings (m=3, r=0.01) were used for the SEn calculations, as recommended by Lake et al.

(2002). For the SEn calculations, force plate data was down sampled to 100 Hz, which allows direct

comparisons of values with previous work (e.g., Kal et al., 2022).

2.5.2. Intermuscular coherence

EMG data from the lower leg was used to estimate intermuscular coherence. EMG signals were high-pass

filtered (cut-off at 20Hz) and rectified using the Hilbert transform, which yields the envelope of the

broadband EMG signal (Boonstra & Breakspear, 2012). Power spectral density and intermuscular

coherence of the EMG envelopes was estimated using the Welch method (window length 1 s, overlap

0.75 s). Intra-limb coherence was estimated between 6 muscle pairs (TA-SOL, TA-GM, TA-PL, SOL-GM,

SOL-PL, and GM-PL) on the left and right side. To determine the frequency bands for statistical analysis,

the 'collapsed localizer' approach was used (Luck & Gaspelin, 2017). The grand-average coherence spectra

were computed by averaging across all four experimental conditions to identify the peaks in IMC.

2.5.3. Manipulation checks

Immediately following the completion of each condition, participants completed a 2-item scale that

assessed the "conscious motor processing" subscale of a shortened state version of the MSRS (Ellmers &

Young, 2018). Total scores ranged from 2-12, with higher scores indicating greater state-CMP. The

subscale was used to determine whether the conscious movement processing manipulations had the

intended effect. Additionally, we also assessed whether the CMP manipulation led to any unintended

feelings of anxiety, as anxiety can have a significant influence on postural control (e.g., Adkin & Carpenter,

2018; Staab et al., 2013). Following the completion of each condition, participants rated the level of

anxiety they experienced during the previous condition on the Anxiety thermometer (range 0-10; higher

values indicate greater anxiety levels). Finally, the cognitive task used during the low-CMP condition was

designed to *distract* people, but we wanted to avoid it being so cognitively demanding that it might impair

performance. To check if this might be the case, all participants rated their perceived mental effort after

each condition using the Rating Scale of Mental Effort (RSME; 0-150, higher values indicate greater

cognitive load).

2.6. Statistical analysis

Demographic data were reported descriptively (mean, standard deviation and range). We first assessed

whether the experimental manipulations were successful: Two separate Wilcoxon-signed ranked tests

were performed to compare state-CMP scores between the high- and low-CMP conditions, one for the

stable and one for the unstable task condition. The experimental manipulations were deemed successful

if the state-CMP is higher in the High-CMP conditions irrespective of the surface condition. Similar tests

were used to compare self-reported anxiety and mental effort scores.

Next, separate repeated-measures ANOVA (analysis of variance) were used to compare how the force

plate outcomes and IMC values were affected by the CMP (high- vs. low-CMP) and surface conditions (stable

vs. unstable). For the COP data we compared RMS, MPF and SEn in the ML and AP direction (6 comparisons)

and for the IMC data we compared 12 muscle pairs in 4 frequency bands (48 comparisons). We used

Benjamini-Hochberg procedure to control the false discovery rate (Benjamini & Hochberg, 1995).

Standardised effect sizes reported were eta squared (η_p^2; ANOVA) or r (Z/√n; pairwise

comparisons). Statistical tests were performed using the IBM SPSS (version 28, IBM® SPSS®). An alpha level

of .05 was set for all tests.

3. Results

272   In total 25 young healthy adults were included. Please see Table 2 for their characteristics.

274   **Table 2**. Participant characteristics (n =25)

Group Characteristics	Mean ± SD	Range
Age (in years)	23.9 ± 0.9	18 - 33
Female/Male	11/14	NA
BMI (kg/m2)	22.5 ± 0.6	18.0 - 26.9
Leg dominance (left/right)	10/15	NA
Cognitive capacity		
Working memory (two-back test)		
% correct responses (max score 100%)	71% ± 5%	10% - 100%
Attention and Concentration (D2 test)		
CP score	201.5 ± 46.1	123 - 287
Conscious motor control preference		
Propensity for reinvestment (MSRS)		
MSRS-total (10-60)	28.7 ± 2.0	12 - 48
MSRS-CMP (5-30)	14.2 ± 5.7	5 - 24
MSRS-MSC (5-30)	14.3 ± 5.4	5 - 24

275   BMI: Body Mass Index; CP: number of correctly marked characters minus the number of
276   incorrectly marked characters (measure of attention span and concentration ability); MSRS: Movement Specific
277   Reinvestment Scale; CMP: Conscious Movement Processing; MSC: Movement Self-Consciousness; NA: not
278   applicable;
279
3.1. Manipulation check

State-CMP was significantly higher during the high- compared to low-CMP condition in both the stable

(high-CMP: *Mdn*=10.00, *IQR*=4; low-CMP: *Mdn*=5.00, *IQR*=5, *z* = -4,03, *p* < .001, *r* = -.570) and unstable

surface condition (high-CMP: *Mdn*=10.00, *IQR*=3, low-CMP: *Mdn*=6.00, *IQR*=5, *z* = -4.12, *p* < .001, *r* = -

.583). The average anxiety level across all conditions was 1.9 ± 0.1 (*range*: 1 to 4). No differences in

anxiety were observed between the high- and low-CMP condition for either the stable (high-CMP: *Mdn*=2,

IQR=1, low-CMP: *Mdn*=2, *IQR*=1, *z* = -.25, *p* = .80, *r* = -.035) or unstable surface condition (high-CMP:

Mdn=2, *IQR*=2, low-CMP: *Mdn*=2, *IQR*=0, *z* = -.97, *p* = .33, *r* = -.14). Finally, the perceived mental effort was

higher in the high- (*Mdn*=35, *IQR*=15) compared to the low-CMP (*Mdn*=25, *IQR*=15) during the unstable

condition, *z* = -2.8, *p* = .006, *r* = -.39. A similar trend was observed when comparing the high- (*Mdn*=30,

IQR=25) to the low-CMP (*Mdn*=25, *IQR*= 17.5) during stable surface condition, however this was not

statistically significant, *z* = 1.4, *p* = .15, *r* = -.39.

3.2. Postural stability

The balance outcomes are visualised in Figure 2, showing that that sway amplitude (RMS) was generally

higher in the high-CMP condition, while sway frequency (MPF) and complexity (SEn) were generally lower

in the high- compared the low-CMP condition. Below the adjusted *p*-values (to account for multiple

comparisons) are presented, the uncorrected *p*-values can be found in supplementary data 1.

Sway amplitude (RMS)

Statistical analyses for sway amplitude (RMS) revealed significant main effects for the CMP manipulation

in both the ML-direction, $F(1,24)$ = 10.4, p_{adj} = 0.006, η_p^2 = .30 and the AP-direction, $F(1,24)$ = 14.5, p_{adj} =

0.004, η_p^2 = .38. Contrasts revealed that sway amplitude was lower in both directions in the low-CMP

condition compared to the high-CMP condition (Figure 2; ΔRMS-ML = -1.3 mm; ΔRMS-AP = -1.9 mm). For

stability, a main effect was observed in the AP-direction, $F(1,24)$ = 22.2, p_{adj} = 0.001 η_p^2= .48, but not in the

ML-direction, $F(1,24)$ = .004, p_{adj} = 0.95, η_p^2< .001. Contrasts revealed that sway amplitude in the AP-

direction was lower in the stable compared to the unstable surface condition (ΔRMS-AP = -1.7 mm). No

significant interaction effect was observed in either direction (ML: $F(1,24)$ = 0.10, p_{adj} = 0.76, η_p^2 = .004;

AP: $F(1,24)$ = 0.33 p_{adj} = 0.57, η_p^2 = .014).

Sway frequency (MPF) and complexity (SEn)

A significant main effect of the CMP manipulation on the frequency of sway (MPF) in the AP-direction was

found, $F(1,24)$ = 6.37, p_{adj} = 0.022, η_p^2 = .21, while no main effect was observed in the ML-direction,

$F(1,24)$ = 1.81, p_{adj} = 0.19, η_p^2 = .07. Contrasts revealed that MPF in AP-direction was significantly higher in

the low-CMP compared to the high-CMP condition (ΔMPF-AP = 0.04 Hz). A main effect of stability was

observed in the ML-direction ($F(1,24)$ = 19.5, p_{adj} = 0.001, η_p^2= .45) and AP-direction ($F(1,24)$ = 17.9, p_{adj} =

0.001, η_p^2 = .43). Contrasts revealed that sway frequency was significantly lower in the stable compared

to the unstable surface condition (ΔMPF-ML = -0.061 Hz; ΔMPF-AP = -0.056 Hz). No significant interaction

effects were observed (ML: $F(1,24)$ = 0.011, p_{adj} = 0.92, η_p^2< .001; AP: $F(1,24)$ = 0.062 p_{adj} = 0.81, η_p^2 =

.003).

A significant main effect of the CMP manipulation on the complexity of sway (SEn) was found in the ML-

direction, $F(1,24)$ = 13.0, p_{adj} = 0.004, η_p^2 = .35, and AP-direction, $F(1,24)$ = 10.2, p_{adj} = 0.006, η_p^2= .30. In

both directions, the complexity of sway was higher in the low-CMP compared to the high-CMP condition

(ΔSEn-MP = 0.067; ΔRMS-AP = 0.067). A main effect of stability was observed in ML-direction, $F(1,24)$ =

6.41, p_{adj} = 0.027, η_p^2= .21, but not in AP-direction, $F(1,24)$ = 3.44, p_{adj} = 0.091, η_p^2 = .13. Contrasts revealed

that in the stable surface condition the complexity of sway in ML direction was significantly lower in the low-CMP compared to the high-CMP condition (ΔSEn-MP = -0.064). No significant interaction effect was observed in either direction (ML: $F(1,24) = 1.08$, $p_{adj} = 0.31$, $\eta_p^2 = 0.043$; AP: $F(1,24) = 0.038$, $p_{adj} = 0.85$, $\eta_p^2 = .002$).

Figure 2. Measures of postural stability in the four experimental conditions: Sway amplitude (RMS in mm) in medio-lateral (**A**) and anterior-posterior (**B**) direction, sway frequency (MPF in Hz) in medio-lateral (**C**) and anterior-posterior (**D**) direction, and sway complexity (SEn) in medio-lateral (**E**) and anterior-posterior

335 (**F**) direction. The bars show the group averages and the 95% CI, with the data from individual participants
336 superimposed. Colours indicate the different experimental conditions.
337

3.3. Intermuscular coherence

Intermuscular coherence was assessed in 12 muscle pairs of the lower leg. Using the collapsed localiser

approach, we identified 4 different frequency bands: 0-5, 5-13, 13-23 and 23-45 Hz (Fig. 3). A peak was

observed in each frequency band for most muscle pairs, but the strength differed between muscle pairs.

For example, the peak in the lowest frequency band (0-5 Hz) was generally stronger in agonistic muscle

pairs (Fig. 3C) than in antagonistic muscle pairs (Fig. 3A). Visual inspection also indicated that

intermuscular coherence was generally stronger in the unstable compared to the stable condition (Fig 3A

and B).

For CMP, some main effects with medium to large effect sizes ($0.13 < \eta_p^2 < 0.23$) were observed especially

in the higher frequencies, across several muscle pairs: PL_d-SOL_d, PL_{nd}-SOL_{nd}, PL_{nd}-GM_{nd}, PL_d-Ta_d, PL_{nd}-TA_{nd},

SOL_d-GM_d and SOL_{nd}-GM_{nd} ($p < .05$). Mainly in the 13-23 Hz frequency, lower coherence was observed in

the high-CMP compared to the low-CMP condition (for visualization see the 13-23 Hz frequency range in

figure 3A). Strongest effects were observed in the dominant ($\eta_p^2 = 0.23$) and non-dominant ($\eta_p^2 = 0.23$)

SOL-GM muscle pair at 13-23 Hz. However, controlling for multiple comparisons rendered these

differences non-statistically significant ($p_{adj} > .05$).

Except from the TA_d-SOL_d and the TA_d-GM_d, significant main effects for stability were observed for all

muscle pairs in the dominant (d) and non-dominant (nd) legs across all frequency ranges ($p_{adj} < .05$;

Supplementary data 2). Consistently, IMC was higher in the unstable compared to the stable condition

(for visualization see figure 3A and C). For most of these main effects, the magnitude of the associated

effect sizes were large ($0.16 < \eta_p^2 < 0.58$). No significant interaction effects were observed for stability and

CMP.

Figure 3. Intermuscular coherence between an antagonist and agonist muscle pair in the dominant leg.
The graphs display the grand-average coherence spectra for the **A)** TA-PL and **C)** GM-PL muscle pairs in
the four experimental conditions. Vertical lines indicate the boundaries of the frequency bands and the
horizontal line the 95% CI. The box plots show the mean (black) and SE (grey) for the **B)** TA-PL and **D)** GM-
PL muscle pairs, in each of the four frequency bands, with individual data points superimposed.

4. Discussion

The aim of the study was to investigate the effects of CMP on IMC and postural control during a quiet

standing task in young healthy adults. In line with our expectations, conditions that promoted CMP led to

increases in sway amplitude (higher RMS) and decreases in frequency (lower MPF) and complexity (lower

SEn) of sway. Further, sway amplitude, frequency and complexity all increased in the unstable compared

to the stable condition. We found some preliminary evidence that IMC may be affected by CMP:

coherence tended to be lower in the 13-23 Hz frequency range for the PL-SOL, PL-GM, PL-TA and SOL-GM

muscle pairs. While these differences had medium to large ($\eta_p^2 = 0.13 - 0.23$) effect sizes, these effects

were not statistically significant when controlling for multiple comparisons ($p_{adj} > .05$). For surface

stability, we observed statistically significant increases in IMC in the unstable conditions for most muscle

combinations and frequency bands. These exploratory analyses suggest that IMC may provide insight into

the neuromuscular effects of conscious processing, but these results need to be replicated before they

can be interpreted more conclusively.

The observed findings related to sway amplitude, and sway frequency and complexity are in line with a

recent study by Kal et al. (2022), who also examined the effects of CMP during static balance within a

group of healthy older adults. Interestingly, these patterns of results are opposite to studies which have

explored the effects of postural threat on balance control (e.g., see Ellmers et al., 2021; Zaback et al.,

2019 or Carpenter et al., 1999). Researchers have previously suggested the 'postural stiffening' strategy

frequently observed during conditions of postural threat may be a consequence of anxiety-related

conscious processing (Adkin & Carpenter, 2018; Young & Williams, 2015). Indeed, increased postural

threat reliably leads to greater conscious control of posture (Huffman et al., 2009; Ellmers et al., 2021;

Zaback et al., 2019). However, these studies typically show a decrease in sway amplitude, combined with

an increase in both frequency and complexity of sway (i.e., opposite patterns of results to those observed

in the present study when CMP was increased). Providing a potential explanation for these seemingly

contradictory patterns of results, Ellmers et al. (2021) have recently suggested that the effects of anxiety

on postural control may be twofold. On the one hand, anxiety leads to postural stiffening (increased sway

frequency and muscular co-contraction) and concurrent increases in movement automaticity (greater

sway complexity). However, on the other hand, anxiety-related conscious processing may serve to

'constrain' these automatic changes (leading to comparatively lower increases in sway frequency and

complexity). Our results support this notion, given that we similarly observed patterns of postural sway

opposite to that of postural stiffening during conditions of conscious processing (e.g., increased sway

amplitude and reduced in sway frequency). Further, the observed patterns of IMC when consciously

processing balance in the present research (e.g., reduced IMC) tended to also be contrary to those

reported previously during conditions of postural threat (Zaback et al., 2022). This provides further

evidence for the notion that conscious processing constrains 'automatic' anxiety-related changes, driving

'opposite' patterns of sway and neuromuscular behaviour as to what is typically seen when balance is

threatened.

The effects of CMP on IMC are less pronounced and not statistically significant when controlling for

multiple comparisons. However, we did observe peaks in IMC in the same frequency bands as previous

studies: 0-5, 5-13, 13-23 and 23-45 Hz (Boonstra et al., 2015; Kerkman et al., 2018). In addition, we

observed a significant increase in IMC across all frequency ranges during the unstable compared to the

stable condition. These effects are in line with previous studies that also observed increased IMC during

tasks which challenge postural control e.g., standing position (e.g., tandem stance) or through inducing

threats (e.g., standing at height; Boonstra et al., 2015; Danna-Dos-Santos et al., 2015; Walker et al., 2020).

These findings indicate that IMC was reliably estimated in the current study. While we did observe

reduced IMC in the 13-23Hz frequency band with a large effect size, the results were not statistically

significant. As these analyses were exploratory investigating differences in coherence in 12 muscle pairs

and 4 frequency bands, it seems likely that the inability to statistically detect these changes in coherence

are due to a lack of statistical power.

While these findings should hence be interpreted with care, it is interesting to note that IMC coherence in

the beta-band was reduced during high compared to low CMP. This was opposite from what we expected

based on studies where IMC was assessed while anxiety was manipulated, leading to 'spontaneous'

increases in conscious balance processing e.g., see Zaback et al (2022). Perhaps previously observed

anxiety-related changes in neuromuscular control of balance (e.g., Zaback et al., 2022) may not be driven

by heightened conscious balance processing. Our results also suggest different underlying mechanisms

for the stability and CMP conditions since for stability the largest effects were observed for lower

frequencies, whereas for CMP large effects were mainly observed at the higher frequencies. IMC at the

lower frequency band is most likely related to co-modulation of muscle activation patterns (Mochizuki et

al., 2006; Saffer et al., 2008) and may also result from resulting in correlated afferent inputs to spinal

motor neurons (Kutch & Valero-Cuevas, 2012). We indeed found that in the unstable condition, IMC

increased together with the increase in sway amplitude (Saffer et al., 2008; Boonstra et al., 2008). In

contrast, IMC in the beta band (15-30Hz) is generally considered to reflect cortical input (Farmer et al,

1993; Grosse et al., 2002), which would agree with reliance of CMP on cortical processes. However, the

apparent reduction of beta-band IMC during the high CMP is unexpected and requires further

confirmation. Participants were more conscious about their balance performance as reflected by higher

state-CMP scores within the high CMP condition, however this self-reported reliance on cortical processes

was not reflected by increased IMC in the beta-band. Potentially, previously studies reported increases in

beta-band IMC within complex tasks and threat conditions (e.g., Zaback et al., 2022) may be driven by

increased arousal rather than specific increases in conscious movement processing.

Although our study was exploratory in nature i.e., the study was not powered for the number of

comparisons made, our results can inform future studies. Following this study, future research should

consider to replicate the effects of CMP on IMC with more statistical power, for example by focusing on

pre-selected muscle pairs and frequency bands. Based on our results we recommend future research to

investigate whether IMC is affected by CMP within the beta band, looking for example at the SOL-GM and

PL-SOL muscle pairs as we observed largest effects for these frequencies and muscle pairs (see

supplementary data 1). Literature also showed that certain personal characteristics such as balance or

cognitive capacity, may influence the effect of CMP (Kal et al., 2022). There is generally much more

variability in these characteristics in patient populations such as older adults, stroke or Parkinson

compared to in young healthy adults. For clinical practice it may therefore be interesting to further

explore the effects of CMP instructions and conditions of postural threat within other (patient)

populations and other (functional) balance tasks e.g., examining a low CMP condition within a threatening

condition and vice versa.

Conclusion

In conclusion, our results demonstrate the presence of common neural input to postural muscles during

bilateral stance while levels of stability and conscious movement processing were challenged. We found

that conscious processing drives behavioural outcomes (i.e., increased postural sway and reduced sway

amplitude), that are opposite to what is typically observed during postural stiffening response (similar to Ellmers et al., 2021, and Kal et al; 2022). Furthermore, exploratory analyses showed reduced IMC in the beta band between several lower limb muscles during high compared to low CMP. While this may suggest that the abovementioned behavioural outcomes are driven by reductions in beta band IMC, these effects were not statistically significant when controlling the false discovery rate and therefore require replication in future studies.

Acknowledgements: The authors would like to thank Oriol Falguera Collboni, Bernadette Bänziger, Pien Alferink, Allanah Smyth, Markus Köpcke and Penelope Edwins who contributed in conducting the experiment and all participants for taking part in the study

Funding: LJ was funded by the Kootstra Talent Fellowship awarded by Maastricht UMC+

Declaration of interest: None

References

Adkin, A. L., & Carpenter, M. G. (2018). New insights on emotional contributions to human postural control. *Frontiers in Neurology, 9*, 1-8. https://doi.org/10.3389/fneur.2018.00789

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing *Journal of the Royal statistical society: series B (Methodological), 57*(1), 289-300.

Boisgontier, M. P., Beets, I. A. M., Duysens, J., Nieuwboer, A., Krampe, R. T., & Swinnen, S. P. (2013). Age-related differences in attentional cost associated with postural dual tasks: Increased recruitment of generic cognitive resources in older adults. *Neuroscience and Biobehavioral Reviews, 37*, 1824-1837. https://doi.org/10.1016/j.neubiorev.2013.07.014

Boonstra, T. (2013). The potential of corticomuscular and intermuscular coherence for research on human motor control. *Frontiers in Human Neuroscience, 7*, 855. https://doi.org/ https://doi.org/10.3389/fnhum.2013.00855

Boonstra, T., & Breakspear, M. (2012). Neural mechanisms of intermuscular coherence: Implications for the rectification of surface electromyography. *Journal of Neurophysiology, 107*, 796-807. https://doi.org/10.1152/jn.00066.2011

Boonstra, T. W., Danna-Dos-Santos, A., Xie, H. B., Roerdink, M., Stins, J. F., & Breakspear, M. (2015). Muscle networks: Connectivity analysis of EMG activity during postural control. *Scientific Reports, 5*, 1-14. https://doi.org/10.1038/srep17830

Boonstra, T. W., Farmer, S. F., & Breakspear, M. (2016). Using computational neuroscience to define common input to spinal motor neurons. *Frontiers in Human Neuroscience, 10*, 2013-2016. https://doi.org/10.3389/fnhum.2016.00313

Boonstra, T. W., Roerdink, M., Daffertshofer, A., Van Vugt, B., Van Werven, G., & Beek, P. J. (2008). Low-alcohol doses reduce common 10- to 15-Hz input to bilateral leg muscles during quiet standing. *Journal of Neurophysiology, 100*, 2158-2164. https://doi.org/10.1152/jn.90474.2008

506   Brickenkamp, R., & Zillmer, E. The d2 Test of Attention. 1998. *Goettingen, Hogrefe*.

508   Carpenter, M. G., Frank, J. S., & Silcher, C. P. (1999). Surface height effects on postural control: A
509   hypothesis for a stiffness strategy for stance. *Journal of Vestibular Research, 9*(4), 277-286.
510   https://doi.org/https://doi.org/10.3233/VES-1999-9405

512   Danna-Dos-Santos, A., Degani, A. M., Boonstra, T. W., Mochizuki, L., Harney, A. M., Schmeckpeper, M.
513   M., Tabor, L. C., & Leonard, C. T. (2015). The influence of visual information on multi-muscle
514   control during quiet stance: a spectral analysis approach. *Experimental Brain Research, 233*,
515   657-669. https://doi.org/10.1007/s00221-014-4145-0

517   Ellmers, T. J., Kal, E. C., & Young, W. R. (2021). Consciously processing balance leads to distorted
518   perceptions of instability in older adults. *Journal of Neurology, 268*, 1374-1384.
519   https://doi.org/10.1007/s00415-020-10288-6

521   Ellmers, T. J., Machado, G., Wong, T. W. L., Zhu, F., Williams, A. M., & Young, W. R. (2016). A validation
522   of neural co-activation as a measure of attentional focus in a postural task. *Gait and Posture,
523   50*, 229-231. https://doi.org/10.1016/j.gaitpost.2016.09.001

525   Ellmers, T. J., & Young, W. R. (2018, 2018/06/01/). Conscious motor control impairs attentional
526   processing efficiency during precision stepping. *Gait & Posture, 63*, 58-62.
527   https://doi.org/https://doi.org/10.1016/j.gaitpost.2018.04.033

529   Farina, D., & Negro, F. (2015). Common synaptic input to motor neurons, motor unit synchronization,
530   and force control. *Exercise and Sport Sciences Reviews, 43*, 23-33.
531   https://doi.org/10.1249/JES.0000000000000032

533   Farmer, S. F., Bremner, F. D., Halliday, D. M., Rosenberg, J. R., & Stephens, J. a. (1993). The frequency
534   content of common synaptic inputs to motoneurones studied during voluntary isometric
535   contraction in man. *The Journal of physiology, 470*(1), 127-155.
536   https://doi.org/https://doi.org/10.1113/jphysiol.1993.sp019851

538   Grosse, P., Cassidy, M. J., & Brown, P. (2002). EEG-EMG, MEG-EMG and EMG-EMG frequency analysis:
539   Physiological principles and clinical applications. *Clinical Neurophysiology, 113*, 1523-1531.
540   https://doi.org/10.1016/S1388-2457(02)00223-7

542   Huffman, J. L., Horslen, B. C., Carpenter, M. G., & Adkin, A. L. (2009). Does increased postural threat
543   lead to more conscious control of posture? *Gait and Posture, 30*, 528-532.
544   https://doi.org/10.1016/j.gaitpost.2009.08.001

546   Kal, E. C., Young, W. R., & Ellmers, T. J. (2022). Balance capacity influences the effects of conscious
547   movement processing on postural control in older adults. *Human Movement Science, 82*,
548   102933. https://doi.org/10.1016/j.humov.2022.102933

550   Kerkman, J. N., Daffertshofer, A., Gollo, L. L., Breakspear, M., & Boonstra, T. W. (2018). Network
551   structure of the human musculoskeletal system shapes neural interactions on multiple time
552   scales. *Science Advances, 4*, 1-11. https://doi.org/10.1126/sciadv.aat0497

554   Kutch, J. J., & Valero-Cuevas, F. J. (2012). Challenges and New Approaches to Proving the Existence of
555   Muscle Synergies of Neural Origin. *PLOS Computational Biology, 8*(5), e1002434.
556   https://doi.org/10.1371/journal.pcbi.1002434

Lake, D. E., Richman, J. S., Griffin, M. P., & Moorman, J. R. (2002). Sample entropy analysis of neonatal
heart rate variability. *American Journal of Physiology-Regulatory, Integrative and Comparative*
Physiology. https://doi.org/https://doi.org/10.1152/ajpregu.00069.2002

Luck, S. J., & Gaspelin, N. (2017). How to get statistically significant effects in any ERP experiment (and
why you shouldn't). *Psychophysiology, 54*, 146-157. https://doi.org/10.1111/psyp.12639

Masters, R. S., Pall, H. S., MacMahon, K. M., & Eves, F. F. (2007). Duration of Parkinson disease is
associated with an increased propensity for "reinvestment". *Neurorehabil Neural Repair, 21*,
123-126. https://doi.org/10.1177/1545968306290728

Masters, R. S. W., Eves, F. F., & Maxwell, J. P. (2005). Development of a movement specific
reinvestment scale. *International Society of Sport Psychology (ISSP) World Congress*.

Mochizuki, G., Semmler, J. G., Ivanova, T. D., & Garland, S. J. (2006, 2006/11/01). Low-frequency
common modulation of soleus motor unit discharge is enhanced during postural control in
humans. *Experimental Brain Research, 175*(4), 584-595. https://doi.org/10.1007/s00221-006-
0575-7

Nandi, T., Hortobágyi, T., van Keeken, H. G., Salem, G. J., & Lamoth, C. J. C. (2019). Standing task
difficulty related increase in agonist-agonist and agonist-antagonist common inputs are driven
by corticospinal and subcortical inputs respectively. *Scientific Reports, 9*, 1-12.
https://doi.org/10.1038/s41598-019-39197-z

Obata, H., Abe, M. O., Masani, K., & Nakazawa, K. (2014). Modulation between bilateral legs and within
unilateral muscle synergists of postural muscle activity changes with development and aging.
Experimental Brain Research, 232, 1-11. https://doi.org/10.1007/s00221-013-3702-2

Orrell, A. J., Masters, R. S. W., & Eves, F. F. (2009). Reinvestment and Movement Disruption Following
Stroke. *Neurorehabilitation and Neural Repair, 23*(2), 177-183.
https://doi.org/10.1177/1545968308317752

Richer, N., & Lajoie, Y. (2020). Automaticity of Postural Control while Dual-tasking Revealed in Young
and Older Adults. *Experimental Aging Research, 46*, 1-21.
https://doi.org/10.1080/0361073X.2019.1693044

Richer, N., Ly, K., Fortier, N., & Lajoie, Y. (2020). Absence of Ankle Stiffening While Standing in Focus
and Cognitive Task Conditions in Older Adults. *Journal of Motor Behavior, 52*, 167-174.
https://doi.org/10.1080/00222895.2019.1599808

Richer, N., Saunders, D., Polskaia, N., & Lajoie, Y. (2017). The effects of attentional focus and cognitive
tasks on postural sway may be the result of automaticity. *Gait and Posture, 54*, 45-49.
https://doi.org/10.1016/j.gaitpost.2017.02.022

Roerdink, M., Hlavackova, P., & Vuillerme, N. (2011). Center-of-pressure regularity as a marker for
attentional investment in postural control: A comparison between sitting and standing
postures. *Human Movement Science, 30*, 203-212.
https://doi.org/10.1016/j.humov.2010.04.005

Saffer, M., Kiemel, T., & Jeka, J. (2008, 2008/02/01). Coherence analysis of muscle activity during quiet stance. *Experimental Brain Research, 185*(2), 215-226. https://doi.org/10.1007/s00221-007-1145-3

Staab, J. P., Balaban, C. D., & Furman, J. M. (2013). Threat assessment and locomotion: Clinical applications of an integrated model of anxiety and postural control. *Seminars in Neurology, 33*, 297-306. https://doi.org/10.1055/s-0033-1356462

Stins, J. F., Roerdink, M., & Beek, P. J. (2011). To freeze or not to freeze? Affective and cognitive perturbations have markedly different effects on postural control. *Human Movement Science, 30*, 190-202. https://doi.org/10.1016/j.humov.2010.05.013

Walker, S., Piitulainen, H., Manlangit, T., Avela, J., & Baker, S. N. (2020). Older adults show elevated intermuscular coherence in eyes-open standing but only young adults increase coherence in response to closing the eyes. *Experimental Physiology, 105*, 1000-1011. https://doi.org/10.1113/EP088468

Watanabe, T., Saito, K., Ishida, K., Tanabe, S., & Nojima, I. (2018). Age-related declines in the ability to modulate common input to bilateral and unilateral plantar flexors during forward postural lean. *Frontiers in Human Neuroscience, 12*, 1-9. https://doi.org/10.3389/fnhum.2018.00254

Young, W. R., & Williams, M. A. (2015). How fear of falling can increase fall-risk in older adults: Applying psychological theory to practical observations. *Gait and Posture, 41*, 7-12. https://doi.org/10.1016/j.gaitpost.2014.09.006

Zaback, M., Adkin, A. L., & Carpenter, M. G. (2019). Adaptation of emotional state and standing balance parameters following repeated exposure to height-induced postural threat. *Scientific Reports, 9*, 1-12. https://doi.org/10.1038/s41598-019-48722-z

Zaback, M., Adkin, A. L., Chua, R., Inglis, J. T., & Carpenter, M. G. (2022). Facilitation and Habituation of Cortical and Subcortical Control of Standing Balance Following Repeated Exposure to a Height-related Postural Threat. *Neuroscience, 487*, 8-25. https://doi.org/10.1016/j.neuroscience.2022.01.012

Zhu, F. F., Poolton, J. M., Wilson, M. R., Maxwell, J. P., & Masters, R. S. W. (2011). Neural co-activation as a yardstick of implicit motor learning and the propensity for conscious control of movement. *Biological Psychology, 87*, 66-73. https://doi.org/10.1016/j.biopsycho.2011.02.004