Evolutionary models predict potential mechanisms of escape from mutational meltdown

Claudia Bank [*,1,2,3], **Mark A. Schmitz** [3] and **Ana Yansi Morales-Arce** [1,2,3]

[1] *Institute of Ecology and Evolution, University of Bern, Bern, Switzerland*
[2] *Swiss Institute of Bioinformatics, Lausanne, Switzerland*
[3] *Gulbenkian Science Institute, Oeiras, Portugal*

Correspondence*:
Claudia Bank
claudia.bank@unibe.ch

2 ABSTRACT

Mutagenic drugs are promising candidates for the treatment of various RNA virus infections.
Increasing the mutation rate of the virus leads to rapid accumulation of deleterious mutation load,
which is proposed to ultimately result in extinction as described by the theoretical concepts of
mutational meltdown and lethal mutagenesis. However, the conditions and potential mechanisms
of viral escape from the effects of mutagenic drugs have not been conceptually explored. Here we
apply a computational approach to quantify the population dynamics and genetics of a population
under high mutation rates and discuss the likelihood of adaptation to a mutagenic drug by means
of three proposed mechanisms: (1) a proportion of "traditional" beneficial mutations that increase
growth/fitness, (2) a mutation rate modifier (i.e., evolution of resistance to the mutagenic drug)
that reduces the mutation rate, and (3) a modifier of the distribution of fitness effects, which
either decreases or increases deleterious effects of mutations (i.e., evolution of tolerance to the
mutagenic drug). We track the population dynamics and genetics of evolving populations and find
that successful adaptations have to appear early to override the increasing mutational load and
rescue the population from its imminent extinction. We highlight that the observed stochasticity
of adaptation, especially by means of modifiers of the distribution of fitness effects, is difficult
to capture in experimental trials, which may leave potential dangers of the use of mutagenic
treatments unexposed.

Keywords: mutation load, adaptation, lethal mutagenesis, mutational meltdown, extinction, eco-evolutionary dynamics, mutagenic
drugs, resistance evolution

1   INTRODUCTION

The search for therapeutics against virus infections remains a challenge in human health research.
Understanding within-host infections and the potential for adaptation (including resistance) to new
drugs requires studying the eco-evolutionary dynamics of virus populations. One heatedly discussed
potential treatment option is particularly rooted in evolutionary theory: mutagenic drugs like favipiravir or
molnupiravir act by inducing mutational meltdown (1, 2, 3) or lethal mutagenesis (4, 5); for a discussion of
the respective terms see Matuszewski et al. (6). Mutational meltdown occurs when the virus population
accumulates so many deleterious mutations that the population growth rate falls below 1, which induces
extinction. Importantly, the evolutionary theory of mutational meltdown rests on the knowledge that most

new mutations are deleterious. Moreover, the action of a mutagenic drug is a population process. Other
drugs act by killing individual cells or viruses, or by stopping their proliferation. Conversely, a key aspect
of mutational meltdown, induced by a mutagenic drug, is that the population as a whole is weakened
through the successive accumulation of deleterious mutations, which ultimately leads to extinction.

Mutagenic treatment is particularly promising against RNA viruses because of their comparably high
mutation rates (7, 8), which suggests that only a slight increase in the mutation rate is sufficient to induce
mutational meltdown (9, 10). Mutagenic drugs such as favipiravir and molnupiravir increase the mutation
rate by targeting the RNA polymerase. After phosphorylation inside the cell their active forms become
incorporated as nucleoside analogues in the newly synthesized RNA chain (11, 12, 13, 14). Experimental
and clinical trials of mutagenic drugs against Sars-Cov-2 have shown promising results (15, 16, 17),
although Sars-Cov-2 has a relatively low mutation rate as compared to other RNA viruses. However, recent
clinical data indicated that their effectiveness was inversely proportional to the severity of symptoms (18) or
completely absent (19). A possible explanation is a counteracting proof reading mechanism called ExonN,
that avoids the incorporation of mutagenic drugs into the virus genome (20, 21). The combination of
different mutagenic drugs, able to overcome a proofreader, to treat Covid-19 patients seems still promising
and is currently under debate (22, 23).

Several examples of adaptation of RNA viruses to mutagenic drugs have been reported (reviewed in (24).
For example, rivabirin resistance was observed *in-vitro* in poliovirus and patient hepatitis C virus strains
(25, 26), whereas remdesivir was shown to induce resistance in *in-vitro* Sars-CoV-2 (27). High doses of the
mutagenic drug favipiravir did not help patients that were infected with the Ebola virus (28) and resistance
was also described in chikungunya virus and in enteroviruses (29, 30). Yet, favipiravir is one of the most
effective compounds inducing extinction in *in-vitro* influenza A virus, foot-and-mouth disease virus, and
West Nile virus (13, 31, 32). Across various screens in the laboratory in influenza virus (12, 33, 34, 35),
only one potential resistance mutant against favipiravir has been identified and experimentally validated to
date (34, 35). Because of its special mode of action, it was hypothesized that the effect of mutagenic drugs
may depend on the viral replication system, the genome structure, and the time at which treatment starts
(10, 23).

Although the foundation for the proposed success of mutagenic treatments lies in evolutionary theory,
evolutionary theory of mutational meltdown also predicts that mutagenic drugs are not without risks
(36). Although most new mutations are deleterious, a mutagenic drug increases not only the chances of
new deleterious but also of new beneficial mutations. Increasing the mutation rate could allow the virus
population to "find" fit genotypes (e.g., including multiple, individually deleterious mutations) that would
be unreachable under the default mutation rate. The longer a virus population survives under mutagenic
drug treatment, the larger the possibility that adaptive genotypes may appear, which could cause damage
not just in the current host, but also when transmitted to new, untreated, hosts (37, 38).

Here we approach this problem by proposing three mechanisms of adaptation to mutagenic drugs from the
view of an evolutionary biologist. Using computer simulations, we quantify the eco-evolutionary dynamics
of a clonal (virus) population under high mutation rates that has access to either of these mechanisms
and show that each can lead to escape from mutational meltdown under different conditions. Because the
actual eco-evolutionary dynamics of a virus population inside a human host are complex and many of the
parameters are unknown, we restrict ourselves to a small and fixed set of parameters. Thus, our study is
meant as a proof-of-concept to spur discussion and future work.

We discuss the escape of a virus population from mutational meltdown through three adaptation
mechanisms:

1. Beneficial, growth-rate increasing mutations. Because accumulation of deleterious mutation load is
unstoppable by such mutations, a continual input of a fraction of growth rate increasing mutations is
necessary to save the population from extinction that strongly depends on the carrying capacity.

2. A mutation rate modifier, which is a mutation that reduces the mutation rate of the virus genome-wide.
We show that this type of mutation has to invade early to lead to escape from mutational meltdown,
and argue that this mechanism qualifies as evolution of *drug resistance*.

3. A modifier of the distribution of fitness effects (in short: DFE modifier), which is a mutation that either
dampens or exaggerates the deleterious effect of all mutations that its carrier is accumulating. We show
that both dampening and exaggerating mutational effects can lead to escape from mutational meltdown
and argue that this mechanism qualifies as evolution of *drug tolerance*.

84 The distinction between resistance and tolerance comes from the field of host-pathogen interactions
85 (39, 40). We have not seen these concepts discussed in the evolutionary literature with respect to different
86 mechanisms of adaptation. We show below that adaptation via evolution of resistance *versus* tolerance
87 lead to fundamentally different implications for the eco-evolutionary dynamics of virus populations in the
88 presence (and absence) of a mutagenic drug.

2   MATERIALS AND METHODS

2.1   Eco-evolutionary dynamics without adaptation

Similar to Lansch-Justen et al. (41), we model the intra-host eco-evolutionary dynamics of a non-
recombining asexual virus population with a carrying capacity of K, which expands from a founding
population size of N_0 mutation-free individuals. The initial absolute growth rate (i.e., the average number
of offspring that a mutation-free genotype produces) is w_0. Each generation, every individual gains a
number of deleterious mutations drawn from a Poisson distribution with mean U_{del}. We consider an
infinite-site model, i.e., every mutation occurs at a new site in the genome; back mutations are neglected.
After mutation, the offspring number, i.e. the number of clonal copies of each individual, is drawn from a
Poisson distribution with its mean set to the absolute growth rate $w(k)$ of the respective individual, and the
individual subsequently dies. We assume that each deleterious mutation has the same effect s_{del}. We consider
a multiplicative model that determines the absolute growth rate of an individual, $w(k) = w_0 \cdot (1 + s_{\text{del}})^k$,
where k is the number of mutations that the individual has accumulated. Finally, if the population size is
above the carrying capacity K, individuals are eliminated randomly to adjust the population size to K.

2.2   Modeling the three adaptation mechanisms

We separately study three different models of adaptation mechanisms. In Section 3.2 we study growth-rate
increasing mutations that occur at mutation rate U_{ben} with effect size $-s_{\text{del}}$. Here, the absolute growth rate
of an individual is $w(k, i) = w_0 \cdot (1 + s_{\text{del}})^k (1 - s_{\text{del}})^i$, where i denotes the number of beneficial mutations
the individual carries. Thus, every beneficial mutation neutralizes one deleterious mutation.

In Section 3.3 we study mutation rate modifiers. A modifier of the mutation rate occurs at rate U_{mutMod}
per genome per generation. The absolute growth rate of individuals remains the same as in the basic model,
but the mutation rate of carriers of at least one mutation modifier is reduced to $U_{\text{del}} \cdot f_{\text{mutMod}}$, where f_{mutMod}
is the effect size of the mutation modifier. The effect size is defined multiplicatively, i.e., if $f_{\text{mutMod}} = 1$,

the mutation rate remains unaltered by the modifier mutation, and if $f_{\mathrm{mutMod}} = 0.1$, the mutation rate is
reduced to 10% of its original value.

In Section 3.4 we study modifiers of the distribution of fitness effects (DFE). A DFE modifier occurs
at rate U_{dfeMod} per genome per generation. Here, the absolute growth rate of a carrier of at least one
DFE modifier mutation is $w^*(k) = w_0 \cdot (1 + s_{\mathrm{del}} \cdot f_{\mathrm{dfeMod}})^k$, where f_{dfeMod} is the effect size of the DFE
modifier. Thus, if $f_{\mathrm{dfeMod}} = 1$, the modifier mutation leaves mutational effects unaltered, whereas DFE
modifiers with $f_{\mathrm{dfeMod}} > 1$ make every existing and new mutation more deleterious, and DFE modifiers
with $f_{\mathrm{dfeMod}} < 1$ make every existing and new mutation less deleterious.

2.3 Simulations and choice of parameters

Throughout the paper, we show results from simulations of the above-described models with carrying
capacities $K \in \{500, 10000\}$ and a founding population size $N_0 = 10$. Each simulation ends when the
population goes extinct, or after 5000 generations. We set the initial absolute growth rate to $w_0 = 2$ and the
deleterious mutation rate per genome per generation to $U_{\mathrm{del}} = 0.2$. In Sections 3.2-3.4, the effect size of a
deleterious mutation is $s_{\mathrm{del}} = -0.023$. In the absence of any adaptation mechanism, the mean extinction
time for our standard parameter set for $K = 500$ is 592.44 ± 69.28. Here, the minimum and maximum
extinction times observed in > 10000 simulations were 384 and 931, respectively. For our standard
parameter set for $K = 10000$, the mean extinction time is 1377.46 ± 157.36. The minimum and maximum
extinction times observed over > 10000 simulations were 947 and 2141 respectively. All results are
based on simulations implemented in Julia v.1.0 or v.1.6.0. The annotated simulation and analysis code is
available at `https://gitlab.com/evoldynamics/ms-mutational-meltdown-2021` and
will be archived on Zenodo upon publication of the manuscript.

3 RESULTS

3.1 Known features of extinction under mutational meltdown

Since the 1990's, many theoretical studies have quantified mutational meltdown (41). We here reiterate
some of these results using our simulations to familiarize the reader with the most important features of
this evolutionary process. As described in Lynch et al. (3), Lansch-Justen et al. (41), the path to extinction
by mutational meltdown consists of three phases. In the first phase, a clonal, non-recombining, deleterious-
mutation-free population invades a new host, where it expands and accumulates deleterious mutations.
Eventually, due to the combined effects of random genetic drift and mutation pressure, the fittest genotype
will be lost, which marks the beginning of the second phase. In the second phase, Muller's ratchet leads to
the step-wise loss of the fittest genotype and an almost linear decrease in mean fitness, until the absolute
growth rate of the population falls below 1. At this point, the third phase of rapid extinction begins, during
which the population size decreases rapidly and selection is ineffective.

[Figure 1 should be placed here.]

For the purpose of this paper, we reiterate three findings from these studies. Firstly, increasing the
mutation rate has the strongest effect on reducing the time to extinction under mutational meltdown (41).
This result is important because it indicates that the meltdown process can be induced by small alterations
in the mutation rate (as attempted by mutagenic drugs) and that the resulting extinction is relatively robust
to other parameters such as the carrying capacity (which is difficult to measure or control inside a host).
Secondly, the variation in extinction times is relatively small for the same set of parameters (see Figure
1; 41). This result is important because it allows us to identify successful escape as an extension of the

extinction time (both in simulations, and potentially also inside the host). Finally, the extinction time is
minimized for intermediately strong deleterious selection coefficients (see Figure 1; 2). This is because
Muller's ratchet leads to the fastest loss in mean fitness when selection coefficients are too small to be
purged immediately but large enough to reduce the mean fitness significantly (42). This result is important
because it determines which types of deleterious mutations drive the meltdown process, and it is the basis
for considering DFE modifiers as a potential rescue mechanism.

157 3.2   A continual input of growth-rate increasing mutations can halt Muller's ratchet

Evolutionary theory often considers adaptation as a process of accumulation of beneficial mutations,
which are mutations with a positive "selection coefficient". Straightforwardly, these can be interpreted
as growth-rate increasing mutations. For example, a resistance mutation often increases the growth rate
of a pathogen in the presence of a drug. Here, we studied how beneficial mutations of equal size as the
deleterious selection coefficient can help the population escape from mutational meltdown. Our simulations
show that mutational meltdown can be prevented when a sufficient ratio of growth-rate increasing mutation
is available. The ratio of beneficial to deleterious mutations that enables escape from mutational meltdown
strongly depends on the carrying capacity of the population (see Figure 2). Here, the population can escape
from mutational meltdown because Muller's ratchet is halted by the continual input of beneficial mutations,
as previously modeled by Silander et al. (43) and Goyal et al. (44).

[Figure 2 should be placed here.]

169 3.3   Reliable escape from mutational meltdown through mutation rate modifiers

Drug resistance occurs when a pathogen manages to "inactivate" the action of the drug such that it can
normally propagate in the presence of the drug. In the case of a mutagenic drug, resistance occurs when a
mutation alters the mutation rate of the virus in the presence of the drug, such that mutational meltdown
is prevented. Here, we model such mutation rate modifiers, which occur at rate U_{mutMod} and which have
effect size f_{mutMod}. The occurrence rate U_{mutMod} determines the mutational target size (how frequently
are such mutations available?) and the effect size indicates by how much the modifier mutation alters the
mutation rate (i.e., $f_{\mathrm{mutMod}} = 0.1$ means that the modifier decreases the mutation rate of its carriers to 10%
of its previous value).

[Figure 3 should be placed here.]

The effect size of the modifier strongly affects whether the population can escape mutational meltdown
(Figure 3). The effect size of the modifier has to be large enough such that mutational meltdown is prevented
at the modified mutation rate, had it been present from the beginning. Moreover, in order to allow for escape
of the population from mutational meltdown, the resistance mutation has to appear early enough to be able
to spread across the population before the lineage carrying the mutation modifier has accumulated too many
deleterious mutations to escape extinction. Therefore, higher carrying capacities K and larger occurrence
rates of the modifier U_{mutMod} facilitate resistance evolution by means of a mutation rate modifier. When
the product of the carrying capacity and the occurrence rate of the modifier become small, whether the
modifier invades and extends the extinction time becomes less deterministic, even for modifiers of large
effect (Figure 3C).

3.4 Both negative and positive DFE modifiers can lead to escape from mutational meltdown

A modifier of the distribution of fitness effects can allow for escape from mutational meltdown in two ways. A DFE modifier can slow down Muller's ratchet either by decreasing the effect of deleterious mutations (i.e., by weakening their effect) or by increasing their effect size (i.e., by making them more deleterious). This is true for intermediate-size deleterious mutations, for which the extinction time is smaller than for small or large-effect deleterious mutations (see Figure 1). We model DFE modifiers that occur at a rate U_{dfeMod} and with effect size f_{dfeMod}. Again, the occurrence rate U_{dfeMod} determines the mutational target size. The effect size determines whether a DFE modifier is a "positive" modifier with $0 < f_{\mathrm{dfeMod}} < 1$, i.e., the effect of deleterious mutations is weakened, or whether it is a "negative" modifier with $f_{\mathrm{dfeMod}} > 1$, i.e., the effect of deleterious mutations becomes stronger.

[Figure 4 should be placed here.]

We find that both positive and negative DFE modifiers can lead to escape from mutational meltdown. The dynamics and probabilities of this escape differ greatly between the two types of modifiers (Figure 4, Supplementary Figure 1). For positive modifiers, a large effect size is necessary to lead to successful escape of the population from mutational meltdown for the whole duration of the simulations. When a positive DFE modifier has a sufficiently large effect size to lead to escape from mutational meltdown, it usually is successful at leading to escape from extinction, because all carriers of the modifier tend to be much fitter than the genotypes without the modifier. Thus, as soon as it appears, a positive DFE modifier provides a large benefit to its carriers. A negative DFE modifier acts fundamentally differently. It initially makes its carrier and their offspring less fit than any genotype with the same number of deleterious mutations that do not carry the DFE modifier. Thus, it initially has a (potentially) large deleterious effect on its carriers. Only when the subpopulation that carries the DFE modifier survives for several generations, the modifier becomes effective by means of efficient purging of new and segregating deleterious mutations. Thus, negative DFE modifiers rarely invade the population. When they invade, they lead to successful escape from mutational meltdown. We observe a strong dependence of the escape probability (and the invasion probability of a negative DFE modifier) on the product of the occurrence rate U_{dfeMod} and the carrying capacity K. That is because a negative DFE modifier has to appear on a genotype with no or very few deleterious mutations in order to be able to spread through the population, and this possibility is increased by maximizing the product of U_{dfeMod} and K.

4 DISCUSSION

Mutagenic drugs are under close investigation regarding potential treatment of RNA virus infections. Their mode of action is rooted deeply in evolutionary theory. Evolutionary theory also predicts mechanisms by which adaptation can occur. Here we propose and discuss three potential and fundamentally different adaptation mechanisms for adaptation to mutagenic drugs. We provide a proof-of-concept simulation study in order to encourage discussion about mechanisms of adaptation to mutagenic drugs and development of methods to infer such mechanisms. In addition, our work delivers a note of caution about potential rare-event adaptations that are almost impossible to predict with current experimental approaches.

4.1 Rethinking 'beneficial' mutations

The three adaptation mechanisms presented here act either by successively increasing the growth rate of the virus, by altering the mutation rate of the virus, or by altering the fitness effects of all deleterious

mutations that the virus is accumulating (and has accumulated). Whereas all three adaptation mechanisms
are ultimately beneficial to the virus population when they successfully lead to survival in the presence
of the drug, only the first category consists of 'beneficial' mutations as they are usually considered in
evolutionary biology.

Growth-rate increasing mutations as presented in the first model, do not provide resistance in the strict
sense, as their continual input is necessary to allow for the survival of the population under high mutation
rates over longer time scales. Notably, our model of growth-rate increasing mutations is greatly simplified
by assuming that all beneficial mutations have the same effect size, which is the same effect size as that of
deleterious mutations. Also, due to the neglect of back mutation in our model, beneficial (and deleterious)
mutations occur at the same rate at all times. Alternative models of fitness increase that incorporate a
fitness optimum, such as Fisher's Geometric Model (see, e.g., 45), for example, are expected to yield
different results regarding escape from mutational meltdown by means of growth-rate increasing mutations
as compared to our simplified model. Largely independently of the underlying model, we expect that for
successful escape, a serial compensation of the continually appearing deleterious mutations by beneficial
mutations has to occur, whereas mutation and DFE modifiers can allow for a significant extension of the
extinction time by means of a single mutation.

A mutation rate modifier that may reverse the mutation rate of the virus to its value in the absence of the
drug can be considered a *drug resistance mutation*. That is because a population that carries a mutation
rate modifier can proliferate in the presence of the mutagenic drug similar to the original genotype in the
absence of the drug. Mutation-rate modifying mutations are frequently detected in experimental studies
of microorganisms, for example in *E. coli* (e.g. 46, 47). However, both theoretical and empirical work are
contentious as to how much selection acts on mutation rates and whether an equilibrium mutation rate will
eventually be obtained (e.g., 48, 49, 50, 51, 52, 53, 54). Interestingly, the most well-studied adaptation to
the mutagenic drug favipiravir, identified in the laboratory, was described as a mutation rate modifier (34).

Finally, a DFE modifier as defined here could be considered a *drug tolerance mutation*, because it
does not alter the action of the drug, yet allows the virus population to survive in its presence. Here,
the mutation rate of the virus remains high. At the same time, the altered selection coefficients of all
mutations change the eco-evolutionary dynamics such that Muller's ratchet becomes less severe and the
population can cope with the increased mutation pressure. A DFE modifier is a specific form of epistasis
(i.e., the genetic-background dependence of mutational effects), where the modifier allele alters mutational
effects genome-wide. Sydykova et al. (55) have recently explored a related model of tunable epistasis,
in which they found that epistasis can evolve in different directions under high mutation rates. Whether
weakening the effects of deleterious mutations can lead to survival has been modeled in the research area
of quasi-species theory, where a so-called "error threshold" determines the condition in which mutation
overwhelms selection (e.g., for finite populations, 56, 57). This leads to the inability of a population to
maintain its genetic information, which can in general be considered similar to mutational meltdown (58,
but see 59). Moreover, the evolution of more severe effects of deleterious mutations has been theoretically
studied with respect to models of drift robustness (60), quasi-species evolution (e.g., 61) and epidemiology
(62).

There is a two-fold potential danger imposed by the successful invasion of a DFE modifier. Firstly, by
changing the distribution of fitness effects of mutations globally, the presence of a DFE modifier may
create a virus that evolves differently than its ancestral strain also in the absence of the drug, and hence in
future infections of non-treated host. Secondly, by surviving in the treated host under high mutation rates,
the virus population may be able to access adaptive genotypes in the sequence space that are not reachable

under normal mutation rates (e.g., because they require multiple mutations). Although any prolonged survival of the virus population poses a danger regarding adaptation of the virus (to both the host immune system and the drug; 36), a DFE modifier is likely to facilitate such adaptation.

We are not aware of any empirical evidence of DFE-modifying adaptations in viruses. Of note, detecting such mutations is not straightforward because of their indirect effects, and a careful experimental design would be necessary to distinguish them from mutation rate modifiers, for example. Moreover, the different adaptation mechanisms described here could occur together (or interfere with each other), or many mutations of small DFE-changing-effect could be involved in adaptation to mutagenic drugs. In higher organisms, mutations in chaperones or global regulators could have the effect of DFE modifiers. For example, mutations of global transcription regulators, such as RpoB in *E. coli*, can change the expression patterns of hundreds or thousands of genes (63, 64). Chaperones (e.g., Hsp90 65 or DnaK 64) buffer the effects of mutations, and by weakening or strengthening the effect of the chaperone through mutation, the distribution of fitness effects would likely be altered either pathway- or genome-wide.

4.2 Negative DFE modifiers would not be detected in experimental screens for adaptations

An important result from the work presented here is that the escape probability for a very efficient adaptation mechanism (efficient if it invades the population) can be very small. Specifically, a negative DFE modifier, i.e., a mutation that makes other mutations more deleterious, has a very low probability to establish in the population in the first place. That is because it initially harms its carriers by reducing their fitness; only in the longer term the more efficient purging of deleterious mutations becomes an advantage. Their rare invasion implies that negative DFE modifiers would in general not be predicted by laboratory evolution experiments that are commonly used to screen for potential adaptations to new drugs (e.g. 12, 33, 34). If the probability of invasion of a negative DFE modifier is 1/10000, thousands of replicates of an evolution experiment would be necessary to observe it in the lab, whereas usually only ≈ 3 replicates are performed in such experiments. However, once millions of patients are treated with the drug, the mutation would have ample opportunity to invade and spread. That is why theoretical and computational discussion and study of adaptation mechanisms are essential to complement experimental studies.

4.3 How to dissect the adaptation mechanism of a candidate mutation

Once candidate mutations of adaptation to mutagenic drugs have been identified (either from laboratory experiments or from the wild), it will be important to dissect the underlying adaptation mechanism. We propose that this could be possible by comparing the eco-evolutionary dynamics of virus population with and without the candidate mutation in the presence and absence of the drug. For example, if a resistance mutation reduces the mutation rate, less segregating and fixed mutations should be observed over time when comparing a population that carries this mutation with a non-mutated reference type. Moreover, if the resistance mutation alters the mutation rate independent of the drug treatment, the same pattern should be observed in non-treated populations. Interestingly, negative DFE modifiers should also display a pattern of reduced mutation accumulation in the presence and absence of the drug, because selection against deleterious mutations becomes more efficient. However, a DFE modifier does not prevent deleterious mutations from appearing, and instead leads to stronger selection against them after their appearance. We thus expect that the observed frequency spectrum of segregating mutations is different between a DFE modifier and a mutation rate modifier, with more low-frequency mutations for the DFE modifier. Conversely, since a positive DFE modifier reduces selection against deleterious mutations while leaving

316 the mutation rate elevated, its signature should include elevated mutation accumulation compared to
317 the reference type. Future work should further explore the predicted signatures of different adaptation
318 mechanisms.

4.4 Limitations and future steps

The goal of this study was to conceptually introduce different types of adaptations to mutagenic drugs
and highlight and discuss their different action from an evolutionary point of view. Our results should
be seen as a qualitative illustration of potential effects of different adaptation mechanisms on looming
extinction, each of which we study separately, but we caution against interpreting the reported numbers at
face value. Our model is a simple depiction of intra-host evolution, in which a small number ($N_0 = 10$)
of mutation-free genotypes infect a host and grow exponentially to a predetermined carrying capacity
($K = 500$ or $K = 10000$) that is maintained until the mean growth rate of the population falls below 1.
The increased mutation rate (as caused by a mutagenic drug) is present from generation zero onwards,
whereas in natural scenarios treatment would only begin after the infection was detected. We consider as
escape from mutational meltdown when a population that would have gone extinct without the adaptation
mechanism in question survives for $t_{max} = 5000$ generations . In natural populations, t_{max} may reflect
the time by which transmission has occurred, or by which the immune system is responding. Because we
do not know the population dynamics of real virus populations in the wild, our parameters are somewhat
arbitrarily chosen. For example regarding the initial population size, recent evidence has pointed towards
infection bottlenecks (in influenza) being very small, with sometimes a single virion infecting a new host
((66, 67)). Moreover, although the census size of virions inside a host is probably on the order of millions,
a much lower "effective" population size could reflect the evolutionary dynamics within a human host.
In laboratory evolution experiments, the constant population size that best describes the evolutionary
dynamics of populations under serial passaging is usually on the order of few hundreds, which is why
we chose $K = 500$ as one of the studied population sizes (68, 33). Reassuringly, our results showed
that the proposed adaptation mechanisms occurred at both tested population sizes, and especially for the
DFE modifier (and surprisingly for both negative and positive DFE modifiers), we observed larger escape
probabilities for larger population sizes, indicating that this escape mechanism is relevant also for (much)
larger populations sizes (and thus, across all regimes of the dynamics of Muller's ratchet, see, e.g. 69). We
further ignore potential back mutations, and we do not consider the ability of some RNA viruses (including
influenza) to rearrange, which is a type of recombination between genomic segments that could slow down
Muller's ratchet and stall extinction. Finally, we only consider a single, constant, deleterious effect that is
common to every deleterious mutation as opposed to the distribution of fitness effects that is observed in
natural and laboratory populations (e.g., 70). Relieving some of these assumptions is an important direction
for future work.

4.5 Relevance for the use of mutagenic drugs in human populations

As argued above, we have studied adaptation to mutagenic drugs in a very simplified model that makes
no attempt at reflecting the complexity of a human host. Nevertheless, our work cautions against the routine
use of mutagenic drugs in human populations by demonstrating that there is much to learn about potential
adaptation to mutagenic drugs. Adding to the arguments proposed by Nelson and Otto (36), our work shows
that unpredictable types of adaptations can occur under wide-spread use of mutagenic drugs. Specifically,
we consider tolerance mutations by means of a negative DFE modifier a type of adaptation that may affect
evolution not only in current but also future (untreated) hosts. Moreover, the probability of adaptations (to
both the drug and the host immune system) occurring and spreading to new hosts further increases when a

mutagenic drug is not administered at sufficiently high doses or until the meltdown process is completed,
as would be the case if a mutagenic drug is approved for household use.

5 CONTRIBUTION TO THE FIELD

Mutagenic drugs are currently discussed regarding their potential use against RNA viruses such as influenza
and Sars-Cov-2. The principle of their action is rooted in the evolutionary theory of mutational meltdown,
according to which the increasing load of deleterious mutations leads to collapse and extinction of a
population. It is contentious whether and how RNA viruses can develop resistance to mutagenic drugs,
and thereby escape mutational meltdown. Here, we present and discuss three potential mechanisms of
adaptation to mutagenic drugs and model the eco-evolutionary dynamics when such mutations are available.
Whereas growth-rate increasing mutations have to be continually available to prevent extinction of the
virus population under drug treatment, modifiers of the mutation rate or the fitness effects of deleterious
mutations can allow escape from mutational meltdown with a single mutation. We highlight the stochastic
nature of this process that will rarely be captured by laboratory experiments. Moreover, we discuss the
different adaptation mechanisms as examples of drug resistances versus drug tolerance.

CONFLICT OF INTEREST STATEMENT

372 The authors declare that the research was conducted in the absence of any commercial or financial
373 relationships that could be construed as a potential conflict of interest.

FUNDING

374 This work was supported by ERC Starting Grant 804569 (FIT2GO), HFSP Young Investigator Grant
375 RGY0081/2020 and EMBO Installation Grant 4152 to CB.

ACKNOWLEDGMENTS

376 The authors are grateful for discussion of the project in the THEE/Evoldynamics lab. Calculations were
377 performed on UBELIX (http://www.id.unibe.ch/hpc), the HPC cluster at the University of Bern.

DATA AVAILABILITY STATEMENT

378 The complete annotated documentation of the computational analyses of this paper will be archived
379 on Zenodo upon publication of the paper and is currently deposited at `https://gitlab.com/`
380 `evoldynamics/ms-mutational-meltdown-2021`.

REFERENCES

1. Lynch M, Gabriel W. Mutation load and the survival of small populations. *Evolution* **44** (1990)
1725–1737.
2. Gabriel W, Lynch M, Bürger R. Muller's ratchet and mutational meltdowns. *Evolution* **47** (1993)
1744–1757.
3. Lynch M, Bürger R, Butcher D, Gabriel W. The mutational meltdown in asexual populations. *J. Hered.*
84 (1993) 339–344.
4. Bull J J, Sanjuán R, Wilke C O. Theory of lethal mutagenesis for viruses. *J. Virol.* **81** (2007) 2930–2939.

5. Bull JJ, Sanjuán R, Wilke CO. CHAPTER 9 - lethal mutagenesis. Domingo E, Parrish CR, Holland JJ, editors, *Origin and Evolution of Viruses (Second Edition)* (London: Academic Press) (2008), 207–218.

6. Matuszewski S, Ormond L, Bank C, Jensen JD. Two sides of the same coin: A population genetics perspective on lethal mutagenesis and mutational meltdown. *Virus Evol* **3** (2017) vex004.

7. Sanjuán R, Nebot MR, Chirico N, Mansky LM, Belshaw R. Viral mutation rates. *J. Virol.* **84** (2010) 9733–9748.

8. Sanjuán R, Domingo-Calap P. Mechanisms of viral mutation. *Cell. Mol. Life Sci.* **73** (2016) 4433–4448.

9. Peck KM, Lauring AS. Complexities of viral mutation rates. *J. Virol.* **92** (2018).

10. Jensen JD, Stikeleather RA, Kowalik TF, Lynch M. Imposed mutational meltdown as an antiviral strategy. *Evolution* **74** (2020) 2549–2559.

11. Furuta Y, Gowen BB, Takahashi K, Shiraki K, Smee DF, Barnard DL. Favipiravir (t-705), a novel viral RNA polymerase inhibitor. *Antiviral Res.* **100** (2013) 446–454.

12. Baranovich T, Wong SS, Armstrong J, Marjuki H, Webby RJ, Webster RG, et al. T-705 (favipiravir) induces lethal mutagenesis in influenza a H1N1 viruses in vitro. *J. Virol.* **87** (2013) 3741–3751.

13. Arias A, Thorne L, Goodfellow I. Favipiravir elicits antiviral mutagenesis during virus replication in vivo. *Elife* **3** (2014) e03679.

14. Kabinger F, Stiller C, Schmitzová J, Dienemann C, Kokic G, Hillen HS, et al. Mechanism of molnupiravir-induced SARS-CoV-2 mutagenesis. *Nat. Struct. Mol. Biol.* **28** (2021) 740–746.

15. Shannon A, Selisko B, Le NTT, Huchting J, Touret F, Piorkowski G, et al. Rapid incorporation of favipiravir by the fast and permissive viral RNA polymerase complex results in SARS-CoV-2 lethal mutagenesis. *Nat. Commun.* **11** (2020) 4682.

16. Finberg RW, Ashraf M, Julg B, Ayoade F, Marathe JG, Issa NC, et al. US201 study: A phase 2, randomized Proof-of-Concept trial of favipiravir for the treatment of COVID-19. *Open Forum Infect Dis* **8** (2021) ofab563.

17. Joshi S, Parkar J, Ansari A, Vora A, Talwar D, Tiwaskar M, et al. Role of favipiravir in the treatment of COVID-19. *Int. J. Infect. Dis.* **102** (2021) 501–508.

18. [Dataset] Ison MG, Scheetz MH. Understanding the pharmacokinetics of favipiravir: Implications for treatment of influenza and COVID-19 (2021).

19. Hassanipour S, Arab-Zozani M, Amani B, Heidarzad F, Fathalipour M, Martinez-de Hoyo R. The efficacy and safety of favipiravir in treatment of COVID-19: a systematic review and meta-analysis of clinical trials. *Sci. Rep.* **11** (2021) 1–11.

20. Moeller NH, Shi K, Demir Ö, Belica C, Banerjee S, Yin L, et al. Structure and dynamics of SARS-CoV-2 proofreading exoribonuclease ExoN. *Proc. Natl. Acad. Sci. U. S. A.* **119** (2022).

21. Smith EC, Blanc H, Surdel MC, Vignuzzi M, Denison MR. Coronaviruses lacking exoribonuclease activity are susceptible to lethal mutagenesis: evidence for proofreading and potential therapeutics. *PLoS Pathog.* **9** (2013) e1003565.

22. Agostini Maria L, Pruijssers Andrea J, Chappell James D, Gribble Jennifer, Lu Xiaotao, Andres Erica L, et al. Small-Molecule antiviral β-d-N4-Hydroxycytidine inhibits a Proofreading-Intact coronavirus with a high genetic barrier to resistance. *J. Virol.* **93** (????) e01348–19.

23. Eloy P, Le Grand R, Malvy D, Guedj J. Combined treatment of molnupiravir and favipiravir against SARS-CoV-2 infection: One + zero equals two? *eBioMedicine* **74** (2021).

24. Hadj Hassine I, Ben M'hadheb M, Menéndez-Arias L. Lethal mutagenesis of RNA viruses and approved drugs with antiviral mutagenic activity. *Viruses* **14** (2022).

25. Pfeiffer JK, Kirkegaard K. A single mutation in poliovirus RNA-dependent RNA polymerase confers resistance to mutagenic nucleotide analogs via increased fidelity. *Proc. Natl. Acad. Sci. U. S. A.* **100** (2003) 7289–7294.

26. Mejer N, Galli A, Ramirez S, Fahnøe U, Benfield T, Bukh J. Ribavirin inhibition of cell-culture infectious hepatitis C genotype 1-3 viruses is strain-dependent. *Virology* **540** (2020) 132–140.

27. Szemiel AM, Merits A, Orton RJ, MacLean OA, Pinto RM, Wickenhagen A, et al. In vitro selection of remdesivir resistance suggests evolutionary predictability of SARS-CoV-2. *PLoS Pathog.* **17** (2021) e1009929.

28. Nguyen THT, Guedj J, Anglaret X, Laouénan C, Madelain V, Taburet AM, et al. Favipiravir pharmacokinetics in Ebola-Infected patients of the JIKI trial reveals concentrations lower than targeted. *PLoS Negl. Trop. Dis.* **11** (2017) e0005389.

29. Delang L, Segura Guerrero N, Tas A, Quérat G, Pastorino B, Froeyen M, et al. Mutations in the chikungunya virus non-structural proteins cause resistance to favipiravir (t-705), a broad-spectrum antiviral. *J. Antimicrob. Chemother.* **69** (2014) 2770–2784.

30. Wang Y, Li G, Yuan S, Gao Q, Lan K, Altmeyer R, et al. In vitro assessment of combinations of enterovirus inhibitors against enterovirus 71. *Antimicrob. Agents Chemother.* **60** (2016) 5357–5367.

31. de Avila AI, Moreno E, Perales C, Domingo E. Favipiravir can evoke lethal mutagenesis and extinction of foot-and-mouth disease virus. *Virus Res.* **233** (2017) 105–112.

32. Escribano-Romero E, Jiménez de Oya N, Domingo E, Saiz JC. Extinction of west nile virus by favipiravir through lethal mutagenesis. *Antimicrob. Agents Chemother.* **61** (2017).

33. Bank C, Renzette N, Liu P, Matuszewski S, Shim H, Foll M, et al. An experimental evaluation of drug-induced mutational meltdown as an antiviral treatment strategy. *Evolution* **70** (2016) 2470–2484.

34. Goldhill DH, Te Velthuis AJW, Fletcher RA, Langat P, Zambon M, Lackenby A, et al. The mechanism of resistance to favipiravir in influenza. *Proc. Natl. Acad. Sci. U. S. A.* **115** (2018) 11613–11618.

35. Goldhill DH, Yan A, Frise R, Zhou J, Shelley J, Gallego Cortés A, et al. Favipiravir-resistant influenza a virus shows potential for transmission. *PLoS Pathog.* **17** (2021) e1008937.

36. Nelson C, Otto S. Mutagenic antivirals: the evolutionary risk of low doses (2021).

37. Hurt AC, Chotpitayasunondh T, Cox NJ, Daniels R, Fry AM, Gubareva LV, et al. Antiviral resistance during the 2009 influenza a H1N1 pandemic: public health, laboratory, and clinical perspectives. *Lancet Infect. Dis.* **12** (2012) 240–248.

38. Rogers MB, Song T, Sebra R, Greenbaum BD, Hamelin ME, Fitch A, et al. Intrahost dynamics of antiviral resistance in influenza a virus reflect complex patterns of segment linkage, reassortment, and natural selection. *MBio* **6** (2015).

39. Schneider DS, Ayres JS. Two ways to survive infection: what resistance and tolerance can teach us about treating infectious diseases. *Nat. Rev. Immunol.* **8** (2008) 889–895.

40. Khan AI, Dinh DM, Schneider D, Lenski RE, Cooper TF. Negative epistasis between beneficial mutations in an evolving bacterial population. *Science* **332** (2011) 1193–1196.

41. Lansch-Justen L, Cusseddu D, Schmitz MA, Bank C. The extinction time under mutational meltdown (2022).

42. Gordo I, Charlesworth B. On the speed of muller's ratchet. *Genetics* (2000).

43. Silander OK, Tenaillon O, Chao L. Understanding the evolutionary fate of finite populations: the dynamics of mutational effects. *PLoS Biol.* **5** (2007) e94.

44. Goyal S, Balick DJ, Jerison ER, Neher RA, Shraiman BI, Desai MM. Dynamic Mutation–Selection balance as an evolutionary attractor. *Genetics* **191** (2012) 1309–1319.

45. Anciaux Y, Chevin LM, Ronce O, Martin G. Evolutionary rescue over a fitness landscape. *Genetics*
209 (2018) 265–279.

46. Notley-McRobb L, Seeto S, Ferenci T. Enrichment and elimination of muty mutators in escherichia
coli populations. *Genetics* **162** (2002) 1055–1062.

47. Ramiro RS, Durão P, Bank C, Gordo I. Low mutational load and high mutation rate variation in gut
commensal bacteria. *PLoS Biol.* **18** (2020) e3000617.

48. Orr HA. The rate of adaptation in asexuals. *Genetics* **155** (2000) 961–968.

49. Lynch M, Ackerman MS, Gout JF, Long H, Sung W, Thomas WK, et al. Genetic drift, selection and
the evolution of the mutation rate. *Nat. Rev. Genet.* **17** (2016) 704–714.

50. Sniegowski PD, Gerrish PJ, Lenski RE. Evolution of high mutation rates in experimental populations
of e. coli. *Nature* **387** (1997) 703–705.

51. Gentile CF, Yu SC, Serrano SA, Gerrish PJ, Sniegowski PD. Competition between high- and higher-
mutating strains of escherichia coli. *Biol. Lett.* **7** (2011) 422–424.

52. Gerrish PJ, Colato A, Perelson AS, Sniegowski PD. Complete genetic linkage can subvert natural
selection. *Proc. Natl. Acad. Sci. U. S. A.* **104** (2007) 6266–6271.

53. Clune J, Misevic D, Ofria C, Lenski RE, Elena SF, Sanjuán R. Natural selection fails to optimize
mutation rates for long-term adaptation on rugged fitness landscapes. *PLoS Comput. Biol.* **4** (2008)
e1000187.

54. Gerrish PJ, Colato A, Sniegowski PD. Genomic mutation rates that neutralize adaptive evolution and
natural selection. *J. R. Soc. Interface* **10** (2013) 20130329.

55. Sydykova DK, LaBar T, Adami C, Wilke CO. Moderate amounts of epistasis are not evolutionarily
stable in small populations. *J. Mol. Evol.* **88** (2020) 435–444.

56. Nowak M, Schuster P. Error thresholds of replication in finite populations mutation frequencies and the
onset of muller's ratchet. *J. Theor. Biol.* **137** (1989) 375–395.

57. Alves D, Fontanari JF. Error threshold in finite populations. *Phys. Rev. E* **57** (1998) 7008–7013.

58. Wilke CO. Quasispecies theory in the context of population genetics. *BMC Evol. Biol.* **5** (2005) 44.

59. Bagnoli F, Bezzi M. Eigen's error threshold and mutational meltdown in a quasispecies model. *Int. J.*
Mod. Phys. C **09** (1998) 999–1005.

60. LaBar T, Adami C. Evolution of drift robustness in small populations. *Nat. Commun.* **8** (2017) 1012.

61. O'Fallon BD, Adler FR, Proulx SR. Quasi-species evolution in subdivided populations favours
maximally deleterious mutations. *Proc. Biol. Sci.* **274** (2007) 3159–3164.

62. Merleau NSC, Pénisson S, Gerrish PJ, Elena SF, Smerlak M. Why are viral genomes so fragile? the
bottleneck hypothesis. *PLoS Comput. Biol.* **17** (2021) e1009128.

63. Rodríguez-Verdugo A, Tenaillon O, Gaut BS. First-Step mutations during adaptation restore the
expression of hundreds of genes. *Mol. Biol. Evol.* **33** (2016) 25–39.

64. González-González A, Hug SM, Rodríguez-Verdugo A, Patel JS, Gaut BS. Adaptive mutations in
RNA polymerase and the transcriptional terminator rho have similar effects on escherichia coli gene
expression. *Mol. Biol. Evol.* **34** (2017) 2839–2855.

65. Zabinsky RA, Mason GA, Queitsch C, Jarosz DF. It's not magic - hsp90 and its effects on genetic and
epigenetic variation. *Semin. Cell Dev. Biol.* **88** (2019) 21–35.

66. Xue KS, Bloom JD. Reconciling disparate estimates of viral genetic diversity during human influenza
infections. *Nat. Genet.* **51** (2019) 1298–1301.

67. McCrone JT, Woods RJ, Martin ET, Malosh RE, Monto AS, Lauring AS. Stochastic processes constrain
the within and between host evolution of influenza virus. *Elife* **7** (2018).

68. Foll M, Poh YP, Renzette N, Ferrer-Admetlla A, Bank C, Shim H, et al. Influenza virus drug resistance:
a time-sampled population genetics perspective. *PLoS Genet.* **10** (2014) e1004185.
69. Gessler DD. The constraints of finite size in asexual populations and the rate of the ratchet. *Genet. Res.*
66 (1995) 241–253.
70. Jiang L, Liu P, Bank C, Renzette N, Prachanronarong K, Yilmaz LS, et al. A balance between inhibitor
binding and substrate processing confers influenza drug resistance. *J. Mol. Biol.* **428** (2016) 538–553.

FIGURES

Figure 1. Deleterious selection coefficients of intermediate size minimize the extinction time. Two different carrying capacities are shown: 500 and 10 000. Each dot represents a simulation run. With larger carrying capacity $K = 10000$ the time to extinction increases. The black vertical line indicates the selection coefficient ($s_{del} = -0.023$) used in the standard parameter set throughout the rest of the paper.

Figure 2. Mutational meltdown can be prevented with sufficient continual input of beneficial mutations. The sharp survival threshold depends strongly on the carrying capacity. The population can escape mutational meltdown when the ratio of beneficial to deleterious mutations is $\approx 10\%$, when the carrying capacity is $K = 500$; a ratio of $\approx 1\%$ is sufficient for escape from mutational meltdown when $K = 10000$. Each dot represents a simulation run.

Figure 3. A mutation modifier of sufficient effect size can lead to escape from mutational meltdown. Each dot in Panels A, C, E, G represents a simulation run; simulations in which the mutation modifier was fixed in the population are shown in color, whereas simulations in which the modifier did not invade are shown in gray. Note that the effect size of the modifier, f_{mutMod}, is defined multiplicatevely: e.g., if $f_{\mathrm{mutMod}} = 0.1$, the mutation rate is reduced to 10% of its original value. B, D, F, H. In general, a mutation modifier that sufficiently decreases the mutation rate is effective at fixing in the population (yellow dots) and increasing the extinction time (black dots). Even when the modifier increases the mutation rate, it becomes fixed in some simulations, caused by random drift during the extinction phase (yellow dots for $f_{\mathrm{mutMod}} > 1$). The graph shows the proportion of simulations for which the modifier allele was fixed (yellow dots), and the proportion of simulations for which it extended the extinction time beyond the maximum observed extinction time without the modifiers (black dots). C. For $K = 500$, escape from mutational meltdown becomes more stochastic with a lower modifier mutation rate. With $U_{\mathrm{mutMod}} = 0.0001$, even for very strong modifiers, there is a small probability that escape is unsuccessful when the modifier allele appears too late. E-H. At higher carrying capacities $K = 10000$, escape from mutational meltdown is possible even with modifiers of relatively small effect, and the escape becomes more deterministic. Each parameter combination was simulated 10000 times.

Figure 4. Both positive and negative DFE modifiers can lead to escape from mutational meltdown. Each dot in Panels A, C, E, G represents a simulation run; simulations in which the mutation modifier was fixed in the population are shown in color, whereas simulations in which the modifier did not invade are shown in gray. For positive DFE modifiers (effect size $f_{\mathrm{dfeMod}} < 1$, i.e., the modifier makes deleterious mutations *less* deleterious), modifier alleles invade and fix reliably, but a large effect size $f_{\mathrm{dfeMod}} << 1$ is necessary to lead to successful escape from mutational meltdown (i.e., survival for 5000 generations). For negative DFE modifiers (effect size $f_{\mathrm{dfeMod}} > 1$, , i.e., the modifier makes deleterious mutations *more* deleterious), modifier alleles invade and fix rarely, especially when the mutation rate to the modifier is low. Interestingly, although a negative DFE modifier initially tends to have a deleterious fitness effect on the lineage it carries, its escape probability increases with the carrying capacity (F, H), unlike what is expected for a deleterious mutation. Each parameter combination was simulated 10000 times.