An α-π transition in S6 shapes the conformational cycle of the bacterial sodium channel NavAb

Koushik Choudhury[1], Rebecca J Howard[2], Lucie Delemotte[1,*]

[1]Science for Life Laboratory, Department of Applied Physics, KTH Royal Institute of Technology, Solna, SE-171 65 Sweden.

[2]Science for Life Laboratory, Department of Biochemistry and Biophysics, Stockholm University, Stockholm, SE-114 19 Sweden

Correspondence to: lucied@kth.se

Abstract

Voltage gated sodium channels play an important role in electrical signaling in excitable cells. In response to changes in membrane potential, they cycle between nonconducting and conducting conformations. With recent advances in structural biology, structures of sodium channels have been captured in several distinct conformations, thought to represent different functional states. However, it has been difficult to capture the intrinsically transient open state. We recently showed that a proposed open state of the bacterial sodium channel NavMs was not conductive, and that a conformational change involving a transition to a π helix in the pore-lining S6 helix converted this structure into a conducting state. However, the relevance of this structural feature in other sodium channels, and its implications for the broader gating cycle, remained unclear. Here, we propose a comparable open state of another class of bacterial channel from *Aliarcobacter butzleri* (NavAb), with characteristic pore hydration, ion permeation and drug binding properties. Furthermore, we show that a π-helix transition is essential for pore opening, and that such a conformational change blocks fenestrations in the inner helix bundle. We also discover that a region in the C-terminal domain can undergo a disordering transition proposed to be important for pore opening. These results support a role for a π-helix transition in the opening of NavAb, enabling new proposals for the structural annotation and drug modulation mechanisms in this important model sodium channel.

Introduction

Voltage-gated sodium ion channels (Navs) are membrane proteins that selectively transport sodium ions across cell membranes. They open in response to membrane depolarization, and, in excitable cells, are responsible for the fast rising phase of the action potential. They are further characterized by rapid inactivation, during which the pore becomes non-conductive despite continued depolarized conditions (1). Eukaryotic Nav channels are pseudotetrameric single-chain proteins consisting of around 2000 residues. Bacterial sodium channels, on the other hand, have a simpler homotetrameric architecture, where each of four identical subunits comprises around 270 residues. Because bacterial channels share important aspects of structure and function with eukaryotic Nav channels, they have been used as prototypes to understand structure-function relationships in this superfamily of channels (2).

Several structures of bacterial (3-20) and eukaryotic Nav channels (21-27) have been determined. A prototypical Nav channel consists of three types of domains - voltage sensing domain (VSD), a pore domain and an intracellular C-terminal domain (CTD, Figure 1.A). Each VSD constitutes a four-helix bundle made of S1-S4, where S4 contains several arginine residues responsible for sensing the changes in potential difference across the membrane. Each channel contains four VSDs, each contributed by one subunit, that are arranged around a central pore. In their resting state, the VSDs' S4 helices are stabilized in an 'inward' or 'down' position. Upon activation, the S4 helices transition into an 'outward' or 'up' position (7). The pore domain consists of a tetrameric inverse cone-shaped arrangement of the S5 and S6 helices from each of the four subunits. The pore is thought to open and close at the so-called intracellular bundle crossing, where an expanded pore likely represents an open state and a constricted pore likely corresponds to a closed state. In the resting state of the channel, the VSDs are in a resting conformation and the pore is non-conductive. In the open state, the VSDs are activated, and the pore is dilated. In the inactivated state, the VSDs remain activated, but the pore is non-conductive. In addition to these transmembrane domains, bacterial Nav channels possess a CTD that extends into the cytoplasm. This domain consists of two main regions - a neck region (proximal to the transmembrane region of the protein) and a distal coiled-coil region (Figure 1.A). Disordering of the neck region of the

CTD favors activation and pore opening in several bacterial sodium channels (19, 20) and is involved in channel inactivation through an unknown mechanism (28).

Structures of the sodium channel from *Aliarcobacter butzleri* (NavAb) have been described as representing three functional states (7,9,10). Based on these structures, the following functional cycle has been proposed (Figure 1B). Starting from a resting state (PDB ID - 6P6W) with VSDs 'down' and the pore in a constricted conformation, the channel transitions to a metastable pre-open state in which the VSDs are activated and the pore non conductive. This transition may occur in two steps, first to a state with a highly constricted pore (PDB ID - 5VB2) followed by a second with a slightly wider pore (PDB ID - 3RVY). After this, the pore transitions to an open state that features an expanded pore (PDB ID - 5VB8). Finally, the channel moves to an inactivated state wherein the VSDs are still activated but the pore is constricted and asymmetrical (C2 symmetry - PDB ID - 4EKW).

Although this proposed cycle is consistent with some functional predictions, most structures of NavAb are of constructs containing one or more mutations, and it is thus unclear how well they represent functional states of wild-type (WT) NavAb. The open state has been especially difficult to capture, given the propensity of these channels to rapidly inactivate. Indeed, the structure of NavAb that is most likely to represent an open state, determined by x-ray crystallography, contains the pore-domain mutation I217C and is truncated after the S6 helix, thus deleting the CTD (10). As molecular dynamics simulations indicated that the pore was hydrated and conducted sodium ions, this structure was assigned as an open state (10). However, functional properties of this structure upon reversion to the WT sequence remained unclear.

In our recent study of the related bacterial channel NavMs (29), we found that a structure initially thought to represent an open state, with an expanded pore with entirely α-helical S6 helices, was dehydrated and non-conductive to sodium ions. However, when the S6 helices transitioned to containing a short π-helical stretch resulting in a rotation of the S6 C-terminus, the pore was hydrated, allowing sodium ion permeation. Since NavMs and NavAb are similar in structure and sequence, we hypothesized that the open state of NavAb may also require a transition to a π-helix in S6.

Local anesthetic drugs act by binding to sodium channels (30), several of them in a state-dependent manner. Bacterial sodium channels are known to be modulated by these drugs in a similar manner as their eukaryotic counterparts, making them useful tools to study state-dependent drug binding (2, 31). Depending on their protonation states, such drugs can access the Nav central cavity either via a hydrophobic pathway involving partitioning of the drugs in the membrane, or via a hydrophilic pathway across the intracellular ion gate (32). Indeed, structures have revealed the presence of fenestrations that connected the Nav central cavity to the membrane (9). These fenestrations were confirmed to play an important role in tonic block, whereby drugs access the central cavity when the pore is closed (5). The generation of a new open model poses the opportunity to study access of local anesthetics to their binding site in various states of NavAb.

In this study, we aimed to revise the proposed conformational cycle of NavAb, and to describe detailed features of the pore domain in resting, open and inactivated states of the channel. To do so, we first used molecular dynamics (MD) simulations to describe the full length structure of WT NavAb in a presumed open state. Reverting the I1217C mutation dehydrated the pore, leading to a model that was impermeable to sodium ions. Given our previous model of NavMs pore opening (29), we then constructed an alternate model of the NavAb pore, in which introducing a π-helix in S6 results in a hydrated and conductive pore. This open-pore model was validated by sodium ion and local-anesthetic permeation free energy calculations. We then used the open model to characterize state-dependent properties of hydrophobic fenestrations and lidocaine binding. Finally, we posited a new opening mechanism for NavAb, in which the formation of a π-helix in S6 and disordering of the neck region of the CTD are prerequisites for pore expansion.

MATERIALS AND METHODS

Model building

Four different NavAb structures were used in this study: a resting state (PDB ID 6P6W), in which the VSDs are at rest, the pore is contracted, and the CTD is truncated; a tentatively assigned open state (PDB ID 5VB8), in which the VSDs are activated, the pore is expanded, and the CTD is again truncated; and two decoupled states, tentatively assigned as pre-open (PDB ID 5VB2) and inactivated (PDB ID 4EKW) in which the VSDs are activated and the pore is contracted. In 5VB2, the pore domain has C4-symmetry and the CTD is ordered, while in 4EKW, the pore domain has only C2-symmetry (partly asymmetric) and the CTD is truncated. All the mutations in the structures were reverted to WT using CHARMM-GUI (33). Incomplete loops were also modeled using CHARMM-GUI (33).

Alternative models of NavAb with contracted or expanded pores were built by introducing a π-helix in a region of the S6 helix of 5VB2 and 5VB8, respectively. This was done using MODELLER 9.22 (34) following a protocol implemented in previous work on NavMs (29). Briefly, this consisted of shifting the structural alignment in S6 of the π- versus template α-models in order to introduce an unpaired backbone carbonyl. Specifically, we introduced one gap in the template (α) sequence prior to T1206, and a second in the target (π) sequence following D1219, thus shifting the region one position upstream from T1206 to the inner end of S6, and resulting in the formation of a π helix around T1206.

For pore-opening free energy calculations, we also built contracted-α and contracted-π NavAb models from PDB ID 5VB2 with the CTD deleted, otherwise following the procedure described above. This approach ensured that the protein length was consistent with the expanded-pore models.

System preparation

Each model was embedded in a homogenous lipid bilayer consisting of 400 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (POPC) using the CHARMM-GUI Membrane builder (33). Lidocaine-protein complexes were prepared by randomly placing the drug in the central cavity of the different models, respectively. Each system was hydrated by adding ~45 Å layers of water to

either side of the membrane. Lastly, the systems were neutralized with 150 mM NaCl. The Charmm36 force field was used to describe interactions between protein (35), lipids (36) and ions. Non-bonded fixes (NBFixes) were considered in the description of interactions of sodium ions with carboxyl and carbonyl groups (37). The TIP3P model was used to describe all water molecules (38). Parameters for lidocaine were taken from our previous study (29).

Simulation details

Each system was minimized for 5000 steps using steepest descent and equilibrated with constant number of particles, pressure and temperature (NPT) for 30 ns, during which position restraints on the protein were gradually released. For the expanded-pore states (with or without lidocaine) simulations, equilibration was done following the protocol reported in (10). Initially the heavy atoms of protein were restrained with a force constant of 1000 kj/mol/nm^2 for 10 ns. Then the protein backbone was restrained with a force constant of 1000 kj/mol/nm^2 for 10 ns. Finally, the protein Cα atoms were restrained with a force constant of 1000 kj/mol/nm^2 for 10 ns. For the other systems, equilibration was carried out in 6 steps according to the default CHARMM-GUI protocol (33). During equilibration, a time step of 2 fs was used, pressure was maintained at 1 bar through Berendsen pressure coupling, temperature was maintained at 300 K through Berendsen temperature coupling (39) with the protein, membrane and solvent coupled and the LINCs algorithm (40) was used to constrain the bonds involving hydrogen atoms. For long range interactions, periodic boundary conditions and particle mesh Ewald (PME) were used (41). For short range interactions, a cut-off of 12 Å was used. Production simulations were run with/without restraints for different systems as shown in Table 1, using Parinello-Rahman pressure coupling (42) and Nose- Hoover temperature coupling (43). Simulations were performed using GROMACS 2020.3 (44,45). Some simulations were performed under a transmembrane voltage of +/- 750 mV, via application of an external electric field. The transmembrane voltage is calculated as the product of electric field and the simulation box length along which the field is applied.

Table 1: Summary table of equilibrium simulations performed. Restrained production simulations (where indicated) were used to prevent pore collapse in the absence of CTD.

System	Equilibration	Production
NavAb expanded α-model (5VB8)	30 ns	185 ns (unrestrained)

NavAb expanded α-model I217C (5VB8)	30 ns	250 ns (C-α restrained)
NavAb expanded α-model (5VB8)	30 ns	200 ns (C-α restrained)
NavAb expanded α-model (+750mV)	30 ns	200 ns (C-α restrained)
NavAb expanded α-model (-750 mV)	30 ns	200 ns (C-α restrained)
NavAb expanded π-model (5VB8)	30 ns	200 ns (C-α restrained)
NavAb expanded α-model + lidocaine	30 ns	200 ns (C-α restrained)
NavAb asymmetric α model + lidocaine	30 ns	200 ns (unrestrained)
NavAb resting α model + lidocaine	30 ns	200 ns (unrestrained)
NavAb expanded π-model + lidocaine	30 ns	200 ns (C-α restrained)
NavAb expanded π-model N1211L	30 ns	200 ns (C-α restrained)
NavAb contracted α-model (5VB2) - 300K	2 ns	60 ns (unrestrained)
NavAb contracted α-model (5VB2) - 320K	2 ns	200 ns (unrestrained)
NavAb contracted α-model (5VB2) - 360K	2 ns	200 ns (unrestrained)
NavAb contracted α-model (5VB2) - 400K	2 ns	200 ns (unrestrained)
NavAb contracted π-model (5VB2)	2 ns	20 ns (unrestrained)

Ion permeation and drug permeation free energy calculations

The free energy profiles of sodium ions and lidocaine permeation along the pore axis were calculated using the accelerated weight histogram (AWH) method (46). In brief, for each equilibrated structure, we applied one independent AWH bias and simulated for 200 ns each with 6 walkers sharing bias data and contributing to the same target distribution. The target distribution chosen was a flat (uniform) distribution. Each bias acts on the z-axis defined using the center-of-mass z-distance between one central sodium ion/lidocaine (all atoms) and all atoms of S1178 residues in NavAb, with a sampling interval corresponding to the entire box length. The rate of

change of each AWH bias was initialized by setting the initial average free energy error to 20 kj/mol and the diffusion constant was set to 0.00005 nm^2/ps. To keep the solute close to the pore entrance, the coordinate radial distance was restrained to stay below 10 Å by adding a flat-bottom umbrella potential. All the Cα atoms of the protein were restrained by imposing harmonic potentials with force constants of 1000 kJ mol^{-1} nm^{-2} to remain as close as possible to the experimental structures.The average potential of mean force (PMF) profile and the error bars were calculated from a single walker since the walkers communicate with each other. For all systems except the expanded π-model with lidocaine, the average PMF profile was calculated by taking the data from the last 100 ns in intervals of 10 ns and the error bars were calculated by calculating the standard deviation. For the expanded π-model with lidocaine, data from the last 50 ns at 10 ns intervals were chosen for calculating the average and standard deviation.

Pore opening free energy calculation

The pore opening free energy calculation was performed using multiple-walker well-tempered metadynamics (49,50) in GROMACS 2020.5 patched with PLUMED 2.7.3 (47, 48). This approach enabled a linear combination of several collective variables, a feature not available in GROMACS at the time we started this work. These simulations were initiated from the equilibrated structure of the constricted α model and constricted π model in absence of CTD. The average of the distance between Cα atoms of I1216/I1217 and I1119 (for the α model and the π model, respectively) across the four subunits was used as the collective variable for the free energy calculations. The hills height and width was set at 1.2 kJ/mol and 0.05 nm respectively. Upper and lower restraints (as shown in the corresponding x-axes of the free energy plots) were applied to restrict the sampling space. The simulations were run for 200 ns per walker, for a total of 6 walkers. The free energy profile was estimated using the sum_hills tool in PLUMED. The average free energy profile was calculated by taking the data from the last 100 ns in intervals of 10 ns and the error bars were estimated by calculating the standard deviation.

Table 2: Summary of the enhanced sampling free energy calculation performed.

System	Enhanced Sampling technique	Production
NavAb (expanded α-model) - Na$^+$	AWH	200 ns × 6 walkers
NavAb (expanded π-model) - Na$^+$	AWH	200 ns × 6 walkers
NavAb (expanded α-model) - lidocaine	AWH	200 ns × 6 walkers
NavAb (resting α model) - lidocaine	AWH	200 ns × 6 walkers
NavAb (asymmetric α model) - lidocaine	AWH	200 ns × 6 walkers
NavAb (expanded π-model) - lidocaine	AWH	200 ns × 6 walkers
NavAb expanded π-model (N214L) - Na$^+$	AWH	139 ns × 6 walkers
Pore opening (α model without CTD)	Well-tempered metadynamics	195 ns × 6 walkers

Pore opening (π model without CTD)	Well-tempered metadynamics	210 ns × 6 walkers

Analysis

The water number density, water hydration free energy and pore radius were calculated using the channel annotation package (CHAP) (51). For the water density analysis, frames were extracted every 100 ps from the trajectories of the respective models. For the fenestration radius analysis, 200-600 equally spaced frames were extracted from the trajectories of the expanded/contracted α and π models.

Representative binding poses of lidocaine were obtained by performing clustering on frames extracted from the AWH calculations corresponding to free energy minima. Clustering was done using gmx cluster, with a cutoff of 0.1 nm. The center of the most populated cluster was then chosen as the representative binding pose. A similar approach was used to extract the most representative structures from the pore opening free energy profiles. Secondary structure was characterized using the do_dssp module in GROMACS.

The files necessary to reproduce the data herein can be found at https://osf.io/u8fg9/

RESULTS

The expanded-α structure of NavAb is non-conductive

To test the relevance to WT channel function of the proposed open-state structure of NavAb, in which the VSDs were captured in an activated state and the pore in an expanded conformation (PDB - 5VB8), we first reverted the I1217C mutations and carried out equilibrium MD simulations, still in absence of the CTD. Because of the α-helical structure of S6, and wide diameter of the pore, we call this model "expanded α". In previous unrestrained simulations of this model, the pore spontaneously collapsed to a dehydrated, nonconducting conformation (10,52). We therefore restrained the C-α positions to maintain the pore in an expanded state. Analyzing the pore hydration by calculating the time-averaged water number density along the central axis showed that the pore is nonetheless dehydrated close to the activation gate (-1 nm along the pore axis, Figure 2A, S1A), likely due to hydrophobic residues (V1213) lining the pore (Figure 2C), and in contrast to the I1217C mutant (Figure S1D).

We next hypothesized that a transmembrane potential difference could affect pore hydration. Previous studies have indeed shown that applying external electric fields can promote hydration of hydrophobic pores (53). For the expanded-α model of NavAb, an external transmembrane voltage of -750 mV did not change pore hydration substantially. Applying a transmembrane voltage of +750mV increased hydration, but with a final water density still lower than that of bulk water (Figure S2). Moreover, the pore was only transiently hydrated, alternating between wetted and dewetted states (Figure S1B,C).

Introducing a π-helix in S6 results in a conductive channel

Inspired by our previous work on NavMs (29), we next hypothesized that a π-helix in S6 might stabilize a conducting state of NavAb. Introducing this modification (see Methods), we produced an "expanded π" model which we interrogated using MD simulations. Simulating this model resulted in a consistently hydrated pore (Figure 2B), in contrast to the expanded-α model (Figure S2). To further confirm that the expanded-π model represents a functional open state, we calculated sodium ion and lidocaine permeation free energy profiles across the channel gate using enhanced sampling. There was a substantial barrier to sodium (~ 40 kT) and lidocaine (~ 20 kT) exit from

the central cavity in the expanded-α model, while there was no barrier in the expanded-π model (Figure 2 D,E, Figure S3-4,S6-7).

The introduction of this π-helix resulted in the rotation of a conserved asparagine, N1211, towards the NavAb pore lumen (Figure 2B). Mutating this pore-facing asparagine to a hydrophobic residue (N1211L) did not affect pore hydration (Figure S2, S5), but increased the barrier to sodium-ion permeation at the site of mutation (Figure 2D). Taken together, these results are consistent with the expanded-π model representing a functional open state, and with N1211 contributing to sodium-ion permeation.

Local anesthetic binding poses and access pathways vary with functional state
We hypothesized that a structural transition in S6 may impact the accessibility of neutral drugs to their binding site in the pore cavity through the pore-lining fenestrations. To test this, we calculated the time-averaged radius of two such fenestrations each in the contracted-α, contracted-π, expanded-α and expanded-π models of NavAb. Comparing the fenestration radii for these models, we found that the π models have narrower fenestrations than the α models (Figure 3A,B). Indeed, the fenestrations are obstructed by large hydrophobic residues F1207 and I1210 in both the contracted- and expanded-π models (Figure 3B). This observation indicates that hydrophobic drugs can access the central cavity only in the context of a fully α-helical S6, irrespective of pore diameter.

We next investigated state-dependent drug access to the central cavity via the hydrophilic pathway by calculating lidocaine permeation free energies across the channel gate in different functional states of NavAb. We particularly compared our presumed-open expanded-π model to two previously reported structures, one with resting VSDs and a symmetrically contracted pore (assigned resting, PDB ID 6P6W), and one with activated VSDs and an asymmetric conformation of the pore (assigned inactivated, PDB ID 4EKW). Unlike the expanded-π model, the resting and inactivated models featured high barriers to drug access from the intracellular side (Figure 3C, Figure S7-9). In all three models, free-energy profiles indicated that the drug binds in the central cavity close to the selectivity filter, forming contacts with T1206 in both the resting and the expanded-π models. However, in the resting model, lidocaine was oriented parallel to the pore

axis with its charged group pointing towards the selectivity filter, while in the expanded-π model it was oriented parallel to the membrane plane (Figure 3D). This difference could be due to the presence of in the expanded-π model of the hydrophilic sidechain of N1211, which accommodates the charged group of lidocaine. In the asymmetric inactivated model, the drug was oriented perpendicular to the membrane plane and shifted further towards the intracellular end of the pore, closer to V1213 (Figure 3D). These results indicate that there is a change in the binding pose of lidocaine as the channel transitions between different functional states, and that the drug can access its binding site via the intracellular gate exclusively in the π model.

Neck region of the C-terminal domain can undergo disordering

Experimental structures and electrophysiology recordings have indicated a disordered segment in the neck region of the CTD in certain states of several bacterial sodium channels (20). This conformational change is proposed to be important for pore opening. To test whether this region can undergo disordering in NavAb, we subjected the full-length constricted-α structure of NavAb (PDB ID - 5VB2) to unrestrained simulations at elevated temperatures (300K, 340K, 360K and 400K). Simulations at 320K and 360K showed limited disordering in the CTD (figure S10A,B) while those at 400K showed a complete disordering of the neck region (Figure 4A,B, Figure S10C) spanning residues I1226 to N1243 (figure 4B). Despite a loss of symmetry, secondary structure was overall preserved in the rest of the protein at 400K (Figure S10C), suggesting that the CTD neck is specifically prone to losing its secondary structure.

A π-helix transition is essential for pore opening

The results above suggested that the NavAb open state features a π-helix in S6, along with an expanded pore. In contrast, the closed experimental structure showed S6 to be contracted and entirely α-helical. Pore opening therefore involves an α- to π-helix transition in S6 on top of pore expansion. To determine whether this secondary structure change occurs before or after expansion of the pore, we calculated the free energy for pore opening in presence of an α- and a π-helix. The average of the distance between Cα atoms of I1216 (α model)/I1217 (π model) and I1119 across the four subunits was used as the collective variable in well-tempered metadynamics simulations (Figure 5A,B), enabling sampling of expanded-pore conformations from simulations initiated from contracted pores (Figure S11). For the α model, we observed two metastable contracted-pore

conformations, while the expanded pore was unstable (Figure 5C, S12). For the π model, on the other hand, conformations corresponding to expanded pores were energetically accessible (Figure 5D, S13). These results indicate that a π helix in S6 is necessary for pore expansion, and that the transition between resting and open should include transitioning via a pre-open state in which the pore is contracted but a region of S6 is π-helical.

In the contracted-α model, N1211 is pointing away from the pore into the peripheral cavity, attracting water from the cytoplasm into peripheral cavities (Figure S14). These water-filled peripheral cavities, as well as the attractive interaction between hydrophobic pore-facing residues at the activation gate could impede pore opening. In the contracted-π model, on the other hand, N1211 faces the pore lumen rather than the peripheral cavities, which are hence no longer hydrated (Figure S14). These observations could constitute a rationale for the low propensity of the α model to expand.

Discussion

Our first observation in this work was that the expanded-α experimental structure of NavAb does not convincingly model a functional open state. Indeed, in simulations of this model, the pore was dehydrated, preventing sodium ion permeation across the gate and lidocaine access to its binding site in the central cavity. However, introducing a short π-helix in S6 produced a consistently hydrated pore which allowed sodium/lidocaine permeation. One contributor to this phenomenon is likely the conserved asparagine (N1211), where mutation to leucine increased the free-energy barrier to ion permeation.

What is then the mechanism of transition between the resting/closed and the activated/open states? We found that the CTD of NavAb has a region proximal to the membrane that can undergo a disordering transition. As similar disordering is thought to facilitate pore opening in other bacterial channels, a similar mechanism may contribute to opening in NavAb. Our pore-opening calculations further showed that it is energetically unfavorable to expand a constricted pore with an α-helical S6. Thus, channel activation should include a transition from α to π-helix in S6, via a state comparable to our contracted-π model. Indeed, the eukaryotic channel NavPas was recently

captured in a conformation consistent with such a pre-open functional assignment, with activated VSDs, a contracted pore, π helices in all of its S6 segments, and a dissociated inactivation particle (54,55). The symmetric expanded-α model of NavAb, which was initially assigned as open (10), may instead represent an inactivated state of the WT channel, possibly contributing to a multi-step inactivation process along with the asymmetric contracted-α structure (9). Resolving this complex inactivation mechanism thus remains an area of active research, since the order in which π to α transitions, pore contraction, and possible changes in symmetry occur is currently unknown.

It has been observed in TRPV1 channels that a conformation with a hydrated pore usually features dehydrated peripheral cavities between the S6 and S4-S5 helices, whereas conformations with a dehydrated pore have hydrated peripheral cavities (56). In the NavAb contracted-α model, N1211 attracts water into the peripheral cavities, possibly impeding pore opening. In the contracted-π model on the other hand, N1211 faces the pore lumen and stabilizing hydrophobic interactions at the gate are no longer present. This remodeling could favor opening of the hydrophobic activation gate, as witnessed in the pore-opening free energy calculation of the π model.

Putting together these results, we propose a modified mechanistic picture for the functional cycle of bacterial sodium channels (Figure 6). Five important processes occur during the transition between the resting and open state - the VSDs activate, an α to π transition occurs in S6 resulting in rotation of the inner half of the helix, peripheral cavities dehydrate, the CTD disorders and the pore expands. Activation of the VSDs in response to membrane depolarization is presumed to be the initiating step in this cycle. Then, we propose that the α to π transition occurs, resulting in dehydration of peripheral cavities, finally followed by pore expansion. We also propose that the CTD disorders during the opening transition, though our data do not definitively indicate at what stage this occurs. We hypothesize that the energy to overcome barriers associated with π-helix formation, CTD disordering and pore opening, is contributed at least in part by coupling to VSD activation, which is favorable thanks to membrane depolarization.

An interesting further consequence of π-helix formation in S6 was observed at the so-called fenestrations, which are thought to provide pathways for hydrophobic drugs such as lidocaine or benzocaine to access binding sites in Nav-channel pores (5). More precisely, hydrophobic drugs

have been proposed to freely access the pore in closed states, but not in open ones (5). In NavAb we found that the fenestrations are closed in π models irrespective of the dimensions (constricted/expanded) of the pore, while they are open in α models. This finding is in agreement with a functional assignment of the eukaryotic channels NavPas and NALCN as open (54,55,57), in which all S6 segments contain π helices and the fenestrations are obstructed. Moreover, recent structures of human Nav1.7 indicated that the presence of a π helix in subunit IV closed the fenestration at the interface with subunit I (58).

Lidocaine binding to different functional states of NavAb revealed that the drug binds near the selectivity filter in resting and π models, in agreement with previous simulations in NavMs and NavPas, respectively (5, 59). Lidocaine binds nearer the intracellular gate in the expanded-α, presumed inactivated model. In the alternative, asymmetric inactivated model, lidocaine binds even further towards the intracellular space, forming contacts with V1213 in addition to T1206. Lidocaine interacts in all states with T1206, suggesting that this residue serves as an important contact for local anesthetics, in agreement with studies on NavAb (5) and NachBac (31) and on eukaryotic sodium channels (60, 61).

Finally, we considered what aspects of the proposed mechanism for bacterial-Nav gating and inactivation may be applicable to eukaryotic channels. Indeed, we have now reported a transition of a region of S6 to a π helix as a factor in opening of two bacterial channels, NavMs and NavAb, raising the possibility that this is a conserved mechanism across evolution. Eukaryotic sodium channels also possess a conserved asparagine residue (N1211) in their S6 helices, and structures have revealed the presence of π-helices in certain states of some isoforms. Most human sodium channels feature a π-helix in domains I and II, while the other two S6 helices (domains III and IV) are α-helical (21,22,24-27). Recently, refined analysis of Nav1.7 densities showed a transient π helix in domain IV (58), indicating that this transition is indeed energetically possible. This conformational change has been linked to inactivation, posing provocative possibilities for future study of helical transitions in the gating cycle of both bacterial and eukaryotic channels.

Acknowledgements

We acknowledge SciLifeLab and the Swedish Research Council to LD (VR 2018-04905) for funding. The MD simulations were performed on resources provided by the Swedish National Infrastructure for Computing (SNIC) at PDC Center for High Performance Computing (PDC-HPC) and the resources provided by Swiss national supercomputing center.

Author contributions

K.C and L.D designed the research. K.C performed the simulations and analyzed the results. K.C, R.J.H and L.D interpreted the data and wrote the manuscript.

Competing interests

The authors declare to have no competing interests.

References

1. Ahern, C.A., J. Payandeh, F. Bosmans, and B. Chanda. 2016. The hitchhiker's guide to the voltage-gated sodium channel galaxy. *The Journal of General Physiology*. 147:1–24.
2. Payandeh, J., and D.L. Minor. 2015. Bacterial Voltage-Gated Sodium Channels (Bac NaVs) from the Soil, Sea, and Salt Lakes Enlighten Molecular Mechanisms of Electrical Signaling and Pharmacology in the Brain and Heart. *Journal of Molecular Biology*. 427:3–30.
3. Irie, K., Y. Haga, T. Shimomura, and Y. Fujiyoshi. 2018. Optimized expression and purification of NavAb provide the structural insight into the voltage dependence. *FEBS Letters*. 592:274–283.
4. Payandeh, J., T. Scheuer, N. Zheng, and W.A. Catterall. 2011. The crystal structure of a voltage-gated sodium channel. *Nature*. 475:353–358.
5. El-Din, T.M.G., M.J. Lenaeus, N. Zheng, and W.A. Catterall. 2018. Fenestrations control resting-state block of a voltage-gated sodium channel. *PNAS*. 115:13111–13116.
6. Jiang, D., T.M. Gamal El-Din, C. Ing, P. Lu, R. Pomès, N. Zheng, and W.A. Catterall. 2018. Structural basis for gating pore current in periodic paralysis. *Nature*. 557:590–594.
7. Wisedchaisri, G., L. Tonggu, E. McCord, T.M.G. El-Din, L. Wang, N. Zheng, and W.A. Catterall. 2019. Resting-State Structure and Gating Mechanism of a Voltage-Gated Sodium Channel. *Cell*. 178:993-1003.
8. McCusker, E.C., C. Bagnéris, C.E. Naylor, A.R. Cole, N. D'Avanzo, C.G. Nichols, and B.A. Wallace. 2012. Structure of a bacterial voltage-gated sodium channel pore reveals mechanisms of opening and closing. *Nature Communications*. 3:1102.
9. Payandeh, J., T.M.G. El-Din, T. Scheuer, N. Zheng, and W.A. Catterall. 2012. Crystal structure of a voltage-gated sodium channel in two potentially inactivated states. *Nature*. 486:135–139.

10. Lenaeus, M.J., T.M.G. El-Din, C. Ing, K. Ramanadane, R. Pomès, N. Zheng, and W.A. Catterall. 2017. Structures of closed and open states of a voltage-gated sodium channel. *PNAS*. 114:E3051–E3060.

11. Tsai, C.-J., K. Tani, K. Irie, Y. Hiroaki, T. Shimomura, D.G. McMillan, G.M. Cook, G.F.X. Schertler, Y. Fujiyoshi, and X.-D. Li. 2013. Two Alternative Conformations of a Voltage-Gated Sodium Channel. *Journal of Molecular Biology*. 425:4074–4088.

12. Zhang, X., W. Ren, P. DeCaen, C. Yan, X. Tao, L. Tang, J. Wang, K. Hasegawa, T. Kumasaka, J. He, J. Wang, D.E. Clapham, and N. Yan. 2012. Crystal structure of an orthologue of the NaChBac voltage-gated sodium channel. *Nature*. 486:130–134.

13. Bagnéris, C., P.G. DeCaen, C.E. Naylor, D.C. Pryde, I. Nobeli, D.E. Clapham, and B.A. Wallace. 2014. Prokaryotic NavMs channel as a structural and functional model for eukaryotic sodium channel antagonism. *PNAS*. 111:8428–8433.

14. Bagnéris, C., P.G. DeCaen, B.A. Hall, C.E. Naylor, D.E. Clapham, C.W.M. Kay, and B.A. Wallace. 2013. Role of the C-terminal domain in the structure and function of tetrameric sodium channels. *Nature Communications*. 4:2465.

15. Sula, A., D. Hollingworth, L.C.T. Ng, M. Larmore, P.G. DeCaen, and B.A. Wallace. 2021. A tamoxifen receptor within a voltage-gated sodium channel. *Molecular Cell*. 81:1160-1169.

16. Naylor, C.E., C. Bagnéris, P.G. DeCaen, A. Sula, A. Scaglione, D.E. Clapham, and B.A. Wallace. 2016. Molecular basis of ion permeability in a voltage-gated sodium channel. *The EMBO Journal*. 35:820–830.

17. Sait, L.G., A. Sula, M.-R. Ghovanloo, D. Hollingworth, P.C. Ruben, and B. Wallace. 2020. Cannabidiol interactions with voltage-gated sodium channels. *eLife*. 9:e58593.

18. Sula, A., J. Booker, L.C.T. Ng, C.E. Naylor, P.G. DeCaen, and B.A. Wallace. 2017. The complete structure of an activated open sodium channel. *Nat Commun*. 8:14205.

19. Shaya, D., F. Findeisen, F. Abderemane-Ali, C. Arrigoni, S. Wong, S.R. Nurva, G. Loussouarn, and D.L. Minor. 2014. Structure of a Prokaryotic Sodium Channel Pore Reveals Essential Gating Elements and an Outer Ion Binding Site Common to Eukaryotic Channels. *Journal of Molecular Biology*. 426:467–483.

20. Arrigoni, C., A. Rohaim, D. Shaya, F. Findeisen, R.A. Stein, S.R. Nurva, S. Mishra, H.S. Mchaourab, and D.L. Minor. 2016. Unfolding of a Temperature-Sensitive Domain Controls Voltage-Gated Channel Activation. *Cell*. 164:922–936.

21. Jiang, D., H. Shi, L. Tonggu, T.M.G. El-Din, M.J. Lenaeus, Y. Zhao, C. Yoshioka, N. Zheng, and W.A. Catterall. 2020. Structure of the Cardiac Sodium Channel. *Cell*. 180:122-134.

22. Jiang, D., L. Tonggu, T.M. Gamal El-Din, R. Banh, R. Pomès, N. Zheng, and W.A. Catterall. 2021. Structural basis for voltage-sensor trapping of the cardiac sodium channel by a deathstalker scorpion toxin. *Nat Commun*. 12:128.

23. Shen, H., Z. Li, Y. Jiang, X. Pan, J. Wu, B. Cristofori-Armstrong, J.J. Smith, Y.K.Y. Chin, J. Lei, Q. Zhou, G.F. King, and N. Yan. 2018. Structural basis for the modulation of voltage-gated sodium channels by animal toxins. *Science*. 362:eaau2596.

24. Li, Z., X. Jin, T. Wu, G. Huang, K. Wu, J. Lei, X. Pan, and N. Yan. 2021. Structural Basis for Pore Blockade of the Human Cardiac Sodium Channel Nav1.5 by the Anti arrhythmic Drug Quinidine**. *Angewandte Chemie International Edition*. 60:11474–11480.

25. Pan, X., Z. Li, Q. Zhou, H. Shen, K. Wu, X. Huang, J. Chen, J. Zhang, X. Zhu, J. Lei,

W. Xiong, H. Gong, B. Xiao, and N. Yan. 2018. Structure of the human voltage-gated sodium channel Nav1.4 in complex with β1. *Science*. 362:eaau2486.

26. Shen, H., D. Liu, K. Wu, J. Lei, and N. Yan. 2019. Structures of human Nav1.7 channel in complex with auxiliary subunits and animal toxins. *Science*. 363:1303–1308.

27. Jiang, D., Banh, R., El-Din, T. M. G., Tonggu, L., Lenaeus, M. J., Pomès, R., … and Catterall, W. A. 2021. Open-state structure and pore gating mechanism of the cardiac sodium channel. *Cell*, *184*(20), 5151-5162.

28. Gamal El-Din, T. M., Lenaeus, M. J., Ramanadane, K., Zheng, N., and Catterall, W. A. 2019. Molecular dissection of multiphase inactivation of the bacterial sodium channel NaVAb. *Journal of General Physiology*, *151*(2), 174-185.

29. Choudhury, K., Kasimova, M. A., McComas, S., Howard, R. J., & Delemotte, L. 2022. An open state of a voltage-gated sodium channel involving a π-helix and conserved pore-facing Asparagine. *Biophysical Journal*, *121*(1), 11-22.

30. Fozzard, H. A., Sheets, M. F., and Hanck, D. A. 2011. The sodium channel as a target for local anesthetic drugs. *Frontiers in pharmacology*, *2*, 68.

31. Lee, S., Goodchild, S. J., and Ahern, C. A. 2012. Local anesthetic inhibition of a bacterial sodium channel. *Journal of General Physiology*, *139*(6), 507-516.

32. Hille, B. 1977. Local anesthetics: hydrophilic and hydrophobic pathways for the drug-receptor reaction. *The Journal of general physiology*, *69*(4), 497-515.

33. Lee, J., D.S. Patel, J. Ståhle, S.-J. Park, N.R. Kern, S. Kim, J. Lee, X. Cheng, M.A. Valvano, O. Holst, Y.A. Knirel, Y. Qi, S. Jo, J.B. Klauda, G. Widmalm, and W. Im. 2019. CHARMM-GUI Membrane Builder for Complex Biological Membrane Simulations with Glycolipids and Lipoglycans. *J. Chem. Theory Comput.* 15:775–786.

34. Webb, B., and A. Sali. 2016. Comparative Protein Structure Modeling Using MODELLER. *Current Protocols in Bioinformatics*. 54:5.6.1-5.6.37.

35. Huang, J., and A.D. MacKerell. 2013. CHARMM36 all-atom additive protein force field: Validation based on comparison to NMR data. *Journal of Computational Chemistry*. 34:2135–2145.

36. Klauda, J.B., R.M. Venable, J.A. Freites, J.W. O'Connor, D.J. Tobias, C. Mondragon Ramirez, I. Vorobyov, A.D. MacKerell, and R.W. Pastor. 2010. Update of the CHARMM All-Atom Additive Force Field for Lipids: Validation on Six Lipid Types. *J. Phys. Chem. B*. 114:7830–7843.

37. Luo, Y., and Roux, B. 2010. Simulation of osmotic pressure in concentrated aqueous salt solutions. *The journal of physical chemistry letters*, *1*(1), 183-189.

38. Jorgensen, W.L., J. Chandrasekhar, J.D. Madura, R.W. Impey, and M.L. Klein. 1983. Comparison of simple potential functions for simulating liquid water. *The Journal of Chemical Physics* 79:926–935.

39. Berendsen, H.J.C., J.P.M. Postma, W.F. van Gunsteren, A. DiNola, and J.R. Haak. 1984. Molecular dynamics with coupling to an external bath. *The Journal of Chemical Physics* 81:3684–3690.

40. Hess, B., H. Bekker, H.J.C. Berendsen, and J.G.E.M. Fraaije. 1997. LINCS: A linear constraint solver for molecular simulations. *J. Comput. Chem.* 18:1463–1472.

41. Darden, T., D. York, and L. Pedersen. 1993. Particle Mesh Ewald - an Nlog(N) method for ewald sums in large systems. *The Journal of Chemical Physics* 98:10089–10092.

42. Parrinello, M., and A. Rahman. 1981. Polymorphic transitions in single crystals: A new molecular dynamics method. *Journal of Applied Physics*. 52:7182–7190.

43. Hoover, W.G. 1985. Canonical dynamics: Equilibrium phase-space distributions. *Physical review A*. 31:1695.

44. Abraham, M.J., T. Murtola, R. Schulz, S. Páll, J.C. Smith, B. Hess, and E. Lindahl. 2015. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX*. 1–2:19–25.

45. Lindahl, Abraham, Hess, and van der Spoel. 2021. GROMACS 2021.2 Manual. .

46. Lindahl, V., J. Lidmar, and B. Hess. 2014. Accelerated weight histogram method for exploring free energy landscapes. *The Journal of Chemical Physics*. 141:044110.

47. Bonomi, M., Branduardi, D., Bussi, G., Camilloni, C., Provasi, D., Raiteri, P., Donadio, D., Marinelli, F., Pietrucci, F., Broglia, R.A. and Parrinello, M., 2009. PLUMED: A portable plugin for free-energy calculations with molecular dynamics. *Computer Physics Communications*, *180*(10), 1961-1972.

48. Tribello, G.A., Bonomi, M., Branduardi, D., Camilloni, C. and Bussi, G., 2014. "PLUMED 2: New feathers for an old bird." *Computer Physics Communications* 185.2: 604-613.

49. Laio, Alessandro, and Michele Parrinello. "Escaping free-energy minima." *Proceedings of the National Academy of Sciences* 99.20 (2002): 12562-12566.

50. Barducci, A., Bussi, G. and Parrinello, M., 2008. "Well-tempered metadynamics: a smoothly converging and tunable free-energy method." *Physical review letters* 100.2 (2008): 020603

51. Klesse, G., S. Rao, M.S.P. Sansom, and S.J. Tucker. 2019. CHAP: A Versatile Tool for the Structural and Functional Annotation of Ion Channel Pores. *Journal of Molecular Biology*. 431:3353–3365.

52. Ke, S., Ulmschneider, M. B., Wallace, B. A., & Ulmschneider, J. P. 2018. Role of the interaction motif in maintaining the open gate of an open sodium channel. *Biophysical journal*, *115*(10), 1920-1930.

53. Klesse, G., Tucker, S. J., & Sansom, M. S. 2020. Electric field induced wetting of a hydrophobic gate in a model nanopore based on the 5-HT3 receptor channel. *ACS nano*, *14*(8), 10480-10491.

54. Shen, H., Li, Z., Jiang, Y., Pan, X., Wu, J., Cristofori-Armstrong, B., Smith, J.J., Chin, Y.K., Lei, J., Zhou, Q. and King, G.F., 2018. "Structural basis for the modulation of voltage-gated sodium channels by animal toxins." *Science* 362.6412: eaau2596.

55. Shen, H., Zhou, Q., Pan, X., Li, Z., Wu, J., and Yan, N. 2017. Structure of a eukaryotic voltage-gated sodium channel at near-atomic resolution. *Science*, *355*(6328), eaal4326.

56. Kasimova, M. A., Yazici, A., Yudin, Y., Granata, D., Klein, M. L., Rohacs, T., and Carnevale, V. 2018. Ion channel sensing: Are fluctuations the crux of the matter?. *The journal of physical chemistry letters*, *9*(6), 1260-1264.

57. Kschonsak, M., Chua, H. C., Noland, C. L., Weidling, C., Clairfeuille, T., Bahlke, O. Ø., ... and Payandeh, J. 2020. Structure of the human sodium leak channel NALCN. *Nature*, *587*(7833), 313-318.

58. Huang, G., Liu, D., Wang, W., Wu, Q., Chen, J., Pan, X., ... and Yan, N. 2022. High-resolution structures of human Nav1. 7 reveal gating modulation through α-π helical transition of S6IV. *Cell Reports*, *39*(4), 110735.

59. Buyan, Amanda, Delin Sun, and Ben Corry. 2018. "Protonation state of inhibitors

determines interaction sites within voltage-gated sodium channels." *Proceedings of the National Academy of Sciences* 115.14: E3135-E3144.

60. Ragsdale, D. S., McPhee, J. C., Scheuer, T., and Catterall, W. A. 1994 . Molecular determinants of state-dependent block of Na+ channels by local anesthetics. *Science*, *265*(5179), 1724-1728.

61. Ragsdale, D.S., McPhee, J.C., Scheuer, T. and Catterall, W.A., 1996. . "Common molecular determinants of local anesthetic, antiarrhythmic, and anticonvulsant block of voltage-gated Na+ channels." *Proceedings of the National Academy of Sciences* 93.17 : 9270-9275.

Figure 1: Structure and proposed functional cycle for NavAb A. Full length structure of NavAb (PDB ID - 5VB2), initially described as a pre-open state, with activated voltage-sensor domains (VSD), contracted pore and ordered C-terminal domain (CTD). Inset shows a cartoon representation of one subunit highlighting the relevant domains - black - VSD, dark grey - Pore domain and light grey - C-terminal domain (CTD) **B**. Currently proposed functional cycle for NavAb based on structures resolved in different functional states.

Figure 2: The π model is a functional open state based on pore hydration, and on sodium ion and lidocaine permeation free energy profiles. A. Time averaged water number density along the central pore axis of the expanded α model. Standard deviation shown in gray shading, the blue dashed line shows the bulk water density and the inset shows a snapshot from the expanded α model simulation wherein the pore is dehydrated close to the activation gate. **B.** Time averaged water number density along the central pore axis of the expanded π model. The inset shows a snapshot from the expanded π model simulation with a completely hydrated pore. **C.** Hydrophobic constriction at V1213 (blue) leading to pore dehydration in the expanded α model (view from the extracellular medium). **D.** Sodium ion permeation free energy profile for the expanded α model (black), and for WT (red) and N1211L (orange) variants of the expanded π model along the central pore axis of the lower half of the S6 helix. **E.** Lidocaine permeation free energy profiles for the expanded α model (black) and expanded π model WT (red) along the central pore axis of the lower half of the S6 helix. Cartoon representations in the background of D and E show the lower half of the S6 helix.

Figure 3: Relationship between fenestrations and π-helix formation. A. Time and subunit averaged radius of fenestrations for different models. For each model, we plot the average bottleneck radius across four subunits. **B.** Surface representation showing the open (yellow circle) and closed fenestrations in the expanded α and π models respectively. The fenestrations in the π models are closed by F1207 (blue) and I1210 (red) residues. **C.** Lidocaine permeation free energy profiles for the resting model (black), expanded π model (red) and asymmetric α model (blue) along the central pore axis of the lower half of the S6 helix. **D.** Lidocaine binding site in the resting model, expanded π model and asymmetric α model. Lidocaine is bound closer to the selectivity filter in the resting model and expanded π model, with a common binding site at T1206. In the asymmetric α model, the drug binding site has shifted more towards the intracellular end away from the selectivity filter.

Figure 4: The neck region of the CTD proximal to the transmembrane helices can disorder.
A. A cartoon representation showing the neck region of CTD disordering upon simulating at 400K.
B. Secondary structure analysis of the CTD of NavAb (PDB ID - 5VB2) over simulations at 400K.

Figure 5: Contracted π model facilitates pore opening A. Distance between I1216 and I1119 used as a collective variable for the α model pore opening well-tempered metadynamics simulation. **B.** Distance between I1217 and I1119 used as a collective variable for the π model pore opening well-tempered metadynamics simulation. **C.** Pore opening free energy profile starting from the constricted α model showing a global minimum at around 1.45 nm, corresponding to a relatively constricted pore. Structural alignment between the most representative structure at around 1.45 nm pore opening (white) and the contracted α model (grey). **D**. Pore opening free energy profile starting from the constricted π model showing a minimum at around 1.09 nm corresponding to a expanded pore. Structural alignment between the most representative structure at around 1.09 nm pore opening (pink) and the expanded π model (red).

Figure 6: Proposed functional cycle for NavAb. Starting from the resting state with α helical S6 and hydrated peripheral cavities, NavAb transitions to a pre-open state with VSDs activated, α to π transition, and dehydration of the peripheral cavities. This conformation allows for favorable pore opening. Pore opening also involves CTD disordering although the point at which this happens remains unclear. Inactivation involves transition from π to α, but whether the pore is expanded, contracted in a symmetric or asymmetric manner, or even transitions between different conformations, remains unclear. Return to the resting state involves deativation of the voltage-sensor domains.