**Microbial community dynamics during a harmful *Chrysochromulina leadbeateri* bloom**

Nerea J. Aalto[1,2], Hannah Schweitzer[1,2], Erlend Grann-Meyer[1], Stina Krsmanovic[1], Jon Brage

Svenning[1], Lars Dalheim[1], Sebastian Petters[1], Richard Ingebrigtsen[1], Chris J. Hulatt[3], Hans C.

Bernstein[1,2]*

**Affiliations:**

*1. Norwegian College of Fisheries Sciences, UiT - The Arctic University of Norway, Tromsø,*

*Norway*

*2. The Arctic Centre for Sustainable Energy, UiT - The Arctic University of Norway, Tromsø,*

*Norway*

*3. Faculty of Biosciences and Aquaculture, Nord University, Bodø, Norway*

**<u>*Correspondence</u>**: Hans C. Bernstein, UiT-The Arctic University of Norway, BFE-fak.,

Postboks 6050 Langnes, 9037 Tromsø, NORWAY*; Phone: +47 776 46 114;  Email:*

Hans.C.Bernstein@uit.no

***Running Title:*** Ecology of the Norwegian harmful algae bloom of 2019

***Keywords:*** Marine Microbiome, Harmful Algae, Aquaculture, Norway, Time Series Analysis

**ABSTRACT**

In late spring 2019 a harmful algae bloom occurred across multiple, interconnected fjords and bays in northern Norway. This was a major event causing severe fish mortality at several salmon aquaculture facilities in the region. The reported focal taxon and suspected agent of fish mortality was a *Chrysochromulina leadbeateri*-like prymnesiophyte that was morphologically identical to the UiO-035 isolate collected from a similar bloom from the same region in 1991. This study reports on succession dynamics of the marine microbiome via detectable changes in abundance of 18S and 16S rRNA gene amplicon sequence variants and closely related taxa of the focal strain *C. leadbeateri* UiO-035. This taxon co-bloomed with a diverse group of microeukaryotes and prokaryotes and their temporal succession dynamics revealed high variability with strong shifts in abundance during the event. These blooming patterns were poorly connected to environmental conditions suggesting that other factors such as biological interactions may be at least as important in shaping the dynamics of this type of harmful algae bloom. A deeper understanding of microbiome succession patterns during these rare but destructive bloom events will help guide future efforts to generate predictive models and anticipate deviations from the natural bloom cycles of the northern Norwegian fjord systems that are home to large scale aquaculture activities.

## INTRODUCTION

Harmful algal blooms (HABs) are common in warm and temperate waters, but also occur in northern cold water ecosystems (Hallegraeff et al., 2021; Karlson et al., 2021). These blooms pose a serious risk for aquaculture activities, because farmed fish cannot escape from developing adverse conditions caused by marine microalgae. Previously, the most infamous northern Norwegian bloom occurred in 1991 in Ofotfjord causing high mortality in affected fish farms (Rey, 1991). Smaller HAB incidents have occurred since then, although they have gained little mainstream attention (Grann-Meyer, 2020). The salmon aquaculture industry has expanded tremendously over the past decades to a landing value of salmon that reached 7 billion USD in 2019 (Statistics Norway, 2020). While this has caused concern, it has not spurred investments in long-term monitoring of algal bloom dynamics or occurrence of potentially harmful algae species in areas that support salmon aquaculture. Therefore, most stakeholders were taken by surprise when a deadly HAB of *Chrysochromulina leadbeateri* (prymnesiophyte) reoccurred and caused massive fish mortalities in the Troms and Nordland regions of costal northern Norway from mid-May to the beginning of June 2019 (Karlsen et al., 2019). The fish loss was ca. 14 500 tonnes, which is the highest amount among fish kill related HABs in Scandinavia since 1988 (Karlsen et al., 2019; Karlson et al., 2021). Thus, knowledge that can eventually inform causation is very limited, especially in the context of seasonal microbial blooming dynamics, detailed taxonomy of co-blooming species, and the environmental conditions during the 2019 incident.

Marine HABs are often associated with altered ratios of inorganic nitrogen-to-phosphate (N:P), coupled with a long period of high irradiance, calm weather, and strong surface water stratification (Nielsen et al., 1990; Anderson et al., 2002; Uronen et al., 2005). However, the role of N:P ratio in determining the magnitude, frequency, and duration of HABs has been questioned (Davidson et al., 2012) as more evidence is showing that abiotic conditions

are not the only influencing factors of microalgal bloom dynamics. The importance of intra-
and interspecific interactions between members of the marine microbiome has become well
recognized as a major factor in structuring the temporal and spatial relationships of
phytoplankton communities (e.g., Needham and Fuhrman, 2016; Martin-Platero et al., 2018;
Aalto et al., 2022). These interactions can be complex and subject to rapid change within a
marine microbial ecosystem. Although they are difficult to disentangle, they have the potential
to stimulate, enhance, and/or perpetuate the bloom of specific taxa associated with HABs.
Therefore, it is important (yet uncommon today) to observe and catalogue the blooming patterns
of whole microbiomes while studying the ecology of focal taxa involved with HAB events.

High-resolution (densely sampled) time series analysis combined with high-resolution
taxonomic screening such as 16S and 18S ribosomal RNA gene amplicon sequencing variants
or operational taxonomic unit-based diversity is an important method for evaluating microbial
succession dynamics and ecological factors that drive blooms at different time scales (Fuhrman
et al., 2015; Martin-Platero et al., 2018). However, knowledge about the nature of taxon-
specific and non-specific interactions with HAB species is poor, because identification of these
"hidden" patterns of interactions in dynamic and species rich communities is challenging.
Therefore, an increasing number of modern analytical tools has been developed to apply with
time series analysis. For example, WaveClust (Martin-Platero et al., 2018) and k-medoid
(Kaufman and Rousseeuw, 2009; Coenen et al., 2020) are applicable and promising approaches
to infer both temporal dynamics and networks of microbial interactions.

The specific mechanisms of how HABs cause fish mortality are often unknown and
difficult to identify, as was the case for the 1991 and 2019 *C. leadbeateri* HABs in northern
Norway (Meldahl et al., 1994; Edvardsen and Paasche, 1998). In fact, the toxicity of many well-
known toxic prymnesiophytes – such as *Chrysochromulina* spp. – are not known to be
dependent on cell density, which makes detection difficult (Edvardsen and Paasche, 1998). The

role that bacteria may play in bloom toxicity is also not clear (Uribe and Espejo, 2003). For example, it has been suggested that toxin biosynthesis in dinoflagellate cells is indirectly mediated by associated bacterial consortia (Green et al., 2004). Heterotrophic bacteria have been shown to increase activity and biomass concentrations during a HAB due to the release of dissolved organic matter from lysed microalgal competitors (Uronen et al., 2007). In order to increase their supply of nutrients, many bacteria also exhibit target-specific or non-specific algicidal activity that inhibit the growth or lyse microalgae; potentially even terminating HABs (Doucette et al., 1999; Meyer et al., 2017). Several known marine algicide bacteria are found in bloom-associated groups such as *Flavobacteria* (*Bacteroidetes*), *Alphaproteobacteria* and *Gammaproteobacteria* (Mayali and Azam, 2004). However, not all bacteria-algae interactions are negative. Several mutualistic and growth enhancing relationships have also been identified, which include metabolite exchange of vitamins and growth hormones from bacteria that trade with microalgae for amino acids and dissolved organic carbon (Cirri and Pohnert, 2019).

The spring-summer phytoplankton seasonal succession along the coast of northern Norway has been extensively studied in the past 40 years with specific focus on spring bloom dynamics. The majority of data collected to-date has used only morphology-based identification of microalgal taxa, which has limited the extent to which the full diversity of these communities has been described (Eilertsen et al., 1981; Reigstad and Wassmann, 1996; Degerlund and Eilertsen, 2010; Aalto et al., 2021). From previous studies, we know that the onset of the "spring bloom" occurs in late March/early April, and it is driven by increased irradiance as the region undergoes a shift from polar night to spring, which results in elevated winter nutrient concentrations and a well-mixed water column (Eilertsen and Taasen, 1984; Eilertsen and Frantzen, 2007; Aalto et al., 2021). This leads to a rapid and intense increase of many chain forming centric diatoms that co-bloom with *Phaeocystis pouchetii* (haptophyte)*,* who often outlasts the diatoms and persists throughout the summer (Degerlund and Eilertsen, 2010). The

spatial and interannual variation of the late bloom (May-June) phytoplankton community can comprise a diverse mix of low-nutrient-tolerant, heterotrophic, and mixotrophic species belonging to dinoflagellates, cryptophytes, ciliates, and other small flagellates (Eilertsen et al., 1981; Aalto et al., 2021). HAB incidents in northern Norway tend to occur after the primary spring diatom bloom, although historical sampling of these events is sparse and the onset and development of HAB conditions is essentially unknown for the region.

Our aim was to contribute to the knowledge of the ecology and microbial succession dynamics associated with the *C. leadbeateri* HAB, which has now proven to be a reoccurring and devastating event in northern Norway. This was accomplished by asking three targeted research questions: i) how did the marine microbiome composition change during the 2019 HAB event in relative relation to the focal taxon?; (ii) what were the major blooming (temporal) patterns among prokaryotes and microeukaryotes and which taxa co-bloomed with the focal taxon?; and iii) which environmental factors were associated with the observable differences in the marine microbiome composition? This study represents the first molecular ecology-based investigation of a *C. leadbeateri* HAB and presents new insights into a seasonal marine microbial ecosystem that can have a major impact on the social-economic welfare of Norway through its relation to the aquaculture industry.

**MATERIALS AND METHODS**

**Study areas and sample collection.** Time series sampling was performed between May 27th (sample day 0) and June 21st (sample day 25) in 2019, at two primary stations (Supplementary Table S1). The first station was named Grøtsund sampling station (**GSSS**, 69.80 N, 19.36 E). The second station, named as Finnfjord sampling station (**FSS**, 69.19 N, 18.03 E), is located 60 km south of GSSS, within the same interconnected fjord-sound system (Figure 1a). Samples were collected from each station twice a week, except station FSS was sampled once in the

second sampling week (starting the week of May 27th, 2019), from fixed sampling depth of 10

m using a 1.7 L Niskin bottle to get approximately 8.5 L of seawater. The collected seawater

was prefiltered for large biomass with a 150 μm plankton mesh, and transported to shore in a

15 L Nalgene plastic carboy. All sampling equipment was rinsed with 2% bleach prior and

between each sampling effort followed by a rinsing with MilliQ water and a minimum of two

volumes of station sampling water prior to collection. Temperature and salinity profiles down

to 15–50 m were recorded with a hand-held AML oceanographic X2 electron conductivity-

temperature-depth (CTD) instrument (AML Oceanographic, Canada). Seawater particles – i.e.

microbial biomass – was collected by vacuum filtration onto 0.22 μm polycarbonate filters (GE

Healthcare Whatman 47 mm Nucleopore Polycarbonate Track Etched Membranes; GE

Healthcare 111106). All filtering was done inside a sterile laminar flow hood, and the total

filtered volume per sample varied from 500 – 750 mL depending on turbidity. Filters were

immediately flash frozen and stored at -80 °C. All filtering equipment was disinfected with 2%

bleach prior and between sampling efforts and thereafter rinsed with MilliQ water. Water

samples for macronutrients: silicate ($Si(OH)_4$), phosphate ($PO_4^{3-}$), and nitrate + nitrite ($NO_3^-$ +

$NO_2^-$), were filtered through 47 mm GF/F glass microfiber filters (Whatman) into Falcon tubes

using disposable syringes and thereafter stored at -20 °C.

In addition to the time series samples from FSS and GSSS, filtered microbial biomass

was collected from eight separate locations as single time points between May 27th and June

21st (Figure 1a and Supplementary Table S1). The data of these samples is not included within

time series analyses but used to compare the 2019 HAB affected area via spatial variation in

the relative abundance of the focal taxon and associated microbial communities. The DNA

extraction and downstream analysis of all samples were performed at the UiT – The Arctic

University of Norway as described below.

**Environmental factors**. The vertical surface water properties from the time series sampling

stations (FSS and GSSS) were inspected using CTD-profile data (Supplementary Figure S1 and

S2). The difference in surface water stratification between sampling stations was determined

by calculating the mixed layer depth (MLD) from each CTD-profile data using potential

density. The depth of 1 m was used as a reference depth as the deployed handheld CTD-sensor

enables measurements near surface. A density threshold of $0.1$ kg m$^{-3}$ was used to define MLD

– i.e. the depth below 1 m where density first exceed the defined threshold (Peralta-Ferriz and

Woodgate, 2015). Temperature and salinity values from the depth of 10 m (corresponding the

depth of collected biomass) as well as MLD values were further used in statistical analyses. As

the CTD-casts between sampling days 11 and 17 were unsuccessful, temperature, salinity, and

MLD values for these days were interpolated using the R function 'na.approx' from the R

package 'zoo' to fill the missing values. Nutrient concentrations were analyzed 20 months later

at the UiT – The Arctic University of Norway with a Quaatro39 autoanalyser (Seal Analytical,

UK). $PO_4^{3-}$ concentrations below the detection limit of $0.04$ μmol L$^{-1}$ were replaced with a small

value (0.001) to enable the estimation of this variable in statistical analysis.

**Unialgal control cultures.** Controls were performed with the *Chrysochromulina leadbeateri*

UiO-035 (Edvardsen et al., 2011) culture obtained from the Norwegian Culture Collection of

Algae (NORCCA) repository and cultivated at 6 °C on f/2 medium (Guillard, 1975), fortified

with 20 nM of selenium. The cultured *C. leadbeateri* was isolated from a toxic bloom of 1991

in Vestfjorden, northern Norway (Eikrem and Throndsen, 1998). Taxonomic identification of

the cultured isolate has been mainly based on scale morphology in comparison to holotype

(Estep, 1984) and other previous reports on *C. leadbeateri*-like cells both from northern and

southern hemisphere (Eikrem and Throndsen, 1998). Several *Chrysochromulina* species are

characterized by species-specific bilayer scales which have been considered a morphotype-

specific character (Eikrem and Throndsen, 1998). Cells were collected and processed

identically to the environmental samples described below. The 18S amplicon analysis assigned

four abundant ASVs comprising these control cultures (>1000 ASV counts in all replicates)

(Supplementary Figure S3). All four of these ASVs had identical taxonomic assignments within

the SILVA v138.1 database: Family = Prymnesiales, Genus = OLI16029, Species = Unknown.

Of these four ASVs, ASV *24a92740c5af4cd5a5ac0db70830fbc0* (hereafter referred to as the

"focal ASV") comprised 88 % of the OLI16029 unialgal control (Supplementary Figure S4).

**Amplicon sequencing**. Genomic DNA was extracted and downstream amplicon sequences

were obtained using the same protocol as described in Aalto et al. (2022). All the samples were

processed together in accordance with the Earth Microbiome Project protocols (Gilbert et al.,

2010) with modifications according to Amaral-Zettler et al. (2009). This included

recommended primers and barcodes for the V4 hypervariable region of the 16S SSU rRNA

gene using the V4 forward (515F) and V4 reverse (806R) primers, and the V9 hypervariable

region of the 18S SSU rRNA was targeted with the V9 forward (1391F) and V9 reverse (EukBr)

primers. Biological replicates, four or five, were sequenced on an Illumina MiSeq instrument

(Illumina, San Diego, CA) according to (Caporaso et al., 2010) at Argonne National Laboratory

(Lemont, IL, USA). The realized length of forward and reverse reads was 151 bp.

**Amplicon analysis.** The amplicon analyses were performed through the QIIME2 environment

and using QIIME2 plugins as previously described (Aalto et al. 2022). Briefly, Illumina reads

and the corresponding barcode files were imported and demultiplexed using the Earth

Microbiome Project paired end flag. All reads were filtered, de-replicated, and chimera checked

using all default parameters in the DADA2 v2021.2.0 (Callahan et al., 2016). Reads were

merged and amplicon sequence variants (ASVs) were determined using DADA2 v2021.2.0.

The DADA2 statistic on sequence reads is provided in Supplementary Data S1 (osf.io). A 16S

and 18S rRNA gene classifier from the SILVA v138.1 database was trained using RESCRIPt

(Quast et al., 2012; Robeson et al., 2020). The ASVs were classified with the self-trained

classifier database. (Yilmaz et al., 2014). It is noted that the major challenge in 16S and 18S

based taxonomy is its dependency on taxonomic resolution limitations of the database. We

chose to use all classifications given from the SILVA v138 database because it is a standardized

approach. Manual curation of taxonomic classification was beyond the scope of this study.

**Statistics, clustering and correlation.** All ASVs not assigned to the expected kingdom

(*Archaea*, *Bacteria* and *Chloroplast*) were removed along with all mitochondria assignments

from 16S data set. ASVs assigned to *Archaea*, *Bacteria*, *Vertebrata* and *Arthropoda* and phyla

*Cnidaria*, *Echninodermata*, *Annelida*, *Nematozoa*, *Nemertea*, *Porifera*, *Mollusca*, *Ctenophore*

and *Tunicata* or phyla belonging to macroalgae and land plants or land fungi such as *Bangiales*,

*Floridephycidae*, *Ochrophyta*, *Phragmoplastophyta*, *Ascomycota*, *Basidiomycota* and

*Myxogastria* were removed from the 18S data set. Additional information on number of ASVs

and sequencing depths before and after removal of unwanted taxa is provided in Supplementary

Table S2. The total number of classified ASVs in FSS for 18S and 16S data set was 900 and

1309, respectively, and in GSSS 2196 and 1865, respectively. As an exception regarding

correction of taxonomic annotations, classified 16S order *Enterobacterales* was manually

changed to closely related order *Alteromondales* as the main genera of this order were

*Pseudoalteromonas*, *Colwellia* and *Glaciecola* that are known to be members of

*Alteromonadales* (Parte et al., 2020).

Downstream analysis was completed in R (R Core Team, 2021), using the 'microeco'

(Liu et al., 2021) and 'vegan' packages (Oksanen et al., 2013). This study also made use of the

K-medoids clustering method adapted from a previously described approach applied to time

series microbiome data to examine distinct temporal trends in blooming patterns within

communities (Coenen et al., 2020). Briefly, ASV counts from each sample type – i.e., sampling

station (FSS or GSSS) and PCR amplification primer type (16S or 18S) – were preprocessed

via z-score and variance stabilizing transformations using the 'DESeq2' package in R (Love et

al., 2014). Pairwise distance matrices were calculated by the Euclidean distance prior to

partitioning each sample-specific ASV into K clusters according to similarity criterion over the

time series. The quality of each cluster per each sample type was assessed via the Calinski-

Harabasz index (Supplementary Figure S5) (Lord et al., 2017). Pearson's correlation was used

to examine the relationship between each temporal dynamic, represented by medoid taxon

(cluster centroids), and environmental measurements but also, to obtain the positively and

negatively co-blooming microalgae and bacterial taxa with focal ASV.

The difference in microbial communities, i.e. beta-diversity, between sampling stations

was measured using unweighted UniFrac distance metric which accounts occurrence -

presence/absence of ASV - and phylogenetic diversity (Lozupone et al., 2011). The beta-

diversity and relationship between community compositions and environmental factors

(temperature, salinity, MLD, $Si(OH)_4$, $PO_4^{3-}$, $NO_3^- + NO_2^-$, and relative abundance of focal

taxon (OLI16029)) was visualized and conducted via distance based redundancy analysis

(dbRDA). The permutational multivariate analysis of variance (PERMANOVA) was

performed to test the (dis)similarity in prokaryotic and microeukaryotic community

compositions between stations (pairwise) and within stations (Anderson, 2001). The Tukey test

was performed on inorganic nutrients to determine if the mean concentration values were

statistically different between stations (Abdi and Williams, 2010).

**Data repository and reproducible analyses.** All sequencing and environmental data is

available on the Open Science Framework (osf.io), along with all R Markdown scripts used for

analyses and graphing (osf.io/4wjhp/). All Supplementary Data S1–S11, referred to in this

study, is available and found on osf.io/4wjhp/.

**RESULTS**

**Geographical distribution of focal taxon in relation to fish mortality.** The focal taxon was

assessed as those ASVs assigned to OLI16029, representing the *Chrysochromulina leadbeateri*

UiO-035 unialgal strain within the class *Prymnesiophyceae*. This taxon was detected during the

2019 harmful *C. leadbeateri* bloom in varying relative abundance across the northern

Norwegian coast and fjords (Figure 1). It´s abundance varied between 0.4 % and 51.8 % of the

microeukaryotic community assessed via 18S amplicon sequencing through the HAB affected

region. The highest relative abundances of the focal taxon – recorded in our molecular sampling

campaign – were obtained in the northern portion of the HAB affected region that included

Middle-Balsfjord, South-Tromsø, GSSS, and Tussøy stations (Figure 1a). The microbial

community composition was relatively similar between Middle-Balsfjord, South-Tromsø, and

GSSS stations with increased relative abundance of 18S ASVs and 16S ASVs indicative of

dinoflagellates (class *Dinophyceae*) and order *Flavobacteriales*, respectively, towards station

GSSS (Figure 1b-c). The relative abundance of ASVs belonging to the focal taxon were low in

Ballangen (Ofotfjord) where the HAB was first reported to originate with high fish mortality in

mid-May (Karlsen et al., 2019). The community structure from Ballangen was also distinct,

compared to other samples, as it was dominated by 18S ASVs indicative of centric diatoms

notably from the genus *Skeletonema* (class *Mediophyceae*) and 16S ASVs classified within the

genera *Pseudoalteromonas* and *Colwellia* (order *Alteromonadales*) (Figure 1b-c and

Supplementary Data S2 and S3 (osf.io)). The relative abundance of the focal taxon was higher

in samples collected from Sagfjord stations (Anevik, Dyping, Veggfjell, and Middle-Sagfjord),

south from Ofotfjord, than in the Ballangen and FSS stations (Figure 1a). The microeukaryotic

community in the Sagfjord location was especially diverse, comprising multiple distinct groups

of dinoflagellates and centric diatoms being the most abundant. In contrast to the other sampling

locations, the 16S prokaryotic ASVs among the Sagfjord stations were predominantly

comprised by the order *Pseudomonadales* (Figure 1b-c).

**Figure 1.** A map showing the geographical area in northern Norway where *Chrysochromulina*
*leadbeateri* -related harmful algal bloom occurred in summer 2019. The main sample locations
are *named* on the map. **A)** Microbial DNA for 18S rRNA and 16S rRNA amplicon sequence

analysis was collected once from the marked stations, except from FSS (Finnfjord sampling
station) and GSSS (Grøtsund sampling station) which are the primary stations of this time series
study and were sampled 7 and 8 times, respectively, between May 27[th] and June 21[st]. The size
of the circles represents the relative abundance of focal taxon (average relative abundance for
FSS and GSSS stations across all samples). Not all of the stations represent localities where
HAB- related fish mortality was experienced. Taxonomic composition of the most abundant **B)**
prokaryotes (16S) and **C)** microeukaryotes (18S) retrieved from Amplicon Sequence Variants
(ASVs) tables specified at the order and class taxonomic rank level, respectively. Note that FSS
and GSSS stations are excluded from **B** and **C** as the taxonomic composition across time series
is shown in Figure 2.

**Microbiome dynamics across the bloom.** The microeukaryotic and prokaryotic community

composition changed across the time series sampling campaign at FSS (PERMANOVA: 16S,

$R^2 = 0.11$ and $p = 0.001$; 18S, $R^2 = 0.19$ and $p = 0.001$) and GSSS (PERMANOVA: 16S, $R^2 =$

$0.13$ and $p = 0.001$; 18S, $R^2 = 0.13$ and $p = 0.001$). These dynamic communities were also

significantly dissimilar between the two time series stations (PERMANOVA: 16S, $R^2 = 0.30$

and $p = 0.001$; 18S, $R^2 = 0.46$ and $p = 0.001$) (Supplementary Table S3). GSSS showed higher

species richness as compared to FSS as measured by the number of observed ASVs per sample

day; 235–288 and 244–358 ASVs for 16S and accordingly 161–265 and 306–415 ASVs for

18S in FSS and GSSS, respectively (Supplementary Figure S6).

Among the most abundant microeukaryotes assigned to the taxonomic level class, only

18S ASVs indicative of *Cryptophyceae*, *Mediophyceae*, *Dinophyceae* and *Prymensiophyceae*

(including the focal taxon) were shared between both stations (Figure 2a-b). The observed

community in FSS was dominated by taxa classified as *Cryptophyceae* (9 – 49 % of the relative

abundance), comprising mainly ASVs assigned to the genus *Teleaulax*, that showed strong

temporal variability in the second half of the time series (Figure 2a and Supplementary Data S4

(osf.io)). The predominant taxon observed from GSSS was classified as *Dinophyceae,* which

accounted for 34–54 % of the relative abundance during the time series (Figure 2b). Most of

the ASVs belonging to the class *Dinophyceae* were not assigned to a lower taxonomic level.

A change in community structure was observed at both stations during the middle of the

observed bloom; between days 7 and 15 in FSS and between days 11 and 16 at the GSSS station

(Figure 2a-b). This shift was observed with an increased relative abundance of ASVs indicative

of centric diatoms (class *Mediophyceae*) in both stations and the class *Coscinodiscophytina* in

FSS which were almost exclusively represented by ASVs indicative within genus

*Leptocylindru*s (Figure 2a-b and Supplementary Data S4). The class *Mediophyceae* comprised

ASVs that were classified within different genera between stations, such as *Acrocellulus* at FSS

and *Thalassiosira* at GSSS (Supplementary Data S4-S5 (osf.io)). In FSS, the contribution of

class *Prymnesiophyceae* to the community composition was minor in comparison to GSSS

(Figure 2a-b) and was composed mainly by ASVs classified to the genus *Chrysochromulina* (2

– 3 % of relative abundance) but also ASVs assigned to the focal taxon, OLI16029 (0.5 – 1 %

of relative abundance) (Figure 2c and Supplementary Data S4-S5 (osf.io)). In GSSS, class

*Prymnesiophyceae* prevailed until day 11 (42 % of the relative abundance) and thereafter

drastically decreased while ASVs assigned to *Syndiniales* concurrently increased (Figure 2b).

The majority of those ASVs belonging to the class *Prymnesiophyceae* were assigned to the

genera *Phaeocystis* and the focal taxon (OLI16029) (Supplementary Data S5 (osf.io)). The

pronounced decrease in abundance of the focal taxon at day 8 coincided with highest relative

abundance of *Phaeocystis* (Figure 2d and Supplementary Data S5 (osf.io)). We note here for

clarity that ASVs assigned to the genus *Chrysochromulina* are not indicative of the focal taxon

(OLI16029), even though the unialgal control has been deposited and frequently referred to as

*Chrysochromulina leadbeateri* UiO-035.

The 16S ASVs – classified at the level of order – showed less temporal variability than

the 18S ASVs (at the class level) across the time series in FSS and GSSS. The taxa with the

highest relative abundances were the same in FSS and GSSS sample stations (Figure 2e-f).

Taxa classified to the order *Flavobacteriales* showed the highest relative abundance with ASVs

belonging *Rhodobacterales* and *Pseudomonadales* being common as well in both time series

stations (Figure 2e-f). The FSS and GSSS locations harbored distinctly different relative

abundances of genera assigned to *Flavobacteriales*. At station GSSS, the dominant genera were

*Polaribacter*, *Ulvibacter*, *Formosa* and the NS5 marine group, as compared to FSS where the

NS5 and NS9 marine groups were dominant (Supplementary Data S6 and S7 (osf.io)). At

GSSS, members of the order *Verrumicrobiales* increased in relative abundance (from 1 % to 8

%**)** within the same time frame as major changes in the microeukaryotes were observed (Figure

2f). Also, the abundance of the *Pseudomonadales* order decreased towards the end of the time

series while the contribution of *Flavobacteriales* simultaneously increased (Figure 2f). The

influence of Archaea to the 16S ASV composition was minor at both stations and accounted

for only 0.1 % and 0.4 % of the relative abundance in FSS and GSSS, respectively

(Supplementary Data S6 and S7 (osf.io)).

Since this study did not include measurements of chlorophyll *a* to indicate changes in

phytoplankton abundance, the 16S ASVs identified as Chloroplast were used as an inference.

The average relative abundance of ASVs assigned to Chloroplast was nearly three times higher

in FSS than in GSSS and the temporal pattern was different: the abundance increased towards

the end of the time series in FSS whereas in GSSS this was opposite (Figure 2e-f).

**Figure 2.** Comparative composition of the most common microeukaryotic and prokaryotic taxa collected from the time series sample stations and determined via 16S and 18S rRNA amplicon analysis. Microeukaryotes classified at the class taxonomic rank level **A)** in FSS and **B)** in GSSS. The temporal change in relative abundance of the focal taxon assigned as genus *OLI16029* (class *Prymnesiophyceae*) **C)** in FSS and **D)** in GSSS. Prokaryotes classified at the order taxonomic rank level **E)** in FSS and **F)** in GSSS.

**Environmental variability during the HAB**. The surface water properties – temperature and salinity – were relatively similar during the HAB, as observed from both time series stations (FSS and GSSS) at 10 m depth from where microbial biomass was collected (Supplementary Table S4). The overlaying surface water was more stratified in FSS than GSSS. This was due

to a low salinity layer (< 30) which persisted near the surface during the entire time series

simultaneously with a strong increase in observed temperature (Supplementary Figure S1 and

S2). The mixed layer depth was observed near the surface (3 m) throughout the study period in

FSS and in the beginning in GSSS time series, where it then sank down to 7 m by the end of

the GSSS time series while the surface water became more homogeneous after day 8

(Supplementary Figure S1 and S2). The mean $NO_3^- + NO_2^-$ concentration was significantly

higher in GSSS than in FSS (0.43 and 0.18 µmol $L^{-1}$, respectively; Tukey, $p < 0.02$) whereas

the $Si(OH)_4$ and $PO_4^{3-}$ concentrations did not show statistically significant differences between

stations (Supplementary Table S4).

**Beta diversity in context with the environment**. Both the microeukarytotic and prokarytotic

components of the HAB-associated microbiome were dissimilar between GSSS and FSS

stations and throughout the observed time series. This was measured via unweighted UniFrac

distances that were ordinated by dbRDA analysis to infer the correlation between selected

environmental measurements to differences in community composition (Figure 3). Results of

beta diversity also corroborated the observed change in community structure during the middle

of the time series that was seen and noted above by inspecting changes in relative taxonomic

abundance (Figure 2). The dbRDA analysis (i.e., sum of RDA1 and RDA2) showed that 77.9

% and 83.7 % of the total variation in the prokaryotic and microeukaryotic community

composition was correlated to environmental measurements during the time series (Figure 3).

Most of this variation was explained by the first RDA component (RDA1: 16S = 60.8 % and

18S = 72.6 %) inferring that the variation of both prokaryotic and microeukaryotic communities

between FSS and GSSS may be driven by response to environmental factors such as MLD,

salinity, and $NO_3^- + NO_2^-$ (Figure 3). Whereas the temporal variability within stations was

 mainly explained by the second RDA component (RDA2) which accounted for 17.1 % and 11.1

 % of the variance in prokaryotic and microeukaryotic communities, respectively.

 **Figure 3.** Dissimilarity in prokaryotic and microeukaryotic community composition between
 sample stations was inferred via beta-diversity as measured with unweighted UniFrac distances
 and related to environmental measurements. Distance-based redundancy analysis (dbRDA)
 biplot of **A)** prokaryotes and **B)** microeukaryotes. Color represents station and shape comprises
 samples (sample scores) within each sample week according to legend information. Arrows
 indicate environmental factors and point to the direction of maximum variation of the respective
 factor. T, temperature; S, salinity; MLD, mixed layer depth, Si, $Si(OH)_4$; PO4, $PO_4^{3-}$; N+N,
 $NO_2^- + NO_3^-$; Focal %, relative abundance of focal taxon in 18S data set.

 **Temporal trends and co-blooming taxa.** The HAB-associated marine microbiome was

 complex and consisted of a minimum of 235 and 161 16S and 18S ASVs, respectively, each

 maintaining its own dynamic trajectory during the observed time series. Examination and

 subsequent clustering of the most predominant blooming patterns was performed via the k-

 medoids clustering method. The resulted number of clusters per data set and time series station,

 as assessed via the Calinski-Harabasz index (normalized ratio for inter-intra-cluster variance)

 (Supplementary Figure S5), was 3, except for 18S at FSS where 4 clusters were selected. The

temporal pattern of each cluster was specified by its medoid taxon as a representative shape for

cluster's blooming dynamics of microeukaryotes (Figure 4a-b) and co-blooming taxa assigned

to 16S ASVs (Figure 5a-b).

The medoid 18S taxon of cluster 1 and 3, from both stations, exhibited approximately

inverse blooming patterns. These clusters also showed strong fluctuation in relative abundances

across the time series, including a strong positive and negative shift, respectively, around the

middle time points and negative correlations to environmental measurements (Figure 4a-d).

The medoid taxon of cluster 2 and 4 in FSS and cluster 2 in GSSS exhibited a short increase in

abundance in the beginning or end of time series (Figure 4a-b).

The focal ASV was included within cluster 3, at both stations (Figure 4a-b). There were

several microalgae that co-bloomed with the focal ASV, but these were distinct between the

FSS and GSSS stations (Figure 4). In FSS the focal ASV co-bloomed with genera *Teleaulax*

(class *Cryptophyceae*) and *Leptocylindrus* (class *Coscinodiscophytina*) which comprised 16 –

53 % and 1 – 53 % of the relative abundance within their respective cluster 3 (Figure 4e and

Supplementary Data S8 (osf.io)). A diverse composition of ASVs indicative of MAST clades

(marine stramenopiles) were also detected, although their relative abundances were minor (< 1

% of relative abundance within the cluster 3). This group was represented with other nano- and

picoplankton such as genera *Telonema* (class *Insertea Sedis*) and *Picomonas* (class

*Picomonadida*) the taxa in cluster 1 with inverse blooming pattern to cluster 3 (Figure 4c,e and

Supplementary Data S9 (osf.io)). ASVs from the genus *Phaeocystis* predominantly co-bloomed

at GSSS with the focal ASV (within cluster 3) together with a less abundant group of ciliates

(class *Intramacronucleata*) and members belonging to genus *Thalassiosira* (class

*Mediophyceae*) (Figure 4d,f and Supplementary Data S10 (osf.io)). ASVs belonging to the class

*Dinophyceae* also included in cluster 3 from GSSS (along with the focal ASV), and these were

composed mainly of members classified within the genus indicative of *Gymnodinium*.

The blooming patterns of 16S ASVs were also clustered. Among these, two clusters (1 and

2 from FSS and GSSS, respectively) showed significant positive correlation (FSS: $r = 0.79$ , $p$

$= 0.036$ and GSSS: $r = 0.71$, $p = 0.047$) with the focal ASV. Only cluster 2 from station FSS

showed a significant negative correlation with the focal ASV ($r = -0.96$, $p < 0.001$) (Figure 5c-

d). The temporal trend represented by the medoid taxon was nearly identical in cluster 3 at both

stations corresponding to the blooming pattern of cluster 2 and inverse blooming pattern of

cluster 4 for microeukaryotes in GSSS and FSS, respectively.

**Figure 4.** Characterization of major blooming patterns in microeukaryotic communities
assessed by K-medoid clustering analysis with z-transformation and detrending of 18S ASVs.
Temporal dynamics of microeukaryotic taxa and number of ASVs per defined cluster **A)** in FSS
and **B)** in GSSS station. The blooming pattern of each cluster is specified by medoid taxon and
drawn with black line. The red line specifies the blooming pattern of focal ASV. Y-axis is a z-
score and a value 0 denotes the mean abundance. Heatmap of Pearson's correlation between
temporal dynamics of medoid taxon (z-scores) of each cluster and focal ASV (z-scores) and
environmental factors **C)** in FSS and **D)** in GSSS. Color indicates Pearson's correlation
coefficient according to the color legend and level of significance is marked with ★ (★, $p \leq$
0.1; ★★, $p \leq 0.05$; ★ ★ ★, $p \leq 0.001$). The grey color for MLD in FSS denotes undefined
correlation as MLD values remained the same through the study period. Focal.dyn, temporal
dynamic of focal ASV; MLD, mixed layer depth; N+N, $NO_2^- + NO_3^-$; PO4, $PO_4^{3-}$; S, salinity;
Si, $Si(OH)_4$; T, temperature. Taxonomic profile of the most common microeukaryotes of each
cluster as relative abundance in a whole community classified at the class taxonomic rank level
**E)** in FSS **F)** in GSSS.

We infer that the bacterial taxa from the GSSS station belonging to cluster 2 co-bloomed

with phototrophic protists in general, because the ASVs assigned to Chloroplasts occupied the

same cluster (Figure 5f and Figure 4f). This cluster was also the only one to show a statistically

significant correlation with phosphate concentrations. The same pattern was not observed in

FSS where approximately two thirds of Chloroplast ASVs were identified within cluster 2 and

rest in cluster 3 (Figure 5e).

Pearson correlation enabled us to infer that different bacterial taxa co-bloomed with the

focal ASV at each time series sampling stations, FSS and GSSS (Figure 5c-d) corresponding to

the observations within 18S microeukaryotes. Cluster 1 at station FSS was dominated by the

order *Flavobacteriales* (35 – 63 % relative abundance within cluster 1) (Figure 5e) and was

predominated by sequences identified within the NS5 marine group but also *Ulvibacter*

(Supplementary Data S11 (osf.io)). ASVs assigned to *Rhodobacterales*, notably evenly

dominating genera *Amylibacter* and *Planktomarina*, also shared the similar temporal dynamics

with the focal ASV (Figure 5e and Supplementary Data S11 (osf.io)). The positive co-blooming

trend between *Flavobacteriales* and *Rhodobacterales* classes and the focal ASV was not

observed at station GSSS.

**Figure 5.** Characterization of major blooming patterns in prokaryotic communities assessed by
K-medoid clustering analysis with z-transformation and detrending of 16S ASVs. Temporal
dynamics of microeukaryotic taxa and number of ASVs per defined cluster **A)** in FSS and **B)**
in GSSS station. The blooming pattern of each cluster is specified by medoid taxon and drawn
with black line. Y-axis is a z-score and a value 0 denotes the mean abundance. Heatmap of
Pearson's correlation between temporal dynamics of medoid taxon (z-scores) of each cluster
and focal ASV (z-scores; Figure 4) and environmental factors **C)** in FSS and **D)** in GSSS. Color
indicates Pearson's correlation coefficient according to the color legend and level of
significance is marked with ★ (★, $p \leq 0.1$; ★★, $p \leq 0.05$; ★ ★ ★, $p \leq 0.001$). The grey color
for MLD in FSS denotes undefined correlation as MLD values remained the same through the
study period. Focal.dyn, temporal dynamic of focal ASV; MLD, mixed layer depth; N+N, $NO_2^-$
+ $NO_3^-$; PO4, $PO_4^{3-}$; S, salinity; Si, $Si(OH)_4$; T, temperature. Taxonomic profile of the most
common prokaryotes of each cluster as relative abundance in a whole community classified at
the class taxonomic rank level **E)** in FSS **F)** in GSSS.

**DISCUSSION**

The results from this study show distinct spatial variation and differences in the dynamic

community composition across the portion of the northern Norwegian coast that was strongly

affected during the 2019 *Chrysochromulina leadbeateri* HAB. This was a major and disruptive

event due to severe fish mortality and hence there were several sampling campaigns launched

simultaneously by public and private stakeholders. To our knowledge, this study represents the

only molecular-based investigation of the holistic marine microbiome during this HAB event.

**Indication for separate focal blooms**.

The 2019 HAB was reported on and monitored daily with focus on afflicted fish farm localities

by the Norwegian Directorate of Fisheries (Karlsen et al., 2019). The results obtained from our

amplicon-based sequencing approach are expected to be more complex and distinct from the

concurrent morphology-based taxonomy and cell counting surveys. However, our results share

some reassuring similarities to other reports, such as the observation that high focal taxon

(OLI16029 as representative ASVs for *C. leadbeateri* UiO-035 unialgal strain) abundance was

found in Balsfjord, which corresponds with microscopy enumerations conducted by Institute of

Marine Research and Akvaplan-Niva (Karlsen et al., 2019; Grann-Meyer, 2020). The current

study did not capture the onset of the HAB due to a reactive sampling effort to an event that is

(at least today) impossible to predict.

According to the prevailing consensus based on reported fish mortality, the toxic bloom

of the focal taxon initiated in Ofotfjord. The last report on fish mortality in Ofotfjord was dated

four days before the sample in Ballangen was taken (Karlsen et al., 2019), thus there was,

presumably, a mismatch between the bloom peak and time of sampling. We do not have data

that represents the community composition during the first reports of fish mortality. This may

explain why we found a surprising high disparity between expected and observed microbial

community composition at Ballangen, which did not correspond with any other sampled

microbial community composition (Figure 1). The microeukaryotic community was dominated

by diatoms, nearly exclusively composed by *Skeletonema*. A similar shift from *C. leadbeateri*

presence to absence with concurrent predominance of diatoms was observed during the 1991

HAB around Ofotfjord and surrounding areas (Hegseth and Eilertsen, 1991; Rey and Aure,

1991). Correspondingly, the sampled prokaryotic community composition in Ballangen does

not presumably represent either the situation with *C. leadbeateri* as a disruptive causative agent

as several genera belonging to the observed predominant bacterial order *Alteromonadales* have

been found to associate with *Skeletonema* sp. (Deng et al., 2021).

*C. leadbeateri* and presumably the cells assigned to the focal taxon in our sequencing

analysis are common and natural members of phytoplankton community along the entire

northern Norwegian coast, although its spatial and interannual variation has not been well

described since the 1991 HAB, as only two following years of monitoring was conducted in

area south and north from Ofotfjord (Hegseth and Eilertsen, 1991; Heidal and Mohus, 1995).

The present study, however, provides evidence for high spatial variation and non-cell-density-

dependent toxicity. As mentioned, the high abundance of our designated focal taxon in the

Balsfjord area was corroborated by manual cell counts and revealed by 18S rRNA amplicon

sequencing samples in this study (Figure 1). This was two weeks after the first reported fish

mortalities from Ofotfjord (Karlsen et al., 2019). Thus, this raises the question whether a bloom

of focal taxon commenced locally in Balsfjord. Due to limited connectivity, it is likely that *C.*

*leadbeateri* blooms developed independently in both fjords since its relative abundance was

low in FSS station (Figure 2c) and no afflicted fish farms were reported near the FSS station,

despite of the affect on fish farms in the area north of Ofotfjord (Karlsen et al., 2019).

Interestingly, despite high abundance of the focal taxon and effective advection-driven

dispersal of cells from the inner part of Balsfjord towards GSSS and Tussøya stations, only one

fish farm near Tussøya was afflicted. However, this farm experienced limited impacts, and its

relation to the HAB remained somewhat questionable (Karlsen et al., 2019). This observation

also lends some indications to the idea that there may have been at least two distinct

physiological states between these two *C. leadbeateri* blooms, and that only one of them was

"harmful".

**Localized microbial communities.**

One of the major questions that drove this study was to ask how the marine microbiome

composition compared across locations during the HAB time series. The results revealed

distinct communities between sample locations, each possessing different predominating

bacterial and eukaryotic microbial groups (Figure 1 and Figure 2). This observation on spatial

heterogeneity provides new information on the development of the late bloom phase within an

interconnected coastal area (Wassmann et al., 1996). GSSS was the only station where

*Phaeocystis* abundance exceeded that of the focal taxon. In fact, the relative abundance of

*Phaeocystis* was less than 1 % at all other stations, which was an unexpected result given

previously reported ubiquity and prolonged prevalence (March-August) in this region (Eilertsen

et al., 1981; Degerlund and Eilertsen, 2010).

Although the predominant microalgae groups found in the two time series stations were different (FSS: cryptophytes, *Leptocylindrus*, heterotrophic nanoflagellates; GSSS; dinoflagellates, prymnesiophyte, *Thalassiosira*), the most abundant bacterial taxa – i.e. orders *Flavobacteriales*, *Rhodobacterales* and *Pseudomonadales* – were essentially consistent. Previously reported studies have also shown that members belonging to these orders commonly accompany microalgae in marine environments, except in the case of *Teleaulax* sp. and *C. leadbeateri,* which may be due to lack of previous records from events such as the 1991 HAB of northern Norway (Teeling et al., 2016; Ajani et al., 2018; Zhou et al., 2018; Aalto et al., 2022). The bacterial community compositions obtained from Middle-Balsfjord and South-Tromsø stations where distinct and corresponded to the highest observed relative abundance of the focal taxon. Specifically, they harboured high relative abundances of SAR11 (mainly clade Ia) and low contributions of *Flavobacteriales,* which is interesting given that SAR11 tend to be negatively correlated with copiotrophic bacterial groups before and after the primary phytoplankton bloom (Needham and Fuhrman, 2016; Teeling et al., 2016). However, the single time point stations present a major limitation for interpreting the HAB microbiome, given that the time series stations (FSS and GSSS) showed pronounced temporal variability during the event.

**Temporal community dynamics.**

Another major aim of this study was to identify the temporal patterns of the HAB associated microbiome and attempt to determine which taxa co-bloomed with the focal ASV. We found that most of the microeukaryotic and bacterial taxa either co-bloomed or inversely co-bloomed with the focal ASV – i.e. the clusters that showed significant positive or negative correlation with the focal ASV also had the highest taxonomic richness (Figure 4 and Figure 5). There has been evidence that the most abundant taxa undergo stronger shifts in their abundance than rare

taxa, which corresponds with our results where the most abundant bacterial orders and microeukaryotic classes were mainly found within the clusters of highest temporal variability (Lindh et al., 2015). However, the high number of taxa in these clusters is somewhat surprising with regard of previous findings of highly coherent communities with well-defined groups of interacting organisms (Martin-Platero et al., 2018). It is likely that the prescribed temporal (sampling) resolution of this study was too coarse to capture the true frequency of taxonomic turnover, which can be very fast (Needham and Fuhrman, 2016; Martin-Platero et al., 2018). However, when an unpredicted and regionally large (250 km wide) HAB occurs, it poses major logistical challenges to react quickly and access the sample locations with frequency that would be the most optimal with biological perspective. The choice of optimal sampling locations is also limited by immediate accessibility to sampling resources, such as suitable water craft.

**The environment.**

The third major question of this study was related to the abiotic environment and how it may have influenced the microbiome composition and presence of the focal taxon. The HAB – as inferred from relative abundance of the focal taxon – was stronger at GSSS than at FSS, yet it occurred concurrently at both stations. The strong decrease in focal taxon abundance in the middle of time series coincided, especially at GSSS, with a change in community composition. However, this was evidenced poorly by environmental factors at both stations. Intriguingly, these results are in accordance with previous findings suggesting lower importance of continuous environmental factors such as temperature, inorganic nutrients and chlorophyll *a* on a change in community composition after seasonal bloom initiation than biotic interactions (Lindh et al., 2015; Needham and Fuhrman, 2016). Whereas, a strong periodical physical forcing (wind, turbulence) can cause a change of entire community (Martin-Platero et al., 2018). The measured water column properties did not indicate strong vertical mixing. However, the

surface water became somewhat more homogenous after the first week in the time series at

GSSS as revealed from CTD-profiles (Supplementary Figure S2).

The difference in microeukaryotic composition between the two time series stations

may be related to environmental factors as revealed by dbRDA analysis (Figure 3). The

phosphate concentration was low at both stations and in addition the $NO_3^- + NO_2^-$ level was low

in FSS, which potentially promoted the development of communities of heterotrophic and

mixotrophic protists. Several MAST-clades (e.g., MAST-1C as dominant in FSS) are known to

be bacterivores and *Teleaulax* has been determined to be mixotrophic, feeding heterotrophically

on bacteria, thus relaxing the need for inorganic nutrients (Massana et al., 2006; Du Yoo et al.,

2017). It is difficult to draw a link between nutrients and the focal taxon with the current data

due to lack of nutrient measurements from stations with high focal taxon abundance such as

Middle-Balsfjord and South-Tromsø. Despite the nutrient information, a plausible connection

would be unlikely as even a 14-year survey in southern Norway on *Chrysochromulina* spp. cell

and nutrients concentration did not provide a clear connection (Dahl et al., 2005). However,

there are indications that *Chrysochromulina* spp. thrive in high N:P ratios during low phosphate

concentration which potentially promotes toxicity if the condition becomes phosphate limited

(Dahl et al., 2005; Edvardsen and Imai, 2006).

**CONCLUSION**

A summary of investigations of the 1991 *Chrysochromulina leadbeateri* related harmful algal

bloom in northern Norway writes that it would not have been possible to predict such an

incident but it would be necessary to launch a coastal monitoring program (Rey, 1991). Three

decades later, we fully agree with the latter statement. We certainly have a long way to go in

order to predict how ecological principles underpin this sporadic event in a highly variable

coastal area and it is difficult to get there by performing investigations that simply react to only

those blooms that have acute economic impacts. It is also difficult to plan to observe such a bloom that happens so infrequently. It has become highly evident that this type of reoccurring bloom is extremely destructive because it causes substantial losses for the fish farmers. In 1991 the fish kill damage linked to *C. leadbeateri* was 742 tonnes with a corresponding direct short term economic cost of 3.5 million USD, while in 2019 these values were increased to 14 500 tonnes and more than 100 million USD (Karlson et al., 2021). Thus, it is expected these numbers will become even higher in the future with increasing aquaculture industry, and there is no reason to believe that this type of HAB will not reoccur. It is challenging to draw any conclusions about exact causation behind the 2019 HAB, especially since the natural bloom cycles of *C. leadbeateri* are not well understood. Therefore, this study relying on molecular-based methods provides a first insight on the spatial and temporal variability of the *C. leadbeateri*-like focal taxon together with its associated marine microbiome. It also supports future molecular-based studies by providing a genetic link with currently available taxonomic annotation between current and year 1991 *C. leadbeateri*. Otherwise the focal taxon may be overlooked due to database limitations – an issue also present in this study. We found that most of the taxa – including the most abundant members of prokaryotes and microeukaryotes and the focal taxon – underwent strong and rapid variability in their succession dynamics. These temporal dynamics were poorly connected to the environmental conditions measured, suggesting that other factors such as biological interactions might drive the late bloom dynamics.

## AUTHOR CONTRIBUTIONS

NJA and HCB conceptualized the study. NJA, EG-M, JBS, LD, CJH and RI carried out the sampling. NJA, HS, SK and SP performed lab work and processed amplicon sequence analyses.

NJA and HCB conducted data analysis and NJA wrote the manuscript draft. All authors

contributed to the final version.

**ACKNOWLEDGMENTS**

We greatly appreciate the logistical support by staff at Lerøy Aurora´s aquaculture facility at

Solheim. We also acknowledge the collaboration between Finnfjord AS and UiT-The Arctic

University of Norway for lending sampling infrastructure and support. We would like thank

Åge Mohus and Vigdis Tverberg at Nord University for their help with sample coordination in

southern portion of the HAB affected area.

**FUNDING**

This study was funded directly by the Faculty of Biosciences, Fisheries and Economics at UiT

– The Arctic University of Norway as the "2019 Harmful Algae Rapid Response Plan". The

authors would like to specifically thank Terje Aspen, Terje Martinussen and Kathrine Tveiterås

for providing rapid support and guidance.

**CONFLICT OF INTEREST**

The authors have no conflicts of interest to declare.

**REFERENCES CITED**

Aalto, N.J., Campbell, K., Eilertsen, H.C., and Bernstein, H.C. (2021). Drivers of atmosphere
707             ocean CO2 flux in northern Norwegian fjords. *Frontiers in Marine Science*, 841. doi:
708             10.3389/fmars.2021.692093.
709     Aalto, N.J., Schweitzer, H., Krsmanovic, S., Campbell, K., and Bernstein, H.C. (2022). Diversity
710             and selection of surface marine microbiomes in the Atlantic-influenced Arctic.
711             *Frontiers in Microbiology*. doi: 10.3389/fmicb.2022.892634/abstract.
712     Abdi, H., and Williams, L.J. (2010). Tukey's honestly significant difference (HSD) test.
713             *Encyclopedia of research design* 3(1), 1-5.
714     Ajani, P.A., Kahlke, T., Siboni, N., Carney, R., Murray, S.A., and Seymour, J.R. (2018). The
715             Microbiome of the Cosmopolitan Diatom Leptocylindrus Reveals Significant Spatial

and Temporal Variability. *Frontiers in Microbiology* 9. doi: 10.3389/fmicb.2018.02758.

Amaral-Zettler, L.A., McCliment, E.A., Ducklow, H.W., and Huse, S.M. (2009). A method for studying protistan diversity using massively parallel sequencing of V9 hypervariable regions of small-subunit ribosomal RNA genes. *PloS one* 4(7), e6372.

Anderson, D.M., Glibert, P.M., and Burkholder, J.M. (2002). Harmful algal blooms and eutrophication: nutrient sources, composition, and consequences. *Estuaries* 25(4), 704-726.

Anderson, M.J. (2001). A new method for non-parametric multivariate analysis of variance. *Austral ecology* 26(1), 32-46.

Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., et al. (2010). QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* 7(5), 335-336.

Cirri, E., and Pohnert, G. (2019). Algae– bacteria interactions that balance the planktonic microbiome. *New Phytologist* 223(1), 100-106.

Coenen, A.R., Hu, S.K., Luo, E., Muratore, D., and Weitz, J.S. (2020). A primer for microbiome time-series analysis. *Frontiers in genetics* 11, 310.

Dahl, E., Bagøien, E., Edvardsen, B., and Stenseth, N.C. (2005). The dynamics of Chrysochromulina species in the Skagerrak in relation to environmental conditions. *Journal of Sea Research* 54(1), 15-24.

Davidson, K., Gowen, R.J., Tett, P., Bresnan, E., Harrison, P.J., McKinney, A., et al. (2012). Harmful algal blooms: How strong is the evidence that nutrient ratios and forms influence their occurrence? *Estuarine, Coastal and Shelf Science* 115, 399-413. doi: https://doi.org/10.1016/j.ecss.2012.09.019.

Degerlund, M., and Eilertsen, H.C. (2010). Main species characteristics of phytoplankton spring blooms in NE Atlantic and Arctic waters (68–80 N). *Estuaries and coasts* 33(2), 242-269.

Deng, Z., Chen, S., Zhang, P., Zhang, X., Adams, J.M., Luo, Q., et al. (2021). Dynamics of Free-Living and Attached Bacterial Assemblages in Skeletonema sp. Diatom Cultures at Elevated Temperatures. *Frontiers in Marine Science* 8. doi: 10.3389/fmars.2021.626207.

Doucette, G.J., McGovern, E.R., and Babinchak, J.A. (1999). ALGICIDAL BACTERIA ACTIVE AGAINST GYMNODINIUM BREVE (DINOPHYCEAE). I. BACTERIAL ISOLATION AND CHARACTERIZATION OF KILLING ACTIVITY1, 3. *Journal of Phycology* 35(6), 1447-1454.

Du Yoo, Y., Seong, K.A., Jeong, H.J., Yih, W., Rho, J.-R., Nam, S.W., et al. (2017). Mixotrophy in the marine red-tide cryptophyte Teleaulax amphioxeia and ingestion and grazing impact of cryptophytes on natural populations of bacteria in Korean coastal waters. *Harmful Algae* 68, 105-117.

Edvardsen, B., Eikrem, W., Throndsen, J., Saez, A.G., Probert, I., and Medlin, L.K. (2011). Ribosomal DNA phylogenies and a morphological revision provide the basis for a revised taxonomy of the Prymnesiales (Haptophyta). *European journal of phycology* 46(3), 202-228.

Edvardsen, B., and Imai, I. (2006). "The ecology of harmful flagellates within Prymnesiophyceae and Raphidophyceae," in *Ecology of harmful algae*. Springer), 67-79.

Edvardsen, B., and Paasche, E. (1998). Bloom dynamics and physiology of Prymnesium and Chrysochromulina. *NATO ASI SERIES G ECOLOGICAL SCIENCES* 41, 193-208.

Eikrem, W., and Throndsen, J. (1998). Morphology of Chrysochromulina leadbeateri
(Prymnesiophyceae) from northern Norway. *Phycologia* 37(4)**,** 292-299.
Eilertsen, H.C., Falk-Petersen, S., Hopkins, C., and Tande, K. (1981). Ecological investigations
on the plankton community of Balsfjorden, northern Norway: program for the
project, study area, topography, and physical environment. *Sarsia* 66(1)**,** 25-34.
Eilertsen, H.C., and Frantzen, S. (2007). Phytoplankton from two sub-Arctic fjords in northern
Norway 2002–2004: I. Seasonal variations in chlorophyll a and bloom dynamics.
*Marine Biology Research* 3(5)**,** 319-332.
Eilertsen, H.C., and Taasen, J. (1984). Investigations on the plankton community of
Balsfjorden, northern Norway. The phytoplankton 1976–1978. Environmental
factors, dynamics of growth, and primary production. *Sarsia* 69(1)**,** 1-15.
Estep, K. (1984). Chloroplast containing microflagellates in natural populations of north
Atlantic nanoplankton, their identification and distribution; including a description of
5 new species of Chrysochromulina (Prymnesiophyceae). *Protistologica* 20**,** 613-634.
Fuhrman, J.A., Cram, J.A., and Needham, D.M. (2015). Marine microbial community
dynamics and their ecological interpretation. *Nature Reviews Microbiology* 13(3)**,**
133-146.
Gilbert, J.A., Meyer, F., Antonopoulos, D., Balaji, P., Brown, C.T., Brown, C.T., et al. (2010).
Meeting report: the terabase metagenomics workshop and the vision of an Earth
microbiome project. *Stand Genomic Sci* 3(3)**,** 243.
Grann-Meyer, E. (2020). *Chrysochromulina leadbeateri - Understanding the presumed causal*
*agent behind the harmful algal bloom of 2019.* Master's thesis, UiT - Arctic University
of Norway.
Green, D.H., Llewellyn, L.E., Negri, A.P., Blackburn, S.I., and Bolch, C.J. (2004). Phylogenetic
and functional diversity of the cultivable bacterial community associated with the
paralytic shellfish poisoning dinoflagellate Gymnodinium catenatum. *FEMS*
*Microbiology Ecology* 47(3)**,** 345-357.
Guillard, R.R. (1975). "Culture of phytoplankton for feeding marine invertebrates," in *Culture*
*of marine invertebrate animals*. Springer), 29-60.
Hallegraeff, G.M., Anderson, D.M., Belin, C., Bottein, M.-Y.D., Bresnan, E., Chinain, M., et al.
(2021). Perceived global increase in algal blooms is attributable to intensified
monitoring and emerging bloom impacts. *Communications Earth & Environment* 2(1)**,**
1-10.
Hegseth, E.N., and Eilertsen, H.C. (1991). The Chrysochromulina leadbeateri bloom in Troms
May/June 1991. Development and causes. . *Fisk. Hav.* 3**,** 45-61.
Heidal, K., and Mohus, A. (1995). The toxic Chrysochromulina-- salmon disaster of 1991 in
northern Norway with some follow-up monitoring records of 1992 and-93.
*LAVOISIER, PARIS(FRANCE).* 163-168.
Karlsen, K.M., Robertsen, R., and Hersoug, B. (2019). Mapping the course of events and
preparedness for algal blooms in spring 2019 - Astafjorden, Ofotfjorden, Vestfjorden
og Tysfjorden. *Nofima AS* 29**,** 1-44.
Karlson, B., Andersen, P., Arneborg, L., Cembella, A., Eikrem, W., John, U., et al. (2021).
Harmful algal blooms and their effects in coastal seas of Northern Europe. *Harmful*
*Algae* 102**,** 101989.
Kaufman, L., and Rousseeuw, P.J. (2009). *Finding groups in data: an introduction to cluster*
*analysis.* New York: John Wiley & Sons.

Lindh, M.V., Sjöstedt, J., Andersson, A.F., Baltar, F., Hugerth, L.W., Lundin, D., et al. (2015).
Disentangling seasonal bacterioplankton population dynamics by high-frequency
sampling. *Environmental microbiology* 17(7)**,** 2459-2476.
Liu, C., Cui, Y., Li, X., and Yao, M. (2021). microeco: an R package for data mining in microbial
community ecology. *FEMS Microbiology Ecology* 97(2)**,** fiaa255.
Lord, E., Willems, M., Lapointe, F.-J., and Makarenkov, V. (2017). Using the stability of
objects to determine the number of clusters in datasets. *Information Sciences* 393**,**
29-46.
Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome biology* 15(12)**,** 1-21.
Lozupone, C., Lladser, M.E., Knights, D., Stombaugh, J., and Knight, R. (2011). UniFrac: an
effective distance metric for microbial community comparison. *The ISME journal* 5(2)**,**
169-172.
Martin-Platero, A.M., Cleary, B., Kauffman, K., Preheim, S.P., McGillicuddy, D.J., Alm, E.J., et
al. (2018). High resolution time series reveals cohesive but short-lived communities in
coastal plankton. *Nature communications* 9(1)**,** 1-11.
Massana, R., Terrado, R., Forn, I., Lovejoy, C., and Pedrós-Alió, C. (2006). Distribution and
abundance of uncultured heterotrophic flagellates in the world oceans.
*Environmental microbiology* 8(9)**,** 1515-1522.
Mayali, X., and Azam, F. (2004). Algicidal bacteria in the sea and their impact on algal blooms
1. *Journal of Eukaryotic Microbiology* 51(2)**,** 139-144.
Meldahl, A.S., Edvardsen, B., and Fonnum, F. (1994). Toxicity of four potentially ichthyotoxic
marine phytoflagellates determined by four different test methods. *Journal of*
*Toxicology and Environmental Health, Part A Current Issues* 42(3)**,** 289-301.
Meyer, N., Bigalke, A., Kaulfuß, A., and Pohnert, G. (2017). Strategies and ecological roles of
algicidal bacteria. *FEMS Microbiology Reviews* 41(6)**,** 880-899.
Needham, D.M., and Fuhrman, J.A. (2016). Pronounced daily succession of phytoplankton,
archaea and bacteria following a spring bloom. *Nature Microbiology* 1(4)**,** 16005. doi:
10.1038/nmicrobiol.2016.5.
Nielsen, T.G., Kiørboe, T., and Bjørnsen, P.K. (1990). Effects of a Chrysochromulina polylepis
subsurface bloom on the planktonic community. *Marine Ecology Progress Series***,** 21-
35.
Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'hara, R., et al. (2013).
Package 'vegan'. *Community ecology package, version* 2(9).
Parte, A.C., Carbasse, J.S., Meier-Kolthoff, J.P., Reimer, L.C., and Göker, M. (2020). List of
Prokaryotic names with Standing in Nomenclature (LPSN) moves to the DSMZ.
*International Journal of Systematic and Evolutionary Microbiology* 70(11)**,** 5607.
Peralta-Ferriz, C., and Woodgate, R.A. (2015). Seasonal and interannual variability of pan-
Arctic surface mixed layer properties from 1979 to 2012 from hydrographic data, and
the dominance of stratification for multiyear mixed layer depth shoaling. *Progress in*
*Oceanography* 134**,** 19-53.
R Core Team (2021). R: A language and environment for statistical computing, R Foundation
for Statistical Computing.
Reigstad, M., and Wassmann, P. (1996). Importance of advection for pelagic-benthic
coupling in north Norwegian fjords. *Sarsia* 80(4)**,** 245-257.
Rey, F. (1991). The Chrysochromulina leadbeateri bloom in Vestfjorden, north Norway, May-
June 1991: Proceedings from a scientific working meeting. *Fisk. Hav.* 3**,** 1-122.

Rey, F., and Aure, J. (1991). The *Chrysochromulina leadbeateri* bloom in the Vestfjord, north
Norway, May-June 1991. Environmental conditions and possible causes. *Fisk. Hav.* 3**,**
13-32.
Statistics Norway (2020). *Aquaculture 2019* [Online]. Statistics Norway. Available:
https://www.ssb.no/jord-skog-jakt-og-fiskeri/statistikker/fiskeoppdrett [Accessed
23.04 2022].
Teeling, H., Fuchs, B.M., Bennke, C.M., Krueger, K., Chafee, M., Kappelmann, L., et al. (2016).
Recurring patterns in bacterioplankton dynamics during coastal spring algae blooms.
*elife* 5**,** e11888.
Uribe, P., and Espejo, R.T. (2003). Effect of associated bacteria on the growth and toxicity of
Alexandrium catenella. *Applied and environmental microbiology* 69(1)**,** 659-662.
Uronen, P., Kuuppo, P., Legrand, C., and Tamminen, T. (2007). Allelopathic Effects of Toxic
Haptophyte Prymnesium parvum Lead to Release of Dissolved Organic Carbon and
Increase in Bacterial Biomass. *Microbial Ecology* 54(1)**,** 183-193. doi:
10.1007/s00248-006-9188-8.
Uronen, P., Lehtinen, S., Legrand, C., Kuuppo, P., and Tamminen, T. (2005). Haemolytic
activity and allelopathy of the haptophyte Prymnesium parvum in nutrient-limited
and balanced growth conditions. *Marine Ecology Progress Series* 299**,** 137-148.
Wassmann, P., Svendsen, H., Keck, A., and Reigstad, M. (1996). Selected aspects of the
physical oceanography and particle fluxes in fjords of northern Norway. *Journal of*
*Marine Systems* 8(1-2)**,** 53-71.
Zhou, J., Richlen, M.L., Sehein, T.R., Kulis, D.M., Anderson, D.M., and Cai, Z. (2018). Microbial
Community Structure and Associations During a Marine Dinoflagellate Bloom.
*Frontiers in Microbiology* 9. doi: 10.3389/fmicb.2018.01201.

880
881

**SUPPLEMENTARY MATERIAL**

895 **Microbial community dynamics during a harmful *Chrysochromulina leadbeateri* bloom**

897 Nerea J. Aalto[1,2], Hannah Schweitzer[1,2], Erlend Grann-Meyer[1], Stina Krsmanovic[1], Jon Brage
898 Svenning[1], Lars Dalheim[1], Sebastian Petters[1], Richard Ingebrigtsen[1], Chris J. Hulatt[3], Hans C.
899 Bernstein[1,2]*

**Affiliations:**

*1. Norwegian College of Fisheries Sciences, UiT - The Arctic University of Norway, Tromsø, Norway*

*2. The Arctic Centre for Sustainable Energy, UiT - The Arctic University of Norway, Tromsø, Norway*

*3. Faculty of Biosciences and Aquaculture, Nord University, Bodø, Norway*

**\*Correspondence**: Hans C. Bernstein, UiT-The Arctic University of Norway, BFE-fak., Postboks 6050 Langnes, 9037 Tromsø, NORWAY*; Phone: +47 776 46 114;  Email:* Hans.C.Bernstein@uit.no

918 **Supplementary Table S1.** Station positions, sample dates, and corresponding sample day and
919 week during time series sample campaign (May 27th = Sample day 0 and June 21st = Sample
920 day 25).

| Station | Position | Sample date | Sample day/Sample week |
|---|---|---|---|
| FSS | 69.19 N, 18.03 E | 27.05.2019 | 0/0 |
| | | 30.05.2019 | 3/0 |
| | | 03.06.2019 | 7/1 |
| | | 11.06.2019 | 15/2 |
| | | 13.06.2019 | 17/2 |
| | | 17.06.2019 | 21/3 |
| | | 20.06.2019 | 24/3 |
| GSSS | 69.80 N, 19.36 E | 28.05.2019 | 1/0 |
| | | 31.05.2019 | 4/0 |
| | | 04.06.2019 | 8/1 |
| | | 07.06.2019 | 11/1 |
| | | 12.06.2019 | 16/2 |
| | | 14.06.2019 | 18/2 |
| | | 18.06.2019 | 22/2 |
| | | 21.06.2019 | 25/3 |
| South-Tromsø | 69.60 N, 18.80 E | 27.05.2019 | 0/0 |
| Middle-Balsfjord | 69.40 N, 19.20 E | 29.05.2019 | 2/0 |
| Tussøy | 69.66 N, 18.18 E | 06.06.2019 | 10/1 |
| Ballangen | 68.41 N, 16.83 E | 05.06.2019 | 9/1 |
| Middle-Sagfjord | 67.98 N, 15.75 E | 14.06.2019 | 18/2 |
| Veggfjell | 67.96 N, 15.80 E | 14.06.2019 | 18/2 |
| Anevik | 68.00 N, 15.14 E | 14.06.2019 | 18/2 |
| Dyping | 67.92 N, 15.34 E | 14.06.2019 | 18/2 |

931 **Supplementary Table S2.** The number of classified ASVs and sequencing depth per sample
932 and data set before and after removal of unwanted taxa. Time series sample set includes
933 stations FSS and GSSS; Single-time-point sample set includes stations Tussøya, South-
934 Tromsø, Middle-Balsfjord, Ballangen, Middle-Sagfjord, Veggfjell, Anevik and Dyping;
935 Unialgal control sample set includes *Chrysochromulina leadbeateri* UiO-035 strain culture.
936

| Condition | Sample set | Data set | # ASVs | Sample depth |
|---|---|---|---|---|
| All taxa | Time series | 16S | 2761 | 54160–217118 |
| | Time series | 18S | 7500 | 72831–34835 |
| | Single-time-point | 16S | 1414 | 83315–168168 |
| | Single-time-point | 18S | 3687 | 143296–469129 |
| | Unialgal control (UiO-035) | 18S | 123 | 134563–273163 |
| Removed unwanted taxa | Time series | 16S | 2644 | 53962–214722 |
| | Time series | 18S | 2674 | 56789–331887 |
| | Single-time-point | 16S | 1368 | 83253–167400 |
| | Single-time-point | 18S | 1682 | 118101–451910 |
| | Unialgal control (UiO-035) | 18S | 15 | 125686–265853 |

**Supplementary Figure S1.** Surface water profiles of temperature and salinity obtained from CTD-casts in FSS time series station. The dashed line denotes calculated mixed layer depth (MLD) based on density gradient and solid line marks the depth of sampled microbial communities (10 m). Note missing CTD-casts in sample days of 15 and 17.

**Supplementary Figure S2.** Surface water profiles of temperature and salinity obtained from CTD-casts in GSSS time series station. The dashed line denotes calculated mixed layer depth (MLD) based on density gradient and solid line marks the depth of sampled microbial communities (10 m). Note missing CTD-casts in sample days of 11 and 16.

954 **Supplementary Figure S2 continues.**

**Supplementary Figure S3.** Number of amplicon sequence reads of each ASV assigned to

OLI16029, in unialgal *Chrysochromulina leadbeateri* UiO-035 control culture.

**Supplementary Figure S4.** Relative abundance of the most abundant ASV

(*24a92740c5af4cd5a5ac0db70830fbc0*, referred as a focal ASV) in unialgal

*Chrysochromulina leadbeateri* UiO-035 control culture among all the ASVs assigned to

OLI16029.

**Supplementary Figure S5.** The Calinski-Harabasz index (y-axis, C-H Stat) was used to
determine the number of K-medoids clusters. C-H Stat is a normalized ratio of the inter and
intra cluster variance calculated for each sample-specific K-medoids cluster. High C-H Stat
value indicates more uniquely portioned clusters. A) Three clusters were chosen to represent
16S ASVs from FSS time series. B) Three clusters were chosen to represent 16S ASVs from
the from GSSS time series. C) Four clusters were chosen to represent 18S ASVs from FSS
time series. D) Three clusters were chosen to represent 18S ASVs from the from GSSS time
series.

987 **Supplementary Table 3.** Results of perMANOVA pariwise comparison between 1) time
988 series sample stations, 2) sample weeks and 3) sample days within each sample station for
989 prokaryotic and microeukaryotic communities, respectively. Unweighted UniFrac was used as
990 a distance metric, and number of permutations was 999.
991

| Groups | R2 | r | p.value | Data type | Station |
|---|---|---|---|---|---|
| FSS vs GSSS | 0.3 | 0.54 | 0.001 | 16S | |
| FSS vs GSSS | 0.46 | 0.68 | 0.001 | 18S | |
| Week_0 vs Week_1 | 0.22 | 0.47 | 0.001 | 16S | FSS |
| Week_0 vs Week_2 | 0.16 | 0.39 | 0.002 | 16S | FSS |
| Week_0 vs Week_3 | 0.21 | 0.45 | 0.001 | 16S | FSS |
| Week_1 vs Week_2 | 0.25 | 0.5 | 0.001 | 16S | FSS |
| Week_1 vs Week_3 | 0.17 | 0.41 | 0.001 | 16S | FSS |
| Week_2 vs Week_3 | 0.13 | 0.36 | 0.001 | 16S | FSS |
| Week_0 vs Week_1 | 0.08 | 0.28 | 0.013 | 16S | GSSS |
| Week_0 vs Week_2 | 0.16 | 0.4 | 0.001 | 16S | GSSS |
| Week_0 vs Week_3 | 0.21 | 0.46 | 0.001 | 16S | GSSS |
| Week_1 vs Week_2 | 0.14 | 0.38 | 0.001 | 16S | GSSS |
| Week_1 vs Week_3 | 0.23 | 0.48 | 0.001 | 16S | GSSS |
| Week_2 vs Week_3 | 0.11 | 0.32 | 0.003 | 16S | GSSS |
| Week_0 vs Week_1 | 0.38 | 0.61 | 0.002 | 18S | FSS |
| Week_0 vs Week_2 | 0.33 | 0.58 | 0.001 | 18S | FSS |
| Week_0 vs Week_3 | 0.34 | 0.58 | 0.001 | 18S | FSS |
| Week_1 vs Week_2 | 0.42 | 0.65 | 0.002 | 18S | FSS |
| Week_1 vs Week_3 | 0.27 | 0.51 | 0.001 | 18S | FSS |
| Week_2 vs Week_3 | 0.24 | 0.49 | 0.001 | 18S | FSS |
| Week_0 vs Week_1 | 0.08 | 0.28 | 0.024 | 18S | GSSS |
| Week_0 vs Week_2 | 0.16 | 0.4 | 0.001 | 18S | GSSS |
| Week_0 vs Week_3 | 0.21 | 0.46 | 0.001 | 18S | GSSS |
| Week_1 vs Week_2 | 0.14 | 0.38 | 0.003 | 18S | GSSS |
| Week_1 vs Week_3 | 0.23 | 0.48 | 0.001 | 18S | GSSS |
| Week_2 vs Week_3 | 0.11 | 0.32 | 0.003 | 18S | GSSS |
| 0 vs 3 | 0.11 | 0.33 | 0.356 | 16S | FSS |
| 0 vs 7 | 0.32 | 0.57 | 0.003 | 16S | FSS |
| 0 vs 15 | 0.24 | 0.49 | 0.01 | 16S | FSS |
| 0 vs 17 | 0.27 | 0.52 | 0.006 | 16S | FSS |
| 0 vs 21 | 0.29 | 0.54 | 0.001 | 16S | FSS |
| 0 vs 24 | 0.36 | 0.6 | 0.007 | 16S | FSS |
| 3 vs 7 | 0.25 | 0.5 | 0.009 | 16S | FSS |
| 3 vs 15 | 0.21 | 0.46 | 0.006 | 16S | FSS |
| 3 vs 17 | 0.24 | 0.49 | 0.007 | 16S | FSS |
| 3 vs 21 | 0.25 | 0.5 | 0.005 | 16S | FSS |
| 3 vs 24 | 0.29 | 0.54 | 0.005 | 16S | FSS |
| 7 vs 15 | 0.36 | 0.6 | 0.004 | 16S | FSS |

| | | | | | |
|---|---|---|---|---|---|
| 7 vs 17 | 0.29 | 0.54 | 0.003 | 16S | FSS |
| 7 vs 21 | 0.29 | 0.54 | 0.006 | 16S | FSS |
| 7 vs 24 | 0.23 | 0.48 | 0.004 | 16S | FSS |
| 15 vs 17 | 0.22 | 0.47 | 0.009 | 16S | FSS |
| 15 vs 21 | 0.22 | 0.46 | 0.004 | 16S | FSS |
| 15 vs 24 | 0.36 | 0.6 | 0.001 | 16S | FSS |
| 17 vs 21 | 0.16 | 0.4 | 0.01 | 16S | FSS |
| 17 vs 24 | 0.27 | 0.52 | 0.003 | 16S | FSS |
| 21 vs 24 | 0.25 | 0.5 | 0.004 | 16S | FSS |
| 1 vs 4 | 0.28 | 0.53 | 0.002 | 16S | GSSS |
| 1 vs 8 | 0.23 | 0.48 | 0.004 | 16S | GSSS |
| 1 vs 11 | 0.31 | 0.56 | 0.006 | 16S | GSSS |
| 1 vs 16 | 0.34 | 0.59 | 0.004 | 16S | GSSS |
| 1 vs 18 | 0.4 | 0.63 | 0.005 | 16S | GSSS |
| 1 vs 22 | 0.29 | 0.54 | 0.002 | 16S | GSSS |
| 1 vs 25 | 0.36 | 0.6 | 0.002 | 16S | GSSS |
| 4 vs 8 | 0.18 | 0.43 | 0.002 | 16S | GSSS |
| 4 vs 11 | 0.14 | 0.38 | 0.009 | 16S | GSSS |
| 4 vs 16 | 0.17 | 0.42 | 0.002 | 16S | GSSS |
| 4 vs 18 | 0.37 | 0.61 | 0.003 | 16S | GSSS |
| 4 vs 22 | 0.31 | 0.56 | 0.004 | 16S | GSSS |
| 4 vs 25 | 0.37 | 0.61 | 0.004 | 16S | GSSS |
| 8 vs 11 | 0.21 | 0.46 | 0.001 | 16S | GSSS |
| 8 vs 16 | 0.2 | 0.44 | 0.002 | 16S | GSSS |
| 8 vs 18 | 0.33 | 0.58 | 0.002 | 16S | GSSS |
| 8 vs 22 | 0.27 | 0.52 | 0.004 | 16S | GSSS |
| 8 vs 25 | 0.32 | 0.56 | 0.004 | 16S | GSSS |
| 11 vs 16 | 0.19 | 0.43 | 0.007 | 16S | GSSS |
| 11 vs 18 | 0.41 | 0.64 | 0.002 | 16S | GSSS |
| 11 vs 22 | 0.34 | 0.58 | 0.002 | 16S | GSSS |
| 11 vs 25 | 0.38 | 0.62 | 0.002 | 16S | GSSS |
| 16 vs 18 | 0.32 | 0.56 | 0.004 | 16S | GSSS |
| 16 vs 22 | 0.27 | 0.52 | 0.001 | 16S | GSSS |
| 16 vs 25 | 0.32 | 0.56 | 0.003 | 16S | GSSS |
| 18 vs 22 | 0.15 | 0.38 | 0.005 | 16S | GSSS |
| 18 vs 25 | 0.24 | 0.49 | 0.003 | 16S | GSSS |
| 22 vs 25 | 0.18 | 0.42 | 0.004 | 16S | GSSS |
| 0 vs 3 | 0.19 | 0.43 | 0.017 | 18S | FSS |
| 0 vs 7 | 0.48 | 0.69 | 0.002 | 18S | FSS |
| 0 vs 15 | 0.5 | 0.71 | 0.008 | 18S | FSS |
| 0 vs 17 | 0.47 | 0.69 | 0.009 | 18S | FSS |
| 0 vs 21 | 0.44 | 0.67 | 0.004 | 18S | FSS |
| 0 vs 24 | 0.56 | 0.75 | 0.001 | 18S | FSS |

| | | | | | |
|---|---|---|---|---|---|
| 3 vs 7 | 0.42 | 0.65 | 0.002 | 18S | FSS |
| 3 vs 15 | 0.44 | 0.66 | 0.003 | 18S | FSS |
| 3 vs 17 | 0.4 | 0.63 | 0.002 | 18S | FSS |
| 3 vs 21 | 0.37 | 0.61 | 0.001 | 18S | FSS |
| 3 vs 24 | 0.49 | 0.7 | 0.004 | 18S | FSS |
| 7 vs 15 | 0.54 | 0.74 | 0.006 | 18S | FSS |
| 7 vs 17 | 0.44 | 0.66 | 0.002 | 18S | FSS |
| 7 vs 21 | 0.41 | 0.64 | 0.005 | 18S | FSS |
| 7 vs 24 | 0.37 | 0.61 | 0.002 | 18S | FSS |
| 15 vs 17 | 0.31 | 0.56 | 0.007 | 18S | FSS |
| 15 vs 21 | 0.35 | 0.59 | 0.002 | 18S | FSS |
| 15 vs 24 | 0.54 | 0.74 | 0.002 | 18S | FSS |
| 17 vs 21 | 0.23 | 0.48 | 0.001 | 18S | FSS |
| 17 vs 24 | 0.44 | 0.66 | 0.004 | 18S | FSS |
| 21 vs 24 | 0.31 | 0.55 | 0.004 | 18S | FSS |
| 1 vs 4 | 0.28 | 0.53 | 0.002 | 18S | GSSS |
| 1 vs 8 | 0.23 | 0.48 | 0.001 | 18S | GSSS |
| 1 vs 11 | 0.31 | 0.56 | 0.004 | 18S | GSSS |
| 1 vs 16 | 0.34 | 0.59 | 0.003 | 18S | GSSS |
| 1 vs 18 | 0.4 | 0.63 | 0.002 | 18S | GSSS |
| 1 vs 22 | 0.29 | 0.54 | 0.003 | 18S | GSSS |
| 1 vs 25 | 0.36 | 0.6 | 0.007 | 18S | GSSS |
| 4 vs 8 | 0.18 | 0.43 | 0.002 | 18S | GSSS |
| 4 vs 11 | 0.14 | 0.38 | 0.008 | 18S | GSSS |
| 4 vs 16 | 0.17 | 0.42 | 0.009 | 18S | GSSS |
| 4 vs 18 | 0.37 | 0.61 | 0.001 | 18S | GSSS |
| 4 vs 22 | 0.31 | 0.56 | 0.003 | 18S | GSSS |
| 4 vs 25 | 0.37 | 0.61 | 0.005 | 18S | GSSS |
| 8 vs 11 | 0.21 | 0.46 | 0.002 | 18S | GSSS |
| 8 vs 16 | 0.2 | 0.44 | 0.005 | 18S | GSSS |
| 8 vs 18 | 0.33 | 0.58 | 0.003 | 18S | GSSS |
| 8 vs 22 | 0.27 | 0.52 | 0.005 | 18S | GSSS |
| 8 vs 25 | 0.32 | 0.56 | 0.004 | 18S | GSSS |
| 11 vs 16 | 0.19 | 0.43 | 0.012 | 18S | GSSS |
| 11 vs 18 | 0.41 | 0.64 | 0.001 | 18S | GSSS |
| 11 vs 22 | 0.34 | 0.58 | 0.004 | 18S | GSSS |
| 11 vs 25 | 0.38 | 0.62 | 0.002 | 18S | GSSS |
| 16 vs 18 | 0.32 | 0.56 | 0.003 | 18S | GSSS |
| 16 vs 22 | 0.27 | 0.52 | 0.006 | 18S | GSSS |
| 16 vs 25 | 0.32 | 0.56 | 0.002 | 18S | GSSS |
| 18 vs 22 | 0.15 | 0.38 | 0.004 | 18S | GSSS |
| 18 vs 25 | 0.24 | 0.49 | 0.006 | 18S | GSSS |
| 22 vs 25 | 0.18 | 0.42 | 0.004 | 18S | GSSS |

992 **Supplementary Table S4.** Summary of physical environmental measurements and analyzed
993 inorganic nutrients as well as the relative abundance of Focal taxon, OLI16029. Mean values
994 and (min, max) are shown.
995

| | Station | |
|---|---|---|
| | FSS | GSSS |
| Temperature (°C)[a] | 6.6 (5.4, 7.4) | 6.2 (5.3, 7.0) |
| Salinity[a] | 33.1 (32.5, 33.4) | 33.4 (33.2, 33.6) |
| MLD (m) | 3 (3.3) | 5 (3, 7) |
| $Si(OH)_4$ (µmol L$^{-1}$) | 1.24 (0.69, 1.87) | 1.56 (0.09, 3.00) |
| $NO_2^- + NO_3^-$ (µmol L$^{-1}$) | 0.18 (0.07, 0.41) | 0.43 (0.09, 1.20) |
| $PO_4^{3-}$ (µmol L$^{-1}$) | 0.06 (< 0.04, 0.12) | 0.07 (< 0.04, 0.39) |
| Focal taxon (%)[b] | 0.7 (0.4, 1.3) | 3.4 (0.02, 17.7) |

996 [a] Values at the depth of 10 m
997 [b] Relative abundance of ASVs assigned to OLI16029
998

1000

1001

1002

1003

1004

1005

**Supplementary Figure S6.** Alpha diversity as observed number of ASVs per sample set: single-time-point stations and time series stations.

1006
1007
1008

1009