Research article

TITLE

Where does the rainbow end? A case study of self-sustaining rainbow trout *Oncorhynchus mykiss* populations in tropical rivers

AUTHOR INFORMATION

Marie Nevoux, Amandine D. Marie, Julien Raitif, Jean-Luc Baglinière, Olivier Lorvelec, Jean-Marc Roussel

DECOD (Ecosystem Dynamics and Sustainability), INRAE, Institut Agro, IFREMER, Rennes, France

Corresponding author:

Marie Nevoux

marie.nevoux@inrae.fr

ORCID: 0000-0003-1451-7732

ABSTRACT

The great physiological, behavioral and morphological plasticity of salmonids contributed to the success of their introduction into a wide range of habitats. However, rainbow trout (*Oncorhynchus mykiss*) is the only salmonid species to live in the tropics. Like for any cold-water species, water temperature is assumed to be a major constraint to the presence and expansion of rainbow trout. In this study, we collected detailed records of the introduction history of rainbow trout at 35 sites across the island over 80 years, and we document the presence of self-sustainable trout at each site. We explored the linear relationship between water temperature and elevation.

KEYWORDS

Elevation, introduction, presence, salmonid, temperature

DECLARATIONS

Conflict of interest

The authors declare no conflict of interest.

INTRODUCTION

Native to the northern hemisphere, salmonid species have been introduced worldwide (Welcomme 1988; Crawford & Muir 2008; Lecomte *et al.* 2013), especially brown trout (*Salmo trutta*), brook trout (*Salvelinus fontinalis*), coho salmon (*Oncorhynchus kisutch*) and rainbow trout (*Oncorhynchus mykiss*). Their great physiological, behavioral and morphological plasticity contributes to the success of their introduction into a wide range of environmental conditions (Sauter *et al.* 2001). Since they are a cold-water species, however, successful introductions have been reported mainly at high latitudes and in temperate regions (Cowx 1997; Stanković *et al.* 2015; Koutsikos *et al.* 2019), since water temperature is a major constraint to their ontogeny and expansion. Among salmonids, rainbow trout has the widest latitudinal distribution in its native range (Fig. 1a), from 23-64°N (MacCrimmon 1971; Welcomme 1988; Crawford & Muir 2008). The population in the Sierra Madre Occidental Mountains of Mexico represents the southernmost native range of any salmonid (Behnke 2010), along with the Formosan landlocked salmon (*Oncorhynchus masou formosanus*) in Asia. These populations reflect rainbow trout's ability to survive under extreme thermal conditions (MacCrimmon 1971; Gall & Crandell 1992). Its optimal range is 10-16°C, but adults can endure peaks of 25°C (Raleigh 1984; Jalabert & Fostier 2010) and even up to 28-30°C (Matthews & Berg 1997; Sloat & Osterback 2013). In contrast, it reproduces in winter to meet requirements for lower temperatures. Spawning and successful egg development have been recorded at temperatures of 4-13°C (Morrison & Smith 1986; Pankhurst *et al.* 1996; Jalabert & Fostier 2010) and even up to 15.5°C (Scott & Crossman 1973). Above 20°C, however, the fecundity rate and embryonic survival decrease greatly (King *et al.* 2003).

Since the middle of the 20[th] century, rainbow trout has been massively introduced worldwide for aquaculture and recreational purposes (Welcomme 1988). According to the FAO (2003), at least 97 countries have introduced it (Fig. 1a), including 38 in the tropics (Table S1). To our knowledge, rainbow trout is the only salmonid species that lives in the tropics. Lethal temperature is the main reason that introductions have failed in Puerto Rico, the Philippines, Trinidad and Mauritius (MacCrimmon 1971). Nevertheless, the existence of self-sustaining populations (i.e. recruitment unsupported by and independent of continued human interactions, *sensu* Copp *et al.* (2005)) has been reported in nearly 70% of tropical countries where the species was introduced (Fig. 1b, Table S1). Tropical populations of rainbow trout are presumably found at high elevations, where the ambient temperature is cool (Therezien 1960; Péfaur & Sierra 1998; Ngugi & Green 2007; Vila *et al.* 2007; Kadye *et al.* 2013; Vimos *et al.* 2015), such as in the Nyanga Mountains (>1,800 m) in Zimbabwe or Lake Titicaca (3,812 m) in Peru and Bolivia. However, the dual influence of water temperature and elevation on the success of introduced rainbow

trout in the tropics has not been clearly determined. In the current context of global warming, understanding how rainbow trout populations respond to high temperatures in tropical regions may help better predict the future of salmonids in their native ranges.

The objective of this study was to investigate how introduction effort and temperature influence the presence of self-sustaining populations of rainbow trout in tropical regions. After reviewing the literature on tropical populations of the species, we focused on the unique case study of Reunion Island (West Indian Ocean), where rainbow trout has been repeatedly introduced since 1940, with varying success. A dense network of rivers that flow from high-elevation volcanic mountains provides a large gradient of water temperatures. Since early life stages of salmonid species are particularly sensitive to high temperatures (Sullivan *et al.* 2000; Benjamin *et al.* 2013), we explored the influence of altitude, as a proxy of temperature as the main driver of successful population establishment after introduction. Warming summer temperatures are identified as a main threat for the viability of rainbow trout populations in its native range in the northwestern United States (Sloat & Osterback 2013; Brewitt & Danner 2014). In the tropics, where summer temperatures are already high, warming winters could reach a critical level that is incompatible with trout reproduction and thus become relatively more limiting than in a temperate climate. We reviewed and collected the stocking history of rainbow trout on Reunion Island to determine whether the introduction effort influenced the presence/absence of trout.

MATERIALS & METHODS

Study site

Reunion Island is a 2,512 km² French overseas department located in the Indian Ocean (21.2°S, 55.6°E). The highest peak of this volcanic island reaches 3,069 m above sea level. Its hydrology is characterized by heavy rain during the cyclone season (December to April). The topography and aquifers generate high spatial and temporal heterogeneity in the quantity of water in rivers (Robert 1985). The island is drained by 13 perennial rivers and 43 transient watersheds characterized by their high degree of natural habitat fragmentation and their torrentiality, which can reach up to 30,000 l/s/km² on a few km² of watershed (Robert 1985). Anthropogenic barriers, which are less common than natural fragmentation, are restricted mainly to lowland areas. The rivers of Reunion Island host 26 fish and 11 decapod crustacean species, of which 20 and 10, respectively, are indigenous catadromous species (Keith 2002). Of the seven species that were introduced to the island, only four established self-sustaining

populations: tilapia (*Oreochromis niloticus*), guppy (*Poecilia reticulata*), swordtail (*Xiphophorus hellerii*) and the non-anadromous form of rainbow trout (Keith 2002).

Introduction history of rainbow trout

The rainbow trout introduced to Reunion Island have come from Madagascar (Robert, 1975), with the initial objective to provide a new source of food for the French colony (Robert, 1978). We reviewed the available reports to update the history of trout stocking on Reunion Island. The main sources of information were publications by R. Robert (Robert 1975, 1976, 1977, 1978) and reports by the National Forest Office and the fishing association. Despite our efforts to identify all introduction events from 1940-2019, some information from 1940-1950 and 1978-1990 may have been missing due to the loss of archival records. The presence of juvenile rainbow trout in the absence of introduction, which indicates wild reproduction, is the criterion used to define a self-sustaining population. The presence of self-sustaining rainbow trout populations is assessed by visual observation (e.g. presence of juveniles, acts of spawning, redds) and electrofishing surveys conducted by the local association for fishing and the protection of aquatic environments (hereafter, fishing association). The presence or absence of a self-sustaining population from 2016-2020 was assessed at each site where trout had been introduced, all sites where its presence was suspected. A site, the smallest spatial unit considered, was defined as a stretch of river a few hundred meters long with homogeneous habitat that is usually delineated by natural discontinuities (waterfalls). We assumed that each site's characteristics and dynamics did not change over the five years of the study. Each site was described by a unique identification number, the river and local name, elevation (Table 2).

Relating temperature to elevation

Autonomous temperature loggers provided detailed measurements of water temperature at the study sites. Water temperature was recorded daily from 2010-2021 by autonomous data loggers or fixed meteorological stations (Table 2). The local water agency (http://www.naiades.eaufrance.fr) and the fishing association provided the water temperature data. Mean monthly temperature at 12 p.m. was used as an explanatory variable in the analysis. However, temporal and spatial coverage was often limited by the loss of loggers due to cyclones or vandalism, by the cost of the devices or by the logistical difficulties involved in deploying scientific equipment in a remote area. Since temperature depends on elevation, we explored the linear relationship between mean monthly water temperature and season/elevation using a GLM with a normal distribution of errors. The model was based on

temperature and elevation data from all 33 sites (data available at https://data.inrae.fr/dataset.xhtml?persistentId=doi:10.57745/9QCPCV).

RESULTS

Introduction history of rainbow trout

We found evidence that rainbow trout were introduced into 8 rivers (i.e. Mât, Roches, Marsouins, Galets, Langevin, Remparts, St Etienne, Ste Suzanne) out of the 13 perennial rivers on Reunion Island (Table 2, Fig. 2). Historical records documented that 340,070 juvenile rainbow trout were released at 38 sites in these rivers or their tributaries from 1940-2019 (Table 2). The introduction effort varied greatly among sites (Fig. 2), with a single introduction of 200 trout in 2006 at site 44, but 72,325 trout introduced over more than 30 events at site 29. The presence of a self-sustaining population of trout was recorded at 10 sites of the 38 introduction sites (Table 2).

Seasonal water temperature profiles

Mean monthly water temperatures at the study sites ranged greatly, from 11.0-22.1°C in winter and 14.6-27.7°C in summer (Fig. 3). Seasonality, which can be described by a decrease in temperature in winter, was distinct at some sites, while temperature profiles remained flat throughout the year at several other sites. Self-sustaining populations of trout were present at sites with low water temperature and high seasonal variability, but some sites with low water temperature had no trout, despite trout introduction. At sites where trout were present, mean temperature in summer and winter were 17.5°C (standard deviation (SD) = 0.8) and 14.9°C (SD = 1.0), respectively. The warmest month on record at a site with trout reached a mean temperature of 19.6°C (SD = 1.04) (site 42 in March 2020).

Using elevation as a proxy of temperature

Mean monthly water temperature correlated strongly with both elevation and season (Table 3), which explained as much as 76.6% of the variance in the data. Adding the interaction between elevation and season did not improve the fit of the model, which indicated that the mean decrease in water temperature for each meter in elevation was similar in winter and summer (Table 3, Fig. 4).

DISCUSSION

Rainbow trout is a stenotherm species that is native to temperate rivers of North America. It has been extensively introduced all over the world and the literature review indicates that tropical regions are no exception (Fig. 1 and S1). Despite the recognized plasticity of trout, its long-term persistence in the tropics seems challenging due to its affinity for cold water. On Reunion Island, the introduction of trout into at least 35 different sites in river stretches across the country illustrates a long-term and widely spread introduction effort across high-elevation perennial rivers. Despite the recorded release of > 400,000 rainbow trout, we are aware of only 10 self-sustaining populations.

We found no evidence of the presence of self-sustaining populations outside sites where trout had been introduced. If trout disperse, they likely do so over very short distances. Several environmental factors can hinder trout expansion across rivers on Reunion Island. First, the island's topography, with steep mountains and many natural waterfalls that result in a high level of habitat fragmentation, strongly limits rainbow trout dispersal from introduction sites. Temporary drying of river stretches due to the highly permeable substrate also limits dispersal. We do not know when the populations established, but based on the long history of introduction (> 80 years), it is likely that some populations have sustained themselves for more than one decade. It is also possible that self-sustaining populations went extinct and re-established themselves at different times in recent history, since trout were introduced into rivers regardless of whether or not they were already present. Nevertheless, there is no evidence for a large expansion of trout from introduction sites, and the risk of expansion from current self-sustaining populations seems low.

The observation of only 10 sites with self-sustaining populations, along with the absence of expansion from these sites, suggest that the environmental conditions that trout encounter after introduction or while dispersing may not meet their ecological requirements well. High water temperature may limit the presence of trout in perennial rivers of Reunion Island, since rainbow trout are sensitive to temperature (Fausch 2007). These tropical rivers have a wide range of water temperatures, with higher variability among sites (spatial variability) than within sites (seasonal variability). As expected, the warmest sites had no trout, while the coldest sites did have them. On Reunion Island, trout experience a mean monthly temperature of 17.5°C in summer, with a maximum of 19.6°C.

This range differs from the optimal temperature range for rainbow trout defined in the literature (i.e. 10-16°C (Raleigh 1984; Jalabert & Fostier 2010)) but lies below the extreme of 30°C recorded in some pools in southern California, United States (Sloat & Osterback 2013). Some of the coldest sites, however, had no trout despite many introduction events. In the trout's native range, increasing summer temperature is a main concern for its persistence under ongoing climate change. It is predicted that higher summer temperatures will decrease the area of thermally suitable riverine trout habitats and cause them to shift upstream (Ruesch et al. 2012; Carlson et al. 2017; Isaak et al. 2018). In contrast, winter temperature requirements are nearly always met in temperate regions (Sloat & Osterback 2013; Brewitt & Danner 2014). Scott and Poynter (1991) suggested that high winter temperatures, rather than high summer temperatures, influenced the northern limit of trout distribution in New Zealand. On Reunion Island, the highly fragmented habitat prevents trout from dispersing upstream, or even migrating seasonally to colder and more suitable habitats.

High summer temperatures may be responsible for the inability of trout to establish themselves at sites where they were introduced, since these temperatures approach the physiological limits of trout. In winter, trout require temperatures no higher than 13°C for several months for successful reproduction (Morrison & Smith 1986; Pankhurst et al. 1996; Jalabert & Fostier 2010), which makes winter a critical period of the life cycle. However, prolonged cold temperatures rarely occur on Reunion Island (8 out of 843 mean monthly temperatures <13°C). The relative rate of increase in summer and winter temperatures will be critical in determining the persistence and extinction risk of trout populations. This has implications for improving understanding of responses of rainbow trout to climate change in its native range. In the future, winter temperatures could become a limiting factor for trout reproduction.

Although detailed water temperature data are rare, the presence of trout populations at high elevations in the tropics, where water temperatures are usually cooler, was no surprise. However, other key requirements may limit the presence of trout, such as natural conditions (e.g. extreme flow regime, spawning habitat, food resources, thermal refugee) and human forcing (e.g. pollution, poaching). Furthermore, although we can model explain 77% of the variance of temperature based on elevation, highly contrasting geologic and topographic features can generate large differences between watersheds. For instance, the Cilaos region of Reunion Island has warmer water due to the presence of geothermal groundwater (sites 23, 36 and 37); thus, it may not be suitable for trout, despite its higher elevation. In contrast, upwelling of cool groundwater in the United States can form thermal refuges that

may become critical habitat that allows trout to persist at higher temperatures (Ebersole *et al.* 2001; Sloat & Osterback 2013; Brewitt & Danner 2014).

AKNOWLEDGMENTS

We are grateful to A. Métro, G. Vienne, J. Maillot, D. Vitry, G. Hoarau and all those who were involved in collecting the data used in this study (FDAAPPMA of Reunion Island, its associated AAPPMA and the water Agency of Reunion Island). The study was funded in part by the French National Office for Biodiversity (TAC-Reunion project), which funded the salary of AM and JR.

REFERENCES

Araki H, Berejikian BA, Ford MJ, Blouin MS (2008) Fitness of hatchery-reared salmonids in the wild. *Evolutionary Applications*, **1**, 342–355.

Behnke R (2010) *Trout and Salmon of North America*. Simon and Schuster.

Benjamin JR, Connolly PJ, Romine JG, Perry RW (2013) Potential effects of changes in temperature and food resources on life history trajectories of juvenile Oncorhynchus mykiss. *Transactions of the American Fisheries Society*, **142**, 208–220.

Bolker BM, Brooks ME, Clark CJ, Geange SW, Poulsen JR, Stevens MHH, White J-SS (2009) Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology & Evolution*, **24**, 127–135.

Brewitt KS, Danner EM (2014) Spatio-temporal temperature variation influences juvenile steelhead (Oncorhynchus mykiss) use of thermal refuges. *Ecosphere*, **5**, 1–26.

Burnham KP, Anderson DR, Huyvaert KP (2011) AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. *Behavioral Ecology and Sociobiology*, **65**, 23–35. https://doi.org/10.1007/s00265-010-1029-6

Carlson AK, Taylor WW, Schlee KM, Zorn TG, Infante DM (2017) Projected impacts of climate change on stream salmonids with implications for resilience-based management. *Ecology of Freshwater Fish*, **26**, 190–204.

Copp GH, Bianco PG, Bogutskaya NG, Er\Hos T, Falka I, Ferreira MT, Fox MG, Freyhof J, Gozlan RE, Grabowska J (2005) To be, or not to be, a non-native freshwater fish? *Journal of Applied Ichthyology*, **21**, 242–262.

Cowx IG (1997) Introduction of fish species into European fresh waters: economic successes or ecological disasters? *Bulletin Francais de la Peche et de la Pisciculture (France)*.

Crawford SS, Muir AM (2008) Global introductions of salmon and trout in the genus Oncorhynchus: 1870–2007. *Reviews in Fish Biology and Fisheries*, **18**, 313–344.

Ebersole JL, Liss WJ, Frissell CA (2001) Relationship between stream temperature, thermal refugia and rainbow trout Oncorhynchus mykiss abundance in arid-land streams in the northwestern United States. *Ecology of freshwater fish*, **10**, 1–10.

FAO (2003) Fishery Records Collections. FIGIS Data Collection.

Fausch KD (2007) Introduction, establishment and effects of non-native salmonids: considering the risk of rainbow trout invasion in the United Kingdom. *Journal of Fish Biology*, **71**, 1–32.

Gall GA, Crandell PA (1992) The rainbow trout. *Aquaculture*, **100**, 1–10.

Isaak DJ, Luce CH, Horan DL, Chandler GL, Wollrab SP, Nagel DE (2018) Global warming of salmon and trout rivers in the Northwestern US: road to ruin or path through purgatory? *Transactions of the American Fisheries Society*, **147**, 566–587.

Jalabert B, Fostier A (2010) *La truite arc-en-ciel: De la biologie à l'élevage*. Editions Quae.

Kadye WT, Chakona A, Marufu LT, Samukange T (2013) The impact of non-native rainbow trout within Afro-montane streams in eastern Zimbabwe. *Hydrobiologia*, **720**, 75–88.

Keith P (2002) Freshwater fish and decapod crustacean populations on Réunion Island, with an assessment of species introductions. *Bulletin Français de la Pêche et de la Pisciculture*, 97–107.

King HR, Pankhurst NW, Watts M, Pankhurst PM (2003) Effect of elevated summer temperatures on gonadal steroid production, vitellogenesis and egg quality in female Atlantic salmon. *Journal of Fish Biology*, **63**, 153–167.

Koutsikos N, Vardakas L, Zogaris S, Perdikaris C, Kalantzi O-I, Economou AN (2019) Does rainbow trout justify its high rank among alien invasive species? Insights from a nationwide survey in Greece. *Aquatic Conservation: Marine and Freshwater Ecosystems*, **29**, 409–423.

Lecomte F, Beall E, Chat J, Davaine P, Gaudin P (2013) The complete history of salmonid introductions in the Kerguelen Islands, Southern Ocean. *Polar Biology*, **36**, 457–475. https://doi.org/10.1007/s00300-012-1281-5

MacCrimmon HR (1971) World Distribution of Rainbow Trout (*Salmo gairdneri*). *Journal of the Fisheries Research Board of Canada*, **28**, 663–704. https://doi.org/10.1139/f71-098

Matthews KR, Berg NH (1997) Rainbow trout responses to water temperature and dissolved oxygen stress in two southern California stream pools. *Journal of Fish Biology*, **50**, 50–67.

Morrison JK, Smith CE (1986) Altering the spawning cycle of rainbow trout by manipulating water temperature. *The Progressive Fish-Culturist*, **48**, 52–54.

Ngugi CC, Green JM (2007) Anthropogenic influences on population sizes, age and growth of naturalized rainbow trout, Oncorhynchus mykiss, in Kenya. *African Journal of Ecology*, **45**, 86–93.

Pankhurst NW, Purser GJ, Van Der Kraak G, Thomas PM, Forteath GNR (1996) Effect of holding temperature on ovulation, egg fertility, plasma levels of reproductive hormones and in vitro ovarian steroidogenesis in the rainbow trout Oncorhynchus mykiss. *Aquaculture*, **146**, 277–290.

Péfaur JE, Sierra NM (1998) Distribución y densidad de la trucha Oncorhynchus mykiss (Salmoniformes: Salmonidae) en los Andes venezolanos. *Revista de biología tropical*, **46**, 775–782.

R Development Core Team (2018) *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.

Raleigh RF (1984) *Habitat suitability information: rainbow trout*. Western Energy and Land Use Team, Division of Biological Services, Research ….

Robert R (1975) Note sur les essais d'aquaculture à la Réunion. *Madagascar : revue de géographie*, **26**, 101–102.

Robert R (1976) L'introduction de la truite arc-en-ciel à La Réunion. *Revue forestière française*, **5**, 386–387.

Robert R (1977) *Introduction et propagation de la truite arc-en-ciel à la réunion.*

Robert R (1978) La salmoniculture à l'Ile de la Réunion. *BOIS & FORETS DES TROPIQUES*, **177**, 65–70.

Robert R (1985) Géographie de l'eau à l'île de la Réunion : essai de distribution régionale de l'alimentation et de l'écoulement. These de doctorat Thesis. Montpellier.

Ruesch AS, Torgersen CE, Lawler JJ, Olden JD, Peterson EE, Volk CJ, Lawrence DJ (2012) Projected climate-induced habitat loss for salmonids in the John Day River network, Oregon, USA. *Conservation Biology*, **26**, 873–882.

Sauter ST, McMillan J, Dunham JB (2001) Salmonid behavior and water temperature. *Seattle, WA: United States, Environmental Protection Agency, Region 10 Office of Water. Final Report to the Policy workgroup of the EPA Region 10 Water Temperature Criteria Guidance Project. EPA 910-D-01-001. 36 p.*

Scott WB, Crossman EJ (1973) *Freshwater fishes of Canada*. Otawa.

Scott D, Poynter M (1991) Upper temperature limits for trout in New Zealand and climate change. *Hydrobiologia*, **222**, 147–151.

Sloat MR, Osterback A-MK (2013) Maximum stream temperature and the occurrence, abundance, and behavior of steelhead trout (Oncorhynchus mykiss) in a southern California stream. *Canadian Journal of Fisheries and Aquatic Sciences*, **70**, 64–73.

Stanković D, Crivelli AJ, Snoj A (2015) Rainbow Trout in Europe: Introduction, Naturalization, and Impacts. *Reviews in Fisheries Science & Aquaculture*, **23**, 39–71. https://doi.org/10.1080/23308249.2015.1024825

Sullivan K, Martin DJ, Cardwell RD, Toll JE, Duke S (2000) An analysis of the effects of temperature on salmonids of the Pacific Northwest with implications for selecting temperature criteria. *Sustainable Ecosystems Institute, Portland, OR*.

Therezien Y (1960) L'introduction de poissons d'eau douce à Madagascar, leur influence sur la modification du biotope. *Bulletin français de pisciculture*, 45–61.

Vila I, Pardo R, Scott S (2007) Freshwater fishes of the Altiplano. *Aquatic Ecosystem Health & Management*, **10**, 201–211.

Vimos DJ, Encalada AC, Ríos-Touma B, Suárez E, Prat N (2015) Effects of exotic trout on benthic communities in high-Andean tropical streams. *Freshwater Science*, **34**, 770–783.

Welcomme RL (1988) *International Introductions of Inland Aquatic Species*. Food and Agriculture Organization of the United Nations.

TABLES

Table 1. Evidence of introductions and the presence of self-sustaining populations of rainbow trout in the tropics.

Country	Origin	Year of first introduction	Motivation	Introduction sites	Altitude	Establishment of self-sustaining populations	Sites where successful	Altitude
Bolivia	U.S.A., Chile	1942	Aquaculture, sport, food	Lake Titicaca (Terrazas 1943 in (Maccrimmon, 1971))	3810 m (Dejoux and Iltis, 1991)	Yes (Maccrimmon, 1971)(FAO 1997)	Lake Titicaca (Terrazas personal communication in (Maccrimmon, 1971))	Unknown
				Lakes and rivers of the Andes: Cordillera Real, Cordillera Oriental, especially the Tunari near Cochabamba, and rivers near Potosi (Terrazas personal communication in (Maccrimmon, 1971))	Unknown	Yes (Aguilera et al., 2006)	Cordillera Real, Cordillera Oriental, especially the Tunari near Cochabamba, and rivers near Potosi Lakes of the Cordillera del Tunari (Aguilera et al., 2006)	4,000 - 4,600 m
Brazil	Denmark, U.S., U.K.	1949	Sport	Bocaina River, Rio de Janeiro, Campos do Jardo, Sao Paulo, Noval Friburgo, Rio de Janeiro; Upper Santa Catarina State (Ripley personal communication in (Maccrimmon, 1971))	Unknown	Yes (Maccrimmon, 1971)	Bocaina River, Rio de Janeiro, Campos do Jardo, Sao Paulo, Noval Friburgo, Rio de Janeiro; Upper Santa Catarina State (Ripley personal communication in (Maccrimmon, 1971))	Unknown

				Silveira stream near São José dos Ausentes, (Winckler-Sosinski et al., 2005)	> 1,200 m	Yes (Lazzarotto et al., 2007)	Upper Macaé River (Lazzarotto et al., 2007)	Unknown
Cameroon	Germany	Unknown	Unknown	Unknown	Unknown	Unknown (Welcomme, 1988)	NA	NA
Colombia	U.S.A.	1926	Aquaculture, sport, food	Lago de Tota, Laguna de La Cocha (Pereira personal communication in (Maccrimmon, 1971))	Unknown	Yes (Maccrimmon, 1971) (Larrahondo Molina, 1993)	Lago de Tota, Laguna de La Cocha (Pereira personal communication in (Maccrimmon, 1971))	Unknown
						Probably not established (Welcomme, 1988)	Embalse del Neusa - Embalse La Regadera - Embalse de Chuza - Embalse de Chisaca - Embalse Los Tunjos (Larrahondo Molina, 1993)	3,269 m - 3,002 m - 2,999 m - 3,146 m - 3,734 m
Costa Rica	U.S.A., Mexico	1927	Aquaculture, sport	Lake in Finca la Georgina in the Central Volcanic Range, Rios Coton, Coto Brus, Macho, Pejiballe, Reventazon, Parrita, Humo, Poas, El Roble, La Paz, Playas, Virilla, Durazno, Savegere (Nanne personal communication in (Maccrimmon, 1971))	Unknown	Yes (Maccrimmon, 1971)	Quebrada del Desengano, Rios Coton, Coto Brus, Macho, Pejiballe, Reventazon, Parrita, Humo, Poas, El Roble, La Paz, Playas, Savegere (Nanne personal communication in (Maccrimmon, 1971))	Unknown
						Yes (Welcomme, 1988)	Unknown	Unknown
Cuba	U.S.A.	1911	Unknown	Unknown	Unknown	Unknown (Maccrimmon, 1971)	NA	NA

Democratic Republic of the Congo	Germany, South Africa	1940	Angling	Tributary of Lake Kivu, Mulungu (De Bont personal communication in (Maccrimmon, 1971)) (FAO 1997)	appr. 1,400 m	No (De Bont personal communication in (Maccrimmon, 1971), (Welcomme, 1988)	NA		NA
Dominican Republic	U.S.A.	1985	Aquaculture	National Park of Valle Nuevo (Sartorio, 2003)	2,600 m	Yes (Sartorio, 2003)	National Park of Valle Nuevo (Sartorio, 2003)		2,600 m
Ecuador	U.S.A., Chile, Columbia, Peru, Argentina	1928	Aquaculture, sport, food	San Pablo Lake (Montenegro personal communication in (Maccrimmon, 1971))	2,660 m	Yes (Maccrimmon, 1971)	Cotopaxi, Punyaru, and Chirimachay, provinces of Carchi, Imbabura, Pichincha, Coxopaxi, Napo, Pastaza, Tungurahua, Chimborazo, Bolivar, Caffar, and Azuay (Montenegro personal communication in (Maccrimmon, 1971))		Unknown
				Lakes and rivers around the Tungurahua Volcano (Montenegro personal communication in (Maccrimmon, 1971))	Unknown	Yes (Jiménez Prado et al., 2015)	In the South: throughout the Paute, Jubones and Cañar basin (Jiménez Prado et al., 2015)		between 1,500 and 4,000 m
							In Alambí, Verdecocha and Carihuaco rivers in the Guayllabamba (Jiménez Prado et al., 2015)		Unknown

Ethiopia	Kenya	1967	Angling	rivers Web, Daka, and Dinchia of Bale Province (Maccrimmon, 1971)		Probable (Maccrimmon, 1971) Yes (Welcomme, 1988; Ahmed et al,, 2014))	Unknown	Unknown
Guatemala	U.S.A.	1982	Unknown	Unknown (FAO 1997)	Unknown	Unknown	NA	NA
Guyana	England	Unknown	Unknown	Unknown (Welcomme 1988)	Unknown	No (Permanent Secretary, personal communication in (Maccrimmon, 1971))	NA	NA
Hawaii	U.S.A.	1920	Angling	Waters of Kauai Island at Kokee, Kawailinana, Kawaikoe, Mohihi, Waiakoali, Kauai, Maui, Hawaii, Oahu and Molokai Islands, Kohala Mountains, Hilo and in the Waipio Valley of Hawaii Island; in the east of Maui Island in Waikamdi and Haipuena streams, Hanawi, Kopiliula, East and West Wailuaki, and East Wailuanui streams (Yoshida personal communication in (Maccrimmon, 1971))	Unknown	Yes (Maccrimmon, 1971))	Waters of Kauai Island at Kokee, Kohala Mountains (Yoshida personal communication in (Maccrimmon, 1971))	915 - 1,220 m
						Yes (Englund and Polhemus, 2001)	Waialae, Koaie, Kauaikinana,Kawaikoi, andWaiakoali, in the western section of the Alakai Plateau (Englund and Polhemus, 2001)	Unknown

Honduras	Costa Rica	Unknown	Angling	Unknown	Unknown	Unknown (Welcomme, 1988)	NA	NA
India	Sri Lanka, U.K., Germany, New Zealand, Scotland	1906	Sport, Angling	Nilgiris Hills (Jhingran personal communication in (Maccrimmon, 1971))	Unknown	Yes (Maccrimmon, 1971)	Nilgiris Hills (Madras State, Avalanche, Emerald, and the Pykara system) (Jhingran personal communication in (Maccrimmon, 1971))	Unknown
						Yes (Petr, 1999)	Nilgiris, Palni Hills and High Range (Petr, 1999)	Unknown
Indonesia	Holland	1929	Unknown	Unknown	Unknown	No (Schuster 1950 in MacCrimmon 1972, (Welcomme, 1988))	NA	NA
Kenya	South Africa	1910	Aquaculture, angling, sport	Nairobi River, on the north slope of Mount Kenya; Nanyuki River; streams in Londiani and Molo on the Mau Escarpment; in the Katamayu River on the Aberdare Mountains; rivers on the east and northeast slopes of Mount Kenya (Smith personal communication in (Maccrimmon, 1971))	Unknown	Yes (Maccrimmon, 1971) (Maccrimmon, 1971) (Ngugi and Green, 2001; Ngugi and Green, 2007)	Mount Kenya, Aberdare Mountains, Mau Escarpment from Nakuru westwards to Kerichs, and Cherangani Mountains and on Mount Elgon (Smith personal communication in (Maccrimmon, 1971))	1,524 m - 2,134 m
							Naro Moru River, Sagana (Mathooko, 1993)	2,035 m
							south-eastern slope of Mt Kenya and Thego stream (Ngugi and Green, 2001; Ngugi and Green, 2007)	4,000 m

Madagascar	France	1922	Angling	Massif of Ankaratra (Thérézien, 1960) and waters of Périnet (Maccrimmon, 1971)	950 m	Yes (Thérézien, 1960 ; Maccrimmon, 1971 ; Welcomme, 1988)	rivers of Antenina and in tributaries of the Onire; Lakera and Ialatsara (Natal personal communication in (Maccrimmon, 1971))	>1500m
Malawi	U.K.	1906	Angling	Zomba Plateau (Crichton personal communication in (Maccrimmon, 1971))	1,220 m	Yes (Maccrimmon, 1971)	Lake Malawi (Crichton personal communication in (Maccrimmon, 1971))	Unknown
				Mlanje Mountain streams (Crichton personal communication in (Maccrimmon, 1971))	1,524 m	Yes (Maccrimmon, 1971)	Likangala River in Zomba Plateau (Eccles personal communication in (Maccrimmon, 1971))	Unknown
				Kaziweziwe stream at Nchenachena and in Nyika Plateau streams and dams (Crichton personal communication in (Maccrimmon, 1971))	2,134 m	Yes (Maccrimmon, 1971)	near Lake Chilwa (Eccles personal communication in (Maccrimmon, 1971))	>1000m
				Streams of the Cameron Highlands in the Telom and Ringlet area (Crichton personal communication in (Maccrimmon, 1971))	Unknown	Yes (Maccrimmon, 1971) (Kadye 2008)	the Telom, the Vertam, the Gulf course area, in Taman Sedia, below Parit falls, upper reaches of the Terla, Ichat, Kial, Bertam, and the Habu (Crichton personal communication in (Maccrimmon, 1971))	Unknown
							Nyika Plateau (Kadye 2008)	Above 1,800 m
Malaysia	Kenya, South Africa,	1907	Unknown	Unknown	Unknown	Probably not established	NA	NA

Country	Source	Year	Reason	Location		Established	Location	Elevation
	Scotland, New Zealand					(Welcomme, 1988)		
Mauritius	South Africa	1934	Aquaculture	Unknown	Unknown	No (Vibert personal communication in (Maccrimmon, 1971))	NA	NA
						Yes (Welcomme, 1988)	unknown	Unknown
Mozambique	Zimbabwe	unknown	unknown	Waters of the Rotanda area (Fonseca personal communication in (Maccrimmon, 1971))	Unknown	No (Fonseca personal communication in (Maccrimmon, 1971)), Unknown (Welcomme, 1988)	NA	NA
Panama	U.S.A.	1925	Sport	Chiriqui and Caldera Rivers (Hildebrand 1935, 1938 in MacCrimmon 1972)	Unknown	Probable (MacCrimmon 1972) Yes (Welcomme, 1988)	Chiriqui and Caldera rivers of the El Volcan de Chiriqui (MacCrimmon 1972)	Unknown
Papua New Guinea	New Zealand, Australia	1952	Aquaculture, angling	Gumanch and Baiyer rivers near Mount Hayen (Rapson personal communication in (Maccrimmon, 1971))	1,830 m	Yes (Maccrimmon, 1971) (Povlsen, 1993)	Gumanch and Baiyer rivers near Mount Hayen, into the Nebilyer system of the western highlands (Conroy personal communication in (Maccrimmon, 1971))	1,830 m

							Kuragamba and Anggura rivers (Povlsen, 1993)	> 2,300 m
Peru	U.S.A.	1927	Food	Lake Titicaca, districts of Junin and Pasco, especially the Rio Mantaro system, waters of the Central Andes, the Rio Mantaro system, and lakes Junrn (Chinchaycocha), Marcapomacocha, Punrun, and others, in all suitable accessible waters of the Andes (Tovar, Vildoso personal communications in (Maccrimmon, 1971))	> 2,000 m	Yes (Maccrimmon, 1971; Dejoux and Iltis, 1991; Vila et al., 2007)	Lake Titicaca and tributary streams, in all suitable accessible waters of the Andes (Tovar personal communication in (Maccrimmon, 1971))	> 2,000 m
							Lake Titicaca (Dejoux 91)	3810 m (Dejoux and Iltis, 1991)
							Altiplano, Andes (Vila et al., 2007)	3,600 to 4,500 m
Philippines	U.S.A.	1962	Unknown	Unknown	Unknown	No (Delmendo personal communication in (Maccrimmon, 1971))	NA	NA
Puerto Rico	U.S.A.	1934	Unknown	Unknown	Unknown	No (Erdman personal communication in (Maccrimmon, 1971) Probably not established	NA	

						(Welcomme, 1988)		
Reunion	Madagascar	1940	Food	Rivière des Marsouins, Salazie Rivers, Cilaos Rivers, Mafate Rivers, Rivière Langevin, Bras Cabot, Rivières des Roches, Bras de la Plaine (Robert, 1978)	Unknown	Yes (this study)	Marsouins River, Salazie Rivers, Mafate Rivers, Rempart River, Ravine des Sept Bras, Bras Ste-Suzanne, Bras Roches Noires (this study)	>550 m
Sri Lanka	U.K.	1889	Angling	Unknown	Unknown	Yes (Maccrimmon, 1971) (Jindasa et al., 2005)(FAO 1997)	Nuwara Eliya stream, Bulu Ellan stream, Portswood Dam, Farr Lakes, Agra Oya, and Gorge Valley rivers, and the Horton Plains streams (Wallis 1969 in (Maccrimmon, 1971))	> 1,220 m
							Tributaries of Belihul Oya only in the Horton Plain (Jindasa et al., 2005)	1,900 m
Sudan	Kenya, U.K.	1947	Sport	Nagishot and Kitiri in the Eastern Equatorial Province (Smith personal communication in (Maccrimmon, 1971))	Unknown	Yes (Smith personal communication in (Maccrimmon, 1971), (Welcomme, 1988)	Unknown	Unknown
Taiwan	Japan	1957	Sport	Inland waters (Welcomme 1988)	Unknown	Yes (Welcomme 1988)	Unknown	Unknown

Thailand	Canada	1973	Sport	Unknown (Welcomme 1988)	Unknown	No (Welcomme 1988)	NA	NA
Tanzania	Scotland	1927	Sport	Streams of the Southern Provinces (Highlands)(Flamwell and Smith personal communications in (Maccrimmon, 1971))	Unknown	Yes (Maccrimmon, 1971), (Welcomme, 1988)	Elton Platau, Mufindi, and Mbeya in the southern Provinces, Mount Meru and the Pare Mountains in the north, and the Usambaras at the coast region (Flamwell personal communication in (Maccrimmon, 1971))	Unknown
				Streams of Mount Kilimanjaro in the Mosho district (Smith personal communication in (Maccrimmon, 1971))	Unknown		Kilimanjaro waters (Smith personal communication in (Maccrimmon, 1971))	> 1,372 m
Trinidad	U.S.A.	1938	Unknown	Unknown	Unknown	No (Wood personal communication in (Maccrimmon, 1971))	NA	NA
Uganda	Scotland, Kenya	1925	Unknown	River Suam on Mount Elgon (Biribonwoha personal communication in (Maccrimmon, 1971))	2,440 to 4,360 m	Yes (Maccrimmon, 1971)	Rivers of Mount Elgon (Biribonwoha personal communication in (Maccrimmon, 1971))	2,440 to 4,360 m

Country	Source	Year	Use	Location (introduction)	Elevation	Established	Location (naturalized)	Elevation
				River Bukwa, Sipi River, Atar and Nyaluti Rivers (Biribonwoha personal communication in (Maccrimmon, 1971))	Unknown	Yes (Maccrimmon, 1971)	Rivers of the Ruwenzori Mountains (Biribonwoha personal communication in (Maccrimmon, 1971))	Unknown
				Mpaga River (Biribonwoha personal communication in (Maccrimmon, 1971))	1,680 to 5,100 m			
Venezuela	U.S.A.	1934	Aquaculture, sport, food	Unknown	Unknown	Yes (Maccrimmon, 1971) (Bastardo, 1994) (Pefaur, 1998)	In suitable waters in the region of the Venezuelan Andes within the states of Trujillo, Mérida, and Tachira (Leon 1966 in (Maccrimmon, 1971))	1,700 - 4,000 m
							Mucunutan stream, Mérida (Bastardo, 1994)	1,900 m
							Chama, Santo Domingo, Motatan, Negro, Grita, Escalante, Mocoties in Andes Venezonalos (Pefaur, 1998)	1700m – 4000m
Zambia	Malawi, South Africa	1942	Unknown	Unknown	Unknown	Unknown (Maccrimmon, 1971)	NA	NA
Zimbabwe	South Africa	1910	Sport	Unknown	Unknown	Yes (Kadye et al., 2013)	Afro-montane streams in Nyanga Mountains (Kadye et al., 2013)	Unknown

Aguilera, X., Declerck, S., De Meester, L., Maldonado, M., and Ollevier, F. (2006). Tropical high Andes lakes: A limnological survey and an assessment of exotic rainbow trout (*Oncorhynchus mykiss*). *Limnologica* 36(4)**,** 258-268. doi: 10.1016/j.limno.2006.08.001.

Ahmed, E.Y., Woldeyes, A., Korra, T., and Laval, G. (2014). Yersiniosis outbreak in rainbow trout at fish farm in Oromia Regional State, Ethiopia. *Ethiopian Veterinary Journal* 18 (2), 35-49.

Bastardo, H. (1994). Food habits of the rainbow trout *Oncorhynchus mykiss* (Salmoniformes: Salmonidae) in an Andean creek of Venezuela. *Revista de Biología Tropical* 42(3)**,** 685-693.

Crawford, S.S., and Muir, A.M. (2008). Global introductions of salmon and trout in the genus *Oncorhynchus*: 1870-2007. *Reviews in Fish Biology and Fisheries* 18(3)**,** 313-344. doi: 10.1007/s11160-007-9079-1.

Dejoux, C., and Iltis, A. (1991). *Lake Titicaca. Synthesis of present limnological knowledge.*

Englund, R.A., and Polhemus, D.A. (2001). Evaluating the Effects of Introduced Rainbow Trout (Oncorhynchus Mykiss) on Native Stream Insects on Kauai Island, Hawaii; Contribution No. 2001-012 to the Hawaii Biological Survey, Bishop Museum. *Journal of Insect Conservation* 5(4)**,** 265-281. doi: 10.1023/A:1013331525088.

FAO. (1997). *FAO database on introduced aquatic species*. FAO Rome.

Jiménez Prado, P., Aguirre, W., Laaz Moncayo, E., Navarrete Amaya, R., Nugra Salazar, F., Rebolledo Monsalve, E., et al. (2015). *Guia de peces para aguas continentales en la vertiente occidental del Ecuador.* Pontificia Universidad Católica del Ecuador Sede EsmeraldasISBN: 978-9942-20-740-1.

Jindasa, J., Kothlawala, A.B., and Herath, H.M.B.K. (2005). Status of Rainbow trout (*Oncorhynchus mykiss*-Waldaum 1792) population after the cessation of stocking in waterways of Horton Plains, Sri Lanka. *Vidyodaya Journal of Science* 12**,** 9-27.

Kadye, W.T., Chakona, A., Marufu, L.T., and Samukange, T. (2013). The impact of non-native rainbow trout within Afro-montane streams in eastern Zimbabwe. *Hydrobiologia* 720(1)**,** 75-88. doi: 10.1007/s10750-013-1624-4.

Larrahondo Molina, M. (1993). "APROVECHAMIENTO ACUICOLA DE EMBALSES EN COLOMBIA," in *AVANCES EN EL MANEJO Y APROVECHAMIENTO ACUICOLA DE EMBALSES EN AMERICA LATINA Y EL CARIBE,* ed. J.R.J.P.a.E. Varsi.).

Lazzarotto, H., Brito, M.F.G., and Caramaschi, É.P. (2007). Threatened fishes of the world: *Pareiorhaphis garbei* (Ihering, 1911) (Ostariophysii: Loricariidae). *Environmental Biology of Fishes* 78(1)**,** 91-92. doi: 10.1007/s10641-006-9036-1.

Maccrimmon, H.R. (1971). World Distribution of Rainbow Trout (*Salmo gairdneri*). *Journal of the Fisheries Research Board of Canada* 28**,** 663-704. doi: 10.1139/f71-098.

Mathooko, J.M. (1993). AVAILABILITY AND SELECTION OF MAYFLY, CAENIS-KUNGU KIMM (EPHEMEROPTERA, CAENIDAE) AS A NATURAL SOURCE OF FOOD BY RAINBOW-TROUT,

ONCORHYNCHUS-MYKISS WALB, IN A TROPICAL RIVER IN KENYA. *African Journal of Ecology* 31(1)**,** 58-65. doi: 10.1111/j.1365-2028.1993.tb00518.x.

Ngugi, C.C., and Green, J.M. (2001). Information from Radio Telemetry on movements and exploitation of naturalized Rainbow trout, *Oncorhynchus mykiss* (Walbaum), in Kenya cold water streams. *African Journal of Tropical Hydrobiology and Fisheries* 10**,** 27-38.

Ngugi, C.C., and Green, J.M. (2007). Anthropogenic influences on population sizes, age and growth of naturalized rainbow trout, *Oncorhynchus mykiss*, in Kenya. *African Journal of Ecology* 45(1)**,** 86-93. doi: 10.1111/j.1365-2028.2006.00684.x.

Pefaur, J.E. (1998). Distribution and density of the trout *Oncorhynchus mykiss* (Salmoniformes : Salmonidae) in the Venezuelan Andes. *Revista de Biología Tropical* 46(3).

Petr, T. (1999). Fish and fisheries at higher altitudes: Asia. . *FAO Fisheries Technical Paper. No. 385. Rome, FAO. 304p.* http://www.fao.org/3/x2614e00.htm.

Povlsen, A.F. (1993). "Observations on the biology and ecology of rainbow trout, *Oncorhynchus mykiss*, and its implications for fisheries in the highlands of Papua New Guinea. ", (ed.) R.p. A report prepared for the Sepik River Fish stock Enhancement Project. Fl:PNG/85/00L Field Doc. 21. FAO.).

Robert, R. (1978). La salmoniculture à l'île de La Réunion. *Revue Bois et Forêts des Tropiques* 77**,** 65-70.

Sartorio, A. (2003). Tropical adaptation of rainbow trout (*Oncorhynchus mykiss*). *Aliens Newsletter of Invasive Species Specialist Group* 17**,** 30-31.

Therezien, Y (1960). L'introduction de poissons d'eau douce à Madagascar, leur influence sur la modification du biotope. *Bulletin Français de Pisciculture* 199, 45-61.

Vila, I., Pardo, R., and Scott, S. (2007). Freshwater fishes of the Altiplano. *Aquatic Ecosystem Health & Management* 10(2)**,** 201-211. doi: 10.1080/14634980701351395.

Welcomme, R.L. 1988. International introductions of inland aquatic species. *FAO, Rome, Fish Tech Pap294* [Online]. Available: http://www.fao.org/docrep/x5628e/x5628e00.htm#Contents.

Winckler-Sosinski, L., Schwarzbold, A., and Schulz, U.H. (2005). Survival of Rainbow Trout *Oncorhynchus mykiss* Walbaum, 1792 (Salmoniformes - Salmonidae) Eggs in an Altitude Stream in Southern Brazil. *Acta Limnol. Bras.* 17(4)**,** 465-472.

71   Table 2**.** Presentation of the study sites in La Reunion Island: elevation, presence of self-sustaining population of

72   trout, trout introduction history (number of introduction, number of trout introduced, period of introduction).

Watershed	Site	ID	Elevation	Presence trout	Number trout	First intro	Last intro	Number intro
Rivière des Marsouins	Bras Chanson	44	1316	0	200	2006	2006	1
Rivière des Marsouins	Grande Ravine	53	608	0	400	2006	2007	2
Rivière du Mât	Bras dAmales	35	1362	0	500	1945	1945	1
Rivière St Etienne	Bras Sec	59	604	0	500	2008	2008	1
Rivière du Mât	Bras des Demoiselles	33	866	0	525	1944	2007	3
Rivière des Galets	Bras Bémal	28	1463	0	900	1980	1983	3
Rivière Ste Suzanne	Niagara	46	93	0	1000	2010	2010	1
Rivière des Galets	Bras Cimendal	27	750	0	1100	1980	1983	3
Rivière des Marsouins	Ravine PK 12	54	665	0	1200	2006	2007	2
Rivière du Mât	Ravine à Jacquot	32	756	0	1200	2019	2019	1
Rivière des Marsouins	Bras Sec	55	701	0	1300	2006	2007	2
Rivière du Mât	Bras Piton	45	874	0	1600	2006	2007	2
Rivière des Roches	Bassin La Mer	56	159	0	1800	2007	2007	1
Rivière du Mât	Rivière Terre Rouge	34	423	0	2000	2006	2007	2
Rivière du Mât	Ravine Camp Pierrot	30	1010	0	2490	2006	2019	5
Rivière du Mât	Bras de Caverne	31	1536	0	2655	1942	2012	6
Rivière des Roches	Ravine Terre Rouge aval	57	94	0	6000	2010	2011	2
Rivière du Mât	Bras des Lianes	11	789	0	6000	2006	2007	2
Rivière des Roches	Ravine Terre Rouge cascade	58	615	0	6500	2017	2018	2
Rivière des Marsouins	Bras Patience	41	836	0	9700	2006	2015	7
Rivière Langevin	Grand Galet	10	409	0	13100	2004	2015	9
Rivière du Mât	Ilet à Vidot	13	688	0	13750	2006	2016	9
Rivière Langevin	La Passerelle	7	242	0	14565	2007	2015	8
Rivière St Etienne	Bras de St Paul	37	922	0	14750	2008	2015	7
Rivière du Mât	Camp Pierrot Grand Sable	20	1105	0	18200	2006	2016	12
Rivière St Etienne	Bras Rouge	23	911	0	32400	1945	2019	14
Rivière St Etienne	Bras de Benjoin	36	915	0	38995	1945	2019	17
Rivière du Mât	Rivière Fleurs Jaunes	29	700	0	72325	1945	2019	30
Rivière des Remparts	Bras Caron	17	827	1	300	1989	1989	1
Rivière St Etienne	Bras des Roches Noires	40	750	1	500	2008	2008	1
Rivière des Marsouins	Barrage Taka1	52	506	1	600	2006	2007	2
Rivière des Marsouins	Bras Cabot	24	1171	1	2820	1945	2007	9
Rivière du Mât	Trou Blanc	18	850	1	3000	2006	2016	14
Rivière des Marsouins	Plaine des Palmistes	25	1297	1	5095	1944	1983	10
Rivière Langevin	Cap Blanc	16	820	1	13500	2004	2015	9
Rivière des Galets	Marla	26	1469	1	13800	2008	2015	8
Rivière du Mât	Camp Pierrot Fleur jaune	22	1043	1	18200	2006	2016	12
Rivière St Etienne	Bras Ste Suzanne	19	890	1	20600	2004	2011	8

74   Table 3. Modeling mean monthly water temperature on Reunion Island as a function of elevation and season.

Model	np	AIC	△AIC	Pseudo-R²
Temp ~ 1	1	5775.1	4426	0.000
Temp ~ Elevation	2	2886.5	1537.4	0.500
Temp ~ Elevation + Season	3	1349.1	0.00	0.766
Temp ~ Elevation × Season	4	1349.7	0.6	0.766

FIGURE CAPTIONS

Fig. 1. Current distribution of rainbow trout worldwide. a) Native-distribution (green), temperate (orange) and

tropical (red) countries in which the species has been introduced (adapted from Crawford and Muir (2008). b)

Results of introducing rainbow trout into tropical countries: evidence of the presence of self-sustaining populations

in 1988 (in the review by Welcomme (1988), light blue) and after 1988 (dark blue), no information available about

the presence of self-sustaining populations (light green), and no evidence of self-sustaining populations (dark

green). See Table 1 for supporting information and references.

Fig. 2. Map of Reunion Island showing perennial rivers and the location and introduction effort at 33 sites where

rainbow trout were introduced from 1940-2019. The number of introduction events, used as a proxy of the effort,

is represented by increasing shades of gray. The star symbol represents an introduction site with unknown effort.

See Table 2 for more information about the sites.

Fig. 3. Mean monthly water temperatures at 35 sites on perennial rivers on Reunion Island. Colors show the sites

where attempts to introduce rainbow trout resulted in self-sustaining populations (green) or were not successful

(red), and sites with no introduction events (gray).

Fig. 4. Relationship between mean monthly water temperature and elevation in winter (blue) and summer (yellow)

on Reunion Island. The solid lines are model estimates (Table 3), and the gray circles are raw data.

a)

98   FIGURES

100   Figure 1

b)

103   Figure 2

107   Figure 3

111   Figure 6