Non-catalyzable denitrification intermediates
induce nitrous oxide reduction in two purple nonsulfur bacteria

Breah LaSarre[1,a], Ryan Morlen[2], Gina C. Neumann[1],
Caroline S. Harwood[2], and James B. McKinlay[1]*

[1] Department of Biology, Indiana University, Bloomington, IN
[2] Department of Microbiology, University of Washington, Seattle, WA

*Corresponding author: jmckinla@iu.edu

Current address:
[a] Department of Plant Pathology and Microbiology, Iowa State University, Ames, Iowa, USA
[b] Benson Hill, St. Louis MO, USA

Abstract

Denitrification is a form of anaerobic respiration wherein nitrate is sequentially reduced via
nitrite, nitric oxide, and nitrous oxide to dinitrogen gas. The pathway has mostly been
characterized in bacteria that have the complete denitrification pathway. However, each
individual step can serve as a form of anaerobic respiration and many bacteria only have partial
denitrification pathways. The conditions under which these partial denitrification pathways are
activated have received comparatively little attention. Here we report the activity of partial
denitrification pathways in two purple nonsulfur bacteria, *Rhodopseudomonas palustris* CGA009
and *Rhodobacter capsulatus* SB1003. These bacteria can use one or more denitrification steps as
an electron sink during phototrophic growth or to transform energy during anaerobic respiration
in the dark. In each case, nitrous oxide reduction required supplementation with a non-
catalyzable denitrification intermediate. Thus, bacteria that maintain partial denitrification
pathways are still subject to regulation by denitrification intermediates that they cannot use.

Importance. Denitrification is a form of microbial respiration wherein nitrate is converted into
dinitrogen gas, a major component of Earth's atmosphere, via several nitrogen oxide
intermediates. Unfortunately, denitrification of nitrate in agricultural fertilizers is often
incomplete, resulting in the emission of the potent greenhouse gas, nitrous oxide. Some bacteria
do not have all the steps for denitrification, but many can still reduce nitrous oxide into harmless
dinitrogen gas. These partial denitrifying bacteria have received little attention. Here we
characterized two partial denitrifying bacteria that are capable of nitrous oxide reduction.
Surprisingly, nitrous oxide reduction was induced by denitrification intermediates that the
bacteria could not respire, suggesting that regulation of nitrous oxide reduction was subject to the
same cues as in complete denitrifiers. This work thus informs on how partial denitrifying
bacteria can contribute to, and potentially combat, greenhouse gas emissions.

Introduction

Denitrification is a multistep respiratory pathway that sequentially reduces nitrate (NO_3^-) via nitrite (NO_2^-), nitric oxide (NO), and nitrous oxide (N_2O) to dinitrogen gas (N_2) (1, 2) (Fig. 1A). While many bacteria have the complete denitrification pathway, the entire pathway is not required; any single denitrification intermediate can potentially support growth via anaerobic respiration (1, 2). Denitrification impacts the health of both humans and the planet (2). Denitrifiers in the human gut help fight pathogens and maintain vascular homeostasis through the generation of NO_2^- and NO (3). Denitrification is also important to the global nitrogen cycle, returning nitrogen to the atmosphere as N_2. Denitrifiers also play important roles in anthropogenic pollution and climate change. Although denitrifiers have the potential to convert NO_3^- in runoff from agricultural fertilizers to harmless N_2 before it can lead to eutrophication of water resources (4), this process often fails to run to completion, even in bacteria capable of complete denitrification. Currently, denitrification is a net generator of the potent greenhouse gas N_2O, driven largely by the wide use of agricultural fertilizers (5).

Denitrifiers can be phototrophs or chemotrophs. Among the bacteria that may be capable of both phototrophic and chemotrophic denitrification are purple nonsulfur bacteria (PNSB). PNSB are perhaps the most versatile organisms ever described. PNSB can grow with either organic or inorganic electron and carbon sources, employing either phototrophy in light or aerobic or anaerobic respiration in the dark. In the dark, anaerobic respiratory pathways serve a critical role in energy transformation (i.e., oxidative phosphorylation). However, during photoheterotrophic growth, wherein light is used for energy and organic substrates are used for carbon and electrons, anaerobic respiratory pathways can help maintain electron balance (6, 7). Electron balance is a challenge to photoheterotrophic growth because more electrons are liberated from many organic substrates than can be incorporated into biomass. Excess electrons must be deposited onto an electron acceptor. CO_2 is commonly used as an electron acceptor under photoheterotrophic conditions, but electron acceptors for anaerobic respiration can also serve this role.

Here we characterized the ability of the PNSB *Rhodopseudomonas palustris* CGA009 and *Rhodobacter capsulatus* SB1003, to carry out denitrification under photoheterotrophic and chemoheterotrophic conditions. In agreement with the genome sequences, we found that CGA009 could use all denitrification intermediates except for NO_3^- and SB1003 could only use N_2O. However, in each case, N_2O utilization required supplementation with other denitrification intermediates, including intermediates that each organism could not reduce. Thus, although each organism only has a partial denitrification pathway, the regulatory schemes resemble those of a complete denitrifier.

Results

R. palustris CGA009 has a partial denitrification pathway. One of the most intensively studied PNSB is *R. palustris* CGA009 (8), yet little is known about its ability to respire anaerobically. Unlike some other model PNSB, CGA009 cannot grow via respiration with dimethylsulfoxide (9-11). However, a partial denitrification pathway was identified in the CGA009 genome (8).

We verified this partial denitrification pathway by aligning query sequences for reductases from
Paracoccus denitrificans PD122 against the CGA009 genome using BlastP (12). We found no
significant similarity to *nar* and *nap* nitrate reductase genes nor to *nirS* nitrite reductase.
However, we found genes with significant sequence identity to those responsible for the last
three steps of denitrification: NO_2^- reduction, NO reduction, and N_2O reduction. The gene
clusters for each of these steps were located at distinct regions in the genome (Fig. 1B). For the
first step of NO_2^- reduction, *R. palustris* CGA009 appears to have two, possibly redundant,
copper-containing NirK NO_2^- reductases encoded far apart on the genome. The latter two steps
of NO reduction and N_2O reduction are each encoded by single gene clusters (Fig. 1B).

Several possible regulatory genes were also identified. A gene that likely encodes NnrR, a
protein that typically regulates NO_2^-, NO, and N_2O reductases in other bacteria in response to
NO (1, 13, 14), was found near the *nirK2* gene (Fig. 1B). Genes with significant sequence
identity to NasTS, which can regulate N_2O reductase in response to NO_3^- (15, 16), were found
far from the other gene clusters, upstream of a NirA-family reductase (RPA3710), which could
be involved in either nitrite or sulfite reduction.

PNSB typically use electron acceptors anaerobically to establish a proton motive force and
generate ATP when incubated in dark. When incubated in light, cells generate ATP by
photophosphorylation and use electron acceptors to enable growth on relatively reduced organic
compounds . Since *R. palustris* grows best in light, we first examined if it could use
denitrification electron acceptors when provided with butyrate, an electron-rich carbon source
that requires an electron sink for photoheterotrophic growth (7, 17).

Fig. 1. General denitrification pathway (A) and the denitrification genes found in the *R. palustris* CGA009 genome (B). Gene products: blue, nitrogen oxide reductases; green, accessory proteins for electron transport, enzyme cofactor assembly, etc.; orange, denitrification regulators; gray, unknown function; black, likely unrelated to denitrification. Numbers indicate the chromosome nucleotide positions. Several CRP/Fnr-family transcriptional regulators with >25% sequence identity to known denitrification regulators are not shown.

In agreement with the genome annotation (Fig. 1B), phototrophic growth on butyrate was not
supported when supplemented with a wide range of $NaNO_3$ concentrations (Fig. 2A). However,
growth on butyrate was observed when CGA009 was provided with 1mM $NaNO_2$, a

concentration that we determined to be near the toxicity limit because it caused a lag in
phototrophic growth on succinate, which does not require supplementation with an electron
acceptor (Fig. 2B). We did not test exogenously-added NO because it is highly toxic and would
likely be impossible to add in amounts that would be practical as an electron sink to yield
observable growth.
123
Based on the amount of phototrophic growth observed on butyrate with NaNO$_2$, we estimated
whether NO$_2^-$ was reduced completely to N$_2$. From our previous work, we estimated that ~6 mM
bicarbonate (HCO$_3^-$) is consumed for an increase in cell density of ~1 OD$_{660}$ during phototrophic
growth on butyrate with NaHCO$_3$ (17). Whereas HCO$_3^-$ can serve as a sink for two electrons,
each molecule of NO$_2^-$ could serve as a sink for three electrons if NO$_2^-$ is reduced sequentially to
N$_2$ but only one electron if NO$_2^-$ is only reduced to NO (Fig. 1A). Based on this stoichiometry,
we predicted a cell density of ~ 0.25 OD$_{660}$ during phototrophic growth with butyrate plus 1 mM
NaNO$_2$ assuming that all the NaNO$_2$ is reduced to N$_2$. Close to our prediction, we observed a
final OD$_{660}$ of 0.28 ± 0.00 (n=3) (Fig. 2A). Thus, we deduce that the latter three reductases for
denitrification (Fig. 1A) are all active in CGA009. However, when CGA009 was provided with a
100% headspace of N$_2$O, no growth was observed within 15 days (Fig. 2A), suggesting that N$_2$O
alone cannot activate N$_2$O reductase.

**Fig. 2. NaNO$_2$ supports phototrophic growth of CGA009 on butyrate within toxicity
limits. A.** Phototrophic growth curves with butyrate and various potential electron acceptors.
B. Phototrophic growth curves on succinate, a condition that readily supports growth without
supplementation with an electron acceptor, with various concentrations of NaNO$_2$ to identify
the toxicity limit. **A, B.** Plots are single representatives of similar trends observed for three
biological replicates except for those exploring different NaNO$_2$ and NaNO$_3$ concentrations.
Cultures had a 100% Ar headspace unless N$_2$O is indicated (100% N$_2$O).

136
N$_2$O reduction by CGA009 is induced by NO$_2^-$ and NO$_3^-$ during photoheterotrophic growth.
In some bacteria, denitrification intermediates other than N$_2$O enhance N$_2$O reduction or are
required to induce N$_2$O reduction (1, 2, 14, 18, 19). NO$_2^-$-induction of N$_2$O reductase can be
mediated by NnrR, though NO$_2^-$ might first need to be converted to NO (14) (Fig. 1B). NO$_3^-$-
induction of N$_2$O reductase can be mediated by NasTS (15, 16) (Fig. 1B). We thus tested
whether NaNO$_2$ or NaNO$_3$ could induce growth with N$_2$O. Micromolar amounts of NaNO$_2$

stimulated growth with N_2O. $NaNO_2$ at 100 µM caused a 4-day lag in growth, probably because
it was toxic at this concentration (Fig. 3A). Final cell densities increased with the amount of
$NaNO_2$ added (Fig. 3A). We conclude that the higher final cell density when N_2O was provided
along with 100 µM $NaNO_2$ is due to the use of N_2O as an electron acceptor rather than NO_2^-
because growth trends with 100 µM $NaNO_2$ alone were much lower (Fig. 3A vs Fig. 2A). Thus,
we speculate that exhaustion of NO_2^- as an electron acceptor eliminates induction of N_2O
reductase, and thus growth on N_2O lasts only as long as the pool of $NaNO_2$.
150
Despite lacking NO_3^- reductase, $NaNO_3$ also induced phototrophic growth on butyrate with N_2O
(Fig. 3B). Similar growth trends were observed between 1 µM to 10 mM $NaNO_3$. Slower growth
and a lower final cell density was observed at $NaNO_3$ concentrations of 0.1 µM or lower (Fig.
3B). The fact that 1 µM $NaNO_3$ resulted in similar growth trends as 10 mM $NaNO_3$ suggests that
NO_3^- is not being reduced and that $NaNO_3$ is relatively non-toxic. Thus, $NaNO_3$ is potentially
useful as a stable inducer of N_2O reductase compared to $NaNO_2$.
157
Initial phototrophic growth on butyrate with $NaNO_3$ was faster than that with bicarbonate
(doubling time ± SD = 13 ± 3 vs 17 ± 2 h, respectively). However, growth with $NaNO_3$
transitioned to slow growth at a cell density of ~0.2 – 0.3 OD_{660} and reached a lower final cell
density than with bicarbonate (ΔOD_{660} ± SD = 0.72 ± 0.05 vs 0.99 ± 0.02, respectively). Flushing
the headspace with N_2O after the transition did not restore the initial growth rate (data not
shown).
164

Fig. 3. $NaNO_2$ and $NaNO_3$ induce phototrophic N_2O utilization by CGA009. A.
Phototrophic growth curves with butyrate +/- N_2O and different concentrations of $NaNO_2$. The
Ar, no N_2O control had a 100% argon headspace in place of N_2O.**B.** Phototrophic growth
curves with butyrate + N_2O and different concentrations of $NaNO_3$. **C.** Phototrophic growth
curves of Calvin cycle deletion mutants with succinate. Cultures had a 100% Ar headspace
unless N_2O is indicated (100% N_2O). **A-C.** Single representatives of similar trends observed for
three biological replicates are shown for each condition except for those exploring different

165
We also tested whether N_2O reduction could serve as an electron sink to rescue *R. palustris*
growth when the CO_2-fixing Calvin cycle is genetically disrupted. *R. palustris* Calvin cycle
mutants cannot grow phototrophically even on relatively oxidized carbon sources like succinate
unless an alternative electron sink is provided (6). For example, NifA* mutations rescue Calvin
cycle mutants by disposing of excess electrons as H_2 gas via constitutively active nitrogenase (6,
17, 20). Here we found that $NaNO_3$ plus N_2O could also rescue an *R. palustris* Calvin cycle

mutant during phototrophic growth on succinate (Fig. 3C). N_2O reduction resulted in more
immediate growth than a NifA* positive control. However, growth eventually slowed and the
culture reached a lower final cell density that the NifA* control, similar to what was observed
during growth on butyrate compared to a bicarbonate-containing control (Fig. 3C vs 2A). Gas
chromatographic analysis of headspace samples confirmed that growth of all cultures with
$NaNO_3$ plus N_2O was not due to spontaneous mutations enabling H_2 production (data not
shown).
179
180 We wondered whether $NaNO_3$ might also improve growth with NO_2^-, perhaps by stimulating
181 NO_2^- reductase activity. However, adding $NaNO_3$ did not affect photoheterotrophic growth
182 trends on butyrate with 1 mM $NaNO_2$, even when $NaNO_3$ was also added to starter cultures as a
183 possible 'pre-inducing' condition (Fig. 4A). The same strategy also did not decrease the lag
184 phase during phototrophic growth on succinate with 1 mM $NaNO_2$.

Fig. 4. $NaNO_3$ does not improve growth trends when NO_2^- is present as an essential electron sink (A) or as a toxic compound (B). Single representatives of similar trends observed for three biological replicates are shown for each condition.

185
N_2O supports anaerobic respiration by CGA009 in the dark when induced with $NaNO_3$.
Without access to light, many PNSB can transform energy via anaerobic respiration. We tested
whether $NaNO_2$ or N_2O could support anaerobic respiration by CGA009 in the dark. Acetate and
butyrate were chosen as two carbon sources that are metabolized via similar pathways but
present the cell with different amounts of electrons (17). Unlike phototrophic conditions,
supplementation with either 0.3 or 1 mM $NaNO_2$ did not lead to observable growth in the dark
with either acetate or butyrate within at least 15 days. However, N_2O plus $NaNO_3$ supported
growth with either acetate or butyrate (Fig. 5). Growth was slower with acetate than with
butyrate (doubling time ± SD = 88 ± 2 h vs 51 ± 3, respectively).

Fig. 5. Anaerobic respiration by CGA009 with N₂O in the dark with acetate (A) or butyrate (B). Single representatives of similar trends observed for three biological replicates are shown for each condition. Cultures had a 100% Ar headspace or 100% N₂O headspace as indicated.

Fig. 6. Denitrification reductase genes in PNSB and related non-phototrophs. A gene was considered present if a predicted gene product exhibited >25% amino acid identity over 50% of its length relative to the query sequence. All subunits for a given enzyme are needed for the activity of a given reductase. The phylogenetic tree was built using PhyloT v2 (phylot.biobyte.de) based on NCBI taxonomy.

***R. capsulatus* SB 1003 requires NO$_2^-$ to induce growth on N$_2$O.** We wondered if induction of
N$_2$O reductase by non-catalyzable denitrification intermediates occurs in other bacteria with
partial denitrification pathways. We thus surveyed the genomes of various PNSB and non-
phototrophic relatives using BlastP analysis with the same query denitrification reductase genes
as for *R. palustris* in Figure 1 (Fig. 6). *R. capsulatus* SB 1003 stood out as an easily cultivatable
and phylogenetically distant PNSB that has N$_2$O reductase but no other denitrification genes
(Fig. 6 and 7A). In agreement with the genome annotation, SB 1003 phototrophic growth on
butyrate was not supported by NaNO$_3$ (Fig. 7B). NaNO$_2$ also did not support growth, though SB
1003 was sensitive to NaNO$_2$ concentrations > 0.5 mM (Fig. 7C), which would likely be needed
to see a clear OD increase. Similar to CGA009, N$_2$O alone did not lead to phototrophic growth
on butyrate. However, supplementation with 100 μM NaNO$_2$, but not NaNO$_3$, stimulated
phototrophic growth on butyrate with N$_2$O (Fig. 7B). The same was true of growth in the dark on
succinate (Fig. 7D), a condition chosen to test whether the same regulatory trend held true during
anaerobic N$_2$O respiration in the dark.
211

**Fig. 7. Activation of the single *R. capsulatus* SB 1003 denitrification step, N$_2$O reduction,
requires NO$_2^-$. A.** SB 1003 N$_2$O reductase genes (*nos*) are on a plasmid. A possible regulator,
NnrR, is encoded on the chromosome. Gene products: blue, N$_2$O reductase; green, accessory
proteins for electron transport, enzyme cofactor assembly, etc.; orange, denitrification
regulators; gray, unknown function. Numbers indicate nucleotide positions. **B.** Phototrophic
growth with butyrate and various denitrification intermediates. **C.** Phototrophic growth with
succinate and various NaNO$_2$ concentrations to determine the toxicity limit. **D.** Chemotrophic
growth with succinate with N$_2$O as an electron acceptor for anaerobic respiration. **B-D.** Single
representatives of similar trends observed for three biological replicates are shown for each
condition except for the test of NaNO$_2$ toxicity in panel C. Cultures had a 100% Ar headspace
unless N$_2$O is indicated (100% N$_2$O).

Discussion

We determined that non-catalyzable denitrification intermediates were required to induce the final denitrification step of N_2O reduction in two phylogenetically distant PNSB. This regulatory response to non-catalyzable intermediates could be the result of gene loss from an ancestor that had a complete denitrification pathway; that is, reductase genes have been lost but the regulatory network remains. For CGA009, gene loss seems more plausible than an alternative scenario of gene gain through horizontal gene transfer given the number of genes present for both catalysis and regulation (Fig. 1B). The spectrum of denitrification activities and gene content also varies for different *R. palustris* strains (21-23) (Fig. 6). However, a combination of gene loss and gene gain, or even gene duplication could have shaped the CGA009 denitrification inventory. Gene duplication could explain the existence of two nitrite reductase genes (Fig. 1B), that share 82% nucleotide sequence identity.

SB 1003 could be an example of gene gain given that the N_2O reductase gene cluster is encoded on a plasmid. How this cluster is regulated in response to NO_2^- remains unclear. Others have reported that unsequenced *R. capsulatus* strains reduce NO_3^- (7), and one isolate carried out complete denitrification (24). If these strains are truly close relatives of SB 1003, then perhaps a common ancestor was capable of denitrification and some regulatory genes were maintained in SB 1003. Alternatively, an unrelated regulatory feature might have evolved to control N_2O reductase activity (25).

The requirement of NO_2^-, but not NO_3^-, for SB 1003 growth with N_2O is unexpected. To the best of our knowledge, there are no clear examples of NO_2^- regulation of N_2O reductase. SB 1003 has several CRP/Fnr-family transcriptional regulator genes encoded in its chromosome with >25% amino acid identity to denitrification regulators like *P. denitrificans* FnrP (RCAP_rcc02493; 74% identity) and *Pseudomonas aeruginosa* Dnr/NnrR (RCAP_rcc00107; 36% identity) (Fig. 7A). However, these regulators respond to NO. It is possible that some of the NO_2^- we added was converted to NO by an unknown, and perhaps non-specific, enzyme activity and then NO activated N_2O reductase via the NnrR regulator (26).

One can also question why bacteria would lack a complete denitrification pathway. Denitrification genes could be lost if their activity is neutral or detrimental to fitness. NO_3^- reduction by Nar could be detrimental as the only denitrification enzyme with a cytoplasmic active site; the rest of the enzymes operate in the periplasm (1, 2, 27). As such, generation of toxic NO_2^- in the cytoplasm by Nar could lead to a selective disadvantage. Operating Nar during phototrophic generation of a proton motive force could also be more susceptible to backpressure (22) than the periplasmic denitrification steps. Nar eliminates protons in the cytoplasm, thus contributing to the proton gradient, whereas the other denitrification steps eliminate protons in the periplasm, detracting from the gradient (27). Thus, light-driven proton translocation could impose a thermodynamic impediment on Nar and thereby slow growth if, for example, the organism is using denitrification to dispose of electrons. However, these explanations do not explain why a phototroph would not have Nap, which reduces NO_3^- in the periplasm. Our survey revealed several examples of Nar or Nap in PNSB (Fig. 6).

Loss of denitrification genes could also be neutral or advantageous when within a community of
denitrifiers. Such scenarios are predicted by the Black Queen Hypothesis, wherein a gene
function that benefits other community members leads to loss of that function in a subset of
beneficiary community members (28). For example, if denitrification intermediates are reliably
generated by a neighbor and not all steps are needed for energy transformation and/or electron
balance, then it could be beneficial to lose other steps rather than incur the cost of synthesizing
the associated enzymes. Dispensing of NO_2^- and NO reductases could also be beneficial if NO_2^-
and NO are reliably removed by other community members. Membership within a denitrifying
community could also explain why non-catalyzable intermediates stimulate N_2O reduction, as
there would be little pressure to evolve responses to N_2O alone if N_2O is always found with other
intermediates of denitrification.
269
Partial denitrification pathways are both variable and widespread in PNSB (Fig. 6). This trend
could very well be true outside of PNSB and their close relatives. Despite decades of
investigation into the activity and regulation of denitrification, there has been comparatively little
research on partial denitrification pathways (2). Further investigation into the regulation of
partial denitrification pathways is needed, as is their potential impact on N_2O greenhouse gas
emissions from agricultural fertilizers.
276
277 **Methods.**
278
Strains. *R. palustris* CGA009 is a chloramphenicol-resistant type strain derived from CGA001
(8). The Calvin cycle mutant Δ*cbbLSMP*::km^R (CGA4008) was constructed by deleting *cbbLS*,
encoding ribulose-1,5-bisphosphate carboxylase (Rubisco) form I, in a previously described
mutant lacking Rubisco form II (Δ*cbbM*; CGA668; (20)) via introduction of the suicide vector
pJQΔ*cbbLS* (29) by conjugation with *E. coli* S17 as described (29, 30). The gene encoding
phosphoribulokinase, *cbbP*, was then deleted in the resulting strain (Δ*cbbLSM*; CGA4006) by
introducing the suicide vector pJQΔ*cbbP*::km^r (29), as above, to generate the Δ*cbbLSMP*::km^R
strain, CGA4008. All strain genotypes were verified by PCR and Sanger sequencing. The NifA*
derivative of CGA4008 with constitutive nitrogenase activity/H_2 production (CGA4011) was
described previously (29). The elimination of three genes unique to the Calvin cycle greatly
decreases the odds of enriching for suppressor mutations. *R. capsulatus* SB1003 was provided
courtesy of Carl Bauer (Indiana University).
291
Growth conditions. Strains were routinely cultivated in 10 ml photosynthetic medium (PM)
(31) in 27-ml anaerobic test tubes. PM was made anaerobic by bubbling tubes with 100% Ar
then sealing with rubber stoppers and aluminum crimps prior to autoclaving. After autoclaving,
tubes were supplemented with either 20 mM sodium acetate, 10 mM sodium butyrate, or 10 mM
sodium succinate from 100X anaerobic stock solutions. SB1003 cultures were also supplemented
with 0.1 µg/ml nicotinic acid, 0.2 µg/ml riboflavin, and 1.3 µg/ml thiamine-HCl. $NaNO_2$ or
$NaNO_3$ were added as indicated in the text. For conditions with N_2O, tubes were flushed with
100% N_2O through a 0.45 µm syringe filter and needle after all liquid supplements were added.
A second needle was used for off-gassing. Cultures were inoculated with a 1% inoculum from
starter cultures grown phototrophically in anaerobic PM with succinate, except for the
experiment testing Calvin cycle mutants (Fig. 3C) in which all starter cultures were grown
aerobically in 3 ml PM with succinate in the dark. Aerobic conditions were chosen for these

starter cultures to accommodate the Δ*cbbLSMP*::km^R mutant (CGA4008) that requires an
electron sink to grow.
306
Analytical procedures. Culture growth was monitored via optical density at 660 nm (OD_{660})
using a Genesys 20 spectrophotometer (Thermo-Fisher, Waltham, MA, USA) directly in culture
tubes without sampling. Specific growth rates were calculated using OD_{660} values between 0.1
and 1.0 where cell density and OD are linearly correlated. H_2 was sampled from culture
headspace using a gas-tight syringe and analysed using a Shimadzu GC-2014 gas chromatograph
as described (32).
313
Bioinformatics. Denitrification reductase gene inventories were determined using NCBI's
protein BLAST (12) to search for query sequences from *P. denitrificans* PD122 (Accession
numbers) NarG (GEK69895), NarH (CAA81215), NarI (CAA81217), NapA (CAA85346),
NapB (GEK66554), NirS (GEK68816), NirK (GEC49504), NorB (BAA32546), NorC
(BAA32545), and NosZ (GEK69878). Denitrification regulatory genes inventories were
similarly determined using query sequences from *P. denitrificans* PD122 FnrP (WP_041529894
(33)) and Nnr (WP_041529894 (33)), *B. japonicum* USDA110 NasT (BAC49838) and NasS
(BAC49837), and *P. aeruginosa* PAO1 Dnr/NnrR (BAA08744). The phylogenetic tree was built
using PhyloT v2 (phylot.biobyte.de) based on NCBI taxonomy and visualized using iTOL v6
(itol.embl.de).
324
325   **Acknowledgements**
326

327   This work was supported in part by a National Science Foundation CAREER
328   award (MCB-1749489NSF), the Division of Chemical Sciences, Geosciences, and Biosciences,
329   Office of Basic Energy Sciences, U.S. Department of Energy (DOE), through Grant DE-FG02-
330   05ER15707, the Office of Science (BER), U.S. Department of Energy, through Grant DE-FG02-
331   07ER64482, and the Indiana University College of Arts and Sciences.

332
333   **References**
334

1.   Zumft WG. 1997. Cell biology and molecular basis of denitrification. Microbiol Mol Biol
Rev 61:533-616.
2.   Zumft WG, Kroneck PM. 2007. Respiratory transformation of nitrous oxide (N_2O) to
dinitrogen by Bacteria and Archaea. Adv Microb Physiol 52:107-227.
3.   Lundberg JO, Weitzberg E, Cole JA, Benjamin N. 2004. Nitrate, bacteria and human
health. Nat Rev Microbiol 2:593-602.
4.   Gold A, Addy K, David M, Schipper L, Needleman B. 2014. Artificial sinks:
opportunities and challenges for managing offsite nitrogen losses. J Contemp Water Res
Educ 151:9-19.
5.   Barnard R, Leadly P, Hungate B. 2005. Global change, nitrification, and denitrification:
A review. Global Biogeochem Cycles 19:327-338.
6.   McCully AL, Onyeziri MC, LaSarre B, Gliessman JR, McKinlay JB. 2020. Reductive
tricarboxylic acid cycle enzymes and reductive amino acid synthesis pathways contribute
to electron balance in a *Rhodospirillum rubrum* Calvin-cycle mutant. Microbiol 166:199-
211.

7.   Richardson DJ, King GF, Kelly DJ, McEwan AG, Ferguson SJ, Jackson JB. 1998. The
role of auxiliary oxidants in maintaining redox balance during phototrophic growth of
Rhodobacter capsulatus on propionate or butyrate. Arch Microbiol 150:131-7.
8.   Larimer FW, Chain P, Hauser L, Lamerdin J, Malfatti S, Do L, Land ML, Pelletier DA,
Beatty JT, Lang AS, Tabita FR, Gibson JL, Hanson TE, Bobst C, Torres JLTy, Peres C,
Harrison FH, Gibson J, Harwood CS. 2004. Complete genome sequence of the
metabolically versatile photosynthetic bacterium *Rhodopseudomonas palustris*. Nat
Biotechnol 22:55-61.
9.   Luxem KE, Kraepiel AML, Zhang L, Waldbauer JR, Zhang X. 2022. Corrigendum.
Carbon substrate re-orders relative growth of a bacterium using Mo-, V-, or Fe-
nitrogenase for nitrogen fixation. Environ Microbiol 24:2170-2176.
10.   Luxem KE, Kraepiel AML, Zhang L, Waldbauer JR, Zhang X. 2020. Carbon substrate
re-orders relative growth of a bacterium using Mo-, V-, or Fe-nitrogenase for nitrogen
fixation. Environ Microbiol 22:1397-1408.
11.   Oda Y, Larimer FW, Chain PSG, Malfatti S, Shin MV, Vergez LM, Hauser L, Land ML,
Braatsch S, Beatty JT, Pelletier DA, Schaefer AL, Harwood CS. 2008. Multiple genome
sequences reveal adaptations of a phototrophic bacterium to sediment
microenvironments. Proc Natl Acad Sci USA 105:18543-18548.
12.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment
search tool. J Mol Biol 215:403-10.
13.   Gaimster H, Alston M, Richardson DJ, Gates AJ, G R. 2018. Transcriptional and
environmental control of bacterial denitrification and N_2O emissions. FEMS Micobiol
Lett 365:fnx277.
14.   Arai H, Mizutani M, Igarashi Y. 2003. Transcriptional regulation of the nos genes for
nitrous oxide reductase in *Pseudomonas aeruginosa*. Microbiol 149:29-36.
15.   Sanchez C, Itakura M, Okubo T, Matsumoto T, Yoshikawa H, Gotoh A, Hidaka M,
Uchida T, Minamisawa K. 2014. The nitrate-sensing NasST system regulates nitrous
oxide reductase and periplasmic nitrate reductase in *Bradyrhizobium japonicum*. Environ
Microbiol 16:3263-74.
16.   Sanchez C, Mitsui H, Minamisawa K. 2017. Regulation of nitrous oxide reductase genes
by NasT-mediated transcription antitermination in *Bradyrhizobium diazoefficiens*.
Environ Microbiol Rep 9:389-396.
17.   McKinlay JB, Harwood CS. 2011. Calvin cycle flux, pathway constraints, and substrate
oxidation state together determine the H2 biofuel yield in photoheterotrophic bacteria.
mBio 2:e00323-10.
18.   Härtig E, Zumft WG. 1999. Kinetics of *nirS* expression (cytochrome cd1 nitrite
reductase) in *Pseudomonas stutzeri* during the transition from aerobic respiration to
denitrification: evidence for a denitrification-specific nitrate- and nitrite-responsive
regulatory system. J Bacteriol 181:161-6.
19.   Sabaty M, Schwintner C, Cahors S, Richaud P, Vermeglio A. 1999. Nitrite and nitrous
oxide reductase regulation by nitrogen oxides in *Rhodobacter sphaeroides* f. sp.
denitrificans IL106. J Bacteriol 181:6028-32.
20.   McKinlay JB, Harwood CS. 2010. Carbon dioxide fixation as a central redox cofactor
recycling mechanism in bacteria. Proc Natl Acad Sci USA 107:11669-11675.

21. Klemme J-H, Chyla I, Preuss M. 1980. Dissimilatory nitrate reduction by strains of the
facultative phototrophic bacterium *Rhodopseudomonas palustris*. FEMS Microbiol Lett
9:137-140.
22. McEwan AG, Greenfield AJ, Wetzstein HG, Jackson JB, Ferguson SJ. 1985. Nitrous
oxide reduction by members of the family *Rhodospirillaceae* and the nitrous oxide
reductase of *Rhodopseudomonas capsulata*. J Bacteriol 164:823-30.
23. Rayyan A, Meyer T, Kyndt J. 2018. Draft whole-genome sequence of the purple
photosynthetic bacterium *Rhodopseudomonas palustris* XCP. Microbiol Resour Announc
7:e00855-18.
24. Richardson DJ, Bell LC, Moir JWB, Ferguson SJ. 1994. A denitrifying strain of
Rhodobacter capsulatus. FEMS Micobiol Lett 120:323-8.
25. Perez JC, Groisman EA. 2009. Evolution of transcriptional regulatory circuits in bacteria.
Cell 138:233-44.
26. Bergaust L, van Spanning RJM, Frostegard A, Bakken LR. 2012. Expression of nitrous
oxide reductase in *Paracoccus denitrificans* is regulated by oxygen and nitric oxide
through FnrP and NNR. Microbiol 158:826-834.
27. Chen J, Strous M. 2013. Denitrification and aerobic respiration, hybrid electron transport
chains and co-evolution. Biochim Biophys Acta 1827:136-44.
28. Morris JJ, Lenski RE, Zinser ER. 2012. The Black Queen Hypothesis: Evolution of
dependencies through adaptive gene loss. mBio 3:e00036-12.
29. Gordon GC, McKinlay JB. 2014. Calvin cycle mutants of photoheterotrophic purple
nonsulfur bacteria fail to grow due to an electron imbalance rather than toxic metabolite
accumulation. J Bacteriol 196:1231-7.
30. Rey FE, Oda Y, Harwood CS. 2006. Regulation of uptake hydrogenase and effects of
hydrogen utilization on gene expression in *Rhodopseudomonas palustris*. J Bacteriol
188:6143-6152.
31. Kim M-K, Harwood CS. 1991. Regulation of benzoate-CoA ligase in
Rhodopseudomonas palustris. FEMS Microbiol Lett 83:199-203.
32. Huang JJ, Heiniger EK, McKinlay JB, Harwood CS. 2010. Production of hydrogen gas
from light and the inorganic electron donor thiosulfate by *Rhodopseudomonas palustris*.
Appl Environ Microbiol 76:7717-7722.
33. van Spanning RJ, De Boer AP, Reijnders WN, Westerhoff HV, Stouthamer AH, van der
Oost J. 1997. FnrP and NNR of *Paracoccus denitrificans* are both members of the FNR
family of transcriptional activators but have distinct roles in respiratory adaptation in
response to oxygen limitation. Mol Microbiol 23:893-907.
429