RESEARCH ARTICLE TITLE

Genomic and transcriptomic analysis of sacred fig (*Ficus religiosa*)

KL Ashalatha [1] , KP Arunkumar[2], Malali Gowda[1, 3*]

*****CORRESPONDENCE:** malali.gene@gmail.com

[1,3]Malali Gowda

The University of Trans-Disciplinary Health Sciences and Technology

Yelahanka, Bengaluru, 560064, India

Abstract

Background: Peepal/Bodhi tree (*Ficus religiosa* L.) is an important, long-lived keystone

ecological species. Communities on the Indian subcontinent have extensively employed the

plant in Ayurveda, traditional medicine, and spiritual practices. The Peepal tree is often

thought to produce oxygen both during the day and at night by Indian folks. The goal of our

research was to produce molecular resources using whole-genome and transcriptome

sequencing techniques.

Results: The whole genome of the Peepal tree was sequenced using two next-generation

sequencers like Illumina HiSeq1000 and MGISEQ-2000. We assembled the draft genome of

406Mb, using a hybrid assembly workflow. The genome annotation resulted in 35,093 protein-

coding genes; 53% of its genome consists of repetitive sequences. To understand the

physiological pathways in leaf tissues, we analyzed photosynthetically distinct conditions:

bright sunny days and nights. The RNA-seq analysis supported the expression of 26,691 genes.

The leaf transcriptomic analysis of the diurnal and nocturnal periods revealed the expression

of significant genes involved in the carbon-fixation pathway.

Conclusions: This study presents a draft hybrid genome assembly for *F. religiosa* and its functional annotated genes. The genomic and transcriptomic data-derived pathways are obtained for future studies on the Peepal tree.

Keywords: *Ficus religiosa*, Peepal, Bodhi, Hybrid genome, Transcriptome, Carbon-fixation pathway

Background

The Peepal tree (*Ficus religiosa* L.) is a sacred fig, hemi- epiphyte that belongs to the Moraceae family and has a diploid sporophytic count (2n = 26) [1]. It is known to be a long-lived deciduous species related to the 755 fig species widespread worldwide [2]. The Peepal tree is a cosmopolitan species, having value for cultural and spiritual practices in Buddhism, Hinduism, and Jainism. It is popularly called the Bodhi tree, where Buddha is believed to have meditated and attained spiritual enlightenment underneath this tree. Hence, the culture is spread across Asia and it has been worshipped. Peepal has several vernacular names, like Pippali, Ashwatha, Arali, and so on; it is frequently found together with the Neem tree near Indian Temple [3]. Generally, the Peepal tree has a special significance in communities across India that are believed to produce oxygen day and night. They have a special type of stomata called sunken, giant, or hydathode at the lower leaf epidermis of *F. religiosa*. These are larger than the normal stomata and occur over the veins or are mixed with normal stomata, it indicates that such stomata hold gaseous and water molecules for a longer time [4]. According to our knowledge, there is no scientific evidence to claim oxygen production during a dark period.

In Ayurveda, the Peepal tree has been classified as a Rasayana (a type of drug), whereby rejuvenators and antioxidants aid in relieving the body's stress [5]. Peepal tree alleviates Pitta

and Kapha (Ayurvedic classifications), hence prescribed for treatment of the disorders like

respiratory and inflammatory disorders, ulcers, stomatitis, hiccup, arthritis, gout, skin diseases,

bone fracture, diabetes, etc., [5]. In animal models such as rats, the Peepal tree has been tested

for the treatment of neurodegenerative disorders such as Parkinson's disease and Huntington's

disease [6], [7], as well as anti-ulcer activity in albino mice [8].

Next-generation sequencing (NGS) technologies have accelerated the generation of draft

genome sequences of Moraceae plant species, the genome size of *Morus notabilis* is 330Mb

[9], 333Mb for *Ficus carica* [10], 436Mb for *F. microcarpa*, and 370Mb for *F. hispida* [11].

The genome sequencing of non-model plant species Peepal was first mentioned in The Neem

Genome book chapter [3]. A recent study has generated the genomic resource of *F.*

religiosa (332Mb) and *F. benghalensis* (392Mb). However, they generated a limited size of

genome assembly and genes (23,929) for the Peepal tree when compared with the present study

[12]. Recently, several studies have attempted sequencing non-model plant species like

pineapple (*Ananas comosus*) and *Kalanchoë* species revealing the gene expressions of the

Crassulacean acid metabolism pathway [13] and [14]. The whole-genome sequencing of

common or crystalline ice plants (*Mesembryanthemum crystallinum*) has been shown to switch

from Calvin- Benson Cycle (C3) to CAM photosynthesis under a salt stress [15]. Using RNA-

seq, the study described by comparing both species with and without the C4 trait and different

tissues within a C4 plant suggests the ways of integration into the underlying C3 metabolism

[16]. These previous study findings and other physiological features of the Peepal tree enabled

us to characterize the pathways in the present work.

Despite its ecological, medicinal, cultural, and historic importance, the molecular biology and

genomics studies on the Peepal tree are scanty. Hence, Peepal is relevant to traditional

medicinal practices and Buddha's meditation, we hypothesized to elucidate the genomic sequence and to study the transcriptome of photosynthetic tissue (leaf tissue) in diurnal and nocturnal periods. The objective of the present study was to generate a genome sequence and annotate genes for the Peepal tree. The transcriptomic analysis has been undertaken to identify the expression of genes in the diurnal and nocturnal periods for photosynthetic activity using a molecular approach. In this study, we aim to characterize the genes involved in various physiological, biochemical metabolic, and other pathways. And also, a comparative genomic study has been carried out to analyze the Peepal tree relationship between closely related species of its Moraceae family.

Results

***De novo* hybrid assembly using Illumina and MGI short reads**

We used two next-generation technology platforms to sequence the whole genome of the fig species, the Peepal tree. Paired-end reads from Illumina HiSeq1000 and MGISEQ-2000 were 266 and 645 million respectively; a total data of 88.44 billion high-quality bases (Quality>20) used for genome assembly. A hybrid assembly was performed using a sequencing depth of 65.5X Illumina reads and 158.86X MGI reads. The raw data details are given (Additional file 1: Table S1.1). We evaluated the distribution of k-mers in both Illumina and MGI reads to estimate the genome size, resulting in a genome size of 393-Mb and 273-Mb respectively. The combination of Illumina and MGI reads was used for assembling the genome. Hybrid genome assembly yielded a genome of 406-Mb. The contig N50 length is 5,817 bp and the largest contig length is 148Kb. The GC content of the Peepal tree genome is 34.23%. The gap-closing step was performed for the hybrid assembly. There were 35,811 (5.5%) misassembled contigs and 6,04,807 (94.4%) truly assembled contig sequences in the final assembled genome. The workflow of the genome assembly is represented in (Additional file 2: Figure S1). The statistics

of assembly contigs and scaffolds are shown (Additional file 3: Table S1.2) and the final

scaffold assembly of the genome is given in Table 1.

The completeness of the Peepal tree genome assembly was assessed with the BUSCO tool. The

results showed that 76.5% (232 out of 303 genes) and 84.1% (1210 out of 1440 genes) of genes

were conserved as single-copy orthologs in eukaryotic and plant universal data sets,

respectively. Out of 232 complete genes in the Eukaryota database, 214 are single-copy

orthologs, 18 are duplicates, 57 are fragmented and 14 are missing; out of 1210 complete genes

in the Embryophyte database 1173 are single-copy orthologs, 37 are duplicates, 105 are

fragmented and 125 are missing (Additional file 4: Figures S2A and S2B). The transcriptome

reads were aligned with the assembled genome and found that 99.46% of reads mapping and

88.25% of paired reads were mapped.

Genome and pathways annotation

We identified 35,093 protein-coding genes with the complete structures in the Peepal tree

genome (Additional file 5: Text file S1.1 and Additional file 6: Text file S1.2). RNA-seq data

from two leaf tissue samples of the Peepal tree and alternative reference ESTs from *Morus*

notabilis and *Arabidopsis thaliana* protein sequences were used as protein homology evidence

during genome annotation. Based on a homology search using BLASTN, out of 35,093 genes

predicted, 32,255 genes (91.9%) were having evidence from transcriptome assembly. About

76.3% of RNA-seq reads from the day and night leaf tissue samples were mapped to the

annotated genes in the Peepal tree.

Based on the sequence similarities, the complete set of annotated genes and their amino acid

sequences were used in the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway

analysis [17]. This result showed the pathways in metabolism, biosynthesis of secondary metabolites, genetic and environmental information processing, and signal transduction pathway were common as several others. The top 5 highest gene count for pathways like Ribosome (123 genes), Spliceosome (96 genes), Oxidative phosphorylation (86 genes), Thermogenesis (82 genes), and RNA transport (74 genes) were found. As importantly candidate genes were also found for human disease pathways like Huntington's disease (68 genes), Parkinson's disease (57 genes), Alzheimer's disease (55 genes), and others listed in (Additional file 7: Table S2).

The plant model Arabidopsis encodes several orthologues of human proteins that function in mechanisms similar to those in other eukaryotes [18] [19]. Previous findings showed that 70% of oncogenes involved in cancer have Arabidopsis, 67% in *D. melanogaster*, 72% in *C. elegans,* and 41% in *S. cerevisiae*. This 'disease gene' similarity is comparable to that observed in other model organisms [20]. Arabidopsis and many other model systems research have led to the discovery or analysis of genes and processes important to human health. Previous studies showed that the Peepal tree has been tested for the treatment of neurodegenerative disorders such as Parkinson's disease and Huntington's disease [6], [7], as well as anti-ulcer activity in animal models. [8]. Another study showed that methanol extract of *F. religiosa* has proven anti-inflammatory properties in LPS-induced activation of BV2 microglial cells, and it might have therapeutic potential for various neurodegenerative diseases [21]. In the present work, we found that Peepal genes show similarities with human disease pathways, which can be utilized to further understand the traditional medicinal practices and Buddha's meditation practices. Thus, plant research opens up new frontiers in terms of drug development and treatment of diseases of great importance to human health. Plants seem to be part of this diverse portfolio of tools necessary to understand fundamental cellular processes.

Protein family and Gene Ontology analysis

The protein family (Pfam) ID and Gene Ontology (GO terms) were assigned to genes using an

InterProScan module [22]. Out of 35,093 genes, 24,163 consisted of Pfam IDs that were

distributed across 3759 types of Pfam domains, and their gene ontology (GO) terms were also

identified. The Pfam domain consisting of proteins that were large in the Peepal tree genome

included 3-Deoxy-D-manno-octulosonic-acid transferase, Ring finger domain, PPR repeat

family, Helix-loop-helix DNA-binding protein, DYW family of nucleic acid deaminases,

Lysine methyltransferase, Putative GTPase activating protein for Arf, Ankyrin repeats and

others were listed in (Additional file 8: Table S3).

Catalase is an antioxidant enzyme known to catalyze H_2O_2 into water and oxygen. The Catalase

gene (FRLM_016351-RA) and its isozyme CAT1 Catalase isozyme 1 (FRLM_016350-RA),

(FRLM_012250-RA) structural gene were identified in the genome of the Peepal tree. Two

catalase genes were identified in differential expression data, KatE enciphers as a

monofunctional catalase and KatG enciphers as a catalase-peroxidase. KatE is known as CatB

expression in the day (FPKM – 937.49) and night (FPKM – 1786.02) transcript abundance.

KatG expression in the day (FPKM – 162.03) and night (FPKM – 81.53) transcript abundance.

KatE gene has been involved in physiological pathways like Glyoxylate and dicarboxylate

metabolism, Tryptophan metabolism, MAPK signaling pathway – plant, FoxO signaling

pathway, serine-pyruvate transaminase, and KatG Tryptophan metabolism, Tyrosine

metabolism, Biosynthesis of secondary metabolism and Drug metabolism.

Identification of homologous, orthologous, and singleton genes

To understand the homologous genes and the orthologous gene evolution, and relationships

across the species, we performed the homologous and orthologous gene detection for the

Peepal tree with the other 5 species. Homologous gene identification and orthologous

clustering of the proteomes of six species, including the model organism *A. thaliana*, *M.*

notabilis closest relative species of Ficus*,* and other closely related species of the Moraceae

family were selected for the analysis. The number of homologous genes identified for *F.*

religiosa proteome sequences (35,093) was 29516 in *Arabidopsis thaliana*, 29924 in *Morus*

notabilis, 29750 in *Cannabis sativa*, 29909 in *Prunus persica*, and 29830 in *Ziziphus jujuba*

based on homology analysis. *F. religiosa, A. thaliana*, *M. notabilis, C. sativa, P. persica*, and

Z. jujuba form a cluster of 24,310 orthologous genes and which are conserved within the

species. The number of specific orthologous gene clusters in *F. religiosa, A. thaliana, M.*

notabilis, C. sativa, P. persica, and *Z. jujuba* were 15016, 16170, 16517, 15921, 16655, and

17235 respectively. However, 1184 single-copy gene clusters were found across the six species

and specific singletons for *F. religiosa, A. thaliana, M. notabilis, C. sativa, P. persica,* and *Z.*

jujuba were 10154, 4469, 2284, 1912, 1802, 4209, respectively (Figure 1).

The identified single-copy clusters were able to suggest the taxonomic and phylogenetic

relationships among a group of species. Based on the similarity of proteomes and single-copy

orthologous clustering, we deduced the phylogenetic tree for *F. religiosa* and the other five

species. A multiple sequence alignment (MSA) and Neighbour-Joining (NJ) method were used

for constructing an evolutionary phylogenetic tree. It was found that *F. religiosa* is closely

related to *M. notabilis* by having more similarities in their proteome as they are evolving from

the Moraceae family, followed by *C. sativa, Z. jujuba, P. persica,* and *A. thaliana* (Additional

file 9: Figure S3)*.*

Comparative analysis of Peepal tree genome

We aligned the genome of the Peepal tree with the three Moraceae family members like *F.*

carica, *F. microcarpa*, and *M. notabilis*. Comparison of our assembly against these genomes

resulted in a mapping of 88.62% to *F. carica*, 89.6% to *F. microcarpa*, and 46.9% to *M.*

notabilis. The results showed the similarity of the Peepal genome is closer to the genus Ficus

(*F. carica* and *F. microcarpa*) and also relatively closer to the genus Morus of the same family.

The statistics of genome sequence alignment of *F. religiosa* against *F. carica*, *M. notabilis*, *F.*

microcarpa genomes (Additional file 10: Text file S2).

Repeats in the genome of the Peepal tree

Repeat library building and repeat identification were performed using the ReapeatModeller

and RepeatMasker tools (www.repeatmasker.org) respectively. *De novo* repeat identification

resulted in 53.55% (269.62Mb) repetitive sequences in the Peepal tree genome. The RNA

elements, long terminal repeats (LTR) high constitute about 5% of repeats and 43.71% of the

repeats did not belong to any of the annotated repeats families. The 53.55% of repetitive

sequences in the Peepal tree genome are closest to its Moraceae family species, 47% are found

in the closest species mulberry (*M. notabilis*), 46.5% in *F. microcarpa*, and 48.9% in *F. hispida*.

The repetitive sequences were classified into known categories, such as LINE1 (0.19%), long

terminal repeat retrotransposon (5.09%), DNA transposons (1.09%), and simple repeats

(3.25%) and unclassified (43.71%) (Additional file 11: Table S4).

Simple sequence repeats (SSRs)

We predicted SSRs from the assembled Peepal tree genome. In total, 7,99,992 SSRs were

identified on 2,67,593 sequences, which are composed of mono- (6,06,169), di- (1,43,113),

tri- (34,327), tetra- (11,791), penta- (2,911), and hexa- (1,681) type repeats (Additional file 12:

Table S5.1). Among mono repeats, the 'A/T' (73.91%) type was the highest followed by 'C/G'

(1.87%). Similarly, the 'AT /TA', 'AG/CT', 'AC/GT', and 'CG/CG' types of di repeats were

in 9.8%, 2.76%, 1.41%, and 0.09% fractions, respectively. 'AAT/ATT', 'AAG/CTT'

'ATA/TAT', 'TTA/TTC', and 'GAA/TAA', were the most abundant tri repeats and 'AAAT'

was most predominant in tetra repeats. The detailed distribution of all types of repeats and

statistics is shown in (Additional file 13: Table S5.2 and Additional file 14: Table S5.3).

Transcription Factors (TFs)

Transcription factors act in regulating gene expression due to several external and internal

signals by activating or suppressing downstream genes. The MAKER annotated protein

sequences of Peepal tree genome assembly were used for BLAST analysis with the Plant

Transcription Factor Database v5.0 [23] using the *A. thaliana* protein sequence as a reference.

A total of 1,264 protein sequences of genome annotated 35,093 protein-coding genes have

evidence for 56 families of Transcription factors (TFs) (Additional file 15: Table S6). The TFs

families include the abundant ERF, M-type MADS, ARF, DBB, MIKC MADS, WOX, C3H,

G2-like, MYB, TALE, B3, HB-other and MYB-related family proteins.

Transcriptome sequencing, assembly, and annotation

De novo transcriptome assembly was performed for the mature leaf samples of the Peepal tree

collected during the day and night periods. The assembly was performed for each sample and

also combined assembly for the reads of both samples. The combined transcriptome assembly

was performed and contributed to 152.8-Mb assembled bases with the N50 length of 2076 bp

and an average transcript length of 13.16 Kb and 42.17% GC content. The transcriptome

assembly and annotation workflow are given in (Additional file 16: Figure S4). The statistics

of assembly contigs and sequence assembled contigs are shown in (Additional file 17: Table

S7).

The *de novo* assembled transcript (1,16,038) sequences were processed for annotation. *De novo*

assembled transcripts were clustered to exclude the redundant transcripts and identified 26,479

unique transcripts sequences. The statistics of unigenes are given in Table 2. *De novo*

assembled transcripts and unigenes were annotated to find the structural and functional genes.

The protein families were identified for the uniquely characterized transcripts of RNA data.

Out of 26,691 transcripts, Pfam IDs for 19175 were distributed across 3977 types of Pfam

domains and their gene ontology terms were also identified (Additional file 18: Table S8).

The differential expressed genes (35,182) from the day and night periods leaf tissue samples

of the Peepal tree are given in (Additional file 19: Table S9.1) and these transcripts were used

for the pathway analysis. The top 272 highly up-regulated differential transcripts were given

for diurnal and nocturnal periods (Additional file 20: Table S9.2).

The transcription factors (TFs) were identified for the differentially expressed transcripts. From

the day sample, the 2 transcripts coded for specific TFs like C3H family protein and nuclear

factor Y, subunit A7 (NF-YA7), and in the night period sample, the 6 transcripts coded for

specific TFs like ERF family protein, CONSTANS-like 2, MYB-related family protein

(Additional file 21: Table S9.3).

Non-coding RNA genes in the Peepal tree genome

Based on a coding potential calculator (CPC), *de novo*-based assembled transcripts (26,691)

were further categorized into protein-coding (19,911) and non-coding (6,780). Based on

BLASTN analysis, out of 6,780 non-coding transcripts, 4219 transcripts targeted genomic

annotated genes and 2,561 remained as non-coding. A total of 30,973 Cufflinks assembled

transcripts (reference-based alignment with genome assembly) were further categorized into

protein coding (7,163) and non-coding (23,810). Out of 23,810 non-coding transcripts, 14605

were having alignment to genome annotated genes using BLASTN. Further, categorization of

specific day and night sample transcripts resulted in protein-coding 6,628 and 7,339 and non-

coding of 20,528 and 25,494 transcripts respectively. From those non-coding transcripts of day

and night, 18,893 and 22,232 transcripts got aligned MAKER-P predicted genes using

BLASTN. The remaining transcripts were considered to be non-coding transcripts, as we did

not find any match to predicted gene evidence to support them.

miRNAs: miRNAs are a major class of non-coding RNAs. Based on the homology search, we

have identified the microRNA precursors using the miRbase database (http://www.mirbase.org).

These microRNAs belong to MIR396, MIR2916, MIR156, MIR164, MIR6236, MIR166,

MIR168, and MIR395 families. Among the identified miRNAs, MIR408 was found specific to

the night period transcripts of the Peepal tree. On the genome, MIR 408 was found on the genes

like TPK5 Two-pore potassium channel 5, prfA peptide chain release factor 1, and also on

proteins of unknown function. MIR408 is a highly conserved microRNA in plants. miR408 has

enriched photosynthesis by mitigating the efficiency of irradiation utilization and the capacity

for carbon dioxide fixation [24].

In the *de novo*-based transcriptome data, the non-redundant unigenes were used to identify the

microRNAs MIR168 and MIR166, and their homologs were found on two of the transcripts.

We identified the miRNAs on genomic scaffolds based on mapping the transcriptome data to

the genome. This provides information on miRNAs specific to the day and night leaf tissue

transcriptome. We identified 23 and 25 pre-miRNA expressions in the day and night period

respectively (Additional file 22: Table S10.1). The statistics of transfer RNAs (tRNA) were

identified and their details are given in the (Additional file 23: Table S10.2).

Elucidation of carbon fixation pathway in Peepal tree

The study was conducted to analyze the gene expression patterns in the leaf tissues of the

Peepal tree under two conditions the diurnal (2 PM) and the nocturnal period (2 AM). Through

the pathway analysis, the candidate genes for carbon fixation pathways like Crassulacean Acid

Metabolism (CAM) pathway, Calvin-Benson cycle (C3) pathway, and C4 - Dicarboxylic

pathway were identified and estimated based on their transcript abundance. The transcriptome

data contained the 20 putative genes involved in the carbon fixation module of CAM, C3, and

C4 including for the key fructose-bisphosphate aldolase class I, fructose-1,6-bisphosphate,

phosphoenolpyruvate carboxylase (PEPC/PPC), phosphoenolpyruvate carboxylase kinase

(PPCK), NAD+ and NADP+, malate dehydrogenase (MDH), pyruvate orthophosphate

dikinase (PPDK) (Additional file 24: Table S11). The complete pathway map was found for

CAM and C4 cycles and three genes were missed from the C3 cycle. The three genes fructose-

6-phosphate phosphoketolase (EC 4.1.2.22) and phosphoketolase (EC 4.1.2.9) were purified in

Acetobacter xylinum [25] and sedoheptulokinase (EC 2.7.1.14) was showed in *Bacillus* species

[26]. These three genes were not found in the Peepal tree for the C3 cycle.

The differentially expressed genes from transcriptomic data were mapped to the reference

carbon fixation in photosynthetic organisms pathway on the KEGG database are provided in

(Additional file 25: Figure S5) [27], [28], and [29]. The diagrammatic representation of the

genes involved in the carbon fixation pathway is shown in (Additional file 26: Figure S6).

Ribulose-bisphosphate carboxylase (RuBP carboxylase or *rubisco*) small chain enzyme is

enriched in leaf tissue collected during the day (2 PM). Rubisco is the most abundant protein

in chloroplasts. The glyceraldehyde-3-phosphate dehydrogenase (NADP+) is enriched in day

sample leaf tissue, the enzyme responsible for the reversible conversion of glyceraldehyde 3-

phosphate to ribulose bisphosphate using ATP, the acceptor for CO2 and completes the Calvin-

Benson Cycle (C3 cycle) [30].

The signature genes responsible for the CAM cycle were expressed in the Peepal tree during

the night. The phosphoenolpyruvate carboxylase kinase (PPCK), NAD(P)-ME (maeB), and

Malate dehydrogenase (MDH) genes were highly enriched in the photosynthetic leaf tissue

collected during the night period than the day. It indicates that the Peepal tree adapts to the

CAM pathway and can fix nocturnal carbon dioxide using the PEP carboxylase (PEPC) enzyme

and accumulate malate by the enzyme malate dehydrogenase in the large vacuoles of their cells.

The transcriptomic genes from the Peepal tree were mapped completely to the KEGG pathway

of the CAM cycle.

In the C4 Dicarboxylic cycle, the high expression of glutamate-glyoxylate aminotransferase

enzyme (GGAT) in the leaf tissue collected during the night period (2 AM) indicates the

photorespiration in the Peepal tree. The carbon fixation begins in the mesophyll cells, where

CO2 is converted into bicarbonate. It adds the 3-carbon acid phosphoenolpyruvate (PEP) by

an enzyme called phosphoenolpyruvate carboxylase. The product of this reaction is the four-

carbon acid oxaloacetate, which is reduced to malate another four-carbon acid [31]. The second

highest expression is NADP-malate dehydrogenase (MDH), which converts the oxaloacetate

generated by PEPC to malate. The genes from Peepal tree differential expression data had a

complete map to the C4 cycle. The gene expression pattern of the carbon-fixation pathway in

the Peepal tree suggests that the plant switches between the C3, C4, and CAM cycles during

the diurnal and nocturnal periods. The Fragments Per Kilobase of transcript per Million

mapped reads (FPKM) and Trimmed Mean of M-values (TMM) values of the differential genes

expressed for carbon fixation pathway are shown in Figures 2A, B, C, and 3A, B, and C.

Discussion

In this study, the genomics and transcriptomics data were generated and annotated for the

keystone species Peepal tree (*F. religiosa*). This plant species is well known for Buddha's

enlightenment while he was meditating underneath this tree. This tree is worshipped as birth-

giving, regenerative, and medicinally valued for many diseases, and this culture is spread

across Asia. We used two Next-Generation Sequencing technologies to sequence the genome

of the Peepal tree and described the hybrid whole genome. The photosynthetic tissues of

distinct conditions (diurnal and nocturnal) were used for transcriptome sequencing and to

understand the genes, proteins, and molecular pathways. The assembled genome resulted in a

size of 406-Mb with 35,093 protein-coding genes, based on *ab initio*, homology-based, and

mRNA evidence used for annotation. The transcriptome analysis resulted in 26,691 unique

transcripts. The completeness of the Peepal tree genome assembly was confirmed based on

BUSCO analysis and comparative analysis of transcriptomic data. A total of 35,093 genes were

predicted, and 32,255 genes (91.9%) were having evidence from transcriptome assembly.

About 76.3% of RNA-seq reads were mapped to the annotated genes. This information has

provided confidence for the downstream analyses. The number of protein-coding genes

identified in the Peepal tree is related to other plant genomes sequenced over the decades.

Similarly, in our laboratory *Santalum album* (Sandalwood) genome was sequenced with the

prediction of 38,119 protein-coding genes based on mRNA and peptide evidence [32].

We performed the downstream analysis of genomic and transcriptomic data to understand

molecular changes in the diurnal and nocturnal leaf tissue samples of the Peepal tree. Analysis

of microRNAs, TFs, and carbon fixation pathways has led to the understanding of

photosynthesis in the Peepal tree. The long non-coding RNAs have important roles in the

metabolic, physiological, and biological processes of the cell. MIR408 has been reported to

target various blue copper protein members and phytocyanin family [33], [34]. In Arabidopsis,

miR408 responds to copper deficiency and light [35]. In *Oryza sativa* MiR408 plants were

efficient at saving and converting light energy into sugars, suggesting that miR408 might

promote grain yield by regulating the phytocyanin (PC) content and photosynthesis by down-

regulating UCL8 uclacyanin (UCL) gene of the phytocyanin family. So, MIR408 found

specific expression in transcripts of night period leaf tissue of the Peepal tree indicating the similar conversion of light energy and accumulation of sugars in the plant during the dark. It also aids in photosynthesis by enhancing carbon fixation.

Transcription factors like C3H family proteins were identified by their motif of three cysteines and one histidine residue. They play an important role in regulating growth, developmental processes, and environmental responses in the plant's [36]. In plants, the nuclear factor-YA has a role in drought stress responses. In rice, NF-YA7 is involved in the drought tolerance pathway which is independent of the Abscisic acid manner [37]. Expression of C3H and NF during the day could influence plant growth and development in the Peepal tree. The Ethylene response factor ERF105 showed the cold-regulated transcription factor gene of Arabidopsis and also functions in the freezing tolerance and cold acclimation [38]. In the Peepal tree, ERF, MYB - related family proteins like REVEILLE 1 (RVE1) and late elongated hypocotyl gene (LHY) are expressed during the night period. RVE1 functions in the circadian clock and auxin pathways [39] and LHY maintains the circadian rhythm in Arabidopsis [40]. Both the RVE1 and LHY are found expressed in night-specific Peepal tree transcripts indicating the active circadian rhythms and pathways during the dark time.

In the current study, Catalase gene expression was found high in the night period transcript abundance of Peepal tree leaf tissue. A previous study on the Peepal tree showed that leaf tissue collected at night time exhibited the scotoactive opening of stomata during the night, which indicates that through the stomatal opening molecular oxygen (O_2) is released by the action of catalase enzyme on hydrogen peroxide (H_2O_2) [41]. Peepal tree plants grown in adverse habitats showed 55% higher H_2O_2 production with about a 30% increase in the peroxidase activity [41]. The physiological interaction between catalase and its substrate H_2O_2 in the plant was determined by quantifying H_2O_2 and assaying the catalase, in which catalase showed a 4-

fold increase in activity, especially during the night. Peepal tree has a higher amount of H_2O_2

deposition during the night than day, indicating pathway switching between carbon fixation

pathways. The transcriptomic sequencing from time-structure diurnal (2 PM) and nocturnal (2

AM) leaf samples showed the gene expression patterns of the carbon fixation pathway. The

gene expression of mRNA in the C3, C4, and CAM cycles indicated that depending on the

carbohydrate, amino acids biosynthesis, metabolism, and environmental conditions the plant

switches between these three cycles in a time-structured manner.

The enzymatic steps involved in the CAM cycle are similar to the C4 plant's [42]. Plants adapt

to the CAM cycle to grow during water constraints and increase the level of carbon dioxide

uptake than their C3 and C4 cycles [43]. The photorespiration in C3 plants dissipates more than

25% of the carbon fixed through the photosynthesis [44]. The day mRNA data showed that the

Peepal tree may carry out the diurnal carbon fixation by the C3 cycle. GGAT plays an important

role in the biosynthesis and metabolism of major amino acids. GGAT is involved in the

photorespiratory process. It catalyzes the reaction of glutamate and glyoxylate in the 2-

oxoglutarate and glycine [45]. High expression of GGAT in the C_4 cycle indicates that there

could be photorespiration in the Peepal tree during the night.

The Peepal tree study provides information on plants using the CAM pathway to fix nocturnal

carbon dioxide using the PEP carboxylase (PEPC) enzyme and the accumulation of malate by

the enzyme malate dehydrogenase in the large vacuoles of their cells. The essential CAM genes

in C3 species expressed in periods suggested that ancestral expression patterns required for

CAM may have pre-dated its origin in *Yucca* [46]. In the *Kalanchoë fedtschenkoi* genome

study, the convergence in protein sequence and re-scheduling of diel transcript expression of

genes was reported to be involved in nocturnal CO_2 fixation, stomatal movement, heat

tolerance, circadian clock, and carbohydrate metabolism with the other CAM species in comparison with non-CAM species [14].

Some of the previous studies in the pineapple genome revealed the gene lineage transition from C3 photosynthesis to CAM, and CAM-related genes exhibit a diel expression pattern in photosynthetic tissues [47]. The evolution of CAM in *Agave* from C3 photosynthesis shows that the core metabolic components required for CAM have ancient genomic origins which could be traceable to non-vascular plants while regulatory proteins required for diel re-programming of metabolism have shared among the recent origin of C3, C4, and CAM species [48].

In summary, the genome data and transcript abundance evidence indicate the molecular switch in the carbon fixation pathway of the Peepal tree (*F. religiosa*) during the day and night periods depending on its physiological and environmental conditions. Our study is a foundation for further experiments to determine the underlying mechanisms in C3, C4, and CAM metabolism.

Conclusions

In this study, we generated the genomic and transcriptomic data for Peepal/Bodhi tree. Genomic data pathway analyses identified the genes associated with several physiological, biochemical, metabolic, and disease pathways. Differential expression data from diurnal and nocturnal leaf tissue samples of Peepal revealed gene expression patterns in the carbon fixation pathway during light and dark. The transcript abundance indicates that plants could switch between the three C3, C4, and CAM pathways. The well-annotated genome for the Peepal tree will have broader implications for studies regarding the physiology, evolution, conservation of species, and human neurological diseases.

Methods

Collection of leaf samples and extraction of nucleic acids

The mature leaves were collected from a cultivated Peepal tree (15 years old) at a Private property, Anuganalu village, Hassan District, India (13.0647° N, 76.0363° E). We have followed a non-invasive method for collecting leaf samples. Genomic DNA was extracted from the leaves using the Qiagen DNeasy Plant Mini kit (Catalog #69106), and the quality and quantity of DNA were confirmed using the Nanodrop. From the same Peepal tree, during the day (2 PM) and night (2 AM) periods, the leaf samples were collected and immediately placed on dry ice. Total RNA was isolated from the leaf samples using the QiagenRNeasy Plant Mini kit (Catalog #74904) method and was treated with RNase-free DNase I (Catalog #M0303S) from New England BioLabs for 30 min at 37 °C to remove residual DNA. RNA integrity and quantity were confirmed on Qubit and Tape station using a dsDNA HS (Catalog #32854) kit from Invitrogen and RNA screen tape from Agilent respectively.

DNA and RNA library preparation and sequencing

Whole-genome shotgun DNA library preparation was performed using the Illumina TrueSeq DNA sample preparation kit (FC-121-2001). The paired-end (PE) (2 x 100 nts) sequencing was carried out using Illumina HiSeq 1000 at the Next Generation Genomics Facility at the Centre for Cellular and Molecular Platforms (C-CAMP), Bengaluru. Also, to increase the size of genome data, we sequenced the genome with paired-end (PE) (2 x 100 nts) using the MGISEQ-2000 platform at Bengaluru Genomics Center (BGC) Pvt. Ltd, Bengaluru.

The RNA libraries were prepared using "TruSeq RNA Library Prep Kit v2 from Illumina®" with Illumina standardized protocol. The RNA libraries were quantified on Qubit (dsDNA HS kit) and validated on the TapeStation instrument (D1000 screen tape). These RNA libraries

were used for sequencing with the Illumina platform at the Bengaluru Genomics Centre (BGC)

Pvt. Ltd, Bengaluru.

Genome size estimation and assembly

Each of the Illumina and MGISEQ-2000 raw reads were processed for a quality check using

the FastQC v0.11.6 tool [8]. Then filtering and trimming of raw reads were done to remove the

low complexity bases using the TrimGalore-0.4.5

(https://www.bioinformatics.babraham.ac.uk/projects/trimgalore/) and reads having quality

value Q>20 and length above 20 bases were taken for constructing the assembly. To estimate

the genome size, filtered reads were taken for the k-mer distribution (different k-mers from

21to 77) and abundance analysis using Jellyfish v1.1.12 [49] and GenomeScope v2.0 [50]. The

separate Illumina and MGI Seq generated raw reads were used to construct the assembly using

the tools SPAdes-3.13.0 [51] and MaSuRCA-3.2.9 [52] respectively. The parameters were the

default k-mer sizes of 21, 33, and 55 for Illumina assembly. The constructed assemblies were

used to build the super scaffolds using the tool SSPACE standard v3.0 [53].

The combined Illumina HiSeq and MGISEQ raw reads were used to construct the hybrid

assembly using the assembler SPAdes-v3.13.0 [51]. The parameters were the default k-mer

sizes 21, 33, and 55, with a 77 mer also set. The gaps in the assembly were closed by GMcloser-

1.6.2. [54]. The assembly statistics were obtained using the tool Quast v4.6.1 [55]. The

completeness and evaluation of the assembly were done by BUSCOv3 tool [56] with the

Embryophyte and Eukaryota database and by aligning the RNA-seq reads to the genome.

Structural gene prediction and functional annotation:

Peepal tree assembled scaffolds were processed for structural and functional gene annotation

using the MAKER-P v.2.31.10 software [57]. The RNA-sequenced data of *Morus notabilis* [9]

consists of expression sequence tags (ESTs) and the GFF (Gene finding format) file which

contains the gene features and structures of genes, protein data of *A. thaliana* and RNA-

sequence data of Peepal tree were imported as evidence for annotation support. The structural

and functional annotation of predicted genes and proteins was performed using BLASTP in the

Uniprot database. The protein family, structures, and gene ontology (GO) terms were identified

for protein-coding genes using InterProScan-V5.27-66.0 [22].

Gene family construction, identification of homologous and orthologous genes

Protein sequences of *A. thaliana*, *M. notabilis*, *P. persica*, *C. sativa*, *Z. jujuba,* and the protein

sequences of the current study *F. religiosa* were taken for the homologous and orthologous

gene identification. The homologous genes were identified in *F. religiosa* proteome sequences

using the BLASTP analysis against the other 5 proteomes of *A. thaliana*, *M. notabilis*, *P.*

persica, *C. sativa*, and *Z. jujuba*. OrthoVenn2 [58] was used for the clustering of orthologous

genes and the identification of the single-copy orthologous genes in all six proteomes. Further,

these single-copy orthologous genes were used for constructing the phylogenetic tree using the

tool MAFFT-v7 [59].

Comparative genome analysis

We downloaded the genomes of *F. carica*, *F. microcarpa*, and *M. notabilis*. We aligned these

genomes against the Peepal tree genome assembly to understand their relationships using the

BWA-V0.7.17 (Burrows-Wheeler Aligner) [60] and Samtools v1.7 [61].

Prediction of repetitive elements: TEs and SSR

The RepeatModeller-open-1.0.11and RepeatMasker-4.0 tools were used for repeat library

building and repeat identification in the assembly respectively. The MicroSAtellite

identification tool (MISA) [62] was used for the identification of SSRs from assembled genome

sequences of *F. religiosa*. The parameters were set to identify perfect di-, tri-, tetra-, penta-,

and hexa nucleotide motifs with a minimum threshold of 6, 5, 5, 5, and 5 repeats, respectively.

Prediction of transcription factor families

The families of transcription factors (TFs) were predicted in genome annotations and

differentially expressed transcripts of the Peepal tree using Plant Transcription Factor Database

v5.0 [23].

Non-coding RNA genes

The transfer RNAs in the Peepal tree genome were found using tRNAscan-SE (v2.0.3) [63]

with the 'eukaryotes' option. tRNAscan-SE software deployed with the covariance models

identifies the primary sequence and secondary structure information of tRNA and gives the

complete tRNA genes for the query genome and transcriptome sequences. tRNAscan-SE

software is integrated with Infernal v1.1 to enhance the tRNA search with better covariance

and other updated models. Using the isotype-specific covariance model provides the functional

classification of tRNAs and in the first pass search cutoff score, 10 is set. The miRbase database

(http://www.mirbase.org) was used for the identification of putative miRNAs in the genome

and unique identified transcripts sequence data based on the homology search. The long non-

coding RNAs (lncRNAs) were identified with the Coding Potential Calculator tools [64].

Transcriptome sequencing, assembly, and annotation

High-quality stranded RNA sequencing (ssRNA-seq) reads were assembled into putative

transcripts using Trinity v2.9.0 (Grabherr et al., 2011). Assembled transcripts were passed

through Transdecoder v5.02 to predict the coding sequences. The transcripts were clustered to

find the unigenes by removing the redundant transcripts using the tool CD-HIT-est v.0.0.1 [65]

with a 95% sequence identity threshold. Transcripts assembled from Trinity and CD-HIT-

v0.0.1 were used in downstream analyses for gene prediction. Unigenes were used to predict

the putative genes using the NCBI non-redundant (nr) database using the BLASTX program

and proteins were predicted from the Uniprot database using the BLASTP program. The Trinity

assembled transcripts were annotated using Trinotate- V3.11. The raw reads were mapped to

scaffold assembled genome using Cufflinks-v2.2.1 [66] and considered as reference assembly.

Transcript quantification and differential gene expression analysis

The estimation of transcripts abundance was determined using RNA-Seq by Expectation-

Maximization (RSEM) tool [67], which quantifies transcript level abundance from RNA-seq

data. RSEM first generates and pre-processes a set of reference transcript sequences and then

aligns reads to reference transcripts followed by an estimation of transcript abundances.

Normalized transcripts obtained from the transcript quantification methods were used in the

next step for analyzing the differential gene expression of genes. FPKM and Trimmed Mean

of M-values (TMM) are calculated to understand the expression levels of genes in day and

night samples of the Peepal tree. For further analysis, the gene expression was estimated using

FPKM and TMM value minimum ≥1. The TMM value was used to cluster the genes according

to their expression pattern using the edgeR package in the R tool. The parameters used in the

differential analysis were a probability value P-value of 0.001 and a fold change value of log2.

The expression value was also determined for assembled transcripts to verify the expression of

genes predicted from gene models. The differentially expressed genes were annotated using

BLAST2GO Annotation software [68].

Pathway Analysis

The annotated genes from the assembled genome and the differentially expressed genes from

the Peepal tree leaf tissues collected during the day (2.00 PM) and night (2.00 AM) were used

for pathway analysis in the KAAS (KEGG Automatic Annotation Server) (KEGG) server [17]

using the BBH (bi-directional best hit) method and the search against a default set of 40

eukaryotic organisms. It provided the list of pathways where the candidate genes were mapped

based on the orthologous homology alignment.

Abbreviations

BUSCO: Benchmarking Universal Single-Copy Orthologous; ESts: Expressed Sequence Tags;

PPR repeat: Pentatricopeptide repeat; FPKM: Fragments Per Kilobase of transcript per Million

mapped reads; MSA: Multiple Sequence Alignment

Statement:

• **Experimental Plant details:** *Ficus religiosa* a cultivated plant was used in this study.

The mature leaves were collected from this plant at Private property, Anuganalu

village, Hassan District, India (13.0647° N, 76.0363° E) were kindly provided by

Prof. Malali Gowda, DNA Life Foundation, Anuganalu village, Hassan District,

India (13.0647° N, 76.0363° E).

• The formal identification of the plant was undertaken by Prof. Malali Gowda, who

owns the place of plant location and is also a corresponding author of this study.

Hence, we had his permission to collect the plant material. The specimen of this

material has not been deposited in any publicly available herbarium.

Declaration

Ethics approval and consent to participate

The study is compiled with relevant institutional, national, and international guidelines and legislation.

Consent for publication

Not Applicable (NA)

Availability of data and materials

The raw sequence reads have been deposited under NCBI Sequence Read Archive (SRA) accession numbers SRR7244210 for Illumina sequenced *F. religiosa* WGS, SRR13827064 for MGISEQ sequenced *F. religiosa* WGS – (https://trace.ncbi.nlm.nih.gov/Traces/study1/?acc=PRJNA474013), SRR7343291 (https://trace.ncbi.nlm.nih.gov/Traces/study1/?acc=SRR7343291) for *F. religiosa* transcriptome). The Whole Genome Shotgun project has been deposited at DDBJ/ENA/GenBank under the accession JAFMPE000000000 (https://www.ncbi.nlm.nih.gov/nuccore/JAFMPE000000000.1/) and Transcriptome Shotgun Assembly project has been deposited at DDBJ/ENA/GenBank under the accession GJAV00000000 (https://www.ncbi.nlm.nih.gov/nuccore/GJAV00000000.1/).

Other published data used in the study: The protein sequences of *Arabidopsis thaliana* (https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/000/001/735/GCF_000001735.4_TAIR10.1/GCF_000001735.4_TAIR10.1_protein.faa.gz), *Morus notabilis* (https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/000/414/095/GCF_000414095.1_ASM41409v2/GCF_000414095.1_ASM41409v2_protein.faa.gz), *Prunus persica* (https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/000/346/465/GCF_000346465.2_Prunus_pers

ica_NCBIv2/GCF_000346465.2_Prunus_persica_NCBIv2_protein.faa.gz), *Cannabis sativa*

(https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/900/626/175/GCF_900626175.2_cs10/GCF_9

[00626175.2_cs10_protein.faa.gz](https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/900/626/175/GCF_900626175.2_cs10/GCF_900626175.2_cs10_protein.faa.gz)),   *Ziziphus*   *jujuba*

(https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/020/796/205/GCF_020796205.1_ASM207962

0v1/GCF_020796205.1_ASM2079620v1_protein.faa.gz).

Competing interests

The authors declare that they have no competing interests.

Funding

Not Applicable (NA)

Author's contribution

AKL performed the DNA and RNA isolation from leaf tissues, genome assembly and

functional annotation, gene prediction, repeat prediction, orthologous gene clustering, Genome

and Transcriptome pathway analysis, submitted WGS and RNA-seq data to NCBI, prepared

the genomic and transcriptomic study tables and figures, wrote the manuscript; MG, AKL, and

AKP designed the genomic and transcriptomic experiments of *Ficus religiosa*; MG conceived

and conceptualized the project; reviewed and edited the manuscript.

Acknowledgment

We would like to thank Tata Education and Development Trust fellowship (01/2018-2019)

provided for Ashalatha K L for her Ph.D. program. We acknowledge the Next Generation

Genomics Facility at the Centre for Cellular and Molecular Platforms (C-CAMP) and

Bengaluru Genomics Center Pvt. Ltd for supporting sequencing and analysis. We are grateful

to the MGI team, in Hong Kong for sequencing the whole genome.

Author's Information:

[1]K L Ashalatha: ashalatha708@gmail.com, ashalatha.kl@tdu.edu.in

[2]K P Arunkumar: arunkumar.kallare@gmail.com, arunkallare.csb@gov.in

[*1,3]Malali Gowda: malali.gene@gmail.com

[1] The University of Trans-Disciplinary Health Sciences and Technology, Yelahanka,

Bengaluru, 560064, India

[2] Central Muga Eri Research and Training Institute (CMER&TI), Central Silk Board, Ministry

of Textiles Lahdoigarh, Jorhat, Assam, 785700, India

[3] DNA Life Organization, Yelahanka, Bengaluru, 560064, India

References

1. IPCN Chromosome Reports, 2014. Index to Plant Chromosome Numbers (IPCN),
Tropicos website. St. Louis, Missouri, USA: Missouri Botanical Garden.
2. Van Noort S, Gardiner AJ, Tolley KA. New records of Ficus (Moraceae) species
emphasize the conservation significance of inselbergs in Mozambique. South African J Bot.
2007;73:642–9.
3. Ashalatha KL, Gowda M. Heritage of Neem–Peepal Tree Resides a Profound Scientific
Facts. 2019. p. 13–9.
4. Chantarasuwan B, Baas P, van Heuven B-J, Baider C, van Welzen PC. Leaf anatomy of
Ficus subsection U rostigma (Moraceae). Bot J Linn Soc. 2014;175:259–81.
5. Singh D, Singh B, Goel RK. Traditional uses, phytochemistry and pharmacology of Ficus
religiosa: A review. J Ethnopharmacol. 2011;134:565–83.
6. Bhangale JO, Acharya SR. Anti-Parkinson Activity of Petroleum Ether Extract of Ficus
religiosa (L.) Leaves. Adv Pharmacol Sci. 2016;2016:1–9.
7. Bhangale JO, Acharya NS, Acharya SR. Protective effect of Ficus religiosa (L.) against 3-
nitropropionic acid induced Huntington disease. Orient Pharm Exp Med. 2016;16:165–74.
8. Gregory M, Divya B, Mary RA, Viji MMH, Kalaichelvan VK, Palanivel V. Anti–ulcer
activity of Ficus religiosa leaf ethanolic extract. Asian Pac J Trop Biomed. 2013;3:554–6.

9. He N, Zhang C, Qi X, Zhao S, Tao Y, Yang G, et al. Draft genome sequence of the
mulberry tree Morus notabilis. Nat Commun. 2013;4:2445.
10. Usai G, Mascagni F, Giordani T, Vangelisti A, Bosi E, Zuccolo A, et al. Epigenetic
patterns within the haplotype phased fig (Ficus carica L.) genome. Plant J. 2020;102:600–14.
11. Zhang X, Wang G, Zhang S, Chen S, Wang Y, Wen P, et al. Genomes of the Banyan
Tree and Pollinator Wasp Provide Insights into Fig-Wasp Coevolution. Cell. 2020;183:875-
889.e17.
12. Abhisek Chakraborty, Shruti Mahajan, Manohar Singh Bisht VKS. Genome sequencing
and comparative analysis of Ficus benghalensis and Ficus religiosa trees reveal evolutionary
mechanisms of longevity. https://doi.org/doi: https://doi.org/10.1101/2021.10.14.464369.
13. Ming R, VanBuren R, Wai CM, Tang H, Schatz MC, Bowers JE, et al. The pineapple
genome and the evolution of CAM photosynthesis. Nat Genet. 2015;47:1435–42.
14. Yang X, Hu R, Yin H, Jenkins J, Shu S, Tang H, et al. The Kalanchoë genome provides
insights into convergent evolution and building blocks of crassulacean acid metabolism. Nat
Commun. 2017;8:1899.
15. Guan Q, Tan B, Kelley TM, Tian J, Chen S. Physiological Changes in
Mesembryanthemum crystallinum During the C3 to CAM Transition Induced by Salt Stress.
Front Plant Sci. 2020;11.
16. Schlüter U, Denton AK, Bräutigam A. Understanding metabolite transport and
metabolism in C4 plants through RNA-seq. Curr Opin Plant Biol. 2016;31:83–90.
17. Moriya Y, Itoh M, Okuda S, Yoshizawa AC, Kanehisa M. KAAS: an automatic genome
annotation and pathway reconstruction server. Nucleic Acids Res. 2007;35 Web
Server:W182–5.
18. Hays JB. Arabidopsis thaliana, a versatile model system for study of eukaryotic genome-
maintenance functions. DNA Repair (Amst). 2002;1:579–600.
19. Spampinato CP, Gomez-Casati DF. Research on Plants for the Understanding of Diseases
of Nuclear and Mitochondrial Origin. J Biomed Biotechnol. 2012;2012:1–12.
20. Jones AM, Chory J, Dangl JL, Estelle M, Jacobsen SE, Meyerowitz EM, et al. The
Impact of Arabidopsis on Human Health: Diversifying Our Portfolio. Cell. 2008;133:939–43.
21. Jung HW, Son HY, Minh C Van, Kim YH, Park Y-K. Methanol extract of Ficus leaf
inhibits the production of nitric oxide and proinflammatory cytokines in LPS-stimulated
microglia via the MAPK pathway. Phyther Res. 2008;22:1064–9.
22. Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, et al. InterProScan 5:
genome-scale protein function classification. Bioinformatics. 2014;30:1236–40.
23. Jin J, Tian F, Yang D-C, Meng Y-Q, Kong L, Luo J, et al. PlantTFDB 4.0: toward a
central hub for transcription factors and regulatory interactions in plants. Nucleic Acids Res.
2017;45:D1040–5.
24. Pan J, Huang D, Guo Z, Kuang Z, Zhang H, Xie X, et al. Overexpression of
microRNA408 enhances photosynthesis, growth, and seed yield in diverse plants. J Integr
Plant Biol. 2018;60:323–40.
25. Schramm M, Klybas V, Racker E. Phosphorolytic cleavage of fructose-6-phosphate by
fructose-6-phosphate phosphoketolase from Acetobacter xylinum. J Biol Chem.
1958;233:1283–8.
26. Ebata M, Sato R, Bak T. The enzymatic phosphorylation of sedoheptulose*. J Biochem.
1955;42:715–25.
27. Kanehisa M. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic Acids Res.
2000;28:27–30.
28. Kanehisa M. Toward understanding the origin and evolution of cellular organisms.
Protein Sci. 2019;28:1947–51.
29. Kanehisa M, Furumichi M, Sato Y, Kawashima M, Ishiguro-Watanabe M. KEGG for

taxonomy-based analysis of pathways and genomes. Nucleic Acids Res. 2022.
https://doi.org/10.1093/nar/gkac963.
30. McFarlane CR, Shah NR, Kabasakal B V., Echeverria B, Cotton CAR, Bubeck D, et al.
Structural basis of light-induced redox regulation in the Calvin–Benson cycle in
cyanobacteria. Proc Natl Acad Sci. 2019;116:20984–90.
31. Cousins AB, Baroli I, Badger MR, Ivakov A, Lea PJ, Leegood RC, et al. The Role of
Phospho enol pyruvate Carboxylase during C4 Photosynthetic Isotope Exchange and
Stomatal Conductance. Plant Physiol. 2007;145:1006–17.
32. Mahesh HB, Subba P, Advani J, Shirke MD, Loganathan RM, Chandana SL, et al. Multi-
Omics Driven Assembly and Annotation of the Sandalwood (Santalum album) Genome.
Plant Physiol. 2018;176:2772–88.
33. Jones-Rhoades MW, Bartel DP, Bartel B. MicroRNAs and their regulatory roles in plants.
Annu Rev Plant Biol. 2006;57:19–53.
34. Sunkar R, Zhu J-K. Novel and Stress-Regulated MicroRNAs and Other Small RNAs
from Arabidopsis[W]. Plant Cell. 2004;16:2001–19.
35. Abdel-Ghany SE, Pilon M. MicroRNA-mediated Systemic Down-regulation of Copper
Protein Expression in Response to Low Copper Availability in Arabidopsis. J Biol Chem.
2008;283:15932–45.
36. Liu C, Xu X, Kan J, Cheng Z ming, Chang Y, Lin J, et al. Genome-wide analysis of the
C3H zinc finger family reveals its functions in salt stress responses of Pyrus betulaefolia.
PeerJ. 2020;8:e9328.
37. Lee D-K, Kim H Il, Jang G, Chung PJ, Jeong JS, Kim YS, et al. The NF-YA transcription
factor OsNF-YA7 confers drought stress tolerance of rice in an abscisic acid independent
manner. Plant Sci. 2015;241:199–210.
38. Bolt S, Zuther E, Zintl S, Hincha DK, Schmülling T. ERF105 is a transcription factor
gene of Arabidopsis thaliana required for freezing tolerance and cold acclimation. Plant Cell
Environ. 2017;40:108–20.
39. Rawat R, Schwartz J, Jones MA, Sairanen I, Cheng Y, Andersson CR, et al.
REVEILLE1, a Myb-like transcription factor, integrates the circadian clock and auxin
pathways. Proc Natl Acad Sci. 2009;106:16883–8.
40. Mizoguchi T, Wheatley K, Hanzawa Y, Wright L, Mizoguchi M, Song H-R, et al. LHY
and CCA1 Are Partially Redundant Genes Required to Maintain Circadian Rhythms in
Arabidopsis. Dev Cell. 2002;2:629–41.
41. Smitha RB, Bennans T, Mohankumar C, Benjamin S. Oxidative stress enzymes in Ficus
religiosa L.: Biochemical, histochemical and anatomical evidences. J Photochem Photobiol B
Biol. 2009;95:17–25.
42. Christin P-A, Arakaki M, Osborne CP, Bräutigam A, Sage RF, Hibberd JM, et al. Shared
origins of a key enzyme during the evolution of C4 and CAM metabolism. J Exp Bot.
2014;65:3609–21.
43. NOBEL PS. Achievable productivities of certain CAM plants: basis for high values
compared with C 3 and C 4 plants. New Phytol. 1991;119:183–205.
44. Sharkey TD. Estimating the rate of photorespiration in leaves. Physiol Plant.
1988;73:147–52.
45. Igarashi D, Miwa T, Seki M, Kobayashi M, Kato T, Tabata S, et al. Identification of
photorespiratory glutamate:glyoxylate aminotransferase (GGAT) gene in Arabidopsis. Plant
J. 2003;33:975–87.
46. Heyduk K, Ray JN, Ayyampalayam S, Moledina N, Borland A, Harding SA, et al. Shared
expression of crassulacean acid metabolism (CAM) genes pre-dates the origin of CAM in the
genus Yucca. J Exp Bot. 2019;70:6597–609.
47. Ming R, VanBuren R, Wai CM, Tang H, Schatz MC, Bowers JE, et al. The pineapple

genome and the evolution of CAM photosynthesis. Nat Genet. 2015;47:1435–42.

48. Yin H, Guo H-B, Weston DJ, Borland AM, Ranjan P, Abraham PE, et al. Diel rewiring
and positive selection of ancient plant proteins enabled evolution of CAM photosynthesis in
Agave. BMC Genomics. 2018;19:588.

49. Marçais G, Kingsford C. A fast, lock-free approach for efficient parallel counting of
occurrences of k-mers. Bioinformatics. 2011;27:764–70.

50. Vurture GW, Sedlazeck FJ, Nattestad M, Underwood CJ, Fang H, Gurtowski J, et al.
GenomeScope: fast reference-free genome profiling from short reads. Bioinformatics.
2017;33:2202–4.

51. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes:
A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. J
Comput Biol. 2012;19:455–77.

52. Zimin A V., Marçais G, Puiu D, Roberts M, Salzberg SL, Yorke JA. The MaSuRCA
genome assembler. Bioinformatics. 2013;29:2669–77.

53. Boetzer M, Henkel C V., Jansen HJ, Butler D, Pirovano W. Scaffolding pre-assembled
contigs using SSPACE. Bioinformatics. 2011;27:578–9.

54. Kosugi S, Hirakawa H, Tabata S. GMcloser: closing gaps in assemblies accurately with a
likelihood-based selection of contig or long-read alignments. Bioinformatics. 2015;:btv465.

55. Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for
genome assemblies. Bioinformatics. 2013;29:1072–5.

56. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics. 2015;31:3210–2.

57. Campbell MS, Law M, Holt C, Stein JC, Moghe GD, Hufnagel DE, et al. MAKER-P: A
Tool Kit for the Rapid Creation, Management, and Quality Control of Plant Genome
Annotations. Plant Physiol. 2014;164:513–24.

58. Xu L, Dong Z, Fang L, Luo Y, Wei Z, Guo H, et al. OrthoVenn2: a web server for whole-
genome comparison and annotation of orthologous clusters across multiple species. Nucleic
Acids Res. 2019;47:W52–8.

59. Katoh K, Rozewicki J, Yamada KD. MAFFT online service: multiple sequence
alignment, interactive sequence choice and visualization. Brief Bioinform. 2019;20:1160–6.

60. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 2009;25:1754–60.

61. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years
of SAMtools and BCFtools. Gigascience. 2021;10.

62. Thiel T, Michalek W, Varshney R, Graner A. Exploiting EST databases for the
development and characterization of gene-derived SSR-markers in barley (Hordeum vulgare
L.). Theor Appl Genet. 2003;106:411–22.

63. Lowe TM, Eddy SR. tRNAscan-SE: A Program for Improved Detection of Transfer RNA
Genes in Genomic Sequence. Nucleic Acids Res. 1997;25:955–64.

64. Kong L, Zhang Y, Ye Z-Q, Liu X-Q, Zhao S-Q, Wei L, et al. CPC: assess the protein-
coding potential of transcripts using sequence features and support vector machine. Nucleic
Acids Res. 2007;35 suppl_2:W345–9.

65. Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of protein
or nucleotide sequences. Bioinformatics. 2006;22:1658–9.

66. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, et al.
Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and
isoform switching during cell differentiation. Nat Biotechnol. 2010;28:511–5.

67. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. BMC Bioinformatics. 2011;12:323.

68. Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M, Robles M. Blast2GO: a
universal tool for annotation, visualization and analysis in functional genomics research.
Bioinformatics. 2005;21:3674–6.

Tables

Table 1: Final assembly and annotation of Peepal genome

Features	Final assembly
Total length of assembled sequence (Mb)	406,103,086
Number of scaffolds/contigs	202,258
Minimum scaffold length (bp)	200
Maximum scaffold length (bp)	148483
GC %	34.23
N50 (bp)	5817
L50	17,605
Number of annotated genes	35093

813 **Table 2:** Statistics of Uni-genes in Peepal Transcripts

Features	Sample (Day - 2 PM)	Sample (Night - 2 AM)	Combined Sample
Number of genes	22597	23360	26479
Number of transcripts	16912	16780	18173
Percent GC	46.37	42.42	46.25
Contig N50 (nts)	1404	2053	1374
Median contig length	813	1011	753
Average Contig (nts)	1060.06	1332.73	1017.44
Total assembled bases	23954145	120608982	27055237

Figure legends

Figure 1: Orthologous clustering of 6 species using proteome data deduced 24310 orthologous

gene clusters and 1184 single-copy gene clusters across the above 6 species.

Figure 2: The candidate genes involved in the C3, CAM, and C4 cycles.

A) Calvin- Benson (C3) cycle B) Crassulacean acid metabolism (CAM) cycle C) C4 cycle. The

X-axis represents the genes involved in pathways, Y-axis is the matrix of normalized

expression trimmed mean of M (TMM) values; the Blue graph - leaf tissue collected during

the day (2 PM), Red graph - leaf tissue collected during the night period (2 AM).

Figure 3: Gene expression pattern of *F. religiosa* carbon fixation genes across the diurnal (2

PM) and nocturnal (2 AM) expression data. A) C3 cycle, B) CAM cycle C) C4 cycle log2-

transformed Fragments Per Kilobase of transcript per Million mapped reads (FPKM) value-

based expression profiles are shown.

838 **Figure 1:** Orthologous clustering of 6 species using proteome data deduced 24310 orthologous

839 gene clusters and 1184 single-copy gene clusters across the above 6 species

Figure 2: The candidate genes involved in the C3, CAM and C4 cycle.

A) Calvin- Benson (C3) cycle B) Crassulacean acid metabolism (CAM) cycle C) C4 cycle. X – axis represents the genes involved pathways, Y – axis is the matrix of normalised expression trimmmed mean of M (TMM) values; Blue graph – leaf tissue collected during day period (2 PM), Red graph – leaf tissue collected during night period (2 AM)

Figure 3: Gene expression pattern of *F. religiosa* carbon fixation genes across the diurnal (2 PM) and nocturnal (2 AM) expression data. A) C3 cycle, B) CAM cycle C) C4 cycle with fold change log2-transformed Fragments Per Kilobase of transcript per Million mapped reads (FPKM) value based expression profiles are shown