Gender and comorbidity moderate the relationship between mood disorder symptoms and effortful avoidance performance

Brandon J. Forys[1], Ryan J. Tomm[1], Dayana Stamboliyska[1], Alex R. Terpstra[1], Luke Clark[1,2], Trisha Chakrabarty[2,3], Stan B. Floresco[1,2], and & Rebecca M. Todd[1,2]

[1] Department of Psychology

University of British Columbia

Vancouver

BC

Canada

[2] Djavad Mowafaghian Centre for Brain Health

University of British Columbia

Vancouver

BC

Canada

[3] Department of Psychiatry

University of British Columbia

Vancouver

BC

Canada

Author Note

The authors declare no competing financial interests. We would like to acknowledge Alan Kingstone for his instrumental feedback on this project, as well as the contributions of Aanandi Sidarth to designing visual stimuli and Ian Daly to designing sounds stimuli for these studies. This research was supported by a Natural Sciences and Engineering Research Council of Canada (NSERC) grant (#F19-05182) and the UBC Djavad Mowafaghian Centre for Brain Health Innovation Fund Kickstart Research Grant (#F19-05932), as well as an NSERC Canada Graduate Scholarship - Master's (CGS M) award to B.J. Forys and a Michael Smith Foundation for Health Research to R.M. Todd.

The authors made the following contributions. Brandon J. Forys: Conceptualization, Investigation, Methodology, Writing - Original Draft Preparation, Writing - Review & Editing; Ryan J. Tomm: Conceptualization, Methodology, Writing - Review & Editing; Dayana Stamboliyska: Conceptualization, Investigation, Methodology, Writing - Review & Editing; Alex R. Terpstra: Conceptualization, Writing - Review & Editing; Luke Clark: Conceptualization, Funding acquisition, Writing - Review & Editing; Trisha Chakrabarty: Conceptualization, Funding acquisition, Writing - Review & Editing; Stan B. Floresco: Conceptualization, Funding acquisition, Writing - Review & Editing; Rebecca M. Todd: Conceptualization, Funding acquisition, Methodology, Project Administration, Resources, Supervision, Writing - Review & Editing.

Correspondence concerning this article should be addressed to Brandon J. Forys, 2136 West Mall, Vancouver, BC Canada V6T 1Z4. E-mail: brandon.forys@psych.ubc.ca

Abstract

We must often decide how much effort to exert or withhold to avoid undesirable outcomes or obtain rewards. In depression and anxiety, levels of avoidance tend to be excessive and reward-seeking is reduced. Yet outstanding questions remain about the links between motivated action/inhibition and anxiety and depression symptoms, and whether they differ between men and women. Here we examined the relationship between anxiety and depression symptoms and performance on effortful active and inhibitory avoidance (Study 1) and reward seeking (Study 2) in humans. Undergraduates and paid online workers ($N_{Avoid} = 545$, $N_{Reward} = 310$; $N_{Female} = 368$, $N_{Male} = 450$, $M_{Age} = 22.58$, $Range_{Age} = 17$-62) were assessed on the Beck Depression Inventory (BDI-II) and the Beck Anxiety Inventory (BAI) and performed an instructed online avoidance or reward-seeking task. Participants had to make multiple presses on active trials and withhold presses on inhibitory trials to avoid an unpleasant sound (Study 1) or obtain points towards a monetary reward (Study 2). Overall, men deployed more effort than women in both avoidance and reward-seeking, and anxiety symptoms were negatively associated with active reward-seeking accuracy. Gender moderated the relationship between anxiety symptoms and inhibitory avoidance, such that women with higher anxiety showed reduced inhibitory avoidance accuracy. Anxiety symptoms predicted poorer active avoidance performance for those high in depression symptoms only. Our results illuminate effects of gender and symptom comorbidity in the relationship between mood disorder symptoms and the motivation to actively and effortfully respond to obtain positive and avoid negative outcomes.

Keywords: avoidance, reward, effort, depression, anxiety

Word count: 6824

Gender and comorbidity moderate the relationship between mood disorder symptoms and effortful avoidance performance

Significance statement

We often need to take effortful action, or withhold action, to avoid unpleasant outcomes or obtain rewards. Depression and anxiety can impact these behaviours' effectiveness, but the role of depression in avoidance and anxiety in reward-seeking is not well understood. Gender differences in avoidance and reward-seeking have also not been examined. Here, we present a task in which participants had to make or withhold button presses to avoid hearing an unpleasant sound or to obtain a reward. Men deployed more effort than women in avoidance, and women with higher levels of anxiety symptoms had lower inhibitory avoidance accuracy than men. Our results illuminate gender differences in how depressive and anxiety symptoms impact our ability to avoid threats and obtain rewards.

Introduction

Avoidance and reward-seeking behaviours

Living organisms are motivated to avoid potential threats or to acquire rewards respectively. Often achieving these goals requires action, but it can also require refraining from action. For example, we may take action to remove a threat's potential harm through active avoidance, or we may decide that withholding action is the best way to let the threat pass by, as in inhibitory avoidance (Krypotos, Effting, Kindt, and Beckers, 2015; LeDoux, Moscarello, Sears, and Campese, 2017). Alternatively, in a situation that offers the possibility of reward, we may take action to approach the reward through active reward-seeking or, instead, inhibit pre-potent reward seeking to wait for a larger reward (Capuzzo and Floresco, 2020; Cools, 2008). Research suggests that the expression of similar behavioral actions (including inhibition) is dependent on the motivational context (aversive

vs. appetitive), which influences the likelihood of selecting a specific action in a specific

motivational context (Wang and Delgado, 2021). However, in neuropsychiatric research,

depressive disorders are often studied with regard to reward-seeking contexts, and anxiety

disorders with regard to avoidance contexts, with little emphasis on the other motivational

context. Symptoms of anxiety and depression have been associated with avoidance,

typically operationalized via active avoidance and via questionnaires, as threats are

overestimated (Bishop and Gagne, 2018; Browning, Behrens, Jocham, O'Reilly, and Bishop,

2015; Cléry-Melin, Schmidt, Lafargue, Baup, Fossati, and Pessiglione, 2011; Mkrtchian,

Aylward, Dayan, Roiser, and Robinson, 2017; Ottenbreit, Dobson, and Quigley, 2014). In

depression, reward-seeking may also be impaired due to a lack of motivation to obtain

rewards (Bishop and Gagne, 2018; Slaney, Perkins, Davis, Penton-Voak, Munafò,

Houghton, and Robinson, 2021). Past research has established the importance of avoidance

and reward-seeking behaviours in helping us navigate our environment and stay safe

(Krypotos, Effting, Kindt, and Beckers, 2015; LeDoux, Moscarello, Sears, and Campese,

2017). However, active vs. inhibitory subtypes of these behaviours have not typically been

distinguished – especially through objective measures of observable behavior. Furthermore,

anxiety and depression are often comorbid (Brown, Campbell, Lehman, Grisham, and

Mancill, 2001), and this comorbidity has been linked to excessive avoidance behaviours

(Ottenbreit, Dobson, and Quigley, 2014). However, the role of comorbidity in affecting

active vs. inhibitory avoidance, or reward-seeking, has not been examined.

Gender may also be an important variable in this relationship. For example, gender

differences have been identified in the presentation and incidence of mood and anxiety

disorders, such that women have higher rates of depression and present more often with

depression than men (Altemus, Sarvaiya, and Neill Epperson, 2014; Kessler, 2006; Parker

and Brotchie, 2010) and have rates of anxiety disorders that are twice as high as those of

men (McLean, Asnaani, Litz, and Hofmann, 2011; Pittig, Treanor, LeBeau, and Craske,

2018). However, we do not know how these gender differences manifest themselves in

avoidance or reward-seeking behaviours. Although mood and anxiety disorders are often comorbid, they also manifest with distinct symptoms and courses that would require distinct strategies to treat in a clinical context (Goldstein-Piekarski, Williams, and Humphreys, 2016; McLean, Asnaani, Litz, and Hofmann, 2011). In the present study, we ask how mood disorder symptoms impact active vs. inhibitory avoidance and reward-seeking behaviours in a population of young adults with depressive and anxiety symptoms ranging from minimal to severe.

The role of mood disorder symptoms and gender differences in avoidance and reward-seeking

It has been proposed that mood and anxiety disorder symptoms shift the perceived value and costs of avoidance and reward-seeking in sub-optimal ways. The Altered Computations Underlying Decision Making (ACDM) framework (Bishop and Gagne, 2018) proposes that anxiety is linked to underestimation of the effort cost in avoiding an aversive outcome and that depression is linked to overestimation of the effort cost in obtaining a reward. These effort costs interact with the perceived value of avoidance or reward-seeking to inform one's decision on whether or not to engage in the behaviour. Past experimental work has also identified impairments in physical effort deployment for reward in populations with depression (Pessiglione, Vinckier, Bouret, Daunizeau, and Le Bouc, 2018; Treadway, Buckholtz, Schwartzman, Lambert, and Zald, 2009; see Culbreth, Moran, and Barch, 2018 for a review) and anxiety (Wang and Delgado, 2021). However, work linking mood and anxiety disorders to impairments in adaptive avoidance and reward-seeking often focuses on these avoidance and reward-seeking behaviours as unitary processes. As such, we still do not know how shifts in perceived effort costs linked to mood and anxiety disorders manifest themselves in active or inhibitory avoidance or reward-seeking.

To better understand the degree to which depressive and anxiety symptoms contribute to active and inhibitory forms of avoidance or reward-seeking a rigorous

assessment of effort deployment in these behaviors is needed. People with Major Depressive

Disorder (MDD) choose high-effort options less often on effort-based decision making tasks

involving reward. This behaviour is potentially symptomatic of a larger-scale motivational

deficit (Pessiglione, Vinckier, Bouret, Daunizeau, and Le Bouc, 2018; Treadway, Bossaller,

Shelton, and Zald, 2012; Treadway, Buckholtz, Schwartzman, Lambert, and Zald, 2009). If

maladaptive effort deployment is a primary characteristic of mood and anxiety disorders,

then we might expect active avoidance and reward-seeking to be impaired more than

inhibitory forms of these behaviours overall (Culbreth, Moran, and Barch, 2018). Anxiety,

especially in the context of depressive symptoms, has been shown to impair our sensitivity

to rewards (Auerbach, Pagliaccio, Hubbard, Frosch, Kremens, Cosby, Jones, Siless, Lo,

Henin, Hofmann, Gabrieli, Yendiki, Whitfield-Gabrieli, and Pizzagalli, 2022; Dillon, Rosso,

Pechtel, Killgore, Rauch, and Pizzagalli, 2014), although whether anxiety's impact on

reward-seeking differs for active or inhibitory behaviours is not yet clear.

Additionally, individual differences in the presentation and severity of mood and

anxiety disorders - beyond the mere presence or absence of the disorder - may manifest

with different patterns of active vs. inhibitory behaviours depending on the motivational

context. Among these differences, gender differences are especially prominent. Women

generally present with more depressive symptoms (Parker and Brotchie, 2010) and

experience depression comorbid with anxiety more often than men (Kessler, 2006; McLean,

Asnaani, Litz, and Hofmann, 2011; Ottenbreit, Dobson, and Quigley, 2014). Thus, the

impact of mood and anxiety disorders on our ability to avoid aversive outcomes and seek

out rewarding outcomes may be linked to gender differences that affect the motivational

deficits these disorders present. If gender differences in depressive and anxiety symptom

presentation - as captured on self-report scales (Beck, Steer, and Brown, 1996) - are

predicted by differences in accuracy and effort deployment on active and inhibitory

avoidance and reward-seeking tasks, this may help elucidate how observed gender

differences in depressive and anxiety symptoms map on to real-life behavioural changes.

₁₇₀   In order to bring our understanding of mood disorder symptoms into a framework

₁₇₁ that acknowledges differences in active vs. inhibitory avoidance and reward-seeking

₁₇₂ behaviours, we must consider both anxiety and depression in a framework that directly

₁₇₃ investigates their impact on these behaviours,, and how they might interact to impair

₁₇₄ effective avoidance and reward-seeking. While the relationships between anxiety and

₁₇₅ avoidance (Bishop and Gagne, 2018; Levita, Hoskin, and Champi, 2012; Norbury, Robbins,

₁₇₆ and Seymour, 2018), and depression and reward-seeking (Rizvi, Pizzagalli, Sproule, and

₁₇₇ Kennedy, 2016; Slaney, Perkins, Davis, Penton-Voak, Munafò, Houghton, and Robinson,

₁₇₈ 2021; Treadway, Bossaller, Shelton, and Zald, 2012) are well established, those between

₁₇₉ anxiety and reward-seeking, as well as depression and avoidance, have yet to be fully

₁₈₀ characterized. As depression and anxiety can be highly comorbid (Goldstein-Piekarski,

₁₈₁ Williams, and Humphreys, 2016; Ottenbreit, Dobson, and Quigley, 2014), parsing which

₁₈₂ aspects of avoidance and reward-seeking behaviors are affected uniquely by depressive

₁₈₃ symptoms, anxiety symptoms, or their common pathophysiological substrates (Insel,

₁₈₄ Cuthbert, Garvey, Heinssen, Pine, Quinn, Sanislow, and Wang, 2010), would be important

₁₈₅ in informing future depression and anxiety treatments.

₁₈₆ **An effortful avoidance and reward-seeking study**

₁₈₇   Despite established gender differences in the prevalence and presentation of mood

₁₈₈ disorder symptoms (Kessler, 2006; Parker and Brotchie, 2010; Thompson and Bland, 2018),

₁₈₉ it is not known how the relationship between mood and anxiety disorder symptoms and

₁₉₀ avoidance and reward-seeking differs by gender. Gender differences in motivational deficits

₁₉₁ may lead to unique patterns in active and inhibitory behaviours, but this has not been

₁₉₂ examined either. As such, in this work, we ask 1) whether anxiety and depression

₁₉₃ symptoms predict accuracy and effort deployment on active/inhibitory avoidance

₁₉₄ vs. reward-seeking and 2) whether the relationship between mood disorder symptoms and

₁₉₅ accuracy is moderated by gender.

To address these questions, the present study examined both avoidance and reward-seeking, each with two non-clinical samples - undergraduates and online workers - with a broad distribution of mood disorder symptoms. Both studies were reverse-translated with modification from a series of rodent studies investigating deficits in active and inhibitory avoidance and reward-seeking behaviours (Capuzzo and Floresco, 2020; Piantadosi, Yeates, and Floresco, 2018). Our studies are the first to combine intermixed active and inhibitory avoidance (Levita, Hoskin, and Champi, 2012) or reward-seeking trials with increasing effort requirements throughout the task, requiring participants to switch between withholding physical effort on inhibitory trials and deploying increasing amounts of effort on active trials in each task. This design allows us to directly compare performance on active and inhibitory trials in the context of increasing effort demands. Increasing effort demands may also pull out differences in choosing an active vs. inhibitory strategy.

Materials and Methods

Participants

We powered each study to detect a moderate-sized main effect of $d = 0.15$ obtained with a previous study of $N = 217$ participants using the `fabs` R package (Biesanz, 2020), resulting in a target sample size of $N = 549$. Demographic information for all studies can be found in Table 1. For each study, we collected data from two samples: an undergraduate population and an online worker population. The study was approved by the University of British Columbia Behavioural Research Ethics Board (BREB) under certificate H20-01388.

Study 1 (Avoidance)

We recruited undergraduate participants from the University of British Columbia Department of Psychology's Human Subject Pool. These participants received one percentage point towards their grade in a psychology course of their choosing for

completing the study. Of these participants, $N = 311$ finished the study, of which $N = 38$ were excluded for not completing the pre-task survey, having below 50% accuracy on active or inhibitory avoidance trials, spending over 100 s on any given attention check, or not responding to all Beck Anxiety Inventory (BAI) questions. As such, data from $N = 273$ participants was used in the data analysis.

Additionally, we recruited paid online workers from around the world ($N = 310$) on the Prolific online study platform (https://www.prolific.co/). These participants received GBP £8.07 for completing the study. Of these participants, $N = 294$ finished the study, of which $N = 22$ were excluded for not completing the pre-task survey, having below 50% accuracy on active or inhibitory avoidance trials, spending over 100 s on any given attention check, or not responding to all Beck Anxiety Inventory (BAI) questions. As such, data from $N = 272$ participants was used in the data analysis.

Study 2 (Reward-seeking)

We recruited undergraduate participants at a location which will be identified if the article is published. These participants received one percentage point towards their grade in a psychology course of their choosing and a CAD $5.00 gift card from Starbucks for completing the study. Of these participants, $N = 83$ finished the study, of which $N = 43$ were part of a separate analysis with different stimuli that is beyond the scope of this paper and $N = 7$ were excluded for not completing the pre-task survey, having below 50% accuracy on active or inhibitory avoidance trials, spending over 100 s on any given attention check, or incorrectly responding to a pre-task attention check. As such, data from $N = 36$ participants was used in the data analysis.

Additionally, we recruited paid online workers from around the world ($N = 309$) on the Prolific online study platform. These participants received GBP £8.07 and a £2.69 bonus for completing the study. Of these participants, $N = 300$ finished the study, of which

$N = 26$ were excluded for not completing the pre-task survey, having below 50% accuracy on active or inhibitory avoidance trials, spending over 100 s on any given attention check, or incorrectly responding to a pre-task attention check. As such, data from $N = 274$ participants was used in the data analysis.

Overall, the excluded sample was 29.17% female and 70.83% male, while the analyzed sample was 45.97% female and 54.03% male.

Materials and Methods

Stimulus presentation

We used PsychoPy 2020.1.2 (RRID: SCR_006571) via the Pavlovia online study platform (Peirce, Gray, Simpson, MacAskill, Höchenberger, Sogo, Kastman, and Lindeløv, 2019). Participants completed the study on their own computers; they were not allowed to complete the study on mobile devices or tablets.

Stimuli

Cues indicating active or inhibitory trials were dark blue squares and circles with a thin black border and were generated by PsychoPy 2020.1.2 (Peirce, Gray, Simpson, MacAskill, Höchenberger, Sogo, Kastman, and Lindeløv, 2019) (Fig. 1); they subtended a visual angle of about 11.5° x 11.5°. All stimuli were presented against a grey background (RGB value [0,0,0] on a scale from -1 to 1). If participants responded incorrectly on any trial in the avoidance studies, an aversive sound was played for 2000 ms. The aversive sounds were randomly selected from a set of eight screeching and scraping sounds created by our lab and ranked as highly aversive by four independent raters and in a pilot study.

Participants completed a series of questionnaires before beginning the main task. These were the State-Trait Anxiety Inventory, form Y-2 (STAI Y-2) (Spielberger, 2008); the Beck Depression Inventory II (BDI-II) (Beck, Steer, and Brown, 1996); the Beck

Anxiety Inventory (BAI) (Steer and Beck, 1997); the Behavioral Activation for Depression Scale (BADS) (Kanter, Mulick, Busch, Berlin, and Martell, 2007); the Generalized Anxiety Disorder Scale (GAD-7) (Spitzer, Kroenke, Williams, and Löwe, 2006); and the Behavioural Inhibition Scale and Behavioural Activation Scale (BIS/BAS) (Carver and White, 1994). In our data analysis, we looked at results from the BDI-II and BAI as these clinically validated scales most directly capture participants' levels of current depressive and anxiety symptoms. The BADS, GAD-7, and BIS/BAS capture specific behavioural facets of depression and anxiety that are less relevant to understanding overall effects of mood and anxiety disorders on avoidance and reward-seeking and were not analyzed in this study. We used the BAI as our primary measure of anxiety symptoms as it is the most widely used and validated among the anxiety scales we included (Fydrich, Dowdall, and Chambless, 1992) and as its structure parallels that of the BDI-II.

Procedure

Avoidance task

A graphical overview of the avoidance task is provided in Fig. 1A.

After calibrating their physical effort capability and the volume of the aversive stimuli in the task, participants read instructions indicating the shape to which they would have to respond with multiple spacebar presses as well as the shape to which they would have to withhold their response. They also heard an example of the aversive sound that would be played if they made an incorrect decision during the task.

Participants then began a series of practice trials in order to gain exposure to the stimuli and types of responses they would have to make (Fig. 1A). This consisted of five trials in which participants had to make an active avoidance response - pressing the spacebar several times to avoid hearing an unpleasant sound; five trials in which participants had to make an inhibitory avoidance response - not pressing the spacebar to

avoid hearing an aversive sound; and ten trials that intermixed these active and inhibitory trials.

On each trial, participants first viewed a grey screen with a white fixation cross for a mean duration of 2000 ms with a standard deviation (SD) of 1200 ms, jittered according to a normal distribution with these parameters on each trial. Participants then saw a visual cue - either a blue circle or a blue square - for 2000 ms. The cues used for active and inhibitory trials were pseudorandomly intermixed between participants. While this cue was on-screen, participants had to press the spacebar multiple times on active avoidance trials or withhold pressing on inhibitory avoidance trials. On active trials, the number of presses required was set according to the average number of presses made during the two effort calibration trials, such that participants who pressed fewer times during the calibration would have to press fewer times to achieve criterion during the task. The initial criterion was 5 presses given an average of 18 or fewer presses during calibration; a criterion of 6 presses given an average of 19-33 presses inclusive during calibration; and 7 presses given an average of 19 or more presses during calibration.

If participants made an incorrect decision (pressing an insufficient number of times on active trials or pressing at all on inhibitory trials), participants heard an aversive sound and saw a fixation cross for 2000 ms. This aversive sound was taken from a set of ten sounds created by our lab and rated as highly aversive. All sounds were scraping sounds that had unpleasant psychoacoustic properties shown to reliably induce aversive responses (Neumann and Waters, 2006) at a variety of frequencies. If participants made a correct decision (pressing a sufficient number of times on active trials or not pressing on inhibitory trials), they saw a fixation cross surrounded by a white border on the edges of the screen for 500 ms.

After completing the practice trials and viewing a final screen reminding them of the instructions, participants began the main task. This consisted of up to 168 active

avoidance trials and 72 inhibitory avoidance trials (70% active and 30% inhibitory),
pseudorandomized such that no more than 6 active trials or 3 inhibitory trials appeared in
a row. On the 15th trial and every 40 trials thereafter, an attention check appeared asking
participants to press a key corresponding to the letter they heard, to ensure that they were
attending to the task and able to hear auditory stimuli. Every 20 trials, the number of
button presses required on active trials increased by one press - this increased the effort
demands on active trials across the task. The task continued until the participant
responded correctly on half or less than half of the last 20 active trials - at this point, the
breakpoint was reached and the participant was thanked for completing the task.

Reward-seeking task

A graphical overview of the reward-seeking task is provided in Fig. 1B.

The design of the reward-seeking task was identical to that of the avoidance task,
with the following exceptions. First, the practice blocks were based on criterion-based
advancement in order to increase consistency with the design of other reward-seeking
studies in our lab. Participants had to achieve at least 80% accuracy in each of the active,
inhibitory, and intermixed reward-seeking trial blocks in order to advance; each block
would repeat until they achieved each criterion. Second, if the participant made a correct
decision during a trial, they would see a screen indicating that they had gained 5 points
along with a sum of their points thus far; if the participant made an incorrect decision
during a trial, they would see a screen indicating that they had gained 0 points along with
a sum of their points thus far. Both screens appeared for 1500 ms. Undergraduate
participants received a CAD $5 gift card as a reward in addition to course credit for
completing the task; online workers received a GBP £2.69 payment as a reward in addition
to their payment for completing the task. Last, as this task did not incorporate audio, no
volume check or audio-based attention check was included.

Data analysis

All data was analyzed using R 4.1.1 "Kick Things" (R Development Core Team, 2011) through RStudio (Booth et al., 2018). On each behavioural task, we measured 1) accuracy on active and inhibitory trials, operationalized as the proportion of correct responses on each trial type; 2) effort on each trial type, operationalized as the number of presses made relative to criterion on each trial; 3) participants' depressive and anxiety symptoms, operationalized as their BDI (Beck, Steer, and Brown, 1996) and BAI (Steer and Beck, 1997) scores respectively; 4) participants' sensitivity (d') and criterion (β); and 5) breakpoint, operationalized as the trial number on which the participant responded correctly on half or less than half of the last 20 active trials. For each study, we combined undergraduate and online worker samples to obtain a gender-balanced sample. All analyses were Bonferroni corrected for multiple comparisons. All confidence intervals were based on 1000 bootstrap replications using the `confintr` R package (Mayer, 2022).

Results

Demographics

Participant's reported gender and sex heavily overlapped, with 97.53% overlap in women and 97.67% overlap in men on the avoidance tasks and 98.84% overlap in women and 97.80% overlap in men on the reward-seeking task. For this reason, the following results are expressed in terms of gender only. Women reported higher levels of depressive symptoms ($t(723.12)$ = -3.84, $p < .001$, $d = 0.27$) and anxiety symptoms ($t(706.44)$ = -4.71, $p < .001$, $d = 0.34$) than men across samples (Table 2, Fig. 2A, 5A). Across both studies, 20.92% of women and 15.11% of men were on medication for depression, and 19.29% of women and 14.89% of men were on medication for anxiety.

Avoidance task

We first compared participants' accuracy on active and inhibitory avoidance trials by gender using a 2x2 within-between ANOVA (Trial Type x Gender) (Fig. 2B) (Table 3). A main effect revealed that, on average, participants were more accurate on inhibitory than on active trials ($F(1, 522) = 405.53$, p $< .001$, $\eta^2 = 0.25$; Table 4). There was a main effect of gender ($F(1, 522) = 5.85$, $p = .02$, $\eta^2 = < .01$) as well as a gender interaction such that men had higher accuracy than women on active but not inhibitory avoidance trials ($F(1, 522) = 4.87$, $p = .03$, $\eta^2 < .01$).

Depression and anxiety symptoms were not significantly associated with inhibitory or active avoidance accuracy (Table 4). However, given observed gender differences in both depression/anxiety disorder symptoms and task performance, we conducted a series of moderation analyses using the PROCESS macro (Hayes, 2017) as implemented in `bruceR` (Bao, 2021) to examine whether gender moderated the relationship between accuracy and anxiety and depressive symptoms (Fig. 3B). We found that gender moderated the relationship between anxiety symptoms (as measured in BAI scores) and inhibitory avoidance accuracy ($F(1, 541) = 4.29$, $p = .04$; Fig. 3A). Increased BAI scores were associated with lower inhibitory avoidance accuracy in females but not males; this relationship was not observed for BDI scores. For the full results of the moderation analysis, see Table 5.

To further examine the relationship between accuracy and anxiety and depressive symptoms, we tested whether BAI scores predicted accuracy in men and women separately. Outside of the moderation analysis after Bonferroni correction, BAI scores overall predicted inhibitory but not active avoidance performance accuracy (Fig. 3B, ($t(543) =$ -3.01, $p = .04$, $r = $-0.13)). Additionally, higher BAI scores predicted lower inhibitory avoidance performance accuracy in women but not men (Fig. 3B; ($t(260) = $-3.73, $p = .01$, $r = $-0.23, *95% CI* [-0.36, -0.09])).

We also conducted an interactive regression analysis to investigate whether the relationship between anxiety symptoms and active and inhibitory avoidance accuracy differed as a function of the level of depressive symptoms (Fig. 4). In active avoidance, the relationship between anxiety symptoms (BAI scores) and accuracy for all participants was a function of their level of depressive symptoms (BDI scores) ($t(541) = -2.65$, $p = .01$); however, this was not the case in inhibitory avoidance ($t(541) = -0.87$, $p = .39$). As such, only those with higher levels of depressive symptoms were impaired in active avoidance, and not those with higher levels of anxiety symptoms but lower levels of depressive symptoms.

To account for participants' bias to deploy effort and engage in active as opposed to inhibitory avoidance in general, we calculated sensitivity (d') and criterion (β) and repeated the moderation analyses on these alternative outcome performance measures. Sensitivity reflects participants' ability to correctly distinguish between active and inhibitory trials and deploy the required amount of effort on active trials only. Criterion reflects participants' bias towards an active or inhibitory response. Gender did not moderate the relationship between anxiety or depressive symptoms and sensitivity or criterion. However, BAI scores predicted sensitivity in women but not men ($t(260) = -3.55$, $p = .01$, $r = -0.21$, *95% CI* [-0.36, -0.08])).

Additionally, we explored the extent to which the amount of effort that participants deployed to avoid aversive outcomes changed across the task (Fig. 8A). A a 2x2 within-between ANOVA (Block x Gender) (Table 6) revealed that participants deployed increasing amounts of effort during the task to meet increasing effort requirements ($F(1, 522) = 405.53$, $p < .001$, $\eta^2 = 0.25$), and that men deployed more effort than women across the task ($F(1, 522) = 5.85$, $p = 0.016$, $\eta^2 = < .01$).

Last, we examined whether participants' breakpoint in the task was explained by their levels of mood and anxiety disorder symptoms in avoidance. After Bonferroni correction, BAI and BDI scores did not predict overall active or inhibitory avoidance

421 performance in either men or women (BAI: ($t(543)$ = -2.44, p = .24, r = -0.10, *95% CI*

422 [-0.19, -0.02]); BDI: ($t(543)$ = -2.15, p = .51, r = -0.09, *95% CI* [-0.17, -0.01])).

Reward-seeking task

We first compared participants' accuracy on active and inhibitory reward-seeking

trials by gender (Fig. 6A) using a 2x2 within-between ANOVA (Trial Type x Gender)

(Table 7). A main effect revealed that, on average, participants were more accurate on

inhibitory than on active trials ($F(1, 292)$ = 58.86, p < .001, η^2 = 0.07). There was no

main effect of gender ($F(1, 292)$ = 2.61, p = .11, η^2 = 0.01) but there was a gender

interaction such that men had higher accuracy than women on active but not inhibitory

reward-seeking trials ($F(1, 292)$ = 7.61, p = .01, η^2 = 0.01).

Given observed gender differences in both depressive/anxiety symptoms and task

performance, we again conducted a series of moderation analyses to see whether gender

moderated the relationship between accuracy and anxiety and depressive symptoms.

Anxiety was associated with reduced active reward seeking accuracy in the total sample.

However, we found no gender moderation of the relationship between anxiety or depressive

symptoms and active or inhibitory reward-seeking accuracy, and BDI scores did not

interact with active or inhibitory reward-seeking accuracy (Table 8, Table 9, Fig. 6A).

To further elucidate the relationship between accuracy and anxiety and depressive

symptoms, we evaluated whether BAI and BDI scores predicted accuracy in men and

women separately. After Bonferroni correction, BAI and BDI scores did not predict overall

active or inhibitory reward-seeking performance in either men or women (Fig. 6B). We also

conducted an interactive regression analysis to investigate whether the relationship

between anxiety symptoms and active and inhibitory reward-seeking accuracy differed as a

function of their level of depressive symptoms (Fig. 7). In active reward-seeking, the

relationship between anxiety symptoms (BAI scores) and accuracy for all participants was

not a function of their level of depressive symptoms (BDI scores) ($t(306) = 0.45$, $p = .66$);

this was also not the case in inhibitory reward-seeking ($t(306) = $ -0.78, $p = .44$).

To evaluate participants' bias to deploy effort and engage in active as opposed to

inhibitory reward-seeking, we again calculated sensitivity (d') and criterion (β) and

performed the moderation analysis using these alternative outcome measures of

performance. Gender did not moderate the relationship between anxiety or depressive

symptoms and sensitivity or criterion.

Additionally, we explored how the effort that participants deployed to obtain reward

changed across the task (Fig. 8B). A a 2x2 within-between ANOVA (Block x Gender)

(Table 10) revealed that participants deployed increasing amounts of effort during the task

to meet increasing effort requirements ($F(1, 292) = 58.86$, p = < .001, $\eta^2 = 0.07$), and

revealed no gender differences in effort deployment ($F(1, 292) = 2.61$, p = .11, $\eta^2 = 0.01$).

Last, we examined whether participants' breakpoint in the task was explained by

their levels of mood and anxiety disorder symptoms in reward-seeking. After Bonferroni

correction, BAI and BDI scores did not predict overall active or inhibitory reward-seeking

performance in either men or women (BAI: ($t(308) = $ -0.31, $p = 1.00$, $r = $ -0.02, *95% CI*

[-0.12, 0.04]); BDI: ($t(308) = $ -0.75, $p = 1.00$, $r = $ -0.04, *95% CI* [-0.12, 0.04])).

Discussion

Summary

In the present study, we investigated gender differences in active and inhibitory

avoidance behaviours. Additionally, we explored how mood and anxiety disorder symptoms

negatively impact active vs. inhibitory avoidance and reward-seeking behaviours. Lastly,

we examined whether accuracy and effort deployment in avoidance and reward-seeking are

associated with depressive and anxiety symptoms, and whether the relationship between

accuracy and mood disorder symptoms is moderated by gender. Anxiety symptoms

interacted with participants' depressive symptoms level in active avoidance but not in active reward-seeking or in inhibitory avoidance or reward-seeking, such that participants' anxiety symptoms negatively impacted their accuracy more if they also had a high level of depressive symptoms. Compared to men, women showed higher levels of self-reported depression and anxiety. Throughout active avoidance and reward-seeking trials, men made more effortful responses than women. Gender moderated the relationship between anxiety symptoms and inhibitory avoidance. Women showed lower performance in inhibitory avoidance than men as a function of higher levels of anxiety symptoms, with no corresponding effect for depressive symptoms. In contrast, higher levels of anxiety were associated with lower active reward-seeking performance in both men and women. Our findings illuminate gender differences in active and inhibitory subtypes of avoidance and reward-seeking behaviours where effort was required to obtain the desired outcome.

Interpretation of results

        Based on previous findings, we can speculate that the gender differences are driven at least in part by mood and anxiety disorder symptoms. In particular, the observed gender difference in inhibitory avoidance performance could reflect more general gender differences in stress tolerance. Parker and Brotchie (2010) argued that women have a higher predisposition (diathesis) to stress than men and, in our study, women reported higher anxiety symptoms than men. In our avoidance task, it is advantageous to act to avoid an aversive outcome - especially given high levels of anxiety symptoms (Bishop and Gagne, 2018). As such, inhibiting responding may be especially difficult given a combination of high anxiety symptoms and a decreased tolerance for stress in women compared to men. Furthermore, this gender difference in diathesis could drive a reduced ability to inhibit effort when needed - an impairment in shifting from an active to an inhibitory strategy given higher levels of anxiety symptoms (Gustavson, Altamirano, Johnson, Whisman, and Miyake, 2017). Given the observed interaction of lower accuracy

at higher depressive and anxiety symptom levels in active avoidance, increased comorbidity of depressive and anxiety symptoms in women could also partly explain these results - although this interaction was not significantly observed in the presence of gender differences in inhibitory avoidance.

Beyond inhibitory avoidance, the observed gender differences in effort and accuracy on active avoidance and reward-seeking trials could reflect gender differences in effort deployment. Men made more effortful responses relative to criterion than women on active avoidance trials. This could be caused by women having smaller wrists with which to generate physical force than men (Morse, Jung, Bashford, and Hallbeck, 2006), as well as increased testosterone in men - which is associated with increased physical effort deployment during competitions (Losecaat Vermeer, Riečanský, and Eisenegger, 2016). Although our tasks did not have competitive elements, participants may still have completed the task with an eye towards maximizing performance. Since deploying more effort in the task would increase one's chance of staying above criterion, this increased effort deployment could explain the increased active trial accuracy for men across the avoidance and reward-seeking tasks. It is important to qualify that gender has a significant cultural component, and cultural factors could also play a role in gender differences in effort deployment - perhaps via effects of a lower tolerance for stress on effort deployment (Parker and Brotchie, 2010).

We also found that the effects of anxiety symptoms on active avoidance accuracy depended on an interaction between anxiety symptoms and depressive symptom level, such that only those with higher levels of depression symptoms who were also high in anxiety were impaired. This suggests a role for comorbidity between anxiety and depressive symptoms in people's ability to deploy effort to actively avoid an unpleasant outcome (Brown, Campbell, Lehman, Grisham, and Mancill, 2001). As such, when those with mood disorder symptoms attempt to avoid aversive outcomes, impairments to effort deployment

necessary for avoidance may be driven by features of anxiety symptoms and depressive symptoms together, as opposed to one mood disorder or another independently. Past work has only predicted increases in effort deployment in avoidance given anxiety and decreases in effort deployment in general given depression (Bishop and Gagne, 2018). As such, the observed impairment given symptoms of both mood disorders sheds light on how comorbid mood disorder symptoms can impact avoidance behaviours. Importantly, this relationship was not observed in inhibitory avoidance or in active or inhibitory reward-seeking. In inhibitory avoidance, mood disorder effects were specific to anxiety symptoms in women only, and only anxiety symptoms significantly impacted active reward-seeking accuracy. As such, it may be that an anxious subtype of depressive symptoms uniquely impacts our ability to deploy effort to avoid an unpleasant outcome. This is inconsistent with previous findings by Wurst, Schiele, Stonawski, Weiß, Nitschke, Hommers, Domschke, Herrmann, Pauli, Deckert, and Menke (2021) suggesting that there were no significant differences in fear learning – which may be a process involved in avoidance – between those with anxious and non-anxious depression. However, as our task's difficulty is bound by increasing effort and not learning, and as we were looking at a non-clinical sample, such effects of comorbidity may not have come out in the context of our study.

   The relationship between higher anxiety symptoms and reduced active reward-seeking performance is novel, as past work has primarily focused on the relationship between depressive symptoms and reward-seeking and has not focused on subtypes of avoidance (Bishop and Gagne, 2018). The ways in which depressive symptoms drive reduced reward sensitivity are well-established (Slaney, Perkins, Davis, Penton-Voak, Munafò, Houghton, and Robinson, 2021), but it was previously not clear whether anxiety symptoms would drive greater success in obtaining rewards (through underestimation of the effort required to obtain a reward) or reduced reward-seeking performance (through inefficient allocation of effort that results in increased fatigue). In our task, participants' performance on effortful active trials was reduced given higher anxiety symptoms,

suggesting that the form of anxiety symptoms participants faced may not have driven

increased effort deployment or reduced effort costs. This finding runs counter to past work

suggesting that trait anxiety is associated with impairments in shifting away from an

effortful task (Gustavson, Altamirano, Johnson, Whisman, and Miyake, 2017). As we

evaluate participants' current as opposed to trait anxiety, this pattern of results could be

explained by anxiety-related impairments to effort deployment that are reflected in

measures of anxiety as a personality trait.

Similarly, the relationship between higher anxiety symptoms and reduced inhibitory

avoidance performance differs from previous predictions of improved avoidance given

anxiety symptoms, such as those of Bishop and Gagne (2018). However, Bishop and Gagne

framed this relationship in terms of active and not inhibitory avoidance, as they predicted

that underestimations of effort cost would drive excessive avoidance behaviours. Anxiety

symptoms may be associated with impairments to inhibitory avoidance precisely because of

this bias towards action given the possibility of aversive outcomes, an effect that could be

driven by a perceived lack of control over outcomes in the task (Wang and Delgado, 2021).

Additionally, we did not observe a relationship between depressive symptoms and accuracy

or effort deployment in reward-seeking, as has previously been observed (Bishop and

Gagne, 2018). The effort demands of the task may not have deterred people with high

depressive symptoms from working for a reward, which could be explained by the

depressive symptoms experienced by participants not impacting effort deployment.

Overall, participants had lower accuracy in avoidance compared to reward-seeking;

this could be a function of differences in motivation to engage in avoidance or

reward-seeking. Motivation to complete the tasks can be driven in part by participants'

valuations of task-relevant stimuli (Bishop and Gagne, 2018). A major difference between

our tasks arises in the outcome of an incorrect decision. In avoidance, an incorrect decision

is associated with an aversive sound; in reward-seeking, it is associated with not receiving

points. Although the salience of an aversive sound may suggest that it is more motivating

and would therefore be associated with increased accuracy, hearing it may also be more

demotivating - especially for participants with mood disorder symptoms. Hearing the

aversive sound repeatedly could be a salient indicator of a lack of control over task

outcomes (Wang and Delgado, 2021).

Limitations

There are some limitations to our interpretation of our findings. First, since the

dichotomy of the task demands is between effortful active trials and inhibitory trials that

require no effort, we cannot compare the effects of high vs. low effort demands on

inhibitory avoidance or reward-seeking behaviours. As such, our interpretation of the

relationship between effort deployment and mood disorder symptoms only extends to

active trials. Accuracy in the task was likely tied to participants' effort capabilities, as

increased effort deployment was required throughout the task on active trials to meet the

criterion level of effort and make the correct response on the trial. However, we calibrated

the criterion to participants' effort ability and considered accuracy on inhibitory as well as

active trials to reduce the reliance of task outcomes on individual differences in effort

deployment. Additionally, as the proportion of active trials was greater than that of

inhibitory trials, participants may have become fatigued on the majority of trials in the

task, increasing over time. This fatigue from effort deployment, combined with boredom

(from the task being repetitive) could be difficult to disentangle from other shifts in

motivation to deploy effort throughout the task (e.g. those related to the value of avoidance

or reward-seeking). However, as fatigue is likely to arise in most physically effortful tasks,

our tasks still reflect real-world physical effort demands. Furthermore, as this study took

place online, the study had to use repeated keyboard presses instead of other, more

continuous or better-controlled measures of physical effort such as a grip squeeze (Aridan,

Malecek, Poldrack, and Schonberg, 2019). However, repeated button presses have been

validated as being physically effortful and have been used in in-person contexts (Gold,

Strauss, Waltz, Robinson, Brown, and Frank, 2013).

Future work

Future studies could build on our findings by investigating how patterns of
information about specific aspects of effortful avoidance and reward-seeking are instantiated
in key brain regions. The posterior anterior cingulate cortex (pACC) and ventral striatum
encode information about prospective gains given physical effort requirements (Aridan,
Malecek, Poldrack, and Schonberg, 2019). These regions - and their homologues in rodents
- have been shown to be differentially necessary for active vs. inhibitory avoidance
(Piantadosi, Yeates, and Floresco, 2018) and reward-seeking (Capuzzo and Floresco, 2020).
Investigating how these regions represent information on prospective gains and losses
relative to effort costs could illuminate how we weigh the benefits and costs of deploying
effort to obtain rewards and avoid aversive outcomes. Additionally, separating out different
factors contributing to effort deployment through a computational modelling approach
would be important to understand the individual contributions of various factors to
participants' performance. These factors could include action biases (Mkrtchian, Aylward,
Dayan, Roiser, and Robinson, 2017), perceived value of avoidance or reward (Bishop and
Gagne, 2018); or fatigue (Pessiglione, Vinckier, Bouret, Daunizeau, and Le Bouc, 2018).
Furthermore, it would be helpful to evaluate whether subscales of mood disorder symptoms
- potentially linked to subtypes such as anxious depression (Wurst, Schiele, Stonawski,
Weiß, Nitschke, Hommers, Domschke, Herrmann, Pauli, Deckert, and Menke, 2021) - pull
out factors that drive participants' behaviours in avoidance and reward seeking. This
analysis could further illuminate our observed gender differences - for example, to evaluate
whether reduced inhibitory avoidance performance in women given increased anxiety
symptoms is reflective of an anxious subtype of depression (Wurst, Schiele, Stonawski,
Weiß, Nitschke, Hommers, Domschke, Herrmann, Pauli, Deckert, and Menke, 2021).

Conclusion

Our studies address outstanding questions of whether accuracy and effort deployment on avoidance and reward-seeking predict mood disorder symptoms, and whether the relationship between accuracy and mood disorder symptoms is moderated by gender. We separate out active and inhibitory avoidance and reward-seeking behaviours in a context that allows for direct comparisons between them, instead of considering avoidance and reward-seeking behaviours as unitary wholes. We highlight gender differences in each of these subtypes of avoidance and reward-seeking given varying levels of mood disorder symptoms, contextualizing past work on gender differences in these symptoms (Parker and Brotchie, 2010). In particular, we are the first to examine these proposed gender differences in an active and inhibitory avoidance and reward-seeking context. These findings could inform clinical interventions to address maladaptive deployment of avoidance and lack of motivation for reward-seeking, targeted by gender. Additionally, we link active avoidance and reward-seeking to motivation for physical effort deployment given varying levels of mood and anxiety disorders. As many tasks in life require physical effort deployment, understanding where it can be impaired is an important pursuit. Our findings underscore the importance of considering individual differences in the ways in which avoidance and reward-seeking can be impaired in life.

Data and code availability

The data and materials for all experiments, as well as the code used to generate this manuscript and conduct all analyses, are available at https://osf.io/2rd3f/.

References

Altemus M, Sarvaiya N, Neill Epperson C (2014) Sex differences in anxiety and depression clinical perspectives. Frontiers in Neuroendocrinology 35:320–330.

Aridan N, Malecek NJ, Poldrack RA, Schonberg T (2019) Neural correlates of effort-based valuation with prospective choices. NeuroImage 185:446–454.

Auerbach RP, Pagliaccio D, Hubbard NA, Frosch I, Kremens R, Cosby E, Jones R, Siless V, Lo N, Henin A, Hofmann SG, Gabrieli JDE, Yendiki A, Whitfield-Gabrieli S, Pizzagalli DA (2022) Reward-Related Neural Circuitry in Depressed and Anxious Adolescents: A Human Connectome Project. Journal of the American Academy of Child & Adolescent Psychiatry 61:308–320.

Bao H-W-S (2021) bruceR: Broadly useful convenient and efficient r functions.

Beck AT, Steer RA, Brown GK (1996) Manual for the beck depression inventory-II. San Antonio, TX: Psychological Corporation.

Biesanz J (2020) Fabs: Functions for applied behavioural sciences in r.

Bishop SJ, Gagne C (2018) Anxiety, depression, and decision making: A computational perspective. Annual Review of Neuroscience 41:371–388.

Booth DS et al. (2018) RStudio: Integrated development for r. Nature.

Brown TA, Campbell LA, Lehman CL, Grisham JR, Mancill RB (2001) Current and lifetime comorbidity of the DSM-IV anxiety and mood disorders in a large clinical sample. Journal of Abnormal Psychology 110:585–599.

Browning M, Behrens TE, Jocham G, O'Reilly JX, Bishop SJ (2015) Anxious individuals have difficulty learning the causal statistics of aversive environments. Nature Neuroscience 18:590–596.

Capuzzo G, Floresco SB (2020) Prelimbic and infralimbic prefrontal regulation of active and inhibitory avoidance and reward-seeking. Journal of Neuroscience 40:4773–4787.

Carver CS, White TL (1994) Behavioral inhibition, behavioral activation, and

affective responses to impending reward and punishment: The BIS/BAS scales.
Journal of Personality and Social Psychology.

Cléry-Melin ML, Schmidt L, Lafargue G, Baup N, Fossati P, Pessiglione M (2011)
Why don't you try harder? An investigation of effort production in major
depression. PLoS ONE 6.

Cools R (2008) Role of Dopamine in the Motivational and Cognitive Control of
Behavior. The Neuroscientist 14:381–395.

Culbreth AJ, Moran EK, Barch DM (2018) Effort-cost decision-making in psychosis
and depression: Could a similar behavioral deficit arise from disparate
psychological and neural mechanisms? Psychological Medicine 48:889–904.

Dillon DG, Rosso IM, Pechtel P, Killgore WDS, Rauch SL, Pizzagalli DA (2014)
Peril and pleasure: An RDOC-inspired examination of threat responses and
reward processing in anxiety and depression. Depression and Anxiety
31:233–249.

Fydrich T, Dowdall D, Chambless DL (1992) Reliability and validity of the beck
anxiety inventory. Journal of Anxiety Disorders 6:55–61.

Gold JM, Strauss GP, Waltz JA, Robinson BM, Brown JK, Frank MJ (2013)
Negative symptoms of schizophrenia are associated with abnormal effort-cost
computations. Biological Psychiatry 74:130–136.

Goldstein-Piekarski AN, Williams LM, Humphreys K (2016) A trans-diagnostic
review of anxiety disorder comorbidity and the impact of multiple exclusion
criteria on studying clinical outcomes in anxiety disorders. Translational
Psychiatry 6:e847–e847.

Gustavson DE, Altamirano LJ, Johnson DP, Whisman MA, Miyake A (2017) Is set
shifting really impaired in trait anxiety? Only when switching away from an
effortfully established task set. Emotion 17:88–101.

Hayes AF (2017) Introduction to mediation, moderation, and conditional process

analysis: A regression-based approach. Guilford publications.

Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, Sanislow C, Wang

P (2010) Research Domain Criteria (RDoC): Toward a New Classification

Framework for Research on Mental Disorders. American Journal of Psychiatry

167:748–751.

Kanter JW, Mulick PS, Busch AM, Berlin KS, Martell CR (2007) The Behavioral

Activation for Depression Scale (BADS): Psychometric Properties and Factor

Structure. Journal of Psychopathology and Behavioral Assessment 29:191–202.

Kessler RC (2006) The epidemiology of depression among women. Women and

Depression: A Handbook for the Social, Behavioral, and Biomedical Sciences

74:22–38.

Krypotos AM, Effting M, Kindt M, Beckers T (2015) Avoidance learning: A review

of theoretical models and recent developments. Frontiers in Behavioral

Neuroscience 9.

LeDoux JE, Moscarello J, Sears R, Campese V (2017) The birth, death and

resurrection of avoidance: A reconceptualization of a troubled paradigm.

Molecular Psychiatry 22:24–36.

Levita L, Hoskin R, Champi S (2012) Avoidance of harm and anxiety: A role for the

nucleus accumbens. NeuroImage 62:189–198.

Losecaat Vermeer AB, Riečanský I, Eisenegger C (2016) Competition, testosterone,

and adult neurobehavioral plasticity, pp213–238. Elsevier.

Mayer M (2022) Confintr: Confidence intervals.

McLean CP, Asnaani A, Litz BT, Hofmann SG (2011) Gender differences in anxiety

disorders: Prevalence, course of illness, comorbidity and burden of illness.

Journal of Psychiatric Research 45:1027–1035.

Mkrtchian A, Aylward J, Dayan P, Roiser JP, Robinson OJ (2017) Modeling

avoidance in mood and anxiety disorders using reinforcement learning.

Biological Psychiatry 82:532–539.

Morse JL, Jung M-C, Bashford GR, Hallbeck MS (2006) Maximal dynamic grip

force and wrist torque: The effects of gender, exertion direction, angular velocity,

and wrist angle. Applied Ergonomics 37:737–742.

Neumann DL, Waters AM (2006) The use of an unpleasant sound as an

unconditional stimulus in a human aversive pavlovian conditioning procedure.

Biological Psychology 73:175–185.

Norbury A, Robbins TW, Seymour B (2018) Value generalization in human

avoidance learning. eLife 7.

Ottenbreit ND, Dobson KS, Quigley L (2014) An examination of avoidance in major

depression in comparison to social anxiety disorder. Behaviour Research and

Therapy 56:82–90.

Parker G, Brotchie H (2010) Gender differences in depression. International Review

of Psychiatry 22:429–436.

Peirce J, Gray JR, Simpson S, MacAskill M, Höchenberger R, Sogo H, Kastman E,

Lindeløv JK (2019) PsychoPy2: Experiments in behavior made easy. Behavior

Research Methods 51:195–203.

Pessiglione M, Vinckier F, Bouret S, Daunizeau J, Le Bouc R (2018) Why not try

harder? Computational approach to motivation deficits in neuro-psychiatric

diseases. Brain 141:629–650.

Piantadosi PT, Yeates DCM, Floresco SB (2018) Cooperative and dissociable

involvement of the nucleus accumbens core and shell in the promotion and

inhibition of actions during active and inhibitory avoidance. Neuropharmacology

138:57–71.

Pittig A, Treanor M, LeBeau RT, Craske MG (2018) The role of associative fear

and avoidance learning in anxiety disorders: Gaps and directions for future

research. Neuroscience & Biobehavioral Reviews 88:117–140.

R Development Core Team R (2011) R: A language and environment for statistical computing.

Rizvi SJ, Pizzagalli DA, Sproule BA, Kennedy SH (2016) Assessing anhedonia in depression: Potentials and pitfalls. Neuroscience and Biobehavioral Reviews 65:21–35.

Slaney C, Perkins AM, Davis R, Penton-Voak I, Munafò MR, Houghton CJ, Robinson ESJ (2021) Objective measures of reward sensitivity and motivation in people with high vs low anhedonia.

Spielberger C (2008) State-trait anxiety inventory for adults. Garden.

Spitzer RL, Kroenke K, Williams JBW, Löwe B (2006) A Brief Measure for Assessing Generalized Anxiety Disorder: The GAD-7. Archives of Internal Medicine 166:1092.

Steer RA, Beck AT (1997) Beck anxiety inventory., pp23–40. Lanham, MD, US: Scarecrow Education.

Thompson AH, Bland RC (2018) Gender similarities in somatic depression and in DSM depression secondary symptom profiles within the context of severity and bereavement. Journal of Affective Disorders 227:770–776.

Treadway MT, Bossaller NA, Shelton RC, Zald DH (2012) Effort-based decision-making in major depressive disorder: A translational model of motivational anhedonia. Journal of Abnormal Psychology 121:553–558.

Treadway MT, Buckholtz JW, Schwartzman AN, Lambert WE, Zald DH (2009) Worth the 'EEfRT'? The effort expenditure for rewards task as an objective measure of motivation and anhedonia. PLoS ONE 4:1–9.

Wang KS, Delgado MR (2021) The Protective Effects of Perceived Control During Repeated Exposure to Aversive Stimuli. Frontiers in Neuroscience 15:625816.

Wurst C, Schiele MA, Stonawski S, Weiß C, Nitschke F, Hommers L, Domschke K, Herrmann MJ, Pauli P, Deckert J, Menke A (2021) Impaired fear learning and

784   extinction, but not generalization, in anxious and non-anxious depression.

785   Journal of Psychiatric Research 135:294–301.

Legend

Figure 1: Trial layout diagram. A diagram of the active and inhibitory avoidance

and reward-seeking tasks. In the avoidance task (A), after an inter-stimulus interval (ISI)

with a fixation cross onscreen, participants were presented with a cue associated with

active or inhibitory avoidance. For the active avoidance cue, participants had to respond

with repeated spacebar presses to avoid hearing an aversive sound. For the inhibitory

avoidance cue, participants had to withhold responding to avoid hearing an aversive sound.

In the reward-seeking task (B), after the ISI, participants were presented with a cue

associated with active or inhibitory reward-seeking. For the active reward-seeking cue,

participants had to respond with repeated spacebar presses to obtain points towards a

monetary reward. For the inhibitory reward-seeking cue, participants had to withhold

responding to obtain points towards a monetary reward. ISI = Inter-stimulus interval.

Figure 2: Avoidance demographics and accuracy measures. (A) Distribution of

anxiety (BAI) and depressive symptom (BDI) proportion scores (score divided by total

possible score) by gender and sample. (B) Accuracy by gender on active and inhibitory

avoidance, and moderation of the relationship between accuracy and anxiety (BAI) scores

by gender. Dots represent average accuracy on each trial type by participant. BAI = Beck

Anxiety Inventory, BDI = Beck Depression Inventory.

Figure 3: Avoidance moderation and correlation analyses. (A) Moderation of the

relationship between accuracy and anxiety (BAI) proportion scores by gender in active and

inhibitory avoidance. Gender significantly moderated the relationship between anxiety

symptoms (BAI proportion scores) and inhibitory avoidance accuracy. (B) Accuracy by

gender on active and inhibitory avoidance. BAI = Beck Anxiety Inventory. Proportion

scores are scores divided by total possible score.

Figure 4: Interactive regression analyses for avoidance. Interactive regression analysis output for active and inhibitory avoidance. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory. Numbers on right indicate coefficients for regression lines at each level of BDI score (centred 1 standard deviation below the mean, at the mean, and 1 standard deviation above the mean).

Figure 5: Reward-seeking demographics and accuracy measures. Distribution of anxiety (BAI) and depressive symptom (BDI) scores by gender and sample. (B) Accuracy by gender on active and inhibitory reward-seeking, and moderation of the relationship between accuracy and anxiety (BAI) scores by gender. Proportion scores are scores divided by total possible score. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Figure 6: Reward-seeking moderation and correlation analyses. (A) Moderation of the relationship between accuracy and anxiety (BAI) proportion scores by gender on active and inhibitory reward-seeking. Anxiety symptoms (BAI proportion scores) were significantly associated with active reward-seeking accuracy. (B) Accuracy by gender on active and inhibitory reward-seeking. BAI = Beck Anxiety Inventory. Proportion scores are scores divided by total possible score.

Figure 7: Interactive regression analyses for reward-seeking. Interactive regression analysis output for active and inhibitory reward-seeking. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory. Numbers on right indicate coefficients for regression lines at each level of BDI score (centred 1 standard deviation below the mean, at the mean, and 1 standard deviation above the mean).

Figure 8: Avoidance and reward-seeking effort deployment. Effort deployment across active avoidance (A) and reward-seeking (B) trials relative to the criterion effort deployment required to avoid an aversive outcome or obtain reward (black line). Each

point represents one trial. N values represent number of participants of each gender who completed the task during that block of trials.

Table 1: Demographic information for all participants.

Table 2: Mood disorder symptom statistics. Mean and SD Beck Depression Inventory II (BDI) and Beck Anxiety Inventory (BAI) proportion scores (score divided by total possible score).

Table 3: Avoidance accuracy statistics. ANOVA of accuracy on active and inhibitory avoidance trials. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Table 4: Avoidance performance and mood disorder statistics. Relationship between accuracy and mood disorder symptoms on active and inhibitory avoidance trials. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Table 5: Avoidance gender moderation statistics. Gender moderation of relationship between accuracy and mood disorder symptoms on active and inhibitory avoidance trials. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Table 6: Avoidance effort deployment statistics by gender. ANOVA of effort deployment on active avoidance trials by gender. Block = group of 28 trials on which an effortful response was required to avoid hearing an unpleasant sound.

Table 7: Reward-seeking accuracy statistics. ANOVA of accuracy on active and inhibitory reward-seeking trials.

Table 8: Reward-seeking performance and mood disorder statistics. Relationship between accuracy and mood disorder symptoms on active and inhibitory reward-seeking trials. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Table 9: Reward-seeking gender moderation statistics. Gender moderation of

relationship between accuracy and mood disorder symptoms on active and inhibitory

reward-seeking trials. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Table 10: Reward-seeking effort deployment statistics by gender. ANOVA of effort

deployment on active reward-seeking trials by gender. Block = group of 28 trials on which

an effortful response was required to obtain a reward.

Table 1

Demographic information for all participants.

Study	$N_{recruited}$	$N_{analyzed}$	N_{female}	N_{male}	N_{other}	M_{age}	$Range_{age}$
1 (Avoidance, undergraduate)	357	273	175	86	12	20.42	17-32
2 (Avoidance, paid global)	310	272	87	176	9	24.06	18-57
3 (Reward-seeking, undergraduate)	114	36	28	8	0	20.67	18-26
4 (Reward-seeking, paid global)	309	274	78	180	16	25.18	18-62

Table 2

Mean and SD Beck Depression Inventory II (BDI) and Beck Anxiety Inventory (BAI) proportion scores (score divided by total possible score).

Task	Gender	$M_{BDIprop}$	$SD_{BDIprop}$	$M_{BAIprop}$	$SD_{BAIprop}$
Avoidance	Female	0.30	0.19	0.30	0.22
	Male	0.23	0.16	0.23	0.19
Reward-seeking	Female	0.29	0.19	0.28	0.21
	Male	0.26	0.17	0.22	0.17

Table 3

ANOVA of accuracy on active and inhibitory avoidance trials. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Effect	df_n	df_d	F	p	sig.	η^2
Gender	1	522	5.85	0.016	*	< .01
Trial Type	1	522	405.53	< .001	***	0.25
Gender:Trial Type	1	522	4.87	0.028	*	< .01

Table 4

Relationship between accuracy and mood disorder symptoms on active and inhibitory avoidance trials. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Trial Type	Comparison	df	t	p	sig.
Active avoidance	BDI~accuracy	541	-0.32	0.75	
	BAI~accuracy	541	-1.37	0.17	
Inhibitory avoidance	BDI~accuracy	541	-0.87	0.38	
	BAI~accuracy	541	-0.73	0.47	

Table 5

Gender moderation of relationship between accuracy and mood disorder symptoms on active and inhibitory avoidance trials. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Trial Type	Comparison	df_n	df_d	t	p	sig.
Active avoidance	Gender moderation of BDI~accuracy	1	541	0.01	0.92	
	Gender moderation of BAI~accuracy	1	541	0.28	0.59	
Inhibitory avoidance	Gender moderation of BDI~accuracy	1	541	1.04	0.31	
	Gender moderation of BAI~accuracy	1	541	4.29	0.04	*

Table 6

ANOVA of effort deployment on active avoidance trials by gender. Block = group of 28 trials on which an effortful response was required to avoid hearing an unpleasant sound.

Effect	df_n	df_d	F	p	sig.	η^2
Gender	3.00	516.00	7.08	$< .001$	***	0.03
Block	3.37	1740.44	294.38	$< .001$	***	0.1
Gender:Block	10.12	1740.44	0.77	0.659		$< .001$

Table 7

ANOVA of accuracy on active and inhibitory reward-seeking trials.

Effect	df_n	df_d	F	p	sig.	η^2
Gender	1	292	2.61	0.107		0.01
Trial Type	1	292	58.86	$< .001$	***	0.07
Gender:Trial Type	1	292	7.61	0.006	**	0.01

Table 8

Relationship between accuracy and mood disorder symptoms on active and inhibitory reward-seeking trials. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Trial Type	Comparison	df	t	p	sig.
Active reward-seeking	BDI~accuracy	306	-0.52	0.61	
	BAI~accuracy	306	-2.56	0.01	*
Inhibitory reward-seeking	BDI~accuracy	306	-0.54	0.59	
	BAI~accuracy	306	-0.83	0.41	

Table 9

Gender moderation of relationship between accuracy and mood disorder symptoms on active and inhibitory reward-seeking trials. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Trial Type	Comparison	df_n	df_d	t	p	sig.
Active reward-seeking	Gender moderation of BDI~accuracy	1	306	0.56	0.46	
	Gender moderation of BAI~accuracy	1	306	3.30	0.07	
Inhibitory reward-seeking	Gender moderation of BDI~accuracy	1	306	0.12	0.73	
	Gender moderation of BAI~accuracy	1	306	0.00	0.96	

Table 10

ANOVA of effort deployment on active reward-seeking trials by gender. Block = group of 28 trials on which an effortful response was required to obtain a reward.

Effect	df_n	df_d	F	p	sig.	η^2
Gender	3.00	379.00	2.03	0.11		0.01
Block	3.39	1283.92	387.02	$< .001$	***	0.16
Gender:Block	10.16	1283.92	0.43	0.936		$< .001$

Figure 1

A diagram of the active and inhibitory avoidance and reward-seeking tasks. In the avoidance task (A), after an inter-stimulus interval (ISI) with a fixation cross onscreen, participants were presented with a cue associated with active or inhibitory avoidance. For the active avoidance cue, participants had to respond with repeated spacebar presses to avoid hearing an aversive sound. For the inhibitory avoidance cue, participants had to withhold responding to avoid hearing an aversive sound. In the reward-seeking task (B), after the ISI, participants were presented with a cue associated with active or inhibitory reward-seeking. For the active reward-seeking cue, participants had to respond with repeated spacebar presses to obtain points towards a monetary reward. For the inhibitory reward-seeking cue, participants had to withhold responding to obtain points towards a monetary reward. ISI = Inter-stimulus interval.

Figure 2

(A) Distribution of anxiety (BAI) and depressive symptom (BDI) proportion scores (score divided by total possible score) by gender and sample. (B) Accuracy by gender on active and inhibitory avoidance, and moderation of the relationship between accuracy and anxiety (BAI) scores by gender. Dots represent average accuracy on each trial type by participant. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Figure 3

(A) Moderation of the relationship between accuracy and anxiety (BAI) proportion scores by gender in active and inhibitory avoidance. Gender significantly moderated the relationship between anxiety symptoms (BAI proportion scores) and inhibitory avoidance accuracy. (B) Accuracy by gender on active and inhibitory avoidance. BAI = Beck Anxiety Inventory. Proportion scores are scores divided by total possible score.

Figure 4

Interactive regression analysis output for active and inhibitory avoidance. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory. Numbers on right indicate coefficients for regression lines at each level of BDI score (centred 1 standard deviation below the mean, at the mean, and 1 standard deviation above the mean).

Figure 5

(A) Distribution of anxiety (BAI) and depressive symptom (BDI) scores by gender and sample.
(B) Accuracy by gender on active and inhibitory reward-seeking, and moderation of the relationship
between accuracy and anxiety (BAI) scores by gender. Proportion scores are scores divided by total
possible score. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Figure 6

(A) Moderation of the relationship between accuracy and anxiety (BAI) proportion scores by gender on active and inhibitory reward-seeking. Anxiety symptoms (BAI proportion scores) were significantly associated with active reward-seeking accuracy. (B) Accuracy by gender on active and inhibitory reward-seeking. BAI = Beck Anxiety Inventory. Proportion scores are scores divided by total possible score.

Figure 7

Interactive regression analysis output for active and inhibitory reward-seeking. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory. Numbers on right indicate coefficients for regression lines at each level of BDI score (centred 1 standard deviation below the mean, at the mean, and 1 standard deviation above the mean).

Figure 8

Effort deployment across active avoidance (A) and reward-seeking (B) trials relative to the criterion effort deployment required to avoid an aversive outcome or obtain reward (black line). Each point represents one trial. N values represent number of participants of each gender who completed the task during that block of trials.

Appendix A

Supplementary material

Effort and audio calibrations

After an introduction screen, participants completed an effort calibration to control for differences in baseline effort ability and keyboard sensitivity. They were instructed to press the spacebar on their computer as many times as possible within a five-second period when a thermometer appeared on screen. Each time they pressed the spacebar, the thermometer would increase in height in order to incentivize participants to press the spacebar as many times as possible. Afterwards, participants repeated this effort calibration. This second calibration was identical to the first except that the thermometer would increase by only half the amount per press that it did for the first calibration, in order to incentivize participants to press more times during the second calibration and thereby better capture the participant's maximum effort capability.

Following the effort calibration, participants completed an audio calibration to control for differences in audio cards and speakers. Here, participants were presented with a series of three one-second 2400 Hz sine tones - spaced by one second - at volumes of -50 dB, -30 dB, and -10 dB from maximum. After listening to these tones, participants were asked whether the first tone was barely heard and the final tone was aversive but not painful (Neumann and Waters, 2006). If this was not the case, participants were asked to adjust the volume on their computer and play the three tones again, repeating the process until the sound met these criteria - equalizing the experience of the sounds across participants. This computer volume was then used for the rest of the task.

Appendix B

Supplementary figures

Figure B1

Sensitivity (d′) and criterion (β) in active and inhibitory avoidance compared to depressive and anxiety symptoms. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

883   Neumann DL, Waters AM (2006) The use of an unpleasant sound as an unconditional

884   stimulus in a human aversive pavlovian conditioning procedure. Biological Psychology

885   73:175–185.

Figure B2

Sensitivity (d′) and criterion (β) in active and inhibitory reward-seeking compared to depressive and anxiety symptoms. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Figure B3

Moderation of the relationship between accuracy and anxiety (BAI) and depressive symptom (BDI) scores by gender in avoidance. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Figure B4

Moderation of the relationship between accuracy and anxiety (BAI) and depressive symptom (BDI) scores by gender in reward-seeking. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Time

A. Avoidance

B. Reward-seeking

ISI
(2000 ms,
jittered 1200 ms)

▲ = 1 key press