Gender impacts the relationship between mood disorder symptoms and effortful avoidance performance

Brandon J. Forys[1], Ryan J. Tomm[1], Dayana Stamboliyska[1], Alex R. Terpstra[1], Luke Clark[1,2], Trisha Chakrabarty[2,3], Stan B. Floresco[1,2], and & Rebecca M. Todd[1,2]

[1] Department of Psychology

University of British Columbia

Vancouver

BC

Canada

[2] Djavad Mowafaghian Centre for Brain Health

University of British Columbia

Vancouver

BC

Canada

[3] Department of Psychiatry

University of British Columbia

Vancouver

BC

Canada

Author Note

The authors declare no competing financial interests. We would like to acknowledge Alan Kingstone for his instrumental feedback on this project, as well as the contributions of Aanandi Sidarth to designing visual stimuli and Ian Daly to designing sounds stimuli for these studies. This research was supported by a Natural Sciences and Engineering Research Council of Canada (NSERC) grant (#F19-05182) to R. M. Todd and the UBC Djavad Mowafaghian Centre for Brain Health Innovation Fund Kickstart Research Grant (#F19-05932), as well as an NSERC Canada Graduate Scholarship - Master's (CGS M) award to B.J. Forys and a Michael Smith Foundation for Health Research Scholar Award to R.M. Todd.

The authors made the following contributions. Brandon J. Forys: Conceptualization, Investigation, Methodology, Writing - Original Draft Preparation, Writing - Review & Editing; Ryan J. Tomm: Conceptualization, Methodology, Writing - Review & Editing; Dayana Stamboliyska: Conceptualization, Investigation, Methodology, Writing - Review & Editing; Alex R. Terpstra: Conceptualization, Writing - Review & Editing; Luke Clark: Conceptualization, Funding acquisition, Writing - Review & Editing; Trisha Chakrabarty: Conceptualization, Funding acquisition, Writing - Review & Editing; Stan B. Floresco: Conceptualization, Funding acquisition, Writing - Review & Editing; Rebecca M. Todd: Conceptualization, Funding acquisition, Methodology, Project Administration, Resources, Supervision, Writing - Review & Editing.

Correspondence concerning this article should be addressed to Brandon J. Forys, 2136 West Mall, Vancouver, BC Canada V6T 1Z4. E-mail: brandon.forys@psych.ubc.ca

Abstract

We must often decide how much effort to exert or withhold to avoid undesirable outcomes or obtain rewards. In depression and anxiety, levels of avoidance can be excessive and reward-seeking may be reduced. Yet outstanding questions remain about the links between motivated action/inhibition and anxiety and depression levels, and whether they differ between men and women. Here we examined the relationship between anxiety and depression scores, and performance on effortful active and inhibitory avoidance (Study 1) and reward seeking (Study 2) in humans. Undergraduates and paid online workers (N_{Avoid} = 545, N_{Reward} = 310; N_{Female} = 368, N_{Male} = 450, M_{Age} = 22.58, $Range_{Age}$ = 17-62) were assessed on the Beck Depression Inventory II (BDI) and the Beck Anxiety Inventory (BAI) and performed an instructed online avoidance or reward-seeking task. Participants had to make multiple presses on active trials and withhold presses on inhibitory trials to avoid an unpleasant sound (Study 1) or obtain points towards a monetary reward (Study 2). Overall, men deployed more effort than women in both avoidance and reward-seeking, and anxiety scores were negatively associated with active reward-seeking performance based on sensitivity scores. Gender interacted with anxiety scores and inhibitory avoidance performance, such that women with higher anxiety showed better avoidance performance. Our results illuminate effects of gender in the relationship between anxiety and depression levels and the motivation to actively and effortfully respond to obtain positive and avoid negative outcomes.

Keywords: avoidance, reward, effort, depression, anxiety

Word count: 8431

Gender impacts the relationship between mood disorder symptoms and effortful avoidance performance

Significance statement

We must often take or withhold effortful action to avoid unpleasant outcomes or obtain rewards. Depression and anxiety can impact these behaviours' effectiveness, but the roles of avoidance in depression and reward-seeking in anxiety are not fully understood. Gender differences in avoidance and reward-seeking have also not been examined. We present a task in which community participants with a range of anxiety and depression levels made or withheld button presses to avoid hearing an unpleasant sound or obtain a reward. Men deployed more effort than women in avoidance, and women with higher anxiety scores had lower avoidance performance than men. We illuminate gender differences in how depressive and anxiety scores impact our ability to avoid threats and obtain rewards.

Introduction

Avoidance and reward-seeking behaviours

Living organisms are motivated to avoid potential threats or to acquire rewards respectively. Often achieving these goals requires action, but it can also require refraining from action. For example, we may take action to remove a threat's potential harm through active avoidance, or we may decide that withholding action is the best way to let the threat pass by, as in inhibitory avoidance (Krypotos, Effting, Kindt, and Beckers, 2015; LeDoux, Moscarello, Sears, and Campese, 2017). Alternatively, in a situation that offers the possibility of reward, we may take action to approach the reward through active reward-seeking or, instead, inhibit pre-potent reward seeking to wait for a larger reward (Capuzzo and Floresco, 2020; Cools, 2008). Research suggests that the expression of similar behavioral actions (including inhibition) is dependent on the motivational context (aversive

vs. appetitive), which influences the likelihood of selecting a specific action in a specific motivational context (Wang and Delgado, 2021). However, in neuropsychiatric research, depressive disorders are often studied with regard to reward-seeking contexts, and anxiety disorders with regard to avoidance contexts, with little emphasis on the other motivational context. Symptoms of anxiety and depression have been associated with avoidance, typically operationalized via active avoidance and via questionnaires, as threats are overestimated (Bishop and Gagne, 2018; Browning, Behrens, Jocham, O'Reilly, and Bishop, 2015; Cléry-Melin, Schmidt, Lafargue, Baup, Fossati, and Pessiglione, 2011; Mkrtchian, Aylward, Dayan, Roiser, and Robinson, 2017; Ottenbreit, Dobson, and Quigley, 2014). In depression, reward-seeking may also be impaired due to a lack of motivation to obtain rewards (Alloy, Olino, Freed, and Nusslock, 2016; Bishop and Gagne, 2018). Past research has established the importance of avoidance and reward-seeking behaviours in helping us navigate our environment and stay safe (Krypotos, Effting, Kindt, and Beckers, 2015; LeDoux, Moscarello, Sears, and Campese, 2017). However, active vs. inhibitory subtypes of these behaviours have not typically been distinguished – especially through objective measures of observable behavior.

Gender - as a culturally defined construct - may also be an important variable in this relationship. For example, gender differences have been identified in the presentation and incidence of mood and anxiety disorders, such that women have higher rates of depression and present more often with depression than men (Altemus, Sarvaiya, and Neill Epperson, 2014; Kessler, 2006; Parker and Brotchie, 2010) and have rates of anxiety disorders that are twice as high as those of men (McLean, Asnaani, Litz, and Hofmann, 2011; Pittig, Treanor, LeBeau, and Craske, 2018). However, we do not know how these gender differences manifest themselves in avoidance or reward-seeking behaviours. Although mood and anxiety disorders are often comorbid, they also manifest with distinct symptoms and courses that would require distinct strategies to treat in a clinical context (Goldstein-Piekarski, Williams, and Humphreys, 2016; McLean, Asnaani, Litz, and

Hofmann, 2011). In the present study, we ask how indices of anxiety and depression levels impact active vs. inhibitory avoidance and reward-seeking behaviours in a community population of young adults with a wide range of depressive and anxiety scores ranging from minimal to severe.

The role of mood disorder symptoms and gender differences in avoidance and reward-seeking

It has been proposed that mood and anxiety disorder symptoms shift the perceived value and costs of avoidance and reward-seeking in sub-optimal ways. The Altered Computations Underlying Decision Making (ACDM) framework (Bishop and Gagne, 2018) proposes that anxiety is linked to underestimation of the effort cost in avoiding an aversive outcome and that depression is linked to overestimation of the effort cost in obtaining a reward. These effort costs interact with the perceived value of avoidance or reward-seeking to inform one's decision on whether or not to engage in the behaviour. Past experimental work has also identified impairments in physical effort deployment for reward in populations with depression (Pessiglione, Vinckier, Bouret, Daunizeau, and Le Bouc, 2018; Treadway, Buckholtz, Schwartzman, Lambert, and Zald, 2009; see Culbreth, Moran, and Barch, 2018 for a review) and anxiety (Wang and Delgado, 2021). However, work linking mood and anxiety disorders to impairments in adaptive avoidance and reward-seeking often focuses on these avoidance and reward-seeking behaviours as unitary processes. As such, we still do not know how shifts in perceived effort costs linked to mood and anxiety disorders manifest themselves in active or inhibitory avoidance or reward-seeking.

To better understand the degree to which depressive and anxiety scores contribute to active and inhibitory forms of avoidance or reward-seeking a rigorous assessment of effort deployment in these behaviors is needed. People with Major Depressive Disorder (MDD) show a reduction in selecting high-effort, high-reward options on effort-based decision making tasks. This behaviour is potentially symptomatic of a larger-scale motivational

deficit (Pessiglione, Vinckier, Bouret, Daunizeau, and Le Bouc, 2018; Treadway, Bossaller,

Shelton, and Zald, 2012; Treadway, Buckholtz, Schwartzman, Lambert, and Zald, 2009). If

maladaptive effort deployment is a primary characteristic of mood and anxiety disorders,

then we might expect active avoidance and reward-seeking to be impaired more than

inhibitory forms of these behaviours overall (Culbreth, Moran, and Barch, 2018). Anxiety,

especially when co-occurring with high levels of depression, has also been shown to impair

our sensitivity to rewards (Auerbach, Pagliaccio, Hubbard, Frosch, Kremens, Cosby, Jones,

Siless, Lo, Henin, Hofmann, Gabrieli, Yendiki, Whitfield-Gabrieli, and Pizzagalli, 2022;

Dillon, Rosso, Pechtel, Killgore, Rauch, and Pizzagalli, 2014); however, whether anxiety's

impact on reward-seeking differs for active or inhibitory behaviours is not yet clear.

Additionally, individual differences in the presentation and severity of mood and

anxiety disorders - beyond the mere presence or absence of the disorder - may manifest

with different patterns of active vs. inhibitory behaviours depending on the motivational

context. Among these patterns, gender differences are especially prominent. Women

generally present with higher levels of depression (Parker and Brotchie, 2010) and

experience depression comorbid with anxiety more often than men (Kessler, 2006; McLean,

Asnaani, Litz, and Hofmann, 2011; Ottenbreit, Dobson, and Quigley, 2014). Thus, the

impact of mood and anxiety disorders on our ability to avoid aversive outcomes and seek

out rewarding outcomes may be linked to gender differences that affect the motivational

deficits these disorders present. If gender differences - looking across a full range of

depressive and anxiety scores captured on self-report scales (Beck, Steer, and Brown, 1996)

- predict differences in performance in a gender-dependent manner, then our study may

help elucidate how gender differences in depressive and anxiety scores translate to changes

in real-life behaviour.

In order to bring our understanding of mood disorder symptoms into a framework

that acknowledges differences in active vs. inhibitory avoidance and reward-seeking

behaviours, we must consider both anxiety and depression in a framework that directly investigates their impact on these behaviours, and how depressive and anxiety symptoms might interact to impair effective avoidance and reward-seeking. While the relationships between anxiety and avoidance (Bishop and Gagne, 2018; Levita, Hoskin, and Champi, 2012; Norbury, Robbins, and Seymour, 2018), and depression and reward-seeking (Alloy, Olino, Freed, and Nusslock, 2016; Rizvi, Pizzagalli, Sproule, and Kennedy, 2016; Treadway, Bossaller, Shelton, and Zald, 2012) are well established, those between anxiety and reward-seeking, as well as depression and avoidance, have yet to be fully characterized.

An effortful avoidance and reward-seeking study

Despite established gender differences in the prevalence and presentation of mood disorder symptoms (Kessler, 2006; Parker and Brotchie, 2010; Thompson and Bland, 2018), it is not known how the relationship between mood and anxiety disorder symptom levels, and avoidance and reward-seeking, differs by gender. Gender differences in motivational deficits may lead to unique patterns in active and inhibitory behaviours, but this has not been examined either. As such, in this work, we ask: 1) whether anxiety and depression symptoms predict accuracy and effort deployment in active/inhibitory avoidance vs. reward-seeking; and 2) whether there are gender differences in the relationship between mood disorder scores and accuracy. We predicted that anxiety scores would significantly predict participants' accuracy and the amount of effort they were able to deploy in avoidance behaviours (Bishop and Gagne, 2018) and that depression scores would significantly predict participants' accuracy and effort in reward-seeking behaviours (Bishop and Gagne, 2018). We also predicted that the relationship between mood disorder scores – especially depression scores – and task performance (accuracy and effort) would be differ by gender, such that women would have lower performance than men in the task given higher depression scores (Parker and Brotchie, 2010).

To address these questions, the present study examined both avoidance and

reward-seeking, each with two community online samples - undergraduates and online

workers - with a broad distribution of mood disorder scores. Both studies were

reverse-translated with modification from a series of rodent studies investigating deficits in

active and inhibitory avoidance and reward-seeking behaviours (Capuzzo and Floresco,

2020; Piantadosi, Yeates, and Floresco, 2018). Our studies are the first to combine

intermixed active and inhibitory avoidance (Levita, Hoskin, and Champi, 2012) or

reward-seeking trials with increasing effort requirements throughout the task, requiring

participants to switch between withholding physical effort on inhibitory trials and

deploying increasing amounts of effort on active trials in each task. This design allows us

to directly compare performance on active and inhibitory trials in the context of increasing

effort demands. Increasing effort demands may also pull out differences in selecting

between active vs. inhibitory strategies.

Materials and Methods

Participants

We powered each study to detect a moderate-sized main effect of $d = 0.15$ obtained

with a previous study of $N = 217$ participants using the `fabs` R package (Biesanz, 2020),

resulting in a target sample size of $N = 549$. Demographic information for all studies can

be found in Table 1. For each study, we collected data from two samples: an undergraduate

population and an online worker population. The study was approved by the University of

British Columbia Behavioural Research Ethics Board (BREB) under certificate H20-01388.

Study 1 (Avoidance)

We recruited undergraduate participants at the University of British Columbia to

participate online in our study. These participants received one percentage point towards

their grade in a psychology course of their choosing for completing the study. Of these

participants, $N = 311$ finished the study, of which $N = 39$ were excluded for not

completing the pre-task survey, having below 50% accuracy on active or inhibitory

avoidance trials, spending over 100 s on any given attention check, or not responding to all

Beck Anxiety Inventory (BAI) questions. As such, data from $N = 272$ participants was

used in the data analysis.

Additionally, we recruited paid online workers from around the world ($N = 310$) on

the Prolific online study platform (https://www.prolific.co/). These participants received

GBP £8.07 for completing the study. Of these participants, $N = 294$ finished the study, of

which $N = 22$ were excluded for not completing the pre-task survey, having below 50%

accuracy on active or inhibitory avoidance trials, spending over 100 s on any given

attention check, or not responding to all Beck Anxiety Inventory (BAI) questions. As such,

data from $N = 272$ participants was used in the data analysis.

Study 2 (Reward-seeking)

We recruited undergraduate participants at the University of British Columbia to

participate online in our study. These participants received one percentage point towards

their grade in a psychology course of their choosing and a CAD $5.00 gift card from

Starbucks for completing the study. Of these participants, $N = 83$ finished the study, of

which $N = 43$ were part of a separate task condition with visual appetitive stimuli that is

beyond the scope of this paper and $N = 4$ were excluded for not completing the pre-task

survey, having below 50% accuracy on active or inhibitory avoidance trials, spending over

100 s on any given attention check, or incorrectly responding to a pre-task attention check.

As such, data from $N = 36$ participants was used in the data analysis.

Additionally, we recruited paid online workers from around the world ($N = 309$) on

the Prolific online study platform. These participants received GBP £8.07 and a £2.69

bonus for completing the study. Of these participants, $N = 300$ finished the study, of which

$N = 26$ were excluded for not completing the pre-task survey, having below 50% accuracy

245 on active or inhibitory avoidance trials, spending over 100 s on any given attention check,

246 or incorrectly responding to a pre-task attention check. As such, data from $N = 274$

247 participants was used in the data analysis.

248   Overall, the excluded sample across both studies was 29.17% female and 70.83%

249 male, while the analyzed sample was 45.90% female and 54.10% male.

Materials and Methods

Stimulus presentation

252   We used PsychoPy 2020.1.2 (RRID: SCR_006571) via the Pavlovia online study

253 platform (Peirce, Gray, Simpson, MacAskill, Höchenberger, Sogo, Kastman, and Lindeløv,

254 2019). Participants completed the study online on their own computers; they were not

255 allowed to complete the study on mobile devices or tablets.

Stimuli

Cues indicating active or inhibitory trials were dark blue squares and circles with a

thin black border and were generated by PsychoPy 2020.1.2 (Peirce, Gray, Simpson,

MacAskill, Höchenberger, Sogo, Kastman, and Lindeløv, 2019) (Fig. 1); they subtended a

visual angle of about 11.5° x 11.5°. All stimuli were presented against a grey background

(RGB value [0,0,0] on a scale from -1 to 1). If participants responded incorrectly on any

trial in the avoidance studies, an aversive sound was played for 2000 ms. The aversive

sounds were randomly selected from a set of eight screeching and scraping sounds created

by our lab and ranked as highly aversive by four independent raters and in a pilot study.

Participants completed a series of questionnaires before beginning the main task.

These were the State-Trait Anxiety Inventory, form Y-2 (STAI Y-2) (Spielberger, 2008);

the Beck Depression Inventory II (BDI) (Beck, Steer, and Brown, 1996); the Beck Anxiety

Inventory (BAI) (Steer and Beck, 1997); the Behavioral Activation for Depression Scale

(BADS) (Kanter, Mulick, Busch, Berlin, and Martell, 2007); the Generalized Anxiety

Disorder Scale (GAD-7) (Spitzer, Kroenke, Williams, and Löwe, 2006); and the

Behavioural Inhibition Scale and Behavioural Activation Scale (BIS/BAS) (Carver and

White, 1994). In our data analysis, we looked at results from the BDI and BAI as these

clinically validated scales most directly capture participants' levels of current depressive

and anxiety scores. The BADS, GAD-7, and BIS/BAS capture specific behavioural facets

of depression and anxiety that are less relevant to understanding overall effects of mood

and anxiety disorders on avoidance and reward-seeking and were not analyzed in this study.

We used the BAI as our primary measure of anxiety scores as it is the most widely used

and validated among the anxiety scales we included (Fydrich, Dowdall, and Chambless,

1992) and as its structure parallels that of the BDI.

Procedure

Avoidance task

A graphical overview of the avoidance task is provided in Fig. 1A.

After an introduction screen, participants completed an effort calibration to control

for differences in baseline effort ability and keyboard sensitivity. They were instructed to

press the spacebar on their computer as many times as possible within a five-second period

when a thermometer appeared on screen. Each time they pressed the spacebar, the

thermometer would increase in height in order to incentivize participants to press the

spacebar as many times as possible. Afterwards, participants repeated this effort

calibration. This second calibration was identical to the first except that the thermometer

would increase by only half the amount per press that it did for the first calibration, in

order to incentivize participants to press more times during the second calibration and

thereby better capture the participant's maximum effort capability.

Following the effort calibration, participants completed an audio calibration to

control for differences in audio cards and speakers. Here, participants were presented with a series of three one-second 2400 Hz sine tones - spaced by one second - at volumes of -50 dB, -30 dB, and -10 dB from maximum. After listening to these tones, participants were asked whether the first tone was barely heard and the final tone was aversive but not painful (Neumann and Waters, 2006). If this was not the case, participants were asked to adjust the volume on their computer and play the three tones again, repeating the process until the sound met these criteria - equalizing the experience of the sounds across participants. This computer volume was then used for the rest of the task.

After calibrating their physical effort capability and the volume of the aversive stimuli in the task, participants read instructions indicating the shape to which they would have to respond with multiple spacebar presses as well as the shape to which they would have to withhold their response. They also heard an example of the aversive sound that would be played if they made an incorrect response during the task.

In order to gain exposure to the stimuli and task contingencies, participants completed a series of practice trials (Fig. 1A). This consisted of five trials in which participants had to make an active avoidance response - pressing the spacebar several times to avoid hearing an unpleasant sound; five trials in which participants had to make an inhibitory avoidance response - not pressing the spacebar to avoid hearing an aversive sound; and ten trials that intermixed these active and inhibitory trials.

On each trial, participants first viewed a grey screen with a white fixation cross for a mean duration of 2000 ms with a standard deviation (SD) of 1200 ms, jittered according to a normal distribution with these parameters on each trial. Participants then saw a visual cue - either a blue circle or a blue square - for 2000 ms. The cues used for active and inhibitory trials were pseudorandomly intermixed between participants. While this cue was on-screen, participants had to press the spacebar multiple times on active avoidance trials or withhold pressing on inhibitory avoidance trials. On active trials, the number of presses

required was set according to the average number of presses made during the two effort

calibration trials, such that participants who pressed fewer times during the calibration

would have to press fewer times to achieve criterion during the task. The initial criterion

was 5 presses given an average of 18 or fewer presses during calibration; a criterion of 6

presses given an average of 19-33 presses inclusive during calibration; and 7 presses given

an average of 34 or more presses during calibration.

If participants made an incorrect response (pressing an insufficient number of times

on active trials or pressing at all on inhibitory trials), participants heard an aversive sound

and saw a fixation cross for 2000 ms. This aversive sound was taken from a set of ten

sounds created by our lab and rated as highly aversive. All sounds were scraping sounds

that had unpleasant psychoacoustic properties shown to reliably induce aversive responses

(Neumann and Waters, 2006) at a variety of frequencies. If participants made a correct

decision (pressing a sufficient number of times on active trials or not pressing on inhibitory

trials), they saw a fixation cross surrounded by a white border that acted as a safety signal

on the edges of the screen for 500 ms.

After completing the practice trials and viewing a final screen reminding them of

the instructions, participants began the main task. This consisted of up to 168 active

avoidance trials and 72 inhibitory avoidance trials (70% active and 30% inhibitory),

pseudorandomized such that no more than 6 active trials or 3 inhibitory trials appeared in

a row. On the 15th trial and every 40 trials thereafter, an attention check appeared asking

participants to press a key corresponding to the letter they heard, to ensure that they were

attending to the task and able to hear auditory stimuli. Every 20 trials, the number of

button presses required on active trials increased by one press - this increased the effort

demands on active trials across the task. The task continued until the participant

responded correctly on half or less than half of the last 20 active trials - at this point, the

breakpoint was reached and the participant was thanked for completing the task.

Reward-seeking task

A graphical overview of the reward-seeking task is provided in Fig. 1B.

The design of the reward-seeking task was identical to that of the avoidance task, with the following exceptions. First, the practice blocks were based on criterion-based advancement in order to increase consistency with the design of other reward-seeking studies in our lab. Participants had to achieve at least 80% accuracy in each of the active, inhibitory, and intermixed reward-seeking trial blocks in order to advance; each block would repeat until they achieved each criterion. Second, if the participant made a correct decision during a trial, they would see a screen indicating that they had gained 5 points along with a sum of their points thus far; if the participant made an incorrect response during a trial, they would see a screen indicating that they had gained 0 points along with a sum of their points thus far. Both screens appeared for 1500 ms. Undergraduate participants received a CAD $5 gift card as a reward in addition to course credit for completing the task; online workers received a GBP £2.69 payment as a reward in addition to their payment for completing the task. Finally, as this task did not incorporate audio, no volume check or audio-based attention check was included.

Data analysis

All data was analyzed using R 4.1.1 "Kick Things" (R Development Core Team, 2011) through RStudio (Booth et al., 2018). On each behavioural task, we measured: 1) participants' sensitivity (d'), operationalized as the rate of correct active trials (hit rate) minus the rate of incorrect inhibitory trials (false alarm rate) and calculated using the `dprime` function in the `psycho` R package (Makowski, 2018) as the Z value of the hit rate minus the Z value of the false alarm rate; 2) effort on each trial type, operationalized as the number of presses made relative to criterion on each trial; 3) participants' depressive and anxiety scores, operationalized as their BDI (Beck, Steer, and Brown, 1996) and BAI (Steer and Beck, 1997) scores respectively; and 4) breakpoint, operationalized as the trial number

on which the participant responded incorrectly on half or less than half of the last 20 active trials. Breakpoint captures the point at which the effort demands of the task are no longer attainable for the participant, which is impaired in people with depression (Hershenberg, Satterthwaite, Daldal, Katchmar, Moore, Kable, and Wolf, 2016). On inhibitory trials, effort reflects the total number of presses made, to capture mistakes in which responses were still made on inhibitory trials. In order to capture all task parameters parsimoniously in our analyses, we analyzed sensitivity for avoidance and reward-seeking through four linear models in which either anxiety scores (BAI scores) or depression scores (BDI scores), together with gender and sample (university undergraduates vs. online workers), were included as predictors. We also analyzed effort for avoidance and reward-seeking through four multi-level models (using the `lmerTest` R package (Kuznetsova, Brockhoff, and Christensen, 2017)) in which either anxiety scores (BAI scores) or depression scores (BDI scores), together with gender, sample, and block of 28 active trials, were included as fixed effects and participants were a random effect. Lastly, we analyzed breakpoint for avoidance and reward-seeking using two linear models in which either anxiety scores (BAI scores) or depression scores (BDI scores), together with gender and sample (university undergraduates vs. online workers), were included as predictors. Linear models were used for sensitivity and breakpoint as these factors did not differ within participants, unlike effort - which differed between active and inhibitory trials, making a multi-level model appropriate. All analyses were Bonferroni corrected for multiple comparisons. All confidence intervals were based on 1000 bootstrap replications using the `confintr` R package (Mayer, 2022). All BDI and BAI scores were divided by the maximum number of points possible on each scale to obtain proportion scores. This was necessary because of the exclusion from our surveys of a BDI question relating to suicidality as the inclusion of such a question was not covered in our ethics, and because of the omission from our surveys of a BAI question that was used as an attention check – rendering raw BDI and BAI scores not comparable to those of other studies.

<center>**Results**</center>

Demographics

Participant's reported gender and sex heavily overlapped. Of those reporting their gender as female, there was a 97.53% overlap with reported sex in women and 97.67% overlap in men on the avoidance tasks. There was a 98.84% overlap with reported sex women and 97.80% overlap in men on the reward-seeking task. For this reason, the following results are expressed in terms of gender only. Women reported higher levels of depressive scores ($t(720.46) = $ -3.83, $p < .001$, $d = 0.27$) and anxiety scores ($t(703.99) = $ -4.68, $p < .001$, $d = 0.34$) than men across samples (Table 2, Figs. 2, 5). Across both studies, 20.98% of women and 15.11% of men were on medication for depression, and 19.35% of women and 14.89% of men were on medication for anxiety. For participants on these medications, BAI and BDI scores reflect their anxiety and depression scores in a medicated state and participants' medication status was not included as a statistical control in our analyses.

Avoidance task

To account for participants' bias to engage in active relative to inhibitory avoidance in general, we first calculated sensitivity (d'). Sensitivity reflects participants' ability to correctly distinguish between active and inhibitory trials and deploy the required amount of effort on active trials only, while withholding effort on inhibitory trials. We additionally present results of active and inhibitory avoidance accuracy analyses at the link to the OSF repository in the data and code availability section (https://osf.io/2rd3f/). As all variance in sensitivity was between subjects, we ran a linear model analysis (Table 3) to evaluate whether d' could be predicted from anxiety scores (BAI scores), gender, and sample in avoidance. Gender in interaction with anxiety scores significantly predicted sensitivity in the avoidance task such that women had lower performance with higher levels of anxiety (Fig. 3A). There was also a BAI x Sample interaction such that participants with high

anxiety in the global sample had lower performance than those with high anxiety in the

undergraduate sample. We ran an additional linear model analysis (Table 4) to evaluate

whether d' could be predicted from depression scores (BDI scores), gender, and sample in

avoidance (Fig. 3B). Depression scores and sample significantly predicted d', as did

interactions between depression scores and gender, and between depression scores and

sample - such that there was a significant interaction with women having lower sensitivity

given higher depression scores. As with anxiety, women had lower performance with higher

levels of depression. Women in the online worker sample had lower performance than those

in the undergraduate sample given higher levels of depression.

Additionally, we explored the extent to which the amount of effort that participants

exerted to avoid aversive outcomes changed across the avoidance task. As effort

deployment could differ both between and within subjects, we conducted a multi-level

model analysis (Table 5) to evaluate whether effort deployment could be predicted from

anxiety (BAI) scores, gender, sample, and block (28 active trials) in avoidance. This

analysis revealed that participants deployed increasing amounts of effort during the task to

meet increasing effort requirements (Fig. 4A). There was also an interaction between

gender and block qualified by a 3-way interaction between gender, BAI score, and block,

indicating that the decrease in effort over time was associated with increased anxiety

primarily in women (Fig. 4B). We ran an additional multi-level model analysis (Table 6) to

evaluate whether effort could be predicted from depression (BDI) scores, gender, sample,

and block in avoidance. Changes in effort across blocks during the task interacted with

participants' BDI scores and with gender, such that women with higher levels of depression

deployed less effort relative to criterion in active avoidance.

Last, we examined whether breakpoint could be predicted from anxiety scores (BAI

scores) or depression scores (BDI scores), gender, and sample in avoidance using two linear

models (Tables 7, 8). None of these factors significantly predicted breakpoint in avoidance.

Reward-seeking task

To evaluate participants' bias to engage in active relative to inhibitory reward-seeking, we again calculated sensitivity (d'). We ran a linear model analysis (Table 9) to evaluate whether d' could be predicted from anxiety scores (BAI scores), gender, and sample in reward-seeking. Only gender significantly predicted sensitivity in the reward-seeking task (Fig. 6). We ran an additional linear model analysis (Table 10) to evaluate whether d' could be predicted from depression scores (BDI scores), gender, and sample in avoidance. None of these factors predicted sensitivity in reward-seeking.

Additionally, we explored the extent to which the amount of effort that participants deployed to obtain reward changed across the reward-seeking task. We ran a multi-level model analysis (Table 11) to evaluate whether effort deployment could be predicted from anxiety scores (BAI scores), gender, and sample in reward-seeking. This analysis revealed that effort decreased relative to criterion as the reward-seeking task progressed (Fig. 7A). There were also effects of gender, with men deploying overall more effort than women, and anxiety, such that those with higher levels of anxiety were less able to deploy effort relative to criterion. These were qualified by an interaction between gender and anxiety such that women increased effort and men decreased it with higher levels of anxiety (Fig. 7B). There were also a number of differences in effort deployment between samples, which interacted with a number of other predictors. We ran an additional multi-level model analysis (Table 12) to evaluate whether effort could be predicted from depression scores (BDI scores), gender, sample, and block in reward-seeking. Only task block and gender predicted effort when depression scores were included as a predictor, such that women with higher depression scores deployed less effort relative to criterion in active reward-seeking.

Last, we examined whether breakpoint could be predicted from anxiety scores (BAI scores) or depression scores (BDI scores), gender, and sample in reward-seeking using two linear models (Tables 13, 14). None of these factors significantly predicted breakpoint in

477 reward-seeking.

Discussion

Summary

In the present study, we investigated effects of gender and anxiety and depression levels in active and inhibitory avoidance and reward-seeking behaviours in a community population. Compared to men, women showed overall higher levels of self-reported depression and anxiety. Gender differences in task performance were in opposite directions depending on whether the task demanded avoidance or reward-seeking. Women showed lower sensitivity (d'), a measure of ability to correctly respond to active *and* inhibitory task demands, when avoiding an aversive outcome than men - but higher sensitivity when seeking reward. During active and inhibitory *avoidance*, gender interacted with anxiety level such that, in women, higher anxiety scores predicted lower sensitivity. There were also gender differences in effort deployment. Overall, throughout active *reward-seeking* trials, men made more effortful responses than women. Yet this finding was qualified by an interaction between gender and anxiety (over time) in both avoidance *and* reward-seeking, such that higher effort levels were associated with anxiety levels for women in both tasks, particularly early in the task. It should be noted that, in effortful reward seeking, effort deployment above criterion differed by sample such that online workers deployed more effort across the task and with increasing anxiety scores than undergraduate participants. Our findings illuminate gender differences in performance when both active and inhibitory responses are required for avoidance and reward-seeking in a context where effort is required to obtain the desired outcome. They also point to important boundary conditions for correlational effects that may vary between populations.

Interpretation of results

Effects of gender and anxiety on avoidance task performance

We observed that, in avoidance, gender differences in performance (sensitivity; d') where men had higher performance than women were moderated by anxiety, where women with higher anxiety had lower performance in the avoidance task. This finding is consistent with previous findings that avoidance behaviours are influenced by mood and anxiety levels (Pittig, Treanor, LeBeau, and Craske, 2018) which may partly explain overall gender differences in this task because of baseline gender differences in the prevalence of depression and anxiety (Parker and Brotchie, 2010). In particular, the observed gender differences in selecting the correct response on active vs. inhibitory trials (sensitivity; d') when avoiding an unpleasant outcome could reflect more general gender differences in stress tolerance. In our study, women reported overall higher anxiety scores than men. Parker and Brotchie (2010) argued that women have a higher predisposition (diathesis) to stress than men and, in a context where an aversive outcome must be avoided, people with higher levels of anxiety scores may have a stronger impulse to act to avoid an aversive outcome (Bishop and Gagne, 2018). Thus, inhibiting a response may be especially difficult given a combination of high anxiety scores and a decreased tolerance for stress in women compared to men. Such a gender difference in diathesis could drive a reduced ability to inhibit effort when needed - an impairment in shifting from an active to an inhibitory strategy given higher levels of anxiety scores (Gustavson, Altamirano, Johnson, Whisman, and Miyake, 2017). Response inhibition can also require cognitive effort. People may have differing tendencies to make physically effortful responses to avoid aversive outcomes or obtain rewards, and to maintain an awareness that a cue is associated with withholding a response. Mood disorders have been associated with undervaluation of the reward of a cognitively effortful outcome (Grahek, Musslick, and Shenhav, 2020). Most of these findings have been associated with depression and reward-seeking. However, the reduced sensitivity with higher BAI scores observed in women in our avoidance task could reflect not only reduced motivation for physical effort on active trials with higher levels of anxiety but also reduced

motivation to deploy cognitive effort to inhibit one's response on inhibitory trials.

Gender differences in avoidance were also moderated by depression scores, such that

women with higher depression scores performed worse on the avoidance task. Effects of

depression scores, as measured by BDI scores, on task performance mostly reflected effects

of anxiety. This is consistent with past findings that negative effects of anxiety and

depression on motivated behaviours like avoidance and reward-seeking are often similar

(Ottenbreit, Dobson, and Quigley, 2014), emphasizing the importance of a transdiagnostic

approach when evaluating the impact of anxiety and depression on these behaviours

(Culbreth, Moran, and Barch, 2018). It may also reflect the extent to which the BAI and

BDI measure overlapping constructs, as reflected by the high correlation between BAI and

BDI scores we observed ($t(852) = 23.86$, $p = < .001$, $r = 0.63$).

Effects of gender and anxiety on effort deployment in reward-seeking and
539
avoidance

In reward-seeking, we observed an opposing relationship between gender and effort

deployment to that in avoidance. Women deployed more effort relative to criterion at

higher anxiety scores than men, while men continued to deploy less effort with higher

anxiety scores. The diathesis effects that may impair effective effort deployment in

avoidance may not be present in reward-seeking, as effects of an error are likely to be less

stressful to participants. Therefore, the observed effort impairments in women may be

valence-specific.

On active reward-seeking trials, men deployed more effort relative to criterion across

the task than women. This could be caused by women having smaller wrists with which to

generate physical force than men (Morse, Jung, Bashford, and Hallbeck, 2006), as well as

men having - on average - higher levels of testosterone levels compared to women - which is

associated with increased physical effort (Losecaat Vermeer, Riečanský, and Eisenegger,

2016) and risk tolerance (Cooper, Goings, Kim, and Wood, 2014). This initial difference in effort deployment capability is reflected in the finding that men pressed significantly more than women in the pre-task effort calibration across all tasks ($t(814.99) = 7.02$, $p = <$ .001). Although our tasks did not have competitive elements, participants may still have completed the task with an eye towards maximizing performance. Since deploying more effort in the task would increase one's chance of staying above criterion, this increased effort deployment in men could explain the increased active trial accuracy for men across the avoidance and reward-seeking tasks. It is important to qualify that gender has a significant cultural component, and cultural factors could also play a role in gender differences in effort deployment - perhaps via effects of a lower tolerance for stress on effort deployment (Parker and Brotchie, 2010). Importantly, in both tasks gender interacted with anxiety levels and, in the avoidance task, gender differences in effort was qualified by a relationship with anxiety early in the task, with women higher in anxiety scores deploying higher effort levels early on. Thus any gender differences in effort are complex, and vary with anxiety levels, motivational context, and likely other boundary conditions as well.

It should be noted that the relationship between higher anxiety scores and reduced avoidance sensitivity, while qualified by gender and sample, differs from previous predictions of improved avoidance in anxiety, such as those of Bishop and Gagne (2018). Bishop and Gagne framed this relationship in terms of active and not inhibitory avoidance, as they predicted that underestimations of effort cost would drive excessive avoidance behaviours. Anxiety scores may be associated with impairments to inhibitory avoidance precisely because of this bias towards action given the possibility of aversive outcomes, an effect that could be driven by a perceived lack of control over outcomes in the task (Wang and Delgado, 2021). Additionally, we did not observe a relationship between depression scores and accuracy or effort deployment in reward-seeking, as has previously been observed (Bishop and Gagne, 2018). As depression scores did not influence effort deployment, we can speculate that, in this task, the effort demands of the task did not

deter those with higher depression scores from working for a reward.

Overall, participants deployed less effort in avoidance compared to reward-seeking; this could be a function of differences in motivation to engage in avoidance or reward-seeking. Motivation to complete the tasks can be driven in part by participants' valuations of task-relevant stimuli (Bishop and Gagne, 2018). A major difference between our tasks arises in the outcome of an incorrect response. In avoidance, an incorrect response is associated with an aversive sound; in reward-seeking, it is associated with not receiving points. Although the salience of an aversive sound may suggest that it is more motivating and would therefore be associated with increased accuracy, hearing it may also be more demotivating - especially for participants with mood disorder symptoms. Hearing the aversive sound repeatedly could be a salient indicator of a lack of control over task outcomes (Wang and Delgado, 2021).

Limitations

There are some limitations to our interpretation of our findings. First, since the dichotomy of the task demands is between effortful active trials and inhibitory trials that require no effort, we cannot compare the effects of high vs. low effort demands on inhibitory avoidance or reward-seeking behaviours. As such, our interpretation of the relationship between effort deployment and mood disorder symptoms only extends to active trials. Accuracy in the task was likely tied to participants' effort capabilities, as increased effort deployment was required throughout the task on active trials to meet the criterion level of effort and make the correct response on the trial. However, we calibrated the criterion to participants' effort ability and considered performance on inhibitory as well as active trials to reduce the reliance of task outcomes on individual differences in effort deployment. Additionally, as the proportion of active trials was greater than that of inhibitory trials, participants may have become increasingly fatigued on the majority of trials in the task. This fatigue from effort deployment, combined with boredom (from the

task being repetitive) could be difficult to disentangle from other shifts in motivation to deploy effort throughout the task (e.g. those related to the value of avoidance or reward-seeking). However, as fatigue is likely to arise in most physically effortful tasks, our tasks still reflect real-world physical effort demands. Furthermore, as this study took place online, the study had to use repeated keyboard presses instead of other, more continuous or better-controlled measures of physical effort such as a grip squeeze (Aridan, Malecek, Poldrack, and Schonberg, 2019). However, repeated button presses have been validated as being physically effortful and have been used in in-person contexts (Gold, Strauss, Waltz, Robinson, Brown, and Frank, 2013).

We found differences between samples in sensitivity in avoidance when BAI or BDI scores were included as a predictor, and in effort in reward-seeking when BAI scores were included as a predictor. These sample differences only arose in the context of specific predictors such as gender. Nevertheless, they may suggest a differing relationship between avoidance performance and depression scores, and reward-seeking effort and anxiety scores, among the younger, female-skewed online undergraduates and older, male-skewed online workers. As online workers were paid in money for participation - as opposed to receiving course credit, as in the case of the undergraduates - motivations for completing the task may have differed between samples. The online workers were also likely a more heterogeneous sample in terms of the devices and contexts in which they completed the task. However, the presence of significant effects that do not interact with sample suggest that sample differences did not explain a large proportion of the variance in our findings.

Future work

Future studies could build on our findings by investigating how patterns of information about specific aspects of effortful avoidance and reward-seeking are instantiated in key brain regions. The posterior anterior cingulate cortex (pACC) and ventral striatum encode information about prospective gains given physical effort requirements (Aridan,

Malecek, Poldrack, and Schonberg, 2019). These regions - and their homologues in rodents

- have been shown to be differentially necessary for active vs. inhibitory avoidance

(Piantadosi, Yeates, and Floresco, 2018) and reward-seeking (Capuzzo and Floresco, 2020).

Investigating how these regions represent information on prospective threats and gains

relative to effort costs could illuminate how we weigh the benefits and costs of deploying

effort to obtain rewards and avoid aversive outcomes. Additionally, separating out different

factors contributing to effort deployment through a computational modelling approach

would be important to understand the individual contributions of various factors to

participants' performance. These factors could include action biases (Mkrtchian, Aylward,

Dayan, Roiser, and Robinson, 2017), perceived value of avoidance or reward (Bishop and

Gagne, 2018), or fatigue (Pessiglione, Vinckier, Bouret, Daunizeau, and Le Bouc, 2018).

Furthermore, it would be helpful to evaluate whether subscales of mood disorder symptoms

- potentially linked to subtypes such as anxious depression (Wurst, Schiele, Stonawski,

Weiß, Nitschke, Hommers, Domschke, Herrmann, Pauli, Deckert, and Menke, 2021) - pull

out factors that drive participants' behaviours in avoidance and reward seeking. This

analysis could further illuminate our observed gender differences - for example, to evaluate

whether reduced avoidance sensitivity in women given increased anxiety scores is reflective

of an anxious subtype of depression (Wurst, Schiele, Stonawski, Weiß, Nitschke, Hommers,

Domschke, Herrmann, Pauli, Deckert, and Menke, 2021).

Conclusion

Our studies address outstanding questions of whether a range of anxiety and

depression scores predict performance (sensitivity) and effort deployment in avoidance and

reward-seeking, and whether the relationship between performance and anxiety/depression

levels is impacted by gender. We elicit both active and inhibitory avoidance and

reward-seeking behaviours in a context that allows for direct comparisons between them,

instead of considering avoidance and reward-seeking behaviours as unitary wholes. We

highlight gender differences in each of these subtypes of avoidance and reward-seeking given varying levels of anxiety and depression scores, contextualizing past work on gender differences (Parker and Brotchie, 2010). In particular, we are the first to examine these proposed gender differences in an active and inhibitory avoidance and reward-seeking context. These findings could inform clinical interventions to address maladaptive deployment of avoidance and lack of motivation for reward-seeking, targeted by gender. Additionally, we link active avoidance and reward-seeking to motivation for physical effort deployment given varying levels of mood and anxiety disorder severity. As many tasks in life require physical effort deployment, understanding where it can be impaired is an important pursuit. Our findings underscore the importance of considering individual differences in the ways in which avoidance and reward-seeking can be impaired in life.

Data and code availability

The data and materials for all experiments, as well as the code used to generate this manuscript and conduct all analyses, are available at https://osf.io/2rd3f/.

References

Alloy LB, Olino T, Freed RD, Nusslock R (2016) Role of Reward Sensitivity and Processing in Major Depressive and Bipolar Spectrum Disorders. Behavior Therapy 47:600–621.

Altemus M, Sarvaiya N, Neill Epperson C (2014) Sex differences in anxiety and depression clinical perspectives. Frontiers in Neuroendocrinology 35:320–330.

Aridan N, Malecek NJ, Poldrack RA, Schonberg T (2019) Neural correlates of effort-based valuation with prospective choices. NeuroImage 185:446–454.

Auerbach RP, Pagliaccio D, Hubbard NA, Frosch I, Kremens R, Cosby E, Jones R, Siless V, Lo N, Henin A, Hofmann SG, Gabrieli JDE, Yendiki A, Whitfield-Gabrieli S, Pizzagalli DA (2022) Reward-Related Neural Circuitry in Depressed and Anxious Adolescents: A Human Connectome Project. Journal of the American Academy of Child & Adolescent Psychiatry 61:308–320.

Beck AT, Steer RA, Brown GK (1996) Manual for the beck depression inventory-II. San Antonio, TX: Psychological Corporation.

Biesanz J (2020) Fabs: Functions for applied behavioural sciences in r.

Bishop SJ, Gagne C (2018) Anxiety, depression, and decision making: A computational perspective. Annual Review of Neuroscience 41:371–388.

Booth DS et al. (2018) RStudio: Integrated development for r. Nature.

Browning M, Behrens TE, Jocham G, O'Reilly JX, Bishop SJ (2015) Anxious individuals have difficulty learning the causal statistics of aversive environments. Nature Neuroscience 18:590–596.

Capuzzo G, Floresco SB (2020) Prelimbic and infralimbic prefrontal regulation of active and inhibitory avoidance and reward-seeking. Journal of Neuroscience 40:4773–4787.

Carver CS, White TL (1994) Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: The BIS/BAS scales.

Journal of Personality and Social Psychology.

Cléry-Melin ML, Schmidt L, Lafargue G, Baup N, Fossati P, Pessiglione M (2011)

Why don't you try harder? An investigation of effort production in major

depression. PLoS ONE 6.

Cools R (2008) Role of Dopamine in the Motivational and Cognitive Control of

Behavior. The Neuroscientist 14:381–395.

Cooper SE, Goings SP, Kim JY, Wood RI (2014) Testosterone enhances risk

tolerance without altering motor impulsivity in male rats.

Psychoneuroendocrinology 40:201–212.

Culbreth AJ, Moran EK, Barch DM (2018) Effort-cost decision-making in psychosis

and depression: Could a similar behavioral deficit arise from disparate

psychological and neural mechanisms? Psychological Medicine 48:889–904.

Dillon DG, Rosso IM, Pechtel P, Killgore WDS, Rauch SL, Pizzagalli DA (2014)

Peril and pleasure: An RDOC-inspired examination of threat responses and

reward processing in anxiety and depression. Depression and Anxiety

31:233–249.

Fydrich T, Dowdall D, Chambless DL (1992) Reliability and validity of the beck

anxiety inventory. Journal of Anxiety Disorders 6:55–61.

Gold JM, Strauss GP, Waltz JA, Robinson BM, Brown JK, Frank MJ (2013)

Negative symptoms of schizophrenia are associated with abnormal effort-cost

computations. Biological Psychiatry 74:130–136.

Goldstein-Piekarski AN, Williams LM, Humphreys K (2016) A trans-diagnostic

review of anxiety disorder comorbidity and the impact of multiple exclusion

criteria on studying clinical outcomes in anxiety disorders. Translational

Psychiatry 6:e847–e847.

Grahek I, Musslick S, Shenhav A (2020) A computational perspective on the roles of

affect in cognitive control. International Journal of Psychophysiology 151:25–34.

Gustavson DE, Altamirano LJ, Johnson DP, Whisman MA, Miyake A (2017) Is set

shifting really impaired in trait anxiety? Only when switching away from an

effortfully established task set. Emotion 17:88–101.

Hershenberg R, Satterthwaite TD, Daldal A, Katchmar N, Moore TM, Kable JW,

Wolf DH (2016) Diminished effort on a progressive ratio task in both unipolar

and bipolar depression. Journal of Affective Disorders 196:97–100.

Kanter JW, Mulick PS, Busch AM, Berlin KS, Martell CR (2007) The Behavioral

Activation for Depression Scale (BADS): Psychometric Properties and Factor

Structure. Journal of Psychopathology and Behavioral Assessment 29:191–202.

Kessler RC (2006) The epidemiology of depression among women. Women and

Depression: A Handbook for the Social, Behavioral, and Biomedical Sciences

74:22–38.

Krypotos AM, Effting M, Kindt M, Beckers T (2015) Avoidance learning: A review

of theoretical models and recent developments. Frontiers in Behavioral

Neuroscience 9.

Kuznetsova A, Brockhoff PB, Christensen RHB (2017) lmerTest package: Tests in

linear mixed effects models. Journal of Statistical Software 82:1–26.

LeDoux JE, Moscarello J, Sears R, Campese V (2017) The birth, death and

resurrection of avoidance: A reconceptualization of a troubled paradigm.

Molecular Psychiatry 22:24–36.

Levita L, Hoskin R, Champi S (2012) Avoidance of harm and anxiety: A role for the

nucleus accumbens. NeuroImage 62:189–198.

Losecaat Vermeer AB, Riečanský I, Eisenegger C (2016) Competition, testosterone,

and adult neurobehavioral plasticity, pp213–238. Elsevier.

Makowski D (2018) The psycho package: An efficient and publishing-oriented

workflow for psychological science. Journal of Open Source Software 3:470.

Mayer M (2022) Confintr: Confidence intervals.

McLean CP, Asnaani A, Litz BT, Hofmann SG (2011) Gender differences in anxiety disorders: Prevalence, course of illness, comorbidity and burden of illness. Journal of Psychiatric Research 45:1027–1035.

Mkrtchian A, Aylward J, Dayan P, Roiser JP, Robinson OJ (2017) Modeling avoidance in mood and anxiety disorders using reinforcement learning. Biological Psychiatry 82:532–539.

Morse JL, Jung M-C, Bashford GR, Hallbeck MS (2006) Maximal dynamic grip force and wrist torque: The effects of gender, exertion direction, angular velocity, and wrist angle. Applied Ergonomics 37:737–742.

Neumann DL, Waters AM (2006) The use of an unpleasant sound as an unconditional stimulus in a human aversive pavlovian conditioning procedure. Biological Psychology 73:175–185.

Norbury A, Robbins TW, Seymour B (2018) Value generalization in human avoidance learning. eLife 7.

Ottenbreit ND, Dobson KS, Quigley L (2014) An examination of avoidance in major depression in comparison to social anxiety disorder. Behaviour Research and Therapy 56:82–90.

Parker G, Brotchie H (2010) Gender differences in depression. International Review of Psychiatry 22:429–436.

Peirce J, Gray JR, Simpson S, MacAskill M, Höchenberger R, Sogo H, Kastman E, Lindeløv JK (2019) PsychoPy2: Experiments in behavior made easy. Behavior Research Methods 51:195–203.

Pessiglione M, Vinckier F, Bouret S, Daunizeau J, Le Bouc R (2018) Why not try harder? Computational approach to motivation deficits in neuro-psychiatric diseases. Brain 141:629–650.

Piantadosi PT, Yeates DCM, Floresco SB (2018) Cooperative and dissociable involvement of the nucleus accumbens core and shell in the promotion and

inhibition of actions during active and inhibitory avoidance. Neuropharmacology
138:57–71.

Pittig A, Treanor M, LeBeau RT, Craske MG (2018) The role of associative fear
and avoidance learning in anxiety disorders: Gaps and directions for future
research. Neuroscience & Biobehavioral Reviews 88:117–140.

R Development Core Team R (2011) R: A language and environment for statistical
computing.

Rizvi SJ, Pizzagalli DA, Sproule BA, Kennedy SH (2016) Assessing anhedonia in
depression: Potentials and pitfalls. Neuroscience and Biobehavioral Reviews
65:21–35.

Spielberger C (2008) State-trait anxiety inventory for adults. Garden.

Spitzer RL, Kroenke K, Williams JBW, Löwe B (2006) A Brief Measure for
Assessing Generalized Anxiety Disorder: The GAD-7. Archives of Internal
Medicine 166:1092.

Steer RA, Beck AT (1997) Beck anxiety inventory., pp23–40. Lanham, MD, US:
Scarecrow Education.

Thompson AH, Bland RC (2018) Gender similarities in somatic depression and in
DSM depression secondary symptom profiles within the context of severity and
bereavement. Journal of Affective Disorders 227:770–776.

Treadway MT, Bossaller NA, Shelton RC, Zald DH (2012) Effort-based
decision-making in major depressive disorder: A translational model of
motivational anhedonia. Journal of Abnormal Psychology 121:553–558.

Treadway MT, Buckholtz JW, Schwartzman AN, Lambert WE, Zald DH (2009)
Worth the 'EEfRT'? The effort expenditure for rewards task as an objective
measure of motivation and anhedonia. PLoS ONE 4:1–9.

Wang KS, Delgado MR (2021) The Protective Effects of Perceived Control During
Repeated Exposure to Aversive Stimuli. Frontiers in Neuroscience 15:625816.

Wurst C, Schiele MA, Stonawski S, Weiß C, Nitschke F, Hommers L, Domschke K,

Herrmann MJ, Pauli P, Deckert J, Menke A (2021) Impaired fear learning and

extinction, but not generalization, in anxious and non-anxious depression.

Journal of Psychiatric Research 135:294–301.

Legend

Figure 1: Trial layout diagram. A diagram of the active and inhibitory avoidance and reward-seeking tasks. In the avoidance task (A), after an inter-stimulus interval (ISI) with a fixation cross onscreen, participants were presented with a cue associated with active or inhibitory avoidance. For the active avoidance cue, participants had to respond with repeated spacebar presses to avoid hearing an aversive sound. For the inhibitory avoidance cue, participants had to withhold responding to avoid hearing an aversive sound. In the reward-seeking task (B), after the ISI, participants were presented with a cue associated with active or inhibitory reward-seeking. For the active reward-seeking cue, participants had to respond with repeated spacebar presses to obtain points towards a monetary reward. For the inhibitory reward-seeking cue, participants had to withhold responding to obtain points towards a monetary reward. ISI = Inter-stimulus interval.

Figure 2: Avoidance demographics. Distribution of anxiety (BAI) and depression (BDI) scores by gender and sample. Proportion scores are scores divided by total possible score. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory II.

Figure 3: Linear model significant effects for sensitivity in avoidance. (A) Gender interacted with anxiety scores (BAI proportion scores) to explain sensitivity (d') in the avoidance task. (B) Gender interacted with depression scores (BDI proportion scores) to explain sensitivity (d') in the avoidance task. BAI = Beck Anxiety Inventory. BDI = Beck Depression Inventory II. Proportion scores are scores divided by total possible score.

Figure 4: Avoidance effort deployment. (A) Multi-level model significant effects and

interactions for effort in avoidance. Effort decreased relative to criterion as the avoidance

task progressed, an effect that (B) interacted with anxiety scores (BAI) and gender.

Proportion scores are scores divided by total possible score.

Figure 5: Reward-seeking demographics. Distribution of anxiety (BAI) and

depressive (BDI) scores by gender and sample. Proportion scores are scores divided by

total possible score. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory II.

Figure 6: Linear model significant effects for sensitivity in reward-seeking. Gender

explained sensitivity (d') in the reward-seeking task. BAI = Beck Anxiety Inventory.

Proportion scores are scores divided by total possible score.

Figure 7: Reward-seeking effort deployment. (A) Multi-level model significant

effects and interactions for effort in reward-seeking. Effort decreased relative to criterion as

the reward-seeking task progressed, an effect that (B) interacted with anxiety scores (BAI)

and gender. Proportion scores are scores divided by total possible score.

Table 1: Demographic information for all participants.

Table 2: Mood disorder symptom statistics. Mean and SD Beck Depression

Inventory II (BDI) and Beck Anxiety Inventory (BAI) proportion scores (score divided by

total possible score).

Table 3: Linear model analysis coefficients and standard errors for sensitivity (d') in

avoidance. BAI (prop. score) = anxiety score on the Beck Anxiety Inventory. Proportion

scores are scores divided by total possible score. AIC = Akaike information criterion, BIC

= Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared

error.

Table 4: Linear model analysis coefficients and standard errors for sensitivity (d') in

avoidance. BDI (prop. score) = depression score on the Beck Depression Inventory II.

Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

Table 5: Multi-level model analysis coefficients and standard errors for effort in avoidance. BAI (prop. score) = anxiety score on the Beck Anxiety Inventory. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, ICC = intraclass correlation, RMSE = root mean squared error.

Table 6: Multi-level model analysis coefficients and standard errors for effort in avoidance. BDI (prop. score) = depression score on the Beck Depression Inventory II. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, ICC = intraclass correlation, RMSE = root mean squared error.

Table 7: Linear model analysis coefficients and standard errors for breakpoint in avoidance. BAI (prop. score) = anxiety score on the Beck Anxiety Inventory. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

Table 8: Linear model analysis coefficients and standard errors for breakpoint in avoidance. BDI (prop. score) = depression score on the Beck Depression Inventory II. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

Table 9: Linear model analysis coefficients and standard errors for sensitivity (d') in reward-seeking. BAI (prop. score) = anxiety score on the Beck Anxiety Inventory.

Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

Table 10: Linear model analysis coefficients and standard errors for sensitivity (d') in reward-seeking. BDI (prop. score) = anxiety score on the Beck Depression Inventory II. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

Table 11: Multi-level model analysis coefficients and standard errors for effort in reward-seeking. BAI (prop. score) = anxiety score on the Beck Anxiety Inventory. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, ICC = intraclass correlation, RMSE = root mean squared error.

Table 12: Multi-level model analysis coefficients and standard errors for effort in reward-seeking. BDI (prop. score) = depression score on the Beck Depression Inventory II. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, ICC = intraclass correlation, RMSE = root mean squared error.

Table 13: Linear model analysis coefficients and standard errors for breakpoint in reward-seeking. BAI (prop. score) = anxiety score on the Beck Anxiety Inventory. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

Table 14: Linear model analysis coefficients and standard errors for breakpoint in reward-seeking. BDI (prop. score) = depression score on the Beck Depression Inventory II.

906 Proportion scores are scores divided by total possible score. AIC = Akaike information

907 criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root

908 mean squared error.

Table 1

Demographic information for all participants.

Study	$N_{recruited}$	$N_{analyzed}$	N_{female}	N_{male}	N_{other}	M_{age}	$Range_{age}$
1A (Avoidance, undergraduate)	357	272	174	86	12	20.42	17-32
1B (Avoidance, paid global)	310	272	87	176	9	24.06	18-57
2A (Reward-seeking, undergraduate)	114	36	28	8	0	20.67	18-26
2B (Reward-seeking, paid global)	309	274	78	180	16	25.18	18-62

Table 2

Mean and SD Beck Depression Inventory II (BDI) and Beck Anxiety Inventory (BAI) proportion scores (score divided by total possible score).

Task	Gender	$M_{BDIprop}$	$SD_{BDIprop}$	$M_{BAIprop}$	$SD_{BAIprop}$
Avoidance	Female	0.30	0.19	0.29	0.22
	Male	0.23	0.16	0.23	0.19
Reward-seeking	Female	0.29	0.19	0.28	0.21
	Male	0.26	0.17	0.22	0.17

Table 3

Linear model analysis coefficients and standard errors for sensitivity (d′) in avoidance. BAI (prop. score) = anxiety score on the Beck Anxiety Inventory. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

	Linear model: BAI pred. Sensitivity (d′)
(Intercept)	2.46***
	(0.09)
BAI (prop. score)	−0.94***
	(0.23)
Gender	−0.07
	(0.11)
Sample	−0.17
	(0.11)
BAI (prop. score) x Gender	0.74*
	(0.31)
BAI (prop. score) x Sample	0.73*
	(0.29)
Gender x Sample	0.06
	(0.15)
BAI (prop. score) x Gender x Sample	−0.69
	(0.46)
Num.Obs.	523
R2	0.050
R2 Adj.	0.037
AIC	769.8
BIC	808.2
Log.Lik.	−375.910
F	3.893
RMSE	0.50

+ p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001

Table 4

Linear model analysis coefficients and standard errors for sensitivity (d′) in avoidance. BDI (prop. score) = depression score on the Beck Depression Inventory II. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

	Linear model: BDI pred. Sensitivity (d′)
(Intercept)	2.52***
	(0.11)
BDI (prop. score)	−1.09***
	(0.29)
Gender	−0.13
	(0.13)
Sample	−0.29*
	(0.13)
BDI (prop. score) x Gender	0.88*
	(0.38)
BDI (prop. score) x Sample	1.11**
	(0.35)
Gender x Sample	0.16
	(0.17)
BDI (prop. score) x Gender x Sample	−0.93+
	(0.53)
Num.Obs.	523
R2	0.042
R2 Adj.	0.029
AIC	774.1
BIC	812.4
Log.Lik.	−378.038
F	3.265
RMSE	0.50

+ p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001

Table 5

Multi-level model analysis coefficients and standard errors for effort in avoidance. BAI (prop. score) = anxiety score on the Beck Anxiety Inventory. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, ICC = intraclass correlation, RMSE = root mean squared error.

	Multi-level model: BAI pred. Effort
(Intercept)	5.74***
	(0.26)
BAI (prop. score)	−0.63
	(0.68)
Block	−0.91***
	(0.02)
Gender	0.29
	(0.32)
Sample	−0.42
	(0.33)
BAI (prop. score) x Block	0.16***
	(0.05)
BAI (prop. score) x Gender	0.97
	(0.92)
Block x Gender	0.04*
	(0.02)
BAI (prop. score) x Sample	−0.65
	(0.87)
Block x Sample	−0.01
	(0.03)
Gender x Sample	−0.19
	(0.44)
BAI (prop. score) x Block x Gender	−0.19**
	(0.06)
BAI (prop. score) x Block x Sample	−0.06
	(0.07)
BAI (prop. score) x Gender x Sample	0.94
	(1.36)
Block x Gender x Sample	0.04
	(0.04)
BAI (prop. score) x Block x Gender x Sample	−0.08
	(0.11)
SD (Intercept participant)	1.42
SD (Observations)	1.91
Num.Obs.	52261
R2 Marg.	0.289
R2 Cond.	0.542
AIC	218248.5
BIC	218408.1
ICC	0.4
RMSE	1.90

+ $p < 0.1$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

Table 6

Multi-level model analysis coefficients and standard errors for effort in avoidance. BDI (prop. score) = depression score on the Beck Depression Inventory II. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, ICC = intraclass correlation, RMSE = root mean squared error.

	Multi-level model: BDI pred. Effort
(Intercept)	5.67***
	(0.32)
BDI (prop. score)	−0.41
	(0.86)
Block	−0.92***
	(0.02)
Gender	0.24
	(0.38)
Sample	−0.40
	(0.38)
BDI (prop. score) x Block	0.19**
	(0.06)
BDI (prop. score) x Gender	1.21
	(1.11)
Block x Gender	0.08**
	(0.03)
BDI (prop. score) x Sample	−0.70
	(1.04)
Block x Sample	0.01
	(0.03)
Gender x Sample	−0.31
	(0.50)
BDI (prop. score) x Block x Gender	−0.34***
	(0.08)
BDI (prop. score) x Block x Sample	−0.12
	(0.08)
BDI (prop. score) x Gender x Sample	1.54
	(1.59)
Block x Gender x Sample	−0.01
	(0.04)
BDI (prop. score) x Block x Gender x Sample	0.12
	(0.13)
SD (Intercept participant)	1.42
SD (Observations)	1.91
Num.Obs.	52261
R2 Marg.	0.289
R2 Cond.	0.542
AIC	218241.9
BIC	218401.4
ICC	0.4
RMSE	1.90

+ p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001

Table 7

Linear model analysis coefficients and standard errors for breakpoint in avoidance. BAI (prop. score) = anxiety score on the Beck Anxiety Inventory. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

	Linear model: BAI pred. Sensitivity (d$'$)
(Intercept)	133.41***
	(7.78)
BAI (prop. score)	−18.55
	(20.00)
Gender	12.52
	(9.52)
Sample	−5.39
	(9.53)
BAI (prop. score) x Gender	42.11
	(27.05)
BAI (prop. score) x Sample	−12.37
	(25.25)
Gender x Sample	5.42
	(12.91)
BAI (prop. score) x Gender x Sample	−30.85
	(39.58)
Num.Obs.	523
R2	0.105
R2 Adj.	0.093
AIC	5440.4
BIC	5478.7
Log.Lik.	−2711.198
F	8.668
RMSE	43.16

+ p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001

Table 8

Linear model analysis coefficients and standard errors for breakpoint in avoidance. BDI (prop. score) = depression score on the Beck Depression Inventory II. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

	Linear model: BDI pred. Sensitivity (d′)
(Intercept)	136.99***
	(9.46)
BDI (prop. score)	−28.79
	(25.30)
Gender	11.13
	(11.28)
Sample	−10.46
	(11.10)
BDI (prop. score) x Gender	42.87
	(32.83)
BDI (prop. score) x Sample	2.96
	(30.35)
Gender x Sample	0.59
	(14.72)
BDI (prop. score) x Gender x Sample	1.39
	(46.40)
Num.Obs.	523
R2	0.101
R2 Adj.	0.089
AIC	5443.1
BIC	5481.4
Log.Lik.	−2712.530
F	8.250
RMSE	43.27

+ p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001

Table 9

Linear model analysis coefficients and standard errors for sensitivity (d') in reward-seeking. BAI (prop. score) = anxiety score on the Beck Anxiety Inventory. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

	Linear model: BAI pred. Sensitivity (d')
(Intercept)	3.12***
	(0.15)
BAI (prop. score)	−0.16
	(0.40)
Gender	0.34*
	(0.17)
Sample	0.15
	(0.27)
BAI (prop. score) x Gender	−0.75
	(0.52)
BAI (prop. score) x Sample	−0.48
	(0.84)
Gender x Sample	−0.53
	(0.42)
BAI (prop. score) x Gender x Sample	3.16
	(2.69)
Num.Obs.	294
R2	0.041
R2 Adj.	0.017
AIC	686.9
BIC	720.1
Log.Lik.	−334.458
F	1.739
RMSE	0.75

+ p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001

Table 10

Linear model analysis coefficients and standard errors for sensitivity (d′) in reward-seeking. BDI (prop. score) = depression score on the Beck Depression Inventory II. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

	Linear model: BDI pred. Sensitivity (d′)
(Intercept)	3.09***
	(0.16)
BDI (prop. score)	−0.05
	(0.46)
Gender	0.26
	(0.20)
Sample	0.07
	(0.30)
BDI (prop. score) x Gender	−0.28
	(0.58)
BDI (prop. score) x Sample	−0.15
	(0.92)
Gender x Sample	−0.35
	(0.50)
BDI (prop. score) x Gender x Sample	1.27
	(2.13)
Num.Obs.	294
R2	0.016
R2 Adj.	−0.008
AIC	694.5
BIC	727.6
Log.Lik.	−338.246
F	0.655
RMSE	0.76

+ p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001

Table 11

Multi-level model analysis coefficients and standard errors for effort in reward-seeking. BAI (prop. score) = anxiety score on the Beck Anxiety Inventory. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, ICC = intraclass correlation, RMSE = root mean squared error.

	Multi-level model: BAI pred. Effort
(Intercept)	5.27***
	(0.26)
BAI (prop. score)	1.44*
	(0.71)
Block	−0.87***
	(0.02)
Gender	1.30***
	(0.31)
Sample	1.08*
	(0.49)
BAI (prop. score) x Block	0.00
	(0.05)
BAI (prop. score) x Gender	−3.35***
	(0.93)
Block x Gender	−0.03
	(0.02)
BAI (prop. score) x Sample	−2.92+
	(1.52)
Block x Sample	−0.12**
	(0.04)
Gender x Sample	−2.05**
	(0.76)
BAI (prop. score) x Block x Gender	0.15*
	(0.07)
BAI (prop. score) x Block x Sample	0.39**
	(0.13)
BAI (prop. score) x Gender x Sample	3.62
	(4.81)
Block x Gender x Sample	0.22***
	(0.06)
BAI (prop. score) x Block x Gender x Sample	−0.79+
	(0.42)
SD (Intercept participant)	1.30
SD (Observations)	1.98
Num.Obs.	37677
R2 Marg.	0.282
R2 Cond.	0.498
AIC	159770.0
BIC	159923.7
ICC	0.3
RMSE	1.98

+ p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001

Table 12

Multi-level model analysis coefficients and standard errors for effort in reward-seeking. BDI (prop. score) = depression score on the Beck Depression Inventory II. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, ICC = intraclass correlation, RMSE = root mean squared error.

	Multi-level model: BDI pred. Effort
(Intercept)	5.64***
	(0.29)
BDI (prop. score)	0.15
	(0.82)
Block	−0.88***
	(0.02)
Gender	0.73*
	(0.35)
Sample	0.44
	(0.53)
BDI (prop. score) x Block	0.03
	(0.06)
BDI (prop. score) x Gender	−1.01
	(1.03)
Block x Gender	0.01
	(0.03)
BDI (prop. score) x Sample	−0.51
	(1.64)
Block x Sample	−0.07
	(0.04)
Gender x Sample	−1.29
	(0.89)
BDI (prop. score) x Block x Gender	0.00
	(0.08)
BDI (prop. score) x Block x Sample	0.17
	(0.13)
BDI (prop. score) x Gender x Sample	1.45
	(3.81)
Block x Gender x Sample	0.14+
	(0.07)
BDI (prop. score) x Block x Gender x Sample	−0.25
	(0.32)
SD (Intercept participant)	1.32
SD (Observations)	1.98
Num.Obs.	37677
R2 Marg.	0.276
R2 Cond.	0.499
AIC	159796.9
BIC	159950.6
ICC	0.3
RMSE	1.98

+ p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001

Table 13

Linear model analysis coefficients and standard errors for breakpoint in reward-seeking. BAI (prop. score) = anxiety score on the Beck Anxiety Inventory. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

	Linear model: BAI pred. Sensitivity (d$'$)
(Intercept)	113.69***
	(3.21)
BAI (prop. score)	7.21
	(8.83)
Gender	6.73+
	(3.84)
Sample	−2.79
	(6.05)
BAI (prop. score) x Gender	−13.91
	(11.57)
BAI (prop. score) x Sample	9.61
	(18.66)
Gender x Sample	3.79
	(9.34)
BAI (prop. score) x Gender x Sample	−3.59
	(59.57)
Num.Obs.	294
R2	0.019
R2 Adj.	−0.005
AIC	2507.8
BIC	2540.9
Log.Lik.	−1244.881
F	0.786
RMSE	16.70

+ p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001

Table 14

Linear model analysis coefficients and standard errors for breakpoint in reward-seeking. BDI (prop. score) = depression score on the Beck Depression Inventory II. Proportion scores are scores divided by total possible score. AIC = Akaike information criterion, BIC = Bayesian information criterion, Log. Lik. = log likelihood, RMSE = root mean squared error.

	Linear model: BDI pred. Sensitivity (d′)
(Intercept)	114.96***
	(3.56)
BDI (prop. score)	2.79
	(10.06)
Gender	6.66
	(4.30)
Sample	−3.31
	(6.45)
BDI (prop. score) x Gender	−12.94
	(12.64)
BDI (prop. score) x Sample	10.64
	(20.01)
Gender x Sample	3.16
	(10.88)
BDI (prop. score) x Gender x Sample	−1.17
	(46.44)
Num.Obs.	294
R2	0.019
R2 Adj.	−0.005
AIC	2507.8
BIC	2541.0
Log.Lik.	−1244.920
F	0.775
RMSE	16.70

+ p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001

Figure 1

A diagram of the active and inhibitory avoidance and reward-seeking tasks. In the avoidance task (A), after an inter-stimulus interval (ISI) with a fixation cross onscreen, participants were presented with a cue associated with active or inhibitory avoidance. For the active avoidance cue, participants had to respond with repeated spacebar presses to avoid hearing an aversive sound. For the inhibitory avoidance cue, participants had to withhold responding to avoid hearing an aversive sound. In the reward-seeking task (B), after the ISI, participants were presented with a cue associated with active or inhibitory reward-seeking. For the active reward-seeking cue, participants had to respond with repeated spacebar presses to obtain points towards a monetary reward. For the inhibitory reward-seeking cue, participants had to withhold responding to obtain points towards a monetary reward. ISI = Inter-stimulus interval.

Figure 2

Avoidance demographics: Distribution of anxiety (BAI) and depressive symptom (BDI) proportion scores (score divided by total possible score) by gender and sample. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory.

Figure 3

Linear models revealed gender interacted with anxiety and depression scores to predict avoidance task performance. (A) Gender interacted with anxiety scores (BAI proportion scores) to explain sensitivity (d′) in the avoidance task. (B) Gender interacted with depression scores (BDI proportion scores) to explain sensitivity (d′) in the avoidance task. BAI = Beck Anxiety Inventory. BDI = Beck Depression Inventory II. Proportion scores are scores divided by total possible score.

Figure 4

Avoidance effort deployment. (A) Multi-level model significant effects and interactions for effort in avoidance. Effort decreased relative to criterion as the avoidance task progressed. (B) Effort deployment interacted with anxiety scores (BAI) and gender. Proportion scores are scores divided by total possible score.

Figure 5

Reward-seeking demographics: Distribution of anxiety (BAI) and depressive symptom (BDI) scores by gender and sample. BAI = Beck Anxiety Inventory, BDI = Beck Depression Inventory II.

Figure 6

*Linear model significant effects for sensitivity in reward-seeking. Gender explained sensitivity (d')
in the reward-seeking task. BAI = Beck Anxiety Inventory. Proportion scores are scores divided by
total possible score.*

Figure 7

Reward-seeking effort deployment. (A) Multi-level model significant effects and interactions for effort in reward-seeking. Effort decreased relative to criterion as the reward-seeking task progressed, an effect that (B) interacted with anxiety scores (BAI) and gender. Proportion scores are scores divided by total possible score.

Time

A. Avoidance

B. Reward-seeking

ISI
(2000 ms,
jittered 1200 ms)

▲ = 1 key press