1   Calcium dependence of both lobes of calmodulin is involved in SK channel gating
2

D. Brent Halling[1], Ashley E. Philpo[1], Richard W. Aldrich[1]

[1]Department of Neuroscience, University of Texas at Austin, Austin, United States

Abstract

KCa2.1-3 Ca^{2+}-activated K^+-channels (SK) require calmodulin to gate in response to cellular
Ca^{2+}. A model for SK gating proposes that the N-terminal domain (N-lobe) of calmodulin is
required for activation, but an immobile C-terminal domain (C-lobe) has constitutive, Ca^{2+}-
independent binding. Although structures support a domain-driven hypothesis of SK gate
activation by calmodulin, only a partial understanding is possible without measuring both
channel activity and protein binding. We measured SK2 (KCa2.2) activity using inside-out patch
recordings. Currents from calmodulin-disrupted SK2 channels can be restored with exogenously
applied calmodulin. We find that SK2 activity only approaches full activation with full-length
calmodulin with both an N- and C-lobe. We measured calmodulin binding to a C-terminal SK
peptide (SKp) using both composition-gradient multi-angle light scattering and tryptophan
emission spectra. Isolated lobes bind to SKp with high affinity, but isolated lobes do not rescue
SK2 activity. Consistent with earlier models, N-lobe binding to SKp is stronger in Ca^{2+}, and C-
lobe binding affinity is strong independent of Ca^{2+}. However, a native tryptophan in SKp is
sensitive to Ca^{2+} binding to both the N- and C-lobes of calmodulin at Ca^{2+} concentrations that
activate SK2, demonstrating that the C-lobe interaction with SKp changes with Ca^{2+}. Our
peptide binding data and electrophysiology show that SK gating models need deeper scrutiny.
We suggest that the Ca^{2+}-dependent associations of both lobes of calmodulin to SKp are crucial
events during gating. Additional investigations are necessary to complete a mechanistic gating
model consistent with binding, physiology, and structure.

Introduction

SK channels modulate cell membrane excitability and help to maintain cytosolic Ca^{2+}
homeostasis. Mutations in an SK gene can cause diseases including Zimmermann-Laband
Syndrome and idiopathic noncirrhotic portal hypertension (Bauer et al., 2019; Koot et al., 2016).
Located on the plasma membrane, SKs open to release cellular K^+ in response to Ca^{2+} influx to
the cell after membrane depolarization. K^+ efflux helps to restore resting membrane potential,
which can result in inactivation of Ca^{2+} entry into the cell (Barrett and Barret, 1976; Hugues et
al., 1982; Moolenaar and Spector, 1979). Thus, SKs operate in the Ca^{2+} feedback cycle to fine
tune intracellular Ca^{2+} signals originating at the plasma membrane. Yet Ca^{2+} does not interact
physically with SK. Ca^{2+} activates the channel via another protein modulator that tethers itself to
diagonal subunits of a homotetrameric SK channel (**Figure 1A, B**).

Calmodulin (**CaM**) directs the Ca^{2+} response in virtually all eukaryotic cells. It binds hundreds of
targets that include ion channels. Different properties from each of its four unique Ca^{2+} binding
sites are highly conserved (Halling et al., 2016) leading to dynamic ranges of Ca^{2+} sensitivity
that can vary depending on which target CaM is bound to (Black et al., 2005; Newman et al.,
2008; Persechini et al., 2000; Westerlund and Delemotte, 2018). The four Ca^{2+} sites are
arranged into two pairs, each pair comprising a domain or 'lobe' of CaM. The two globular lobes
form a 'dumbbell' like appearance (Babu et al., 1985) (**Figure 1D**). The two Ca^{2+} sites within
each lobe may tune each lobe to respond differently to Ca^{2+} stimuli (Liang et al., 2003).

Each CaM target has evolved to use different properties of a versatile CaM architecture (**Figure**
1C-F). A common challenge encountered with mechanistic studies on CaM's targets is knowing
to what extent the lobes of CaM sense Ca^{2+} independently (Banerjee et al., 2018; Schumacher
et al., 2001; Tadross et al., 2008; Williams et al., 2018) or together as intercommunicating units
to carry out a function (Evans et al., 2011; Mukherjea et al., 1996; Persechini et al., 1996).
Changes in cytosolic Ca^{2+} alter how a tethered CaM binds to SK (Xia et al., 1998). For SK
gating, a simple model that was based on structures, electrophysiology, and some binding data
suggests that the C-terminal domain (C-lobe) of CaM is constitutively bound to SK independent
of Ca^{2+} (Keen et al., 1999; Lee and MacKinnon, 2018; Schumacher et al., 2001). The N-terminal
domain (N-lobe) forms the Ca^{2+} sensor of SK and only binds tightly in the presence of Ca^{2+}. We
refer to domain dependent mechanism as the 'N-sensor hypothesis' for SK gating (**Figure 1G**).
Although the N-sensor hypothesis has an experimental basis, it has not been thoroughly tested.

Certain properties must hold to validate the N-sensor hypothesis for SK: 1) The C-lobe is
insensitive to Ca^{2+} when bound to SK. 2) The N- and C-lobes act independently and carry out
distinct functions, i.e., the N-lobe is the Ca^{2+} sensor, whereas the C-lobe is the Ca^{2+}-
independent glue to keep CaM bound to SK. 3) The N-lobe and C-lobe bind at different sites so
that their unique functions can be carried out. In contrast, many CaM binding domains bind both
lobes in proximity and interact with other either directly or through peptide contacts (Meador et
al., 1992; Villarroel et al., 2014). Recent structures of SK4 with bound CaM show multiple
conformations of the N-lobe at trace Ca^{2+}, but a better resolved interaction at high Ca^{2+} (Lee and
MacKinnon, 2018). At high Ca^{2+} the N- and C-lobes of CaM bind to different domains of SK4 in
a unique conformation for CaM that has not been observed previously (**Figure 1A, B**). Given
the diversity of interactions in the structures, careful binding studies will be needed to sort out
how each domain of CaM interacts with SK at low and at high Ca^{2+}.

Although CaM modulation of SK has been studied for decades, there are gaps in our
fundamental understanding that still need to be worked out. Transduction of Ca^{2+} binding to
channel gating is a dynamic, multistep process. It includes Ca^{2+} binding to CaM, CaM binding to
SK, and the intrinsic motions of SK through different gating states.

SK function can be studied by exchanging WT-CaM with mutant CaM

WT-CaM has a high affinity for SK2 at both high and trace Ca^{2+} (Xia et al., 1998). In a
heterologous system, WT-CaM is always present, so we developed a method that allowed us to
exchange WT-CaM for a mutant CaM (Li et al., 2009) (**Figure 2A**). At 10 µM Ca^{2+}, SK2
approaches maximal open probability in the presence of WT-CaM. To study how different CaM
constructs can restore SK2 currents, we first co-expressed the SK channel with a mutant CaM
(E1Q) that has a glutamate instead of a glutamine at the conserved position 12 in the N-lobe EF
hand 1, which is residue 31 in the protein sequence. This mutation leads to only one functional
EF hand in the N-Lobe (Maune et al., 1992). We demonstrated (Li et al., 2009) that co-
expression of E1Q with SK2 in *Xenopus laevis* oocytes leads to a rapid rundown of SK2 current.
Current rundown can be recovered permanently upon application of exogenous WT-CaM, that
is SK2 current no longer runs down once WT-CaM associates with the channel. We assume
that exogenous CaM binds to empty CaM sites on SK from E1Q dissociation, or alternatively
E1Q leaves the C-lobe bound and that exogenous CaM can displace the mutant. These findings
gave us a tool to test whether SK channel activity could be recovered by various non-WT CaMs.

Binding studies provide information on a Ca^{2+} dependence of interactions

SK2 gating requires CaM to bind its C-terminus (Schumacher et al., 2001; Xia et al., 1998).
Solution binding studies can be performed on protein domains of SK that bind CaM. A peptide
of SK2 that comprises the CaM binding domain is referred to as **SKp** (**residues 396-487 of rat
KCa2.2**) has been used in multiple binding and structure studies (Halling et al., 2014; Nam et
al., 2017; Schumacher et al., 2001; Zhang et al., 2013, 2012a, 2012b). We used multi-angle
light scattering to show that full-length WT-CaM has unique properties with SKp that isolated,
separated, or duplicated lobes of CaM cannot reproduce.

Intrinsic fluorescence from a peptide, such as SKp, can be used to analyze CaM binding (Keen
1999). Tryptophan (W) has the strongest fluorescent properties of all the amino acids
(Lakowicz, 2006). Its fluorescence is also sensitive to its environment. Although tryptophan
emission spectra are subject to several factors, the comparisons of gross changes in its
amplitude and the location of its peak intensity can be a reporter for molecular binding. In
general, a spectrum with a peak at longer wavelengths suggests a solvent exposed tryptophan,
whereas a peak at shorter wavelengths describes a tryptophan in a more polar environment,
such as being buried in a protein. We used SKp-W432 spectra to show that either lobe of CaM
can bury this residue.

We use both electrophysiology and peptide binding data to show some properties of CaM that
are necessary for CaM to bind SKp, and how these findings relate to CaM dependent SK
function.

Results

WT CaM can recover SK channel activity in HEK293 cells

We wanted to explore what components of mammalian wild-type calmodulin (WT-CaM) are
necessary for SK channel activation. In control experiments, patches excised from cells that
were transfected with either SK2 alone or both SK2 and WT-CaM had minimal increase in SK
current in 10 μM Ca^{2+} when exogenous WT-CaM was applied for 297 ± 6 seconds or 278 ± 41
seconds respectively (SK alone, 21 ± 14%; n=4; SK + WT-CaM 32 ± 13%; n=3) (**Figure 2B, C**).
Since adding exogenous CaM does not substantially increase current, all available SK channels
in the excised patch are likely fully saturated with WT-CaM.

We conducted electrophysiological experiments to determine if non-wild-type CaMs could
support SK channel activation at saturating (10 μM) calcium. Because our expression system
currently is in HEK293 cells and the results from Li *et al.*, were produced from oocytes (Li et al.,
2009), we first validated the same approach in HEK293 cells. As expected, patches excised
from cells that were co-transfected with SK and E1Q had an average of a ~6-fold increase in
current level when exogenous 20 μM WT-CaM was applied for 300 seconds (n=3) (**Table 1,**
Figure 2D).

Like Oocytes, HEK293 cells produce endogenous CaM that transfected E1Q protein must
outcompete to produce the observed current rundown. If SK2 binds E1Q with lower affinity than
WT-CaM, one explanation is that the exogenously applied WT-CaM binds to sites that were
vacated by E1Q when the patch was excised. In the absence of E1Q, endogenous WT-CaM is
sufficient to form functional units with SK2.

Although we did not quantify the extent of SK channel run down after patch excision, like what
was done in oocytes, we are confident that as in oocytes the increase in SK current with the
application of exogenous WT CaM was due to the recovery of SK channel activity that had been
lost by the over expression of E1Q.

 SKp binds WT-CaM with different conformations at low and at saturating Ca^{2+}

The C-terminus of SK is required for CaM binding. We use a peptide to represent this site
(**SKp**). Complex formation between SKp and CaM constructs in solution was measured using
composition-gradient multi-angle light scattering (CG-MALS). CG-MALS measures the weight
average molar mass of particles in solution as a function of molar ratio of the molecules of
interest (Attri and Minton, 2005). Light-scattering signals are directly converted into a weight
average molar mass if protein concentrations and instrument parameters are precisely known
(Wyatt, 1993), thus quantifying higher order stoichiometries is a designed capability of this
approach. Our prior work shows that CaM and SKp form different stoichiometries that depend
on the molar ratio concentrations of CaM to SKp in aqueous solution (Halling et al., 2014). In a
'cross-over' experiment, the sample applied to the flow cell in the light chamber varies using a
series of injections to step from high molar concentrations of SKp relative to CaM crossing over
to high molar ratios of CaM relative to SKp. Complex formation is measured at each step of
varying protein ratios, thus capturing a wide range of possible stoichiometries in a single
experiment. For simplicity, we refer to the stoichiometries as follows: 1) When the molar
concentration of SKp exceeds WT-CaM, a complex with 2 SKp and 1 CaM (2SKp/1CaM) forms,
which we call "P-C-P". 2) At equimolar concentrations of SKp and WT-CaM it forms a
1SKp/1CaM complex, which we call "P-C". 3) When WT-CaM molar concentrations exceed
SKp, 1 SKp and 2 WT-CaMs (1SKp/2CaM) form a complex we call "C-P-C".

SKp has a molar mass of 11.1 kDa. We repeated SKp binding measurements with WT-CaM at
low and at high Ca^{2+} as a control to draw comparisons with different CaM constructs (**Figure
3A**). WT-CaM has a molar mass of 16.7 kDa. At 5 mM Ca^{2+}, the characteristic appearance of
the calculated weight-averaged molar masses shows an 'M' shape as a function of the molar
ratio of WT-CaM protein to SKp. The 'M' shape is fortuitous in our studies because the features,
i.e., peaks and valleys, constrain models that determine what stoichiometries WT-CaM and SKp
can form in solution. The light-scattering data are decomposed into molar fractions of free and
bound protein that are present in solution giving rise to the features. The data confirm that CaM
forms three different complexes (**Figure 3K**), P-C-P, C-P, and C-P-C.

Fitting the data to a model provides ranges on macroscopic association constants. Since these
data are weight averages, the C-P-C peak is smaller because it forms with weaker affinity than
the P-C-P complex, with a log K_A of 16.0 and 17.0 (p<0.01) respectively (**Table 2**). We note one
difference from our previous results (Halling et al., 2014), fitting for incompetent fraction did not
provide substantial improvements to our fits, so our data this time were fit with simpler models
and with fewer parameters. Eliminating the incompetent fraction allowed us to use the minimum
number of variables to avoid parameter compensation that could lead to non-unique solutions
for association constants.

Metal chelators, such as EDTA and EGTA are commonly used to reduce free metal in solution,
but the quantities of free metal ions in solutions are usually calculated or assumed. When we
use EGTA, we consider it a "trace Ca^{2+}" condition with Ca^{2+} concentrations assumed to be
nanomolar, but metal contaminants are unknown. At trace Ca^{2+}, there are fewer distinguishing
features (**Figure 3A, F**). Notably P-C-P is at much smaller quantities, and C-P-C is absent as
observed previously (Halling et al., 2014). Just as we observed with Ca^{2+}, in EGTA P-C has a
high affinity. Since the P-C complex has negligibly different masses at high and at trace Ca^{2+},
another approach is needed to observe whether Ca^{2+} causes a change to the interaction of the
peptide and CaM.
177

Intrinsic tryptophan fluorescence of SKp confirms calcium sensitivity of calmodulin.

In the absence of a binding partner, the SKp W432 emission spectrum has a peak position at or
near 348 nm in all buffers used (**Figure 4A-C**). The appearance of this emission spectrum is
consistent with earlier reports, and it suggests that in the absence of CaM the tryptophan in SKp
is solvent exposed (Keen et al., 1999).

In the presence of WT-CaM equimolar with SKp, the emission peak from W432 of SKp shifts to
shorter wavelengths (**Figure 4A-C**). At traceCa^{2+}, with 5 mM EGTA, the W432 peak is at 334
nm, and it has a 230% greater amplitude than SKp alone (**Figure 4A**).

In our electrophysiology experiments, SK2 has maximal opening at 10 µM free Ca^{2+}. On the
other hand, structures are usually solved at millimolar concentrations of Ca^{2+}. To draw
comparisons to both electrophysiology and structure data, we measured Ca^{2+}-bound CaM
binding to SKp at both 10 µM Ca^{2+} (**Figure 4B**) and 5 mM Ca^{2+} (**Figure 4C**). The spectrum of
W432 at either Ca^{2+} concentration is virtually the same. Compared to trace Ca^{2+} the amplitude
appears 30% weaker, and the peak is shifted towards shorter wavelengths (329 nm). We
suspect a 310 nm cut-on emission filter contributes to the decrease in amplitude as the peak
shifts towards smaller wavelengths, so the decrease in amplitude could be an artifact of our
optics. Still, the pronounced change in spectral properties indicates both that W432 senses
protein binding and that protein binding is sensitive to Ca^{2+}. These major differences in spectral
shapes strongly suggest that the P-C complex has different conformations at low and at high
Ca^{2+}.

We can use these features as we study other CaM constructs to determine what properties of
CaM are required for SK2 activity and for Ca^{2+} dependent interactions with SKp.

SK requires CaM with both an N-lobe and a C-lobe connected for activity to be recovered

Given that we could recover SK channel activity with the application of exogenous WT-CaM
protein in HEK293 cells, we then tested whether the non-WT CaM constructs could also recover
SK activity. For each experiment we excised inside-out patches from cells that were
cotransfected with SK and E1Q. We took each patch and calculated what fraction of the total
recovered SK current was due to the application of our various mutant CaM constructs (**Figure
5A**). The constructs included an N-lobe only, a C-lobe only, a mixture of N- and C-lobes, a
double N-lobe, a double C-lobe, and CaM with Ca^{2+} binding disrupted in the C-lobe CaM
(E34Q). The CaM (E34Q) mutant contains glutamate to glutamine (E to Q) substitutions at
positions 104 and 140 (sites 3 and 4) in the C-lobe, CaM (E34).

We found that none of the non-WT CaMs, except for the mutant CaM (E34Q), lead to a
magnitude of recovery comparable to WT-CaM (**Figure 5B, C Table 1**).
212

SKp binds N- and C-lobe with high affinity

The absence of SK current recovery by a non-WT CaM construct could indicate either that CaM
can still bind SK but requires its full protein sequence to activate the channel, or that CaM
binding is disrupted and the absence of CaM renders SK Ca^{2+} insensitive. Binding
measurements are necessary to determine whether our mutant CaM constructs can still bind to
SK.

N-CaM has a molar mass of 9.0 kDa. Combined with SKp, which has a molar mass of 11.1 kDa,
the molar mass of a 1/1 complex would be about 20 kDa. We used CG-MALS to determine if
this stoichiometry, or other combinations, can form at high and at trace Ca^{2+}. Unlike the features
seen with WT-CaM at high Ca^{2+}, N-CaM binding to SKp has a simple, triangular shape in CG-
MALS data (**Figure 6A**). The apex of triangular feature approaches the predicted molar mass of
a 1/1 complex. There are no bulges at other molar ratios that would indicate other
stoichiometries can form. The triangular shape with the apex at the predicted molar mass shows
a strong K_A for a 1/1 complex and that the P-C complex is dominant in the solution (**Figure 6A,**
E, Table 2). Similar extrapolated K_A values between N-CaM and WT-CaM binding to SKp might
indicate similar binding properties, but the interpretation could be misleading without additional
work. WT-CaM has four Ca^{2+} sites, but N-CaM only has two. Careful Ca^{2+} binding
measurements are needed to compare the K_A values of mutant CaMs with different numbers of
Ca^{2+} sites to WT-CaM in binding to SKp. The key comparisons of the MALS data at high Ca^{2+} is
that WT-CaM has more ways to bind to SKp than N-CaM, and that both CaMs can bind SKp
with strong affinity. In contrast at trace Ca^{2+}, N-CaM binding to SKp is noticeably weaker (lower
K_A) and mostly unbound (**Figure 6A, C, Table 2**).

C-CaM forms the same stoichiometry regardless of Ca^{2+} (**Figure 6B, D, F**). The association of
C-CaM with SKp is strong, it is also independent of Ca^{2+}. N-CaM forms only a 1/1 complex with
SKp when Ca^{2+} is present. Additional approaches are needed to determine whether Ca^{2+} alters
the configuration of P-C for SKp binding to C-CaM.

Both lobes of CaM have Ca^{2+} dependent associations with SKp

We showed in **Figure 4A, B** that Ca^{2+} alters the interaction of SKp with full-length CaM. We
wanted to use this observation to assay whether Ca^{2+} changes the configuration of N-CaM or C-
CaM. We compared the spectra of SKp + WT-CaM with spectra of SKp + either N-CaM or C-
CaM. At trace Ca^{2+}, the W432 spectra obtained with SKp bound to C-CaM is indistinguishable
from when SKp is bound to WT-CaM (**Figure 4D**). In contrast, when N-CaM is present, the
W432 spectrum is indistinguishable from when SKp is by itself, which is consistent with N-
CaM's low affinity for SKp at trace Ca^{2+}. Thus, we conclude that the environment of W432
appears to be dominated by the C-lobe of WT-CaM at trace Ca^{2+}.
248

10 µM Ca^{2+} is sufficient to fully activate SK2 channels (Li et al., 2009). Binding experiments at
10 µM Ca^{2+}show that SKp bound to either isolated N-CaM or isolated C-CaM produces spectra
that are very close to SKp + WT-CaM (**Figure 4 B, E**), but there is a subtle, 10-20% smaller
amplitude when the C-lobe is bound. The spectrum with SKp bound to N-CaM appears slightly
more like the spectrum of SKp bound to WT-CaM, but not enough to conclude that it is the N-
lobe of CaM that binds W432 when full length WT-CaM binds SKp. For binding measurements,
we wanted to make sure 10 µM Ca^{2+} is saturating. 5 mM Ca^{2+} does not change the spectra
further (**Figure 4 C, F**), indicating that 10 µM Ca^{2+} is sufficient for maximal effect of Ca^{2+} with
either or both lobes.

C-P-C only forms with full-length CaM

At high Ca^{2+} SKp forms the C-P-C complex with WT-CaM (**Figure 3A, F**). The C-P-C complex is
not observed with either N-CaM or C-CaM using the cross-over approach (**Figure 6**). We
wanted to make sure that a C-P-C complex with N- or C-CaM was not missed due to weaker
binding. To sample data at high molar ratios of CaM relative to SKp, we designed experiments
that would emphasize a Ca^{2+} dependent site. We used the high-affinity P-C complex of SKp and

WT-CaM as a tool. Beginning with the preformed P-C and holding the P-C concentration at a
constant value, we titrated increasing concentrations of WT-CaM. At low additions of WT-CaM
there is a slight increase in the weight-averaged molar mass followed by a decrease (**Figure**
7A). The increase in measured mass is due to the formation of C-P-C as WT-CaM binds to the
preformed SKp/WT-CaM complex. The decrease in weight-averaged molar mass at higher
concentrations of added WT-CaM is expected due to the rising levels of unbound WT-CaM,
which is much smaller than C-P-C but dominates the composition of the solution.

Using preformed P-C with WT-CaM, we next added N-CaM or C-CaM. Instead of showing an
increase in mass at low concentrations of added "half-CaM", the measured weight average
molar mass drops off with increasing mutant CaM (**Figure 7B**). The decrease in measured
mass follows the predicted attenuation in weight average molar mass if the P-C complex
remains intact, but the free N-CaM or C-CaM neither convincingly binds to SKp at another site
nor replaces WT-CaM in the preformed complex. Taken together, **Figure 7A, B** show that C-P-
C requires two full-length CaMs.

SKp binds one lobe of CaM if lobes are not joined

In **Figure 2**, combining N-CaM and C-CaM does not restore SK2 current. We wanted to know if
unconnected lobes CaM can bind SKp at the same time. N-CaM and C-CaM individually bind
SKp at high Ca^{2+}; however, a second unattached lobe does not measurably bind to SKp, i.e.,
there is no C-P-C with the half-CaMs. If we mixed both N-CaM and C-CaM together, could the
pair bind to SKp? In a cross-over CG-MALS experiment a sample with equimolar concentrations
of N-CaM and C-CaM is varied against SKp. The results again indicate high-affinity binding of
exactly one domain of CaM, not both, to SKp (**Figure 7C**). This experiment cannot distinguish
how much of P-C is SKp bound to N-CaM vs C-CaM. Unconnected N-CaM and C-CaM can only
bind one at a time, suggesting that once bound to SKp, the other lobe is prevented from binding
if it is not attached to the other lobe.

CaM (E34Q) rescues SK2 current like WT-CaM, but NN-CaM and CC-CaM do not

Unconnected lobes of CaM do not activate SK, nor do they bind in the other's presence. We
also wanted to know to what extent mutant full-length CaMs can activate SK. Previous work has
shown that the CaM (E34Q) construct can substitute for WT-CaM to make functioning SK2
channels (Keen et al., 1999; Li et al., 2009). We confirmed those results here, while showing
that fractional recovery of current is at least 50% of that of WT-CaM (**Figure 5A, B**). Due to
variability in channel density and unknown occupancy by endogenous WT-CaM from the HEK
cells, we are not able to confirm whether CaM (E34Q) rescues at a different rate or to a different
saturation level than WT-CaM. Qualitatively, their abilities to rescue SK2 current are the same
(**Figure 5C**).

NN-CaM and CC-CaM are approximately the same size as WT-CaM. If a CaM site on the C-
terminus of SK can bind either lobe, an N-sensor view of CaM activation might suggest that at
high Ca^{2+}, both lobes would be able to bind SK2, and perhaps, gate the channel. Neither NN-
CaM nor CC-CaM activate SK2 convincingly (**Figure 5**). Thus, size is not the only factor, SK2
gating requires both an N- and a C-lobe.

E to Q CaM constructs have different interactions with SKp than WT-CaM

CaM (E34Q) can activate SK2. Although SK2 activation suggests that CaM (E34Q) appears to
function like WT-CaM, we currently do not know whether SK2 gating is the same with either
CaM bound to it.

CG-MALS data show that WT-CaM forms three different stoichiometries at high Ca^{2+} (**Figure**
3A, K). The profile for CaM (E34Q) looks different (**Figure 3D**). The characteristic 'M' shape is
gone. Instead, there is a stronger peak at equimolar ratios of CaM (E34Q):SKp, and the right
hump is absent. This can best be explained by a change in stoichiometries that form when SKp
binds WT-CaM compared to CaM (E34Q). Whereas there is support for a P-C-P complex, as
seen for WT-CaM, differences are seen at increasing molar ratios of CaM (E34Q). At equimolar
ratios of WT-CaM:SKp, A 1/1 complex of CaM/SKp is observed with 28 kDa. However, the
weight average molar mass at equimolar concentrations of CaM (E34Q):SKp is about 34 kDa
(**Table 2**). The best solution to fit the data includes a small quantity of a 2/2 complex. We call
this 2/2 complex the 2x-P-C (**Figure 3N**). At higher molar ratios of CaM (E34Q):SKp, there is no
evidence for C-P-C, and in fact the weight average molar mass falls off quickly with increasing
CaM (E34Q) arguing against a C-P-C with CaM (E34Q) (**Figure 3N**). In contrast, there are no
obvious differences at trace Ca^{2+} (**Figure 3F, I**).

We have shown that mutating only one N-lobe Ca^{2+} binding site is sufficient to disrupt CaM
binding to SK2 enough so that we can exchange CaMs in our rescue experiments. CaM (E12Q)
forms high affinity P-C with SKp at both low and high Ca^{2+}, but with weaker evidence for a P-C-
P and no C-P-C (**Table 2**) (**Figure 3E, J, O**).

We also used W432 spectra to evaluate the Ca^{2+} dependent changes of these E to Q
complexes. At trace Ca^{2+}, both CaM (E12Q) and CaM (E34Q) produce spectra similar to WT-
CaM (**Figure 4G**). At 10 μM Ca^{2+}, the W432 spectrum with CaM (E12Q) bound is comparable in
amplitude (<15%) and peak position (both 329 nm) to WT-CaM (**Figure 4H**). If Ca^{2+} binding to
the N-lobe is disrupted in E12Q as intended, the spectral overlap with WT-CaM strongly
suggests that Ca^{2+} binding to C-lobe is at least partially responsible for the spectral shift
observed with WT-CaM (**Figure 4E**).

On the other hand, the spectrum of W432 with CaM (E34Q) bound has weaker Ca^{2+}
dependence (**Figure 4H**). Ca^{2+} binding to the intact N-lobe of E34Q is not enough to produce a
strong spectral change (≤ 3 nm shift and <20% from **Figure 4G to 4H**)) that would overlap with
the WT-CaM in Ca^{2+} (≥ 4 nm and >30% from **Figure 4A to 4B**). 5 mM Ca^{2+} does not shift the
curve farther to smaller wavelengths indicating that higher Ca^{2+} cannot drive a weaker
interaction caused by the E to Q mutations (**Figure 4I**).

Summarizing the CaM (E34Q) results, 1) E34Q CaM rescues like WT-CaM, 2) CaM (E34Q) has
different Ca^{2+} dependent interactions with SKp than WT-CaM, and 3) Ca^{2+} is less capable of
altering the W432 environment when SKp is bound to CaM (E34Q) than when bound to WT-
CaM. All these argue that the mutant CaM (E34Q) operates differently than WT-CaM to produce
similar functional outcomes.

NN-CaM and CC-CaM have Ca^{2+} dependent interactions with SKp

The double-lobe constructs were used to draw out more differences between the lobes. Neither
construct can substitute for WT-CaM and restore channel activity (**Figure 5**). We used these
constructs as tools to evaluate Ca^{2+} dependent binding to SKp.

N-CaM by itself has a low affinity for SKp at trace Ca^{2+} (**Figure 6A, C**). In contrast, NN-CaM
binds with stronger affinity (p<0.01) (**Figure 3B, Table 2**). Although data clearly display binding,
the weight average molar mass is less than a P-C complex indicating a significant mole fraction
of unbound protein in solution (**Figure 3G**). We observe that P-C-P may be present, but our
confidence in higher order complexes is low since multiple models for oligomers can produce a

flatter shaped curve with weak binding. It's only when the weight average molar mass exceeds
that of a 1/1 complex across molar ratios that models become more constrained for higher
stoichiometries.

At high Ca^{2+}, P-C-P and C-P-C both appear at lower mole fractions with NN-CaM than with WT-
CaM (**Figure 3B, L**). The weight average molar masses exceed what would be expected by the
P-C complex alone, so there we have some confidence that P-C-P and C-P-C are detected. At
trace Ca^{2+}, the weight averaged masses are less than P-C, and there are more solutions
involving different combinations of weaker affinities (**Figure 3B, G**). On one hand, the similarity
in the appearance of the data might argue that C-P-C exists even at trace Ca^{2+}; however, we
are more confident using a model with fewer variables and suggest that at a minimum the P-C is
present with lesser amounts of C-P-C. The key observation is that NN-CaM binds stronger than
N-CaM at trace Ca^{2+} and that there is a decreased Ca^{2+} sensitivity of complex formation
compared with WT-CaM.

Except for geometric constraints, both lobes of CC-CaM are expected to bind SKp to make P-C-
P. C-CaM by itself does not show Ca^{2+} sensitivity for stoichiometry formation, but the spectral
data shows that C-CaM binding to SKp is Ca^{2+} dependent (**Figure 4D, E**). Using CG-MALS,
CC-CaM shows a small Ca^{2+} dependent change (**Figure 3C**). P-C-P shows up weaker at trace
Ca^{2+} than at high Ca^{2+}(**Figure 3H, M**). Just like NN-CaM, CC-CaM shows a subtle but
consistent hint of C-P-C at high Ca^{2+}.

For a bimolecular interaction, e.g., P-C, CG-MALS approaches a limit of detection for solving
the K_A. For bimolecular fitting with no higher order oligomers, log K_A values above ~7.0 are not
well constrained (**Table 2**). Other stoichiometries being present can broaden the range for the
limit of detection. At high Ca^{2+}, WT-CaM, CC-CaM, NN-CaM, and CaM (E34Q) all have features
in the data that expand the range for solving for K_A of P-C to higher values than 7.0 (**Figure 3A-
D**). For example, the dip in the M-shape for WT-CaM can only be observed if there is a K_A
greater than 10. Even in this case there is some ambiguity about whether the limit was crossed,
and it could be stronger yet. Thus, we limit our interpretations to be qualitative.

NN-CaM and CC-CaM binding confirm that both lobes retain Ca^{2+} sensitivity when bound to
SKp and that each lobe has different properties. Both N-lobe and C-lobe have Ca^{2+} sensitive
interactions that SK2 decodes during function.

Discussion

The C-terminus of SK with bound CaM is a Ca^{2+} sensor for SK

CaM is tightly bound to SK regardless of Ca^{2+}. It is required for SK to decode Ca^{2+} signals that
vary in amplitude, location, and duration. CaM has an extremely versatile structure (**Figure 1C-
F**). It can position its lobes, tune the Ca^{2+} sensitivity of its diverse partners, compete with other
proteins, and have dynamic associations.

An early study narrowed the essential CaM modulatory domain to SK2's C-terminus, and there
was no indication of another region of SK2 involved with CaM binding (Xia et al., 1998).
Although recent work suggests that other domains of SK may play roles in gating and need to
be considered in complete gating models (Lee and MacKinnon, 2018; Nam et al., 2021) (**Figure
1A, B**), multiple lines of evidence show that the C-terminus is the critical Ca^{2+}-sensing
component in SK gating. Engineered mutations in the C-terminus affect trafficking and gating of
channels in a heterologous system (Keen et al., 1999; Lee et al., 2003; Nam et al., 2021),

whereas cells from different tissue control SK activity through splice variations in the C-terminus
of SK (Zhang et al., 2001, 2012a).

In addition to SK2's C-terminus, a second CaM-N-lobe binding site has been shown on SK4 at
the S4-S5 linker (Lee and MacKinnon, 2018). Additional studies of heterologously expressed
SK2 channels show that mutations in either the S4-S5 linker or C-terminus can make SK2 more
sensitive to Ca^{2+} (Nam et al., 2021, 2017; Orfali et al., 2022). Although there are mutations in
S4-S5 that make SK2 gate at lower Ca^{2+}, more work is needed to determine whether the
mutations alter an opened or closed state of SK2 so that it opens easier or if the mutations alter
CaM interactions such that Ca^{2+} binds the complex at lower concentrations. Binding
measurements of CaM to other SK2 domains, such as the 4-5 linker, are difficult given the
folding of the S4-S5 linker and have yet to be completed.

Lessons from structures

If Ca^{2+} affects C-lobe interactions at 10 µM concentrations, why then is Ca^{2+} not seen as density
in the C-lobe binding sites of the structures solved at higher Ca^{2+}? High-resolution x-ray
structures would be able to detect ions as strong electron densities, and there are methods to
identify which metal ion is coordinated (Echols et al., 2014; Harding, 2004). The absence of
Ca^{2+} in the C-lobe is an experimental result that depends on the conditions used to arrive at the
data. A major challenge in crystallography is getting a protein to form a crystal, which requires
high protein concentrations in a somewhat arbitrarily determined medium. CaM co-crystallizes
with SKp in acidic solutions that range from pH 4.6 to 5.6 at high ionic strength (~1M Li^+)
(Schumacher et al., 2001; Zhang et al., 2013, 2012a). Both proton concentration and ionic
strength alter the affinity of CaM for Ca^{2+} (Linse et al., 1991; Ogawa and Tanokura, 1984;
Steiner et al., 1983). Even so, in one instance the C-lobe of CaM still is Ca^{2+} bound under acidic
conditions when the SK peptide contains an additional three residues as a splice variant,
showing that there are conditions at extreme pH (pH ~4.7) when the C-lobe is still Ca^{2+}
occupied (Zhang et al., 2012a). Interestingly, CaM was found with a C-shape in the original
structure (**Figure 1E**), but with an S-shape with the splice variant (**Figure 1D**).

Recently, structures at low and at high Ca^{2+} were determined for a full SK channel under
different experimental conditions (Lee and MacKinnon, 2018): 1) it used the full-length channel
as a tetramer with 4 bound CaMs with W-shapes (**Figure 1A-C**), 2) It used a more distantly
related SK family member, the SK4 channel, 3) the media is at more physiological ionic strength
(0.15M) and at pH 8, and 4) electron densities were from cryogenic electron microscopy (cryo-
EM). The structure obtained at high Ca^{2+} shows that on CaM one C-lobe site is Ca^{2+} bound and
the other is Ca^{2+} free. Cations are difficult to assign at worse than 3 angstrom resolution under
conditions where electron densities are approachable experimentally (Echols et al., 2014;
Harding, 2004). Both resolution limitations and the lack of anomalous scattering in modern
electron cryo-EM make unambiguously assigning ions a challenge, but methods are emerging
to locate and define bound metal ions (Elad et al., 2017). A vital question that remains is
whether both C-lobe Ca^{2+} are occupied at 10 µM, or if only one site is occupied as reported (Lee
and MacKinnon, 2018). Our results do not distinguish occupancy of one vs two Ca^{2+} sites within
a lobe. To our knowledge, there are no reports of lobe specific Ca^{2+} binding measurements of
CaM bound SK.

Even with a clearer picture of the architecture, high resolution details remain elusive. Cryo-EM is
an ensemble method that averages proteins across many states. Even though resolution is
often the primary goal, this approach can explore different states (Orlova and Saibil, 2010; Roh

et al., 2017; Zhang et al., 2019). Data are class averaged representations of thousands of
particles. Class averaging minimizes differences while amplifying molecular similarities. Less
frequent observed states are not necessarily minor functional states, and conversely more
frequent observed states may not be functionally relevant. However, structures of a subset of
states can generate testable hypotheses. Continued studies using cryo-EM on SK/CaM
complexes would be valuable.

Limits of the N-sensor hypothesis in SK gating

Across structures, W432 of SK2 binds the C-lobe of CaM (Lee and MacKinnon, 2018;
Schumacher et al., 2001; Zhang et al., 2012a). The C-lobe was modeled with a putative Ca^{2+}
ion bound to one of its sites in the Ca^{2+}-bound cryo-EM structure. The similarity of the C-lobe
structures at both trace and high Ca^{2+} forms the basis for the N-sensor model. The basis for the
N-sensor hypothesis is that the C-lobe is tethered independent of Ca^{2+}. In contrast to C-lobe
binding, the N-lobe of CaM in subsets of cryo-EM class averaged data was observed in multiple
conformations at trace Ca^{2+}(Lee and MacKinnon, 2018). The flexibility of N-lobe binding could
indicate different binding modes with SK in the absence of Ca^{2+}. Models from structures,
however, are strongly biased towards what can be observed either in a crystal or through class
averaging of thousands of molecules in cryo-EM. After particle image class averaging, a small
percentage (~40%) of total observed molecules is included in the final, highest 3.5 Å resolution
structure of SK. There is an opportunity to learn what information is contained in the rest of the
datasets about CaM binding.

The N-sensor hypothesis argues that the C-lobe has a weak Ca^{2+} dependence, and therefore
has weaker binding for Ca^{2+}. For binding, this would infer that the C-lobe would not alter the
environment about W432 of SKp with added Ca^{2+}. Weaker Ca^{2+} binding to the C-lobe is not
consistent with our data.

At minimally saturating Ca^{2+}, the spectra for CaM (E34Q) should bind SK like WT-CaM and
CaM (E12Q) should appear like WT-CaM does at trace Ca^{2+}. Our data show the opposite. Our
results are consistent with what is expected if both lobes bind Ca^{2+} and better support the NC-
sensor hypothesis. CaM has four Ca^{2+} binding sites, and the NC-sensor hypothesis is more
intuitive from an energetic scheme that includes all ligand binding sites.

The N-sensor hypothesis for SK gating by WT-CaM (**Figure 1**) is not supported by binding data,
but the mutant CaMs show results that could have been considered as consistent with the N-
sensor hypothesis. CaM (E34Q), whether overexpressed (Keen et al., 1999) or reconstituted (Li
et al., 2009) (**Figure 5**), activates SK2. If the E to Q mutations only alter Ca^{2+} binding to CaM,
the CaM (E34Q) data suggest that Ca^{2+} binding to N-lobe is required for SK gating, but not to
the C-lobe. The assumption was that CaM (E34Q) activates SK2 the same as WT-CaM.
However, our protein binding data challenges that assumption. CaM (E34Q) and WT-CaM do
not interact with SK in the same way and CaM (E34Q) clearly has a different binding mode.

SK gating requires both lobes of CaM

How does Ca^{2+} open SK channels? The answer is maybe it doesn't; Ca^{2+} just keeps SK from
closing. Hirschberg *et al.*, used single channel recordings of SK2 to show that the SK2 has
many closed and open states (Hirschberg et al., 1998). A key finding was that only a closed
state with long durations is unambiguously Ca^{2+} sensitive; none of the open states or short
duration closed states are Ca^{2+} sensitive. This observation sets a premise for SK gating: Ca^{2+}
must do work to prevent SK from visiting the closed state that has the longest dwell time. When

SK is activated, this essentially means that long closed times are diminished. More binding data,
electrophysiology, and structures are essential to learn how SK traverses different states in its
response to Ca^{2+}.

SK gating by CaM requires various properties of CaM. From our electrophysiology (**Figure 5**)
we find that the following are necessary for CaM to gate SK: 1) CaM must have two joined
lobes. 2) CaM must have both an N- and a C-lobe. 3) The N-lobe needs intact Ca^{2+} sites. We
infer that CaM with mutations in C-lobe, i.e., CaM (E34Q), can activate via an alternate
mechanism.

Our binding data reveal that CaM has dynamic associations with a peptide in response to Ca^{2+}.
One limitation of our study is that a peptide is free to bind CaM in configurations that are less
constrained than it would be in a membrane bound channel. Further binding data with CaM to
full channels are needed. Regardless, the peptide binding data still informs that Ca^{2+} alters
binding as expected for a sensor. We learn from the C-P-C configuration that Ca^{2+} opens a site
on SKp that is different than when there is no Ca^{2+} present. In the context of a full channel, there
might not be room for CaM molecules to associate (Lee and MacKinnon, 2018), but in terms of
binding it means that at high Ca^{2+}, CaM has more ways to interact with SK.

C-P-C binds strongest with WT-CaM and only in Ca^{2+} (**Figure 8**). The double-lobe mutants CC-
CaM and NN-CaM show a suggestion of C-P-C, but with WT-CaM C-P-C is clearly stronger.
With half-CaMs, neither N-CaM nor C-CaM form C-P-C, although either one binds SKp with
high affinity. If the N- and C-lobes bind at different specialized sites, there would be an
expectation that both the N-CaM and C-CaM can bind SKp together, but they do not (**Figure
7C**). However, C-P-C requires both an N- and a C-lobe. One alternative that has not yet been
tested is that the 'linker' between lobes contains the necessary requirements for C-P-C
formation. N-CaM and C-CaM each possess part of the linker and overlap at residue positions
78-80 (DTD). The presence of part of the linker on each construct somewhat tests the
hypothesis that the linker may need to be present to form C-P-C, although, because there is not
a full and intact continuous linker present, half-CaM's with variable linkers would be needed for
testing whether the linker is required for C-P-C formation.

For C-P-C, both lobes of both CaMs exchange near a common binding site that CaM only
senses in the presence of Ca^{2+}. There are two possible arrangements that our data cannot
distinguish that we present as scheme 1 and scheme 2 (**Figure 8**). In scheme 1, two CaMs bind
directly to SKp and have no association with the other CaM. In Scheme 2, SKp binds to a CaM
oligomer. We favor scheme 1 because our binding measurements have not detected any CaM
dimers in our solution measurements.

Testing these constraints requires having probes that can measure the N- and C-lobe positions
on the peptide and the conformation of the P-C complex. Our hypothesis predicts multiple
conformations of the P-C complex to account for the binding behavior of WT-CaM. Data that
show distributions of different conformations would support this conjecture. In contrast, CaM
(E34Q) binding to SKp potentially can be explained with fewer states.

For P-C-P we speculate that both lobes of a single CaM molecule participate with binding to
both SKp peptides, possibly in a more globular form but not necessarily in a "bridging"
conformation (**Figure 8**). Indeed, CaM binds many targets with an O-shape (**Figure 1F**), except
that the O-shape has not yet been observed in structures with SK. We do not detect dimers of
SKp during our measurements using purified peptide, yet we note that impurities in the solution

may cause the peptide to precipitate. Since we don't observe SKp dimers, we favor scheme 1 in
Figure 8 to describe P-C-P.

Finally, at trace Ca^{2+}, we only consider a strong P-C signature as convincing (**Figure 8**). Further
investigation is needed to find out whether P-C-P exists at trace Ca^{2+} and how the weaker
binding would fit into a gating model. In our current model, both lobes of CaM interact with SK at
trace Ca^{2+} in the P-C state.

Predictions of SK gating from CaM binding data

A more general model is needed to describe how SK decodes Ca^{2+} binding to both lobes of
CaM. The 'NC-sensor hypothesis' proposes that binding of Ca^{2+} to both lobes of CaM in
complex with SKp are energetically coupled processes, i.e., the entire CaM molecule works as a
unit and *not* as a sum of its parts. Although both lobes have unique qualities as they do in the N-
sensor hypothesis, binding data suggest that they work as a unit by communicating through SK.
The NC-sensor hypothesis has testable predictions. Both WT-CaM and CaM (E34Q) activate
SK, but since one lobe of CaM (E34Q) is less Ca^{2+} sensitive, SK must have different dwell times
in the opened or closed states. Single channel recordings of SK current can probe different
activation states of SK. A careful study of distributions of SK states would distinguish whether
different opened and closed states are prevalent if CaM (E34Q) is bound instead of WT-CaM.

Single channel recordings of SK2 show that in trace Ca^{2+} the open probability is dominated by a
long-lived closed state (Hirschberg et al., 1998). Rare bursts of activity show openings that are
marked with flickers to closed states. **At high Ca^{2+}, the channel is open about 80% of the**
time. The 'open' channel still has frequent flickers to a closed state. The increase in open
probability seems to be entirely due to a drastic reduction in the long-lived closed state and not
an increase in time spent in the open state, i.e., there is not a decrease in 'flickering'. Thus, at
high Ca^{2+}, SK2 shows extremely dynamic behavior. At equilibrium, a distribution of different
states of SK2 should be expected. Whether CaM (E34Q) is bound is expected to modify this
behavior.

More structures of SK with bound CaM captured in identifiable closed and open intermediates
would help to interpret how CaM alters SK activity. We know of at least three closed and two
open states for SK activity from physiology (Hirschberg et al., 1998). Our stoichiometry data
shows that SK has multiple ways to bind CaM, and there is ample structural support displaying
how flexible CaM is when bound to targets. Our binding and functional work are only edge
cases for concentrations and there is much more work to be done to solve what the protein
experiences dynamically during gating.

A generalized hypothesis for SK gating

CaM binding to peptide may be different from how it binds to a full-length protein such
as a channel, especially if another domain such as the 3-4 linker interacts with a lobe of
CaM. This is an extraordinarily difficult question to answer with just one binding study,
yet incorrect aspects of models can quickly be eliminated. We observe that the C-lobe
of CaM is calcium sensitive when binding to a peptide. How can the full-length channel
binding to the C-lobe be Ca^{2+} insensitive? It would require that the full-length channel
blocks the Ca^{2+} C-lobe configuration while preserving the Ca^{2+} free configuration, but
this has not been demonstrated. Considering that the channel has at least 3 closed
states and 2 open states, C-lobe Ca^{2+} sensing is highly likely to be important during

gating. From current structure analyses, we have at best an ensemble view of two out of
five channel/CaM states.

Although the N-sensor hypothesis is elegant, the binding data show that a more complex model
is likely necessary to truly represent the mechanism for SK gating by WT-CaM (**Figures 1G,8**).
Both lobes of CaM are energetically coupled through binding to SK. The complex of SK and
CaM decodes a Ca^{2+} signal, with both lobes in complex with SK acting as sensors. The C-lobe
of CaM is a Ca^{2+} sensor; we show that it is not insensitive to Ca^{2+} as described in the N-sensor
hypothesis. We propose that the mechanism for the NC-sensor hypothesis requires solving
energetic values for Ca^{2+} binding to four ligand sites of CaM, multiple CaMs binding to SK, and
SK open probability at different liganded states. The challenge of solving energetic states is far
greater and more general in the NC-sensor hypothesis, but it provides a realistic starting point
for solving the mechanism of SK activation.

Methods

Materials and constructs

The mammalian WT-CaM protein sequence (**NCBI: NP_059022**) expression clone from rat
DNA (**Rattus norvegicus calmodulin 2 (Calm2) NCBI: NM_017326.3**) was provided by S.
Hamilton (Baylor College of Medicine, Houston, TX), which we cloned into the Novagen pET21a
vector (Merck KGaA, Darmstadt, Germany) for expression in BL21(DE3) bacteria (New England
Biolabs (NEB), Ipswich, MA). Double lobe CaM DNA constructs, NN-CaM and CC-CaM were
codon optimized for bacterial expression and synthesized by Genewiz (South Plainfield, NJ). J.
Adelman (Vollum Institute, Portland, OR) provided the clone for the Rattus norvegicus **SKp**
(**residues 396-487 rat KCa2.2, UniProt accession P70604.1**) of SK2, and was cloned without
a tag as described previously (Halling et al., 2014). The human protein sequence of SK2 is
identical in this region. Reagents used for buffers were minimal American Chemical Society
quality and purchased from Sigma-Aldrich unless otherwise indicated. All solutions were made
with 18.2 MΩ water from a Milli-Q® Integral 5 System (Burlington, MA).

Cloning of proteins for expression

The N-CaM construct was designed by cloning in a stop codon (TAA) into the WT-CaM
construct in the pET21a vector using standard PCR mutagenesis after residue D80 of the
protein sequence using primers ordered from Sigma-Aldridge (Merck). C-CaM was cloned using
PCR amplification of the C-terminus of WT-CaM with added flanks of a 5' NdeI overhang and a
3' BamHI overhang and ligated into pET21a vector (Merck). Insertion of PCR product into
pET21a was achieved with standard protocols (NEB).

The CaM constructs we tested fall into four categories. 1) WT-CaM is used as our standard. 2)
Isolated CaM domains, i.e., isolated N-terminal domain (N-CaM) or C-terminal domain (C-CaM),
were used to determine if Ca^{2+} binding to an isolated domain of CaM by itself could activate
SK2. In some experiments, we used N-CaM and C-CaM in the same mixture to see if the parts
of CaM could restore SK2 activity without both lobes being joined. 3) Double-lobe CaMs that
have repeated domains were used to determine if the domains of CaM have compensating
roles in SK2 activation. We made CaM with two N-lobes but no C-lobe (NN-CaM) and CaM with
two C-lobes but no N-lobe (CC-CaM). 4) Finally, we used full-length CaM mutants that are
deficient for Ca^{2+} binding to one domain, but the opposite domain has normal Ca^{2+} binding. The

CaM (E12Q) mutant CaM has glutamines (Q) in place of glutamates (E) and positions 31 and
67 in the N-lobe of CaM. The E to Q mutations in the N-lobe were reported to disrupt Ca^{2+}
binding to sites 1 and 2 (Mukherjea et al., 1996). Similarly, the CaM (E34Q) mutant, which has
E to Q mutations at positions 104 and 140, was reported to disrupt binding to sites 3 and 4 in
the C-lobe.

For CaM protein sequence, the N-terminal methionine gets cleaved in the final protein product
and the next alanine is acetylated (**Sasagawa Biochemistry 1982**). In our protein constructs,
the N-terminal methionine, which is subject to cleavage during expression, is referred to as
position M0. The N-CaM protein sequence (residues M0-D80 of WT-CaM) is

MADQLTEEQIAEFKEAFSLFDKDGDGTITTKELGTVMRSLGQNPTEAELQDMINEVDADGNGTI
DFPEFLTMMARKMKDTD-

The C-CaM protein sequence (M76-K148) is

MKDTDSEEEIREAFRVFDKDGNGYISAAELRHVMTNLGEKLTDEEVDEMIREADIDGDGQVNY
EEFVQMMTAK-

NN-CaM and CC-CaM were purchased as described in materials and arrived in pUC57-
Kanamycin vector. The gene sequences were flanked by restriction enzyme sites for 5' NdeI
and 3' EcoRI (NEB). Genes were cloned into pET21a using standard protocols from NEB. Due
to a need for a start codon, the CC-CaM positions 1 and 2 (residues MA) do not align with WT-
CaM. Residues D78-D81 of WT-CaM were retained as the flexible part of the linker in both
constructs of NN-CaM and CC-CaM. Some effort was made to make the linkers of NN-CaM and
CC-CaM of similar length.

The NN-CaM protein sequence (residues M0-D81, Q8-K77) is
MADQLTEEQIAEFKEAFSLFDKDGDGTITTKELGTVMRSLGQNPTEAELQDMINEVDADGNGTI
DFPEFLTMMARKMKDTDQIAEFKEAFSLFDKDGDGTITTKELGTVMRSLGQNPTEAELQDMINE
VDADGNGTIDFPEFLTMMARK

The CC-CaM protein sequence (residues S81-K148, D79-K148) is

MASEEEIREAFRVFDKDGNGYISAAELRHVMTNLGEKLTDEEVDEMIREADIDGDGQVNYEEFV
QMMTAKDTDSEEEIREAFRVFDKDGNGYISAAELRHVMTNLGEKLTDEEVDEMIREADIDGDG
QVNYEEFVQMMTAK

Protein preparation

A single colony of a protein construct in BL21(DE3) cells is selected with a toothpick and put
into 20 mL of Luria Broth (Fisher Scientific Company L.L.C., Pittsburg, PA) with appropriate
antibiotic for the construct, either carbenicillin or kanamycin. These 'starter cultures' are grown
for 7-8 hours, but no more than 8 hours, at 37°C and stored in the fridge overnight. The next
day, the starter culture is used to inoculate 750 mL LB in a 2 L flask and the cultures are grown
at 37°C in an orbital shaker to an optical density of 0.6 absorbance units at a wavelength of 600
nm. The temperature of the shaker is either kept at 37°C for SKp protein or changed to 25°C for
CaM constructs. Cultures are then induced with 0.5 mM isopropyl β-D-1-thiogalactopyranoside
and induced for 2-3 hours for SKp or for 16-18 hours for CaM constructs. Proteins were purified
using established procedures (Halling et al., 2014). Only proteins that migrate as a single band
on a 20% Laemmli gel, i.e., negligible contaminants and degradation in sample, are used in
experiments.

 HEK293 Cell Maintenance and Transfection.

HEK293 cells (ATCC, Manassas, VA) were grown according to standard procedures. Cells were
cultured at 37°C in a 5% CO2 atmosphere in Dulbecco's modified Eagle's medium without
sodium pyruvate and with l-glutamine, 1% penicillin/streptomycin, 1% GlutaMAX, and 10% fetal
bovine serum (ThermoFisher Scientific). Cell lines were split with trypsin/EDTA in Hanks'
balanced salt solution (ThermoFisher Scientific) up to 20–25 cycles. For experimental
electrophysiological recordings, cells were transiently cotransfected (Lipofectamine 3000;
Invitrogen) with the rat SK2 channel and the E1Q calmodulin mutant. For control experiments,
cells were transfected with either the rat SK2 channel and wildtype calmodulin, or the rat SK2
channel alone. Enhanced green fluorescent protein was also cotransfected as a marker in all
cases. Electrophysiological recordings were done 16–72 hours after transfection.

Patch-Clamp Recordings.

Patch clamp recordings of SK currents were performed at room temperature (22–24°C) using
an inside-out configuration. Recording electrodes were fabricated with borosilicate glass pipets
(A-M Systems) and pressure/fire-polished before use. Open electrode resistance was 2–4 MΩ
in the bath solution. An Axopatch 200A amplifier (Molecular Devices San Jose, CA) coupled
with an ITC-18 interface (HEKA) was used to record data. Acquisition software
(PATCHMASTER; HEKA; Lambrecht, Germany) was used to sample data at 50 kHz, which
were low-pass filtered at 1 kHz using the built-in filter of the amplifier. SK currents were elicited
by stepping to a voltage of -60 mV for 1 second every 3 seconds until recording was paused for
solution changes.

For all recordings, the extracellular (pipette) solution contained the following: 2 mM KCl, 136
mM KOH, 20 mM HEPES, and 2 mM $MgCl_2$, adjusted to pH 7.2 with $MeSO_3H$.The intracellular
(bath) solution contained the following: 6 mM KCl, 132 mM KOH, and 20 mM HEPES, adjusted
to pH 7.2 with $MeSO_3H$. Ca^{2+} chelators were used in the intracellular solution to control free
Ca^{2+} and (+)-(Crown-6)-2,3,11,12-tetracarboxylic acid (Synquest Laboratories) was used to
chelate any contaminating barium to prevent barium block of channels. 3.06 mM $CaCl_2$ and 5
mM HEDTA were added to the internal solution for an estimated 10 μM free Ca^{2+}(MaxChelator)
and was verified by measurement with a Ca^{2+}-sensitive electrode. Solution with 5 mM EGTA
and no added Ca^{2+} was considered as nominally Ca^{2+} free and was used to estimate leak
current levels. All solutions containing calmodulin protein were made using 10 μM free Ca^{2+}
internal solution and the concentration of either WT-CaM or non-WT constructs was 20 μM. For
solution changes, a pressurized SmartSquirt (Automate Scientific) micro-perfusion system was
used. When switching between Ca^{2+} solutions with or without exogenous calmodulin protein, the
patch pipette was moved in close proximity to the SmartSquirt perfusion pencil (250 μM
opening) which was immersed in the bath chamber.

The experimental protocol was the following: patches were excised from cells immersed in a
bath chamber containing 10μM free Ca^{2+}. Using the SmartSquirt, the solution was changed
between Ca^{2+} free (to estimate leak and confirm that current was Ca^{2+}-dependent) and 10μM
free Ca^{2+} until the current was stable across solution changes. Then, the exogenous CaM
protein was perfused on using the SmartSquirt, which varied in the amount of time from patch to
patch. Early experiments had more variability than later experiments, in the time CaM protein
was perfused on, due to the need to determine how long it took to see a measurable effect on
current levels. If it was an experiment where two CaM constructs were being applied to the
patch, the non-WT CaM always was applied first, and the solution was changed to Ca^{2+} free

solution and then back to 10µM free Ca^{2+} before finally perfusing on WT CaM. All
measurements of current change, in experiments where both non-WT CaM and WT-CaM were
applied to the patch, were done over the same amount of time for both CaM constructs.
However, the amount of time each construct was being applied varied from patch to patch due
to the need to conserve exogenous protein and the variability of patch lifetime.

In some experiments, where SK current ran up slightly during the course of the experiment, a
drift correction was performed by fitting a line to the drift and subtracting it from the average
current level.

Light scattering

Stock protein samples were centrifuged at 17 kg prior to dilution into the working buffer. Protein
concentrations were measured in a cuvette with 1cm path length and using a UV-2600 UV-
spectrophotometer (Shimadzu Scientific Instruments, Inc, Columbia, MD) to collect protein
spectra. The average absorbance was subtracted by baseline averages from 330nm to 350 nm.
Protein absorbance was converted to concentrations using predicted extinction coefficients (ε)
based on numbers of tyrosines and tryptophans using either ProtParam (Gasteiger et al., 2005)
or phenylalanines as follows. Free phenylalanine ε was determined by weighing desiccated L-
phenylalanine with milligram accuracy, dissolving into 1 mL of 18 MΩ water and collecting
spectra from 350 nm to 250 nm. The ε was computed by the relation ε=A/Lc, where
A=Absorbance, L is the path-length = 1 cm, c is the molar concentration of L-phenylalanine. L-
phenylalanine has a spectral maximum at 258 nm, and ε was measured at this wavelength for
phenylalanine. We assume a linear relation of the number of phenylalanines in the protein with
total ε. The values we used for ε in units of $M^{-1} \cdot cm^{-1}$ (at given wavelength) are ε = 6990 SKp
(280 nm), 3020 WT-CaM (277 nm), 975 N-CaM (258 nm), 3020 C-CaM (277 nm), 1950 NN-
CaM (258 nm), 5960 CC-CaM (277 nm), and 3020 for CaM (E34Q)- and CaM (E12Q) (277 nm).

Working solutions for light scattering were filtered through Whatman® 0.02 µm aluminum oxide
filters (GE Healthcare Bio-Sciences, Pittsburgh, PA). All polystyrene bottles (Nalgene,
Rochester, NY) for solution storage were rinsed 10 times in 18.2 MΩ water. Buffer reagents
were purchased from Sigma-Aldrich (subsidiary of Merck). Azide was added as a preservative.
Low Ca^{2+} (< 5nM free Ca^{2+}) buffer contained 100 mM NaCl, 10 mM HEPES, 5 mM EGTA,
0.02% NaN_3 pH 7.0. High Ca^{2+} buffer contained 100 mM NaCl, 10 mM HEPES, 5 mM $CaCl_2$,
0.02% NaN_3 pH 7.0. Some controls were performed in 10 µM free Ca^{2+} buffer which contained
100 mM NaCl, 10 mM HEPES, 5 mM HEDTA, 2.8 mM total $CaCl_2$, 0.02% NaN_3 pH 7.0. The
final free Ca^{2+} concentration was verified with an Orion 93-20 calcium half-electrode (Thermo
Electron Corporation, Waltham, MA).

Stock protein solutions were diluted into a working buffer, 0.096 mg/mL SKp and 0.09 mg/mL for
any CaM construct. Diluted samples were again centrifuged for 6.5 kg for 20 minutes.
Composition gradient multi-angle light scattering experiments were carried out as previously
described (Halling et al., 2014) on a Calypso-II (Wyatt Technology, Santa Barbara, CA) pump
delivery system to a Dawn Heleos-II (Wyatt) multi-angle 18 detector system held at 25°C. An
inline UV monitor 170S was used to monitor protein absorbance at 280 nm (Bio-Rad, Hercules,
CA). Monomeric protein concentration was evaluated in each experiment by stepping
concentrations and monitoring UV absorbance and light-scattering intensity. The molar masses
of monomeric SKp and CaMs are known and used to determine if any small adjustments were
needed to correct SKp or CaM concentrations. After SKp and CaM concentrations are verified,
analysis of the 'cross-over' phase of the experiment was carried out using Calypso® software

(Wyatt) as described previously (Halling et al., 2014), except that we determined that fitting for
'incompetent fraction' was unnecessary. For small molecules relative to the light wavelength
and at low concentrations, light scattering signals are not angular dependent and corrections
from virial coefficients at high concentrations are negligible. Under these simplified conditions,
the weight averaged molar mass of all particles in solution, $M_z = R(\theta,c)/(P(\theta) \cdot K^* \cdot c)$, is directly
computed (Wyatt, 1993). $R(\theta,c)$ is the excess Rayleigh ratio of the light-scattering caused by the
dissolved protein in a solvent. It is the ratio of protein plus solvent to solvent alone. For small
molecules $R(\theta,c)$ is virtually angle independent ($P(\theta) \approx 1$), but changes as a function of protein
concentration, c. The value $K^* = 4\pi^2(dn/dc)^2 n_o^2/(N_A \lambda_o^2)$ is constant. N_A is Avagadro's number, λ_o
is the laser wavelength in a vacuum (λ_o=660 nm), n_o is the solvent refractive index ($n_o \approx 1.33$),
and dn/dc is the proteins refractive index increment. We measured the dn/dc of CaM to be
0.186 on an Optilab rEX refractometer (Wyatt) using Astra 7.1.1.3 software, which is consistent
with published experimental (0.185) and computed values (0.190) for proteins (Khago et al.,
2018; Zhao et al., 2011).

Analyses on light-scattering samples were performed in the software Calypso-II Version
2.2.4.32 (Wyatt). Observed light-scattering intensity at different molar ratios of protein and
ligand was used to model the proportions of free vs bound proteins in the sample. The total
concentrations of added protein are known and verified before and after experiments as part of
the method. Their monomeric molar masses are known from sequence and were verified by
mass-spectrometry on an AB Voyager-DE PRO MALDI-TOF instrument (Applied Biosystems
now ThermoFisher). Stoichiometries are assumed to be comprised of known molar masses of
SKp and CaM. Free and bound protein concentrations were modelled across molar ratios as
previously described (Halling et al., 2014). The concentrations of free protein and complexed
protein were fitted to M_z by Levenberg-Marquardt fitting of the equilibrium association constants,
K_A, for each complex.

Tryptophan emission spectra

SKp contains a single, native tryptophan at position 432, whereas CaM has no tryptophan.
Between the two proteins, there are 3 tyrosines, but tryptophan fluorescence is dominant over
tyrosine. W432 has been shown to contact the C-lobe of CaM in multiple structures (Nam et al.,
2017; Schumacher et al., 2001; Zhang et al., 2013, 2012a), and its spectra have been used to
assess WT-CaM binding to SKp (Keen et al., 1999). Because of its position and potential role in
forming a CaM binding site, W432 spectra could be used as a probe in assessing CaM binding
under more conditions.

1 μM SKp (W432) plus/minus 1 μM CaM (no tryptophan) were dissolved in the same working
solutions used for CG-MALS. Fluorescence spectra were collected in a 1 cm pathlength
Spectrosil® quartz cuvette (Starna, Atascadero, CA) on a QM-400 PTI spectrofluorometer
(Horiba Scientific, Kyoto, Japan) containing a 75 W xenon arc lamp and an R928 side-on
photomultiplier (PMT) detector (Hamamatsu Photonics K.K., Shizuoka, Japan). The light path
included an excitation monochromator, a 2 nm bandwidth excitation slit, a cuvette with sample,
a 305 nm cut-on emission filter, a 1 nm bandwidth emission slit, emission monochromator, and
PMT. Fluorescence was excited at 265 nm and its emission was scanned in 1 nm steps from
315-400 nm with 1 second signal integrations. All scans are buffer background subtracted.
CaMs, including those with tyrosines, were measured for controls. Averaged spectra were
subjected to binomial smoothing using Igor Version 5.0.5.7 (Wavemetrics Inc., Lake Oswego,
OR).

Figures

Figure 1-CaM as an SK Ca²⁺ sensor. (A) Cartoon representation of the SK4 K⁺ ion channel
shown as cylindrical helices and molecular surface depictions of CaM (pdb 6CNN). For clarity,
only 2 of 4 subunits and CaMs are shown. Subunit 1 is yellow and subunit 3 is green. The N-
lobe surface of CaM is vermillion, and the C-lobe surface of CaM is blue. Transmembrane
helices are shown as S1-S6, and SK C-domain helices are shown as HA-HC. The peptide used
in the binding studies, SKp, includes HA, HB, and part of HC. Diagonal subunits of SK4 bind a
CaM in EM structure. Intracellular helix 4-5A between S4 and S5 of transmembrane segments
bind in a pocket of the N-lobe of CaM. C-domain helices HA and HB of the cross-subunit bind
the C-lobe of CaM. (B) Bottom views of SK4 showing all 4 subunits showing CaMs bound, left
side, or with CaM models removed, right side. The colors are the same as in panel A. The
subunits that are not depicted in A are grey. On the right side, the SK sites that bind CaM are
colored the same as the CaM lobes they contact. Panels (C-F) demonstrate the lobes of CaM
can reorient to form diverse conformation states. The lobes of CaM can when bound to SK4
(pdb 6CNN) form a W-shape in C, when not bound to protein (pdb 1CLL) form an S-shape in D,
when bound to an SK2 peptide (pdb 1G4Y) form a C-shape in E, or when bound to an MLCK
peptide (pdb 1CDL) form an O-shape in F. (G) The N-sensor hypothesis of SK activation by
CaM claims the N-lobe of CaM to be the Ca²⁺ sensor. The N-lobe binds a Ca²⁺ dependent site
when Ca²⁺ binds to it. The C-lobe remains anchored to a site with no local change. In the NC-
sensor sensor hypothesis, both lobes of CaM are functional Ca²⁺ sensors. At trace Ca²⁺, the C-
lobe dominates the interaction. At high Ca²⁺, a different binding mode is revealed that binds both
Ca²⁺ bound lobes. The binding sites for each lobe of CaM need not be on the same sub-unit and
the depiction is meant to emphasize that SK responds to Ca²⁺ binding to both lobes of CaM.

Figure 2-*WT-CaM recovers SK2 current in HEK293 cells*. (A) cartoon depicts patch excision at
time zero (a) and moving an electrode into the flow path from a fast solution exchanger (b-d).
For simplicity, only 3 of 8 solution positions are depicted. The illustration in (A) best describes
the conditions for the experiment in panel (D). The scale bars in B-D indicate 0.13 nA and 50 s
for horizontal and vertical lines respectively. (B-D) Average current level at -60 mV measured
every 3 s from a patch excised from HEK 293 cells transfected with rSK2 only in (B),
cotransfected with excess WT-CaM in (C), or cotransfected with excess mutant CaM (E1Q) in
(D). Note that in (B) rSK2 contacted endogenous CaM, and that in both (C) and (D) WT- and
mutant CaM are massively overexpressed relative to endogenous CaM. Traces show rSK2
current before (black) and after (blue) application of exogenous WT-CaM protein. Time point
labels in (B-D) are the same as in (A): a. patch excision, b. movement of electrode to the Ca^{2+}
solution-exchange stream, c. First application of EGTA to find basal patch current, d. Timepoint
at which purified, recombinant WT-CaM applied shown in blue trace, e. Timepoint at which
fractional recovery is measured. WT-CaM is indicated by a dumbbell with an N- and a C-lobe.
E1Q is depicted as a dumbbell with an 'x' in the upper lobe to indicate where Ca^{2+} binding is
disrupted.

Figure 3-CG-MALS assays of SKp binding full-length CaM constructs. A 1/1 complex of
SKp/WT-CaM is 28 kDa and indicated by blue arrows on the left panels. Each row assays the
complex formation of SKp (P) with a specific CaM construct indicated by labels: Row A) WT-
CaM, B) Double N-lobe CaM, C) Double C-lobe CaM, D) C-lobe Ca^{2+} mutant CaM, and E) N-
lobe Ca^{2+} mutant CaM. A-E) Weight average molar mass as a function of molar ratio of
CaM:SKp, black circles 5 mM EGTA, vermillion circles 2 mM $CaCl_2$. Error bars show standard
error of the mean. Solid curves connecting data points are simulations using association
constants derived from fits. Gray lines show what points would follow if there was no interaction.
Dashed lines in B, C, D and E are the same curves as in panel A for comparison. Panels F-J) in
5 mM EGTA and K-O) in 2 mM Ca^{2+} show fitted mole fractions of different complexes required
in the experimental solution to produce the data in A-E. Shaded regions are within the standard
error of the mean. The gray regions (P) and (C) correspond to fraction of unbound SKp (P) or

851   CaM (C) protein. The arrows along the bottom indicate where the total molar ratios of SKp:CaM

852   are 2:1 green, 1:1 blue, and 1:2 purple.

Figure 4-Both lobes can alter the environment of W432 of SKp. Shown are averaged fluorescent
emission spectra from excitation at 265 nm. Shaded regions are within the standard error of the
mean. Rows of panels have the same concentration of Ca^{2+}. Top row was measured in 5 mM
EGTA (<5 nM Ca^{2+}), middle row at 10 µM free Ca^{2+}, bottom row in 5 mM Ca^{2+}. A-C) SKp (P)
emission spectra are shaded gray. SKp emission spectra with equimolar CaM are shaded blue.
D-F) SKp emission spectra with equimolar N-CaM (gray) or C-CaM (blue). G-I) SKp emission
spectra with equimolar CaM (E12Q) (gray) or CaM (E34Q) (blue). Controls of CaM emission
without SKp present are included in most panels. Purple diamonds and nearby values indicate
the wavelength (nm) where the peak average intensity was located.

Figure 5-*Comparison of fractional SK2 current recovery with various CaM constructs.* Column
A) Average current levels at -60 mV measured every 3 s from patches excised from cells
cotransfected (CT) with rSK2 and mutant CaM (E1Q). Basal EGTA (open black circles) and
Ca^{2+} (filled black circles) currents are established, exogenous non-WT CaM protein is applied
(R1-vermillion traces), washed out (black traces again), WT-CaM protein is applied (R2-blue
traces), then washed out (black traces). Each row (i-vi) is an experiment with the non-WT CaM
as indicated by the vermillion labels under R1. Column B) Each bar represents a different patch.
The blue portion of the bar represents the fraction of current recovered by exogenous WT-CaM,
the vermillion portion represents the fraction recovered by exogenous non-WT CaM, and the
unfilled portion represents the fraction of drift in current at 10µM free Ca^{2+} after application of
non-WT CaM and before application of WT-CaM. The grey rectangles to the right of certain
experiments indicate that they deviated slightly from the experimental protocol described in the
Materials and Methods section, primarily in solution changes before and after non-WT CaM.
Column C) Fraction of SK current recovered with the application of non-WT CaM constructs
relative to WT-CaM. Each open black circle represents a different patch, and the black vertical
bar is the median fractional recovery. In B, fractional recovery extended beyond 1 and below 0
in some experiments due to current drifting up or down slightly across solution changes and
over the course of recovery. Similarly, in C, fractional recovery for some experiments was below
0 due to current decreasing slightly during application of non-WT CaM construct. In paired T-
tests, rows i-v were not significantly different from each other (each p>0.01), but row vi was
different from each other row (each p<0.01).

Figure 6-CG-MALS assays of SKp binding to half-CaM molecules. A-B) Weight average molar mass as a function of molar ratio of half-CaM:SKp, black circles 5 mM EGTA, vermillion circles 2 mM CaCl$_2$. Error bars show standard error of the mean. Lines connecting data points are simulations using association constants calculated from fits. Panels C-D) in 5 mM EGTA and E-F) in 2 mM Ca^{2+} show fitted mole fractions of free or bound proteins using the data in A-B. Shaded regions around the curve are within the standard error of the mean.

Figure 7-Only full-length CaM can bind to form the C-P-C complex. A-B) Weight average molar
mass of complex as a function of added CaM. Note that the weight average molar mass
decreases at higher concentrations of unbound protein relative to a complex. Full-length, WT-
CaM was premixed with SKp at a 1:1 molar ratio to form the P-C complex. The same amount of
P-C complex is mixed with increasing concentrations of WT-CaM (A) or half-CaM (B), N-CaM
open blue circles, C-CaM closed vermillion circles. Solid line shows predicted weight average
molar mass of C-P-C at higher concentrations of CaM using association constants used from
fits to Figure 3A. The dashed line shows prediction if P-C stays bound but no additional
molecules bind. C) Weight average molar mass, closed circles, as a function of molar ratio of
total CaM:SKp at 2 mM Ca^{2+}. CaM_T (total CaM) has equal parts of N-CaM and C-CaM. blue line
is the simulation N-CaM binding to SKp from Figure 6A, vermillion line C-CaM binding to SKp
from Figure 6B, and purple line is a simulation if both N-CaM and C-CaM bind SKp together.

Figure 8-Shematics showing different configurations of stoichiometries between WT-CaM and
SKp. SKp is represented as a black square; CaM is represented as a gray circle. Different
stoichiometries are favored at different molar ratios of peptide and CaM in solution. Ca^{2+} allows
peptide and CaM to bind with additional stoichiometries

.   922

Table 1. Quantification of effects exogenous CaM protein has on SK current recovery

CaM Construct	n	Fraction of Recovered Current (Median)	Fraction of Recovered Current (Mean)	Standard Deviation
WT	25	.95	.86	.21
CaM (E34Q)	4	.54	.50	.16
N-CaM	4	.02	.05	.09
C-CaM	4	.007	.03	.08
N-CaM + C-CaM	5	.05	.07	.06
Double N-lobe	4	.09	.08	.04
Double C-lobe	4	.21	.19	.12

Table 2. Fitted association constants, K_A, from CG-MALS, shown as 'mean logK_A (standard error) n', where n = number of trials

SKp + CaM construct	1SKp/1CaM (P-C)		2SKp/1CaM (P-C-P)		1SKp/2CaM (C-P-C)		2SKp/2CaM (2x P-C)	
	$-Ca^{2+}$	$+Ca^{2+}$	$-Ca^{2+}$	$+Ca^{2+}$	$-Ca^{2+}$	$+Ca^{2+}$	$-Ca^{2+}$	$+Ca^{2+}$
WT-CaM	6.9(0.2)5	10.4(0.2)5	12.2(0.1)5	17.0(0.2)5	--	16.0(0.3)5	--	--
NN-CaM	*6.4(0.1)3	7.0(0.1)4	11.5(0.1)3	12.5(0.1)4	--	12.3(0.1)4	--	--
CC-CaM	9.9(0.1)3	9.7(0.9)3	15.1(0.5)3	16.5(1.0)3	--	14.8(0.9)3	--	--
N-CaM	5.1(0.1)3	8.9(0.7)3	--	--	--	--	--	--
C-CaM	7.6(0.2)3	8.1(0.7)3	--	--	--	--	--	--
CaM (E12Q)	8.2(0.7)3	8.8(0.8)3	13.2(0.7)3	14.0(0.9)3	--	--	--	--
CaM (E34Q)	8.3(0.8)3	9.1(0.1)6	13.1(0.9)3	15.8(0.1)6	--	--	--	23.3(0.1)6

* Italicized values fall outside the limit of detection for modelling K_A, or fits are ill constrained

References

Attri, A.K., Minton, A.P., 2005. Composition gradient static light scattering: A new technique for rapid detection and quantitative characterization of reversible macromolecular hetero-associations in solution. Anal. Biochem. 346, 132–138. https://doi.org/10.1016/j.ab.2005.08.013

Babu, Y.S., Sack, J.S., Greenhough, T.J., Bugg, C.E., Means, A.R., Cook, W.J., 1985. Three-dimensional structure of calmodulin. Nature 315, 37–40. https://doi.org/10.1038/315037a0

Banerjee, R., Yoder, J.B., Yue, D.T., Amzel, L.M., Tomaselli, G.F., Gabelli, S.B., Ben-Johny, M., 2018. Bilobal architecture is a requirement for calmodulin signaling to $Ca_V1.3$ channels. Proc. Natl. Acad. Sci. 115, E3026. https://doi.org/10.1073/pnas.1716381115

Barrett, E.F., Barret, J.N., 1976. Separation of two voltage-sensitive potassium currents, and demonstration of a tetrodotoxin-resistant calcium current in frog motoneurones. J. Physiol. 255, 737–774. https://doi.org/10.1113/jphysiol.1976.sp011306

Bauer, C.K., Schneeberger, P.E., Kortüm, F., Altmüller, J., Santos-Simarro, F., Baker, L., Keller-Ramey, J., White, S.M., Campeau, P.M., Gripp, K.W., Kutsche, K., 2019. Gain-of-Function Mutations in KCNN3 Encoding the Small-Conductance Ca2+-Activated K+ Channel SK3 Cause Zimmermann-Laband Syndrome. Am. J. Hum. Genet. 104, 1139–1157. https://doi.org/10.1016/j.ajhg.2019.04.012

Black, D.J., Halling, D.B., Mandich, D.V., Pedersen, S.E., Altschuld, R.A., Hamilton, S.L., 2005. Calmodulin interactions with IQ peptides from voltage-dependent calcium channels. Am. J. Physiol.-Cell Physiol. 288, C669–C676. https://doi.org/10.1152/ajpcell.00191.2004

Echols, N., Morshed, N., Afonine, P.V., McCoy, A.J., Miller, M.D., Read, R.J., Richardson, J.S., Terwilliger, T.C., Adams, P.D., 2014. Automated identification of elemental ions in macromolecular crystal structures. Acta Crystallogr. D Biol. Crystallogr. 70, 1104–1114. https://doi.org/10.1107/S1399004714001308

Elad, N., Bellapadrona, G., Houben, L., Sagi, I., Elbaum, M., 2017. Detection of isolated protein-bound metal ions by single-particle cryo-STEM. Proc. Natl. Acad. Sci. 114, 11139. https://doi.org/10.1073/pnas.1708609114

Evans, T.I.A., Hell, J.W., Shea, M.A., 2011. Thermodynamic linkage between calmodulin domains binding calcium and contiguous sites in the C-terminal tail of CaV1.2. Spec. Issue 25th Anniv. Gibbs Conf. Biothermodynamics 159, 172–187. https://doi.org/10.1016/j.bpc.2011.06.007

Gasteiger, E., Hoogland, C., Gattiker, A., Duvaud, S., Wilkins, M.R., Appel, R.D., Bairoch, A., 2005. Protein Identification and Analysis Tools on the ExPASy Server, in: The Proteomics Protocols Handbook. Humana Press, pp. 571–607.

Halling, D.B., Kenrick, S.A., Riggs, A.F., Aldrich, R.W., 2014. Calcium-dependent stoichiometries of the $KCa_{2.2}$ (SK) intracellular domain/calmodulin complex in solution. J. Gen. Physiol. 143, 231. https://doi.org/10.1085/jgp.201311007

Halling, D.B., Liebeskind, B.J., Hall, A.W., Aldrich, R.W., 2016. Conserved properties of individual Ca^{2+}-binding sites in calmodulin. Proc. Natl. Acad. Sci. 113, E1216. https://doi.org/10.1073/pnas.1600385113

Harding, M.M., 2004. The architecture of metal coordination groups in proteins. Acta Crystallogr. D Biol. Crystallogr. 60, 849–859. https://doi.org/10.1107/S0907444904004081

Hirschberg, B., Maylie, J., Adelman, J.P., Marrion, N.V., 1998. Gating of Recombinant Small-Conductance Ca-activated K^+ Channels by Calcium. J. Gen. Physiol. 111, 565. https://doi.org/10.1085/jgp.111.4.565

Hugues, M., Romey, G., Duval, D., Vincent, J.P., Lazdunski, M., 1982. Apamin as a selective blocker of the calcium-dependent potassium channel in neuroblastoma cells: voltage-clamp and biochemical

characterization of the toxin receptor. Proc. Natl. Acad. Sci. 79, 1308.
https://doi.org/10.1073/pnas.79.4.1308
Keen, J.E., Khawaled, R., Farrens, D.L., Neelands, T., Rivard, A., Bond, C.T., Janowsky, A., Fakler, B.,
Adelman, J.P., Maylie, J., 1999. Domains Responsible for Constitutive and Ca^{2+}-Dependent
Interactions between Calmodulin and Small Conductance Ca^{2+}-Activated Potassium Channels. J.
Neurosci. 19, 8830. https://doi.org/10.1523/JNEUROSCI.19-20-08830.1999
Khago, D., Bierma, J.C., Roskamp, K.W., Kozlyuk, N., Martin, R.W., 2018. Protein refractive index
increment is determined by conformation as well as composition. J. Phys. Condens. Matter 30,
435101. https://doi.org/10.1088/1361-648x/aae000
Koot, B.G.P., Alders, M., Verheij, J., Beuers, U., Cobben, J.M., 2016. A de novo mutation in KCNN3
associated with autosomal dominant idiopathic non-cirrhotic portal hypertension. J. Hepatol. 64,
974–977. https://doi.org/10.1016/j.jhep.2015.11.027
Lakowicz, J.R. (Ed.), 2006. Protein Fluorescence, in: Principles of Fluorescence Spectroscopy. Springer US,
Boston, MA, pp. 529–575. https://doi.org/10.1007/978-0-387-46312-4_16
Lee, C.-H., MacKinnon, R., 2018. Activation mechanism of a human SK-calmodulin channel complex
elucidated by cryo-EM structures. Science 360, 508. https://doi.org/10.1126/science.aas9466
Lee, W.-S., Ngo-Anh, T.J., Bruening-Wright, A., Maylie, J., Adelman, J.P., 2003. Small Conductance Ca2+-
activated K+ Channels and Calmodulin: CELL SURFACE EXPRESSION AND GATING. J. Biol. Chem.
278, 25940–25946. https://doi.org/10.1074/jbc.M302091200
Li, W., Halling, D.B., Hall, A.W., Aldrich, R.W., 2009. EF hands at the N-lobe of calmodulin are required
for both SK channel gating and stable SK–calmodulin interaction. J. Gen. Physiol. 134, 281.
https://doi.org/10.1085/jgp.200910295
Liang, H., DeMaria, C.D., Erickson, M.G., Mori, M.X., Alseikhan, B.A., Yue, D.T., 2003. Unified Mechanisms
of Ca2+ Regulation across the Ca2+ Channel Family. Neuron 39, 951–960.
https://doi.org/10.1016/S0896-6273(03)00560-9
Linse, S., Helmersson, A., Forsén, S., 1991. Calcium binding to calmodulin and its globular domains. J.
Biol. Chem. 266, 8050–8054.
Maune, J.F., Beckingham, K., Martin, S.R., Bayley, P.M., 1992. Circular dichroism studies on calcium
binding to two series of calcium binding site mutants of Drosophila melanogaster calmodulin.
Biochemistry 31, 7779–7786. https://doi.org/10.1021/bi00149a006
Meador, W., Means, A., Quiocho, F., 1992. Target enzyme recognition by calmodulin: 2.4 A structure of a
calmodulin-peptide complex. Science 257, 1251. https://doi.org/10.1126/science.1519061
Moolenaar, W.H., Spector, I., 1979. The calcium current and the activation of a slow potassium
conductance in voltage-clamped mouse neuroblastoma cells. J. Physiol. 292, 307–323.
https://doi.org/10.1113/jphysiol.1979.sp012852
Mukherjea, P., Maune, J.F., Beckingham, K., 1996. Interlobe communication in multiple calcium-binding
site mutants of Drosophila calmodulin. Protein Sci. 5, 468–477.
https://doi.org/10.1002/pro.5560050308
Nam, Y.-W., Cui, M., Orfali, R., Viegas, A., Nguyen, M., Mohammed, E.H.M., Zoghebi, K.A., Rahighi, S.,
Parang, K., Zhang, M., 2021. Hydrophobic interactions between the HA helix and S4-S5 linker
modulate apparent Ca2+ sensitivity of SK2 channels. Acta Physiol. 231, e13552.
https://doi.org/10.1111/apha.13552
Nam, Y.-W., Orfali, R., Liu, T., Yu, K., Cui, M., Wulff, H., Zhang, M., 2017. Structural insights into the
potency of SK channel positive modulators. Sci. Rep. 7, 17178. https://doi.org/10.1038/s41598-
017-16607-8
Newman, R.A., Van Scyoc, W.S., Sorensen, B.R., Jaren, O.R., Shea, M.A., 2008. Interdomain cooperativity
of calmodulin bound to melittin preferentially increases calcium affinity of sites I and II. Proteins
Struct. Funct. Bioinforma. 71, 1792–1812. https://doi.org/10.1002/prot.21861

Ogawa, Y., Tanokura, M., 1984. Calcium Binding to Calmodulin: Effects of Ionic Strength, Mg^{2+}, pH and
Temperature. J. Biochem. (Tokyo) 95, 19–28.
https://doi.org/10.1093/oxfordjournals.jbchem.a134584
Orfali, R., Nam, Y.-W., Nguyen, H.M., Rahman, M.A., Yang, G., Cui, M., Wulff, H., Zhang, M., 2022.
Channelopathy-causing mutations in the S45A/S45B and HA/HB helices of KCa2.3 and KCa3.1
channels alter their apparent Ca2+ sensitivity. Cell Calcium 102, 102538.
https://doi.org/10.1016/j.ceca.2022.102538
Orlova, E.V., Saibil, H.R., 2010. Chapter Twelve - Methods for Three-Dimensional Reconstruction of
Heterogeneous Assemblies, in: Jensen, G.J. (Ed.), Methods in Enzymology. Academic Press, pp.
321–341. https://doi.org/10.1016/S0076-6879(10)82013-0
Persechini, A., White, H.D., Gansz, K.J., 1996. Different Mechanisms for Ca Dissociation from Complexes
of Calmodulin with Nitric Oxide Synthase or Myosin Light Chain Kinase. J. Biol. Chem. 271, 62–
67. https://doi.org/10.1074/jbc.271.1.62
Persechini, A., Yano, K., Stemmer, P.M., 2000. Ca2+ Binding and Energy Coupling in the Calmodulin-
Myosin Light Chain Kinase Complex. J. Biol. Chem. 275, 4199–4204.
https://doi.org/10.1074/jbc.275.6.4199
Roh, S.-H., Hryc, C.F., Jeong, H.-H., Fei, X., Jakana, J., Lorimer, G.H., Chiu, W., 2017. Subunit
conformational variation within individual GroEL oligomers resolved by Cryo-EM. Proc. Natl.
Acad. Sci. 114, 8259. https://doi.org/10.1073/pnas.1704725114
Schumacher, M.A., Rivard, A.F., Bächinger, H.P., Adelman, J.P., 2001. Structure of the gating domain of a
Ca2+-activated K+ channel complexed with Ca2+/calmodulin. Nature 410, 1120–1124.
https://doi.org/10.1038/35074145
Steiner, R.F., Lambooy, P.K., Sternberg, H., 1983. The dependence of the molecular dynamics of
calmodulin upon pH and ionic strength. Arch. Biochem. Biophys. 222, 158–169.
https://doi.org/10.1016/0003-9861(83)90513-1
Tadross, M.R., Dick, I.E., Yue, D.T., 2008. Mechanism of Local and Global Ca2+ Sensing by Calmodulin in
Complex with a Ca2+ Channel. Cell 133, 1228–1240. https://doi.org/10.1016/j.cell.2008.05.025
Villarroel, A., Taglialatela, M., Bernardo-Seisdedos, G., Alaimo, A., Agirre, J., Alberdi, A., Gomis-Perez, C.,
Soldovieri, M.V., Ambrosino, P., Malo, C., Areso, P., 2014. The Ever Changing Moods of
Calmodulin: How Structural Plasticity Entails Transductional Adaptability. J. Mol. Biol. 426, 2717–
2735. https://doi.org/10.1016/j.jmb.2014.05.016
Westerlund, A.M., Delemotte, L., 2018. Effect of Ca2+ on the promiscuous target-protein binding of
calmodulin. PLOS Comput. Biol. 14, e1006072. https://doi.org/10.1371/journal.pcbi.1006072
Williams, B., Haeseleer, F., Lee, A., 2018. Splicing of an automodulatory domain in $Ca_v1.4$ Ca^{2+} channels
confers distinct regulation by calmodulin. J. Gen. Physiol. 150, 1676.
https://doi.org/10.1085/jgp.201812140
Wyatt, P.J., 1993. Light scattering and the absolute characterization of macromolecules. Anal. Chim. Acta
272, 1–40. https://doi.org/10.1016/0003-2670(93)80373-S
Xia, X.-M., Fakler, B., Rivard, A., Wayman, G., Johnson-Pais, T., Keen, J.E., Ishii, T., Hirschberg, B., Bond,
C.T., Lutsenko, S., Maylie, J., Adelman, J.P., 1998. Mechanism of calcium gating in small-
conductance calcium-activated potassium channels. Nature 395, 503–507.
https://doi.org/10.1038/26758
Zhang, B.-M., Kohli, V., Adachi, R., López, J.A., Udden, M.M., Sullivan, R., 2001. Calmodulin Binding to the
C-Terminus of the Small-Conductance Ca2+-Activated K+ Channel hSK1 Is Affected by Alternative
Splicing. Biochemistry 40, 3189–3195. https://doi.org/10.1021/bi001675h
Zhang, C., Cantara, W., Jeon, Y., Musier-Forsyth, K., Grigorieff, N., Lyumkis, D., 2019. Analysis of discrete
local variability and structural covariance in macromolecular assemblies using Cryo-EM and
focused classification. 75th Birthd. Christ. Colliex 85th Birthd. Arch. Howie 75th Birthd. Hannes

Lichte PICO 2019 - Fifth Conf. Front. Aberration Corrected Electron Microsc. 203, 170–180.
1082   https://doi.org/10.1016/j.ultramic.2018.11.016
1083   Zhang, M., Abrams, C., Wang, L., Gizzi, A., He, L., Lin, R., Chen, Y., Loll, P.J., Pascal, J.M., Zhang, J., 2012a.
1084   Structural Basis for Calmodulin as a Dynamic Calcium Sensor. Structure 20, 911–923.
1085   https://doi.org/10.1016/j.str.2012.03.019
1086   Zhang, M., Pascal, J.M., Schumann, M., Armen, R.S., Zhang, J.-F., 2012b. Identification of the functional
1087   binding pocket for compounds targeting small-conductance Ca2+-activated potassium channels.
1088   Nat. Commun. 3, 1021.
1089   Zhang, M., Pascal, J.M., Zhang, J.-F., 2013. Unstructured to structured transition of an intrinsically
1090   disordered protein peptide in coupling Ca^{2+}-sensing and SK channel activation. Proc. Natl. Acad.
1091   Sci. 110, 4828. https://doi.org/10.1073/pnas.1220253110
1092   Zhao, H., Brown, P.H., Schuck, P., 2011. On the Distribution of Protein Refractive Index Increments.
1093   Biophys. J. 100, 2309–2317. https://doi.org/10.1016/j.bpj.2011.03.004

A SK4 channel

Subunit 1
S1-S6

Subunit 3
S1-S6

K+

Subunit 3
C-domain

N-lobe

C-lobe

S2 S1 P S5 S6 P S1 S2 S4
S4 S5 S6
S3
4-5 B HA 4-5 B
4-5 A HB
HA HC
N HB HC
C C

B Bottom, cytosolic view

4 CaMs on SK4   Sites on SK4 for CaM

HA-HB S1-S6
S1-S6 HA-HB

C W-shape

N-lobe   C-lobe

D S-shape

E C-shape

F O-shape

G

Apo-CaM   Ca²⁺-CaM

N-sensor Theory

SK N-lobe site   SK C-lobe site

Apo-State

Ca²⁺-State

N-C Sensor Theory

Apo-State   Ca²⁺-State

Calculated and Simulated Molar Mass

Simulated Mole Fraction